1   Dichloroacetate improves hepatic and systemic energy metabolism during sepsis
2

3   Rabina Mainali[1*], Manal Zabalwi[2*], David Long[2], Nancy Buechler[1], Shalini Jain[3], Ellen
4   Quillen[2], Chia-Chi Key[2], Xuewei Zhu[2], Peter W. Stacpoole[4], Jennifer Martinez[5], Charles
5   E. McCall[2#], Matthew A, Quinn[1,2,5#]
6

[1]Department of Pathology, Section on Comparative Medicine Wake Forest School of
Medicine, Winston-Salem, North Carolina

[2]Department of Internal Medicine, Section of Molecular Medicine Wake Forest School of
Medicine, Winston-Salem, North Carolina

[3]Department of Internal Medicine, Section of Endocrinology and Metabolism Wake Forest
School of Medicine, Winston-Salem, North Carolina

[4]Division of Endocrinology, Diabetes and Metabolism and Department of Biochemistry
and Molecular Biology, University of Florida College of Medicine, Gainesville, Florida

[4]Immunity, Inflammation, and Disease Laboratory, National Institute of Environmental
Health Sciences (NIEHS), National Institutes of Health (NIH), Research Triangle Park,
NC
18
19   [5]Lead Contact
20
21
* these authors contributed equally
# denotes co-corresponding author

Corresponding Authors:

Matthew A. Quinn, Ph.D.
Wake Forest School of Medicine
Medical Center Blvd
Winston-Salem, NC 27157
Office: 336-713-1995
mquinn@wakehealth.edu
31
Charles E. McCall, M.D.
Wake Forest School of Medicine
Medical Center Blvd
Winston-Salem, NC 27157
Office: 336 413 8034
chmccall@wakehealth.edu
38
39
40

41   **Keywords:** sepsis, mitochondria function, liver, inflammation, metabolism, steatosis

Abstract

Dramatic metabolic reprogramming between an anabolic resistance and catabolic toler-ance occurs within the immune system in response to systemic infection with the sepsis syndrome. While metabolic tissues such as the liver are subject to end-organ damage during sepsis and are the primary cause of sepsis death, how their metabolic and energy reprogramming during sepsis state ensures survival is unclear. Employing comprehen-sive metabolomic screening, targeted lipidomic screening, and transcriptional profiling in a mouse model of septic shock, we show that hepatocyte lipid metabolism, mitochondrial TCA energetics, and redox balance are significantly reprogramed after cecal ligation and puncture (CLP). We identify increases in TCA cycle metabolites citrate, cis-aconitate, and itaconate with reduced fumarate, elevated triglyceride synthesis, and lipid droplet accumulation in the septic hepatocytes. Transcription analysis of liver tissue supports and extends the hepatocyte findings. Plasma metabolomics show systemic hypoglycemia and increased concentrations of free fatty acids, ketones and corticosterone in parallel with liver reprogramming. Strikingly, the administration of the pyruvate dehydrogenase kinase (PDK) inhibitor dichloroacetate (DCA) reverses dysregulated hepatocyte and systemic metabolism and mitochondrial dysfunction. DCA administered during sepsis arrests ano-rexia and weight loss, restores V02 levels as an index of increased carbohydrate oxida-tion and promotes physical activity. We suggest that sepsis inflicts an energy demand and supply crisis with distinct shifts in hepatocyte and systemic mitochondrial function. Targeting the mitochondrial PDC/PDK energy homeostat rebalances life-threatening en-ergy deregulation caused by bacterial sepsis.

Introduction

Sepsis represents a life-threatening condition that results from dysregulated inflammatory responses to disseminated and uncontrolled infection. If not diagnosed and treated early after its onset, sepsis may limit survival by inducing early shock and subsequent immunometabolic paralysis. Sepsis accounts for one in three hospital deaths in the U.S. and millions of deaths each year globally, highlighting its hazard to public health [1,2]. The high mortality rate associated with sepsis reflects the lack of a clinically viable molecular-based therapeutic target. Therefore, understanding the pathogenesis of sepsis at both the molecular and organismal levels is major gap in public health and of the utmost importance.

Sepsis is a biphasic phenomenon, with clearly defined acute and chronic phases [3,4]. The acute phase of sepsis is driven by high energy consumption and hyper-inflammation characterized by oxidative stress, followed by cellular reprogramming to a low energy, anti-inflammatory state of immunometabolic paralysis, with accompanying organ failure.[2,5]. In addition to inflammatory responses, dysregulation of the host's metabolism during sepsis plays a vital role in its progression and severity [6-8]. Compared to healthy individuals, energy homeostasis is significantly altered in septic patients [7,8]. The early phase of sepsis is a hypermetabolic condition marked by increased catabolism of fats, proteins and carbohydrates, associated with high rates of oxygen consumption and ATP synthesis [9-11]. Following the acute phase is a hypometabolic state where ATP production and mitochondrial respiration decrease [9]. Immune cells and some organ cells enter an energy

conserving "hibernation-like" state as a protective mechanism to lower the metabolic de-

mands of the cell and help with its recovery [6,12,13]. However, staying in this hypometabolic

state for a prolonged period can lead to organ dysfunction and failure [12,13].

Of particular interest is that, during the hyper-inflammatory anabolic phase of sep-

sis, an increase in the expression and activity of pyruvate dehydrogenase kinase 1

(PDK1) consistently occurs [4]. This enzyme is one of four PDK isoforms that reversibly

phosphorylates serine residues on pyruvate dehydrogenase complex (PDC) E1α subunit,

inhibiting the conversion of pyruvate to acetyl coenzyme A (acetyl CoA) [14]. Inhibition of

this important enzymatic activity that connects glycolysis to tricarboxylic cycle (TCA), ox-

idative phosphorylation (OXPHOS) and the lipogenic pathway is thought to be one of the

main mechanisms that is driving dysfunction of mitochondrial respiration and cell bioen-

ergetics observed during sepsis [15,16], suggesting that PDK may be a novel, druggable

target for treatment of sepsis..

There a few reports that indicate changes in hepatic metabolism during sepsis but

these studies are not conclusive [17]. Therefore, there is limited knowledge regarding the

metabolic changes and infection related liver dysfunction in response to sepsis. There-

fore, we set out to characterize the global hepatic manifestations of sepsis, with the over-

all goal of identifying master regulators of the hepatic metabolic response to polymicrobial

infection.

Results

Sepsis Impairs Hepatic Mitochondrial Metabolism

To test the long-term hepatic transcriptional changes elicited by sepsis we performed RNA-seq in whole livers thirty-hours post cecal ligation puncture (CLP). At this time point, septic mice exhibit tolerance, in which immunometaboic paralysis and end organ dysfunctions decrease survival. Ingenuity pathway analysis (IPA) of the top physiological pathways subject to alteration in response to sepsis revealed a significant increase in the acute phase response pathway, highlighting a persistent inflammatory state in the liver into the chronic phase of sepsis (Fig. 1a). Of particular interest were the findings that oxidative phosphorylation and mitochondrial dysfunction were the top two enriched pathways in the liver of septic mice (Fig. 1a). To gain further insight into the effects of sepsis on the transcriptional regulation of mitochondrial function we performed gene set enrichment analysis (GSEA) of the oxidative phosphorylation pathway (GO:0006119). Septic mice at 30 h decreased the transcriptional output of OXPHOS components, as evidenced by a negative GSEA enrichment score (Fig. 1b). Given the connection between oxidative phosphorylation and the tricarboxylic acid (TCA) cycle, we next asked if polymicrobial infection would elicit similar alterations in this pathway. In accordance with the changes observed in the oxidative phosphorylation pathway, we found a negative enrichment score for TCA cycle enzymes (GO:0006099) in the liver of CLP mice (Fig. 1c). Thus, our transcriptome data indicate that sepsis impairs mitochondrial metabolism in the liver.

Next we wanted to assess if the transcriptional changes elicited by sepsis would manifest in altered hepatic TCA cycle metabolism. Therefore, we performed global

unbiased metabolomic screening in isolated hepatocytes from control and septic mice by

Ultrahigh Performance Liquid Chromotography-Tandem Mass Spectroscopy (UPLC-

MS/MS). In line with altered transcriptional regulation of the TCA cycle, we found that

sepsis altered the relative abundance of multiple metabolites involved in the TCA cycle

(Fig. 1d). In particular, significant elevation of citrate and cis-aconitate was observed in

septic hepatocytes at 30h (Fig. 1e&f). Unlike macrophages, which shift their ratio of

succinate and α-ketoglutarate to favor succinate accumulation over α-ketoglutarate [18-20],

we found no changes in the levels of these metabolites in hepatocytes (Fig. 1G&H).

Furthermore, unlike macrophages [21], levels of fumarate decreased, but malate levels

were unchanged in hepatocytes from septic animals (Fig. 1I&J). Monocytes,

macrophages and dendritic cells reprogram the TCA cycle during acute inflammation to

a catabolic tolerance phenotype by shunting cis-aconitate to itaconate through the

enzymatic action of aconitase decarboxylase (ACOD1; also known as immune

responsive gene 1 [IRG1]) [21-24]. Accordingly, we also found that sepsis significantly

induces *ACOD1* and elevates itaconate in isolate hepatocyte preparations (Fig. 1K&L).

Sepsis Impairs Hepatic Redox Balance and Promotes Oxidative Stress

Perturbations in oxidative phosphorylation have been shown to promote the

production of reactive oxygen species (ROS) [25]. Furthermore, itaconate has been shown

to induce genes involved in oxidative stress regulation through activation of the master

NRF2 KEAP1 antioxidative pathway in macrophages [23]. Therefore, we examined if sepsis

invokes a gene expression profile involved in regulation of ROS. GSEA of the ROS

metabolic process pathway (GO:0072593) revealed a positive enrichment, highlighting

the transcriptional induction of genes involved in the regulation of oxidative stress in livers

of septic mice (Fig. 2a). Given the induction of ROS metabolic genes we wanted to

determine if sepsis alters key metabolites involved in redox balance, specifically the

cysteine-glutathione transsulfuration redox regulatory cycle. Hepatocytes isolated during

sepsis-induced tolerance did not adjust intracellular homocysteine levels (Fig. 2b),

however, cystathionine significantly decreased (Fig. 2c). Furthermore, we found a trend

for decreased cysteine levels in the liver (p=0.0534) (Fig. 2d) and significantly decreased

glycine (Fig. 2e). Most dramatically, septic mice depleted hepatocyte glutathione (Fig. 2f).

During oxidative stress, the pool of glutathione shifts from reduced glutathione (GSH) to

oxidized glutathione (GSSG). Additional evidence of increased hepatic oxidative stress

was a significant shift in the cellular glutathione pools, favoring GSSG over GSH in CLP

mice compared to sham controls (Fig. 2g).

169   Sepsis promotes hepatic steatosis

Thus far, our data revealed that sepsis induces significant impairments in hepato-

cyte OXPHOS, alterations in the TCA cycle, and induction of oxidative stress, findings

that are also metabolic hallmarks of fatty liver disease [26]. Because perturbations in global

lipid profiles occur in septic patients [11], we evaluated the effects of CLP on the hepatic

lipid metabolism pathway. In support of our hypothesis that sepsis alters hepatic lipid

metabolism, transcriptome data indicate significant repression of lipid metabolism in livers

of septic mice (Fig. 3a). Sepsis also decreased transcription output that supports fatty

acid metabolic process components in the liver, as evidenced by a negative GSEA en-

richment score (Fig. 3b).

Since hepatic lipid metabolism significantly changes during sepsis, we further in-

vestigated the effect of sepsis on both the fatty acid oxidation and biosynthetic processes

pathways. Slight changes occurred in the expression profiles of genes involved in fatty

acid oxidation in response to CLP after 30 h. In contrast, we found a striking increase in

genes involved in fatty acid biosynthesis (Fig. 3c). We then asked whether the transcrip-

tional alterations elicited by sepsis would be manifested in hepatic lipid metabolites. Con-

sistent with our transcriptional profiling, our unbiased metabolomic screening in hepato-

cytes identified lipid metabolism as the top metabolic pathway altered in sepsis (Fig.3d).

Our metabolomic screen identified increases in virtually all fatty acids and acylcarnitine

derivatives assayed (Fig. 3e). Given the consistency between our transcriptome data and

metabolomics panel, we performed lipidomic screening in the liver of sham vs CLP mice

to capture the global consequences of sepsis on hepatic lipid metabolism. Strikingly, we

find over 500 lipid species increased in septic hepatocytes (Fig.3f). As a consequence of

alterations in lipid metabolism, we observe an increase in lipid droplet formation in liver

tissues of CLP mice (Fig. 3g), confirming the presence of hepatic steatosis.

Sepsis-induced anorexia increases plasma free fatty acids

The onset of anorexia and a switch from glucose metabolism to alternative fuel

sources during sepsis has been well established [11,27,28]. Consistent with this principle [29],

septic mice housed in metabolic chambers exhibited onset of anorexia, as evidenced by

a marked decrease in total food intake (Fig. 4a). While anorexia induced by systemic

inflammation is highly conserved, the extent to which sepsis-associated anorexia is as-

sociated with altered plasma metabolites is unclear. To address this knowledge gap, we

performed global metabolomic screen in plasma from sham and CLP mice 30 hours post-

surgery. We find plasma metabolites known to be altered in the fasting state are indeed

altered in response to polymicrobial infection. Specifically, sepsis induces a significant

decrease in the relative abundance of plasma glucose compared to sham mice (Fig. 4b).

Similar to previous reports [30], increased circulating levels of corticosterone, a primary

mediator of the fasting response also occurred (Fig. 4c). During nutrient deprivation, ke-

togenesis increases circulating ketones [31]. Likewise, we found elevated levels of circulat-

ing ketone bodies 3-hydroxybutyrate, 2-hydroxybutyrate, and 2-aminobutyric in septic

mice (Fig. 4d-f). (Fig. 4d-f). Fasting also liberates free fatty acids and increases their

levels in plasma [32]. Consistent with a fasting response, sepsis increased 75 lipid species

in the circulation (Fig. 4g). Collectively, these data indicate sepsis induces a systemic

starvation state, concomitant with repressed mitochondrial energetics and nutrient

changes in hepatocytes.

PDK inhibition attenuates sepsis-induced transcriptional and metabolic changes

in the liver

218   Given the establishment of hepatocyte mitochondrial dysfunction and steatosis

during sepsis, we asked whether pharmacological targeting of the hepatic PDK/PDC axis

would mitigate the disruption of key metabolic and bioenergetic processes induced by

sepsis. We previously reported that PDK inhibition by the pyruvate analog and pan-PDK

inhibitor dichloroacetate (DCA) promotes PDC-dependent immunometabolic adaptations

to sepsis and increases survival [15,33]. Therefore, we hypothesized that PDK inhibition

would also ameliorate sepsis-induced metabolic dysfunction in the liver. To test this pos-

tulate, we administered DCA to septic mice 24 h after CLP onset and assessed the global

transcription profile of liver 6 h later via RNA-seq. DCA treatment of septic mice reversed

the majority of sepsis-regulated gene networks (Fig. 5a). DCA also restored to nearly

sham levels the top physiological pathways altered in response to sepsis, such as inflam-

mation mediators, lipid triglycerides synthesis, redox control and mitochondrial TCA func-

tion (Fig. 5b).

Given our reversal of the sepsis-regulated hepatic transcriptome by DCA, we next

determined whether it would have functional effects on the hepatic metabolome. We re-

peated our unbiased metabolomic profiling in isolated hepatocytes from septic mice and

found that TCA metabolites that accumulate in hepatocytes during sepsis— citrate, cis-

aconitate, and itaconate—returned almost to sham levels after DCA treatment (Fig. 5c).

Of particular note, we also observed that DCA treatment did not restore fumarate levels,

but rather decreased this TCA intermediate decreased even further than measured in

septic livers not exposed to the drug (Fig. 5c).

Next, we focused on metabolites proximal to the glutathione synthesis redox path-

way, given the transcriptional reversal of the conjugation to glutathione pathway in DCA

treated septic mice and distal to methionine (Fig. 5b). We found that DCA not only caused

a reversal of cystathionine and taurine low levels in septic mice, but also increased accu-

mulation of these metabolites in the liver after DCA (Fig. 5d). No significant differences

occurred in glycine. The increased levels of transsulfuration redox metabolites in DCA-

treated mice resulted in a reduced oxidative burden in hepatocytes during sepsis, as ev-

idenced by an increased GSSG/GSH ratio (Fig. 5e).

Finally, given the pronounced steatotic phenotype present during sepsis, and the

restoration of mitochondrial function and redox balance in response to DCA treatment,

we next determined whether PDK inhibition would also reverse the fatty liver phenotype

of sepsis. Metabolomics revealed broad reductions in hepatic mono-, di- and triglycerides

in septic mice treated with DCA (Fig. 5f), indicating that PDK activation can control the

development of hepatic steatosis during sepsis (Fig. 5g).

DCA rescues systemic energy crisis during polymicrobial infection

The above findings strongly suggest that PDK inhibition reverses many sepsis-

induced metabolic and transcriptional changes in the liver. We next investigated the sys-

temic effects of DCA treatment on energy metabolism in CLP mice. Plasma metabolite

screening in CLP mice revealed alterations in over 250 metabolites in response to CLP,

with the majority of these metabolites decreasing after DCA treatment (Fig. 6a&b). When

we further examined the 198 sepsis-dysregulated metabolites normalized by DCA treat-

ment, we found that the majority involve pathways of glycine and methionine metabolism

that fuel glutathione (Fig. 6c) and promote the tolerant and antioxidant catabolic phase of

sepsis [10]. Consistent with our findings in livers from septic mice, these data indicate that

DCA restores circulating metabolite levels comparable to concentrations measured in

sham controls and identify a reversal in sepsis-induced systemic energy depletion follow-

ing PDK inhibition.

Given that DCA restores levels of sepsis-dysregulated circulating metabolites, we

hypothesized that PDK inhibition might also reverse the sepsis-induced anorexia re-

sponse and starvation. To test this hypothesis, we housed septic mice treated with or

without DCA in metabolic cages for after sepsis onset. Consistent with previous reports

[34-36], CLP induced a dramatic decrease in body weight within 24 h of surgery (Fig. 6d).

Strikingly, DCA administration completely normalized body weight in septic mice (Fig. 6d)

and increased physical activity (data not shown). DCA-induced body weight correction

may result from decreasing anorexia, as food intake concomitantly increased in DCA-

treated septic mice (Fig. 6e). While modestly increases in respiratory exchange ratio

(RER) occurred, DCA restored VO2 to sham values. We interpret the VO2 increase as

indicating that PDK inhibition readjusts carbohydrate oxidation during sepsis starvation,

consistent with PDC activation (Fig. 6f&g). Together, our findings identify PDK as a sig-

nificant driver of systemic energy imbalance during sepsis and a therapeutic target for its

reversal by increasing PDC oxidation of glucose carbons. Contributing to this effect are

reversing anorexia, increasing food intake and restoring mitochondrial function (Fig. 6h).

Discussion

Considerable attention has been focused recently on the immunometabolic consequences of inmflammation in immune cells [37]. However, significant gaps exist in our understanding of the metabolic adaptations induced by systeminc inflammation from sepsis in vital organs. Addressing this limitation is critical because sepsis survival depends on restoring both organ and immune cell homeostasis following dysregulated and disseminated inflammation [13]. The present study reveals major disruption of several key aspects of hepatic mitochondrial metabolic and bioenergy functions in septic mice, including TCA cycle activity, OXPHOS, redox balance and lipid metabolism. In addition, at the organismal level, anorexia increase as an early response to sepsis followed by an altered circulating lipid profile and increased levels of corticosterone, a pattern consistent with a state of fasting or starvation. Importantly, PDK inhibition by DCA highlights the PDK/PDC as a critical gatekeeper of energy metabolism during life-threatening systemic infection. Within one hour of its administration, DCA reversed the starvation phenotype and, within six hours, restored overall mitochondrial capacity to meet the demands used to regenerate immune and organ physiologic competence. Overall, these findings support the notion that dysregulated mitochondrial reprogramming in hepatocytes contributes to a sepsis-induced systemic energy supply and demand crisis and the vunlerability of PDC to significant post-transcriptional inhibition during sepsis.

Upregulation of PDKs inactivates PDC to skew carbohydrate metabolism away from mitochondrial oxidation and promotes cytoplasmic reduction of glycolytically-derived pyruvate to lactate, as the central feature of the Warburg response. Upregulating oxidative phosphorylation also shifts immune cells to Warburg glycolysis during cell activation

[8,38], and alters levels of TCA cycle citrate, cis-aconitate, succinate, and fumarate metab-

olite signaling properties [8,20,38,39]. For example, succinate primes inflammation through

succinate dehydrogenase (SDH)-mediated ROS generation and ATP synthesis [20,38].

SDH-derived ROS enhances IL-1β production by activating HIF-1α and activating the

NLRP3 inflammasome [20]. Inhibition of SDH activity increases IL-10 production and skews

inflamed immune cells towards an anti-inflammatory response [20]. Fumarate, on the other

hand, possesses anti-inflammatory properties as it inhibits pro-inflammatory cytokine pro-

duction either by activating the NRF2 pathway or via inhibiting pathways like NF-κB and

MAPK that lie downstream of TLR signaling in immune cells [40,41]. Decreased fumarate

did not align with succinate in hepatocytes at the time point assayed.

Cis-aconitate decarboxylation to itaconate represses glycolysis [22,42] and dampens

inflammation in monocytes and macrophages [21,23]. In response to LPS stimulation,

macrophages significantly upregulate IRG1 and accumulate itaconate. Direct inhibition of

SDH by itaconate limits IL-1β levels [20]. Transcriptional regulation of the NRF2/Keap1 and

IκBζ-ATF3 pathways by itaconate also mute IL-1β expression [23,43,44]. In line with what

occurs in macrophages, we observed an increase in IRG1, citrate, cis-aconitate, and

itaconate in septic hepatocytes at 30 h. While we did not fully characterize the hepatic

function of itaconate, a recent paper demonstrates its anti-inflammatory effects during

Ischemia-reperfusion (IR) injury in the liver [45]. The deletion of IRG1 heightens

inflammation and liver damage and renders hepatocytes susceptible to oxidative injury

after I/R injury. Further, itaconate administration reduces liver damage and inflammation

associated with I/R in IRG1 KO mice, also emphasizing its anti-inflammatory and

hepatoprotective effects [45]. Given the immunomodulatory effects of itaconate and its

accumulation in the liver in our study, we hypothesize that itaconate directs hepatocyte

shifts in both TCA cycle metabolism and mitochondrial energetics. DCA reduces itaconate

levels, which we also find in a human monocyte model of sepsis [16].

ROS also may direct hepatic responses to sepsis . Similar to immune cells, liver

cells increase ROS production during sepsis [46,47]. We found that the thiol antioxidant

glutathione decreased, concomitant with an increase in oxidized glutathione, in

hepatocytes during sepsis. Overall, our findings support the notion that sepsis disrupts

hepatic redox balance, as reflected in alterations in glycine, cystathionine and taurine

levels in the liver.

Sepsis also profoundly alters lipid metabolism, resulting in significant fatty acid

and glycerol metabolic shifts in plasma, changes reported to predict prognosis in septic

patients [48]. Sirt1 and Sirt6 NAD-dependent deacetylases play an essential role in

regulating the balance between glycolysis and fatty acid oxidation [49]. Specifically, Sirt1

upregulates PGC-1-dependent fatty acid oxidation during the shift from the acute to the

late phase of sepsis. However, we do not observe a fatty acid oxidation transcription

profile in septic livers. Yet another vital lipid metabolic pathway in immune cells is *de novo*

lipogenesis [50]. Macrophages generate malonyl-CoA and subsequent malonylation of

glycolytic enzymes [51]. While we did not measure malonyl-CoA and acetyl-CoA levels, our

metabolomics screening did show an increase in hepatic citrate levels. Furthermore, our

transcriptional studies identify a cohort of genes in livers from septic animals that are

involved in fatty acid biosynthesis. Our data are consistent with human data where

patients who died from sepsis had evidence of hepatic steatosis affecting 5%-80% of liver

parenchyma [52]. These data emphasize the robust steatotic phenotype conserved

between mice and humans.

The combination of anorexia and weight loss is an important clinical manifestation

of sepsis in humans. Representative animal models demonstrated significant reductions

in food intake and body weight in response to infection [11,27,28]. Consistent with a fasting

state, we find a circulating metabolomic signature enriched in ketone bodies and

glucocorticoids with decreased glucose levels [53-57]. During the fasting period, free fatty

acids released from adipose tissues are taken up at a nonsaturable rate by the liver,

where they are packaged into triglycerides and transported out and into circulation by

very low-density lipoproteins (VLDL), which make them available as an energy source

during periods of energy crisis [54,58]. In addition to the fasting metabolomics signature

observed in septic mice, we found that sepsis significantly elevates levels of 75 lipid

species in the plasma. We propose nutrient depletion as another mechanism contributing

to the development of sepsis-induced steatosis.

PDC is a master metabolic regulator controlling the conversion of pyruvate to

acetyl-CoA in the mitochondria [14] . A part of the metabolic reprogramming that occurs in

immune cells in response to inflammatory signaling is inactivation of PDC [14,59]. PDK is a

negative regulator of the PDC, as it phosphorylates PDC and inhibits the conversion of

pyruvate to acetyl-CoA [14]. During sepsis, the expression and activity of PDK1 in immune

cells is heightened, contributing to the dysfunction of mitochondria metabolism. One

mechanism driving this observed increase is through pro-inflammatory mediators such as

LPS and interferon gamma (IFN-γ) [60]. Another possible mechanism contributing to PDK

activation in the context of sepsis is through glucocorticoid signaling [61]. In fact, starvation

is well established to activate the PDK pathway [62-66]. Based on our findings of elevated

glucocorticoid levels in both the circulation and the liver, we hypothesize the stress

hormone pathway as a contributor to PDK activation during sepsis.

We reported in a sepsis monocyte model that DCA reduced TCA cycle catabolic

effects, concurrent with increasing amino acid anaplerotic catabolism of branched-chain

amino acids, leading to increased TCA-driven anabolic energetics [59]. In the present

study, we find restoration of TCA metabolites, decreased triglyceride accumulation,

lessening of lipid synthesis and oxidative stress rebalances in septic mice after DCA

administration. Not only does DCA reverse sepsis-induced gene expressions and

metabolic profiles to baseline levels, it also leads to an increase in antioxidants, body

weight and food intake, suggesting systemic improvements in response to DCA and

reversal of anorexia. Overall we demonstrate that hepatic dysfunction and a systemic

energy crisis during sepsis improves after targeting the PDC/PDK axis with DCA.

Furthermore, we show sepsis not only affects immunity, but it dysregulates hepatic and

organism-wide lipid metabolism and redox homeostasis in the liver. The net results is an

energy crisis between low nutrient supply and high energy demand for immunity and

organ function.

Further work elucidating the mechanistic pathways involved in the dysfunction of

lipid metabolism and the impact of TCA metabolite alterations is warranted. In particular,

further investigation specifically into the PDK pathway to determine which isoform

underlies the hepatic manifestions of sepsis. This is warranted given multiple isoforms of

PDK expressed in the liver [67] and DCA inhibits multiple isoforms of PDK [68,69]. The results

of this study fill a gap in understanding how sepsis at the molecular level compromises

liver function. It also further informs the novel therapeutic targeting potential of the

PDC/PDK homeostat.

Material and Methods

Animal Experiments

Male C57BL/6J mice aged 8-10 weeks were purchased from The Jackson Laboratory

(Bar Harbor, ME). All animals were subject to a 12:12 hour dark/light cycle with *ad libitum*

access to standard rodent chow and water. Cecal ligation puncture model was used to

induce sepsis as previously described [33,70]. Briefly, cecum was ligated and punctured two

times with a 22-gauge needle. Contents were then returned and incision was closed in

two layers (peritoneum and skin). Sham operation where abdominal incision was made

but cecum not ligated or punctured was used as a control. Subcutaneous fluids (1ml

normal saline) were given to each animal. Mice were euthanized 30 hours post-surgery

for tissue collection. Dichloroacetate (DCA) (Sigma; MO, USA) was administered

(25mg/kg) intraperitoneally at various time points post-surgery: for metabolomic

screening and RNA-seq DCA was administered 24 hours post-surgery and tissues

collected 6 hours post DCA administration (30 hours post-surgery), for metabolic cages

DCA was administered immediately post-surgery.

Hepatocyte Isolation

Hepatocytes were isolated via portal vein perfusion and collagenase digestion as

previously described [71]. Following perfusion, liver cells were liberated by gentle

dissocation in DMEM (ThermoFisher; CA, USA). Cells were then filtered through nylon

mesh to remove cellular debris and connective tissue and resulting cells pelleted by

centrifugation at 50g for 1 minute. After three washes with DMEM cells were counted and

viability assessed via Trypan Blue exclusion.

RNA-sequencing

RNA was isolated from whole liver tissues using Trizol and the RNeasy RNA isolation kit

(Qiagen; MD, USA) according to manufacturer's protocol. One microgram of high quality

RNA (RIN>8) was used as a template for library generation using the Illumina TruSeq

RNA Sample Prep Kit v2 (Illumina; CA, USA) according to the manufacturer's protocol.

Generated libraries were then poly(A) enriched for mRNA prior to sequencing. Indexed

samples were sequenced at 100bp-paired-end protocol with the NovaSeq 6000

(Illumina), generating approximately 20-30 million reads per sample. Sequenced reads

were aligned to the University of California Santa Cruz (UCSC) mm10 reference genome

using STAR v2.5 as previously described [72]. The mapped read counts were quantified by

Subread feature Counts v1.5.0-p1 [73]. Differentially expressed genes (DEGs) were

determined by DESeq2 v1.14.1 [73] using a false discory rate of 0.05. Ingenuity Pathway

Analysis (Qiagen) and Gene Set Enrichment Analysis v4.0.3 (GSEA) were further used

as previously described [74].

Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy

Hepatocytes were isolated described above for metabolomic screening via Ultrahigh

Performance Liquid Chromotography-Tandem Mass Spectroscopy (UPLC-MS/MS)

(Metabolon; NC, USA). Briefly, 150-200 microliter cell pellets per animal were used as

starting material. Samples were prepared using the automated MicroLab STAR system

(Hamilton; NV, USA). Proteins were precipitated using methanol under vigorous shaking

for 2 minutes followed by centrifugation. Prior to analysis organic solvents were removed

with TurboVap (Zymark; MA, USA) and overnight storage under nitrogen. Dried samples

were reconstituted with solvents compatible with the three following analytical methods:

1) reverse phase (RP)/UPLC-MS/MS methods with positive ion mode electrospray

ionization (ESI), 2) RP/UPLC-MS/MS with negative ion mode ESI and 3) HILIC/UPLC-

MS/MS with negative ion mode ESI. Resulting samples were analyzed with the ACQUITY

UPLC (Waters; MA, USA) and a Q-Exactive high resolution/accurate mass spectrometer

(ThermoScientific; MA, USA) interfaced with a heated electrospray ionization (HESI-II)

source and Orbitrap mass analyzer operated at 35,000 mass resolution. An acidic positive

ion condition was used on an aliquot optimized to detect more hydrophilic compounds.

Another acidic positive ion condition was ran but chromotographically optimized for

hydrophobic compounds. Basic negative ion optimized conditions was also used on a

separate C18 column. Resulting raw data was extracted and peaks identified using

Metabolon's hardware and software. Compounds were identified by comnparing to known

library entries of purified standards or recurrent unknown entities. A lirary of authenticated

standards containing retention time/index (RI), mass to charge ratio (m/z) and

chromotographic data (MS/MS spectral data) on all library compounds. Three criteria are

used to identify chemicals: 1) RI within a narrow window of proposed identification, 2)

accurate mass match to the library ± 10 ppm and 3) MS/MS forward and reverse scores

between the experimental data and authentic standards. Peaks were quantified using

area under the curve (AUC). For complex lipid panel lipids were extracted with

methanol:dichloromethane in the presence of internal standards. Extracts were

concentrated under nitrogen and reconstituted in 250ml of 10mM ammonium acetate

dichloromethane:methanol (50:50). Mass spec analysis was performed in a Shimazdu LC

with nano PEEK tubing and the Sciex SlexIon-5500 QTRAP (Sciex; MA, USA). Both

negative and positive mode electrospray was used. Individual lipid species were

quantified by taking the peak area ratios of target compounds and their assigned internal

standards then multiplying by the concentration of added internal standards. Lipid class

concentrations were calculated from the sum of all molecular species within a class, and

fatty acid compositions were determined by calculating the proportion of each class

comprised by individual fatty acids.

Oil Red O Staining

Livers were frozen in optimal cutting temperature (OCT) media (FisherScientific; NH,

USA) and sectioned with a cryostat (Leica; Wetzler, Germany) at 4mm. Liver sections

were fixed at room temperature for 15 minutes with 4% paraformaldehyde followed by

washing three times with phosphate buffered saline (PBS). Liver sections were then

incubated with Oil Red O as previously described [74,75]. Liver sections were then washed

three times with PBS and cover slipped with Fluoroshield Mounting Media (Sigma).

Stained slides were visualized with a brightfield microscope (Nikon; NY, USA).

Indirect Calorimetry

To measure whole body energy expenditure in live animals, mice were housed

individually in metabolic chambers of PhenoMaster (Indirect calorimetry system; TSE

systems, Germany) and acclimatized for 3 days with free access to food and water. The

Energy expenditure (Oxygen consumption (VO2), Respiratory exchange ratio (RER) and

food intake were obtained continuously during a 12 h light and 12 h dark cycle for four

days and measured for last 24 h after adaptation.

Acknowledgements

The authors of this manuscript would like to thank the National Institute of Environmental

Health Sciences Epigenomics and DNA Sequencing Core for their help with RNA-

sequencing and Dr. Sara Grimm for her analysis of transcriptome data. This work was

supported by NIH Intramural Program 1ZIAES10328601 (J.M.), NIH R01 HL132035

(X.Z.), NIH K01 DK117069 (C.K), NIH K01 AG056663 (E.Q.), NIH R01 AI065791

(C.E.M.), R01 GM102497 (C.E.M.), R35 GM126922 (C.E.M.).

Citations

1. Buchman, T. G. *et al.* Sepsis Among Medicare Beneficiaries: 1. The Burdens of Sepsis, 2012–2018. *Critical Care Medicine* **48**, 276 (2020).

2. Rhee, C. *et al.* Incidence and trends of sepsis in US hospitals using clinical vs claims data, 2009-2014. *Jama* **318**, 1241-1249 (2017).

3. Mira, J. C. *et al.* Sepsis Pathophysiology, Chronic Critical Illness, and Persistent Inflammation-Immunosuppression and Catabolism Syndrome. *Crit Care Med* **45**, 253-262, doi:10.1097/CCM.0000000000002074 (2017).

4. Vachharajani, V. & McCall, C. E. Epigenetic and metabolic programming of innate immunity in sepsis. *Innate Immun* **25**, 267-279, doi:10.1177/1753425919842320 (2019).

5. Netea, M. G. *et al.* Defining trained immunity and its role in health and disease. *Nat Rev Immunol* **20**, 375-388, doi:10.1038/s41577-020-0285-6 (2020).

6. Lewis, A. J., Billiar, T. R. & Rosengart, M. R. Biology and metabolism of sepsis: innate immunity, bioenergetics, and autophagy. *Surgical infections* **17**, 286-293 (2016).

7. Bermudes, A. C. G. *et al.* Changes in lipid metabolism in pediatric patients with severe sepsis and septic shock. *Nutrition* **47**, 104-109 (2018).

8. Cheng, S.-C. *et al.* Broad defects in the energy metabolism of leukocytes underlie immunoparalysis in sepsis. *Nature immunology* **17**, 406 (2016).

9. Singer, M. The role of mitochondrial dysfunction in sepsis-induced multi-organ failure. *Virulence* **5**, 66-72 (2014).

10. Druml, W., Heinzel, G. & Kleinberger, G. Amino acid kinetics in patients with sepsis. *The American journal of clinical nutrition* **73**, 908-913 (2001).

11. Mecatti, G. C., Messias, M. C. F. & de Oliveira Carvalho, P. Lipidomic profile and candidate biomarkers in septic patients. *Lipids in Health and Disease* **19**, 1-9 (2020).

12. Quinones, Q. J., Ma, Q., Zhang, Z., Barnes, B. M. & Podgoreanu, M. V. Organ protective mechanisms common to extremes of physiology: a window through hibernation biology. *American Zoologist* **54**, 497-515 (2014).

13. Singer, M. *et al.* The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* **315**, 801-810, doi:10.1001/jama.2016.0287 (2016).

14. Stacpoole, P. W. The pyruvate dehydrogenase complex as a therapeutic target for age-related diseases. *Aging cell* **11**, 371-377 (2012).

15. McCall, C. E. *et al.* Pyruvate dehydrogenase complex stimulation promotes immunometabolic homeostasis and sepsis survival. *JCI insight* **3** (2018).

16. Zhu, X. *et al.* Stimulating pyruvate dehydrogenase complex reduces itaconate levels and enhances TCA cycle anabolic bioenergetics in acutely inflamed monocytes. *J Leukoc Biol* **107**, 467-484, doi:10.1002/JLB.3A1119-236R (2020).

17. Dahn, M. S., Mitchell, R. A., Lange, M. P., Smith, S. & Jacobs, L. A. Hepatic metabolic response to injury and sepsis. *Surgery* **117**, 520-530 (1995).

18. Liu, P. S. *et al.* alpha-ketoglutarate orchestrates macrophage activation through metabolic and epigenetic reprogramming. *Nat Immunol* **18**, 985-994, doi:10.1038/ni.3796 (2017).

19. Mills, E. & O'Neill, L. A. Succinate: a metabolic signal in inflammation. *Trends Cell Biol* **24**, 313-320, doi:10.1016/j.tcb.2013.11.008 (2014).

20. Mills, E. L. *et al.* Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to Drive Inflammatory Macrophages. *Cell* **167**, 457-470 e413, doi:10.1016/j.cell.2016.08.064 (2016).

21   Lampropoulou, V. *et al.* Itaconate Links Inhibition of Succinate Dehydrogenase with Macrophage
Metabolic Remodeling and Regulation of Inflammation. *Cell Metab* **24**, 158-166,
doi:10.1016/j.cmet.2016.06.004 (2016).
22   Liao, S. T. *et al.* 4-Octyl itaconate inhibits aerobic glycolysis by targeting GAPDH to exert anti-
inflammatory effects. *Nat Commun* **10**, 5091, doi:10.1038/s41467-019-13078-5 (2019).
23   Mills, E. L. *et al.* Itaconate is an anti-inflammatory metabolite that activates Nrf2 via alkylation of
KEAP1. *Nature* **556**, 113-117, doi:10.1038/nature25986 (2018).
24   Daniels, B. P. *et al.* The Nucleotide Sensor ZBP1 and Kinase RIPK3 Induce the Enzyme IRG1 to
Promote an Antiviral Metabolic State in Neurons. *Immunity* **50**, 64-76 e64,
doi:10.1016/j.immuni.2018.11.017 (2019).
25   Indo, H. P. *et al.* Evidence of ROS generation by mitochondria in cells with impaired electron
transport chain and mitochondrial DNA damage. *Mitochondrion* **7**, 106-118,
doi:10.1016/j.mito.2006.11.026 (2007).
26   Simoes, I. C. M., Fontes, A., Pinton, P., Zischka, H. & Wieckowski, M. R. Mitochondria in non-
alcoholic fatty liver disease. *Int J Biochem Cell Biol* **95**, 93-99, doi:10.1016/j.biocel.2017.12.019
(2018).
27   Brooks, H., Osabutey, C., Moss, R., Andrews, P. & Davies, D. Caecal ligation and puncture in the
rat mimics the pathophysiological changes in human sepsis and causes multi-organ dysfunction.
Metabolic brain disease **22**, 353-373 (2007).
28   Okin, D. & Medzhitov, R. Evolution of inflammatory diseases. *Curr Biol* **22**, R733-740,
doi:10.1016/j.cub.2012.07.029 (2012).
29   Weis, S. *et al.* Metabolic Adaptation Establishes Disease Tolerance to Sepsis. *Cell* **169**, 1263-1275
e1214, doi:10.1016/j.cell.2017.05.031 (2017).
30   Quinn, M. A. & Cidlowski, J. A. Endogenous hepatic glucocorticoid receptor signaling coordinates
sex-biased inflammatory gene expression. *FASEB J*, doi:10.1096/fj.15-278309 (2015).
31   Dąbek, A., Wojtala, M., Pirola, L. & Balcerczyk, A. Modulation of Cellular Biochemistry,
Epigenetics and Metabolomics by Ketone Bodies. Implications of the Ketogenic Diet in the
Physiology of the Organism and Pathological States. *Nutrients* **12**, 788 (2020).
32   DiMarco, N. M., Beitz, D. C. & Whitehurst, G. B. Effect of fasting on free fatty acid, glycerol and
cholesterol concentrations in blood plasma and lipoprotein lipase activity in adipose tissue of
cattle. *J Anim Sci* **52**, 75-82, doi:10.2527/jas1981.52175x (1981).
33   McCall, C. E. *et al.* Pyruvate dehydrogenase complex stimulation promotes immunometabolic
homeostasis and sepsis survival. *JCI Insight* **3**, doi:10.1172/jci.insight.99292 (2018).
34   Breuille, D. *et al.* A sustained rat model for studying the long-lasting catabolic state of sepsis.
Infect Immun **67**, 1079-1085 (1999).
35   Laudanski, K. *et al.* The clinical and immunological performance of 28 days survival model of
cecal ligation and puncture in humanized mice. *PLoS One* **12**, e0180377,
doi:10.1371/journal.pone.0180377 (2017).
36   Pugh, A. M., Auteri, N. J., Goetzman, H. S., Caldwell, C. C. & Nomellini, V. A Murine Model of
Persistent Inflammation, Immune Suppression, and Catabolism Syndrome. *Int J Mol Sci* **18**,
doi:10.3390/ijms18081741 (2017).
37   Buck, M. D., Sowell, R. T., Kaech, S. M. & Pearce, E. L. Metabolic Instruction of Immunity. *Cell*
169, 570-586, doi:10.1016/j.cell.2017.04.004 (2017).
38   Williams, N. C. & O'Neill, L. A. A role for the Krebs cycle intermediate citrate in metabolic
reprogramming in innate immunity and inflammation. *Frontiers in immunology* **9**, 141 (2018).
39   Zaslona, Z. & O'Neill, L. A. J. Cytokine-like Roles for Metabolites in Immunity. *Mol Cell*,
doi:10.1016/j.molcel.2020.04.002 (2020).

40   McGuire, V. A. *et al.* Dimethyl fumarate blocks pro-inflammatory cytokine production via
inhibition of TLR induced M1 and K63 ubiquitin chain formation. *Scientific reports* **6**, 31159
(2016).
41   Della Giustina, A. *et al.* Dimethyl fumarate modulates oxidative stress and inflammation in
organs after Sepsis in rats. *Inflammation* **41**, 315-327 (2018).
42   Qin, W. *et al.* S-glycosylation-based cysteine profiling reveals regulation of glycolysis by
itaconate. *Nat Chem Biol* **15**, 983-991, doi:10.1038/s41589-019-0323-5 (2019).
43   Bambouskova, M. *et al.* Electrophilic properties of itaconate and derivatives regulate the IκBζ–
ATF3 inflammatory axis. *Nature* **556**, 501-504 (2018).
44   Hooftman, A. & O'Neill, L. A. The immunomodulatory potential of the metabolite itaconate.
Trends in immunology (2019).
45   Yi, Z. *et al.* IRG1/Itaconate Activates Nrf2 in Hepatocytes to Protect Against Liver Ischemia-
Reperfusion Injury. *Hepatology*, doi:10.1002/hep.31147 (2020).
46   Giustina, A. D. *et al.* Vitamin B6 reduces oxidative stress in lungs and liver in experimental sepsis.
An Acad Bras Cienc **91**, e20190434, doi:10.1590/0001-3765201920190434 (2019).
47   Chen, J. *et al.* Protective effects of melatonin on sepsis-induced liver injury and dysregulation of
gluconeogenesis in rats through activating SIRT1/STAT3 pathway. *Biomed Pharmacother* **117**,
109150, doi:10.1016/j.biopha.2019.109150 (2019).
48   Langley, R. J. *et al.* An integrated clinico-metabolomic model improves prediction of death in
sepsis. *Sci Transl Med* **5**, 195ra195, doi:10.1126/scitranslmed.3005893 (2013).
49   Liu, T. F., Vachharajani, V. T., Yoza, B. K. & McCall, C. E. NAD+-dependent sirtuin 1 and 6 proteins
coordinate a switch from glucose to fatty acid oxidation during the acute inflammatory
response. *Journal of biological chemistry* **287**, 25758-25769 (2012).
50   Galván-Peña, S. *et al.* Malonylation of GAPDH is an inflammatory signal in macrophages. *Nature
communications* **10**, 1-11 (2019).
51   Galvan-Pena, S. *et al.* Malonylation of GAPDH is an inflammatory signal in macrophages. *Nat
Commun* **10**, 338, doi:10.1038/s41467-018-08187-6 (2019).
52   Koskinas, J. *et al.* Liver histology in ICU patients dying from sepsis: a clinico-pathological study.
World journal of gastroenterology: WJG **14**, 1389 (2008).
53   Balasse, E. O. & Féry, F. Ketone body production and disposal: effects of fasting, diabetes, and
exercise. *Diabetes/metabolism reviews* **5**, 247-270 (1989).
54   Newman, J. C. & Verdin, E. Ketone bodies as signaling metabolites. *Trends in Endocrinology &
Metabolism* **25**, 42-52 (2014).
55   OWEN, O. E. & Schramm, V. L.   (Portland Press Ltd., 1981).
56   Barcellos, L., Marqueze, A., Trapp, M., Quevedo, R. & Ferreira, D. The effects of fasting on
cortisol, blood glucose and liver and muscle glycogen in adult jundiá Rhamdia quelen.
Aquaculture **300**, 231-236 (2010).
57   Nakamura, K. *et al.* Effect of food-restriction stress on immune response in mice. *Journal of
neuroimmunology* **30**, 23-29 (1990).
58   Xu, C. *et al.* Direct effect of glucocorticoids on lipolysis in adipocytes. *Molecular endocrinology*
23, 1161-1170 (2009).
59   Zhu, X. *et al.* Stimulating pyruvate dehydrogenase complex reduces itaconate levels and
enhances TCA cycle anabolic bioenergetics in acutely inflamed monocytes. *Journal of Leukocyte
Biology* (2020).
60   Min, B. K. *et al.* Pyruvate Dehydrogenase Kinase Is a Metabolic Checkpoint for Polarization of
Macrophages to the M1 Phenotype. *Front Immunol* **10**, 944, doi:10.3389/fimmu.2019.00944
(2019).

61 Huang, B., Wu, P., Bowker-Kinley, M. M. & Harris, R. A. Regulation of pyruvate dehydrogenase
kinase expression by peroxisome proliferator-activated receptor-alpha ligands, glucocorticoids,
and insulin. *Diabetes* **51**, 276-283, doi:10.2337/diabetes.51.2.276 (2002).
62 Hutson, N. J. & Randle, P. J. Enhanced activity of pyruvate dehydrogenase kinase in rat heart
mitochondria in alloxan-diabetes or starvation. *FEBS Lett* **92**, 73-76, doi:10.1016/0014-
5793(78)80724-8 (1978).
63 Baxter, M. A. & Coore, H. G. The mode of regulation of pyruvate dehydrogenase of lactating rat
mammary gland. Effects of starvation and insulin. *Biochem J* **174**, 553-561,
doi:10.1042/bj1740553 (1978).
64 Fuller, S. J. & Randle, P. J. Reversible phosphorylation of pyruvate dehydrogenase in rat skeletal-
muscle mitochondria. Effects of starvation and diabetes. *Biochem J* **219**, 635-646,
doi:10.1042/bj2190635 (1984).
65 Denyer, G. S., Kerbey, A. L. & Randle, P. J. Kinase activator protein mediates longer-term effects
of starvation on activity of pyruvate dehydrogenase kinase in rat liver mitochondria. *Biochem J*
239, 347-354, doi:10.1042/bj2390347 (1986).
66 Wu, P. *et al.* Starvation increases the amount of pyruvate dehydrogenase kinase in several
mammalian tissues. *Arch Biochem Biophys* **381**, 1-7, doi:10.1006/abbi.2000.1946 (2000).
67 Klyuyeva, A., Tuganova, A., Kedishvili, N. & Popov, K. M. Tissue-specific kinase expression and
activity regulate flux through the pyruvate dehydrogenase complex. *J Biol Chem* **294**, 838-851,
doi:10.1074/jbc.RA118.006433 (2019).
68 Bao, H., Kasten, S. A., Yan, X. & Roche, T. E. Pyruvate dehydrogenase kinase isoform 2 activity
limited and further inhibited by slowing down the rate of dissociation of ADP. *Biochemistry* **43**,
13432-13441, doi:10.1021/bi049488x (2004).
69 Pratt, M. L. & Roche, T. E. Mechanism of pyruvate inhibition of kidney pyruvate dehydrogenasea
kinase and synergistic inhibition by pyruvate and ADP. *J Biol Chem* **254**, 7191-7196 (1979).
70 Vachharajani, V. T. *et al.* SIRT1 inhibition during the hypoinflammatory phenotype of sepsis
enhances immunity and improves outcome. *J Leukoc Biol* **96**, 785-796,
doi:10.1189/jlb.3MA0114-034RR (2014).
71 Chen, Z., Fitzgerald, R. L., Averna, M. R. & Schonfeld, G. A targeted apolipoprotein B-38.9-
producing mutation causes fatty livers in mice due to the reduced ability of apolipoprotein B-
38.9 to transport triglycerides. *J Biol Chem* **275**, 32807-32815, doi:10.1074/jbc.M004913200
(2000).
72 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
73 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
74 Quinn, M. A., Xu, X., Ronfani, M. & Cidlowski, J. A. Estrogen Deficiency Promotes Hepatic
Steatosis via a Glucocorticoid Receptor-Dependent Mechanism in Mice. *Cell Rep* **22**, 2690-2701,
doi:10.1016/j.celrep.2018.02.041 (2018).
75 Mehlem, A., Hagberg, C. E., Muhl, L., Eriksson, U. & Falkevall, A. Imaging of neutral lipids by oil
red O for analyzing the metabolic status in health and disease. *Nat Protoc* **8**, 1149-1154,
doi:10.1038/nprot.2013.055 (2013).

Figure Legends

Figure 1: Sepsis Impairs hepatic mitochondrial metabolism. (A) Top 5 canonical pathways subject to transcriptional alterations in the liver identified by Ingenuity Pathway Analysis of RNA-seq of sham versus CLP mice (n = 4 mice per group). Blue represents a negative z-score and orange represents a positive z-score. Shading indicates intensity of pathway activation/inhibition. (B) GSEA of the oxidative phosphorylation pathway (GO:0006119) indicating a negative normalized enrichment score (NES = -0.853). (C) GSEA of the TCA cycle pathway (GO:0006099) indicating a negative enrichment score (NES = -1.1956). (D) Schematic representation of hepatic TCA cycle metabolites altered during chronic sepsis. Red denotes a metabolite increased in response to sepsis; green indicates a metabolite decreased in response to sepsis; black indicates a metabolite un-changed in response to sepsis; grey indicates a metabolite not measured in our metabo-lomic screening. (E-J) Relative metabolite levels measured by Ultrahigh Performance Liq-uid Chromatography-Tandem Mass Spectroscopy (UPLC-MS/MS) from livers of sham and CLP mice 30 hours post-surgery (n = 5 sham; 8 CLP). (K) rlog transformed counts from RNA-seq of sham and CLP mice 30 hours post-surgery. (L) Relative itaconate levels in livers of sham and CLP mice measured by UPLC-MS/MS (n = 5 sham; 8 CLP). * de-notes $p<0.05$.

Figure 2: Impaired hepatic redox balance in septic mice. (A) GSEA of the ROS met-abolic pathway (GO:0072593) from RNA-seq from livers comparing sham to CLP showing a positive enrichment score (NES = 1.117) (n = 4 mice per group). (B-F) Relative metab-olite levels of metabolites involved in redox balance measured by UPLC-MS/MS (n = 5

sham; 8 CLP). (G) % of glutathione pool that is either oxidized or reduced in livers from

sham and CLP mice 30 hours post-surgery. * denotes $p<0.05$, **** $p<0.0001$.

Figure 3: Sepsis promotes hepatic steatosis. (A) Top 5 induced and repressed physi-

ological pathways in the liver of sham versus CLP mice identified by IPA of RNA-seq. (B)

GSEA of the fatty acid metabolic pathway (GO:0019395) from RNA-seq from livers com-

paring sham to CLP showing a negative enrichment score (NES = -1.018) (n = 4 mice per

group). (C) Heatmap representation of average log2 fold change of genes involved in fatty

acid oxidation (GO:0019395) and fatty acid biosynthetic process (GO:0006633) in CLP

versus sham operated mice (n = 5 sham; n = 8 CLP). (D) IPA of top 5 metabolic pathways

significantly altered in the liver in response to sepsis identified by global metabolomic

screening. (E) Heatmap representation of log2 fold change of different lipid species in

sham and CLP mice measured by UPLC-MS/MS. (F) Targeted lipidomic screening by

UPLC-MS/MS of livers from sham and CLP mice 30 hours post-surgery (n = 5 sham; n =

8 CLP). (G) Oil red O staining of frozen liver sections from sham and CLP mice 30 hours

post-surgery. Scale bar is 50μm. Image is representative of 3 independent mice per

group.

Figure 4: Sepsis-induced anorexia increases plasma free fatty acid levels. (A) Cu-

mulative food intake over 24-hour period following sham or CLP surgery. (B) Relative

abundance of (B) glucose, (C) corticosterone, (D) 3-hydroxybutyrate, (E) 2-hydroxybutyr-

ate and (F) 2-aminobutyrate measured by UPLC-MS/MS in plasma from sham and CLP

mice 30 hours post-surgery (n = 6 sham; n = 10 CLP). (G) Heatmap representation of fold

change of different lipid species in plasma from sham and CLP mice 30 hours post-sur-

gery (n = 6 sham; n = 10 CLP). * denotes p<0.05, ** p<0.01, p<0.001, **** p<0.0001.

Figure 5: PDK inhibition restores hepatic metabolism in septic mice. (A) Venn dia-

gram of differentially expressed genes (DEGs) assessed by RNA-seq in CLP versus

sham compared to CLP+DCA versus sham 30 hours post-surgery (left). Heatmap of av-

erage log2 fold change of DEGs in sham, CLP and CLP+DCA (right) (n = 4 mice per

group). (B) Heatmap depiction of z-scores of top canonical pathways identified by IPA in

CLP versus sham and CLP+DCA versus sham. (C) Heatmap depiction of average log2

fold change in metabolite levels involved in TCA cycle in sham, CLP and CLP+DCA 30

hours post-surgery measured by UPLC-MS/MS (n = 5 sham; n = 8 CLP and CLP+DCA).

(D) Heatmap depiction of average log2 fold change in metabolite levels involved in redox

balance in sham, CLP and CLP+DCA 24 hours post-surgery measured by UPLC-MS/MS

(n = 5 sham; n = 8 CLP and CLP+DCA). (E) GSSG/GSH ratio in sham, CLP and

CLP+DCA measured by UPLC-MS/MS. (F) Average fold change in monoacylglycerol,

diacylglycerol and triacylglycerol in CLP and CLP+DCA relative to sham measured by

UPLC-MS/MS. (G) Oil red O staining of frozen liver sections from sham and CLP mice 30

hours post-surgery. Scale bar is 50μm * denotes p<0.05, ** p<0.01, *** p<0.001, ****

p<0.0001.

Figure 6: DCA rescues systemic energy crisis during polymicrobial infection. (A) Heatmap of average log2 fold change in plasma metabolite levels in sham, CLP and CLP+DCA (left) assessed by UPLC-MS/MS. Venn diagram of altered plasma metabolites in CLP versus sham compared to CLP+DCA versus sham 30 hours post-surgery (right) (n = 6 sham; n = 10 CLP and CLP+DCA). (C) Pathway analysis of 198 unique plasma metabolites altered in CLP mice. Left: % overlap of metabolites changed in CLP mice with metabolic pathway. Right: -log(p-value) of pathways altered. (D) Change in body weight in sham, CLP and CLP+DCA 30 hours post-surgery. Total food intake after 30 hours post-surgery in sham, CLP and CLP+DCA. (F) Respiratory exchange ratio (RER) in VO2 in sham, CLP and CLP+DCA 24 hours-post surgery. (H) Schematic representation of hepatic and systemic energy consequences of sepsis and PDK inhibition. * denotes $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 1: Sepsis Impairs Hepatic Mitochondrial Metabolism

Figure 2: Impaired Hepatic Redox Balance in Septic Mice

Figure 3: Sepsis promotes hepatic steatosis

Figure 4: : Sepsis-induced anorexia increases plasma free fatty acid levels

Figure 5: PDK inhibition restores hepatic metabolism in septic mice

Figure 6: DCA rescues systemic energy crisis during polymicrobial infection