Dichloroacetate reverses sepsis-induced hepatic metabolic dysfunction

Rabina Mainali[1*], Manal Zabalawi[2*], David Long[2], Nancy Buechler[1], Ellen Quillen[2], Chia-Chi Key[2], Xuewei Zhu[2], John S. Parks[2], Cristina Furdui[2], Peter W. Stacpoole[3], Jennifer Martinez[4], Charles E. McCall[2#], Matthew A, Quinn[1,2,5#]

[1]Department of Pathology, Section on Comparative Medicine Wake Forest School of Medicine, Winston-Salem, North Carolina

[2]Department of Internal Medicine, Section on Molecular Medicine Wake Forest School of Medicine, Winston-Salem, North Carolina

[3]Division of Endocrinology, Diabetes and Metabolism, Department of Medicine and Department of Biochemistry and Molecular Biology, University of Florida College of Medicine, Gainesville, Florida

[4]Immunity, Inflammation, and Disease Laboratory, National Institute of Environmental Health Sciences (NIEHS), National Institutes of Health (NIH), Research Triangle Park, NC

[5]Lead Contact

* these authors contributed equally
# denotes co-corresponding author

Corresponding Authors:

Matthew A. Quinn, Ph.D.
Wake Forest School of Medicine
Medical Center Blvd
Winston-Salem, NC 27157
Office: 336-713-1995
mquinn@wakehealth.edu

Charles E. McCall, M.D.
Wake Forest School of Medicine
Medical Center Blvd
Winston-Salem, NC 27157
Office: 336 413 8034
chmccall@wakehealth.edu

Keywords: sepsis, mitochondria function, liver, inflammation, metabolism, steatosis

Abstract

Dramatic metabolic reprogramming between an anabolic resistance and catabolic toler­ance state occurs within the immune system in response to systemic infection with the sepsis syndrome. While metabolic tissues such as the liver are subject to end-organ dam­age during sepsis and are the primary cause of sepsis death, how their metabolic and energy reprogramming during sepsis state ensures survival is unclear. Employing com­prehensive metabolomic screening, targeted lipidomic screening, and transcriptional pro­filing in a mouse model of septic shock, we show that hepatocyte lipid metabolism, mito­chondrial TCA energetics, and redox balance are significantly reprogramed after cecal ligation and puncture (CLP). We identify increases in TCA cycle metabolites citrate, cis-aconitate, and itaconate with reduced fumarate and triglyceride accumulation in septic hepatocytes. Transcription analysis of liver tissue supports and extends the hepatocyte findings. Strikingly, the administration of the pyruvate dehydrogenase kinase (PDK) in­hibitor dichloroacetate (DCA) reverses dysregulated hepatocyte metabolism and mito­chondrial dysfunction. Our data indicate sepsis promotes hepatic metabolic dysfunction. Furthermore, our data indicate that targeting the mitochondrial PDC/PDK energy home­ostat rebalances transcriptional and metabolic manifestations of sepsis within the liver.

Sepsis is a potentially life-threatening condition that occurs due to the body's overwhelming response to an infection. This reaction leads to aberrant immune responses and if not diagnosed and treated early after its onset, may limit survival by inducing coagulopathy, altered microvasculature, and dysregulation of the host's metabolism and organ function [1-7]. Sepsis accounts for one in three hospital deaths in the U.S. and millions of deaths each year globally, highlighting its hazard to public health [8,9]. The high mortality rate associated with sepsis reflects the lack of a clinically viable molecular-based therapeutic target. Therefore, understanding the pathogenesis of sepsis at both the molecular and organismal level is of utmost importance to address major gaps in knowledge regarding sepsis.

The response to sepsis has classically been characterized as a biphasic phenomenon, where the acute phase is characterized by high energy consumption and hyper-inflammation associated with oxidative stress, followed by cellular reprogramming to a low energy, anti-inflammatory state of immunometabolic paralysis, with accompanying organ failure [8,10-12]. Additionally, during the early phase of sepsis increased catabolism of fats, proteins, and carbohydrates, associated with high rates of oxygen consumption and ATP synthesis is observed [13-16]. Following the acute phase is a hypometabolic state where ATP production and mitochondrial respiration decreases [13]. It is postulated that this is a protective mechanism to overall lower the metabolic demands of the cell and help with its recovery [1,17,18].

Of particular interest to us, during the hyper-inflammatory anabolic phase of sepsis, an increase in the expression and activity of pyruvate dehydrogenase kinase 1 (PDK1) consistently occurs [19,20]. This enzyme is one of four PDK isoforms that reversibly phosphorylates serine residues on pyruvate dehydrogenase complex (PDC) E1a subunit, inhibiting the conversion of pyruvate to acetyl coenzyme A (acetyl CoA) [21]. Inhibition of this important enzymatic activity that connects glycolysis to tricarboxylic acid (TCA) cycle, oxidative phosphorylation (OXPHOS), and the lipogenic pathway is thought to be an important mechanism driving the dysfunction of mitochondrial respiration and cell bioenergetics observed during sepsis [19,22]. Therefore, PDK serves as a potential therapeutic target as it can allow for the downstream oxidation of glucose to continue, restoring

OXPHOS and mitochondrial functioning that we know to be altered in both immune and
non-immune cells during sepsis [16,19,23,24].

The liver is an important metabolic and immune organ due to its role in nutrient
metabolism and production of acute phase proteins. However, our understanding of tran-
scriptional alterations and subsequent metabolic manifestations elicited by sepsis still re-
main limited [16,25-28]. Furthermore, the effects of DCA at restoring hepatic metabolic func-
tion in the context of sepsis is completely unknown. Hence, we set out to characterize the
hepatic manifestations of sepsis, with the overall goal of identifying global metabolic path-
ways subject to dysregulation and whether these pathways are restored by DCA treat-
ment.

Results

Sepsis Impairs Hepatic Mitochondrial Metabolism

To test the long-term hepatic transcriptional changes elicited by sepsis, we performed RNA-seq in whole livers thirty-hours post cecal ligation puncture (CLP). At this time point, septic mice exhibit tolerance, in which immunometaboic paralysis and end organ dysfunctions decrease survival. Ingenuity pathway analysis (IPA) of the top physiological pathways subject to alteration in response to sepsis revealed a significant increase in the acute phase response pathway, highlighting a persistent inflammatory state in the liver into the chronic phase of sepsis (Fig. 1a). Of particular interest were the findings that oxidative phosphorylation and mitochondrial dysfunction were the top two enriched pathways in the liver of septic mice (Fig. 1a). To gain further insight into the effects of sepsis on the transcriptional regulation of mitochondrial function, we performed gene set enrichment analysis (GSEA) of the oxidative phosphorylation pathway (GO:0006119). Septic mice at 30 h decreased the transcriptional output of OXPHOS components, as evidenced by a negative GSEA enrichment score (Fig. 1b). Given the connection between oxidative phosphorylation and the tricarboxylic acid (TCA) cycle, we next asked if polymicrobial infection would elicit similar alterations in this pathway. In accordance with the changes observed in the oxidative phosphorylation pathway, we found a negative enrichment score for TCA cycle enzymes (GO:0006099) in the liver of CLP mice (Fig. 1c). Thus, our transcriptome data indicate that sepsis impairs mitochondrial metabolism in the liver.

Next, we wanted to assess if the transcriptional changes elicited by sepsis would manifest in altered hepatic TCA cycle metabolism. Therefore, we performed global

unbiased metabolomic screening in isolated hepatocytes from control and septic mice by

Ultrahigh Performance Liquid Chromotography-Tandem Mass Spectroscopy (UPLC-

MS/MS). In line with altered transcriptional regulation of the TCA cycle, we found that

sepsis altered the relative abundance of multiple metabolites involved in the TCA cycle

(Fig. 1d). In particular, significant elevation of citrate and cis-aconitate was observed in

septic hepatocytes at 30h (Fig. 1e&f). Unlike macrophages, which shift their ratio of

succinate and α-ketoglutarate to favor succinate accumulation over α-ketoglutarate [29-31],

we found no changes in the levels of these metabolites in hepatocytes (Fig. 1G&H).

Furthermore, unlike macrophages [32], levels of fumarate decreased, but malate levels

were unchanged in hepatocytes from septic animals (Fig. 1I&J). Monocytes,

macrophages and dendritic cells reprogram the TCA cycle during acute inflammation to

a catabolic tolerance phenotype by shunting cis-aconitate to itaconate through the

enzymatic action of aconitase decarboxylase (ACOD1; also known as immune

responsive gene 1 [IRG1]) [32-35]. Accordingly, we also found that sepsis significantly

induces *ACOD1* and elevates itaconate in isolate hepatocyte preparations (Fig. 1K&L).

Sepsis Impairs Hepatic Redox Balance and Promotes Oxidative Stress

Perturbations in oxidative phosphorylation have been shown to promote the

production of reactive oxygen species (ROS) [36]. Furthermore, itaconate has been shown

to induce genes involved in oxidative stress regulation through activation of the master

NRF2 KEAP1 antioxidative pathway in macrophages [34]. Therefore, we examined if sepsis

invokes a gene expression profile involved in regulation of ROS. GSEA of the ROS

metabolic process pathway (GO:0072593) revealed a positive enrichment, highlighting

the transcriptional induction of genes involved in the regulation of oxidative stress in livers

of septic mice (Fig. 2a). Given the induction of ROS metabolic genes, we wanted to

determine if sepsis alters key metabolites involved in redox balance, specifically the

cysteine-glutathione transsulfuration redox regulatory cycle. Sepsis did not alter

intracellular homocysteine levels in isolated hepatocytes (Supplemental Fig. 1), however,

cystathionine decreased significantly (Fig. 2b). Furthermore, we found a trend for

decreased cysteine levels in the liver (p=0.0534) (Fig. 2c) and significantly decreased

glycine (Fig. 2d). Most dramatically, septic mice depleted hepatocyte glutathione (Fig. 2e)

and led to decreases in oxidized glutathione (GSSG) (Fig. 2f). Ophthalamate has recently

been shown to be a biomarker of oxidative stress and signals consumption of hepatic

GSH [37]. Functionally, we observe a significant accumulation of hepatic ophthalamate

levels in response to sepsis, signifying a state of oxidative injury (Fig. 2g).

Sepsis promotes hepatic steatosis

Thus far, our data has revealed that sepsis induces significant impairments in

hepatocyte OXPHOS, alterations in the TCA cycle, and induction of oxidative stress, find-

ings that are also metabolic hallmarks of fatty liver disease [38]. Because perturbations in

global lipid profiles occur in septic patients [15], we evaluated the effects of CLP on the

hepatic lipid metabolism pathway. In support of our hypothesis that sepsis alters hepatic

lipid metabolism, transcriptome data indicate significant repression of lipid metabolism in

livers of septic mice (Fig. 3a). Sepsis also decreased transcription output that supports

fatty acid metabolic process components in the liver, as evidenced by a negative GSEA

enrichment score (Fig. 3b). In particular, we find alterations in genes involved in both fatty

acid oxidation as well as fatty acid biosynthesis (Supplemental Fig. 2).

Since transcriptional regulation of hepatic lipid metabolism is significantly altered

during sepsis, we further investigated the functional manifestation of these gene expres-

sion profile. Consistent with our transcriptional profiling studies, our unbiased metabo-

lomic screening in hepatocytes also identified lipid metabolism as the top metabolic path-

way altered in sepsis (Fig.3c). We observe increases in virtually all fatty acids and acyl-

carnitine derivatives assayed (Fig. 3d). The dysregulation in hepatic lipid metabolism in

response to sepsis ultimately culminates in the development of steatosis, as evidenced

by increased free fatty acids, mono-, di- and triglycerides (Fig. 3e).

Sepsis remodels the hepatic lipidome

Given the severe dysregulation in hepatic lipid metabolism triggered by sepsis, we

next sought to characterize the global consequences on hepatic lipid profiles. This was

achieved by targeted lipidomic screening. Parallel to our unbiased metabolomic

screening, we find sepsis leads to profound accumulation of almost all lipid species

surveyed (Fig. 4a). We next evaluated the hepatic lipid composition to discern which lipid

species are must vulnerable to sepsis-induced alterations. We find very little alterations

to the ceramide pool in septic hepatocytes (Fig. 4b) with only modest changes in

dihydroceramide (Supplemental Fig 3.). However, the hepatic fatty acid fractions

displayed significant remodelling in response to sepsis (Fig. 4b). Specifically, we find

increases in free fatty acid species such as 16:0, 18:1 and 18:2 in septic hepatocytes

(Supplemental Fig. 4a). Morever, hepatic phospholipid composition showed robust

alterations following CLP (Fig. 4b). Of the phospholipids (PLs) most significantly affected,

we find accumulation of phosphotidylcholines (PCs), phosphotidylethanolamine (PE) and

sphingomyelin (SM) (Fig. 4c). Of the hepatic PC pool we find the most abundant species

(16:0/18:2 & 16:0/18:1) to show increased accumulion in response to sepsis

(Supplemental Fig. 4b). We also observed significant increases in the most abundant PE

species (18:0/20:4, 16:0/22:6 & 18:0/22:6) and SM species (22:0, 16:0 & 20:0) to show

significant accumulation in septic hepatocytes (Supplemental Fig. 4c&d). We did observe

a decrease in one phospholipid species, lysophosphatidylcholine, however, the hepatic

concentration of this phospholipid species is signifcantly lower than other PL species

assayed (Fig. 4c). Finally, we detected no differences in other PL species such as

phosphotidylinositol (PI) and lysophosphatidylethanolamine (LPE) and no differences in

hepatic cholesterol ester levels (Supplemental Fig. 5). Our data collectively detail the

impact of sepsis on hepatic lipid composition and robustly show a global rise in nearly all

lipid species.

PDK inhibition attenuates sepsis-induced transcriptional and metabolic changes

in the liver

We have previously reported that PDK inhibition by the pyruvate analog and pan-

PDK inhibitor dichloroacetate (DCA) promotes PDC-dependent immunometabolic adap-

tations to sepsis and increases survival [39]. Given the establishment of hepatocyte mito-

chondrial dysfunction and steatosis during sepsis, we asked whether pharmacological

targeting of the hepatic PDK/PDC axis would mitigate the disruption of key metabolic and

bioenergetic processes induced by sepsis. To test this postulate, we administered DCA

to septic mice 24 h after CLP onset and assessed the global transcription profile of liver

6 h later via RNA-seq. DCA treatment of septic mice reversed the majority of sepsis-

regulated gene networks (Fig. 5a). Most notably, pathways involved in redox homeosta-

sis, lipid metabolism and mitochondrial dysfunction are returned to control levels with the

administration of DCA in septic mice (Fig. 5b).

Given our previous results indicating the transcriptional changes associated with

sepsis in the liver underlie functional alterations in metabolism, we wanted to determine

if the DCA reversal of transcriptional changes mounts to differential metabolomes. Given

our RNA-seq data showing a positive enrichment score for production of ROS along with

negative enrichment for glutathione conjugation, we started off with evaluating the redox

pathway metabolites. Strikingly, the depletion of redox components cystathionine, cyste-

ine, glycine, hypotaurine, taurine and glutathione elicited by sepsis was completely re-

versed by administration of DCA (Fig. 5c). More importantly, we observed a significant

decrease in hepatic ophthalmate levels, signifying a functional decrease in oxidative

stress (Fig. 5c). The restoration in redox metabolites in septic mice following DCA admin-

istration prompted us to investigate the mitochondrial consequences of this pharmaco-

logical intervention. With the exception of fumarate, DCA completely restores TCA me-

tabolite levels to that of sham levels (Fig. 5d). Finally, we wished to define the effects of

DCA treatment on the hepatic lipidome. Lipidomic analysis shows a protective effect of

DCA in ameliorating sepsis-induced steatosis. In sum, these data reveal pharmacological

inhibition of the PDK pathway restores transcriptional and metabolic changes associated

with sepsis in the liver.

Discussion

Considerable attention has been focused recently on the immunometabolic consequences of inmflammation in immune cells [40]. However, significant gaps exist in our understanding of the metabolic adaptations induced by systeminc inflammation from sepsis in vital organs. Addressing this limitation is critical because sepsis survival depends on restoring both organ and immune cell homeostasis following dysregulated and disseminated inflammation [41]. In the present study, we reveal severe disruption of several key hepatic metabolic pathways in septic mice, including TCA cycle activity, OXPHOS, redox balance and lipid metabolism. Importantly, DCA administration restored TCA cycle flux, restored rerdox balance and ameliorated sepsis-induced steatosis. Our study highlights the clinical potential of targeting the PDC/PDK axis in restoring liver function/metabolism in the context of sepsis.

Remodelling of TCA cycle flux is well appreciated to support immune cell effector function in the face of inflammatory challenges [3,31,42,43]. For example, succinate primes inflammation through succinate dehydrogenase (SDH)-mediated ROS generation and ATP synthesis [31,42]. SDH-derived ROS enhances IL-1β production by activating HIF-1α and activating the NLRP3 inflammasome [20]. Inhibition of SDH activity increases IL-10 production and skews inflamed immune cells towards an anti-inflammatory response [31]. Fumarate, on the other hand, possesses anti-inflammatory properties as it inhibits pro-inflammatory cytokine production either by activating the NRF2 pathway or via inhibiting pathways like NF-κB and MAPK that lie downstream of TLR signaling in immune cells [44,45]. Our findings extend the concept that the TCA cycle is subject to reprogramming in response to sepsis in non-immune cells such as hepatocytes. Furthermore, we found

similarities as well as differences in terms of TCA cycle remodelling in hepatocytes com-

pared to immune cell effects, clearly indicating this pathway is regulated in a cell-type and

context-dependent manner during sepsis.

In response to LPS stimulation, macrophages significantly upregulate IRG1 and

accumulate itaconate from the decarboxylation of cis-aconitate [32,34]. One of the primary

functions itaconate serves in immune cells is to limit IL-1 β production through direct

inhibition of SDH and activation of the NRF2/Keap1 and IκBζ-ATF3 pathways [32,34,46]. In

line with what occurs in macrophages, we observed an increase in IRG1 and itaconate in

septic hepatocytes at 30h. While we did not fully characterize the hepatic function of

itaconate, a recent paper demonstrates its anti-inflammatory effects during Ischemia-

reperfusion (IR) injury in the liver [47]. The deletion of IRG1 heightens inflammation and

liver damage and renders hepatocytes susceptible to oxidative injury after I/R injury.

Further, itaconate administration reduces liver damage and inflammation associated with

I/R in IRG1 KO mice, also emphasizing its anti-inflammatory and hepatoprotective effects

[47]. Given the immunomodulatory effects of itaconate and its accumulation in the liver in

our study, we hypothesize that itaconate directs hepatocyte shifts in both TCA cycle

metabolism and mitochondrial energetics. DCA reduces itaconate levels, which we also

find in a human monocyte model of sepsis [48].

Similar to immune cells, liver cells increase ROS production during sepsis [49,50].

We offer a potential mechanism whereby sepsis leads to significant depletion of redox

metabolites within the glutathione pathway ultimately culminating in loss of cellular

glutathione levels resulting in ophthalamate accumulation in hepatocytes.

Sepsis profoundly alters lipid metabolism, resulting in significant increases in

plasma fatty acid and glycerol concentrations, changes reported to predict prognosis in

septic patients [51]. Furthermore, patients who died from sepsis had evidence of hepatic

steatosis affecting 5%-80% of liver parenchyma [52]. Human data supports a robust hepatic

steatosis phenotype subsequent to sepsis. One vital lipid metabolic pathway which can

contribute to steatosis is *de novo* lipogenesis. While we did not directly measure hepatic

lipogenesis, our metabolomics screening did show an increase in hepatic citrate and free

fatty acid levels suggesting an increased lipogenic drive during sepsis. Functionally,

immune cells exploit the lipogenic pathway for the generation of malonyl-CoA and

subsequent malonylation of glycolytic enzymes, which ultimately contributes to the

metabolic adaptation of macrophages to inflammatory cues [53]. Future studies dissecting

the role of lipogenesis in altering hepatocyte function during systemic infection are

warranted.

In addition to accumulation of free fatty acids and triglycerides in the liver, our

lipidomic analysis revealed significant increases in phospholipids. In the present study,

we   find   hepatic   levels   of   the   phospholipids   phosphatidylcholine,

phosphatidylethanolamine, and sphingomyelin to be heightened during sepsis, indicating

abnormal metabolism of phospholipids. PC and PE are the most abundant phospholipids

that make up cellular membranes and an abnormally high ratio of PC and PE has been

reported to influence energy metabolsm and linked to fatty liver disease and impairement

in liver regeneration after injury [54-56]. Altered PC: PE ratio has also been shown to

influence the dynamics and regulation of lipid droplets contributing towards steatosis, a

phenotype we also report in our study [54,55,57]. The dysregulation of hepatic phospholipid

metabolism may be a contributing mechanism underlying steatosis, which is a topic of

future investigation. Additionally, phospholipids serve as precursor molecules to bioactive

lipids, which are involved in numerous signal transduction cascades [58]. Investigating

whether signaling properties of bioactive lipids in the liver during sepsis is pathogenic or

contributes to sepsis resolution is also needed in the future.

PDC is a master metabolic regulator controlling the conversion of pyruvate to

acetyl-CoA in the mitochondria [59] . A part of the metabolic reprogramming that occurs in

immune cells in response to inflammatory signaling is inactivation of PDC [22,59]. PDK is a

negative regulator of the PDC, as it phosphorylates PDC and inhibits the conversion of

pyruvate to acetyl-CoA [59]. During sepsis, the expression and activity of PDK1 in immune

cells is heightened, contributing to the dysfunction of mitochondria metabolism. One

mechanism driving this observed increase is through pro-inflammatory mediators such as

LPS and interferon gamma (IFN-γ) [60]. Another possible mechanism contributing to PDK

activation in the context of sepsis is through glucocorticoid signaling [61]. In fact, starvation

is well established to activate the PDK pathway [62-66]. Based on our findings of elevated

glucocorticoid levels in both the circulation and the liver, we hypothesize the stress

hormone pathway as a contributor to PDK activation during sepsis.

We reported in a sepsis monocyte model that DCA reduced TCA cycle catabolic

effects concurrent with increasing amino acid anaplerotic catabolism of branched-chain

amino acids, leading to increased TCA-driven anabolic energetics [22]. In the present

study, we find restoration of hepatic TCA metabolites, decreased triglyceride

accumulation, lessening of lipid synthesis and oxidative stress rebalances in septic mice

after DCA administration. Overall, we demonstrate that hepatic transcriptional and

metabolic dysfunction improves after targeting the PDC/PDK axis with DCA.

Further work elucidating the mechanistic pathways involved in the dysfunction of

lipid metabolism and the impact of TCA metabolite alterations is warranted. In particular,

further investigation specifically into the PDK pathway to determine which isoform

underlies the hepatic manifestions of sepsis. This is warranted given multiple isoforms of

PDK expressed in the liver [67] and DCA inhibits multiple isoforms of PDK [68,69]. The results

of this study fill a gap in understanding how sepsis at the molecular level compromises

liver function. It also further informs the novel therapeutic targeting potential of the

PDC/PDK homeostat.

376
377

Material and Methods

Animal Experiments

Male C57BL/6J mice aged 8-10 weeks were purchased from The Jackson Laboratory

(Bar Harbor, ME). All animals were subject to a 12:12 hour dark/light cycle with *ad libitum*

access to standard rodent chow and water. Cecal ligation puncture model was used to

induce sepsis as previously described [39,70]. Briefly, cecum was ligated and punctured two

times with a 22-gauge needle. Contents were then returned and incision was closed in

two layers (peritoneum and skin). Sham operation where abdominal incision was made

but cecum not ligated or punctured was used as a control. Subcutaneous fluids (1ml

normal saline) were given to each animal. Mice were euthanized 30 hours post-surgery

for tissue collection. Dichloroacetate (DCA) (Sigma; MO, USA) was administered

(25mg/kg) intraperitoneally at various time points post-surgery: for metabolomic

screening and RNA-seq DCA was administered 24 hours post-surgery and tissues

collected 6 hours post DCA administration (30 hours post-surgery), for metabolic cages

DCA was administered immediately post-surgery.

Hepatocyte Isolation

Hepatocytes were isolated via portal vein perfusion and collagenase digestion as

previously described [71]. Following perfusion, liver cells were liberated by gentle

dissocation in DMEM (ThermoFisher; CA, USA). Cells were then filtered through nylon

mesh to remove cellular debris and connective tissue and resulting cells pelleted by

centrifugation at 50g for 1 minute. After three washes with DMEM cells were counted and

viability assessed via Trypan Blue exclusion.

RNA-sequencing

RNA was isolated from whole liver tissues using Trizol and the RNeasy RNA isolation kit (Qiagen; MD, USA) according to manufacturer's protocol. One microgram of high quality RNA (RIN>8) was used as a template for library generation using the Illumina TruSeq RNA Sample Prep Kit v2 (Illumina; CA, USA) according to the manufacturer's protocol. Generated libraries were then poly(A) enriched for mRNA prior to sequencing. Indexed samples were sequenced at 100bp-paired-end protocol with the NovaSeq 6000 (Illumina), generating approximately 20-30 million reads per sample. Sequenced reads were aligned to the University of California Santa Cruz (UCSC) mm10 reference genome using STAR v2.5 as previously described [72]. The mapped read counts were quantified by Subread feature Counts v1.5.0-p1 [73]. Differentially expressed genes (DEGs) were determined by DESeq2 v1.14.1 [73] using a false discory rate of 0.05. Ingenuity Pathway Analysis (Qiagen) and Gene Set Enrichment Analysis v4.0.3 (GSEA) were further used as previously described [74].

Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy

Hepatocytes were isolated described above for metabolomic screening via Ultrahigh Performance Liquid Chromotography-Tandem Mass Spectroscopy (UPLC-MS/MS) (Metabolon; NC, USA). Briefly, 150-200 microliter cell pellets per animal were used as starting material. Samples were prepared using the automated MicroLab STAR system (Hamilton; NV, USA). Proteins were precipitated using methanol under vigorous shaking for 2 minutes followed by centrifugation. Prior to analysis organic solvents were removed

with TurboVap (Zymark; MA, USA) and overnight storage under nitrogen. Dried samples

were reconstituted with solvents compatible with the three following analytical methods:

1) reverse phase (RP)/UPLC-MS/MS methods with positive ion mode electrospray

ionization (ESI), 2) RP/UPLC-MS/MS with negative ion mode ESI and 3) HILIC/UPLC-

MS/MS with negative ion mode ESI. Resulting samples were analyzed with the ACQUITY

UPLC (Waters; MA, USA) and a Q-Exactive high resolution/accurate mass spectrometer

(ThermoScientific; MA, USA) interfaced with a heated electrospray ionization (HESI-II)

source and Orbitrap mass analyzer operated at 35,000 mass resolution. An acidic positive

ion condition was used on an aliquot optimized to detect more hydrophilic compounds.

Another acidic positive ion condition was ran but chromotographically optimized for

hydrophobic compounds. Basic negative ion optimized conditions was also used on a

separate C18 column. Resulting raw data was extracted and peaks identified using

Metabolon's hardware and software. Compounds were identified by comnparing to known

library entries of purified standards or recurrent unknown entities. A lirary of authenticated

standards containing retention time/index (RI), mass to charge ratio (m/z) and

chromotographic data (MS/MS spectral data) on all library compounds. Three criteria are

used to identify chemicals: 1) RI within a narrow window of proposed identification, 2)

accurate mass match to the library ± 10 ppm and 3) MS/MS forward and reverse scores

between the experimental data and authentic standards. Peaks were quantified using

area under the curve (AUC). For complex lipid panel lipids were extracted with

methanol:dichloromethane in the presence of internal standards. Extracts were

concentrated under nitrogen and reconstituted in 250ml of 10mM ammonium acetate

dichloromethane:methanol (50:50). Mass spec analysis was performed in a Shimazdu LC

with nano PEEK tubing and the Sciex SlexIon-5500 QTRAP (Sciex; MA, USA). Both

negative and positive mode electrospray was used. Individual lipid species were

quantified by taking the peak area ratios of target compounds and their assigned internal

standards then multiplying by the concentration of added internal standards. Lipid class

concentrations were calculated from the sum of all molecular species within a class, and

fatty acid compositions were determined by calculating the proportion of each class

comprised by individual fatty acids.

470 **Acknowledgements**
471
The authors of this manuscript would like to thank the National Institute of Environmental

Health Sciences Epigenomics and DNA Sequencing Core for their help with RNA-

sequencing and Dr. Sara Grimm for her analysis of transcriptome data. This work was

supported by NIH Intramural Program 1ZIAES10328601 (J.M.), NIH R01 HL132035

(X.Z.), NIH R01 HL119962 (J.S.P.), NIH K01 DK117069 (C.K), NIH K01 AG056663

(E.Q.), NIH R01 AI065791 (C.E.M.), R01 GM102497 (C.E.M.), R35 GM126922 (C.E.M.).

Citations
479

1   Lewis, A. J., Billiar, T. R. & Rosengart, M. R. Biology and metabolism of sepsis: innate immunity,
bioenergetics, and autophagy. *Surgical infections* **17**, 286-293 (2016).
2   Bermudes, A. C. G. *et al.* Changes in lipid metabolism in pediatric patients with severe sepsis and
septic shock. *Nutrition* **47**, 104-109 (2018).
3   Cheng, S.-C. *et al.* Broad defects in the energy metabolism of leukocytes underlie
immunoparalysis in sepsis. *Nature immunology* **17**, 406 (2016).
4   Levi, M. & van der Poll, T. Coagulation and sepsis. *Thrombosis research* **149**, 38-44 (2017).
5   Schneck, E. *et al.* Plasma DNA and RNA differentially impact coagulation during abdominal
sepsis—an explorative study. *Journal of Surgical Research* **210**, 231-243 (2017).
6   McDonald, B. *et al.* Platelets and neutrophil extracellular traps collaborate to promote
intravascular coagulation during sepsis in mice. *Blood, The Journal of the American Society of*
Hematology **129**, 1357-1367 (2017).
7   Kidokoro, A., Iba, T., Fukunaga, M. & Yagi, Y. Alterations in coagulation and fibrinolysis during
sepsis. *Shock (Augusta, Ga.)* **5**, 223-228 (1996).
8   Rhee, C. *et al.* Incidence and trends of sepsis in US hospitals using clinical vs claims data, 2009-
2014. *Jama* **318**, 1241-1249 (2017).
9   Buchman, T. G. *et al.* Sepsis Among Medicare Beneficiaries: 1. The Burdens of Sepsis, 2012–
2018. *Critical Care Medicine* **48**, 276 (2020).
10   Osuchowski, M. F., Craciun, F., Weixelbaumer, K. M., Duffy, E. R. & Remick, D. G. Sepsis
chronically in MARS: systemic cytokine responses are always mixed regardless of the outcome,
magnitude, or phase of sepsis. *The Journal of Immunology* **189**, 4648-4656 (2012).
11   Chiswick, E. L., Mella, J. R., Bernardo, J. & Remick, D. G. Acute-phase deaths from murine
polymicrobial sepsis are characterized by innate immune suppression rather than exhaustion.
The Journal of Immunology **195**, 3793-3802 (2015).
12   Xiao, H., Siddiqui, J. & Remick, D. G. Mechanisms of mortality in early and late sepsis. *Infection*
and immunity **74**, 5227-5235 (2006).
13   Singer, M. The role of mitochondrial dysfunction in sepsis-induced multi-organ failure. *Virulence*
5, 66-72 (2014).
14   Druml, W., Heinzel, G. & Kleinberger, G. Amino acid kinetics in patients with sepsis. *The*
American journal of clinical nutrition **73**, 908-913 (2001).
15   Mecatti, G. C., Messias, M. C. F. & de Oliveira Carvalho, P. Lipidomic profile and candidate
biomarkers in septic patients. *Lipids in Health and Disease* **19**, 1-9 (2020).
16   Seymour, C. W. *et al.* Metabolomics in pneumonia and sepsis: an analysis of the GenIMS cohort
study. *Intensive care medicine* **39**, 1423-1434 (2013).
17   Quinones, Q. J., Ma, Q., Zhang, Z., Barnes, B. M. & Podgoreanu, M. V. Organ protective
mechanisms common to extremes of physiology: a window through hibernation biology.
American Zoologist **54**, 497-515 (2014).
18   Singer, M., De Santis, V., Vitale, D. & Jeffcoate, W. Multiorgan failure is an adaptive, endocrine-
mediated, metabolic response to overwhelming systemic inflammation. *The Lancet* **364**, 545-
548 (2004).
19   McCall, C. E. *et al.* Pyruvate dehydrogenase complex stimulation promotes immunometabolic
homeostasis and sepsis survival. *JCI insight* **3** (2018).
20   Vary, T. C. Increased pyruvate dehydrogenase kinase activity in response to sepsis. *American*
Journal of Physiology-Endocrinology and Metabolism **260**, E669-E674 (1991).
21   Stacpoole, P. W. The pharmacology of dichloroacetate. *Metabolism* **38**, 1124-1144 (1989).
22   Zhu, X. *et al.* Stimulating pyruvate dehydrogenase complex reduces itaconate levels and
enhances TCA cycle anabolic bioenergetics in acutely inflamed monocytes. *Journal of Leukocyte*
Biology (2020).

23   Kelly, B. & O'neill, L. A. Metabolic reprogramming in macrophages and dendritic cells in innate
immunity. *Cell research* **25**, 771-784 (2015).
24   Eyenga, P. *et al.* Time course of liver mitochondrial function and intrinsic changes in oxidative
phosphorylation in a rat model of sepsis. *Intensive care medicine experimental* **6**, 31 (2018).
25   Ilaiwy, A. *et al.* Identification of Metabolic Changes in Ileum, Jejunum, Skeletal Muscle, Liver, and
Lung in a Continuous IV Pseudomonas aeruginosa Model of Sepsis Using Nontargeted
Metabolomics Analysis. *The American journal of pathology* **189**, 1797-1813 (2019).
26   Dahn, M. S., Mitchell, R. A., Lange, M. P., Smith, S. & Jacobs, L. A. Hepatic metabolic response to
injury and sepsis. *Surgery* **117**, 520-530 (1995).
27   Levy, E., Slusser, R. J. & Ruebner, B. Hepatic changes produced by a single dose of endotoxin in
the mouse. Electron microscopy. *The American journal of pathology* **52**, 477 (1968).
28   Han, X., Fink, M. P., Uchiyama, T., Yang, R. & Delude, R. L. Increased iNOS activity is essential for
hepatic epithelial tight junction dysfunction in endotoxemic mice. *American Journal of*
Physiology-Gastrointestinal and Liver Physiology **286**, G126-G136 (2004).
29   Liu, P. S. *et al.* alpha-ketoglutarate orchestrates macrophage activation through metabolic and
epigenetic reprogramming. *Nat Immunol* **18**, 985-994, doi:10.1038/ni.3796 (2017).
30   Mills, E. & O'Neill, L. A. Succinate: a metabolic signal in inflammation. *Trends Cell Biol* **24**, 313-
320, doi:10.1016/j.tcb.2013.11.008 (2014).
31   Mills, E. L. *et al.* Succinate Dehydrogenase Supports Metabolic Repurposing of Mitochondria to
Drive Inflammatory Macrophages. *Cell* **167**, 457-470 e413, doi:10.1016/j.cell.2016.08.064
(2016).
32   Lampropoulou, V. *et al.* Itaconate Links Inhibition of Succinate Dehydrogenase with Macrophage
Metabolic Remodeling and Regulation of Inflammation. *Cell Metab* **24**, 158-166,
doi:10.1016/j.cmet.2016.06.004 (2016).
33   Liao, S. T. *et al.* 4-Octyl itaconate inhibits aerobic glycolysis by targeting GAPDH to exert anti-
inflammatory effects. *Nat Commun* **10**, 5091, doi:10.1038/s41467-019-13078-5 (2019).
34   Mills, E. L. *et al.* Itaconate is an anti-inflammatory metabolite that activates Nrf2 via alkylation of
KEAP1. *Nature* **556**, 113-117, doi:10.1038/nature25986 (2018).
35   Daniels, B. P. *et al.* The Nucleotide Sensor ZBP1 and Kinase RIPK3 Induce the Enzyme IRG1 to
Promote an Antiviral Metabolic State in Neurons. *Immunity* **50**, 64-76 e64,
doi:10.1016/j.immuni.2018.11.017 (2019).
36   Indo, H. P. *et al.* Evidence of ROS generation by mitochondria in cells with impaired electron
transport chain and mitochondrial DNA damage. *Mitochondrion* **7**, 106-118,
doi:10.1016/j.mito.2006.11.026 (2007).
37   Soga, T. *et al.* Differential metabolomics reveals ophthalmic acid as an oxidative stress
biomarker indicating hepatic glutathione consumption. *J Biol Chem* **281**, 16768-16776,
doi:10.1074/jbc.M601876200 (2006).
38   Simoes, I. C. M., Fontes, A., Pinton, P., Zischka, H. & Wieckowski, M. R. Mitochondria in non-
alcoholic fatty liver disease. *Int J Biochem Cell Biol* **95**, 93-99, doi:10.1016/j.biocel.2017.12.019
(2018).
39   McCall, C. E. *et al.* Pyruvate dehydrogenase complex stimulation promotes immunometabolic
homeostasis and sepsis survival. *JCI Insight* **3**, doi:10.1172/jci.insight.99292 (2018).
40   Buck, M. D., Sowell, R. T., Kaech, S. M. & Pearce, E. L. Metabolic Instruction of Immunity. *Cell*
169, 570-586, doi:10.1016/j.cell.2017.04.004 (2017).
41   Singer, M. *et al.* The Third International Consensus Definitions for Sepsis and Septic Shock
(Sepsis-3). *JAMA* **315**, 801-810, doi:10.1001/jama.2016.0287 (2016).
42   Williams, N. C. & O'Neill, L. A. A role for the Krebs cycle intermediate citrate in metabolic
reprogramming in innate immunity and inflammation. *Frontiers in immunology* **9**, 141 (2018).

43 Zaslona, Z. & O'Neill, L. A. J. Cytokine-like Roles for Metabolites in Immunity. *Mol Cell*,
doi:10.1016/j.molcel.2020.04.002 (2020).
44 McGuire, V. A. *et al.* Dimethyl fumarate blocks pro-inflammatory cytokine production via
inhibition of TLR induced M1 and K63 ubiquitin chain formation. *Scientific reports* **6**, 31159
(2016).
45 Della Giustina, A. *et al.* Dimethyl fumarate modulates oxidative stress and inflammation in
organs after Sepsis in rats. *Inflammation* **41**, 315-327 (2018).
46 Bambouskova, M. *et al.* Electrophilic properties of itaconate and derivatives regulate the
IkappaBzeta-ATF3 inflammatory axis. *Nature* **556**, 501-504, doi:10.1038/s41586-018-0052-z
(2018).
47 Yi, Z. *et al.* IRG1/Itaconate Activates Nrf2 in Hepatocytes to Protect Against Liver Ischemia-
Reperfusion Injury. *Hepatology*, doi:10.1002/hep.31147 (2020).
48 Zhu, X. *et al.* Stimulating pyruvate dehydrogenase complex reduces itaconate levels and
enhances TCA cycle anabolic bioenergetics in acutely inflamed monocytes. *J Leukoc Biol* **107**,
467-484, doi:10.1002/JLB.3A1119-236R (2020).
49 Giustina, A. D. *et al.* Vitamin B6 reduces oxidative stress in lungs and liver in experimental sepsis.
An Acad Bras Cienc **91**, e20190434, doi:10.1590/0001-3765201920190434 (2019).
50 Chen, J. *et al.* Protective effects of melatonin on sepsis-induced liver injury and dysregulation of
gluconeogenesis in rats through activating SIRT1/STAT3 pathway. *Biomed Pharmacother* **117**,
109150, doi:10.1016/j.biopha.2019.109150 (2019).
51 Langley, R. J. *et al.* An integrated clinico-metabolomic model improves prediction of death in
sepsis. *Sci Transl Med* **5**, 195ra195, doi:10.1126/scitranslmed.3005893 (2013).
52 Koskinas, J. *et al.* Liver histology in ICU patients dying from sepsis: a clinico-pathological study.
World journal of gastroenterology: WJG **14**, 1389 (2008).
53 Galvan-Pena, S. *et al.* Malonylation of GAPDH is an inflammatory signal in macrophages. *Nat
Commun* **10**, 338, doi:10.1038/s41467-018-08187-6 (2019).
54 van der Veen, J. N. *et al.*   (Elsevier, 2017).
55 Ling, J., Chaba, T., Zhu, L. F., Jacobs, R. L. & Vance, D. E. Hepatic ratio of phosphatidylcholine to
phosphatidylethanolamine predicts survival after partial hepatectomy in mice. *Hepatology* **55**,
1094-1102 (2012).
56 Li, Z. *et al.* The ratio of phosphatidylcholine to phosphatidylethanolamine influences membrane
integrity and steatohepatitis. *Cell metabolism* **3**, 321-331 (2006).
57 Listenberger, L. *et al.* Decreasing phosphatidylcholine on the surface of the lipid droplet
correlates with altered protein binding and steatosis. *Cells* **7**, 230 (2018).
58 Sunshine, H. & Iruela-Arispe, M. L. Membrane lipids and cell signaling. *Current opinion in
lipidology* **28**, 408 (2017).
59 Stacpoole, P. W. The pyruvate dehydrogenase complex as a therapeutic target for age-related
diseases. *Aging cell* **11**, 371-377 (2012).
60 Min, B. K. *et al.* Pyruvate Dehydrogenase Kinase Is a Metabolic Checkpoint for Polarization of
Macrophages to the M1 Phenotype. *Front Immunol* **10**, 944, doi:10.3389/fimmu.2019.00944
(2019).
61 Huang, B., Wu, P., Bowker-Kinley, M. M. & Harris, R. A. Regulation of pyruvate dehydrogenase
kinase expression by peroxisome proliferator-activated receptor-alpha ligands, glucocorticoids,
and insulin. *Diabetes* **51**, 276-283, doi:10.2337/diabetes.51.2.276 (2002).
62 Hutson, N. J. & Randle, P. J. Enhanced activity of pyruvate dehydrogenase kinase in rat heart
mitochondria in alloxan-diabetes or starvation. *FEBS Lett* **92**, 73-76, doi:10.1016/0014-
5793(78)80724-8 (1978).

63   Baxter, M. A. & Coore, H. G. The mode of regulation of pyruvate dehydrogenase of lactating rat
mammary gland. Effects of starvation and insulin. *Biochem J* **174**, 553-561,
doi:10.1042/bj1740553 (1978).
64   Fuller, S. J. & Randle, P. J. Reversible phosphorylation of pyruvate dehydrogenase in rat skeletal-
muscle mitochondria. Effects of starvation and diabetes. *Biochem J* **219**, 635-646,
doi:10.1042/bj2190635 (1984).
65   Denyer, G. S., Kerbey, A. L. & Randle, P. J. Kinase activator protein mediates longer-term effects
of starvation on activity of pyruvate dehydrogenase kinase in rat liver mitochondria. *Biochem J*
239, 347-354, doi:10.1042/bj2390347 (1986).
66   Wu, P. *et al.* Starvation increases the amount of pyruvate dehydrogenase kinase in several
mammalian tissues. *Arch Biochem Biophys* **381**, 1-7, doi:10.1006/abbi.2000.1946 (2000).
67   Klyuyeva, A., Tuganova, A., Kedishvili, N. & Popov, K. M. Tissue-specific kinase expression and
activity regulate flux through the pyruvate dehydrogenase complex. *J Biol Chem* **294**, 838-851,
doi:10.1074/jbc.RA118.006433 (2019).
68   Bao, H., Kasten, S. A., Yan, X. & Roche, T. E. Pyruvate dehydrogenase kinase isoform 2 activity
limited and further inhibited by slowing down the rate of dissociation of ADP. *Biochemistry* **43**,
13432-13441, doi:10.1021/bi049488x (2004).
69   Pratt, M. L. & Roche, T. E. Mechanism of pyruvate inhibition of kidney pyruvate dehydrogenasea
kinase and synergistic inhibition by pyruvate and ADP. *J Biol Chem* **254**, 7191-7196 (1979).
70   Vachharajani, V. T. *et al.* SIRT1 inhibition during the hypoinflammatory phenotype of sepsis
enhances immunity and improves outcome. *J Leukoc Biol* **96**, 785-796,
doi:10.1189/jlb.3MA0114-034RR (2014).
71   Chen, Z., Fitzgerald, R. L., Averna, M. R. & Schonfeld, G. A targeted apolipoprotein B-38.9-
producing mutation causes fatty livers in mice due to the reduced ability of apolipoprotein B-
38.9 to transport triglycerides. *J Biol Chem* **275**, 32807-32815, doi:10.1074/jbc.M004913200
(2000).
72   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
73   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
74   Quinn, M. A., Xu, X., Ronfani, M. & Cidlowski, J. A. Estrogen Deficiency Promotes Hepatic
Steatosis via a Glucocorticoid Receptor-Dependent Mechanism in Mice. *Cell Rep* **22**, 2690-2701,
doi:10.1016/j.celrep.2018.02.041 (2018).

661   **Figure Legends**

Figure 1: Sepsis Impairs hepatic mitochondrial metabolism. (A) Top 5 canonical pathways subject to transcriptional alterations in the liver identified by Ingenuity Pathway Analysis of RNA-seq of sham versus CLP mice (n = 4 mice per group). Blue represents a negative z-score and orange represents a positive z-score. Shading indicates intensity of pathway activation/inhibition. (B) GSEA of the oxidative phosphorylation pathway (GO:0006119) indicating a negative normalized enrichment score (NES = -0.853). (C) GSEA of the TCA cycle pathway (GO:0006099) indicating a negative enrichment score (NES = -1.1956). (D) Schematic representation of hepatic TCA cycle metabolites altered during chronic sepsis. Red denotes a metabolite increased in response to sepsis; green indicates a metabolite decreased in response to sepsis; black indicates a metabolite unchanged in response to sepsis; grey indicates a metabolite not measured in our metabolomic screening. (E-J) Relative metabolite levels measured by Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy (UPLC-MS/MS) from livers of sham and CLP mice 30 hours post-surgery (n = 5 sham; 8 CLP). (K) rlog transformed counts from RNA-seq of sham and CLP mice 30 hours post-surgery. (L) Relative itaconate levels in livers of sham and CLP mice measured by UPLC-MS/MS (n = 5 sham; 8 CLP). * denotes $p < 0.05$.

Figure 2: Impaired hepatic redox balance in septic mice. (A) GSEA of the ROS metabolic pathway (GO:0072593) from RNA-seq from livers comparing sham to CLP showing a positive enrichment score (NES = 1.117) (n = 4 mice per group). (B-F) Relative metabolite levels of metabolites involved in redox balance measured by UPLC-MS/MS (n = 5

sham; 8 CLP). (G) Relative levels of ophthalmate involved in redox balance measured by

UPLC-MS/MS. * denotes $p<0.05$, ** $p<0.01$, **** $p<0.0001$.

Figure 3: Sepsis promotes hepatic steatosis. (A) Top 5 induced and repressed physi-

ological pathways in the liver of sham versus CLP mice identified by IPA of RNA-seq. (B)

GSEA of the fatty acid metabolic pathway (GO:0019395) from RNA-seq from livers com-

paring sham to CLP showing a negative enrichment score (NES = -1.018) (n = 4 mice per

group). (C) IPA of top 5 metabolic pathways significantly altered in the liver in response

to sepsis identified by global metabolomic screening. (D) Heatmap representation of log2

fold change of different lipid species in sham and CLP mice measured by UPLC-MS/MS.

(E) Quantification of lipids by UPLC-MS/MS in hepatocytes isolated from sham and CLP

mice 30 hours post-surgery (n = 5 sham; n = 8 CLP). * denotes $p<0.05$, ** $p<0.01$, ***

$p<0.001$.

Figure 4: Sepsis reprograms the hepatic lipidome. (A) Volcano plot of significantly

altered lipids from sham and CLP hepatocytes measured by UPLC-MS/MS (n = 5 sham;

n = 8 CLP). (B) Pie chart of lipid compositions from sham and CLP mice. (C) Quantifica-

tion of lipids by UPLC-MS/MS in hepatocytes isolated from sham and CLP mice 30 hours

post-surgery (n = 5 sham; n = 8 CLP). ** denotes $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure 5: PDK inhibition restores hepatic metabolism in septic mice. (A) Venn dia-

gram of differentially expressed genes (DEGs) assessed by RNA-seq in CLP versus

sham compared to CLP+DCA versus sham 30 hours post-surgery (left). Heatmap of av-

erage log2 fold change of DEGs in sham, CLP and CLP+DCA (right) (n = 4 mice per

group). (B) Heatmap depiction of z-scores of top canonical pathways identified by IPA in

CLP versus sham and CLP+DCA versus sham. (C) Heatmap depiction of average log2

fold change in metabolite levels involved in redox balance in sham, CLP and CLP+DCA

24 hours post-surgery measured by UPLC-MS/MS (n = 5 sham; n = 8 CLP and

CLP+DCA). (D) Heatmap depiction of average log2 fold change in metabolite levels in-

volved in TCA cycle in sham, CLP and CLP+DCA 30 hours post-surgery measured by

UPLC-MS/MS (n = 5 sham; n = 8 CLP and CLP+DCA). (E) Heatmap depiction of average

log2 fold change in MAG, DAG and TAG levels in sham, CLP and CLP+DCA 30 hours

post-surgery measured by UPLC-MS/MS (n = 5 sham; n = 8 CLP and CLP+DCA). (F)

Heatmap depiction of average log2 fold change in phospholipids in sham, CLP and

CLP+DCA 30 hours post-surgery measured by UPLC-MS/MS (n = 5 sham; n = 8 CLP

and CLP+DCA.

Figure 1: Sepsis Impairs Hepatic Mitochondrial Metabolism

Figure 2: Impaired Hepatic Redox Balance in Septic Mice

Figure 3: Sepsis promotes hepatic steatosis

Figure 4: Sepsis reprograms the hepatic lipidome

Figure 5: DCA reverses transcriptional and metabolic changes elicited by sepsis in the liver

Supplemental figure 1: Hepatic homocysteine levels in sham and CLP mice. Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy measurement of homocysteine from isolated hepatocytes 30 hours post-surgery. n=5 sham, n=8 CLP.

Supplemental figure 2: Fatty acid oxidation and biosynthetic pathways in septic livers. Heatmap representation of average log2 fold change of genes involved in fatty acid oxidation (GO:0019395) and fatty acid biosynthetic process (GO:0006633) in CLP versus sham operated mice (n = 5 sham; n = 8 CLP).

Supplemental figure 3: Hepatic ceramide levels in sham and CLP mice. Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy measurement of **(A)** dihydroceramide, **(B)** ceramide, **(C)** hexosylceramide and **(D)** lactosylceramide from isolated hepatocytes 30 hours post-surgery. n=5 sham, n=8 CLP. * denotes $p<0.05$ determined by Student's T-test.

Supplemental figure 4: Phosphatidylcholine, phosphatidylethanolamine, and sphingomyelin quantifications.
Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy measurement of **(A)** free fatty acid species, **(B)** phosphatidylcholine species, **(C)** phosphatidylethanolamine species and **(D)** sphingomyelin species from isolated hepatocytes 30 hours post-surgery. n=5 sham, n=8 CLP. **** denotes $p<0.0001$ determined by two-way ANOVA followed by multiple comparisons test.

Supplemental figure 5: PI, LPE and CE levels in septic livers. Ultrahigh Performance Liquid Chromatography-Tandem Mass Spectroscopy measurement of **(A)** phosphatidylinositol, **(B)** lysophosphatidylethanolamine species and **(C)** cholesterol ester from isolated hepatocytes 30 hours post-surgery. n=5 sham, n=8 CLP. **** denotes *p*<0.0001 determined by two-way ANOVA followed by multiple comparisons test.