Towards robust and replicable sex differences in the intrinsic brain

function of autism

Dorothea L. Floris*[1,2], José O. A. Filho*[3], Meng-Chuan Lai[4,5,6,7,8], Steve

Giavasis[3], Marianne Oldehinkel[9], Maarten Mennes[1], Tony Charman[10], Julian

Tillmann[10,21], Guillaume Dumas[11], Christine Ecker[12,13], Flavio Dell'Acqua[12,14],

Tobias Banaschewski[15], Carolin Moessnang[16], Simon Baron-Cohen[7], Sarah

Durston[17], Eva Loth[12,14], Declan G. M. Murphy[12,14], Jan K. Buitelaar[1,2,18],

Christian F. Beckmann[1,2,19], Michael P. Milham[3,20], Adriana Di Martino[3]

*equal contribution

[1] Donders Center for Brain, Cognition and Behavior, Radboud University Nijmegen, Nijmegen, The Netherlands

[2] Department for Cognitive Neuroscience, Radboud University Medical Center Nijmegen, Nijmegen, The

Netherlands

[3] Child Mind Institute, New York City, New York, USA

[4] The Margaret and Wallace McCain Centre for Child, Youth & Family Mental Health, Azrieli Adult

Neurodevelopmental Centre, and Campbell Family Mental Health Research Institute, Centre for Addiction and

Mental Health, Toronto, Canada

[5] Department of Psychiatry and Autism Research Unit, The Hospital for Sick Children, Toronto, Canada

[6] Department of Psychiatry, Faculty of Medicine, University of Toronto, Toronto, Canada

[7] Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, United Kingdom

[8] Department of Psychiatry, National Taiwan University Hospital and College of Medicine, Taipei, Taiwan

[9] Turner Institute for Brain and Mental Health, School of Psychological Sciences, Monash University, Victoria,

Australia

[10] Department of Psychology, Institute of Psychiatry, Psychology, and Neuroscience, King's College London,

London, United Kingdom

[11] Human Genetics and Cognitive Functions, Institut Pasteur, UMR3571 CNRS, Université de Paris, Paris

(75015), France

[12] Sackler Institute for Translational Neurodevelopment, Institute of Psychiatry, Psychology, and Neuroscience,

King's College London, London, United Kingdom

[13] Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital

Frankfurt am Main, Goethe University, Frankfurt, Germany

[14] Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, Psychology, and

Neuroscience, King's College London, London, United Kingdom

[15] Child and Adolescent Psychiatry, Department of Psychiatry and Psychotherapy, Central Institute of Mental

Health, University of Heidelberg, Mannheim, Germany

[16] Department of Psychiatry and Psychotherapy, Central Institute of Mental Health, University of Heidelberg,

Mannheim, Germany

[17] Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, the

Netherlands

[18] Karakter Child and Adolescent Psychiatry University Centre, Nijmegen, the Netherlands

[19] Centre for Functional MRI of the Brain, University of Oxford, Oxford, United Kingdom

[20] Nathan Kline Institute for Psychiatric Research, Orangeburg, New York, USA

[21] Department of Applied Psychology: Health, Development, Enhancement, and Intervention, University of

Vienna, Vienna, Austria

Corresponding author:

Adriana Di Martino, MD

Child Mind Institute,

Autism Center

101 E 56 Street

NY, NY 10026

Email: Adriana.Dimartino@childmind.org

Keywords

Autism spectrum disorder, resting-state functional connectivity, sex differences, replication,

robustness, voxel-mirrored homotopic connectivity

Abstract

Background: Marked sex differences in autism prevalence accentuate the need to understand the role of biological sex-related factors in autism. Efforts to unravel sex differences in the brain organization of autism have, however, been challenged by the limited availability of female data. **Methods**: We addressed this gap by using the largest sample of males and females with autism and neurotypical (NT) control individuals, to date (ABIDE; Autism: 362 males, 82 females; NT: 409 males, 166 females; 7-18 years). Discovery analyses examined main effects of diagnosis, sex and their interaction across five resting-state fMRI (R-fMRI) metrics (voxel-level $Z > 3.1$, cluster-level $p < 0.01$ gaussian random field corrected). Secondary analyses assessed the robustness of the results to different pre-processing approaches and their replicability in two independent samples: the EU-AIMS Longitudinal European Autism Project (LEAP) and the Gender Explorations of Neurogenetics and Development to Advance Autism Research (GENDAAR). **Results**: Discovery analyses in ABIDE revealed significant main effects across the intrinsic functional connectivity (iFC) of the posterior cingulate cortex, regional homogeneity and voxel-mirrored homotopic connectivity (VMHC) in several cortical regions, largely converging in the default network midline. Sex-by-diagnosis interactions were confined to the dorsolateral occipital cortex, with reduced VMHC in females with autism. All findings were robust to different pre-processing steps. Replicability in independent samples varied by R-fMRI measures and effects with the targeted sex-by-diagnosis interaction being replicated in the larger of the two replication samples – EU-AIMS LEAP. **Limitations**: Given the lack of *a priori* harmonization among the discovery and replication datasets available to date, sample-related variation remained and may have affected replicability. **Conclusions**: Atypical cross-hemispheric interactions are neurobiologically relevant to autism. They likely result from the combination of sex-dependent and sex-independent factors with a differential effect across cortical functional

103   networks. Systematic assessments of the factors contributing to replicability are needed and

104   necessitate coordinated large-scale data collection across studies.

Background

Autism spectrum disorder (autism) is characterized by a marked male preponderance in prevalence with three times more males being diagnosed than females [1]. This pronounced sex-differential prevalence implies that sex-related biological factors are likely implicated in the neurobiology of autism. However, little is known about the differential underlying neural expressions in males and females with autism. Such knowledge could widen our understanding of potential underlying mechanisms of autism and related neurodevelopmental conditions [2].

This has motivated research into the impact of biological sex on brain organization in autism [2–5]. With the widely accepted view that the neurobiology of autism involves differences in large-scale brain networks [6,7], resting-state functional magnetic resonance imaging (R-fMRI) has proven to be a valuable complementary tool for investigating atypicalities in intrinsic functional connectivity (iFC). While the exact nature of the intrinsic brain organization in autism remains to be established [6], research on the impact of biological sex differences in autism is just beginning to emerge.

Several R-fMRI studies have focused on autism-related sex differences in iFC [2,8–17]. They vary on the extent of the functional networks and intrinsic properties examined. Most of them examined the strength of iFC between one or more regions/networks selected *a priori* [8–11,13,14], or via data-driven analyses [17]. A few others investigated either local or homotopic iFC across the whole brain [2,12,16]. Across these different efforts, the pattern of findings have also been mixed; some studies supported the predictions from the 'extreme male brain theory' [11,12] (i.e., shifts towards maleness in both sexes [18]), others supported

those from the 'gender-incoherence' theory [8,10,13,14,17] (i.e., androgynous patterns in the

sexes [19]), and one other supported the coexistence of both, at least in males [2].

Specifically, both Ypma et al. [11] and Kozhemiako et al. [12,16] observed results consistent

with a shift towards maleness in autism. In contrast, other studies [8,10,13], reported findings

consistent with 'gender-incoherence' where the results largely revealed hyper-connectivity in

females with autism similar to NT males and hypo-connectivity in males with autism similar

to NT females. Such seeming inconsistencies in the pattern of sex-related differences were in

part addressed by Floris et al. [2] who showed that, at least in males with autism, distinct

patterns of atypical sex-differentiation coexist, and vary as a function of the neural networks

involved. However, the intrinsic brain organization in females with autism has remained

largely unclear. Scarce availability of female datasets in most studies may have contributed to

this variability in findings [20,21].

Here, to examine sex-related atypicalities in both males and females in autism relative to NT

controls, we used, as discovery sample, a large autism R-fMRI sample from the Autism Brain

Imaging Data Sharing Exchange (ABIDE) [21,22] of both males and females of autism and

controls. By aggregating neuroimaging datasets from multiple sources, this data sharing

initiative has begun to provide a means to address the challenge of underrepresentation of

female datasets in autism research. Examining both sexes in both autism and controls allows

to directly capture not only sex differences that are common across individuals (i.e.,

regardless of their diagnosis [main effect of sex]), but also those that are specific to autism

and point towards atypical autism-specific sex differential patterns (i.e., sex-by-diagnosis

interaction effects) [4]. Unlike most prior work that focused on specific networks or circuits

selected *a priori,* we investigated autism-specific sex differences in the whole-brain across

multiple R-fMRI metrics in the same study. Specifically, we selected R-fMRI metrics

capturing unique aspects of the intrinsic brain organization [23], during typical development

[24]. Most germane to this study, their selection was motivated by the consistent involvement

in typical sex differences and to be affected by autism. They comprised: 1) posterior

cingulate cortex (PCC)-iFC – e.g., [2,11,22,25–31]; 2) voxel-mirrored homotopic

connectivity (VMHC) [32] – e.g., [12,31–34]; 3) regional homogeneity (ReHo) [35] – e.g.,

[16,31,36,37]; 4) network degree centrality (DC) [38] – e.g., [31,38–40]; and 5) fractional

amplitude of low frequency fluctuations (fALFF) [22,41] – e.g., [22,31,42].

Beside the role of small female samples, prior inconsistencies in autism-related sex

differences in R-fMRI can be due to other factors that can impact reproducibility. For

example, while there are growing concerns on the role of pre-processing strategies [43,44], a

recent study showed that their impact on autism-related mean group-differences is minimal

[45]. Additionally, while several studies have reported some degree of consistency on R-

fMRI findings across either independent, or partially overlapping, samples [40,46–48], results

from others have raised concerns on the replicability of group-mean diagnostic effects

[45,49]. However, none of these studies have explicitly examined robustness and replicability

of sex-by-diagnosis interaction effects, which account for a potentially relevant source of

variability in autism – biological sex. Thus, we conducted secondary analyses to assess the

extent to which the pattern of findings obtained in our discovery analyses were also observed

a) after applying different nuisance pre-processing steps that have been previously validated,

though used inconsistently in the autism literature [45], and *b)* across two independent,

multisite R-fMRI datasets: the EU-AIMS Longitudinal European Autism Project (LEAP)

[50,51] and the Gender Explorations of Neurogenetics and Development to Advance Autism

Research (GENDAAR) dataset [52] – i.e., *robustness* and *replicability*.

Methods

Discovery sample: ABIDE I and II

For discovery analyses, we examined the R-fMRI dataset with the largest number of females

and males in both the autism and the NT groups available to date, selected from the Autism

Brain Imaging Data Exchange (ABIDE) repositories ABIDE I and II [21,22]. For specific

selection criteria, see Additional file 2 and Additional file 4. The final ABIDE I and II dataset

of N=1,019 included N=82 females with autism, N=362 males with autism, N=166

neurotypical females (NT F), and N=409 neurotypical males (NT M), aggregated across 13

sites. Details on the sample selection process and criteria are provided in Supplementary

methods (Additional file 2) and Additional file 4. Briefly, we selected cases between 7 -18

years of age (the ages most represented across ABIDE sites), with MRI data successfully

completing brain image co-registration and transformation to standard space, with FIQ

between 70-148 and with mean framewise displacement (mFD) [53] within three times the

interquartile range (IQR) + the third quartile (Q3) of the sample (i.e., 0.39 mFD). Further

steps included matching for mean age *across* groups and for mFD and IQ *within* diagnostic

groups. This latter step limited the number of exclusions while keeping average group motion

low (mFD<.2mm) and bypassing sampling biases resulting from matching

neurodevelopmental conditions to NT around intrinsic features such as IQ [54,55]. At each

step, any sites with less than three individual datasets per diagnostic/sex groups were

excluded. Demographics and characteristics of this sample are summarized in Additional file

1: Table 1 and Additional file 2.

Discovery analysis pre-processing pipeline

We examined five whole-brain R-fMRI metrics previously reported to reflect typical sex

differences and found to be atypical in autism, including 1) PCC-iFC, 2) VMHC, 3) ReHo, 4)

DC and 5) fALFF (see Additional file 2).

R-fMRI image pre-processing steps included: slice time correction, 24 motion parameters

regression [56], component-based noise reduction (CompCor) [57], removal of linear and

quadratic trends, and band-pass filtering (0.01-0.1 Hz, for all metrics but fALFF). Functional-

to-anatomical co-registration was achieved by Boundary Based Registration (BBR) using

FSL FLIRT [58]. Linear and nonlinear spatial normalization of functional echo planar images

(EPIs) to Montreal Neurological Institute 152 (MNI152) stereotactic space ($2mm^3$ isotropic)

was done using ANTS registration (Advanced Neuroimaging Tools) [59]. Computation of

voxel-mirrored homotopic connectivity (VMHC) followed registration to a symmetric

template. All R-fMRI derivatives were smoothed by a 6mm FWHM Gaussian kernel. To

account for site and collection time variability across each of the data collections in ABIDE I

and II data repositories, site effects were removed using the ComBat function available in

python [60] (https://github.com/brentp/combat.py). This approach has been shown to

effectively account for scanner-related variance in multi-site R-fMRI data [60]. For further

details see Additional file 2.

Group-level analyses

Statistical Z-maps were generated within study-specific functional volume masks including

all voxels in MNI space present across all subjects. Sex differences were explored by fitting a

general linear model (GLM) including diagnosis or sex as the regressors of interest, and age

and mean framewise displacement (mFD) [53] as nuisance covariates. We did not include

FIQ as a covariate as this is not optimal when comparing groups selected from populations

carrying intrinsic IQ differences such as autism and NT [55]. We assessed the main effects of

diagnosis and sex, along with their interaction effects. We applied gaussian random field

theory correction based on strict voxel-level threshold of $Z > 3.1$ as recommended by [61]

and cluster level, $P < 0.01$, given the assessment of five R-fMRI metrics in the same study

(i.e., P 0.05/5 R-MRI metrics=0.01).

Functional relevance of sex differences in autism

Post-hoc analyses were conducted to functionally characterize the sex-by-diagnosis

interaction result(s). First, to explore the cognitive domains implicated in the cluster(s), we

quantified the percentage of its overlap with 12 cognitive ontology maps [62] thresholded at

$P = 1e–5$. We labelled these components based on the top five tasks each component recruits

[2]. Second, we used the Neurosynth Image Decoder (http://neurosynth.org/decode/) [63] to

visualize the terms most strongly associated with the significant cluster. After excluding

anatomical and redundant terms (synonyms, plurals or noun/adjective/adverb equivalents),

we visualized the top 27 terms showing correlations with the cluster map between $r = 0.64$

and $r = 0.10$. Third, to explore potential clinical relevance of the significant cluster, we

explored brain-behavior relationships as a function of sex within the autism group.

Specifically, we ran a GLM examining the interaction between biological sex and available

ADOS calibrated severity total score (CSS) [64], as well as social-affect (SA) and restricted,

repetitive behavior (RRB) subscores (see Additional file 2) with the dependent variable(s)

being the R-fMRI metric(s) extracted from the cluster mask(s) showing a statistically

significant sex-by-diagnosis effect(s).

Robustness and Replicability

Robustness. We assessed whether patterns of results from the discovery analyses were

observable with two other nuisance regression pipelines that include commonly used

preprocessing steps. One pipeline was global signal regression (GSR) [65] which has often

been used in autism studies, and the other was Independent Component Analysis - Automatic

Removal of Motion Artifacts (ICA-AROMA) [66] which is a relatively novel but

increasingly utilized approach [45]. Given the scope of the present study, unlike prior work

focusing on a wide range of individual preprocessing pipelines [45], we selected GSR and

ICA-AROMA as examples of previously validated approaches thought to have impact on

motion and physiological noise [44]. We conducted group-level analyses on the R-fMRI

metrics means extracted from the masks corresponding to the clusters showing significant

effects in discovery analyses and regressed out the effect of inter-individual differences in

age, and mFD across all effects tested; for main diagnostic and sex comparisons, sex or

diagnosis, respectively, were also regressed out.

Replicability. Similarly, we assessed whether the group patterns observed in significant

clusters identified in discovery analyses, were observed in two relatively large-scale,

independent datasets selected from *a)* the EU-AIMS Longitudinal European Autism Project

(LEAP), a large multi-site European initiative aimed at identifying biomarkers in autism

[50,51] and *b)* the Gender Explorations of Neurogenetics and Development to Advance

Autism Research (GENDAAR) dataset collected by the GENDAAR consortium and shared

in the National Database for Autism Research [52]. For details on autism and NT inclusion

and exclusion criteria for these samples, as well as our selection process, see Additional file 2

[51,52]. The resulting EU-AIMS LEAP (N=309) R-fMRI datasets comprised N=133 males

and N=43 females with autism as well as N=85 NT males, and N=48 NT females (see

Additional file 3: Table S1); resulting GENDAAR (N=196) R-fMRI datasets comprised

N=43 males and N=44 females with autism, as well as N=56 NT males and N=53 NT

females (see Additional file 3: Table S2). For a comparison of demographic and clinical

information between ABIDE, EU-AIMS LEAP and GENDAAR, see Additional file 3: Table

S3, S4 and S5. After applying the same ComBat and pre-processing pipeline used in the

ABIDE-based discovery analyses to adjust for inter-site variability within EU-AIMS LEAP

and GENDAAR, we extracted each of the R-fMRI metrics means from the Z-maps in masks

corresponding to the clusters showing statistically significant effects in discovery analyses.

We then computed group-mean comparisons and their interactions of the residuals data

resulting from regressing out the same nuisance variables used in primary analyses (i.e., mFD

and age across all tests, and sex or diagnosis for diagnostic or sex comparisons, respectively).

For these exploratory analyses, discovery findings were determined to be robust and/or

replicable (R+) based on two criteria: *1)* the group mean difference(s) observed were in the

same direction as those identified in the findings from discovery analyses [67] and *2)* their

effects were not negligible as defined by Cohen' *d* [68] (i.e., any effects, small, medium or

large as indexed by Cohen's $d \geq |0.2|$) which is also consistent with prior work [40].

Results

Discovery analyses – ABIDE

Main effect of diagnosis. Analyses revealed clusters showing a significant effect of diagnosis

(voxel-level $Z > 3.1$; cluster-level $P < 0.01$, corrected) for three of the five R-fMRI metrics:

PCC-iFC, VMHC and ReHo. These were mainly evident in anterior and posterior regions of

the default network (DN) across at least two or all three R-fMRI metrics (Figure 1,

Additional file 5, and Additional file 3: Table S6). Autism-related hypo-connectivity was

present for: a) PCC-iFC, VMHC and ReHo within bilateral paracingulate cortex and frontal

pole, b) VMHC and ReHo in the bilateral PCC and precuneus, and c) ReHo in right insula

and central operculum (Figure 1, Additional file 5 and Additional file 3: Table S6). Autism-

related hyper-connectivity was only evident for PCC-iFC with left superior lateral occipital

cortex, temporal occipital fusiform cortex and occipital fusiform gyrus (Additional file 5). To

verify that these findings were not driven by particular acquisition site(s), post-hoc analyses

computed group means for diagnostic subgroups for the R-fMRI metrics extracted at the

cluster-level masks excluding one out of the 13 ABIDE sites at a time. The pattern of results

was essentially unchanged (Additional file 7: Figure S2a)6.

Main effect of sex. Analyses revealed clusters showing statistically significant main sex

differences (voxel-level $Z > 3.1$; cluster-level $P < 0.01$, corrected) for three metrics out of

five: PCC-iFC, VMHC and ReHo. Findings involved lateral and medial portions of the DN

with bilateral PCC and precuneus showing the highest overlap (see Figure 1 and Additional

file 5). Specifically, regardless of diagnosis, relative to females, males showed decreased

PCC-iFC with paracingulate cortex and frontal pole, right middle frontal gyrus, bilateral

superior lateral occipital cortex and bilateral PCC and precuneus. Males also showed

decreased VMHC and ReHo localized in PCC and precuneus. Decreased ReHo was also

evident in the left angular gyrus and lateral occipital cortex in females relative to males

(Figure 1, Additional file 5, and Additional file 3: Table S6). Post-hoc analyses assessing the

consistency of these findings across sites, as described above, revealed a similar pattern of

results (Additional file 7: Figure S2b).

Sex-by-diagnosis interaction. Statistically corrected voxel-wise analyses (voxel-level $Z >$

3.1; cluster-level $P < 0.01$) revealed a cluster of significant sex-by-diagnosis interaction only

for VMHC which was localized in the dorsolateral occipital cortex (Figure 2a and Additional

file 3: Table S6). Post-hoc cluster-level group means showed that NT females had higher

VMHC than the three other groups, whereas autism females had lower VMHC than the three

other groups (Figure 2a). Similar to the main effects, analyses assessing the consistency of

these findings across sites, as described above, showed a similar pattern of results (Additional

file 8).

Functional relevance of autism-related sex differences

Post-hoc analyses to functionally characterize this VMHC sex-by-diagnosis interaction

indicated that the VMHC cluster in superior lateral occipital cortex overlapped with cognitive

maps involved in higher-order visual, oculomotor, cognitive flexibility and language-related

processes (Figure 3a). Further, as shown in Figure 3b, the most common terms were

primarily related to lower-order visual processing and higher-order visual cognition, such as

'visuospatial' and 'spatial attention.' To explore potential clinical relevance of the VMHC

dorsolateral occipital cluster, we explored brain-behavior relationships as a function of sex,

within the autism group using three available ADOS scores (calibrated severity total score,

and non-calibrated social affect and RRB subscores; see Supplemental Material). Although

not surviving a strict Bonferroni correction for multiple testing (i.e., 0.05/3=0.02), an

interaction effect was observed for ADOS social affect scores. It revealed that more severe

social deficits ($F_{(1,311)}$=4.44, p=0.036) were associated with decreased VMHC in females with

autism (r=-0.29), but not in males with autism (r=0.03). Given that ABIDE data were

aggregated and released when calibrated social affect scores [69] were not available to assess

potential differences in language abilities and age, analyses were repeated after including

ADOS module (ADOS Module 2 to 4) as a nuisance covariate: results remained unchanged

($F_{(1,306)}$=5.0, p=0.026) as they did also after removing the few data with the less represented

ADOS module 2 (see Additional file 2: Supplementary methods). There were no significant

findings, with regard to the CSS total score and non-calibrated RRB sub-score (Figure 3c),

even at an exploratory statistical threshold of $P<0.05$.

Robustness

The same pattern of results identified in primary analyses was observed in the results pre-

processed using GSR or ICA-AROMA, across the three R-fMRI metrics in all the clusters

identified in primary analyses across effects (Cohen's d range=|0.2-0.6|; Figure 2b, Figure 4,

Additional file 6, Additional file 3: Table S6).

Replicability

Main effects of diagnosis. Main effects of diagnosis showed higher replicability (i.e., non-

negligible Cohen's d effects showing a similar group differences pattern as observed in

discovery analyses) in GENDAAR than in EU-AIMS LEAP. Specifically, across the three R-

fMRI metrics that were significant in discovery analyses, four of the seven clusters (57%;

Cohen'd range=|0.2-0.4|) in GENDAAR, while only one (14%; Cohens' d=|0.2|) in EU-

AIMS LEAP, were replicated. Nevertheless, this one cluster showing decreased ReHo in

ASD vs. NT in insula and central operculum was replicated across all samples (Figure 4,

Additional file 9: Figure S3a, and Additional file 3: Table S6).

Main effects of sex. Across all three R-fMRI metrics, the main effects of sex observed in

primary analyses was evident in both independent samples, for most clusters in the EU-AIMS

LEAP (90%; 9/10; Cohen' d range=|0.2-0.5|) and GENDAAR (60%; 6/10; Cohen' d

range=|0.2-0.3|); (Figure 4, Additional file 9: Figure S3b and Additional file 3: Table S6).

Notably, the pattern of typical sex differences localized along the default network midline

(i.e., decreased VMHC, ReHo and PCC-iFC) was replicated across both independent samples

(Figure 6b).

Sex-by-diagnosis interaction effect. The pattern of autism-related VMHC sex differences

observed in discovery analyses in the superior lateral occipital cortex was clearly observed in

the EU-AIMS LEAP dataset (Cohen's *d*=0.34); (Figure 2c; Figure 4, Additional file 3: Table

S6). In the GENDAAR dataset, while group means in males with autism, NT males and NT

females showed a similar direction as in the ABIDE discovery findings, females with autism

differed in magnitude and in the direction of group differences, resulting in a negligible effect

with Cohen's *d*=0.09 (Figure 2c).

Discussion

We examined autism-related sex differences for intrinsic functional brain organization across

multiple R-fMRI metrics in a large discovery sample of males and females with autism

relative to age-group matched NT selected from the ABIDE repositories [21,22]. Analyses

revealed significant main effects of sex and diagnosis across intrinsic functional connectivity

(iFC) of the posterior cingulate cortex, regional homogeneity and voxel-mirrored homotopic

connectivity (VMHC) in several cortical regions. Notably, effects converged along the

midline of the default network. In contrast, sex-by-diagnosis interactions were limited to

VMHC in the superior lateral occipital cortex. This result, placed in the context of sex and

diagnostic main effects on interhemispheric homotopic connectivity in cortical regions,

suggests that atypical interhemispheric interactions are pervasive in autism but reflect a

combination of sex-independent (i.e., main effect of diagnosis common across both sexes)

and sex-dependent (i.e., sex-by-diagnosis interaction) effects, each specific to different

functional cortical system. Sex-by-diagnosis interaction was robust to distinct pre-processing

strategies as those observed for main effects. Further, despite the lack of *a priori*

harmonization for data acquisition among the three samples, it was replicable in the larger of

the two independent samples (i.e., EU-AIMS LEAP). These findings together with largely

replicable main effects of sex with variable replicability of main diagnostic effects by sample,

suggest on the one hand that inter-sample replicability of R-fMRI can be feasible in autism

when sources of variability in diagnostic groups are accounted for in samples sized properly

to account for such variability. On the other hand, they highlight the urgent need to obtain

multiple harmonized datasets properly powered to systematically address and understand

sources of heterogeneity, including and beyond the role of biological sex.

Sex-dependent and independent atypical interhemispheric interactions in autism

VMHC reflects inter-hemispheric homotopic relations. Its strength has been suggested to

index coordinated cross-hemispheric processing: *stronger* VMHC index weaker hemispheric

specialization and vice versa [32,70]. Several lines of evidence support the notion that the

neurobiology of autism is related to atypical hemispheric interactions, including homotopic

connectivity and hemispheric lateralization [34,71–80]. VMHC and functional hemispheric

lateralization have also been shown to be sex-differential in NT [32,81,82]. The dorsolateral

occipital association cortex identified in our discovery analyses is known to serve

hemispherically specialized processes, such as visuospatial coordination [83]. Thus, our

findings of NT males' VMHC in dorsolateral lateral occipital cortex being lower than NT

females are consistent with the notion of increased hemispheric lateralization in this cortical

region in NT males relative to NT females. In our data, females with autism instead showed

even lower VMHC than NT males, while males with autism showed slightly higher VMHC

than NT males. This pattern is indicative of 'gender-incoherence' [19] as males and females

with autism display the opposite pattern expected in NT per their biological sex. Findings of

'gender incoherence' have been reported in earlier neuroimaging studies of autism using

different modalities [11–13,84–86]. Among them, several R-fMRI studies explicitly focusing

on detecting diagnosis-by-sex interactions (i.e., the regression model included a sex-by-

diagnosis interaction term) yielded a pattern of results consistent with 'gender incoherence'.

In contrast, other studies [11–13] reported a pattern consistent with the 'extreme male brain

theory' [18] – i.e., a shift towards maleness in both females and males with autism. While the

seemingly diverging conclusions of these two sets of studies may be attributed to

methodological differences, such as the extent of brain networks explored and the statistical

modelling employed, findings from our prior work suggest that both shifts towards either

maleness or femaleness co-occur in the intrinsic brain of males with autism, in a network-

specific manner [2]. However, such prior work did not include female data, thus by not

directly assessing sex-by-diagnosis interactions, unlike the present study, results could not

point to patterns affecting diagnostic differences between the sexes vs. those common to

autism across both sexes [4]. This is relevant for efforts focusing on identifying underlying

mechanisms of autism. Findings resulting from sex-by-diagnosis interactions may shed light

on sex differential mechanisms that go awry in autism and may reflect sex-specific

susceptibility mechanisms. On the other hand, diagnostic atypicalities common for both sexes

may reflect factors central to the emergence of autism, regardless of whether they overlap

with patterns known to be differential between sexes [87]. Interestingly, a recent study based

on a sample selected from GENDAAR [15] revealed that the iFC between the nucleus

accumbens (selected *a priori*) and in a region of the dorsolateral occipital cortex partially

overlapping with that identified by our VMHC analyses, was differentially modulated by the

aggregate number of oxytocin receptor risk alleles in females with autism vs. NT females and

vs. males with autism. Although VMHC was not directly tested in that earlier study [15], its

result in dorsolateral occipital cortex is consistent with our observation of atypical sexual

differentiation of this region of the visual network and suggest the need for future whole-

brain studies of oxytocin effects in autism.

Along with the sex-dependent autism patterns, our analyses found statistically significant

main effects of diagnosis in inter-hemispheric interactions indexed by VMHC. These were

localized along the midline of the DN (paracingulate/frontal cortex consistently and

PCC/precuneus) where main effects of PCC-iFC and ReHo also converged. Our results are

consistent with prior reports of atypical intrinsic organization of the DN in autism

[11,22,25,88–90]. Together they support the role for a common, sex-independent DN

atypicality in autism. This is also supported by a recent autism neurosubtyping study that

identified three latent iFC factors, all sharing DN atypicalities along with their neurosubtype-

specific patterns [91]. Building on this evidence to disentangle the specific roles of each of

the factors affecting autism in sex-independent and sex-dependent ways, a necessary next

step is to engage in novel large-scale data collection efforts including more female data.

Robustness, replicability and sources of variability

The growing awareness of the replication crisis in neuroscience [92–94] motivated our

analyses examining robustness and replicability of findings. While a comprehensive and

systematic reproducibility assessment is beyond the scope of the study, here we focused on

examining whether the findings observed in the discovery analyses were also seen after using

different preprocessing pipelines – *robustness* – as well as in fully independent, albeit

convenience (i.e., not harmonized *a priori* with each other), samples – *replicability*. To this

extent, given the lack of consensus on quantitative metrics of replicability and to avoid

chasing after p values, we opted to consider findings to be robust and/or replicable for any

non-negligible effects per Cohen's *d* [68], including small, but not negligible effects (i.e.,

Cohen's $> d = 0.2$). Furthermore, given their distributed and heterogenous nature [6], it is

plausible that atypicalities in the autism connectome stem from a combination of differently

sized effects, as shown for autism in other biological domains such as genetics [95,96].

With this in mind, across the two preprocessing methods examined here, the patterns of

findings were consistent with those observed in discovery analyses across all R-fMRI metrics

and effects. These robustness results are consistent with a prior study by He and colleagues

[45] reporting that differences in a wider range of pre-processing pipelines have marginal

effects on variation in diagnostic group average comparisons. Our study confirms and builds

on this earlier report by extending findings of robustness to sex group mean differences and

their interactions with diagnosis.

A more nuanced picture emerged from the inter-sample analyses as replicability varied by

sample, across the effects and R-fMRI metrics examined. Specifically, while inter-sample

main effects of sex were moderately to largely replicable across R-fMRI metrics on both

independent samples (~60 to 90% of the clusters in GENDAAR and EU-AIMS-LEAP,

respectively), replicability of diagnostic effects significantly varied by sample (57 to 14%)

across R-FMRI metrics. This is at least in part consistent with findings by King et al. [49]

that, depending on the R-fMRI feature examined, diagnostic group differences varied across

samples. Even in this scenario, King et al. also reported that findings of decreased homotopic

connectivity in autism were relatively more stable than other R-fMRI metrics [49]. This

combined with replicability of our VMHC sex-by-diagnosis interaction findings in the larger

of the two independent samples (EU-AIMS LEAP), suggests that measures of homotopic

connectivity may have a relevant biological relevance for autism. It is also possible that given

its moderate to high test-retest reliability, VMHC is more suitable in efforts assessing

replicability [97,98].

The striking clinical and biological heterogeneity in autism should be considered as a major

contributor to discrepancies in findings of studies focusing on the main effects of diagnostic

group means contrasts/interactions [99–101]. We interpret our replicability findings on

diagnostic effects and, in turn, diagnosis-by-sex interactions against this background. Inter-

sample differences may have contributed to the more variable results of replicability on the

diagnosis main effects. These include IQ, autism symptom level, and age. For example, the

EU-AIMS LEAP sample was on-average older, had lower VIQ and most notably, lower

symptom severity across all subscales of the ADOS and ADI-R than the ABIDE sample. On

the other hand, the GENDAAR sample (which has greater number of replicable diagnostic

mean group patterns) did not differ from ABIDE in these variables, except for mean age.

Furthermore, a fact that is often neglected, is that the NT groups may also present with

considerable sample heterogeneity between studies [99,102]. For instance, our NT controls in

the EU-AIMS LEAP sample had lower VIQ than both ABIDE and GENDAAR NT controls.

This has potentially influenced the low replicability of diagnosis main effects in EU-AIMS

LEAP.

In contrast, sex-by-diagnosis effect on VMHC in the dorsolateral occipital cortex was

replicable in the larger samples, the EU-AIMS LEAP, but not in GENDAAR. Small samples

introduce larger epistemic variability (i.e., greater variation related to known and unknown

confounds) [103]. Increasing the number of subjects/data allows mitigating epistemic

variability and, thus, capturing the underlying variability of interest. Thus, although the rate

of EU-AIMS LEAP replicability for the main effect of diagnosis was limited, accounting for

biological sex, a known key source variability in autism, may have contributed to a replicable

sex-by-diagnosis pattern in this larger sample. In line with sample size concerns, using four

datasets sized between 36 and 44 individuals selected from the ABIDE repository, He et al.

[45] found low similarity rates of diagnostic group-level differences on the strength of iFC

edges in contrast with the largely similar pattern of results across pipelines. Of note, unlike

prior efforts [45,49], we controlled for site effects within each of the samples (i.e., ABIDE,

GENDAAR and EU-AIMS LEAP), using ComBat. These results are consistent with earlier

reports of reproducible imaging biomarkers even when accounting for inter-site differences in

multisite datasets such as the ABIDE I repository [46]. Future large-scale harmonized data

collections are needed to control and assess the impact of inter-sample variability. Taken

together, these findings highlight that sample differences can impact replicability.

Beyond clinical and biological sources of variation, samples may differ in MRI acquisition

methods, as well as in approaches used to mitigate head motion during data collection and its

impact on findings [104]. Adequately controlling for head motion remains a key challenge for

future studies assessing inter-sample replicability. For the present study, we excluded

individuals with high motion, retained relatively large samples with group average low

motion (mean $\pm$ standard deviation range mFD range=0.09-0.16 $\pm$ 0.06-0.10 mm) as well as

included mFD at the second-level analyses as a nuisance covariate. Overall, the extent to

which each sample-related factors affect replicability needs to be systematically examined in

future well-powered studies. Only this type of studies will allow for emerging subtyping

approaches to dissect heterogeneity by brain imaging features using a range of data-driven

methods [105], including normative modelling [106–108].

Inter-sample differences and methodological differences, beyond nuisance regression, may

have contributed to some differences in findings between the present and earlier studies,

conducted with independent or partially overlapping samples [10,22,40]. For example,

Alaerts et al. [10] also examined sex-by-diagnosis interaction in PCC iFC in a dataset

selected from ABIDE I only. Although the pattern of results was consistent with the 'gender

incoherence' model, the resulting clusters were distinct from those identified in the present

study. Along with differences in samples selected from the same data repositories, other

studies differed with ours in the inclusion of sex-by-diagnosis interaction [16], the extent of

the whole-brain voxel-based analyses [14,15], or the statistical threshold utilized [22].

Nevertheless, it is remarkable that even in light of these differences, consistent results have

emerged including the role of the 'gender incoherence' model for females with autism,

atypical inter-hemispheric interactions in autism, and sex-dependent and sex-independent

atypical intrinsic brain function across distinct functional networks.

Limitations

Along with the inter-sample differences resulting from the lack of sufficiently available

harmonized inter-site replication datasets in the field, other limitations of this study should be

addressed in future efforts. One regards the lack of measures differentiating the effects of sex

vs. that of gender so as to disentangle their relative roles (e.g., gender-identity and gender-

expression) in the intrinsic brain properties [109]. Further, in-depth cognitive measures to

directly characterize the role of VMHC findings were not available. Additional behavioral

measures are needed to establish whether our result in VMHC of the dorsolateral occipital

cortex mainly applies to low-level (bottom-up) visual processing differences or higher-level

(top-down) attentional/controlled processes in males and females with autism. As a

neurodevelopmental disorder, autism shows striking inter-individual differences in clinical

and developmental trajectories, as well as outcomes. Thus, age may influence symptom

presentation [110,111] and neurobiology [12,106,112]. Despite the considerable size of the

samples available for this study, it is still difficult to sufficiently cover a broad age range

across both males and females and diagnostic groups across contributing sites and to evaluate

age effects appropriately. Even larger cross-sectional samples are needed to derive

meaningful age-related information that ultimately require confirmation in longitudinal study

designs.

Conclusions

The present work revealed sex differences in the intrinsic brain of autism, particularly in

dorsolateral occipital interhemispheric interactions, which were robust to pre-processing

pipeline decisions and replicable in the larger of the two independent samples. While

differences in nuisance regression pipelines have little influence on the consistency of

findings, sample heterogeneity represents a challenge for replicability of findings. Lateralized

cognitive functions and cross-hemispheric interactions should be further explored in relation

to sex differences in autism while addressing this challenge with future harmonized data

acquisition efforts with even larger samples.

List of abbreviations

ABIDE: Autism Brain Imaging Data Exchange

ASD: Autism spectrum disorder

DC: Degree centrality

DN: Default network

EPI: Echo-planar image

LEAP: Longitudinal European Autism Project

fALFF: Fractional amplitude of low frequency fluctuations

FIQ: Full-scale IQ

GENDAAR: Gender Explorations of Neurogenetics and Development to Advance Autism

Research

GSR: Global Signal Regression

ICA-AROMA: Independent Component Analysis - Automatic Removal of Motion Artifacts

iFC: Intrinsic functional connectivity

mFD: Mean framewise displacement

NDAR: National Database for Autism Research

NT: Neurotypical

PCC: Posterior cingulate cortex

ReHo: Regional Homogeneity

R-fMRI: Resting-state functional magnetic resonance imaging

VMHC: Voxel-mirrored homotopic connectivity

Ethics approval and consent to participate

ABIDE I and II: All contributions were based on studies approved by the local Institutional

Review Boards, and data were fully anonymized (removing all 18 HIPAA (Health Insurance

Portability and Accountability)-protected health information identifiers, and face information

from structural images).

EU-AIMS LEAP: Ethical approval was obtained through ethics committees at each site.

GENDAAR: Informed assent and consent were obtained from all participants and their legal

guardians, and the experimental protocol was approved by the Institutional Review Board at

each participating site.

Consent for publication

Not applicable

Availability of data and materials

ABIDE I and II: Data are freely accessible in the publicly available Autism Brain Imaging

Data Exchange repository (http://fcon_1000.projects.nitrc.org/indi/abide).

EU-AIMS LEAP: Data are currently only available for sites involved in data collection, but

data (starting with the first wave) will be made available on request.

GENDAAR: Anonymized data are publicly available through the National Database for

Autism Research (NDAR) and access can be requested via https://nda.nih.gov/about.html.

Competing interests

ADM receives royalties from the publication of the Italian version of the Social

Responsiveness Scale—Child Version by Organization Speciali, Italy. JKB has been a

consultant to, advisory board member of, and a speaker for Takeda/Shire, Medice, Roche,

and Servier. He is not an employee of any of these companies and not a stock shareholder of

any of these companies. He has no other financial or material support, including expert

testimony, patents, or royalties. CFB is director and shareholder in SBGneuro Ltd. TC has

received consultancy from Roche and Servier and received book royalties from Guildford

Press and Sage. DM has been a consultant to, and advisory board member, for Roche and

Servier. He is not an employee of any of these companies, and not a stock shareholder of any

of these companies. TB served in an advisory or consultancy role for Lundbeck, Medice,

Neurim Pharmaceuticals, Oberberg GmbH, Shire, and Infectopharm. He received conference

support or speaker's fee by Lilly, Medice, and Shire. He received royalties from Hogrefe,

Kohlhammer, CIP Medien, Oxford University Press; the present work is unrelated to these

652   relationships. JT is a consultant to Roche. The remaining authors declare no competing

653   interests.

Funding

Work for this study has been partly supported by a Postdoctoral Training Award from the

Autism Science Foundation (to DLF/ADM); by NIMH (R21MH107045, R01MH105506,

R01MH115363 to ADM); by gifts to the Child Mind Institute from Phyllis Green, Randolph

Cowen, and Joseph Healey, and by UO1 MH099059 (to MPM); by the Ontario Brain

Institute via the Province of Ontario Neurodevelopmental Disorders Network (IDS-I l-02),

the Slifka-Ritvo Award for Innovation in Autism Research from the International Society for

Autism Research and the Alan B. Slifka Foundation, the Academic Scholars Award from the

Department of Psychiatry, University of Toronto, the O'Brien Scholars Program in the Child

and Youth Mental Health Collaborative at the Centre for Addiction and Mental Health

(CAMH) and The Hospital for Sick Children, the Slaight Family Child and Youth Mental

Health Innovation Fund from CAMH Foundation, and the Canadian Institutes of Health

Research Sex and Gender Science Chair (GSB 171373) (to M-CL). We also acknowledge the

contributions of all members of the EU-AIMS LEAP group. EU-AIMS LEAP has received

funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement

No 115300 (for EU-AIMS) and No 777394 (for AIMS-2-TRIALS). This joint undertaking

receives support from the European Union's Horizon 2020 research and innovation program

and EFPIA and AUTISM SPEAKS, Autistica, SFARI. DM is also supported by the NIHR

Maudsley Biomedical Research Centre. SBC was supported by the Autism Research Trust

during the period of this work.

Authors' contributions (in alphabetical order by contribution)

ADM and DLF have contributed to the study's conception; ADM, DLF, JOAF, M-CL, MPM

have contributed to distinct aspects of the study design and interpretation of all findings; DLF

and JOAF conducted the analyses; SG has provided support in all data analyses requiring

CPAC; ADM, DLF, JOAF have generated figures and tables; ADM and DLF drafted the

manuscript; ADM, DLF, JOAF, M-CL, and MPM have revised and edited multiple versions

of the manuscript; CFB, MM, MO, have organized the EU-AIMS LEAP data and edited

latest manuscript versions and its revisions; CFB, CE, CM, DGMM, EL, FDA, GD, JKB, JT,

SB-C, TC, TB, SD have contributed to the coordination, data acquisition and coordination of

the EU-AIMS LEAP project, as well as edited later versions of the manuscript. All authors

read and approved the manuscript.

Acknowledgements

The authors thank all investigators and contributors to the Gender Explorations of

Neurogenetics and Development to Advance Autism Research (GENDAAR) for collecting

and sharing their data as well as addressing question related to the data, the Autism Brain

Imaging Data Exchange, and the contributors to EU-AIMS Longitudinal European Autism

Project for their efforts in data collection and sharing. The GENDAAR Consortium

comprises, in alphabetical order, Elizabeth H. Aylward, Raphael A. Bernier, Susan Y.

Bookheimer, Mirella Dapretto, Nadine Gaab, Daniel H. Geschwind, Andrei Irimia, Allison

Jack, Charles A. Nelson, Kevin A. Pelphrey, Matthew W. State, John D. Van Horn, Pamela

Ventola, and Sara J. Webb. We thank all participants and their families for participating in

the respective study.

References

1. Loomes R, Hull L, Mandy WPL. What Is the Male-to-Female Ratio in Autism Spectrum

Disorder? A Systematic Review and Meta-Analysis. J Am Acad Child Adolesc Psychiatry

[Internet]. 2017 [cited 2018 Sep 12];56:466–74. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/28545751

2. Floris DL, Lai MC, Nath T, Milham MP, Di Martino A. Network-specific sex

differentiation of intrinsic brain function in males with autism. Mol Autism. 2018;

3. Lai M-C, Lombardo M V, Suckling J, Ruigrok AN V, Chakrabarti B, Ecker C, et al.

Biological sex affects the neurobiology of autism. Brain. 2013;136:2799–815.

4. Lai M-C, Lerch JP, Floris DL, Ruigrok ANV, Pohl A, Lombardo M V., et al. Imaging

sex/gender and autism in the brain: Etiological implications. J Neurosci Res [Internet].

Wiley-Blackwell; 2017 [cited 2018 Sep 12];95:380–97. Available from:

http://doi.wiley.com/10.1002/jnr.23948

5. Ecker C. Notice of Retraction and Replacement: Ecker et al. Association between the

probability of autism spectrum disorder and normative sex-related phenotypic diversity in

brain structure. JAMA Psychiatry. 2017;74(4):329-338. JAMA Psychiatry. 2019.

6. Picci G, Gotts SJ, Scherf KS. A theoretical rut: revisiting and critically evaluating the

generalized under/over-connectivity hypothesis of autism. Dev Sci. 2016;

7. Geschwind DH, Levitt P. Autism spectrum disorders: developmental disconnection

syndromes. Curr. Opin. Neurobiol. 2007.

8. Smith REW, Avery JA, Wallace GL, Kenworthy L, Gotts SJ, Martin A. Sex Differences in

Resting-State Functional Connectivity of the Cerebellum in Autism Spectrum Disorder. Front

Hum Neurosci. 2019;13.

9. Cummings KK, Lawrence KE, Hernandez LM, Wood ET, Bookheimer SY, Dapretto M, et

al. Sex Differences in Salience Network Connectivity and its Relationship to Sensory

Over￼Responsivity in Youth with Autism Spectrum Disorder. Autism Res. 2020;

10. Alaerts K, Swinnen SP, Wenderoth N. Sex differences in autism: A resting-state fMRI

investigation of functional brain connectivity in males and females. Soc Cogn Affect

Neurosci. 2016;

11. Ypma RJF, Moseley RL, Holt RJ, Rughooputh N, Floris DL, Chura LR, et al. Default

Mode Hypoconnectivity Underlies a Sex-Related Autism Spectrum. Biol Psychiatry Cogn

Neurosci Neuroimaging. 2016;

12. Kozhemiako N, Vakorin V, Nunes AS, Iarocci G, Ribary U, Doesburg SM. Extreme male

developmental trajectories of homotopic brain connectivity in autism. Hum Brain Mapp.

2019;

13. Lee JK, Amaral DG, Solomon M, Rogers SJ, Ozonoff S, Nordahl CW. Sex Differences in

the Amygdala Resting-State Connectome of Children With Autism Spectrum Disorder. Biol

Psychiatry Cogn Neurosci Neuroimaging. 2020;5.

14. Lawrence KE, Hernandez LM, Bowman HC, Padgaonkar NT, Fuster E, Jack A, et al. Sex

Differences in Functional Connectivity of the Salience, Default Mode, and Central Executive

Networks in Youth with ASD. Cereb Cortex. 2020;

15. Hernandez LM, Lawrence KE, Padgaonkar NT, Inada M, Hoekstra JN, Lowe JK, et al.

Imaging-genetics of sex differences in ASD: distinct effects of OXTR variants on brain

connectivity. Transl Psychiatry. 2020;10.

16. Kozhemiako N, Nunes AS, Vakorin V, Iarocci G, Ribary U, Doesburg SM. Alterations in

Local Connectivity and Their Developmental Trajectories in Autism Spectrum Disorder:

Does Being Female Matter? Cereb Cortex. 2020;

17. Olson LA, Mash LE, Linke A, Fong CH, Müller RA, Fishman I. Sex-related patterns of

intrinsic functional connectivity in children and adolescents with autism spectrum disorders.

Autism. 2020;

18. Baron-Cohen S. The extreme male brain theory of autism. Trends Cogn Sci. 2002;

19. Bejerot S, Eriksson JM, Bonde S, Carlström K, Humble MB, Eriksson E. The extreme

male brain revisited: Gender coherence in adults with autism spectrum disorder. Br J

Psychiatry. 2012;

20. Watkins E. ScholarWorks at WMU The Gender of Participants in Published Research

Involving People with Autism Spectrum Disorders [Internet]. 2014. Available from:

http://scholarworks.wmich.edu/masters_theseshttp://scholarworks.wmich.edu/masters_theses/

482

21. Di Martino A, O'Connor D, Chen B, Alaerts K, Anderson JS, Assaf M, et al. Enhancing

studies of the connectome in autism using the autism brain imaging data exchange II. Sci

Data. 2017;

22. Di Martino A, Yan C-G, Li Q, Denio E, Castellanos FX, Alaerts K, et al. The autism

brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain

architecture in autism. Mol Psychiatry [Internet]. Nature Publishing Group; 2014 [cited 2018

Sep 12];19:659–67. Available from: http://www.nature.com/articles/mp201378

23. Yan CG, Yang Z, Colcombe SJ, Zuo XN, Milham MP. Concordance among indices of

intrinsic brain function: Insights from inter-individual variation and temporal dynamics. Sci

Bull. Elsevier B.V.; 2017;62:1572–84.

24. Di Martino A, Fair DA, Kelly C, Satterthwaite TD, Castellanos FX, Thomason ME, et al.

Unraveling the miswired connectome: A developmental perspective. Neuron. 2014.

25. Assaf M, Jagannathan K, Calhoun VD, Miller L, Stevens MC, Sahl R, et al. Abnormal

functional connectivity of default mode sub-networks in autism spectrum disorder patients.

Neuroimage. 2010;

26. Lynch CJ, Uddin LQ, Supekar K, Khouzam A, Phillips J, Menon V. Default mode

network in childhood autism: Posteromedial cortex heterogeneity and relationship with social

deficits. Biol Psychiatry. 2013;74.

27. Lau WKW, Leung MK, Lau BWM. Resting-state abnormalities in Autism Spectrum

Disorders: A meta-analysis. Sci Rep. 2019;9.

28. Biswal BB, Mennes M, Zuo X-N, Gohel S, Kelly C, Smith SM, et al. Toward discovery

science of human brain function. Proc Natl Acad Sci U S A. 2010;

29. Dumais KM, Chernyak S, Nickerson LD, Janes AC. Sex differences in default mode and

dorsal attention network engagement. PLoS One. 2018;13.

30. Scheinost D, Finn ES, Tokoglu F, Shen X, Papademetris X, Hampson M, et al. Sex

differences in normal age trajectories of functional brain networks. Hum Brain Mapp. 2015;

31. Yan CG, Craddock RC, Zuo X-N, Zang YF, Milham MP. Standardizing the intrinsic

brain: Towards robust measurement of inter-individual variation in 1000 functional

connectomes. Neuroimage. 2013;

32. Zuo X-N, Kelly C, Martino A Di, Mennes M, Margulies DS, Bangaru S, et al. Growing

Together and Growing Apart: Regional and Sex Differences in the Lifespan Developmental

Trajectories of Functional Homotopy. J Neurosci. 2010;

33. Dinstein I, Pierce K, Eyler L, Solso S, Malach R, Behrmann M, et al. Disrupted Neural

Synchronization in Toddlers with Autism. Neuron. 2011;

34. Hahamy A, Behrmann M, Malach R. The idiosyncratic brain: Distortion of spontaneous

connectivity patterns in autism spectrum disorder. Nat Neurosci. 2015;18.

35. Zang Y, Jiang T, Lu Y, He Y, Tian L. Regional homogeneity approach to fMRI data

analysis. Neuroimage. 2004;

36. Paakki JJ, Rahko J, Long X, Moilanen I, Tervonen O, Nikkinen J, et al. Alterations in

regional homogeneity of resting-state brain activity in autism spectrum disorders. Brain Res.

2010;

37. Shukla DK, Keehn B, Müller RA. Regional homogeneity of fMRI time series in autism

spectrum disorders. Neurosci Lett. 2010;476.

38. Zuo X-N, Ehmke R, Mennes M, Imperati D, Castellanos FX, Sporns O, et al. Network

centrality in the human functional connectome. Cereb Cortex. 2012;

39. Di Martino A, Zuo XN, Kelly C, Grzadzinski R, Mennes M, Schvarcz A, et al. Shared

and distinct intrinsic functional network centrality in autism and attention-

deficit/hyperactivity disorder. Biol Psychiatry. 2013;74.

40. Holiga Š, Hipp JF, Chatham CH, Garces P, Spooren W, D'Ardhuy XL, et al. Patients

with autism spectrum disorders display reproducible functional connectivity alterations. Sci

Transl Med. 2019;

41. Zou QH, Zhu CZ, Yang Y, Zuo XN, Long XY, Cao QJ, et al. An improved approach to

detection of amplitude of low-frequency fluctuation (ALFF) for resting-state fMRI:

Fractional ALFF. J Neurosci Methods. 2008;

42. Itahashi T, Yamada T, Watanabe H, Nakamura M, Ohta H, Kanai C, et al. Alterations of

local spontaneous brain activity and connectivity in adults with high-functioning autism

spectrum disorder. Mol Autism. 2015;6.

43. Ciric R, Wolf DH, Power JD, Roalf DR, Baum GL, Ruparel K, et al. Benchmarking of

participant-level confound regression strategies for the control of motion artifact in studies of

functional connectivity. Neuroimage. 2017;

44. Parkes L, Fulcher B, Yücel M, Fornito A. An evaluation of the efficacy, reliability, and

sensitivity of motion correction strategies for resting-state functional MRI. Neuroimage.

2018;

45. He Y, Byrge L, Kennedy DP. Nonreplication of functional connectivity differences in

autism spectrum disorder across multiple sites and denoising strategies. Hum Brain Mapp.

2020;1–17.

46. Abraham A, Milham MP, Di Martino A, Craddock RC, Samaras D, Thirion B, et al.

Deriving reproducible biomarkers from multi-site resting-state data: An Autism-based

example. Neuroimage. 2017;

47. Alaerts K, Nayar K, Kelly C, Raithel J, Milham MP, Di martino A. Age-related changes

in intrinsic function of the superior temporal sulcus in autism spectrum disorders. Soc Cogn

Affect Neurosci. 2015;

48. Yahata N, Morimoto J, Hashimoto R, Lisi G, Shibata K, Kawakubo Y, et al. A small

number of abnormal brain connections predicts adult autism spectrum disorder. Nat

Commun. 2016;

49. King JB, Prigge MBD, King CK, Morgan J, Weathersby F, Fox JC, et al. Generalizability

and reproducibility of functional connectivity in autism. Mol Autism. 2019;

50. Charman T, Loth E, Tillmann J, Crawley D, Wooldridge C, Goyard D, et al. The EU-

AIMS Longitudinal European Autism Project (LEAP): Clinical characterisation. Mol Autism.

2017;

51. Loth E, Charman T, Mason L, Tillmann J, Jones EJH, Wooldridge C, et al. The EU-

AIMS Longitudinal European Autism Project (LEAP): Design and methodologies to identify

and validate stratification biomarkers for autism spectrum disorders. Mol Autism. 2017;

52. Irimia A, Torgerson CM, Jacokes ZJ, Van Horn JD. The connectomes of males and

females with autism spectrum disorder have significantly different white matter connectivity

densities. Sci Rep. 2017;

53. Jenkinson M, Bannister P, Brady M, Smith S. Improved optimization for the robust and

accurate linear registration and motion correction of brain images. Neuroimage. 2002;

54. Lefebvre A, Beggiato A, Bourgeron T, Toro R. Neuroanatomical Diversity of Corpus

Callosum and Brain Volume in Autism: Meta-analysis, Analysis of the Autism Brain

Imaging Data Exchange Project, and Simulation. Biol Psychiatry. 2015;78.

55. Dennis M, Francis DJ, Cirino PT, Schachar R, Barnes MA, Fletcher JMJM. Why IQ is

not a covariate in cognitive studies of neurodevelopmental disorders. J. Int. Neuropsychol.

Soc. 2009.

56. Friston KJ, Williams S, Howard R, Frackowiak RSJ, Turner R. Movement-related effects

in fMRI time-series. Magn Reson Med. 1996;

57. Behzadi Y, Restom K, Liau J, Liu TT. A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. Neuroimage. 2007;

58. Greve DN, Fischl B. Accurate and robust brain image alignment using boundary-based

registration. Neuroimage. 2009;

59. Avants B, Tustison N, Song G. Advanced Normalization Tools (ANTS). Insight J. 2009;

60. Nielson DM, Pereira F, Zheng CY, Migineishvili N, Lee JA, Thomas AG, et al. Detecting

and harmonizing scanner differences in the ABCD study - annual release 1.0. bioRxiv. 2018;

61. Eklund A, Nichols T. How open science revealed false positives in brain imaging.

Significance. 2017;14.

62. Yeo BT, Krienen FM, Eickhoff SB, Yaakub SN, Fox PT, Buckner RL, et al. Functional

specialization and flexibility in human association cortex. Cereb Cortex. 2015;

63. Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. Large-scale automated

synthesis of human functional neuroimaging data. Nat Methods. 2011;

64. Gotham K, Pickles A, Lord C. Standardizing ADOS scores for a measure of severity in

autism spectrum disorders. J Autism Dev Disord. 2009;

65. Murphy K, Birn RM, Handwerker DA, Jones TB, Bandettini PA. The impact of global

signal regression on resting state correlations: Are anti-correlated networks introduced?

Neuroimage. 2009;

66. Pruim RHR, Mennes M, van Rooij D, Llera A, Buitelaar JK, Beckmann CF. ICA-

AROMA: A robust ICA-based strategy for removing motion artifacts from fMRI data.

Neuroimage. 2015;

67. Kong XZ, Consortium E, C F. An illustration of reproducibility in neuroscience research

in the absence of selective reporting. bioRxiv. 2020;

68. Cohen J. Statistical power analysis for the behavioural sciences. Hillside. NJ Lawrence

Earlbaum Assoc. 1988.

69. Hus V, Lord C. The autism diagnostic observation schedule, module 4: Revised algorithm

and standardized severity scores. J Autism Dev Disord. 2014;

70. Stark DE, Margulies DS, Shehzad ZE, Reiss P, Kelly AMC, Uddin LQ, et al. Regional

variation in interhemispheric coordination of intrinsic hemodynamic fluctuations. J Neurosci.

2008;

71. Floris DL, Howells H. Atypical structural and functional motor networks in autism. Prog

Brain Res. 2018.

72. Floris DL, Wolfers T, Zabihi M, Holz NE, Zwiers MP, Charman T, et al. Atypical brain

asymmetry in autism – a candidate for clinically meaningful stratification. Biol Psychiatry

Cogn Neurosci Neuroimaging. 2020;

73. Floris DL, Lai M-C, Auer T, Lombardo M V., Ecker C, Chakrabarti B, et al. Atypically

rightward cerebral asymmetry in male adults with autism stratifies individuals with and

without language delay. Hum Brain Mapp. 2016;

74. Floris DL, Chura LR, Holt RJ, Suckling J, Bullmore ET, Baron-Cohen S, et al.

Psychological correlates of handedness and corpus callosum asymmetry in autism: The left

hemisphere dysfunction theory revisited. J Autism Dev Disord. 2013;

75. Floris DL, Barber AD, Nebel MB, Martinelli M, Lai MC, Crocetti D, et al. Atypical

lateralization of motor circuit functional connectivity in children with autism is associated

with motor deficits. Mol Autism. 2016;

76. De Fossé L, Hodge SM, Makris N, Kennedy DN, Caviness VS, McGrath L, et al.

Language-association cortex asymmetry in autism and specific language impairment. Ann

Neurol. 2004;56:757–66.

77. Herbert MR, Ziegler D a, Deutsch CK, O'Brien LM, Kennedy DN, Filipek P a, et al.

Brain asymmetries in autism and developmental language disorder: a nested whole-brain

analysis. Brain. 2005;128:213–26.

78. Escalante-Mead PR, Minshew NJ, Sweeney JA. Abnormal brain lateralization in high-

functioning autism. J Autism Dev Disord. 2003;33:539–43.

79. Flagg EJ, Cardy JEO, Roberts W, Roberts TPL. Language lateralization development in

children with autism: insights from the late field magnetoencephalogram. Neurosci Lett.

2005;386:82–7.

80. Lindell AK, Hudry K. Atypicalities in cortical structure, handedness, and functional

lateralization for language in autism spectrum disorders. Neuropsychol Rev. 2013;23:257–70.

81. Zaidel E, Aboitiz F, Clarke J. Sexual dimorphism in interhemispheric relations:

Anatomical-behavioral convergence. Biol Res. 1995.

82. Proverbio AM, Brignone V, Matarazzo S, Del Zotto M, Zani A. Gender differences in

hemispheric asymmetry for face processing. BMC Neurosci. 2006;

83. Vogel JJ, Bowers CA, Vogel DS. Cerebral lateralization of spatial abilities: A meta-

analysis. Brain Cogn. 2003;

84. Kirkovski M, Enticott PG, Hughes ME, Rossell SL, Fitzgerald PB. Atypical Neural

Activity in Males But Not Females with Autism Spectrum Disorder. J Autism Dev Disord.

2016;

85. Lai MC, Lombardo M V., Chakrabarti B, Ruigrok ANV, Bullmore ET, Suckling J, et al.

Neural self-representation in autistic women and association with 'compensatory

camouflaging.' Autism. 2019;

86. Lai MC, Lombardo M V., Suckling J, Ruigrok ANV, Chakrabarti B, Ecker C, et al.

Biological sex affects the neurobiology of autism. Brain. 2013;

87. Lai M-C, Lombardo M V., Auyeung B, Chakrabarti B, Baron-Cohen S. Sex/Gender

Differences and Autism: Setting the Scene for Future Research. J. Am. Acad. Child Adolesc.

Psychiatry. 2015.

88. Moseley RL, Ypma RJF, Holt RJ, Floris D, Chura LR, Spencer MD, et al. Whole-brain

functional hypoconnectivity as an endophenotype of autism in adolescents. NeuroImage Clin.

2015;

89. Kennedy DP, Courchesne E. Functional abnormalities of the default network during self-

and other-reflection in autism. Soc Cogn Affect Neurosci. 2008;

90. Hull J V., Jacokes ZJ, Torgerson CM, Irimia A, Van Horn JD, Aylward E, et al. Resting-

state functional connectivity in autism spectrum disorders: A review. Front. Psychiatry. 2017.

91. Tang S, Sun N, Floris DL, Zhang X, Di Martino A, Yeo BTT. Reconciling Dimensional

and Categorical Models of Autism Heterogeneity: a Brain Connectomics & Behavioral

Study. Biol Psychiatry. 2019;

92. Barch DM, Yarkoni T. Introduction to the special issue on reliability and replication in

cognitive and affective neuroscience research. Cogn. Affect. Behav. Neurosci. 2013.

93. Poldrack RA, Poline JB. The publication and reproducibility challenges of shared data.

Trends Cogn. Sci. 2015.

94. Gorgolewski KJ, Nichols T, Kennedy DN, Poline JB, Poldrack RA. Making replication

prestigious. Behav. Brain Sci. 2018.

95. Gratten J, Wray NR, Keller MC, Visscher PM. Large-scale genomics unveils the genetic

architecture of psychiatric disorders. Nat. Neurosci. 2014.

96. Sanders SJ, He X, Willsey AJ, Ercan-Sencicek AG, Samocha KE, Cicek AE, et al.

Insights into Autism Spectrum Disorder Genomic Architecture and Biology from 71 Risk

Loci. Neuron. 2015;87.

97. Zuo XN, Xing XX. Test-retest reliabilities of resting-state FMRI measurements in human

brain functional connectomics: A systems neuroscience perspective. Neurosci. Biobehav.

Rev. 2014.

98. Zuo XN, Xu T, Milham MP. Harnessing reliability for neuroscience research. Nat. Hum.

Behav. 2019.

99. Feczko E, Miranda-Dominguez O, Marr M, Graham A, Nigg J, Fair D. The

Heterogeneity Problem: Approaches to Identify Psychiatric Subtypes. Trends Cogn Sci.

2019;23.

100. Marquand AF, Wolfers T, Mennes M, Buitelaar J, Beckmann CF. Beyond Lumping and

Splitting: A Review of Computational Approaches for Stratifying Psychiatric Disorders. Biol

Psychiatry Cogn Neurosci Neuroimaging. Elsevier; 2016;1:433–47.

101. Lombardo M V., Lai MC, Baron-Cohen S. Big data approaches to decomposing

heterogeneity across the autism spectrum. Mol Psychiatry. 2019;

102. Yokota S, Takeuchi H, Hashimoto T, Hashizume H, Asano K, Asano M, et al.

Individual differences in cognitive performance and brain structure in typically developing

children. Dev Cogn Neurosci. 2015;

103. Marquand AF, Kia SM, Zabihi M, Wolfers T, Buitelaar JK, Beckmann CF.

Conceptualizing mental disorders as deviations from normative functioning. Mol Psychiatry

[Internet]. 2019; Available from: http://www.nature.com/articles/s41380-019-0441-1

104. van Dijk KRA, Sabuncu MR, Buckner RL. The influence of head motion on intrinsic

functional connectivity MRI. Neuroimage. 2012;59.

105. Hong S-J, Vogelstein JT, Gozzi A, Bernhardt BC, Yeo BTT, Milham MP, et al.

Towards Neurosubtypes in Autism. Biol Psychiatry. 2020;

106. Zabihi M, Oldehinkel M, Wolfers T, Frouin V, Goyard D, Loth E, et al. Dissecting the

Heterogeneous Cortical Anatomy of Autism Spectrum Disorder Using Normative Models.

Biol Psychiatry Cogn Neurosci Neuroimaging. 2019;

107. Floris DL, Wolfer T, Zabihi M, Holz N, Zwiers M, Charman T, et al. Atypical brain

asymmetry in autism - a candidate for clinically meaningful stratification. BioRxiv. 2020;

108. Marquand AF, Rezek I, Buitelaar J, Beckmann CF. Understanding Heterogeneity in

Clinical Cohorts Using Normative Models: Beyond Case-Control Studies. Biol Psychiatry.

2016;

109. Strang JF, van der Miesen AI, Caplan R, Hughes C, daVanport S, Lai M-C. Both sex-

and gender-related factors should be considered in autism research and clinical practice.

Autism. 2020;

110. Charman T, Taylor E, Drew A, Cockerill H, Brown JA, Baird G. Outcome at 7 years of

children diagnosed with autism at age 2: Predictive validity of assessments conducted at 2

and 3 years of age and pattern of symptom change over time. J Child Psychol Psychiatry

Allied Discip. 2005;

111. Fecteau S, Mottron L, Berthiaume C, Burack JA. Developmental changes of autistic

symptoms. Autism. 2003;

112. Lin HY, Ni HC, Lai M-C, Tseng WYI, Gau SSF. Regional brain volume differences

between males with and without autism spectrum disorder are highly age-dependent. Mol

Autism. 2015;

Figure Captions

Fig 1. Overlap across R-fMRI metrics for main effects of diagnosis and sex

Upper panel: the surface inflated maps depict the extent of overlap across any of three resting state fMRI (R-fMRI) metrics for clusters showing significant main effects of diagnosis (left) and sex (right). Purple clusters represent areas of significant group differences emerging for only one of any of the three R-fMRI measures, orange and yellow clusters indicate measures with overlap among 2 and 3 R-fMRI measures (see Additional file 5 for statistical maps of main effects for each R-fMRI metric). Cluster masks are overlaid on inflated brain maps generated by BrainNet Viewer. Lower panel: for each of the yellow and orange clusters in panel A, the table lists the cluster's anatomical area label based on the Harvard Oxford atlas, the specific R-fMRI metrics involved, and the group difference direction (ASD<NT or M<F in blue, ASD>NT or M>F in red; see Additional file 3: Table S6 for more details).
Abbreviations: L= Left hemisphere; R= Right hemisphere, PCG/FP=Paracingulate cortex/frontal pole, ACC=Anterior cingulate cortex, PCC/Prec=Posterior cingulate cortex/precuneus, ASD=autism spectrum disorder, NT=neurotypical, M=Males, F=Females).

Fig 2. Sex-by-diagnosis interaction effect, its robustness and replicability

a) On the right, the surface maps show the cluster with a significant ($Z > 3.1$, $P<0.01$) sex-by-diagnosis interaction for voxel-mirrored homotopic connectivity (VMHC) resulting from discovery analyses in the ABIDE sample using the component-based noise reduction (CompCor) pipeline. The statistical Z maps are overlaid on inflated brain maps generated by BrainNet Viewer. **b)** The upper panels show the pattern of VMHC group means in males and females by each diagnostic group (ASD and NT) extracted from the same cluster in pre-processed data following two alternative denoising pipelines, Global Signal Regression

(GSR, left) and Independent Component Analysis – Automatic Removal of Motion Artifacts

(ICA-AROMA, right). Results show a pattern similar to the primary results observed in

discovery analyses with moderate to high effect sizes (Cohen's d range= 0.5-0.6). **2c)** The

lower graph shows replicability in two independent samples: the Gender Explorations of

Neurogenetics and Development to Advance Autism Research (GENDAAR) and the EU-

AIMS Longitudinal European Autism Project (LEAP). The pattern of results was replicable

in the EU-AIMS LEAP (N=309) with a moderate effect size (Cohen's $d = 0.34$) and had a

negligible effect size in GENDAAR (N=196; Cohen's $d = 0.09$). For all graphs VMHC data

are shown as residuals obtained after regressing out mean framewise displacement and age

effects. **Abbreviations:** L=left, R=right, A=anterior, P=posterior.

Fig 3. Functional relevance of sex-by-diagnosis interaction in VMHC

a) The radar plot shows the percentage (0-80%) of overlap between the voxels in the

dorsolateral occipital cluster showing a significant VMHC sex-by-diagnosis interaction in

discovery analyses and the 12 Yeo cognitive ontology probability maps [62] (probability

threshold at $P = 1e-5$) for cognitive components C1-C12. As in Floris et al. [2], we labelled

each component based on the top five tasks reported to be most likely recruited by a given

component. **b)** Word cloud based on the top 27 terms showing correlations between r=0.64 to

r=0.10 associated with the same VMHC cluster based on the Neurosynth Image Decoder. **c)**

Sex-differential association between each individual's VMHC at the cluster showing a

significant sex-by-diagnosis interaction in primary analyses and available ADOS social-affect

uncalibrated sub-scores in males and females with ASD. VMHC data are shown as residuals

obtained after regressing out mean framewise displacement and age effects. While males

showed no significant associations at corrected and uncorrected thresholds, females with

lower dorsolateral occipital VMHC showed more severe social-affect symptoms at

uncorrected statistical threshold ($F_{(1,311)}$=4.44, p=0.036).

Fig 4. Robustness and replicability summary

a) The histogram summarizes the percentage of clusters showing a robust and replicable

pattern of results as that observed in discovery analyses in the ABIDE sample for main

effects of diagnosis (Dx; green; N=10 clusters), sex (yellow; N=7 clusters) and their

interaction (blue; N=1 cluster) across three R-fMRI metrics. All findings were robust to

different preprocessing pipelines. Across R-fMRI metrics, main sex effects were moderately

(60%) to largely (90%) replicable across independent samples: Gender Explorations of

Neurogenetics and Development to Advance Autism Research (GENDAAR) and the EU-

AIMS Longitudinal European Autism Project (LEAP), respectively. Replicability for main

effects of diagnosis was moderately replicable in GENDAAR (57%) and minimally

replicable in EU-AIMS LEAP (14%). The VMHC pattern observed for sex-by-diagnosis

interaction in primary analyses was replicated in EU-AIMS LEAP only. **b)** Surface

conjunction maps show the clusters replicated in GENDAAR only (G, purple), EU-AIMS

LEAP only (E, blue) and in both samples (G&E, red) for each effect separately. Cluster

masks are overlaid on inflated brain maps generated by BrainNet Viewer.

Additional files

Additional file 1

Title: Table 1

File format: docx

Description: Characterization of sample merged across ABIDE I and II

Additional file 2

Title: Supplemental Material

File format: docx

Description: Supplemental Methods

Additional file 3

Title: Supplementary Tables

File format: docx

Description: Characterization of EU-AIMS LEAP and GENDAAR samples, comparison

between samples and summary table of main effects.

Additional file 4

Title: Selection flowchart for the ABIDE sample

File format: tif

Description: The flowchart illustrates the selection process resulting in the final ABIDE I

and II combined sample of 1019 subjects; each step is detailed in Supplementary material.

Abbreviations: ASD=autism spectrum disorder, NT=neurotypical, A I=ABIDE I.

Additional file 5

Title: Main effects of diagnosis and sex in the ABIDE discovery sample

File format: png

Description: Significant results ($Z > 3.1$, $P < 0.01$, corrected) of voxel-wise discovery

analyses conducted in the ABIDE dataset for main effects (ME) of diagnosis (left) and sex

(right) for seed-based intrinsic functional connectivity of the posterior cingulate cortex-

(PCC), voxel mirror homotopic connectivity (VMHC), and Regional Homogeneity (ReHo).

Significant clusters are overlaid on inflated brain maps generated by BrainNet Viewer. No

significant effects were detected for degree centrality or fractional amplitude of low

frequency fluctuations. **ME Diagnosis**: <u>PCC-iFC:</u> bilateral paracingulate cortex and frontal

pole (PCG/FP), superior lateral occipital cortex (sLOC), temporal occipital fusiform cortex

and occipital fusiform gyrus (TOFC/OFC); <u>VMHC</u>: bilateral posterior cingulate gyrus and

precuneus (PCC/Prec), PCG/FP; <u>ReHo</u>: PCG/FP, central operculum and insula (CO/Ins). **ME**

Sex: <u>PCC-iFC</u>: bilateral sLOC, middle frontal gyrus (MFG), bilateral PCC/Prec, bilateral

PCG/FP; <u>VMHC</u>: bilateral PCC/Prec, bilateral anterior cingulate cortex (ACC); <u>ReHo</u>:

bilateral PCC, angular gyrus and lateral occipital cortex (AnG/LOC). See Additional file 3:

Table S6 for details on each cluster sizes. *Due to processing failure of two subjects for

VMHC, the sample size comprised 1017 subjects instead of 1019.

Additional file 6

Title: Robustness to nuisance corrections of main effects of diagnosis and sex

File format: png

Description: Cluster-level replication of the results emerging from the voxel-wise discovery

analyses in the ABIDE dataset preprocessed using CompCor for the statistically significant

main effects of diagnosis (**Figure S1a)** and sex (**Figure S1b**) after preprocessing with GSR

and with ICA-AROMA. The second column on the left shows the clusters ($Z > 3.1$, $P < 0.01$,

corrected) with significant diagnostic and sex effects for posterior cingulate cortex intrinsic

functional connectivity (PCC-iFC), voxel-mirrored homotopic connectivity (VMHC), and

regional homogeneity (ReHo). Results are overlaid on inflated brain maps generated by

BrainNet Viewer. The bar plots represent the residual means resulting from regressing out

diagnosis or sex effects depending on the desired main effect from the cluster means. **Figure**

S1a) The ABIDE GSR and ABIDE ICA-AROMA columns illustrate, for each of these R-

fMRI indices, the diagnostic group mean pattern across clusters with a diagnostic effect size

of $d \geq 0.2$. Color codes: Red=ASD; Green=NT. <u>PCC-iFC</u>: bilateral paracingulate cortex and

frontal pole (PCG/FP), superior lateral occipital cortex (sLOC), temporal occipital fusiform

1112 cortex and occipital fusiform gyrus (TOFC/OFC); <u>VMHC</u>: bilateral posterior cingulate gyrus

1113 and precuneus (PCC/Prec), PCG/FP; <u>ReHo</u>: PCG/FP, central operculum and insula (CO/Ins).

1114 ABIDE GSR: 7 out of 7 main effects of diagnosis replicated (100%). ABIDE ICA-AROMA:

1115 7 out of 7 main effects of diagnosis replicated (100%) **Figure S1b)** The ABIDE GSR and

1116 ABIDE ICA-AROMA columns illustrate, for each of these R-fMRI indices, the sex group

1117 mean pattern across clusters with an effect size of Cohen' $d \geq 0.2$. Color codes: Blue=males;

1118 Pink=females. <u>PCC-iFC</u>: bilateral sLOC, middle frontal gyrus (MFG), bilateral PCC/Prec,

1119 bilateral PCG/FP; <u>VMHC</u>: bilateral PCC/Prec, bilateral anterior cingulate cortex (ACC);

1120 <u>ReHo</u>: bilateral PCC, angular gyrus and lateral occipital cortex (AnG/LOC). ABIDE GSR: 10

1121 out of 10 main effects of diagnosis replicated (100%). ABIDE ICA-AROMA: 10 out of 10

1122 main effects of diagnosis replicated (100%). Due to processing failure of two subjects for

1123 VMHC, the sample size comprised 1017 subjects instead of 1019. VMHC data are shown as

1124 residuals obtained after regressing out mean framewise displacement and age effects.

1125 **Abbreviations:** ASD=autism spectrum disorder, NT=neurotypical, M=males, F=females,

1126 CompCor=component base noise reduction, GSR=Global Signal Regression, ICA-

1127 AROMA=independent component analysis – automatic removal of motion artifacts, PCC-

1128 iFC=posterior cingulate cortex intrinsic functional connectivity (x=0, y=-53, z=26),

1129 VMHC=voxel-mirrored homotopic connectivity, ReHo=regional homogeneity, L=left,

1130 R=right, R+=replication based on same direction of results and Cohen's d $\geq$ 0.2, R-=non-

1131 replication of results (displayed in gray plots).

1132 **Additional file 7**

1133 **Title:** Stability of main effects

1134 **File format:** png

1135 **Description:** Inter-site stability was assessed after extracting group means at masks

1136 corresponding to the clusters showing significant main effects of diagnosis (**Figure S2a**) and

sex (**Figure S2b**) in the discovery analyses and then deriving the group mean when leaving

one acquisition site out at the time. The pattern of results was unchanged. Different ABIDE

sites are color-coded on legend on the side. Due to processing failure of two subjects for

VMHC, the sample size comprised 1017 subjects. **Abbreviations**: ASD=autism spectrum

disorder, NT=neurotypical, PCC-iFC=posterior cingulate cortex intrinsic functional

connectivity (x=0, y=-53, z=26), VMHC=voxel-mirrored homotopic connectivity,

ReHo=regional homogeneity, L=left, R=right. Different sites in ABIDE are color-coded on

the top left. Data are shown as residuals obtained after regressing out mean framewise

displacement and age effects.

Additional file 8

Title: Stability of sex-by-diagnosis interaction effect

File format: tif

Description: Inter-site stability of the sex-by-diagnosis interaction pattern was assessed after

extracting group means at the mask corresponding to the clusters showing a significant

interaction in the discovery analyses and then deriving the group mean when leaving one

acquisition site out at the time. The pattern of results was unchanged. Different sites in

ABIDE are color-coded in the legend on the right. Due to processing failure of two subjects

for VMHC, the sample size comprised 1017 subjects. **Abbreviations**: ASD=autism spectrum

disorder, NT=neurotypical, PCC-iFC=posterior cingulate cortex intrinsic functional

connectivity (x=0, y=-53, z=26), VMHC=voxel-mirrored homotopic connectivity,

ReHo=regional homogeneity, L=left, R=right. Different sites in ABIDE are color-coded on

the top left.

Additional file 9

Title: Replicability of main effects of diagnosis and sex

File format: png

Description: Cluster-level replication of the results emerging from the voxel-wise analyses in the ABIDE dataset for main effects of diagnosis (**Figure S3a**) and sex (**Figure S3b**) in the Gender Explorations of Neurogenetics and Development to Advance Autism Research (GENDAAR) and the EU-AIMS Longitudinal European Autism Project (LEAP) samples. The ABIDE column on the left shows the clusters ($Z > 3.1$, $P < 0.01$, corrected) with significant diagnostic and sex effects for posterior cingulate cortex intrinsic functional connectivity (PCC-iFC), voxel-mirrored homotopic connectivity (VMHC), and regional homogeneity (ReHo). No significant effects were detected for degree centrality and fractional amplitude of low frequency fluctuations. Results are overlaid on inflated brain maps generated by BrainNet Viewer. The bar plots represent the residual means resulting from the linear Gaussian regression for each group. **Figure S3a)** The GENDAAR and EU-AIMS LEAP columns illustrate, for each R-fMRI index, the diagnostic group mean pattern across clusters with a diagnostic effect size of $d \geq 0.2$. Color codes: Red=ASD; Green=NT. <u>PCC-iFC</u>: bilateral paracingulate cortex and frontal pole (PCG/FP), superior lateral occipital cortex (sLOC), temporal occipital fusiform cortex and occipital fusiform gyrus (TOFC/OFC); <u>VMHC</u>: bilateral posterior cingulate gyrus and precuneus (PCC/Prec), PCG/FP; <u>ReHo</u>: PCG/FP, central operculum and insula (CO/Ins). <u>GENDAAR</u>: 4 out of 7 main effects of diagnosis replicated (57.1%); <u>EU-AIMS LEAP</u>: 1 out of 7 main effects of diagnosis replicated (14.3%). **Figure S3b)** The GENDAAR and EU-AIMS LEAP columns illustrate, for each of these R-fMRI index, the sex group mean pattern across clusters with an effect size of $d \geq 0.2$. Color codes: Blue=males, Pink=females. <u>PCC-iFC</u>: bilateral sLOC, middle frontal gyrus (MFG), bilateral PCC/Prec, bilateral PCG/FP; <u>VMHC</u>: bilateral PCC/Prec, bilateral anterior cingulate cortex (ACC); <u>ReHo</u>: bilateral PCC, angular gyrus and lateral occipital cortex (AnG/LOC). <u>GENDAAR</u>: 5 out of 10 main effects of sex replicated (50%); <u>EU-AIMS</u>

<u>LEAP</u>: 9 out of 10 main effects of sex replicated (90%). *Due to processing failure of two

subjects for VMHC, the sample size comprised 1017 subjects instead of 1019.

Abbreviations: ASD=autism spectrum disorder, NT=neurotypical, M=males, F=females,

PCC-iFC=posterior cingulate cortex intrinsic functional connectivity (x=0, y=-53, z=26),

VMHC=voxel-mirrored homotopic connectivity, ReHo=regional homogeneity, L=left,

R=right, R+=replication based on same direction of results and Cohen's $d \geq 0.2$, R-=non-

replication of results.

1193

1194 Figure 1

1195

1196

1197   Figure 2

1205　　Figure 3

a) Overlap of VMHC cluster with cognitive ontology maps

b) Neurosynth terms correlation with VMHC cluster

c) Correlation between social-affect ADOS scores and VMHC cluster

1206

1207   Figure 4

1208