Living in the concrete jungle: carnivore spatial ecology in urban parks

Siria Gámez[1], Nyeema C. Harris[1]

[1]Applied Wildlife Ecology Lab, Ecology and Evolutionary Biology, University of Michigan
1101 N. University Ave, Ann Arbor, Michigan 48106

Corresponding author: Siria Gámez (sgamez@umich.edu)

ABSTRACT

People and wildlife are living in an increasingly urban world, replete with unprecedented human

densities, sprawling built environments, and altered landscapes. Such anthropogenic pressures

can affect processes at multiple ecological scales from individuals to ecosystems, yet few studies

integrate two or more levels of ecological organization. We tested two competing hypotheses,

humans as shields versus humans as competitors, to characterize how humans directly affect

carnivore spatial ecology across three scales. From 2017-2020, we conducted the first camera

survey of city parks in Detroit, Michigan, and obtained spatial occurrence data of the local native

carnivore community which included coyotes (*Canis latrans*), red foxes (*Vulpes vulpes*), gray

foxes (*Urocyon cinereoargenteus*), raccoons (*Procyon lotor*), and striped skunks (*Mephitis*

mephitis). We constructed single-species occupancy models to discriminate parks into areas of

low and high human presence using the average human occupancy as the threshold. At the

community level, carnivores were spatially aggregated at both high and low human use sites,

though aggregation was stronger at sites with high human occupancy, indicating that other

factors such as available resources rather than human presence might influence community

segregation. Three carnivore species pairs changed from positive association at low human

occupancy to negative association at high human occupancy, two subordinate species pairs

aligning with the humans as competitors hypothesis, and an apex-subordinate pair consistent

with the humans as shields hypothesis. At the individual level, human hotspots of high site use

did not overlap with any carnivore activity hotspots, similarly lending support to the humans as

competitors hypothesis. Overall, we found inference on anthropogenic impacts on carnivore

spatial ecology varied depending on the scale of ecological organization. Our findings

demonstrate how urban carnivores are exploiting spatial refugia in the cityscape.

Keywords: city, co-occurrence, distribution, community structure, camera survey, Detroit,

human shield, coyote, overlap

INTRODUCTION

Cities are highly heterogeneous landscapes of risk and reward, borne of unique

interactions between anthropogenic and ecological processes (Alberti et al., 2003; Liu et al.,

2007). As urbanization and land cover conversion rates continue to increase worldwide, cities

have emerged as a new and unique habitat for wildlife. By 2050, over half of the global human

population will live in a city while urban development is projected to grow by 120 million

hectares globally by 2030 (McDonald et al., 2018; United Nations, 2018). Cities can be a source

or a sink for mammal species, a duality driven by both increases in availability of food sources

and risks of mortality (Bateman, & Fleming, 2012; Lepczyk et al., 2017). For example, black

bears (*Ursus americanus*) in urban areas of Nevada had higher birth rates than their rural and

protected area counterparts, but also experienced higher age-specific mortality rates, which

significantly decreased fitness (Beckmann, & Lackey, 2008). Cougars (*Puma concolor*) in an

urban-wildland system in Colorado successfully exploited anthropogenic food sources, yet faced

a 6.5% increase in mortality risk in developed areas (Moss et al., 2016). Wildlife responses to the

built environment are unsurprisingly driven by humans themselves and their induced

modifications to landscapes through food provisioning, artificial habitat and light, and roads

(Clucas, & Marzluff, 2011; Gaston et al., 2017; Riley et al., 2014).

Anthropogenic pressures affect wildlife at multiple levels of ecological organization,

from community structure and intraguild interactions down to behavioral shifts in individual

species. Perturbations to higher trophic levels can have cascading impacts on ecosystem

processes, which underscores the need to understand how carnivores respond to human activities

(Ripple et al., 2014b; Terborgh, 2010). A study in the city of Chicago found that raccoons

(*Procyon lotor*) comprised a larger relative proportion of the mesopredator community in urban

compared to rural sites, irrespective of patch size (Prange, & Gehrt, 2004). Intraguild interactions

are also mediated by anthropogenic pressure the human shield effect (Berger, 2007; Gallo et al.,

2019; Muhly et al., 2011). Red foxes (*Vulpes vulpes*) have been shown to exploit highly

developed core urban areas as spatial refugia to avoid their dominant coyote (*Canis latrans*)

competitors (Moll et al., 2018). Individual species responses to human activity are varied and

depend on each species' biology and tolerance to human encounters. Cottontail rabbits

(*Sylvilagus floridanus*) in an urban area were more vigilant at sites where coyotes were absent,

suggesting humans are a third "player" in a predator-prey-human system (Gallo et al., 2019).

Striped skunk (*Mephitis mephitis*) occupancy was greatest near urban areas, but this positive

association weakened as the percentage of urban land cover increased, signaling a sensitivity to

within-city metrics of anthropogenic pressure (Ordeñana et al., 2010). Evidently, urban systems

produce a suite of complex and synergistic changes to processes at multiple ecological scales.

Despite evidence that human activity induces complex responses in urban wildlife, there

is a dearth of studies that to quantify these effects at multiple scales of ecological organization,

particularly for terrestrial carnivores. Further, the field of urban ecology is relatively new and the

majority of approaches to characterizing effects on wildlife have been dominated by the use

landscape-level metrics of the built environment as proxies for human activity (Wu, 2014). A

meta-analysis of urban ecology studies found that only 10.2% of 244 studies quantified large

mammal responses to urbanization and only 6% of all urbanization metrics employed in these

studies explicitly considered humans (Moll et al., 2019). Worldwide population declines and

range contraction in carnivores highlight the urgency to assess how spaces dominated by humans

alter ecological interactions at community, population, and individual levels (Ceballos, &

Ehrlich, 2002; Ripple et al., 2014a).

We leveraged a North American native carnivore guild comprised of coyotes, raccoons,

red foxes, gray foxes (*Urocyon cinereoargenteus)*, and striped skunks to investigate how human

occupancy influences spatial ecology at the individual, pairwise, and community level.

Specifically, we implemented the first camera survey of city parks in Detroit, Michigan from

2017-2020 to study the city's carnivores. By directly measuring human activity and not proxies

of human pressure such as housing density, we explicitly disentangled the effects of humans on

wildlife from those related to the built environment.

In characterizing human effects on the carnivore community, two theoretical frameworks

emerge with distinct expectations for mammalian community response to fine-scale human

activity (Figure 1). The humans as shields hypothesis (HSH) argues that humans differentially

exert top-down pressure on the apex predator in the system, indirectly benefitting the subordinate

competitors and facilitating greater spatial overlap between humans and subordinate species

(Moll et al., 2018; Shannon et al., 2014). A contrasting approach frames humans as competitors

(HCH), asserting that anthropogenic pressure affects multiple species irrespective of their trophic

level or dominance hierarchy (Chapron, & López-Bao, 2016; Farris et al., 2017). The HCH

asserts that human presence is functionally similar to antagonism from another competitor in the

guild, resulting in increased vigilance, competitive exclusion, and spatial avoidance across the

entire community (Gallo et al., 2019).

Here, we addressed the following questions to test whether effects from anthropogenic

pressures on native urban carnivores align with expectations of the HSH and HCH at different

levels of ecological organization: 1) how does human activity in city parks affect the community

structure? 2) how are pairwise interactions affected by human activity within a competing

carnivore guild? 3) how do individual species of carnivores respond to areas of high human

activity? We evaluate the ecological role of humans in urban carnivore communities according to

these two contrasting frameworks:

Humans as shields hypothesis (HSH)

With HSH, subordinate mesopredators will exploit the spatial refugia created by human

top-down pressures on apex predators and spatially overlap with humans at the park scale

(Geffroy et al., 2015; Moll et al., 2018). 1) At the community level, we expected spatial

avoidance (i.e. low co-occurrence among carnivores) at low human activity versus spatial

aggregation (i.e. increased spatial co-occurrence) at high human activity owing to multiple

subordinate species exploiting spatial refugia (Amarasekare, 2003; Gehrt et al., 2013). 2) Within

pairwise interactions, we expected increased apex carnivore activity and reduced subordinate

carnivore activity at parks with low human activity versus reduced apex activity and increased

subordinate activity at parks with high human activity (Smith et al., 2018). 3) At the individual

level, we expected less spatial overlap between human and dominant carnivore hotspots as well

as greater overlap between human and subordinate carnivore hotspots (Berger, 2007; Muhly et

al., 2011).

Humans as competitors hypothesis (HCH)

Conversely, humans will function like a superior competitor, reduce the niche space and

thus spatially displace carnivores regardless of whether they are apex or subordinate with HCH

(Everatt et al., 2019). 1) At the community level, we expect greater co-occurrence and spatial

aggregation of carnivores at low human activity versus low co-occurrence and spatial avoidance

at parks with high human activity (Schuette et al., 2013). 2) Within pairwise interactions, we

expect increased activity and greater co-occurrence for all carnivore species in parks with low

human activity versus reduced activity and low co-occurrence for all carnivore species in parks

with increased human activity. 3) At the individual level, we expect separate, non-overlapping

hotspots for humans and all other carnivore species (George, & Crooks, 2006).

MATERIALS & METHODS

Study Area

We surveyed the carnivore community at 24 urban parks throughout Detroit, a ~ 370 km²

city in southeastern Michigan, USA using remotely triggered cameras (Figure 2). Parks sampled

as part of our study represent 51% of the total area of the green space in city parks (City of

Detroit, 2015). The Detroit River runs along the city's southern boundary and the Rouge River

runs through the southwestern districts including an automotive manufacturing plant. Over the

last 70 years, Detroit has experienced a substantial population decline, dropping from 1.8 million

residents at the height of its industrial era in 1950 to approximately 673,000 residents in 2016

(U.S. Census Bureau, 2016). Current human population density is 1,819/km compared to 2,374/

km found in the smaller mid-western city of Milwaukee, Wisconsin. The number of empty lots

peaked at roughly 120,000 vacant lots (33% of total parcels) and 48,000 abandoned buildings

(13% of parcels) in 2010 (Detroit Residential Parcel Survey, 2010; Raleigh, & Galster, 2015).

Over time, many empty lots have progressed through early successional stages and have even

developed enough vegetative cover to support small mammal populations, a key prey source for

urban carnivores (Bateman, & Fleming, 2012).

Camera Survey

We conducted a 3-year, non-invasive survey by installing motion-triggered trail cameras

(Reconyx© PC 850, 850C, 900, 900C) throughout city parks during the fall-winter season in

Detroit (November 2017-March 2018, November 2018-February 2019, November 2019 – March

2020). We deployed 39 stations across 23 city parks in 2017, 41 stations across 24 parks in 2018,

and 36 stations at 23 parks in 2019; the first such camera survey in Detroit. We selected parks to

ensure representation of ecological and anthropogenic features such as park size, vegetation

cover, distance to water, trails, and built infrastructure such as visitor pavilions and playgrounds.

To improve the detection probability of mammals, camera placement was informed animal sign

such as tracks, scat, or natural trails. Unbaited cameras were placed approximately 0.5 m from

the ground on trees >10 cm in diameter, following standard protocol for mesocarnivore camera

trap studies (Cove et al., 2012). Camera settings were set to high sensitivity, with three images

captured per trigger at 1s intervals, and 15s delay between triggers. For parks with > 1 camera

station, the average distance between cameras was 1,416 m, while the average distance between

parks was 3,200 m.

After camera retrieval, at least two members of the [redacted for peer review] classified

images to species and confirmed accuracy. Any unresolved photos were classified as "unknown"

and removed from the analysis. We implemented a 30-minute quiet period to account for

pseudoreplication and improve independence for analysis, given some animals tend to remain in

front of the camera and trigger it multiple times. We categorized skunks, raccoons, red foxes,

and gray foxes as subordinate, based on both their relatively smaller body sizes (< 10 kg) and a

shared history of intraguild killing and antagonistic interactions with the dominant carnivore in

this system, the coyote (Fedriani et al., 2000). Domestic dogs and cats were excluded from the

analysis as we could not differentiate between feral animals who form part of the local carnivore

community and those which were temporarily off-leash or roaming from their owners. Focal

species in this study are relatively common in the eastern United States and comprise a guild that

is hierarchically structured, ideal for investigating the effects of human presence on the space use

of a carnivore community in urban environments.

Modeling Human Occupancy

To quantify human activity at each park, we constructed single-species, single-season

occupancy models with year as a covariate (MacKenzie et al., 2003; Mackenzie, & Royle, 2005).

We generated weekly detection histories (i.e. presence '1' or absence '0' of humans at each

camera location) using the R package 'camtrapR' (Niedballa, 2016). The occupancy modeling

framework considers the maximum likelihood of both the process of detection (p) and the state

of occupancy (φ) given the data, each with covariates which might potentially explain

heterogeneity (MacKenzie et al., 2003). To build the detection model while holding occupancy

constant, we first included covariates such as year (YR), understory vegetation at camera (VEG),

number of trap nights (TN). We then used the top detection model to build the occupancy model,

with distance to nearest school (DSCH), and park size (AREA) as covariates explaining

occupancy (Table S1). Top models were selected using an information theoretic approach using

Akaike's Information Criterion (AIC) to identify the model with the lowest ΔAIC and greatest

weight (w). Model goodness-of-fit was assessed using a chi-squared discrepancy method

(MacKenzie, & Bailey, 2004).

Human occupancy estimates were then calculated at the camera level; for parks

containing >1 camera station, we averaged occupancy model estimates for all cameras within the

park. We calculated the global average of human occupancy for all the parks in the study to

produce the threshold value for categorizing parks as having either "high" (above average

occupancy) versus "low" (below average occupancy) human activity.

Community-Level Response to Human Activity

To compare the carnivore community structure at varying levels of human occupancy, we

first pooled detections at camera stations according to park and subsequently created a species'

detections per park matrix, allowing for park-level analyses of community composition and

spatial structure. We then used the R package 'EcoSimR' to test for significant spatial

aggregation or segregation in community assemblages. *EcoSimR* uses a null-model approach and

generates random matrices (n=1000) using Markov chain Monte Carlo (MCMC) methods. The

null-model matrix approach assumed: 1) species rarity and species richness vary at each site; and

2) for each iteration, both the number of parks each species occurs at as well the number of

species occurring at each park was held constant (Gotelli et al., 2015). We used "Simulation 10"

in the EcoSimR package where each species' occurrence at a park was weighted by its overall

prevalence throughout the study area, such that the presence of a rare species was not equivalent

to the presence of a common one (Gotelli et al., 2015).

We calculated c-scores for parks at low vs high human occupancy, which were then

compared to the aggregated c-score for the randomly generated null model matrices (Stone, &

Roberts, 1990). Negative c-scores or values close to zero indicate clustering or greater co-

occurrence among all species in the community than expected by random chance. In contrast,

large observed c-scores indicate spatial avoidance or lower co-occurrence than expected. If the

community structure was clustered significantly at high human occupancy and segregated at low

occupancy, our interpretation was that humans were facilitating positive spatial co-occurrence

among subordinate species consistent with the HSH (Berger, 2007). Alternatively, we interpreted

segregation of the community structure at a high human activity and aggregation at low activity

as evidence of spatial partitioning and competition avoidance, following the HCH (de Satgé et

al., 2017).

Two additional scenarios exist: segregation at both low and high human occupancy,

indicating spatial avoidance between competing carnivores irrespective of human activity as well

as aggregation at both low and high human occupancy, indicating that carnivores are spatially

clustered within city parks regardless of human activity. Both of these latter scenarios would

suggest that the spatial structure of the carnivore community is determined not by the degree of

human activity, but by other park and landscape attributes such as park size, vegetation, access to

water, or food availability (Gompper et al., 2016).

Pairwise Interspecific-Level Response to Human Activity

To understand how human activity affects paired interspecific spatial relationships, we

compared independent detections for four species (coyote, raccoon, red fox, skunk) at parks

designated as high human occupancy versus low. Grey foxes were only detected at 1 park and

were thus excluded from this component of our analysis. Using the species detections matrix

from the community-level analysis where each park is a spatial replicate, we calculated

Spearman's R correlation coefficients for each species pair to test for significant positive or

negative associations. In support of the HSH, we expected humans and subordinate carnivores to

be positively correlated while for humans and coyotes to be negatively correlated at parks where

human occupancy was high. Additionally, we expected the subordinate species to be negatively

correlated with coyotes at low human occupancy, indicating competitive spatial partitioning in

the absence of the "shield", though we expected the raccoon and skunk response to be weaker

(Gehrt, & Prange, 2006; Gosselink et al., 2003; Prange, & Gehrt, 2007; Prugh et al., 2009).

Conversely, we expected humans and all carnivore species to be negatively correlated at parks

with high human occupancy, with a weaker negative or non-significant correlation at parks

where human occupancy was low in support of the HCH. If humans are negatively correlated

with species across the community, it would indicate that humans do not simply exert top-down

pressure on the apex carnivore, but that these finer patch-scale effects are felt for other species as

well (George, & Crooks, 2006).

To test for changes in associations between species pairs in response to human activity,

we transformed the Spearman's *R* correlation coefficient to Fisher's Z-statistic and compared

"high" vs "low" groups at α=0.05 significance (Myers, & Sirois, 2014).

Species-Level Response to Human Activity

Finally, we quantified individual species-level response to human occupancy, by

generating activity heatmaps for human, coyote, red fox, skunk, and raccoon occurrence in

Detroit city parks with kernel density estimation (KDE) in ArcMap software (v 10.6.1, ESRI

2010). KDE heatmaps measured the frequency of activity in an area scaled by total activity

(min: 0, max: 1), where 1 indicates the highest activity for a species at a site. We then used the

kernel density values at each park to calculate a Getis-Ord Gi* statistic (Z-score) and test for

areas of significant activity for each species (i.e. "hotspots"). Determining which sites were

hotspots for each carnivore species and comparing them to where hotspots of human activity

occurred enabled a spatially explicit analysis of each individual carnivore species' use of urban

parks.

Given that Detroit is classified as a major metropolitan area and that > 50% of its 309 city

parks are designated as "mini" (i.e. < 3 acres) and embedded within neighborhoods with high

foot traffic, we anticipated that the majority of city parks would be hotspots of high human

activity (U.S. Census Bureau, 2016). With respect to the HSH versus HCH hypothetical

framework, we expected hotspots of human and subordinate species (i.e. red fox, skunk and

raccoon) to occur at the same parks in support of the HSH. Alternatively, if the data supported

the HCH, we expected the parks where human and carnivore hotspots occurred to be distinct.

RESULTS

Our 12,106-trap night survey yielded detections of coyotes (n=220), raccoons (n=1,496),

red (n=88) and grey (n=11) foxes, and striped skunks (n=38) in a highly urbanized landscape

from 2017-2020. The proportion of sites occupied (uncorrected for imperfect detection) varied

by species and year (Table 1). Contrary to our expectations, sites of significant densities of

human detections were localized to a small proportion of parks.

Human Occupancy Estimation

We recorded 1,103 human detections at 25 parks (naïve occupancy 0.64 for combined

2017-2020 data). The top model (lowest ΔAIC and highest *w*) revealed that the number of trap

nights (TN, β=0.01, *p*=0.03), understory vegetation (VEG, β=-0.62, *p*<0.01), camera type

(CAM, β=-0.57, *p*<0.01), and park size (AREA, β=-0.0009, *p*<0.01) best explained human

detection. Housing density within 500 meters (HOUSE, β=0.001, *p*=0.03) best explained human

occupancy (Table 2). Using the top performing model, we found the average occupancy across

all sites was $\varphi=0.68$. Neither the distance to the nearest school nor year covariates appeared in

any of the top candidate models, and thus did not influence human occupancy.

Community-Level Response to Human Activity

Using the threshold value of "high" and "low" from the top human occupancy model, we

found significant clustering at both levels of human presence: (Observed C-Score$_{Low}$: 0.0714,

$p=0.026$, Observed C-Score$_{High}$: 0.396, $p=0.008$). Therefore, the emergent trend was that the

carnivores' spatial structure was aggregated at both high and low human occupancy, supporting

the alternate scenario where habitat-related resources such as vegetation cover, prey availability,

or water drive significant spatial overlap in the limited city green space (Gompper et al., 2016).

Though we found spatial aggregation across levels of human occupancy, the degree of observed

clustering was higher at parks with high human occupancy.

Pairwise Interspecific Level Response to Human Activity

At both levels of human occupancy, all species dyads had non-significant Spearman's

correlation coefficients (Figure 3). We did not find any significant differences between

correlation coefficients going from low to high human occupancy for any species pair, though

several dyads changed the direction of their association in response to human activity. Coyote-

skunk ($R=0.16$, $R=-0.33$), raccoon-fox ($R=0.32$, $R=-0.44$), and skunk-fox ($R=0.48$, $R=-0.02$)

correlated positively at low human Ψ and negatively at high human Ψ.

Species-Level Response to Human Activity

We found distinct areas of high kernel density activity for humans, coyotes, raccoons,

striped skunk, and red foxes across urban parks in Detroit (Figure 4). Of the 24 city parks

sampled, 4 (17.39%) were significant hotspots based on intensity of use by humans. However,

none of the carnivore species' hotspots significantly overlapped with human hotspots, indicating

spatial avoidance of human activity in support of the HCH. Surprising for an urban environment,

humans were the only species in the system to have a "coldspot" (Getis-Ord Gi* z-score < 0)

meaning activity was significantly lower than the expected value. One park contained significant

hotspots for coyote, red fox, and skunk, the only such location with > 2 species hotspots. Two

parks were hotspots for two carnivores pairs: coyote-raccoon and raccoon-red fox.

DISCUSSION

Our study provides necessary insights into carnivore spatial responses to human

occupancy in parks at multiple levels of ecological organization. Common proxies for human

activity derived from landscape-level metrics of urbanization such as housing density, percent

impervious surfaces, or road density may not capture the resolution necessary to determine fine-

scale consequences for wildlife (Tablado, & Jenni, 2017). The inclusion of indices of direct

human activity is therefore needed in future urban ecology studies to disentangle the effects of

humans from the built environment (Nickel et al., 2020). Further, the design of parks and urban

green spaces should include considerations of how human presence shapes animal communities

to adequately balance the needs of people and wildlife.

Support for either the HSH or HCH frameworks depended upon the scale of ecological

complexity. At the community level, carnivores were spatially aggregated and co-occurred

significantly at parks with low human activity and high human activity, indicating that factors

other than humans were driving space-use in the parks. Key resources such as den sites and prey

availability may be intrinsically concentrated in urban green spaces, given that they are

essentially habitat fragments embedded in an urban matrix (Marzluff, 2005). Carnivores in our

study being aggregated, even at parks with significant human activity, highlights the importance

of city parks for both people and wildlife. Measures to both improve habitat quality and reduce

human-wildlife conflict are of paramount importance for urban natural resource managers in

order to ensure both the long-term persistence of wildlife and access to beneficial outdoor

recreation opportunities for city dwellers.

Whether the species pair was apex-subordinate or subordinate-subordinate changed the

interpretation of whether the results supported the HSH or HCH at the pairwise interspecific

correlation level. Coyote-skunk and multiple subordinate-subordinate species pairs changed the

direction of their association in response to human occupancy, though these changes were not

statistically significant. It is unclear whether this is due to low site sample size and limited

statistical power or a case where biological and statistical significance are incongruent

(Martínez-Abraín, 2008). Both coyotes and skunks are particularly effective urban exploiters,

given their generalist dietary breadth and ability to capitalize on anthropogenic food sources

(Gehrt et al., 2011; Murray et al., 2015; Prange, & Gehrt, 2004). Because humans increase

nocturnality in urban wildlife, the change from positive to negative association between coyotes

and skunks in response to human activity may be evidence of spatial avoidance, given their

increased temporal overlap (Gaynor et al., 2018).

Following results from pairwise species correlations, none of the four focal carnivore

species shared hotspots of significant activity with humans at the individual level, lending

support to the HCH. Raccoons, red foxes, and coyotes are emblematic of organisms who are

spatially associated with urbanization at a coarse scale, but our results illustrate that in spite of

being embedded in a highly urbanized cityscape, these species are still finding pockets of spatial

refugia away from people (Prange et al., 2004). Results from the paired interspecific spatial

correlations appear to contradict those at the individual species level, which we believe is a result

of methodological differences in our approach to quantifying each level's response. For paired

species correlations, we categorically binned parks based on high versus low human occupancy,

while the geospatial tool was informed by continuous relative activity data (i.e. the number of

detections). A solution for future studies could be a before-after-controlled-impact (BACI)

approach to experimentally manipulate human activity at each park and thus compare

community aggregation, paired species associations, and individual hotspots at each location in

response to a similarly measured variable of human activity.

Patterns in nature are coupled with the scale at which they are observed, thus scale is a

critical component in ecological studies and we further corroborate this tenet for urban

environments (Levin, 1992; Sayre, 2005). The effect of human activity did not scale with levels

of ecological organization, suggesting that the causal mechanism through which human pressures

illicit responses from wildlife is different for individuals, species pairs, and communities. For

example, reduction of mortality through spatial avoidance of people could determine individual-

level responses to human occupancy (Loveridge et al., 2017; Ordiz et al., 2011). In contrast,

human activity might alter higher order interactions through changes in temporal activity in

dominant carnivores, facilitating spatial overlap at the community-level (Billick, & Case, 1994;

Dröge et al., 2017).

If humans function as shields and facilitate spatial overlap in the carnivore community,

then urban areas could serve as key refugia in cities and increase co-existence in an otherwise

highly competitive guild. Paradoxically, the human shield hypothesis indicates that urbanization

does not inherently result in biodiversity loss at the patch scale, given that subordinate species

can exploit refugia (Lewis et al., 2019; Moll et al., 2018). However, biodiversity loss due to

urbanization at both the landscape and global scale remains a concern for conservation efforts

(Lewis et al., 2015; McDonald et al., 2013; McIntyre, 2014).

In addition to wild carnivores, domestic dogs and cats were commonly detected and often

without human company. Human affiliates such as dogs and cats are widely recognized as

having significant detrimental effects on wildlife (Lenth et al., 2008; Loss et al., 2013; Vanak, &

Gompper, 2009). Despite this, the distinction between free roaming, feral, or simply temporarily

off leash remains unresolved in our study system. As a result, we were unable to determine

whether these human affiliates were true long-term members of the local carnivore assembly and

thus were excluded from the analysis. Leashed dogs sometimes harass and injure wildlife,

including some of the focal species for this study (Hughes, & Macdonald, 2013). However, how

these antagonistic interactions determine the composition, structure, and distribution of carnivore

communities in urban spaces are not well understood. Because dogs are one of the most widely

distributed terrestrial carnivores, filling this knowledge gap should be a key consideration for

future studies to better inform natural resource managers seeking to mitigate their effects on

wildlife (Gehrt et al., 2010).

Our study provides a multilevel analysis of an urban carnivore community in a mid-sized

city whose population has declined over the last 70 years. Notably, this emigration of people

from the city of Detroit is complex and tied to various historical socioeconomic biases. Further,

we recognize that how people are distributed in the city and who has access to green spaces is

not equitable and a consequence of discriminatory housing policies (Watkins, & Gerrish, 2018).

This economic and racial segregation of neighborhoods introduces a bias to understanding

human-wildlife interactions in cities, requiring further study to disentangle species coexistence

patterns, resource availability, and socioeconomic factors.

Examples abound in the literature of generalizations about biodiversity loss and

homogenization of community assemblages based primarily in large cities (Groffman et al.,

2014; Pearse et al., 2018). Homogenization patterns in urban systems do not necessarily scale

down to smaller cities. Moreover, the historical trajectory and relationship between of population

decline, housing vacancy, and vegetation varies by city (Schwarz et al., 2018). We therefore

anticipate that future multi-city comparisons will need to incorporate locations of varying

combinations of urban area and human population density (Magle et al., 2019; Steele, & Wolz,

2019). As the world continues to develop and urbanize, our work informs how carnivore

communities may respond to living in a human-dominated landscape.

Finally, our study could inform the way natural resource managers and city planners

approach urban design. Given that urban carnivores seek spatial refuge from human activity

hotspots, future park designs could incorporate wildlife zones where the use of walking trails

diverts human foot traffic around rather than through important habitat. Urban planners are thus

tasked with promoting access to natural areas for the public, while still conserving habitat for

wildlife. City parks are an important resource for urbanites and provide recreational, cultural,

psychological and physiological benefits to visitors (Soga, & Gaston, 2020). Therefore, finding a

balance between the wellbeing of people and wildlife is a fundamental challenge of the 21[st]

century (Chawla, 2015; Liu et al., 2017; Rigolon, 2016).

ACKNOWLEDGEMENTS

First, we recognize implementing our field research with camera traps was conducted on lands

originally belonging to the People of the Three Fires. Our work is not human subjects research

requiring IRB review, though we remain grateful to authorities granting permission for our

research and their efforts to manage coupled human-natural ecosystems. Our sincere thanks to

members past and present of the Applied Wildlife Ecology (AWE) Lab at the University of

Michigan, specifically K. Mills, R. Malhotra, S. Lima, S. Bower, and G. Gadsden who

contributed to the data collection, image sorting, ArcGIS expertise, and logistical support of this

project. We thank our partners at the Detroit Zoological Society for their financial support and

the City of Detroit for collaboration, permits, and access to the parks in our study. We thank all

of our volunteers for their assistance with camera checks as well as the *Michigan ZoomIN* online

community for their contribution to image classification.

REFERENCES

Alberti, M., et al. (2003). Integrating Humans into Ecology: Opportunities and Challenges for
Studying Urban Ecosystems. *BioScience, 53*(12), 1169-1179. doi:10.1641/0006-
3568(2003)053[1169:IHIEOA]2.0.CO;2
Amarasekare, P. (2003). Competitive coexistence in spatially structured environments: a
synthesis. *Ecology Letters, 6*(12), 1109-1122. doi:10.1046/j.1461-0248.2003.00530.x
Bateman, P. W., et al. (2012). Big city life: carnivores in urban environments. *Journal of
Zoology, 287*(1), 1-23. doi:doi:10.1111/j.1469-7998.2011.00887.x
Beckmann, J. P., et al. (2008). Carnivores, urban landscapes, and longitudinal studies: a case
history of black bears. *Human-Wildlife Conflicts, 2*(2), 168-174.
Berger, J. (2007). Fear, human shields and the redistribution of prey and predators in protected
areas. *Biology Letters, 3*(6), 620-623. doi:10.1098/rsbl.2007.0415
Billick, I., et al. (1994). Higher Order Interactions in Ecological Communities: What Are They
and How Can They be Detected? *Ecology, 75*(6), 1530-1543. doi:10.2307/1939614
Ceballos, G., et al. (2002). Mammal Population Losses and the Extinction Crisis. *Science,
296*(5569), 904. doi:10.1126/science.1069349
Chapron, G., et al. (2016). Coexistence with Large Carnivores Informed by Community Ecology.
Trends in Ecology & Evolution, 31(8), 578-580.
doi:https://doi.org/10.1016/j.tree.2016.06.003

Chawla, L. (2015). Benefits of Nature Contact for Children. *Journal of Planning Literature,*
30(4), 433-452. doi:10.1177/0885412215595441
City of Detroit. (2015). *Detroit Open Data Portal*. Retrieved from: https://data.detroitmi.gov/
Clucas, B., et al. (2011). Coupled Relationships between Humans and other Organisms in Urban
Areas. In J. Niemela (Ed.), *Urban Ecology: Patterns, Processes, and Applications*:
Oxford University Press.
Cove, M. V., et al. (2012). Use of Camera Traps to Examine the Mesopredator Release
Hypothesis in a Fragmented Midwestern Landscape. *The American Midland Naturalist,*
168(2), 456-465. doi:10.1674/0003-0031-168.2.456
de Satgé, J., et al. (2017). Competition and coexistence in a small carnivore guild. *Oecologia,*
184(4), 873-884. doi:10.1007/s00442-017-3916-2
Detroit Residential Parcel Survey. (2010). *Citywide report for vacant and occupied housing*.
Retrieved from http://www.detroitparcelsurvey.org/
Dröge, E., et al. (2017). Spatial and temporal avoidance of risk within a large carnivore guild.
Ecology and Evolution, 7(1), 189-199. doi:10.1002/ece3.2616
Everatt, K. T., et al. (2019). Africa's apex predator, the lion, is limited by interference and
exploitative competition with humans. *Global Ecology and Conservation, 20*, e00758.
doi:https://doi.org/10.1016/j.gecco.2019.e00758
Farris, Z. J., et al. (2017). Threats to a rainforest carnivore community: A multi-year assessment
of occupancy and co-occurrence in Madagascar. *Biological Conservation, 210*, 116-124.
doi:https://doi.org/10.1016/j.biocon.2017.04.010
Fedriani, J. M., et al. (2000). Competition and intraguild predation among three sympatric
carnivores. *Oecologia, 125*(2), 258-270. doi:10.1007/s004420000448
Gallo, T., et al. (2019). Urbanization alters predator-avoidance behaviours. *Journal of Animal*
Ecology, 88(5), 793-803. doi:10.1111/1365-2656.12967
Gaston, K. J., et al. (2017). Impacts of Artificial Light at Night on Biological Timings. *Annual*
Review of Ecology, Evolution, and Systematics, 48(1), 49-68. doi:10.1146/annurev-
ecolsys-110316-022745
Gaynor, K. M., et al. (2018). The influence of human disturbance on wildlife nocturnality.
Science, 360(6394), 1232. doi:10.1126/science.aar7121
Geffroy, B., et al. (2015). How Nature-Based Tourism Might Increase Prey Vulnerability to
Predators. *Trends in Ecology & Evolution, 30*(12), 755-765.
doi:https://doi.org/10.1016/j.tree.2015.09.010
Gehrt, S., et al. (2011). Is the Urban Coyote a Misanthropic Synanthrope? The Case from
Chicago. *Cities and the Environment (CATE), 4*(1).
Gehrt, S. D., et al. (2006). Interference competition between coyotes and raccoons: a test of the
mesopredator release hypothesis. *Behavioral Ecology, 18*(1), 204-214.
doi:10.1093/beheco/arl075
Gehrt, S. D., et al. (Eds.). (2010). *Urban Carnivores: ecology, conflict, and conservation*.
Baltimore, Maryland: The Johns Hopkins University Press.
Gehrt, S. D., et al. (2013). Population Ecology of Free-Roaming Cats and Interference
Competition by Coyotes in Urban Parks. *PLOS ONE, 8*(9), e75718.
doi:10.1371/journal.pone.0075718
George, S. L., et al. (2006). Recreation and large mammal activity in an urban nature reserve.
Biological Conservation, 133(1), 107-117.
doi:https://doi.org/10.1016/j.biocon.2006.05.024

Gompper, M. E., et al. (2016). Differential Habitat Use or Intraguild Interactions: What
Structures a Carnivore Community? *PLOS ONE, 11*(1), e0146055.
doi:10.1371/journal.pone.0146055
Gosselink, T. E., et al. (2003). Temporal Habitat Partitioning and Spatial Use of Coyotes and
Red Foxes in East-Central Illinois. *The Journal of Wildlife Management, 67*(1), 90-103.
doi:10.2307/3803065
Gotelli, N. J., et al. (2015). EcoSimR: Null model analysis for ecological data (Version 0.1.0).
Groffman, P. M., et al. (2014). Ecological homogenization of urban USA. *Frontiers in Ecology*
and the Environment, 12(1), 74-81. doi:10.1890/120374
Hughes, J., et al. (2013). A review of the interactions between free-roaming domestic dogs and
wildlife. *Biological Conservation, 157*, 341-351.
doi:https://doi.org/10.1016/j.biocon.2012.07.005
Lenth, B. E., et al. (2008). The Effects of Dogs on Wildlife Communities. *Natural Areas*
Journal, 28(3), 218-227, 210.
Lepczyk, C. A., et al. (2017). Biodiversity in the City: Fundamental Questions for Understanding
the Ecology of Urban Green Spaces for Biodiversity Conservation. *BioScience, 67*(9),
799-807. doi:10.1093/biosci/bix079
Levin, S. A. (1992). The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur
Award Lecture. *Ecology, 73*(6), 1943-1967. doi:10.2307/1941447
Lewis, A. D., et al. (2019). Does Nature Need Cities? Pollinators Reveal a Role for Cities in
Wildlife Conservation. *Frontiers in Ecology and Evolution, 7*(220).
doi:10.3389/fevo.2019.00220
Lewis, J. S., et al. (2015). Interspecific interactions between wild felids vary across scales and
levels of urbanization. *Ecology and Evolution, 5*(24), 5946-5961. doi:10.1002/ece3.1812
Liu, H., et al. (2017). The relationships between urban parks, residents' physical activity, and
mental health benefits: A case study from Beijing, China. *Journal of Environmental*
Management, 190, 223-230. doi:https://doi.org/10.1016/j.jenvman.2016.12.058
Liu , J., et al. (2007). Complexity of Coupled Human and Natural Systems. *Science, 317*(5844),
1513-1516. doi:10.1126/science.1144004
Loss, S. R., et al. (2013). The impact of free-ranging domestic cats on wildlife of the United
States. *Nature Communications, 4*(1), 1396. doi:10.1038/ncomms2380
Loveridge, A. J., et al. (2017). The landscape of anthropogenic mortality: how African lions
respond to spatial variation in risk. *Journal of Applied Ecology, 54*(3), 815-825.
doi:10.1111/1365-2664.12794
MacKenzie, D. I., et al. (2004). Assessing the fit of site-occupancy models. *Journal of*
Agricultural, Biological, and Environmental Statistics, 9(3), 300-318.
doi:10.1198/108571104X3361
MacKenzie, D. I., et al. (2003). Estimating Site Occupancy, Colonization, and Local Extinction
When a Species is Detected Imperfectly. *Ecology, 84*(8), 2200-2207. doi:doi:10.1890/02-
3090
Mackenzie, D. I., et al. (2005). Designing occupancy studies: general advice and allocating
survey effort. *Journal of Applied Ecology, 42*(6), 1105-1114. doi:doi:10.1111/j.1365-
2664.2005.01098.x
Magle, S. B., et al. (2019). Advancing urban wildlife research through a multi-city collaboration.
Frontiers in Ecology and the Environment, 17(4), 232-239. doi:10.1002/fee.2030

Martínez-Abraín, A. (2008). Statistical significance and biological relevance: A call for a more
cautious interpretation of results in ecology. *Acta Oecologica, 34*(1), 9-11.
doi:https://doi.org/10.1016/j.actao.2008.02.004
Marzluff, J. M. (2005). Island biogeography for an urbanizing world: how extinction and
colonization may determine biological diversity in human-dominated landscapes. *Urban
Ecosystems, 8*(2), 157-177. doi:10.1007/s11252-005-4378-6
McDonald, R. I., et al. (2018). Conservation priorities to protect vertebrate endemics from global
urban expansion. *Biological Conservation, 224*, 290-299.
doi:https://doi.org/10.1016/j.biocon.2018.06.010
McDonald, R. I., et al. (2013). Urbanization and Global Trends in Biodiversity and Ecosystem
Services. In T. Elmqvist, M. Fragkias, J. Goodness, B. Güneralp, P. J. Marcotullio, R. I.
McDonald, S. Parnell, M. Schewenius, M. Sendstad, K. C. Seto, & C. Wilkinson (Eds.),
*Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities: A
Global Assessment* (pp. 31-52). Dordrecht: Springer Netherlands.
McIntyre, N. E. (2014). Wildlife Responses to Urbanization: Patterns of Diversity and
Community Structure in Built Environments. In R. A. McCleery, C. E. Moorman, & M.
N. Peterson (Eds.), *Urban Wildlife conservation: Theory and Practice* (pp. 103-115).
Boston, MA: Springer US.
Moll, R. J., et al. (2018). Humans and urban development mediate the sympatry of competing
carnivores. *Urban Ecosystems, 21*(4), 765-778. doi:10.1007/s11252-018-0758-6
Moll, R. J., et al. (2019). What does urbanization actually mean? A framework for urban metrics
in wildlife research. *Journal of Applied Ecology, 56*(5), 1289-1300. doi:10.1111/1365-
2664.13358
Moss, W. E., et al. (2016). Quantifying risk and resource use for a large carnivore in an
expanding urban–wildland interface. *Journal of Applied Ecology, 53*(2), 371-378.
doi:10.1111/1365-2664.12563
Muhly, T. B., et al. (2011). Human Activity Helps Prey Win the Predator-Prey Space Race.
PLOS ONE, 6(3), e17050. doi:10.1371/journal.pone.0017050
Murray, M., et al. (2015). Greater consumption of protein-poor anthropogenic food by urban
relative to rural coyotes increases diet breadth and potential for human–wildlife conflict.
Ecography, 38(12), 1235-1242. doi:10.1111/ecog.01128
Myers, L., et al. (2014). Spearman Correlation Coefficients, Differences between. *Wiley
StatsRef: Statistics Reference Online*. doi:doi:10.1002/9781118445112.stat02802
Nickel, B. A., et al. (2020). Human presence and human footprint have non-equivalent effects on
wildlife spatiotemporal habitat use. *Biological Conservation, 241*, 108383.
doi:https://doi.org/10.1016/j.biocon.2019.108383
Niedballa, J., Sollmann, R. , Courtiol, A., Wilting, A. (2016). camtrapR: an R package for
efficient camera trap data management (Version 1.1). Methods Ecol Evol.
Ordeñana, M. A., et al. (2010). Effects of urbanization on carnivore species distribution and
richness. *Journal of Mammalogy, 91*(6), 1322-1331. doi:10.1644/09-MAMM-A-312.1
Ordiz, A., et al. (2011). Predators or prey? Spatio-temporal discrimination of human-derived risk
by brown bears. *Oecologia, 166*(1), 59-67. doi:10.1007/s00442-011-1920-5
Pearse, W. D., et al. (2018). Homogenization of plant diversity, composition, and structure in
North American urban yards. *Ecosphere, 9*(2), e02105. doi:10.1002/ecs2.2105
Prange, S., et al. (2004). Changes in mesopredator-community structure in response to
urbanization. *Canadian Journal of Zoology, 82*(11), 1804-1817. doi:10.1139/z04-179

Prange, S., et al. (2007). Response of Skunks to a Simulated Increase in Coyote Activity. *Journal*
of Mammalogy, 88(4), 1040-1049. doi:10.1644/06-MAMM-A-236R1.1
Prange, S., et al. (2004). Influences of Anthropogenic Resources on Raccoon (Procyon lotor)
Movements and Spatial Distribution. *Journal of Mammalogy, 85*(3), 483-490.
doi:10.1644/1383946
Prugh, L. R., et al. (2009). The Rise of the Mesopredator. *BioScience, 59*(9), 779-791.
doi:10.1525/bio.2009.59.9.9
Raleigh, E., et al. (2015). Neighborhood Disinvestment, Abandonment, and Crime Dynamics.
Journal of Urban Affairs, 37(4), 367-396. doi:10.1111/juaf.12102
Rigolon, A. (2016). A complex landscape of inequity in access to urban parks: A literature
review. *Landscape and Urban Planning, 153*, 160-169.
doi:https://doi.org/10.1016/j.landurbplan.2016.05.017
Riley, S. P. D., et al. (2014). Wildlife Friendly Roads: The Impacts of Roads on Wildlife in
Urban Areas and Potential Remedies. In R. A. McCleery, C. E. Moorman, & M. N.
Peterson (Eds.), *Urban Wildlife conservation: Theory and Practice* (pp. 323-360).
Boston, MA: Springer US.
Ripple, W. J., et al. (2014a). Status and Ecological Effects of the World's Largest Carnivores.
Science, 343(6167). doi:10.1126/science.1241484
Ripple, W. J., et al. (2014b). Status and Ecological Effects of the World's Largest Carnivores.
Science, 343(6167), 1241484. doi:10.1126/science.1241484
Sayre, N. F. (2005). Ecological and geographical scale: parallels and potential for integration.
Progress in Human Geography, 29(3), 276-290. doi:10.1191/0309132505ph546oa
Schuette, P., et al. (2013). Occupancy patterns and niche partitioning within a diverse carnivore
community exposed to anthropogenic pressures. *Biological Conservation, 158*, 301-312.
doi:https://doi.org/10.1016/j.biocon.2012.08.008
Schwarz, K., et al. (2018). Green, but not just? Rethinking environmental justice indicators in
shrinking cities. *Sustainable Cities and Society, 41*, 816-821.
doi:https://doi.org/10.1016/j.scs.2018.06.026
Shannon, G., et al. (2014). Behavioral Responses Associated with a Human-Mediated Predator
Shelter. *PLOS ONE, 9*(4), e94630. doi:10.1371/journal.pone.0094630
Smith, J. A., et al. (2018). Human activity reduces niche partitioning among three widespread
mesocarnivores. *Oikos, 127*(6), 890-901. doi:doi:10.1111/oik.04592
Soga, M., et al. (2020). The ecology of human–nature interactions. *Proceedings of the Royal*
Society B: Biological Sciences, 287(1918), 20191882. doi:10.1098/rspb.2019.1882
Steele, M. K., et al. (2019). Heterogeneity in the land cover composition and configuration of US
cities: implications for ecosystem services. *Landscape Ecology, 34*(6), 1247-1261.
doi:10.1007/s10980-019-00859-y
Stone, L., et al. (1990). The checkerboard score and species distributions. *Oecologia, 85*(1), 74-
79. doi:10.1007/BF00317345
Tablado, Z., et al. (2017). Determinants of uncertainty in wildlife responses to human
disturbance. *Biological Reviews, 92*(1), 216-233. doi:10.1111/brv.12224
Terborgh, J. E., J. A. (Ed.) (2010). *Trophic Cascades: Predators, Prey, and the Changing*
Dynamics of Nature: Island Press.
U.S. Census Bureau. (2016). Decennial Census of Population and Housing.

United Nations. (2018). *World Urbanization Prospects: The 2019 Revision*. Retrieved from
https://www.un.org/en/events/citiesday/assets/pdf/the_worlds_cities_in_2018_data_bookl
et.pdf
Vanak, A. T., et al. (2009). Dogs Canis familiaris as carnivores: their role and function in
intraguild competition. *Mammal Review, 39*(4), 265-283. doi:10.1111/j.1365-
2907.2009.00148.x
Watkins, S. L., et al. (2018). The relationship between urban forests and race: A meta-analysis.
Journal of Environmental Management, 209, 152-168.
doi:10.1016/j.jenvman.2017.12.021
Wu, J. (2014). Urban ecology and sustainability: The state-of-the-science and future directions.
Landscape and Urban Planning, 125, 209-221.
doi:https://doi.org/10.1016/j.landurbplan.2014.01.018

676   **Table 1** Summary of 2017-2020 DMP camera survey including trap nights, number of sampling
677   sites, detections for all species, and number and proportion of sites occupied at parks with low
678   versus high human occupancy.

	Low Human Ψ	High Human Ψ	Total
Trap Nights	5,572	6,534	12,106
Sites	8	18	26
Summary of DMP 2017-2020 Camera Detections			
Human	99	1,004	1,103
Coyote	122	98	220
Red Fox	72	16	88
Grey Fox	9	2	11
Raccoon	1,044	452	1,496
Skunk	19	19	38
Domestic Dog	245	355	600
Domestic Cat	139	300	439
Number (Proportion) Sites Occupied			
Human	7 (0.86)	18 (1)	25 (0.96)
Coyote	7 (0.86)	9 (0.50)	16 (0.62)
Red Fox	3 (0.38)	4 (0.22)	7 (0.27)
Grey Fox	1 (0.13)	1 (0.06)	2 (0.08)
Raccoon	7 (0.86)	12 (0.67)	19 (0.73)
Skunk	3 (0.38)	3 (0.17)	6 (0.23)
Domestic Dog	8 (1)	16 (0.89)	24 (0.92)
Domestic Cat	7 (0.86)	16 (0.89)	23 (0.88)

698   **Table 2** Summary of candidate human occupancy models with covariates including trap nights
699   (TN), distance to nearest school (DSCH), understory vegetation at camera (VEG), park size
700   (AREA), and year (YR). Dot models (.) denote null (i.e. random) occupancy or detection.
701   Akaike's Information Criterion (AIC), delta AIC, AIC model weight, model goodness of fit (X^2),
702   and overdispersion ($\hat{c}$) are listed.

Model	AIC	ΔAIC	AIC Weight	Goodness of Fit (X^2)	$\hat{c}$
ρ(TN+AREA+VEG +CAM) Ψ (HOUSE)	1509.94	0.00	0.351	$p=0.756$	0.985
ρ(TN+AREA+VEG+CAM) Ψ (AREA)	1511.18	1.24	0.189	$p=0.451$	1.000
ρ(TN+AREA+VEG+CAM) Ψ (HOUSE + AREA)	1511.27	1.33	0.181	$p=0.215$	1.015
ρ(TN+AREA+VEG+ CAM) Ψ (.)	1511.36	1.42	0.173	$p=0.474$	1.002
ρ (AREA+VEG+ CAM) Ψ (HOUSE)	1513.23	3.28	0.068	$p=0.085$	0.975
ρ (AREA+VEG+ CAM) Ψ (AREA)	1514.33	4.38	0.039	$p=0.089$	1.009

Figure Legends

Figure 1. Conceptual framework for the effects of humans on carnivores at community (1),
pairwise interspecific interactions (2), and species level (3) under two hypotheses: humans as
shields (HSH) and humans as competitors (HCH). HSH1: Humans exert top-down pressure on
dominant carnivore, facilitating spatial aggregation at high human occupancy (Ψ). HSH2:
Positive association between subordinate carnivores at high human Ψ, negative association at
low human Ψ, while apex-subordinate pairs negative at both. HSH3: Overlap between human
and subordinate carnivore hotspots. HCH1: Increased activity from human competitors reduces
available niche space for all carnivore species, community is spatially segregated at high human
Ψ, aggregated at low human Ψ. HCH2: Negative association between subordinate carnivores at
high human Ψ and positive association at low human Ψ, while apex-subordinate pairs negative at
both. HCH3: No overlap between human and subordinate carnivore hotspots.

Figure 2. Study area – City of Detroit, Michigan. Shaded polygons represent the 24 city parks in
the Detroit Metro Parks system included in the analysis while black dots denote camera stations
in the study.

Figure 3. Correlogram of pairwise interspecific associations using Spearman's R at: A) low
human occupancy versus B) high human occupancy. Values in white backgrounds denote non-
significant correlation at 95% confidence while values in color tiles denote $p<0.05$. Asterisk
indicates a significant change in Spearman's R from low to high human occupancy.

Figure 4. Hotspots (Getis-Ord Gi* at 95% confidence) of significant activity of carnivore
species and humans in Detroit Metro Parks from camera survey, 2017-2020. Hotspots are
denoted by unique symbols for each species and/or combination.

Humans as Shields Hypothesis (HSH)

Humans as Competitors Hypothesis (HCH)

1. Community

2. Pairwise

3. Species

 Apex

 Subordinate

 ✓ Supported by Data

 High Human Ψ / Low Human Ψ

Detroit Camera Survey 2017-2020

- Camera Station
- City Parks
- City of Detroit Boundary

0 2.25 4.5 9 13.5 18
Kilometers

Species Hotspots
(Getis Ord-Gi*)

⬟ Human 👫

◉ Skunk 🦨

▼ Coyote 🦊

● Red Fox, Coyote 🦊🦊

▲ Raccoon, Coyote 🦝🦊

★ Raccoon, Red Fox, Skunk 🦝🦊🦨