MethylationToActivity: a deep-learning framework that reveals

promoter activity landscapes from DNA methylomes in individual

tumors

Justin Williams[1], Beisi Xu[2], Daniel Putnam[1], Andrew Thrasher[1], Chunliang Li[3], Jun

Yang[4] and Xiang Chen[1]

[1]Department of Computational Biology, St. Jude Children's Research Hospital, Memphis,

Tennessee, United States

[2]Center for Applied Bioinformatics, St. Jude Children's Research Hospital, Memphis, Tennessee,

United States

[3]Department of Tumor Cell Biology, St. Jude Children's Research Hospital, Memphis,

Tennessee, United States

[4]Department of Surgery, St. Jude Children's Research Hospital, Memphis, Tennessee, United

States

Corresponding author: Xiang Chen

Department of Computational Biology

St. Jude Children's Research Hospital

262 Danny Thomas Place

Mail Stop 1135

Memphis, TN 38105

U.S.A.

Tel: +1 901 595 7074

Fax: +1 901 595 7100

Email: xiang.chen@stjude.org

Abstract

Although genome-wide DNA methylomes have demonstrated their clinical value as reliable

biomarkers for tumor detection, subtyping, and classification, their direct biological impacts at

the individual gene level remain elusive. Here we present MethylationToActivity (M2A), a

machine learning framework that uses convolutional neural networks to infer promoter activities

(H3K4me3 and H3K27ac enrichment) from DNA methylation patterns for individual genes.

Using publicly available datasets in real-world test scenarios, we demonstrate that M2A is highly

accurate and robust in revealing promoter activity landscapes in various pediatric and adult

cancers, including both solid and hematologic malignant neoplasms.

Keywords:

40 DNA methylation, histone modifications, convolutional neural network, transfer learning

Background

Transcriptional regulation is fundamental to the identity and function of cells. Deregulation of

gene expression is a defining feature of common diseases, including cancers. Promoters, the

regulatory regions surrounding the transcription starting sites (TSSs), integrate signals from

distal enhancers and local histone modifications (HMs) to initiate transcription. Almost half of

human protein-coding genes harbor multiple TSSs; consequently, promoter activities determine

both the level of transcription and the transcript isoforms that are expressed, with the latter

potentially having different translation efficiencies and encoding different protein sequences [1].

Tumors frequently use alternative promoters to increase the isoform diversity [2, 3], to activate

oncogenes that are normally repressed [1-3], and to evade host immune attacks by

immunoediting [3, 4]. Compared to cancers in adults, pediatric tumors harbor fewer mutations [5,

6] and use epigenetic deregulation to promote tumorigenesis and progression [7].

Promoter activities can be determined experimentally through transcriptomic approaches,

such as CAP analysis gene expression (CAGE), or through epigenomic approaches, including

chromatin immunoprecipitation followed by sequencing (ChIP-seq) [8]. Because of transcript

degradation by 5′ RNA exonucleases, ChIP-seq approaches for specific HMs have been the

gold standard for studying promoter activities [9]. Several studies [10-12] have demonstrated

that HMs and other epigenetic features can be used to predict gene expression. Using a linear

regression model, Karlić et al. show that approximately 50%–60% of the variation in gene

expression can be accounted for, and that ~50% of the variation in gene expression can be

modeled by promoter H3K27ac enrichment alone [10]. Subsequent work by Dong et al. further

explained 69% of gene expression variance using a hybrid random forest/linear regression

model with features derived from 11 HMs, one histone variant, and DNase I hypersensitivity

[11]. More recently, Singh et al. used deep-learning models on five HMs to predict gene

expression status (high/low) and achieved an average AUC of 0.80 [12]. However, the scarcity

of pediatric tumors, the limited amounts of fresh starting material available, and the

extensive workload involved in acquiring the promoter activity landscapes constrain their

interrogation for individual patient tumors [13, 14].

DNA methylation (DNAm) is a well-studied, relatively stable, and inheritable epigenetic

regulatory mechanism that involves transferring a methyl group to cytosine (C) to form 5-

methylcytosine (5mC), mostly in the CpG context. In contrast to HMs, DNAm can be accurately

and robustly profiled in various tissues, including archival formalin-fixed, paraffin-embedded

(FFPE) tumor samples, through both array [15, 16] and sequencing [17] platforms; therefore, it

has exceptional applicability to studying epigenetic deregulation in tumors. Consequently,

genome-wide DNAm profiles represent a widely available epigenetic asset for studying

epigenetic abnormalities in primary tumors.

The DNAm pattern is mechanistically connected with transcription factor binding and HMs

[18-25]. It also plays critical roles in establishing the chromatin structure in physiologic and

pathologic conditions [26, 27]. Moreover, recent applications of machine learning to genome-

wide DNAm patterns have demonstrated that DNAm can accurately predict the patterns of

chromatin packaging (A/B compartments, the square of the Pearson correlation coefficient R^2 =

0.50–0.66) [28-30] and can reveal distinct subgroups with prognostic significance among

patients with cancer [31, 32]. Recently, DNAm signature–based molecular classifiers were

shown to improve diagnostic accuracy, as compared to that of traditional schemes, further

demonstrating the critical regulatory roles of DNAm in tumor development [33, 34]. However,

unlike HMs, where established biological interpretations of various marks have resulted in a

general "histone code" hypothesis [35, 36], the relation between DNAm signatures and their

transcriptional regulatory roles is complex and nonlinear. In many cases, even promoter DNAm

may positively and negatively correlate with gene expression depending on the genomic

structure involved in a given tumor [37]. Consequently, with few exceptions (e.g.,

hypermethylation of the promoters of *RB1*, *CDKN2A*, and *MGMT*) [38], the contribution of

DNAm to the regulation of expression of individual genes remains largely elusive [39-41].

Recent attempts to use DNAm signatures to account for gene expression levels have had

limited success, with the best model (binomial distribution probit regression [BPR] model)

capturing 25%–49% of the expression variations [42]. Undoubtedly, the lack of interpretability

of the DNAm pattern at the individual gene level has severely hampered our understanding of

the biological significance of DNAm signatures.

To address these challenges, we have developed MethylationToActivity (M2A), a deep-

learning framework. The central hypothesis of M2A is that the complex relation between DNAm

signatures and promoter activities (measured as H3K4me3 and H3K27ac enrichment in the

TSS ± 1 kb region) can be captured by incorporating both summary statistics extracted from

window-based CpG methylation levels and high-order spatial information from these windows in

the promoter and flanking regions (up to 25 kb from the TSS). Using a cohort of six pediatric

neuroblastoma (NBL) orthoptic patient-derived xenograft (O-PDX) samples profiled in the

Pediatric Cancer Genome Project (PCGP) [43], we trained the model using whole-genome

bisulfite sequencing (WGBS) data to predict the enrichment of H3K4me3 and H3K27ac (two

HMs critical for promoter [10]) for genome-wide annotated promoters. We validated the

predictive accuracy of the model in the remaining NBL samples (N = 10, WGBS) from the

same cohort. We further confirmed its accuracy and generalizability in diverse tumor types from

four publicly available datasets representing real-world applications, including (1) pediatric

rhabdomyosarcoma (RMS) O-PDX tumors profiled in the Pediatric Cancer Genome Project

(N = 16, WGBS) [44]; (2) a set of commonly used cell lines profiled in ENCODE (N = 9,

WGBS) [45]; (3) primary acute myeloid leukemias (AMLs) profiled by the BLUEPRINT

consortium (N = 19, WGBS) [46]; and (4) a large primary Ewing sarcoma (EWS) cohort using

reduced representation bisulfite sequencing (RRBS) (N = 140) [22]. These

applications demonstrate that M2A can accurately reveal promoter activities from DNAm

patterns, which will be of great use not only in functionally interpreting differential DNAm

patterns but also in profiling promoter usage in individual patient tumors. This will facilitate

precision medicine by tailoring treatments based on both genetic variants and epigenetic

deregulations.

Results

Extensive diversity of promoter activity among *MYCN*-amplified NBL cell lines and O-PDX models

To date, most cancer HM profiling studies have made use of tumor models, including cell

lines, xenografts, and more recently, organoids. Technical limitations and challenges when

working with human tumor tissues prevent the generation of high-quality ChIP-seq profiles for

primary patient specimens [47]. Despite the documented epigenetic heterogeneity [48], a

common practice in deciphering major HM deregulations in various cancers is to extrapolate the

epigenetic profiles from related cancer models (surrogate models). Many studies [43, 44, 49-52]

have compared model systems to primary tumors with respect to characteristics such as

mutations, gene expression, and DNAm signatures. In this study, we began by evaluating the

level of promoter activity diversity in closely related NBL models. Specifically, we evaluated

promoter activity, as measured by the H3K27ac level, in three O-PDX models (SJNBL046,

SJNBL108, and SJNBL013763) and three cell line models (IMR-32, NB-5, and SKNBE2) that

harbor *MYCN* amplification with no other major oncogenic mutations. All samples displayed a

bimodal distribution of promoter H3K27ac levels across the genome (Additional file 1: Figure

S1), and O-PDX models had a marginally higher fraction of active promoters (mean: 31.9%,

range: 27.6%–36.1%) than did cell line models (mean: 26.1%, range: 25.6%–26.7%)

($P = 0.14$, Student's *t*-test). However, there were extensive variations in the promoter

activities in both the cell line models (Figures 1a and 1d) and the O-PDX models (Figures 1b

and 1e). Moreover, greater divergence was observed between a cell line model and an O-PDX

model (mean: 34.9%, range: 29.9%–39.0%) than between two cell line models (mean: 31.0%,

range: 29.2%–32.1%; $P = 0.44$, Student's t-test) or between two O-PDX models (mean:

31.0%, range: 22.9%–37.0%; $P = 0.02$, Student's t-test) (Figure 1f). Variations in promoter

activity may play a significant role in the transcriptional deregulation of individual tumors, as a

substantial fraction of established cancer consensus genes (22.4% in O-PDX models and 31.1%

in cell line models, including *APOBEC3B*, *TGFBR2*, *PAX7*, *HOXA11*, *PDCD1LG2*, *PTK6*,

BCL11B, *FAS*, and *MYC*; (Additional file 2: Table S1) displayed heterogeneous promoter

activities in the surveyed tumor models. Therefore, we sought to develop a computational

approach to infer the promoter activity landscape for individual tumors.

M2A: a deep-learning framework to reveal promoter activities from DNA

methylation

DNAm plays a critical role in determining the framework of gene expression for a given

cell/cellular state. However, the highly complex and non-linear relations between DNAm

patterns and HMs severely hamper the interpretability of the biological impact of differential

DNAm patterns. Previous studies have shown the usefulness of extracting higher-order

methylation features [42], for predicting gene expression. Moreover, recent studies applied

deep-learning approaches to infer DNAm states from their local sequence composition and

adjacent DNAm states [53]. We hypothesize that these high-level DNAm features (that capture

the spatial information from DNAm patterns in the promoter and regions in its vicinity) could also

provide an opportunity to infer promoter activities such as H3K27ac and H3K4me3 enrichment

accurately. We propose to use a convolutional neural network (CNN)–based deep-learning

framework to extract such features.

The M2A conceptual framework and workflow is shown in Figure 2. M2A starts with raw

DNAm feature extraction from around individual TSSs (Figure 2a). This is followed by high-level

feature extraction through the CNN layers and mapping between the generalized feature and

the final output (i.e., the H3K4me3 and H3K27ac of the promoter) in the fully connected (FC)

layers. The vanilla model described in this report was trained on six NBL PDX tumors

(SJNBL046_X, SJNBL013761_X1, SJNBL012401_X1, SJNBL013762_X1, SJNBL013763_X1,

and SJNBL015724_X1; Figure 2b) for which comprehensive genomic and epigenomic profiling

data are available, including the results of whole-genome sequencing, whole-exome sequencing,

RNA sequencing, WGBS, and ChIP-seq of eight histone marks (H3K4me1, H3K4me2,

H3K4me3, H3K27me3, H3K27ac, H3K36me3, H3K9/14ac, and H3K9me3), CTCF, BRD4, and

RNA polymerase II (PolII).

We started with an analysis of the information content in DNAm patterns by examining the input

feature distribution in different windows, among active (high H3K27ac), poised (high H3K4me3

and low H3K27ac), and inactive promoters (low H3K4me3 and low H3K27ac) in the six NBL O-

PDX training samples. These features show distinct patterns among the three promoter

categories (Additional File 1: Figure S2), indicating the feasibility of modeling promoter activities

from DNAm patterns. Although the interpretability of CNN extracted features remains an active

field of research in deep-learning [54], we examined the efficacy of CNN extracted features in

modeling the promoter activities. We first compared the square of Pearson's correlation (R^2)

between each feature (both raw input and CNN extracted features) and the response variable

(H3K27ac) in the training set, The analysis revealed that CNN-extracted features have

significantly higher R^2 with the response (250 bp: $P = 1.5 \times 10^{-11}$, 2500 bp $P = 3.9 \times 10^{-5}$,

Wilcoxon signed-rank test, Additional File 1: Figure S3). We further evaluated the best features

for both raw input and CNN-extracted features in the validation samples and again the CNN-

extracted features significantly outperformed the raw input features (250 bp: $P = 1.1 \times 10^{-5}$,

2500 bp $P = 1.1 \times 10^{-5}$, Wilcoxon signed-rank test, Additional File 1: Figure S3).

M2A produces a highly accurate landscape of promoter activity in pediatric NBL

To evaluate the performance of M2A, we first explored its performance in the remaining NBL

samples in the cohort (the validation set), including one O-PDX tumor, one primary autopsy

tumor, and eight cell lines. Using the validation set, we compared the performance of the M2A

framework of three CNN layers and two FC layers (Figure 2) with three frequently used

statistical and machine learning approaches (baseline models), namely multivariate adaptive

regression splines (MARS), random forest, and artificial neural network (ANN) consisting of only

two FC layers. In every instance, the M2A framework outperformed baseline models (Additional

file 2: Table S2). From a qualitative perspective, M2A correctly revealed the bimodal distribution

of the promoter activities for both H3K4me3 and H3K27ac in all samples, and from a

quantitative perspective, the inferred genome-wide promoter activity landscape was

highly accurate for individual samples for both H3K4me3 ($R^2 = 0.933 \pm 0.019$; RMSE =

0.621 ± 0.072) (Figures 3a and 3d) and H3K27ac ($R^2 = 0.799 \pm 0.053$; RMSE =

0.644 ± 0.074) (Figures 3b, and 3c). Moreover, the addition of CNN layers was merited, as

there was a decrease in the prediction error (measured as $1 - R^2$) from the next highest

performer by 17.8% ($P = 0.0020$, Wilcoxon signed-rank test) and 12.4% ($P = 0.0020$,

Wilcoxon signed-rank test) for the model topologies for H3K4me3 and H3K27ac, respectively

(Additional file 2: Table S2).

Our analysis of *MYCN*-amplified NBL cell line and O-PDX models has revealed substantial

variations in their promoter activities, which is a potential caveat to the practice of surrogate

model (representing primary tumor epigenomes by a few profiled models). Conversely,

M2A produced highly accurate promoter activity landscapes, significantly outperforming the

observed consistency between training and testing samples for both H3K4me3 ($R^2 =$

0.891 ± 0.023, $P = 2.3 \times 10^{-5}$, Wilcoxon rank-sum test) and H3K27ac (R^2 =

0.720 ± 0.045, $P = 9.5 \times 10^{-5}$, Wilcoxon rank-sum test; Additional file 2: Table S3).

Remarkably, in nine (of 10) test samples, the accuracy of the M2A-inferred promoter H3K27ac

activity was better than the highest similarity attained by any individual training sample

($P = 0.027$, Wilcoxon signed-rank test). The same pattern was observed for H3K4me3 levels,

with M2A being more accurate for nine of 10 samples ($P = 0.037$, Wilcoxon signed-rank test),

demonstrating the accuracy of M2A in revealing individual tumor promoter activity landscapes.

Finally, the predictive accuracy of M2A was comparable to the experimental consistency

observed between replicates from the same cell lines profiled in ENCODE for H3K4me3 (R^2

$= 0.933 ± 0.018$ for M2A vs. $0.922 ± 0.056$ for ENCODE replicates [$N = 25$]; $P = 0.55$,

Wilcoxon rank-sum test) (Figure 3a; Additional file 2: Table S4). The accuracy of M2A also

approached the replicate consistency for H3K27ac ($R^2 = 0.799 ± 0.050$ for M2A vs.

$0.849 ± 0.047$ for ENCODE replicates [$N = 26$]; $P = 0.0078$, Wilcoxon rank-sum test)

(Figure 3b: Additional file 2: Table S4). Measurement of the root mean square error (RMSE)

revealed a similar pattern (Additional file 2: Table S4).

M2A is generalizable and scalable

Aside from the model accuracy, there are two additional requirements with practical importance

for deploying a machine learning model (such as M2A) in real-world applications: (1)

generalizability, i.e., M2A needs to achieve a similar performance with a set of unseen test

samples, including tumor/tissue types not used in the model training; and (2)

scalability, i.e., M2A must be able to be applied efficiently to external data.

We first demonstrated the accuracy, generalizability, and scalability of M2A by

using test samples from rhabdomyosarcoma (RMS) O-PDX tumors. The RMS O-PDX dataset

consists of 16 pediatric RMS tumors (11 embryonal, four alveolar, and one spindle subtype,

termed ERMS, ARMS, and spindle subtypes, respectively). As with the NBL cohort, each RMS

sample was extensively profiled, including by WGBS, RNA-seq, and ChIP-seq of H3K4me3 and

H3K27ac. Using the vanilla M2A model (the 3CNN-FC model trained on the six NBL PDX

samples), M2A achieved an overall predictive accuracy with the RMS dataset that was

comparable to that of the NBL test group for both H3K4me3 ($R^2 = 0.937 \pm 0.017$, $P = 0.30$;

RMSE $= 0.639 \pm 0.119$, $P = 0.82$, Wilcoxon rank-sum test) (Figures 3a and 3f; Additional file 2:

Table S5) and H3K27ac ($R^2 = 0.790 \pm 0.037$, $P = 0.44$; RMSE $= 0.589 \pm 0.084$, $P = 0.058$,

Wilcoxon rank-sum test) (Figures 3b and 3e; Additional file 2: Table S5), which was comparable

to or significantly outperformed the observed similarities between two different RMS tumors for

H3K4me3 ($R^2 = 0.917 \pm 0.028$, $P = 0.0020$; RMSE $= 0.646 \pm 0.133$, $P = 0.64$, Wilcoxon

rank-sum test) and H3K27ac ($R^2 = 0.780 \pm 0.066$, $P = 0.43$; RMSE $= 0.550 \pm 0.095$, $P =$

0.14, Wilcoxon rank-sum test). The accuracy of the inferred H3K4me3 activity was comparable

to the inter-replicate consistency of the ENCODE samples ($P = 0.83$, Wilcoxon rank-sum test).

By definition, generalizability can be achieved only in the absence of over-fitting (or

"memorization" of the training data). Neural networks often fall victim to this problem through a

combination of factors, including relatively small training datasets and/or over-parameterization.

The relatively consistent expression of housekeeping genes across different tissues may lead to

an inaccurate (often inflated) interpretation of the performance measurement in such a model,

as evidenced by the relatively high R^2 value (0.663 ± 0.040) between the promoter H3K27ac

level of a random RMS test tumor and the most similar NBL training tumor (Additional file 1:

Figure S4a). Therefore, we focused on the set of genes that are differentially expressed (DE) in

RMS and NBL PDX samples [51], for which an over-fitted or memorized model would perform

poorly. Not surprisingly, the average correlative consistency between the NBL validation

samples and the most similar NBL training sample dropped from 0.755 to 0.599 when the

measurement was restricted to promoters encoding the DE genes (Additional file 1: Figure S4b),

whereas a sharp decline (from 0.663 to 0.259) was also observed for RMS test tumors

(Additional file 1: Figure S4b). Conversely, the six-PDX NBL-trained M2A model maintained

high accuracy for promoters of DE genes in both the NBL validation set (R^2 = 0.729 ± 0.071

and the RMS test set (R^2 = 0.715 ± 0.044) (Additional file 1: Figure S4b), further

demonstrating the generalizability of M2A.

M2A is efficient and scalable. For a local implementation of M2A (source code, built models

and a Docker image available at https://github.com/chenlab-sj/M2A), the training of the vanilla

M2A model (with six NBL O-PDX tumors) takes approximately 16 min (using a Tesla P100-

16GB GPU). The feature extraction and promoter activity prediction from WGBS data (as a

genome-wide DNAm level file in a tab-delimited text format) can be executed on a personal

computer (in this case, we used a MacBook Air with a 2.2-GHz Intel Core i7 and 8-GB 1600-

MHz DDR3 RAM) and takes 15–19 min. Moreover, we have implemented a cloud version of

M2A (https://platform.stjude.cloud/workflows/methylation-to-activity), available to the general

research community.

Transfer learning further improves the performance of M2A with minimal

additional input in the target domain.

Although we have demonstrated the generalizability of M2A in the RMS dataset, the fact that

epigenetic genes are frequently mutated in pediatric tumors [7] raises the possibility that

individual tumor types carry a type-specific interpretation of the DNAm patterns. When ChIP-seq

measurement is available for sufficient samples, a type-specific model is desirable. However,

although pediatric solid tumors as a group constitute a rare disease, they comprise many

different tumor types, and it is rare to have sufficiently profiled samples available for many of

them. In addressing this challenge, we hypothesize that a fixed feature-extraction strategy

(transfer learning) can achieve the goal of deriving an efficient tumor type–specific model by

using a small labeled dataset. A primary assumption here is that generalized features extracted

based on a large dataset are similarly informative for apparently different tasks. The feature

learning and selection characteristics of CNNs provide exceptional portability in various tasks

with extremely small labeled datasets.

In M2A, the CNN layers capture generalized DNAm features and the FC layers learn the

mapping function between the DNAm features and the promoter activities. Here we start with

the pretrained vanilla M2A model, fix the feature-extraction layers (CNN layers), and use a

single sample from the target tumor type to update the mapping function (the weights and

biases of the FC layers). Because the consistency of M2A for H3K4me3 approached the inter-

replicate consistency in both NBL and RMS datasets, we focused on H3K27ac inference for

transfer learning. Upon performing transfer learning with a single sample in the RMS dataset,

we observed significantly improved accuracy (R^2 = 0.813 ± 0.038, P = 3.1 × 10^{-5},

Wilcoxon signed-rank test) (Figures 3b, 3g, and 3h; Additional file 2: Table S5). Moreover, this

model significantly outperformed a single RMS sample model with the identical model

architecture, in which both the CNN layer and the FC layers were derived from the RMS training

sample (P = 9.2 × 10^{-5}, Wilcoxon signed-rank test) (Additional file 1: Figure S5; Additional

file 2: Table S6) and marginally outperformed the observed similarities between different RMS

tumors (P = 0.053, Wilcoxon rank-sum test). This analysis demonstrated the value of both the

pretrained CNN layers for general feature extraction and a single profiled sample in the target

domain. Consequently, we applied transfer learning to both the EWS and AML datasets.

However, transfer was not feasible in the ENCODE dataset because those cell lines were

derived from different tissues.

314   **M2A accurately reveals promoter activity landscapes in adult tumors and in**

315   **hematologic malignant neoplasms**

316   We next evaluated the performance of M2A in independently collected datasets, including ones

317   for adult tumors and hematologic malignant neoplasms. Upon analyzing nine ENCODE cell

318   lines, we found that differences in antibody usage and experimental protocols between the

319   ENCODE and NBL datasets resulted in different signal-to-noise profiles (Additional file 1: Figure

320   S6), with higher RMSE values between the model predictions and the actual observations

321   (H3K4me3: 0.961 ± 0.238, P = .0019; H3K27ac: 0.918 ± 0.188, P = 2.6 × 10^{-4},

322   Wilcoxon rank-sum test). However, the predicted activities remained highly correlated with the

323   experimental measurements (H3K4me3: R^2 = 0.895 ± 0.027; H3K27ac: R^2 = 0.680 ± 0.149)

324   (Additional file 1: Figure S7). Although the accuracy of H3K4me3 was relatively uniform, H1-

325   ESC and SK-N-SH were outliers with a substantially lower accuracy of H3K27ac inference (by

326   the boxplot-based method [55]) (Additional file 2: Table S4; Additional file 1; Figure S8a). An

327   investigation of the promoter activity (the measured and inferred H3K27ac levels) and the

328   measured gene expression in H1-ESC revealed that a subset of actively transcribed genes

329   showed little or no H3K27ac levels in their promoters, where M2A inferred relatively strong

330   promoter activity (Additional file 1: Figure S8b). Consequently, the inference of promoter activity

331   by M2A outperformed the actual measurement in terms of both the quantitative consistency with

332   the gene expression level (R^2 = 0.536 for the M2A-inferred H3K27ac level vs. 0.435 for the

333   measured H3K27ac level) (Additional file1: Figures S8b and S8c) and the accuracy in predicting

334   expressed genes (AUC = 0.891 for the M2A-inferred H3K27ac level and 0.881 for observed

335   H3K27ac level) (Additional file 1: Figure S9d). We also observed a small fraction of inferred

336   active promoters without strong expression; these may represent genes subject to

337   transcriptional pausing (where transcription is initiated but there is no elongation), a distinctive

feature of undifferentiated stem cells [56]. Similarly, better consistency with gene expression

levels was observed in SK-N-SH (Additional file 2: Table S7).

We further evaluated the performance of M2A in revealing promoter activities in 19 acute

myeloid leukemia (AML) primary patient samples collected by the BLUEPRINT consortium,

which is part of the International Human Epigenome Consortium (IHEC) [46]. Analyses of

observed promoter activity (measured by H3K27ac level) and gene expression (measured in

fragments per kilobase of transcript per million mapped reads [FPKM]) in the same sample

revealed non-uniform qualities with a wide range of consistency (mean R^2 = 0.539, range:

0.178–0.720) (Additional file 2: Table S8). Similarly, the M2A-inferred promoter activity

landscape displayed substantial variability with respect to the observed activities among these

samples (mean R^2 = 0.473, range: 0.031–0.729). Strikingly, the consistency between the

observed promoter activity and gene expression was highly predictive of the performance of

M2A with individual samples (R^2 = 0.975, P = 4.1 $\times$ 10^{-15}, Pearson's correlation test)

(Additional file 1: Figure S10). Finally, although gene expression was not used in model

generation with M2A (for the vanilla model or the transfer learning step), the promoter activity

inferred by M2A showed uniform consistency with gene expression (mean R^2 = 0.628, range:

0.541–0.684) (Additional file 2: Table S8).These results jointly suggest that the ChIP-seq library

quality is a potential confounding factor for both the consistency between the observed (from

ChIP-seq) and the M2A-inferred promoter activity landscapes and the consistency between the

observed promoter activity and gene expression in these samples.

Promoter activity landscape inferred by M2A faithfully recapitulates the subtype

difference between embryonal and alveolar rhabdomyosarcomas

Identifying recurrent epigenetic deregulations (epi-drivers) is a primary research focus in cancer

epigenome studies [57]. To this end, we investigated whether subtype-specific epigenetic

deregulation was captured in M2A-revealed promoter landscapes in the RMS O-PDX tumors. A

t-SNE embedding using M2A-inferred promoter activity landscapes (from the NBL-trained model)

recapitulated the clear separation of ARMS and ERMS tumors (Figure 4a) in the DNAm profiles

(data not shown), which further demonstrates the generalizability of the CNN-extracted high-

order DNAm features. Importantly, when focusing on the promoters of DE genes in the ARMS

and ERMS subtypes, the vanilla M2A model faithfully retained the subtype-specific promoter

activity patterns (R^2 = 0.713 for DE genes with a single annotated promoter; 0.621 when all

annotated promoters for DE genes were included) (Additional file 1: Figures S11a and S11c).

Transfer learning using data from a single RMS further improved the consistency ($R^2 = 0.758$

and 0.673, respectively) (Additional file 1: Figures S11b and S11d).

GAS2 is a gene selectively expressed in ERMS [44]. Although promoter hypomethylation

was found in the ARMS tumors, the M2A model correctly predicted significantly stronger

promoter activities in ERMS tumors ($P = 0.01$, Wilcoxon rank-sum test) (Figures 4b and 4d).

Similarly, although both ERMS and ARMS tumors had *NOS1-005* promoter hypomethylation,

strong promoter activity was predicted in ARMS tumors only ($P = 0.0015$, Wilcoxon rank-sum

test) (Figures 4c and 4e), consistent with the ChIP-seq measurement.

M2A reveals the contribution of differentially methylated regions to promoter

activities

Although DNA methylation patterns, including differentially methylated regions (DMRs), are well-

established biomarkers for diverse diseases and have revealed molecularly and clinically

different subtypes for many cancers, their functional importance in individual gene regulation is

less clear [39-41]. For example, many cancer-specific CpG island hypermethylation regions

occur in genes that normally are not expressed or are expressed at only a low level [58].

Moreover, even in genes that are both differentially expressed and differentially methylated in

different subtypes, up-regulated samples can be associated with hypomethylation,

hypermethylation, or both [37, 44]. The observed nonlinear relation between DNA methylation

and gene expression complicated the functional interpretation of specific DMRs. In this analysis,

we interpret the functional roles of DMRs based on the promoter activities of their associated

DE genes.

To summarize unambiguously the contribution of DMRs to differential promoter activities, we

focused on 371 genes in ERMS and ARMS that have a single annotated promoter and that are

both differentially expressed (197 are over-expressed in ERMS, 172 are over-expressed in

ARMS) and differentially methylated (169 are hypomethylated, 128 are hypermethylated, and 74

have both hypomethylation and hypermethylation) in the two major RMS subtypes (Additional

file 2: Tables S9 and S10) [44]. Among these genes, 140 promoters showed significantly higher

H3K27ac measurements in the over-expressed subtype (FDR < 0.1, Wilcoxon rank-sum test),

whereas 206 promoters had measurements that were significantly higher when the DNAm-

based H3K27ac activity was measured. These 206 promoters included 118 of the 140

promoters identified using the observed signals. These results suggest that M2A can reveal the

role of DMRs in modulating the promoter activities of affected genes in a context-specific

manner.

M2A identifies subtype-specific promoter usage encoding different protein

isoforms in rhabdomyosarcoma

Alternative promoter usage is an important pretranslational mechanism for tissue-specific

regulation as it affects the diversity of isoforms available. Recently, light was shed on the

pervasiveness of alternative promoter usage in cancer; in some cases, promoter usage is a

more accurate reflection of patient survival than is gene expression [2]. Among 10,835 active

genes with multiple annotated promoters in the RMS dataset, we found 2,584 genes (24%) with

alternative primary promoter usage among 16 samples. We focused on 562 genes that 1) were

active in both the ERMS and ARMS subtypes and 2) had subtype-specific promoter usage. We

explored the accuracy of M2A in predicting alternative promoter usage in ARMS and ERMS

(Additional file 2: Table S11).

Based on measured promoter activities, 428 genes exhibited significant usage difference

between the two subtypes (FDR < 0.1, Wilcoxon rank-sum test [used as the ground truth]). The

M2A-inferred promoter activity landscape revealed 276 genes for which there was a significant

difference in promoter usage between the subtypes (FDR < 0.1, Wilcoxon rank-sum test), and

210 of them matched the ground truth (precision = 0.76, recall = 0.49, F1 score = 0.60)

(Additional file 2: Table S11).

PDZ Domain Containing Ring Finger 3 (*PDZRN3*) is a known target of the PAX3/7–FOXO1

fusion protein [59-61], which blocks terminal differentiation in myogenesis [62]. M2A predicted a

subtype-specific promoter usage pattern in *PDZRN3*. Functional studies have shown that

PDZRN3 regulates myoblast differentiation into myotubes through transcriptional and

posttranslational regulation of Id2 [62]. Its over-expression in ARMS was primarily driven by the

fusion protein binding adjacent to an alternative promoter (*PDZRN3-006*) located 191 kbp

downstream of the canonical promoter (*PDZRN3-001*) (Figure 5a). The subtype-specific isoform

usage is accompanied by DMRs of the alternative promoter and its immediate downstream

regions and is further confirmed by RNA-seq read alignment (Figure 5a). Compared to the

canonical isoform expressed in ERMS, the ARMS-preferred *PDZRN3-006* isoform lacks the

RING-finger and Sina domains in the N-terminus and harbors a shorter PDZ domain. The

isoform difference, as well as the differences in expression level, between subtypes may

contribute to the impairment of myogenesis at different stages in the development of ARMS and

ERMS tumors.

M2A identifies alternative promoter usages with potential prognostic values in

Ewing sarcoma

We next examined the predicted epigenetic promoter activities in 140 EWS samples with DNAm

data assayed by RRBS [22]. Three of these samples had matching ChIP-seq profiles for

H3K4me3 and H3K27ac. To interrogate this dataset, we applied the pretrained vanilla model

with transfer learning, as detailed above, to recalibrate the weights mapping the high-level

features to the promoter activities in the EWS cohort. Despite the difference in DNAm platforms

(WGBS for the NBL training model and RRBS for the EWS samples), the inferred promoter

activity landscape was accurate (R^2 = 0.718, 0.628, and 0.702 for the three samples with HM

profiles, using leave-one-out prediction) (Additional file 2: Table S12).

Ewing sarcomas with mutations in *TP53* and *STAG2* have a particularly dismal prognosis

[63]. We explored whether the promoter activity had additional prognostic value in 72 samples

for which survival data was available. Because of the limited sample size, the initial analysis

revealed a significant association between poor clinical outcomes and *TP53* mutations

(P = 0.00047, log-rank test) (Additional file 1: Figure S12a) but not *STAG2* mutations

(P = 0.67, log-rank test) (Additional file 1: Figure S12b). We identified 21 active genes that

showed a potential difference (absolute mean difference of log-scaled activity ≥ 1) between

TP53 mutant tumors and wild-type tumors, and we applied Cox proportional hazards models to

evaluate their potential contributions to patient survival that are independent of *TP53* or *STAG2*

mutation status (Additional file 2: Table S13). We performed the same analysis for 45 genes

with potential different promoter usage in tumors with and without *TP53* mutations (Additional

file 2: Table S14). Finally, we derived a multivariate Cox proportional hazards model including

both *TP53* and *STAG2* mutation status, one candidate gene with differential promoter activity

(*CALCB*), and five candidate gene isoforms (*CASZ1*, ENST00000496432; *RET*,

ENST00000479913; *TEX40*, ENST00000328404; TNS1, ENST00000446903; and SLC27A6,

ENST00000508645), and we followed this with backwards stepwise model selection. The final

model (Figure 5b) revealed potential protective roles for one candidate transcript *TNS1*

(ENST00000446903), a marginally protective role for candidate transcript *RET*

(ENST00000479913), and a candidate transcript associated with a poor prognosis, *SLC27A6*

(ENST00000508645).

TNS1 encodes the well-studied protein Tensin 1, and is involved in several key aspects of

cell function, including extracellular matrix formation, actin cytoskeleton formation, and signal

transduction [64, 65]. More recently, the up-regulation of TNS1 in colorectal cancer was found to

be associated with poor overall survival in patients [66], although previous studies have shown

suppression of *TNS1* expression [67] is associated with metastatic cancers. Our results suggest

that *TNS1* is a candidate prognostic indicator for EWS. Further studies are needed to draw

more attention to the functional roles of these genes/transcripts in EWS progression.

476   **Discussion**

Although epigenetic studies in disease models (cell lines, xenografts, and organoids) and in a

limited number of primary tumor samples have demonstrated the oncogenic contributions of

epigenetic deregulations to cancer initiation, progression, and response to treatment [48, 68, 69],

genome-wide profiling of promoter activities by using standard approaches (e.g., ChIP-seq or

CAGE) has not been carried out in large patient tumor cohorts, despite the continuous efforts of

large epigenome consortia [45, 46, 70]. Our analyses of *MYCN*-amplified NBL tumors revealed

both commonly active promoters and promoters that were active in some tumors but not in

others. Moreover, these sample-specific active promoters are functionally important, as they

drive the expression of several cancer consensus genes, including *MYC*. This observation is

consistent with recent reports of heterogeneous enhancer activities of cell line–defined super-

enhancers in primary gastric cancers [71], emphasizing the critical importance of deriving

sample-specific epigenomic signatures. To bridge the gap between the extensive epigenomic

resources in disease models and the limited ChIP-seq profiles of primary patient tumors, we

developed MethylationToActivity (M2A), a deep-learning framework, to characterize the

promoter activity landscape (both H3K4me3 and H3K27ac levels) in individual tumors by using

DNAm data, which is the most extensively documented epigenetic modification for patient

tumors and can be robustly and accurately profiled in FFPE archived retrospective samples.

M2A demonstrated excellent performance across various tumor types, with accuracy

comparable to that of ChIP-seq measurements of replicate samples from high-quality cohorts

(Figure 3).

Although our framework was strictly trained on HM levels, the inferred promoter activity was

highly correlated with the transcript-based gene expression levels quantified by RNA-seq

(Additional file 2: Tables S5, S7 and S8). The correlation between gene expression and inferred

promoter activities (mean R^2 = 0.668 for ENCODE data, 0.722 for K562, 0.705 for GM12878,

and 0.536 for H1-ESC) surpassed that with the state-of-art BPR model [42], which was

developed for predicting gene expression levels from DNAm patterns (the best reported R^2

values were 0.49 for K562, 0.37 for GM12878, and 0.25 for H1-ESC). Strikingly, the (indirect)

predictive accuracy of M2A for gene expression across nine ENCODE cell lines (average R^2 =

0.668) was comparable to the predictive accuracy of a model built on 11 HMs, one histone

variant, and DNase I hypersensitivity [11]. Similarly, the accuracy of binary prediction of

expressed genes (average AUC = 0.941 and 0.931 for the ENCODE and AML datasets,

respectively) surpassed that of DeepChrome (average AUC = 0.80), a state-of-art deep-learning

algorithm trained to predict expressed genes by using five core HMs (H3K4me3, H3K4me1,

H3K36me3, H3K9me3, and H3K27me3) [12]. These results further validated our framework.

Finally, both the M2A and BRP models suggest that it is insufficient to represent DNAm

information by using a simple average methylation level in the promoter region. To properly

reveal the regulatory roles of DNAm, we need to derive high-order features that capture spatial

relations among CpG probes (or window-based derived features calculated from them) in

promoters and in their vicinity. M2A uses the feature learning and selection characteristics of

CNNs to achieve its exceptional performance, thus demonstrating the rich information content of

DNAm signatures at both the genome-wide and local gene levels.

Analysis of the deep-learning framework revealed that M2A derives 1) high-level features

from DNAm patterns that are common among different tumors and 2) tumor subtype–specific

mapping functions from the mapping of high-level DNAm features to promoter activities in

individual tumor subtypes by using transfer learning (when feasible). Although our deep-learning

model cannot establish a causal relation between DNAm and promoter activities, these findings

nevertheless shed light on both the general and tumor subtype–specific rules for interpreting

DNAm patterns.

In evaluating our predictions, we found that several samples (the "poor performers") showed

abnormally low predictive accuracy with both the ENCODE and BLUEPRINT datasets.

Investigations of the promoter H3K27ac levels revealed that the fraction of active promoters in

these samples was substantially lower than in other samples. Furthermore, joint analyses with

RNA-seq data from the matching samples indicated that 1) in contrast to the predictive accuracy

of H3K27ac levels, the "poor performers" achieved comparable consistency between the M2A-

inferred promoter activity and gene expression; 2) the "poor performers" showed significantly

lower correlation between the measured promoter H3K27ac level and gene expression; 3) the

correlation between the measured promoter H3K27ac level and gene expression was highly

predictive of the accuracy of the H3K27ac prediction (Additional file 1: Figure S10),and 4) in

ENCODE samples with replicates, the replicate with better consistency between H3K27ac and

gene expression also showed significantly higher correlation between the actual and measured

H3K27ac level ($P = 0.0039$, Wilcoxon signed rank test, Additional file 2: Table S15, an example

of H3K27ac signal discrepancy between H1-ESC replicates shown in Additional file1: Figure

S8d). Although we cannot unequivocally rule out the possibility that these "poor performers"

share a distinct biological mechanism where promoter H3K27ac level is no longer a stronger

predictor for gene activities, these results suggested that the "poor performers" could reflect the

quality of the ChIP-seq results included in the test data. This observation emphasizes the value

of conducting a preliminary analysis to evaluate the data quality before incorporating public data.

It also suggests that M2A can provide a robust surrogate for promoter activities when the ChIP-

seq experimental data is questionable.

Alternative promoter usage increases the transcriptomic diversity during normal tissue

development and oncogenesis. Recent work demonstrated that an alternative promoter of

ERBB2 is predictive of a poor clinical outcome but that the canonical promoter shows no

significant association with survival in patients with low-grade glioma [2]. Whereas earlier

studies focused on identifying alternative promoter usage through CAGE or RNA-seq data, our

research has shown that alternative promoter usage can be extensively studied by using DNAm

profiles from diverse samples, including retrospective FFPE tumor samples, for which traditional

approaches (CAGE, ChIP-seq, and RNA-seq) are technically challenging. Our analysis of a

large EWS cohort (without matching RNA-seq data) revealed promoter activities for several

specific isoforms that are independently associated with clinical outcomes, including a specific

isoform of the *TNS1* gene (ENST00000446903). This demonstrates the importance of analyzing

alternative promoter usage in epigenomic studies.

Finally, our analysis quantitatively emphasizes that promoter activity is one of the

mechanisms that regulate the final transcriptional output: 1) both the observed and predicted

promoter activities account for 50%–70% of the variation in gene expression, and 2)

approximately 40% of the genes differentially expressed in ERMS and ARMS tumors show

significant differences in their promoter activities. In addition to promoter activities, other

epigenetic mechanisms, including enhancer activities, contribute substantially to gene regulation

[72]. Recent work has demonstrated the roles of DNAm in regulating enhancer activities [73]

and aberrant cancer-specific DNAm patterns in super-enhancers [74]. Consequently, we aim to

expand our M2A framework to infer enhancer activities from DNAm patterns in the future.

Conclusion

We have demonstrated that MethylationToActivity overcomes the unique challenges of

systematically characterizing promoter activities from DNA methylation signatures. It achieved

an accurate, robust and generalizable performance in various pediatric and adult cancers,

including both solid and hematologic malignant neoplasms. MethylationToActivity will serve as a

valuable tool to provide functional interpretation of DNAm deregulation, characterize promoter

activity differences from DNAm patterns, and reveal alternate promoter usage in patient tumors,

which will facilitate precision medicine by tailoring treatments based on both genetic variants

and epigenetic deregulation.

Methods

Datasets

Five separate publicly available datasets were used in this study, including a pediatric NBL O-

PDX dataset (N□=□16) [40]; an RMS O-PDX dataset (N□=□16) [42]; ENCODE datasets with

matching H3K27ac and H3K4me3 histone mark ChIP-seq, RNA-seq, and WGBS experimental

data (N□=□9) [43]; a DCC BLUEPRINT AML dataset (N□=□19) [44]; and a pediatric EWS

dataset (N□=□140) [19] (Additional file 2: Table S16). Of the 140 samples in the EWS cohort,

only three had matching ChIP-seq and reduced-representation bisulfite sequencing (RRBS)

data available; for the remaining 137 samples, only RRBS data was available. All other cohort

datasets (i.e., the RMS, NBL, ENCODE, and AML datasets) contained matching H3K27ac and

H3K4me3 profiles, along with RNA-seq and WGBS experimental data.

Feature processing

All datasets were evaluated using GENCODE annotation definitions (www.gencodegenes.org/);

the NBL, RMS, ENCODE, and EWS datasets were evaluated using GENECODE GRCh37.p13

(release 19), and the AML dataset was evaluated using GENCODE GRCh38.p13 (release 32).

Promoter regions are defined as the TSS ± 1 kbp, where the TSS is defined as each unique

transcript start position. To avoid using identical or near-identical promoter regions in training

and baseline performance, only TSSs with promoter regions with less than 50% overlap were

considered. Gene orientation was taken into account, and any promoters with overlying regions

but opposite orientations were not considered as overlapping. Because of differences in sex

amongst samples, all chromosome X and Y promoter regions were removed from consideration.

This resulted in a total of 96,756 and 104,722 non-overlapping promoter regions from the

annotation files GRCh37.p13 and GRCh38.p13, respectively. For gene expression analysis, we

followed the definition in [11] and retained all 141,152 and 147,980 autosomal promoters for

protein-coding genes from the annotation files GRCh37.p13 and GRCh38.p13, respectively.

M2A uses only one variable feature type: DNA methylation. For WGBS/RRBS data, the M-value

was calculated as follows:

$$Mval_k = log_2 \left(\frac{Methylated_k + \alpha}{Unmethylated_k + \alpha} \right),$$

where $Methylated_k$ and $Unmethylated_k$ correspond to the number of methylated and unmethylated

reads of the k^{th} CpG site, respectively. By default, the offset α was set to 0.5, and a global M-value

threshold was set to a maximum value of $log_2(65)$ and a minimum value of $log_2(\frac{1}{65})$. CpG sites with

coverage of less than five reads were removed.

M2A uses a promoter region–based windowed approach, comprising 20 windows of two sizes (250

bp and 2.5 kbp) and a step size equal to the window size, centered on a given TSS. For instance,

$\boldsymbol{W}_{ij} = \{W_{i1}, W_{i2}, W_{i3} ..., W_{i20}\}$ is the vector of windows where $\boldsymbol{i}$ represents a particular TSS and $\boldsymbol{j}$

represents a particular window corresponding with the $\boldsymbol{i^{th}}$ TSS. This means that W_{i10} and W_{i11}

represent the windows immediately downstream and upstream of the $\boldsymbol{i^{th}}$ TSS. Therefore,

$$W_{i1} = TSS_i - 10^{th} \; window,$$

619 $$W_{i2} = TSS_i - 9^{th} \; window, \; ...$$

$$W_{i19} = TSS_i + 9^{th} \; window,$$

$$W_{i20} = TSS_i + 10^{th} \; window$$

623   Each feature was calculated in this manner. The DNA methylation features, including the windowed

624 M-value mean, variance, and the fraction of the SSD of M-values (FSSD), were calculated and

625   represented by the feature vectors $Mave_i$, $Mvar_i$, and $Mfssd_i$. Therefore, the features for a particular

626   window, denoted as W_{ij}, would be calculated as follows:

$$Mave_{ij} = \frac{1}{n_{ij}} \sum Mval_{ik(j)},$$

$$Mvar_{ij} = \frac{1}{n_{ij}} \sum (Mave_{ij} - Mval_{ik(j)})^2,$$

627   and

$$Mfssd_{ij} = \frac{Mssd_{ij}}{\sum (Mave_i - Mval_k)^2},$$

628   where

$$Mave_i = \frac{1}{n_i} \sum Mval_k.$$

and

$Mssd_{ij} = \sum (Mave_{ij} - Mval_{ik(j)})^2,$

Here, i represents the promoter, j represents a specific window for a particular promoter, and

$Mval_k$ represents the Mval for individual CpGs in a region where $Mval_{k(j)}$ is the Mval for an individual

CpG in a specific window. Each feature was interleaved by window size to provide model input wherein

each window contained a number of "channels" equal to the number of features, with the feature array

shape being (*N*, 2, 20, 3), where *N* represents the total number of TSSs in a sample, 2 represents the

number of window sizes (250 bp and 2.5 kbp), 20 represents the number of windows, and 4 represents

the number of features per window. All features were scaled from 0.1 to 1 (using MinMaxScaler with

default values from sklearn version 0.22); in instances where windows overlapped regions without

methylation data, resulting in NaNs (such as chromosomal boundaries, telomeric regions, and

centromeric regions), these feature values were marked as 0.

Calculating histone modification enrichment

The response variable was calculated for each non-overlapping promoter region, a 2000-bp region

centered on each TSS. Histone modification enrichment (HM) for the i^{th} promoter region is calculated

as follows:

$$HM_i = log_2 \frac{\sum_k ChIP_{ik} + \alpha}{\sum_k Input_{ik} + \alpha},$$

where $\sum_k ChIP_{ik}$ represents the sum of either the H3K27ac or the H3K4me3 read signal mapped to the

promoter region at each position, $\sum_k Input_{ik}$ represents the sum of the control read signal mapped to

the promoter region, and α represents the 25th percentile of the $\sum_k Input_{ik}$ calculated for a given

sample.

M2A topology

M2A is a machine learning framework that leverages canonical deep-learning strategies, including

convolutional neural network (CNN) and fully connected (FC) layers. Each layer employs a LeakyReLU

(alpha□=□0.1) and a kernel constraint by L2-normalization using maxnorm(3). CNN layers are two-

dimensional, with zero padding, a stepsize of (1,1), and a kernel size of (1, 3) to maintain feature space

and prevent convolutions across features from different window sizes. To test the efficacy of this

approach, we compared the performance of a traditional artificial neural network (ANN) consisting of

two FC layers versus three CNN layers in addition to two FC layers. During transfer learning, weights

corresponding to each of the three CNN layers of the six–NBL O-PDX M2A model trained previously

were frozen; only the weights corresponding to the two FC layers were optimized. A summary of all

model topologies and parameters can be found in Additional file 2: Table S17. To train and test each

model topology, we used Keras (v2.2.4) and Tensorflow (v2.1.0) in Python 3.6.5.

M2A parameter tuning

Parameters such as the window size, batch size, and kernel size were optimized using the validation

NBL data set (n=10). Each parameter configuration was tested holding all other parameters constant,

and models with a numerical performance advantage were chosen. For batch size, three configurations

were tested (64, 128, 256). Four kernel size configurations ([1,2], [1,3], [1,4], [1,5]) were tested. and

window configurations ([100bp, 1000bp], [250bp, 2500bp], [500bp, 5000bp]) were considered. Due

to >50% uninformative features in the 100bp window resolution, only the (250bp, 2500bp) model and

(500bp, 5000bp) model performances were compared (Additional file 1: Figure S13; Additional file 2:

Table S18).

Training M2A

The core M2A model (without transfer) training set consisted of six O-PDX samples from the 16-

sample NBL cohort (Additional file 2: Table S19); we trained separate models for H3K27ac and

H3K4me3 HMs with the same WGBS features as the input. After the base models were trained,

transfer learning was employed for three separate datasets, namely the RMS, AML, and EWS

datasets. Each transfer learning model was trained using one sample from the cohort, for a total

of N models, where N equals the number of samples in the cohort. For the RMS and EWS

cohorts, an ensemble approach was used, whereby an averaged prediction from N−1 models

was generated after transfer learning with each sample. The same approach was used with the

AML cohort, except that only samples with $R^2 \geq 0.60$ between FPKM and H3K27ac were used

for transfer learning.

For each training scheme, the same parameters were used, including an 80/20 training/validation

split and a batch size of 64 (Additional file 2: Table S17). All sample input was randomized before

training. The Keras implementation of adadelta (default parameters) minimizing the mean squared error

(MSE) was used to optimize M2A. To prevent overtraining, the EarlyStopping method was employed by

monitoring validation loss for 10 epochs without at least a minimal gain in performance (min_delta =

0.0001) for a maximum of 80 epochs. In no case was the maximum number of epochs reached.

Determining promoter diversity in NBL models

Promoter region–based H3K27ac distributions clearly show a bimodal distribution (see

Additional file 1: Figure S1); therefore, to determine class occupancy (active versus inactive), a

Gaussian mixture model (GaussianMixture from sklearn version 0.22 with n_components = 2)

was applied for each individual sample. To determine the percentage of differentially active

promoters among all active promoters, we used a pairwise comparison approach for all samples.

Cancer consensus genes were downloaded from COSMIC (https://cancer.sanger.ac.uk/census)

(accessed on February 1, 2020). To avoid artificially inflated values from genes with multiple

TSSs, only cancer consensus genes with a single TSS according to GENCODE GRCh37.p13

(release 19) definitions were considered.

Evaluating M2A performance

When determining prediction performance, two primary metrics were considered, namely the R^2 and

the root mean squared error (RMSE). To measure the accuracy of M2A in predicting expressed genes,

we calculated the AUC-ROC by using roc_curve and the precision-recall curve AUC by using

average_precision_score from sklearn 0.22. Paired analyses were tested for statistical significance by

using a Wilcoxon signed-rank test (R v3.4.1). To determine outliers, a median-based method was

implemented using the "outlier" function in the R package GmAMisc [55]. To ensure that low-

mappability regions were not a confounding factor, we used the

wgEncodeCrgMApabilityAlign100mer.bw file downloaded from the UCSC Genome Browser

(http://genome.ucsc.edu/cgi-bin/hgFileUi?db=hg19&g=wgEncodeMapability). For GRCh38.p13

annotations, we used liftOver from UCSC (http://hgdownload.cse.ucsc.edu/admin/exe/linux.x86_64/) to

convert the mappability track to GRCh38. For all performance-related analyses, the average value of

717   this track within each non-overlapping 2000-bp promoter region was calculated, and promoter regions

718   with mappability > 0.75 were retained for evaluation.

719   In analyses between H3K27ac levels (actual or predicted) and gene expression, we followed the

720   gene filtering steps described in [11]: all autosomal protein-coding gene promoters were considered

721   and for genes with multiple promoters, we use the maximum promoter activity to represent the gene.

Baseline models

724   Multivariate adaptive regression splines (MARS) was implemented by "earth" package in R (all default

725   values were used), and the random forest baseline model was implemented by the

726   "sklearn.ensemble.RandomForestRegressor" package in python 3.7.0 (max_features='sqrt'). For

727   comparison purposes, each model tested used identical feature input to M2A.

M2A captures the impact of DMRs on promoter region activity

730   The lists of genes that are differentially expressed in ERMS and ARMS samples and genes with DMRs

731   were previously reported in reference [42].

Alternative promoter usage analyses

To infer alternative promoter usage that was specific to the RMS subtypes ARMS and ERMS, we first

delineated the "active" vs. "inactive" promoters in a subtype (ERMS or ARMS) by applying a threshold

of $mean(H3K27ac) > 1$ for the average samples in the subtype. Next, primary promoters from multi-

promoter genes were determined by the average H3K27ac level within a specific subtype, and the

promoter with the maximum activity from a given gene was counted as the primary promoter. The

differences in promoter usage between two subtypes was defined as the difference between the activity

sum of the primary promoters and the activity sum of the secondary promoters in the two subtypes.

Analysis of alternative promoter usage in EWS

In the same manner as the RMS subtype analysis, EWS alternative promoter usage was determined in

two patient sample groups: *TP53* mutant and *TP53* wild-type groups (for sample status, see Additional

file 2: Table S18). Candidate genes were identified as active genes (H3K27ac ≥ 1 in at least one group)

with potential differential promoter activities (absolute difference ≥ 1) in the *TP53* wild-type and *TP53*

mutant groups. Candidate genes with alternative promoter usage were identified as genes that used

different active primary promoters in the wild-type and mutant groups and had an average promoter

usage difference of at least 0.4 between groups. Both alternative promoters and differentially active

promoters were considered in prognostic analyses (univariate screening incorporating both *TP53* and

STAG2 mutation status, followed by a multivariate analysis) using Cox proportional hazard models (R

3.4.1). The final model was derived from backward stepwise selection from a Cox proportional hazards

model including *TP53* and *STAG2* mutation status and all potential markers (all genes or promoters

with an FDR < 0.05 in the univariate analysis).

Analysis of M2A feature input and extracted features

To determine the merit of a CNN-based approach, each feature average for a particular window

position and window size were plotted with a 95% confidence interval. The plotted feature distribution

was calculated from all M2A vanilla model training data input (NBL, N=6), and stratified by promoter

status. Promoter status was determined by class occupancy of both H3K27ac and H3K4me3, (active

versus inactive), where (H3K27ac=active), (H3K27ac=inactive, H3K4me3=active), and

(H3K27ac=inactive, H3K4me3=inactive), represents active, poised, and inactive promoters,

respectively. Class occupancy was determined by applying a Gaussian mixture model

(GaussianMixture from sklearn version 0.22 with n_components = 2).

The efficacy of the CNN-based feature extraction was tested by 1) training sample input feature

predictive performance as compared to CNN-extracted feature performance, calculated by Pearson's

767   R^2 for all features across the entire training set (each window size considered separately; NBL, N=6)

768   and 2) The performance (Pearson's R^2) of the best performing feature identified in 1) when applied to

769   each sample in the validation set (NBL, N=10; Additional File 1: Figure S3).

771   M2A code availability

772   The latest M2A models, feature generation, prediction pipeline, and a Docker image of the M2A

773   environment pre-loaded are available for download at https://github.com/chenlab-sj/M2A. Additionally,

774   source code with detailed instructions for transfer learning using the M2A model with input samples

775   from other domains is available. The cloud-based implementation of M2A is available to anyone with a

776   (free) St. Jude Cloud account (https://platform.stjude.cloud/workflows/methylation-to-activity).

778   Declarations

779   Ethics approval and consent to participate
780   Not applicable

781   Consent for publication
782   Not applicable

783   Availability of data and materials
784   The datasets generated and/or analyzed during the current study are publicly available, as detailed in

785   Additional File 2: Table S16.

786   Competing interests
787   The authors declare that they have no known competing financial interests or personal relationships

788   that could have appeared to influence the work reported in this paper.

789   Funding

Authors' contributions

794 **JW**: Methodology, software, data analysis, result interpretation, writing - original draft, writing - review &

795 editing. **BX**: Resources, data curation. **DP**: Software, resources. **AT**: Software, resources. **CL**: Result

796 interpretation, writing – review & editing. **JY**: Result interpretation, writing – review & editing. **XC**:

797 Conceptualization, methodology, data analysis, result interpretation, writing - review & editing,

798 supervision.

Acknowledgements

800 We thank Keith A. Laycock, PhD, ELS, for editing the manuscript.

References

1. Davuluri RV, Suzuki Y, Sugano S, Plass C, Huang TH: **The functional consequences of**
alternative promoter use in mammalian genomes. *Trends Genet* 2008, **24:**167-177.
2. Demircioglu D, Cukuroglu E, Kindermans M, Nandi T, Calabrese C, Fonseca NA,
Kahles A, Lehmann KV, Stegle O, Brazma A, et al: **A Pan-cancer Transcriptome**
Analysis Reveals Pervasive Regulation through Alternative Promoters. *Cell* 2019,
178:1465-1477 e1417.
3. Qamra A, Xing M, Padmanabhan N, Kwok JJT, Zhang S, Xu C, Leong YS, Lee Lim AP,
Tang Q, Ooi WF, et al: **Epigenomic Promoter Alterations Amplify Gene Isoform and**
Immunogenic Diversity in Gastric Adenocarcinoma. *Cancer Discov* 2017, **7:**630-651.
4. Sotillo E, Barrett DM, Black KL, Bagashev A, Oldridge D, Wu G, Sussman R, Lanauze
C, Ruella M, Gazzara MR, et al: **Convergence of Acquired Mutations and Alternative**
Splicing of CD19 Enables Resistance to CART-19 Immunotherapy. *Cancer Discov*
2015, **5:**1282-1295.
5. Ma X, Liu Y, Liu Y, Alexandrov LB, Edmonson MN, Gawad C, Zhou X, Li Y, Rusch
MC, Easton J, et al: **Pan-cancer genome and transcriptome analyses of 1,699**
paediatric leukaemias and solid tumours. *Nature* 2018, **555:**371-376.
6. Grobner SN, Worst BC, Weischenfeldt J, Buchhalter I, Kleinheinz K, Rudneva VA,
Johann PD, Balasubramanian GP, Segura-Wang M, Brabetz S, et al: **The landscape of**
genomic alterations across childhood cancers. *Nature* 2018, **555:**321-327.

7.   Huether R, Dong L, Chen X, Wu G, Parker M, Wei L, Ma J, Edmonson MN, Hedlund
823   EK, Rusch MC, et al: **The landscape of somatic mutations in epigenetic regulators**
824   **across 1,000 paediatric cancer genomes.** *Nat Commun* 2014, **5:**3630.
825   8.   Andersson R, Sandelin A: **Determinants of enhancer and promoter activities of**
826   **regulatory elements.** *Nat Rev Genet* 2020, **21:**71-87.
827   9.   Kelley DZ, Flam EL, Izumchenko E, Danilova LV, Wulf HA, Guo T, Singman DA,
828   Afsari B, Skaist AM, Considine M, et al: **Integrated Analysis of Whole-Genome ChIP-**
829   **Seq and RNA-Seq Data of Primary Head and Neck Tumor Samples Associates HPV**
830   **Integration Sites with Open Chromatin Marks.** *Cancer Res* 2017, **77:**6538-6550.
831   10.   Karlic R, Chung HR, Lasserre J, Vlahovicek K, Vingron M: **Histone modification levels**
832   **are predictive for gene expression.** *Proc Natl Acad Sci U S A* 2010, **107:**2926-2931.
833   11.   Dong X, Greven MC, Kundaje A, Djebali S, Brown JB, Cheng C, Gingeras TR, Gerstein
834   M, Guigo R, Birney E, Weng Z: **Modeling gene expression using chromatin features**
835   **in various cellular contexts.** *Genome Biol* 2012, **13:**R53.
836   12.   Singh R, Lanchantin J, Robins G, Qi Y: **DeepChrome: deep-learning for predicting**
837   **gene expression from histone modifications.** *Bioinformatics* 2016, **32:**i639-i648.
838   13.   Kagohara LT, Stein-O'Brien GL, Kelley D, Flam E, Wick HC, Danilova LV, Easwaran H,
839   Favorov AV, Qian J, Gaykalova DA, Fertig EJ: **Epigenetic regulation of gene**
840   **expression in cancer: techniques, resources and analysis.** *Brief Funct Genomics* 2018,
841   **17:**49-63.
842   14.   Zhang P, Lehmann BD, Shyr Y, Guo Y: **The Utilization of Formalin Fixed-Paraffin-**
843   **Embedded Specimens in High Throughput Genomic Studies.** *Int J Genomics* 2017,
844   **2017:**1926304.
845   15.   Moran S, Vizoso M, Martinez-Cardus A, Gomez A, Matias-Guiu X, Chiavenna SM,
846   Fernandez AG, Esteller M: **Validation of DNA methylation profiling in formalin-fixed**
847   **paraffin-embedded samples using the Infinium HumanMethylation450 Microarray.**
848   *Epigenetics* 2014, **9:**829-833.
849   16.   de Ruijter TC, de Hoon JP, Slaats J, de Vries B, Janssen MJ, van Wezel T, Aarts MJ, van
850   Engeland M, Tjan-Heijnen VC, Van Neste L, Veeck J: **Formalin-fixed, paraffin-**
851   **embedded (FFPE) tissue epigenomics using Infinium HumanMethylation450**
852   **BeadChip assays.** *Lab Invest* 2015, **95:**833-842.
853   17.   Gu H, Bock C, Mikkelsen TS, Jager N, Smith ZD, Tomazou E, Gnirke A, Lander ES,
854   Meissner A: **Genome-scale DNA methylation mapping of clinical samples at single-**
855   **nucleotide resolution.** *Nat Methods* 2010, **7:**133-136.
856   18.   Ziller MJ, Gu H, Muller F, Donaghey J, Tsai LT, Kohlbacher O, De Jager PL, Rosen ED,
857   Bennett DA, Bernstein BE, et al: **Charting a dynamic DNA methylation landscape of**
858   **the human genome.** *Nature* 2013, **500:**477-481.
859   19.   Charlet J, Duymich CE, Lay FD, Mundbjerg K, Dalsgaard Sorensen K, Liang G, Jones
860   PA: **Bivalent Regions of Cytosine Methylation and H3K27 Acetylation Suggest an**
861   **Active Role for DNA Methylation at Enhancers.** *Mol Cell* 2016, **62:**422-431.
862   20.   Onuchic V, Lurie E, Carrero I, Pawliczek P, Patel RY, Rozowsky J, Galeev T, Huang Z,
863   Altshuler RC, Zhang Z, et al: **Allele-specific epigenome maps reveal sequence-**
864   **dependent stochastic switching at regulatory loci.** *Science* 2018, **361**.
865   21.   Stadler MB, Murr R, Burger L, Ivanek R, Lienert F, Scholer A, van Nimwegen E,
866   Wirbelauer C, Oakeley EJ, Gaidatzis D, et al: **DNA-binding factors shape the mouse**
867   **methylome at distal regulatory regions.** *Nature* 2011, **480:**490-495.

22.   Sheffield NC, Pierron G, Klughammer J, Datlinger P, Schonegger A, Schuster M, Hadler
J, Surdez D, Guillemot D, Lapouble E, et al: **DNA methylation heterogeneity defines a**
disease spectrum in Ewing sarcoma. *Nat Med* 2017, **23:**386-395.
23.   Zhu H, Wang G, Qian J: **Transcription factors as readers and effectors of DNA**
methylation. *Nature Reviews Genetics* 2016, **17:**551-565.
24.   Kondo Y: **Epigenetic cross-talk between DNA methylation and histone modifications**
in human cancers. *Yonsei Med J* 2009, **50:**455-463.
25.   Rothbart SB, Strahl BD: **Interpreting the language of histone and DNA modifications.**
Biochim Biophys Acta 2014, **1839:**627-643.
26.   Hashimshony T, Zhang J, Keshet I, Bustin M, Cedar H: **The role of DNA methylation**
in setting up chromatin structure during development. *Nat Genet* 2003, **34:**187-192.
27.   Moore LD, Le T, Fan G: **DNA methylation and its basic function.**
Neuropsychopharmacology 2013, **38:**23-38.
28.   Jenkinson G, Pujadas E, Goutsias J, Feinberg AP: **Potential energy landscapes identify**
the information-theoretic nature of the epigenome. *Nat Genet* 2017, **49:**719-729.
29.   Fortin JP, Hansen KD: **Reconstructing A/B compartments as revealed by Hi-C using**
long-range correlations in epigenetic data. *Genome Biol* 2015, **16:**180.
30.   Simmonds P, Loomis E, Curry E: **DNA methylation-based chromatin compartments**
and ChIP-seq profiles reveal transcriptional drivers of prostate carcinogenesis.
Genome Med 2017, **9:**54.
31.   Mack SC, Witt H, Piro RM, Gu L, Zuyderduyn S, Stutz AM, Wang X, Gallo M, Garzia L,
Zayne K, et al: **Epigenomic alterations define lethal CIMP-positive ependymomas of**
infancy. *Nature* 2014, **506:**445-450.
32.   Teodoridis JM, Hardie C, Brown R: **CpG island methylator phenotype (CIMP) in**
cancer: causes and implications. *Cancer Lett* 2008, **268:**177-186.
33.   Capper D, Jones DTW, Sill M, Hovestadt V, Schrimpf D, Sturm D, Koelsche C, Sahm F,
Chavez L, Reuss DE, et al: **DNA methylation-based classification of central nervous**
system tumours. *Nature* 2018, **555:**469-474.
34.   Perez E, Capper D: **Invited Review: DNA methylation-based classification of**
paediatric brain tumours. *Neuropathol Appl Neurobiol* 2020.
35.   Strahl BD, Allis CD: **The language of covalent histone modifications.** *Nature* 2000,
403:41-45.
36.   Musselman CA, Lalonde ME, Cote J, Kutateladze TG: **Perceiving the epigenetic**
landscape through histone readers. *Nat Struct Mol Biol* 2012, **19:**1218-1227.
37.   Spainhour JC, Lim HS, Yi SV, Qiu P: **Correlation Patterns Between DNA**
Methylation and Gene Expression in The Cancer Genome Atlas. *Cancer Inform* 2019,
18:1176935119828776.
38.   Baylin SB: **DNA methylation and gene silencing in cancer.** *Nature Clinical Practice*
Oncology 2005, **2:**S4-S11.
39.   Jones PA: **Functions of DNA methylation: islands, start sites, gene bodies and**
beyond. *Nature Reviews Genetics* 2012, **13:**484-492.
40.   Lay FD, Liu Y, Kelly TK, Witt H, Farnham PJ, Jones PA, Berman BP: **The role of DNA**
methylation in directing the functional organization of the cancer epigenome.
Genome Research 2015, **25:**467-477.
41.   Kagohara LT, Stein-O'Brien GL, Kelley D, Flam E, Wick HC, Danilova LV, Easwaran
H, Favorov AV, Qian J, Gaykalova DA, Fertig EJ: **Epigenetic regulation of gene**

expression in cancer: techniques, resources and analysis. *Briefings in Functional*
Genomics 2018, **17:**49-63.
42.   Kapourani CA, Sanguinetti G: **Higher order methylation features for clustering and**
prediction in epigenomic studies. *Bioinformatics* 2016, **32:**i405-i412.
43.   Zeineldin M, Federico S, Chen X, Fan Y, Xu B, Stewart E, Zhou X, Jeon J, Griffiths L,
Nguyen R, et al: **MYCN amplification and ATRX mutations are incompatible in**
neuroblastoma. *Nat Commun* 2020, **11:**913.
44.   Stewart E, McEvoy J, Wang H, Chen X, Honnell V, Ocarz M, Gordon B, Dapper J,
Blankenship K, Yang Y, et al: **Identification of Therapeutic Targets in**
Rhabdomyosarcoma through Integrated Genomic, Epigenomic, and Proteomic
Analyses. *Cancer Cell* 2018.
45.   Consortium EP: **An integrated encyclopedia of DNA elements in the human genome.**
Nature 2012, **489:**57-74.
46.   Stunnenberg HG, International Human Epigenome C, Hirst M: **The International**
Human Epigenome Consortium: A Blueprint for Scientific Collaboration and
Discovery. *Cell* 2016, **167:**1145-1149.
47.   Singh AA, Schuurman K, Nevedomskaya E, Stelloo S, Linder S, Droog M, Kim Y,
Sanders J, van der Poel H, Bergman AM, et al: **Optimized ChIP-seq method facilitates**
transcription factor profiling in human tumors. *Life Sci Alliance* 2019, **2:**e201800115.
48.   Seligson DB, Horvath S, Shi T, Yu H, Tze S, Grunstein M, Kurdistani SK: **Global**
histone modification patterns predict risk of prostate cancer recurrence. *Nature*
2005, **435:**1262-1266.
49.   Chen X, Stewart E, Shelat AA, Qu C, Bahrami A, Hatley M, Wu G, Bradley C, McEvoy
J, Pappo A, et al: **Targeting oxidative stress in embryonal rhabdomyosarcoma.**
Cancer Cell 2013, **24:**710-724.
50.   Stewart E, Shelat A, Bradley C, Chen X, Federico S, Thiagarajan S, Shirinifard A,
Bahrami A, Pappo A, Qu C, et al: **Development and characterization of a human**
orthotopic neuroblastoma xenograft. *Dev Biol* 2015, **407:**344-355.
51.   Stewart E, Federico SM, Chen X, Shelat AA, Bradley C, Gordon B, Karlstrom A,
Twarog NR, Clay MR, Bahrami A, et al: **Orthotopic patient-derived xenografts of**
paediatric solid tumours. *Nature* 2017, **549:**96-100.
52.   Murphy AJ, Chen X, Pinto EM, Williams JS, Clay MR, Pounds SB, Cao X, Shi L, Lin T,
Neale G, et al: **Forty-five patient-derived xenografts capture the clinical and**
biological heterogeneity of Wilms tumor. *Nat Commun* 2019, **10:**5806.
53.   Angermueller C, Lee HJ, Reik W, Stegle O: **DeepCpG: accurate prediction of single-**
cell DNA methylation states using deep learning. *Genome Biol* 2017, **18:**67.
54.   Barredo Arrieta A, Díaz-Rodríguez N, Del Ser J, Bennetot A, Tabik S, Barbado A,
Garcia S, Gil-Lopez S, Molina D, Benjamins R, et al: **Explainable Artificial**
Intelligence (XAI): Concepts, taxonomies, opportunities and challenges toward
responsible AI. *Information Fusion* 2020, **58:**82-115.
55.   Wilcox RR: **The Normal Curve and Outlier Detection.** In *Fundamentals of Modern*
Statistical Methods. 2001: 31-47
56.   Golob JL, Kumar RM, Guenther MG, Pabon LM, Pratt GA, Loring JF, Laurent LC,
Young RA, Murry CE: **Evidence that gene activation and silencing during stem cell**
differentiation requires a transcriptionally paused intermediate state. *PLoS One*
2011, **6:**e22416.

57. Berdasco M, Esteller M: **Clinical epigenetics: seizing opportunities for translation.** *Nat Rev Genet* 2019, **20:**109-127.

58. Hinoue T, Weisenberger DJ, Lange CPE, Shen H, Byun HM, Van Den Berg D, Malik S, Pan F, Noushmehr H, van Dijk CM, et al: **Genome-scale analysis of aberrant DNA methylation in colorectal cancer.** *Genome Research* 2011, **22:**271-282.

59. Cao L, Yu Y, Bilke S, Walker RL, Mayeenuddin LH, Azorsa DO, Yang F, Pineda M, Helman LJ, Meltzer PS: **Genome-wide identification of PAX3-FKHR binding sites in rhabdomyosarcoma reveals candidate target genes important for development and cancer.** *Cancer research* 2010, **70:**6497-6508.

60. Laé M, Ahn E, Mercado G, Chuai S, Edgar M, Pawel B, Olshen A, Barr F, Ladanyi M: **Global gene expression profiling of PAX-FKHR fusion-positive alveolar and PAX-FKHR fusion-negative embryonal rhabdomyosarcomas.** *The Journal of Pathology* 2007, **212:**143-151.

61. Marshall AD, Grosveld GC: **Alveolar rhabdomyosarcoma – The molecular drivers of PAX3/7-FOXO1-induced tumorigenesis.** *Skeletal Muscle* 2012, **2:**25.

62. Honda T, Inui M: **PDZRN3 regulates differentiation of myoblasts into myotubes through transcriptional and posttranslational control of Id2.** *Journal of Cellular Physiology* 2019, **234:**2963-2972.

63. Tirode F, Surdez D, Ma X, Parker M, Le Deley MC, Bahrami A, Zhang Z, Lapouble E, Grossetete-Lalami S, Rusch M, et al: **Genomic Landscape of Ewing Sarcoma Defines an Aggressive Subtype with Co-Association of STAG2 and TP53 Mutations.** *Cancer Discov* 2014, **4:**1342-1353.

64. Lo SH, An Q, Bao S, Wong WK, Liu Y, Janmey PA, Hartwig JH, Chen LB: **Molecular cloning of chick cardiac muscle tensin. Full-length cDNA sequence, expression, and characterization.** *J Biol Chem* 1994, **269:**22310-22319.

65. Chen H, Duncan IC, Bozorgchami H, Lo SH: **Tensin1 and a previously undocumented family member, tensin2, positively regulate cell migration.** *Proc Natl Acad Sci U S A* 2002, **99:**733-738.

66. Zhou H, Zhang Y, Wu L, Xie W, Li L, Yuan Y, Chen Y, Lin Y, He X: **Elevated transgelin/TNS1 expression is a potential biomarker in human colorectal cancer.** *Oncotarget* 2018, **9:**1107-1113.

67. Hall EH, Daugherty AE, Choi CK, Horwitz AF, Brautigan DL: **Tensin1 requires protein phosphatase-1alpha in addition to RhoGAP DLC-1 to control cell polarization, migration, and invasion.** *J Biol Chem* 2009, **284:**34713-34722.

68. Shen H, Laird PW: **Interplay between the cancer genome and epigenome.** *Cell* 2013, **153:**38-55.

69. De Craene B, Berx G: **Regulatory networks defining EMT during cancer initiation and progression.** *Nat Rev Cancer* 2013, **13:**97-110.

70. Roadmap Epigenomics C, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang J, et al: **Integrative analysis of 111 reference human epigenomes.** *Nature* 2015, **518:**317-330.

71. Ooi WF, Xing M, Xu C, Yao X, Ramlee MK, Lim MC, Cao F, Lim K, Babu D, Poon LF, et al: **Epigenomic profiling of primary gastric adenocarcinoma reveals super-enhancer heterogeneity.** *Nat Commun* 2016, **7:**12983.

72. Pennacchio LA, Bickmore W, Dean A, Nobrega MA, Bejerano G: **Enhancers: five essential questions.** *Nat Rev Genet* 2013, **14:**288-295.

1006   73.   Weigel C, Chaisaingmongkol J, Assenov Y, Kuhmann C, Winkler V, Santi I, Bogatyrova
1007   O, Kaucher S, Bermejo JL, Leung SY, et al: **DNA methylation at an enhancer of the**
1008   **three prime repair exonuclease 2 gene (TREX2) is linked to gene expression and**
1009   **survival in laryngeal cancer.** *Clin Epigenetics* 2019, **11:**67.
1010   74.   Heyn H, Vidal E, Ferreira HJ, Vizoso M, Sayols S, Gomez A, Moran S, Boque-Sastre R,
1011   Guil S, Martinez-Cardus A, et al: **Epigenomic analysis detects aberrant super-**
1012   **enhancer DNA methylation in human cancer.** *Genome Biol* 2016, **17:**11.

1013

Figure Legends

Figure 1 Variations of promoter H3K27ac levels among *MYCN* amplified NBL samples.

(a–c) Promoters were classified as active or inactive based on H3K27ac levels in individual NBL

tumors: **(a)** NBL cell line samples, **(b)** NBL O-PDX samples, and **(c)** NBL cell line and O-PDX

samples. Each promoter region plotted spans TSS ± 5,000 bp binned by non-overlapping 250

bp windows; the color bar represents the scaled windowed H3K27ac enrichment, from 0 (lowest)

to 3 (highest). Horizontal dotted lines delineate shared promoters (active in all samples) and

sample-specific promoters (promoter activity in at least one sample was different from the

remaining samples). Promoters were sorted by average descending H3K27ac enrichment

across all samples within each group. **(d, e)** Venn diagram indicating the number of shared and

sample-specific H3K27ac promoter activities between NBL cell line samples (**d**) and O-PDX

samples **(e)**. **(f)** The promoter activity variation is highlighted by the proportion of sample-

specific active promoters among all H3K27ac active promoters, as compared within cell line or

O-PDX NBL samples, and between cell line and O-PDX NBL samples.

1028

Figure 2. M2A feature processing and training workflow.

The M2A framework hinges on the feature processing pipeline. **(a)** First, windowed features (20

total non-overlapping windows for each of two sizes including 250 bp and 2,500 bp) centered

around the TSS are calculated from WGBS data for each unique promoter region, extending up

to 2,500 bp (250 bp window), and 25 kbp (2,500 bp window) away from the TSS. Response

variables (H3K27ac and H3K4me3) for separate model training were generated for matching

promoter regions (TSS ± 1 kbp). The matching window features and response variables serve

as input to the model topologies, where M2A first extracts high-level features by using a series

of convolutional layers then maps these features to response variables in fully-connected (FC)

layers. Transfer learning with M2A leverages pretrained feature extraction (frozen CNN layers

indicated in blue), training only the FC layers. **(b)** The overall workflow for training, validation,

and testing M2A is detailed, as well as an overview of the analyses performed to validate M2A

performance in different real-world applications. M2A models for H3K4me3 and H3K27ac were

trained separately, indicated by blue (H3K4me3) and green (H3K27ac).

1043

Figure 3. M2A performance in NBL and RMS cohorts.

(a, b) Analysis of the performance of M2A in NBL and RMS cohorts with **(a)** H3K4me3 inference

and **(b)** H3K27ac inference. ENCODE replicate consistencies were calculated as Pearson's

correlation squared (R^2) between replicates (two replicates per sample). ENCODE sample

KMS-11 was excluded as an apparent outlier R^2=0.016, RMSE = 1.869. Prediction accuracy

was measured by R^2 between the actual measurement and M2A's prediction. **(c–g)** Individual

examples of median M2A performers in **(c)** NBL cell line H3K27ac inference, **(d)** NBL cell line

H3K4me3 inference, **(e)** RMS H3K27ac inference (pre-transfer), **(f)** RMS H3K4me3 inference,

and **(g)** RMS H3K27ac inference (post-transfer). To indicate the density of data points where

individual data points cannot be resolved, a KDE was applied, called from 1 (highest) to 0

(lowest). **(h)** The boost to M2A performance (measured by RMSE) due to transfer learning is

shown, as applied and tested in RMS samples.

1056

Figure 4. M2A recapitulates subtype differences in RMS.

(a) A t-distributed Stochastic Neighbor Embedding (tSNE) analysis of observed (left), M2A

inferred (center: pre-transfer, right: post-transfer) H3K27ac promoter levels. Embryonal and

alveolar RMS subtypes are well separated in each analysis, demonstrating that M2A inferred

H3K27ac levels maintain the delineation of RMS subtypes, consistent with the observed

H3K27ac tSNE analysis. **(b–d)** The subtype-specific genes *GAS2* **(b)** and *NOS1* **(c)** show

subtype distinct patterns of DNAm, H3K27ac, and H3K4me3 levels. The windowed average

DNAm feature (2,500 bp windows, over the genomic region TSS ± 25 kb) is shown as example

(partial) M2A input. These subtype differences were faithfully recapitulated by the M2A

H3K27ac inferences for *GAS2* **(d)** and *NOS1* **(e)**.

1067

Figure 5. M2A reveals alternate promoter usage in RMS and EWS.

(a) An analysis of alternate primary promoter usage between RMS subtypes ERMS and ARMS

shows that M2A appropriately predicts subtype–specific promoter usage in *PDZRN3*, a known

target of the PAX3/7–FOXO1 fusion protein, which is consistent at the observed values of

H3K27ac, H3K4me3, RNA-seq, and DNA methylation. Partial M2A input (DNAm 2,500 bp

windowed average) is shown to emphasize the DNAm patterns in the genomic region

surrounding the TSS ± 25 kb. **(b)** Alternate promoter usage in EWS patient samples with and

without *TP53* mutations were incorporated in a Cox proportional hazards model, highlighting the

potential prognostic value of the isoforms identified by M2A.

1077

Additional Files

Additional file 1 (file type .PDF, 7 MB):

Figure S1. NBL sample H3K27ac promoter distribution.

A comparison of promoter H3K27ac enrichment in *MYCN* amplified (MNA) NBL cell line and O-

PDX samples shows a clear bimodal distribution, delineating "active" and "non-active"

promoters.

1084

Figure S2. DNAm input feature pattern analysis.

The DNAm features show promoter-status-specific patterns between active, poised, and

inactive promoters, emphasizing the utility of positional-based relationships in a windowed

DNAm feature approach. For both 250 bp and 2,500 bp window sizes, the average scaled input

feature was plotted in relationship to the TSS (ribbons represent the 95% confidence interval),

stratified by the promoter status. Promoter status was determined by class occupancy of both

H3K27ac and H3K4me3, either "active" or "inactive", where (H3K27ac=active),

(H3K27ac=inactive, H3K4me3=active), and (H3K27ac=inactive, H3K4me3=inactive), represents

active, poised, and inactive promoters, respectively.

1094

Figure S3. Feature performance comparison: Input features vs CNN mapped features.

Using all samples included in training the vanilla M2A model (NBL, N=6), the individual feature

performance (as determined by Pearson's R^2 between the feature and the response variable,

H3K27ac) for each feature was plotted, comparing the distribution of performances between raw

input training feature and the CNN mapped features at a particular window size (250 bp or

2,500 bp). The best feature from this analysis for each window size and feature type (input or

CNN mapped) was used to determine Pearson's R^2 with H3K27ac from each sample in the NBL

validation set (N=10).

1103

Figure S4. M2A prediction generalizability analysis.

(a) A comparison between performance of the M2A model (R^2 of observed H3K27ac promoter

levels and predicted levels in the test sample) with the surrogate model (represented by the

highest R^2 of observed H3K27ac promoter levels in the test sample and the observed H3K27ac

promoter levels in any training sample). M2A extracts generalizable features capable of out-

performing the surrogate model in both the NBL validation set and the RMS test set, further

highlighted in **(b),** a comparison of surrogate model and M2A model performance using

promoters from DE genes between NBL and RMS.

1112

1113 **Figure S5. M2A with transfer learning outperforms a vanilla M2A model of the same**

1114 **cancer type.**

1115 An M2A model with transfer learning (initially trained with six NBL O-PDX samples and then

1116 transferred with a single RMS sample) consistently outperforms an M2A model trained with a

1117 single RMS sample.

1118

1119 **Figure S6. Signal-to-noise analysis of ENCODE and NBL datasets.**

1120 Comparison at the observed H3K27ac promoter level **(a)** and the H3K4me3 **(b)** promoter levels

1121 revealed different signal-to-noise profiles between the ENCODE dataset and the NBL datasets,

1122 which results in a highly correlated prediction with larger RMSE values.

1123

1124 **Figure S7. M2A ENCODE cohort performance.**

1125 The distribution of M2A prediction performance (R^2), shows that M2A accurately infers both

1126 H3K27ac and H4K4me3 promoter levels in the publicly available ENCODE dataset.

1127

Figure S8. Analysis of outlier H1-ESC.

(a) When inferring H3K27ac promoter levels, M2A was substantially less accurate in the

ENCODE sample H1-ESC, which is an outlier in the ENCODE cohort. **(b, c)** M2A-predicted

H3K27ac promoter levels **(b)** are more consistent with, and more predictive of, H1-ESC gene

expression than are the actual observed H1-ESC H3K27ac promoter levels **(c)**. (**d**) The

hypomethylated region surrounding the promoters of genes *PSMA7* and *SS18L1* (often

indicative of H3K27ac enrichment) showed inconsistent H3K27ac levels between H1-ESC

ChIP-seq replicates from ENCODE.

1136

1137

Figure S9. Predicting gene expression in the ENCODE dataset.

(a–i) The indirect ability of M2A to predict gene expression (i.e., expressed vs. not expressed) on the basis of both M2A-predicted and observed H3K27ac promoter levels for each sample was determined by comparing the AUCs of the receiver operating characteristic (ROC) curves.

Figure S10. Consistency of gene expression and H3K27ac promoter levels in the AML cohort.

The consistency, as determined by Pearson's R^2, between the observed values for gene expression and the H3K27ac promoter levels is remarkably predictive of the performance M2A in predicting H3K27ac promoter levels in samples from the AML cohort.

Figure S11. M2A accurately determines subtype differences between embryonal and alveolar RMS.

(a) The promoter activities of single-promoter, differentially expressed genes in the RMS subtypes ERMS and ARMS are accurately inferred by an M2A base model (trained with six O-PDX NBL samples). **(b)** The predictive performance of M2A is further boosted by transfer learning with one RMS sample. **(c, d)** The performance of M2A declines slightly when the model is applied to all promoters of differentially expressed genes **(c)**, but it recovers when an M2A model with transfer learning with only one RMS training sample is applied **(d)**.

Figure S12. Kaplan–Meier log-rank analysis by mutation status in EWS.

The prognostic ability of **(a)** *TP53* or **(b)** *STAG2* mutation status in the EWS cohort was determined by the log-rank test and visualized using the Kaplan–Meier survival curve. The *STAG2* mutation status showed no significant difference in overall survivability, thus only *TP53*

1162   mutation status was considered when forming the univariate and multivariate Cox proportional

1163   hazards model.

1164

Figure S13. CpG distribution by window relative to the TSS.

To achieve feature input that is informative to the model, M2A window size selection was

partially based on the number of CpGs captured by window size. Each analysis consists of 20

windows surrounding each TSS at a particular window size, representing the theoretical CpG

input to M2A for that particular resolution. Three different window configurations were

considered, comprised of two window sizes: 1) 100 bp and 1,000 bp, 2) 250 bp and 2,500 bp,

and 3) 500 bp and 5,000 bp. Due to NaNs in feature windows calculated with fewer than 2

CpGs, the [100 bp, 1000 bp] model was removed from consideration (> 50% NaNs).

1173

Additional file 2 (file type: .XLSX, 179 KB).

Table S1: H3K27ac active cancer consensus genes in 3 NBL cell lines, and 3 NBL O–PDX

samples.

Table S2: Baseline models vs. vanilla M2A predictive performance comparison.

Table S3: M2A predictive performance in NBL cell line samples.

Table S4: Observed H3K27ac and H3K4me3 ENCODE replicate consistencies.

Table S5: M2A predictive performance in RMS O–PDX samples.

Table S6: M2A RMS transfer model predictive performance in RMS O–PDX samples.

Table S7: M2A predictive performance in ENCODE dataset.

Table S8: M2A predictive performance in AML samples.

Table S9: ERMS vs. ARMS DMRs and associated genes (overexpressed in ERMS).

Table S10: ERMS vs. ARMS DMRs and associated genes (overexpressed in ARMS).

Table S11: M2A alternate promoter usage predictive performance between ARMS and

ERMS samples.

Table S12: M2A predictive performance in EWS samples, before and after transfer.

Table S13: A univariate survival analysis of differential H3K27ac promoter activity

between TP53 mutant and TP53 wild–type EWS tumors.

Table S14: A univariate survival analysis of alternate promoter usage between TP53

mutant and TP53 wild–type EWS tumors.

Table S15: ENCODE H3K27ac replicate consistency with gene expression.

Table S16: Dataset availability.

Table S17: M2A model topologies.

Table S18: Parameter tuning: mean performance in the NBL validation set (R^2)

Table S19: Sample summary information.

(a)

Observed promoter activity landscape of RMS tumors

ARMS
ERMS
Spindle

tSNE-2

tSNE-1

Promoter activity landscape of RMS tumors inferred from M2A

tSNE-2

tSNE-1

Promoter activity landscape of RMS tumors inferred from M2A with transfer

tSNE-2

tSNE-1

(b)

SJRHB013757_X1 (ARMS)

WGBS
Feature (ave)
H3K27ac
H3K4me3

2,500bp

GAS2

SJRHB013758_X2 (ERMS)

WGBS
Feature (ave)
H3K27ac
H3K4me3

GAS2

(c)

SJRHB013757_X1 (ARMS)

WGBS
Feature (ave)
H3K27ac
H3K4me3

2,500bp

NOS1

SJRHB013758_X2 (ERMS)

WGBS
Feature (ave)
H3K27ac
H3K4

NOS1

(d)

GAS2

H3K27ac level

ARMS   ERMS
RMS Subtype

(e)

NOS1-005

H3K27ac level

ARMS   ERMS
RMS Subtype

Observed ARMS
Observed ERMS
Predictions from M2A

(a)

SJRHB010468_X1 (ARMS)

WGBS

Feature
(2,500 bp M-ave)

H3K27ac

H3K4me3

RNA-seq

PDZRN3-006

50 kbp

SJRHB000026_X1 (ERMS)

WGBS

Feature
(2,500 bp M-ave)

H3K27ac

H3K4me3

PDZRN3-001

RNA-seq

PDZRN3

(b)

	hazard ratio		p-value
TP53 mutation	3.01 (1.03 – 8.80)		0.044*
TNS1 (ENST00000446903.1)	0.43 (0.24 – 0.77)		0.004 **
RET (ENST00000479913.1)	0.56 (0.30 – 1.05)		0.073
SLC27A6 (ENST00000508645.1)	1.72 (1.05 – 2.81)		0.031 *

0.2   0.5   1   2   5   10

N=72; #Events: 31; Global p-value (Log-Rank): 1.0375e-05
AIC: 211.96; Concordance Index: 0.77