1 **MethylationToActivity: a deep-learning framework that reveals**

2 **promoter activity landscapes from DNA methylomes in individual**

3 **tumors**

Justin Williams[1], Beisi Xu[2], Daniel Putnam[1], Andrew Thrasher[1] and Xiang Chen[1]

[1]Department of Computational Biology, St. Jude Children's Research Hospital, Memphis,

Tennessee, United States

[2]Center for Applied Bioinformatics, St. Jude Children's Research Hospital, Memphis, Tennessee,

United States

Corresponding author: Xiang Chen

Department of Computational Biology

St. Jude Children's Research Hospital

262 Danny Thomas Place

Mail Stop 1135

Memphis, TN 38105

U.S.A.

Tel: +1 901 595 7074

Fax: +1 901 595 7100

Email: xiang.chen@stjude.org

Abstract

Although genome-wide DNA methylomes have demonstrated their clinical value as reliable

biomarkers for tumor detection, subtyping, and classification, their direct biological impacts at

the individual gene level remain elusive. Here we present MethylationToActivity (M2A), a

machine learning framework that uses convolutional neural networks to infer promoter activities

(H3K4me3 and H3K27ac enrichment) from DNA methylation patterns for individual genes.

Using publicly available datasets in real-world test scenarios, we demonstrate that M2A is highly

accurate and robust in revealing promoter activity landscapes in various pediatric and adult

cancers, including both solid and hematologic malignant neoplasms.

Keywords:

DNA methylation, histone modifications, convolutional neural network, transfer learning

Background

Transcriptional regulation is fundamental to the identity and function of cells. Deregulation of

gene expression is a defining feature of common diseases, including cancers. Promoters, the

regulatory regions surrounding the transcription starting sites (TSSs), integrate signals from

distal enhancers and local histone modifications (HMs) to initiate transcription. Almost half of

human protein-coding genes harbor multiple TSSs; consequently, promoter activities determine

both the level of transcription and the transcript isoforms that are expressed, with the latter

potentially having different translation efficiencies and encoding different protein sequences [1].

Tumors frequently use alternative promoters to increase the isoform diversity [2, 3], to activate

oncogenes that are normally repressed [1–3], and to evade host immune attacks by

immunoediting [3, 4]. Compared to cancers in adults, pediatric tumors harbor fewer mutations [5,

6] and use epigenetic deregulation to promote tumorigenesis and progression [7].

Promoter activities can be determined experimentally through transcriptomic approaches,

such as CAP analysis gene expression (CAGE), or through epigenomic approaches, including

chromatin immunoprecipitation followed by sequencing (ChIP-seq) [8]. Because of transcript

degradation by 5′ RNA exonucleases, ChIP-seq approaches for specific HMs have been the

gold standard for studying promoter activities [9]. However, the scarcity of pediatric tumors, the

limited amounts of fresh starting material available, and the extensive workload involved in

acquiring the promoter activity landscapes constrain their interrogation for individual patient

tumors [10, 11].

DNA methylation (DNAm) is a well-studied, relatively stable, and inheritable epigenetic

regulatory mechanism that involves transferring a methyl group to cytosine (C) to form 5-

methylcytosine (5mC), mostly in the CpG context. In contrast to HMs, DNAm can be accurately

and robustly profiled in various tissues, including archival formalin-fixed, paraffin-embedded

(FFPE) tumor samples, through both array [12, 13] and sequencing [14] platforms; therefore, it

has exceptional applicability to studying epigenetic deregulation in tumors. Consequently,

genome-wide DNAm profiles represent a widely available epigenetic asset for studying

epigenetic abnormalities in primary tumors.

The DNAm pattern is mechanistically connected with transcription factor binding and HMs

[15–22]. It also plays critical roles in establishing the chromatin structure in physiologic and

pathologic conditions [23, 24]. Moreover, recent applications of machine learning to genome-

wide DNAm patterns have demonstrated that DNAm can accurately predict the patterns of

chromatin packaging (A/B compartments, R^2 = 0.50–0.66) [25–27] and can reveal distinct

subgroups with prognostic significance among patients with cancer [28, 29]. Recently,

DNAm signature–based molecular classifiers were shown to improve diagnostic accuracy, as

compared to that of traditional schemes, further demonstrating the critical regulatory roles

of DNAm in tumor development [30, 31]. However, unlike HMs, where established biological

interpretations of various marks have resulted in a general "histone code" hypothesis [32, 33],

the relation between DNAm signatures and their transcriptional regulatory roles is complex and

nonlinear. In many cases, even promoter DNAm may positively and negatively correlate with

gene expression depending on the genomic structure involved in a given tumor [34].

Consequently, with few exceptions (e.g., hypermethylation of the promoters of *RB1*, *CDKN2A*,

and *MGMT*) [35], the contribution of DNAm to the regulation of expression of individual genes

remains largely elusive [36–38]. Recent attempts to use DNAm signatures to account for gene

expression levels have had limited success, with the best model capturing 25%–49% of the

expression variations [39]. Undoubtedly, the lack of interpretability of the DNAm pattern at the

individual gene level has severely hampered our understanding of the biological significance of

DNAm signatures.

To address these challenges, we have developed MethylationToActivity (M2A), a deep-

learning framework. The central hypothesis of M2A is that the complex relation between DNAm

signatures and promoter activities can be captured by incorporating both individual CpG

methylation levels and high-order spatial information from different CpGs in the promoter and

flanking regions. Using a cohort of six pediatric neuroblastoma (NBL) orthoptic patient-derived

xenograft (O-PDX) samples profiled in the Pediatric Cancer Genome Project (PCGP) [40], we

trained the model to predict the enrichment of H3K4me3 and H3K27ac (two HMs critical for

promoter transcriptional activity [41]) for genome-wide annotated promoters. We validated the

predictive accuracy of the model in the remaining NBL samples (N $=$ 10) from the same

cohort. We further confirmed its accuracy and generalizability in diverse tumor types from four

publicly available datasets representing real-world applications, including (1) pediatric

rhabdomyosarcoma (RMS) O-PDX tumors profiled in the Pediatric Cancer Genome Project

(N□=□16) [42]; (2) a set of commonly used cell lines profiled in ENCODE (N□=□9) [43]; (3)

primary acute myeloid leukemias (AMLs) profiled by the BLUEPRINT consortium (N□=□19) [44];

and (4) a large primary Ewing sarcoma (EWS) cohort (N□=□140) [19]. These

applications demonstrate that M2A can accurately reveal promoter activities from DNAm

patterns, which will be of great use not only in functionally interpreting differential DNAm

patterns but also in profiling promoter usage in individual patient tumors. This will facilitate

precision medicine by tailoring treatments based on both genetic variants and epigenetic

deregulations.

Results

Extensive diversity of promoter activity among *MYCN*-amplified NBL cell lines

and O-PDX models

To date, most cancer HM profiling studies have made use of tumor models, including cell

lines, xenografts, and more recently, organoids. Technical limitations and challenges when

working with human tumor tissues prevent the generation of high-quality ChIP-seq profiles for

primary patient specimens [45]. Despite the documented epigenetic heterogeneity [46], a

common practice in deciphering major HM deregulations in various cancers is to extrapolate the

epigenetic profiles from related cancer models. Many studies [40, 42, 47–50] have compared

model systems to primary tumors with respect to characteristics such as mutations, gene

expression, and DNAm signatures. In this study, we began by evaluating the level of promoter

activity diversity in closely related NBL models. Specifically, we evaluated promoter activity, as

measured by the H3K27ac level, in three O-PDX models (SJNBL046, SJNBL108, and

SJNBL013763) and three cell line models (IMR-32, NB-5, and SKNBE2) that harbor *MYCN*

amplification with no other major oncogenic mutations. All samples displayed a bimodal

distribution of promoter H3K27ac levels across the genome (Additional file 1: Figure S1), and O-

PDX models had a marginally higher fraction of active promoters (mean: 31.9%, range: 27.6%–

36.1%) than did cell line models (mean: 26.1%, range: 25.6%–26.7%) ($P = 0.14$, Student's *t*-

test). However, there were extensive variations in the promoter activities in both the cell line

models (Figures 1a and 1d) and the O-PDX models (Figures 1b and 1e). Moreover, greater

divergence was observed between a cell line model and an O-PDX model (mean: 34.9%, range:

29.9%–39.0%) than between two cell line models (mean: 31.0%, range: 29.2%–32.1%;

$P = 0.44$, Student's *t*-test) or between two O-PDX models (mean: 31.0%, range: 22.9%–

37.0%; $P = 0.02$, Student's *t*-test) (Figure 1f). Variations in promoter activity may play a

significant role in the transcriptional deregulation of individual tumors, as a substantial fraction of

established cancer consensus genes (22.4% in O-PDX models and 31.1% in cell line models,

including *APOBEC3B*, *TGFBR2*, *PAX7*, *HOXA11*, *PDCD1LG2*, *PTK6*, *BCL11B*, *FAS*, and *MYC*;

(Additional file 2: Table S1) displayed heterogeneous promoter activities in the surveyed tumor

models. Therefore, we sought to develop a computational approach to infer the promoter activity

landscape for individual tumors.

M2A: a deep-learning framework to reveal promoter activities from DNA

methylation

DNAm plays a critical role in determining the framework of gene expression for a given

cell/cellular state. However, the highly complex and non-linear relations between DNAm

patterns and HMs severely hamper the interpretability of the biological impact of differential

DNAm patterns. We hypothesize that high-level DNAm features that capture the spatial

information from different CpGs in the promoter and regions in its vicinity provide an opportunity

to infer promoter activities accurately. We propose to use a convolutional neural network

(CNN)–based deep-learning framework to extract such features.

The M2A conceptual workflow is shown in Figure 2. M2A starts with raw DNAm feature

extraction from around individual TSSs (Figure 2a). This is followed by high-level feature

extraction through the CNN layers and mapping between the generalized feature and the final

output (i.e., the H3K4me3 and H3K27ac of the promoter) in the fully connected (FC) layers

(Figure 2b). The baseline model described in this report was trained on six NBL PDX tumors

(SJNBL046_X, SJNBL013761_X1, SJNBL012401_X1, SJNBL013762_X1, SJNBL013763_X1,

and SJNBL015724_X1) for which comprehensive genomic and epigenomic profiling data are

available, including the results of whole-genome sequencing, whole-exome sequencing, RNA

sequencing, whole-genome bisulfite sequencing (WGBS), and ChIP-seq of eight histone marks

(H3K4me1, H3K4me2, H3K4me3, H3K27me3, H3K27ac, H3K36me3, H3K9/14ac, and

H3K9me3), CTCF, BRD4, and RNA polymerase II (PolII).

156   M2A produces a highly accurate landscape of promoter activity in pediatric NBL

To evaluate the performance of M2A, we first explored its performance in the remaining NBL

samples in the cohort, including one O-PDX tumor, one primary autopsy tumor, and eight cell

lines. Using this test group, we compared the performance of a traditional artificial neural

network (ANN) model consisting of only two FC layers to that of a deep-learning framework with

three CNN layers and two FC layers (Figure 2b). From a qualitative perspective, M2A correctly

revealed the bimodal distribution of the promoter activities for both H3K4me3 and H3K27ac in

all samples, and from a quantitative perspective, the inferred genome-wide promoter activity

landscape was highly accurate for individual samples for both H3K4me3 (R^2 = 0.937 ± 0.019;

RMSE = 0.596 ± 0.083) (Figures 3a and 3d) and H3K27ac (R^2 = 0.797 ± 0.047; RMSE =

0.644 ± 0.071) (Figures 3b, and 3c). Moreover, the addition of CNN layers was merited, as

there was a decrease in the prediction error (measured as $1 - R^2$) by 12.0% (P = 0.002,

Wilcoxon signed-rank test) and 12.2% ($P = 0.0039$, Wilcoxon signed-rank test) for the model

topologies for H3K4me3 and H3K27ac, respectively (Additional file 2: Table S2).

Our analysis of *MYCN*-amplified NBL cell line and O-PDX models has revealed substantial

variations in their promoter activities, which is a potential caveat to the practice of representing

primary tumor epigenomes by a few profiled models. Conversely, M2A produced

highly accurate promoter activity landscapes, significantly outperforming the observed

consistency between training and testing samples for both H3K4me3 ($R^2 = 0.891 \pm 0.023$,

$P = 1.5 \times 10^{-5}$, Wilcoxon rank-sum test) and H3K27ac ($R^2 = 0.720 \pm 0.045$,

$P = 9.5 \times 10^{-5}$, Wilcoxon rank-sum test; Additional file 2: Table S3). Remarkably, in eight (of

10) test samples, the accuracy of the M2A-inferred promoter H3K27ac activity was better than

the highest similarity attained by any individual training sample ($P = 0.048$, Wilcoxon signed-

rank test). The same pattern was observed for H3K4me3 levels, with M2A being more accurate

for nine of 10 samples ($P = 0.013$, Wilcoxon signed-rank test), demonstrating the accuracy of

M2A in revealing individual tumor promoter activity landscapes. Finally, the predictive accuracy

of M2A was comparable to the experimental consistency observed between replicates from the

same cell lines profiled in ENCODE for H3K4me3 ($R^2 = 0.937 \pm 0.019$ for

M2A vs. 0.922 ± 0.056 for ENCODE replicates [$N = 25$]; $P = 0.90$, Wilcoxon rank-sum test)

(Figure 3a; Additional file 2: Table S4). The accuracy of M2A also approached the replicate

consistency for H3K27ac ($R^2 = 0.797 \pm 0.047$ for M2A vs. 0.849 ± 0.047 for ENCODE

replicates [$N = 26$]; $P = 0.0088$, Wilcoxon rank-sum test) (Figure 3b: Additional file 2: Table

S4). Measurement of the root mean square error (RMSE) revealed a similar pattern (Additional

file 2: Table S4).

M2A is generalizable and scalable

Aside from the model accuracy, there are two additional requirements with practical importance

for deploying a machine learning model (such as M2A) in real-world applications: (1)

generalizability, i.e., M2A needs to achieve a similar performance with a set of unseen test

samples, including tumor/tissue types not used in the model training; and (2)

scalability, i.e., M2A must be able to be applied efficiently to external data.

We first demonstrated the accuracy, generalizability, and scalability of M2A by

using test samples from rhabdomyosarcoma (RMS) O-PDX tumors. The RMS O-PDX dataset

consists of 16 pediatric RMS tumors (11 embryonal, four alveolar, and one spindle subtype,

termed ERMS, ARMS, and spindle subtypes, respectively). As with the NBL cohort, each RMS

sample was extensively profiled, including by WGBS, RNA-seq, and ChIP-seq of H3K4me3 and

H3K27ac. Using the baseline model (the 3CNN-FC model trained on the six NBL PDX

samples), M2A achieved an overall predictive accuracy with the RMS dataset that was

comparable to that of the NBL test group for both H3K4me3 (R^2 = 0.938 ± 0.018, P = 0.78;

RMSE = 0.643 ± 0.125, P = 0.45, Wilcoxon rank-sum test) (Figures 3a and 3f; Additional file 2:

Table S5) and H3K27ac (R^2 = 0.790 ± 0.038, P = 0.78; RMSE = 0.590 ± 0.093, P = 0.06,

Wilcoxon rank-sum test) (Figures 3b and 3e; Additional file 2: Table S5), which was comparable

to or significantly outperformed the observed similarities between two different RMS tumors for

H3K4me3 (R^2 = 0.917 ± 0.028, P = 0.0011; RMSE = 0.646 ± 0.134, P = 0.69, Wilcoxon

rank-sum test) and H3K27ac (R^2 = 0.780 ± 0.028, P = 0.42; RMSE = 0.550 ± 0.096, P =

0.15, Wilcoxon rank-sum test). The accuracy of the inferred H3K4me3 activity was comparable

to the inter-replicate consistency of the ENCODE samples (P = 0.95, Wilcoxon rank-sum test).

By definition, generalizability can be achieved only in the absence of over-fitting (or

"memorization" of the training data). Neural networks often fall victim to this problem through a

combination of factors, including relatively small training datasets and/or over-parameterization.

The relatively consistent expression of housekeeping genes across different tissues may lead to

an inaccurate (often inflated) interpretation of the performance measurement in such a model,

as evidenced by the relatively high R^2 value (0.663 ± 0.040) between the promoter H3K27ac

level of a random RMS test tumor and the most similar NBL training tumor (Additional file 1:

Figure S2a). Therefore, we focused on the set of genes that are differentially expressed (DE) in

RMS and NBL PDX samples [49], for which an over-fitted or memorized model would perform

poorly. Not surprisingly, the average correlative consistency between the NBL validation

samples and the most similar NBL training sample dropped from 0.755 to 0.600 when the

measurement was restricted to promoters encoding the DE genes (Additional file 1: Figure S2b),

whereas a sharp decline (from 0.663 to 0.264) was also observed for RMS test tumors

(Additional file 1: Figure S2b). Conversely, the six-PDX NBL-trained M2A model maintained

high accuracy for promoters of DE genes in both the NBL validation set (R^2 = 0.724 ± 0.066)

and the RMS test set (R^2 = 0.718 ± 0.046) (Additional file 1: Figure S2b), further

demonstrating the generalizability of M2A.

M2A is efficient and scalable. On average, the training of the baseline M2A model (with six

NBL O-PDX tumors) takes approximately 16 min (using a Tesla P100-16GB GPU). The feature

extraction and promoter activity prediction from WGBS data (as a genome-wide DNAm level file

in a tab-delimited text format) can be executed on a personal computer (in this case, we used a

MacBook Air with a 2.2-GHz Intel Core i7 and 8-GB 1600-MHz DDR3 RAM) and takes 15–19

min.

Transfer learning further improves the performance of M2A with minimal

additional input in the target domain.

Although we have demonstrated the generalizability of M2A in the RMS dataset, the fact that

epigenetic genes are frequently mutated in pediatric tumors [7] raises the possibility that

individual tumor types carry a type-specific interpretation of the DNAm patterns. When ChIP-seq

measurement is available for sufficient samples, a type-specific model is desirable. However,

although pediatric solid tumors as a group constitute a rare disease, they comprise many

different tumor types, and it is rare to have sufficiently profiled samples available for many of

them. In addressing this challenge, we hypothesize that a fixed feature-extraction strategy

(transfer learning) can achieve the goal of deriving an efficient tumor type–specific model by

using a small labeled dataset. A primary assumption here is that generalized features extracted

based on a large dataset are similarly informative for apparently different tasks. The feature

learning and selection characteristics of CNNs provide exceptional portability in various tasks

with extremely small labeled datasets.

In M2A, the CNN layers capture generalized DNAm features and the FC layers learn the

mapping function between the DNAm features and the promoter activities. Here we start with

the pretrained M2A baseline model, fix the feature-extraction layers (CNN layers), and use a

single sample from the target tumor type to update the mapping function (the weights and

biases of the FC layers). Because the consistency of M2A for H3K4me3 approached the inter-

replicate consistency in both NBL and RMS datasets, we focused on H3K27ac inference for

transfer learning. Upon performing transfer learning with a single sample in the RMS dataset,

we observed significantly improved accuracy (R^2 = 0.814 ± 0.037, P = 9.2 × 10^{-5},

Wilcoxon signed-rank test) (Figures 3b, 3g, and 3h; Additional file 2: Table S5). Moreover, this

model significantly outperformed a single RMS sample model with the identical model

architecture, in which both the CNN layer and the FC layers were derived from the RMS training

sample (P = 0.0042, Wilcoxon signed-rank test) (Additional file 1: Figure S3; Additional file 2:

Table S6) and from the observed similarities between different RMS tumors (P = 0.042,

Wilcoxon rank-sum test). This analysis demonstrated the value of both the pretrained CNN

layers for general feature extraction and a single profiled sample in the target domain.

Consequently, we applied transfer learning to both the EWS and AML datasets. However,

transfer was not feasible in the ENCODE dataset because those cell lines were derived from

different tissues.

M2A accurately reveals promoter activity landscapes in adult tumors and in

hematologic malignant neoplasms

We next evaluated the performance of M2A in independently collected datasets, including ones

for adult tumors and hematologic malignant neoplasms. Upon analyzing nine ENCODE cell

lines, we found that differences in antibody usage and experimental protocols between the

ENCODE and NBL datasets resulted in different signal-to-noise profiles (Additional file 1: Figure

S4), with higher RMSE values between the model predictions and the actual observations

(H3K4me3: 1.013 ± 0.273, $P = 4.1 \times 10^{-4}$; H3K27ac: 0.966 ± 0.226, $P = 2.6 \times 10^{-4}$,

Wilcoxon rank-sum test). However, the predicted activities remained highly correlated with the

experimental measurements (H3K4me3: $R^2 = 0.900 ± 0.024$; H3K27ac: $R^2 = 0.666 ± 0.152$)

(Additional file 1: Figure S5). Although the accuracy of H3K4me3 was relatively uniform, H1-

ESC was an outlier with a substantially lower accuracy of H3K27ac inference (by the median-

based method [51]) (Additional file 2: Table S4; Additional file 1; Figure S6a). An investigation of

the promoter activity (the measured and inferred H3K27ac levels) and the measured gene

expression in H1-ESC revealed that a subset of actively transcribed genes showed little or no

H3K27ac levels in their promoters, where M2A inferred relatively strong promoter activity

(Additional file 1: Figure S6b). Consequently, the inference of promoter activity by M2A

outperformed the actual measurement in terms of both the quantitative consistency with the

gene expression level ($R^2 = 0.586$ for the M2A-inferred H3K27ac level vs. 0.460 for the

measured H3K27ac level) (Additional file1 : Figures S6b and S6c) and the accuracy in

predicting expressed genes (AUC = 0.919 for the M2A-inferred H3K27ac level and 0.880 for

observed H3K27ac level) (Additional file 1: Figure S7d). We also observed a small fraction of

inferred active promoters without strong expression; these may represent genes subject to

transcriptional pausing (where transcription is initiated but there is no elongation), a distinctive

feature of undifferentiated stem cells [52]. Similarly, better consistency with gene expression

levels was observed in two extra samples with low accuracies of H3K27ac inference (SK-N-SH

and GM23248) (Additional file 2: Table S7).

We further evaluated the performance of M2A in revealing promoter activities in 19 acute

myeloid leukemia (AML) primary patient samples collected by the BLUEPRINT consortium,

which is part of the International Human Epigenome Consortium (IHEC) [44]. Analyses of

observed promoter activity (measured by H3K27ac level) and gene expression (measured in

fragments per kilobase of transcript per million mapped reads [FPKM]) in the same sample

revealed non-uniform qualities with a wide range of consistency (mean R^2 = 0.448, range:

0.121–0.603) (Additional file 2: Table S8). Similarly, the M2A-inferred promoter activity

landscape displayed substantial variability with respect to the observed activities among these

samples (mean R^2 = 0.471, range: 0.027–0.730). Strikingly, the consistency between the

observed promoter activity and gene expression was highly predictive of the performance of

M2A with individual samples (R^2 = 0.974, P = 6.4 ± 10^{-15}, Pearson's correlation test)

(Additional file 1: Figure S8). Finally, although gene expression was not used in model

generation with M2A (for the baseline model or the transfer learning step), the promoter activity

inferred by M2A showed significantly greater and uniform consistency with gene expression

(mean R^2 = 0.568, range: 0.539–0.593; P = 0.0048, Wilcoxon signed-rank test) (Additional file

2: Table S8). These results jointly suggest that the ChIP-seq library quality is a potential

confounding factor for both the consistency between the observed (from ChIP-seq) and the

M2A-inferred promoter activity landscapes and the consistency between the observed promoter

activity and gene expression in these samples.

Promoter activity landscape inferred by M2A faithfully recapitulates the subtype

difference between embryonal and alveolar rhabdomyosarcomas

Identifying recurrent epigenetic deregulations (epi-drivers) is a primary research focus in cancer

epigenome studies [53]. To this end, we investigated whether subtype-specific epigenetic

deregulation was captured in M2A-revealed promoter landscapes in the RMS O-PDX tumors. A

t-SNE embedding using M2A-inferred promoter activity landscapes showed clear separation of

ARMS and ERMS tumors (Figure 4a). Importantly, when focusing on the promoters of DE

genes in the ARMS and ERMS subtypes, the baseline M2A model faithfully recapitulated the

subtype-specific promoter activity patterns (R^2 = 0.730 for DE genes with a single annotated

promoter; 0.617 when all annotated promoters for DE genes were included) (Additional file 1:

Figures S9a and S9c). Transfer learning using data from a single RMS further improved the

consistency ($R^2 = 0.767$ and 0.672, respectively) (Additional file 1: Figures S9b and S9d).

GAS2 is a gene selectively expressed in ERMS [42]. Although promoter hypomethylation

was found in the ARMS tumors, the M2A model correctly predicted significantly stronger

promoter activities in ERMS tumors ($P = 0.01$, Wilcoxon rank-sum test) (Figures 4b and 4d).

Similarly, although both ERMS and ARMS tumors had *NOS1-005* promoter hypomethylation,

strong promoter activity was predicted in ARMS tumors only ($P = 0.0015$, Wilcoxon rank-sum

test) (Figures 4c and 4e), consistent with the ChIP-seq measurement.

M2A reveals the contribution of differentially methylated regions to promoter

activities

Although DNA methylation patterns, including differentially methylated regions (DMRs), are well-

established biomarkers for diverse diseases and have revealed molecularly and clinically

different subtypes for many cancers, their functional importance in individual gene regulation is

less clear [36–38]. For example, many cancer-specific CpG island hypermethylation regions

occur in genes that normally are not expressed or are expressed at only a low level [54].

Moreover, even in genes that are both differentially expressed and differentially methylated in

different subtypes, up-regulated samples can be associated with hypomethylation,

hypermethylation, or both [34, 42]. The observed nonlinear relation between DNA methylation

and gene expression complicated the functional interpretation of specific DMRs. In this analysis,

we interpret the functional roles of DMRs based on the promoter activities of their associated

DE genes.

To summarize unambiguously the contribution of DMRs to differential promoter activities, we

focused on 371 genes in ERMS and ARMS that have a single annotated promoter and that are

both differentially expressed (197 are over-expressed in ERMS, 172 are over-expressed in

ARMS) and differentially methylated (169 are hypomethylated, 128 are hypermethylated, and 74

have both hypomethylation and hypermethylation) in the two major RMS subtypes (Additional

file 2: Tables S9 and S10) [42]. Among these genes, 140 promoters showed significantly higher

H3K27ac measurements in the over-expressed subtype (FDR < 0.1, Wilcoxon rank-sum test),

whereas 152 promoters had measurements that were significantly higher when the DNAm-

based H3K27ac activity was measured. These 152 promoters included 100 of the 140

promoters identified using the observed signals. These results suggest that M2A can reveal the

role of DMRs in modulating the promoter activities of affected genes in a context-specific

manner.

M2A identifies subtype-specific promoter usage encoding different protein

isoforms in rhabdomyosarcoma

Alternative promoter usage is an important pretranslational mechanism for tissue-specific

regulation as it affects the diversity of isoforms available. Recently, light was shed on the

pervasiveness of alternative promoter usage in cancer; in some cases, promoter usage is a

more accurate reflection of patient survival than is gene expression [2]. Among 10,835 active

genes with multiple annotated promoters in the RMS dataset, we found 2,584 genes (24%) with

alternative primary promoter usage among 16 samples. We focused on 562 genes that 1) were

active in both the ERMS and ARMS subtypes and 2) had subtype-specific promoter usage. We

explored the accuracy of M2A in predicting alternative promoter usage in ARMS and ERMS

(Additional file 2: Table S11).

Based on measured promoter activities, 389 genes exhibited significant usage difference

between the two subtypes (FDR < 0.1, Wilcoxon rank-sum test [used as the ground truth]). The

M2A-inferred promoter activity landscape revealed 236 genes for which there was a significant

difference in promoter usage between the subtypes (FDR < 0.1, Wilcoxon rank-sum test), and

196 of them matched the ground truth (precision = 0.83, recall = 0.50, F1 score = 0.63)

(Additional file 2: Table S11).

PDZ Domain Containing Ring Finger 3 (*PDZRN3*) is a known target of the PAX3/7–FOXO1

fusion protein [55–57], which blocks terminal differentiation in myogenesis [58]. M2A predicted a

subtype-specific promoter usage pattern in *PDZRN3*. Functional studies have shown that

PDZRN3 regulates myoblast differentiation into myotubes through transcriptional and

posttranslational regulation of Id2 [58]. Its over-expression in ARMS was primarily driven by the

fusion protein binding adjacent to an alternative promoter (*PDZRN3-205*) located 191 kbp

downstream of the canonical promoter (*PDZRN3-201*) (Figure 5a). The subtype-specific isoform

usage is accompanied by DMRs of the alternative promoter and its immediate downstream

regions and is further confirmed by RNA-seq read alignment (Figure 5a). Compared to the

canonical isoform expressed in ERMS, the ARMS-preferred *PDZRN3-205* isoform lacks the

RING-finger and Sina domains in the N-terminus and harbors a shorter PDZ domain. The

isoform difference, as well as the differences in expression level, between subtypes may

contribute to the impairment of myogenesis at different stages in the development of ARMS and

ERMS tumors.

M2A identifies alternative promoter usages with potential prognostic values in

Ewing sarcoma

We next examined the predicted epigenetic promoter activities in 140 EWS samples with DNAm

data assayed by reduced representation bisulfite sequencing (RRBS) [19]. Three of these

samples had matching ChIP-seq profiles for H3K4me3 and H3K27ac. To interrogate this

dataset, we applied the pretrained baseline model with transfer learning, as detailed above, to

recalibrate the weights mapping the high-level features to the promoter activities in the EWS

cohort. Despite the difference in DNAm platforms (WGBS for the NBL training model and RRBS

for the EWS samples), the inferred promoter activity landscape was accurate (R^2 = 0.721, 0.634,

and 0.705 for the three samples with HM profiles, using leave-one-out prediction) (Additional file

2: Table S12).

Ewing sarcomas with mutations in *TP53* and *STAG2* have a particularly dismal prognosis

[59]. We explored whether the promoter activity had additional prognostic value in 72 samples

for which survival data was available. Because of the limited sample size, the initial analysis

revealed a significant association between poor clinical outcomes and *TP53* mutations

($P = 0.00047$, log-rank test) (Additional file 1: Figure S10a) but not *STAG2* mutations

($P = 0.67$, log-rank test) (Additional file 1: Figure S10b). We identified 20 active genes that

showed a potential difference (absolute mean difference of log-scaled activity ≥ 1) between

TP53 mutant tumors and wild-type tumors, and we applied Cox proportional hazards models to

evaluate their potential contributions to patient survival that are independent of *TP53* mutation

status (Additional file 2: Table S13). We performed the same analysis for 24 genes with

potential different promoter usage in tumors with and without *TP53* mutations (Additional file 2:

Table S14). Finally, we derived a multivariate Cox proportional hazards model including *TP53*

mutation status and one candidate gene with differential promoter activity (*CLDN16*) and four

candidate gene isoforms (*CASZ1*, ENST00000496432; *RET*, ENST00000479913; *TEX40*,

ENST00000535981; and ENST00000328404), and we followed this with backwards stepwise

model selection. The final model (Figure 5b) revealed potential protective roles for three

candidate transcripts (*CASZ1*, ENST00000496432; *RET*, ENST00000479913; and *TEX40*,

ENST00000535981). *CASZ1* encodes a zinc-finger transcription factor, and its down-regulation

is associated with poor prognosis in NBL [60], hepatocellular carcinoma [61], and clear-cell

renal cell carcinoma [62]. Our results suggest that *CASZ1* is a new prognostic indicator for EWS.

Further studies are needed to draw more attention to the functional roles of these

genes/transcripts in EWS progression.

Discussion

Although epigenetic studies in disease models (cell lines, xenografts, and organoids) and in a

limited number of primary tumor samples have demonstrated the oncogenic contributions of

epigenetic deregulations to cancer initiation, progression, and response to treatment [46, 63, 64],

genome-wide profiling of promoter activities by using standard approaches (e.g., ChIP-seq or

CAGE) has not been carried out in large patient tumor cohorts, despite the continuous efforts of

large epigenome consortia [43, 44, 65]. Our analyses of *MYCN*-amplified NBL tumors revealed

both commonly active promoters and promoters that were active in some tumors but not in

others. Moreover, these sample-specific active promoters are functionally important, as they

drive the expression of several cancer consensus genes, including *MYC*. This observation is

consistent with recent reports of heterogeneous enhancer activities of cell line–defined super-

enhancers in primary gastric cancers [66], emphasizing the critical importance of deriving

sample-specific epigenomic signatures. To bridge the gap between the extensive epigenomic

resources in disease models and the limited ChIP-seq profiles of primary patient tumors, we

developed MethylationToActivity (M2A), a deep-learning framework, to characterize the

promoter activity landscape (both H3K4me3 and H3K27ac levels) in individual tumors by using

DNAm data that is widely available for patient tumors. M2A demonstrated excellent

performance across various tumor types, with accuracy comparable to that of ChIP-seq

measurements of replicate samples from high-quality cohorts (Figure 3).

Although our framework was strictly trained on HM levels, the inferred promoter activity was

highly correlated with the transcript-based gene expression levels quantified by RNA-seq

(Additional file 2: Tables S5, S7 and S8). The correlation between gene expression and inferred

promoter activities (mean R^2 = 0.618 for ENCODE data, 0.622 for K562, 0.638 for GM12878,

and 0.586 for H1-ESC) surpassed that with the state-of-art model binomial distribution probit

regression [BPR] model [39], which was developed for predicting gene expression levels from

DNAm patterns (the best reported R^2 values were 0.49 for K562, 0.37 for GM12878, and 0.25

for H1-ESC). Strikingly, the (indirect) predictive accuracy of M2A for gene expression across

nine ENCODE cell lines (average R^2 = 0.618) was comparable to the predictive accuracy of a

model built on 11 HMs, one histone variant, and DNase I hypersensitivity [67]. Similarly, the

accuracy of binary prediction of expressed genes (average AUC = 0.933 and 0.919 for the

ENCODE and AML datasets, respectively) surpassed that of DeepChrome (average AUC =

0.80), a state-of-art deep-learning algorithm trained to predict expressed genes by using five

core HMs (H3K4me3, H3K4me1, H3K36me3, H3K9me3, and H3K27me3) [68]. These results

further validated our framework. Finally, both the M2A and BRP models suggest that it is

insufficient to represent DNAm information by using a simple average methylation level in the

promoter region. To properly reveal the regulatory roles of DNAm, we need to derive high-order

features that capture spatial relations among different CpG probes in promoters and in their

vicinity. M2A uses the feature learning and selection characteristics of CNNs to achieve its

exceptional performance, thus demonstrating the rich information content of DNAm signatures

at both the genome-wide and local gene levels.

Analysis of the deep-learning framework revealed that M2A derives 1) high-level features

from DNAm patterns that are common among different tumors and 2) tumor subtype–specific

mapping functions from the mapping of high-level DNAm features to promoter activities in

individual tumor subtypes by using transfer learning (when feasible). Although our deep-learning

model cannot establish a causal relation between DNAm and promoter activities, these findings

nevertheless shed light on both the general and tumor subtype–specific rules for interpreting

DNAm patterns.

In evaluating our predictions, we found that several samples (the "poor performers") showed

abnormally low predictive accuracy with both the ENCODE and BLUEPRINT datasets.

Investigations of the promoter H3K27ac levels revealed that the fraction of active promoters in

these samples was substantially lower than in other samples. Furthermore, joint analyses with

RNA-seq data from the matching samples indicated that 1) in contrast to the predictive accuracy

of H3K27ac levels, the "poor performers" achieved comparable consistency between the M2A-

inferred promoter activity and gene expression; 2) the "poor performers" showed significantly

lower correlation between the measured promoter H3K27ac level and gene expression; and 3)

the correlation between the measured promoter H3K27ac level and gene expression was highly

predictive of the accuracy of the H3K27ac prediction (Additional file 1: Figure S8). Collectively,

these results suggested that the "poor performers" could reflect the quality of the ChIP-seq

results included in the test data. This observation emphasizes the value of conducting a

preliminary analysis to evaluate the data quality before incorporating public data. It also

suggests that M2A can provide a robust surrogate for promoter activities when the ChIP-seq

experimental data is questionable.

Alternative promoter usage increases the transcriptomic diversity during normal tissue

development and oncogenesis. Recent work demonstrated that an alternative promoter of

ERBB2 is predictive of a poor clinical outcome but that the canonical promoter shows no

significant association with survival in patients with low-grade glioma [2]. Whereas earlier

studies focused on identifying alternative promoter usage through CAGE or RNA-seq data, our

research has shown that alternative promoter usage can be extensively studied by using DNAm

profiles from diverse samples, including retrospective FFPE tumor samples, for which traditional

approaches (CAGE, ChIP-seq, and RNA-seq) are technically challenging. Our analysis of a

large EWS cohort (without matching RNA-seq data) revealed promoter activities for several

specific isoforms that are independently associated with better clinical outcomes, including a

specific isoform of the *CASZ1* gene (ENST00000496432). This demonstrates the critical

importance of analyzing alternative promoter usage in epigenomic studies, because the activity

of the primary *CASZ1* promoter (ENST00000344008) neither differed substantially between

patients with different *TP53* mutation status nor suggested a significant association with clinical

outcome beyond *TP53* mutation status. Moreover, all three specific isoforms retained in the final

model are transcripts of the intron retention class (*CASZ1*, ENST00000496432; *RET*,

ENST00000479913; and *TEX40*, ENST00000535981), suggesting another layer of gene

regulation [69].

Finally, our analysis quantitatively emphasizes that promoter activity is one of the

mechanisms that regulate the final transcriptional output: 1) both the observed and predicted

promoter activities account for 50%–60% of the variation in gene expression, and 2)

approximately 40% of the genes differentially expressed in ERMS and ARMS tumors show

significant differences in their promoter activities. In addition to promoter activities, other

epigenetic mechanisms, including enhancer activities, contribute substantially to gene regulation

[70]. Recent work has demonstrated the roles of DNAm in regulating enhancer activities [71]

and aberrant cancer-specific DNAm patterns in super-enhancers [72]. Consequently, we aim to

expand our M2A framework to infer enhancer activities from DNAm patterns in the future.

Conclusion

We have demonstrated that MethylationToActivity overcomes the unique challenges of

systematically characterizing promoter activities from DNA methylation signatures. It achieved

an accurate, robust and generalizable performance in various pediatric and adult cancers,

including both solid and hematologic malignant neoplasms. MethylationToActivity will serve as a

valuable tool to provide functional interpretation of DNAm deregulation, characterize promoter

activity differences from DNAm patterns, and reveal alternate promoter usage in patient tumors,

which will facilitate precision medicine by tailoring treatments based on both genetic variants

and epigenetic deregulation.

Methods

Datasets

Five separate publicly available datasets were used in this study, including a pediatric NBL O-

PDX dataset ($N = 16$) [40]; an RMS O-PDX dataset ($N = 16$) [42]; ENCODE datasets with

matching H3K27ac and H3K4me3 histone mark ChIP-seq, RNA-seq, and WGBS experimental

data ($N = 9$) [43]; a DCC BLUEPRINT AML dataset ($N = 19$) [44]; and a pediatric EWS

dataset ($N = 140$) [19] (Additional file 2: Table S15). Of the 140 samples in the EWS cohort,

only three had matching ChIP-seq and reduced-representation bisulfite sequencing (RRBS)

data available; for the remaining 137 samples, only RRBS data was available. All other cohort

datasets (i.e., the RMS, NBL, ENCODE, and AML datasets) contained matching H3K27ac and

H3K4me3 profiles, along with RNA-seq and WGBS experimental data.

Feature processing

All datasets were evaluated using GENCODE annotation definitions (www.gencodegenes.org/);

the NBL, RMS, ENCODE, and EWS datasets were evaluated using GENECODE GRCh37.p13

(release 19), and the AML dataset was evaluated using GENCODE GRCh38.p13 (release 32).

Promoter regions are defined as the TSS ± 1 kbp, where the TSS is defined as each unique

transcript start position. To avoid using identical or near-identical promoter regions, only TSSs

with promoter regions with less than 50% overlap were considered. Gene orientation was taken

into account, and any promoters with overlying regions but opposite orientations were not

considered as overlapping. Because of differences in sex amongst samples, all chromosome X

and Y promoter regions were removed from consideration. This resulted in a total of 96,756 and

104,722 non-overlapping promoter regions from the annotation files GRCh37.p13 and

GRCh38.p13, respectively.

M2A uses only one variable feature type: DNA methylation. For WGBS data, the M-value was

calculated as follows:

$$Mval_k = log_2 \left(\frac{Methylated_k + \alpha}{Unmethylated_k + \alpha} \right),$$

where $Methylated_k$ and $Unmethylated_k$ correspond to the number of methylated and unmethylated

reads of the k^{th} CpG site, respectively. By default, the offset α was set to 0.5, and a global M-value

threshold was set to a maximum value of $\log_2(65)$ and a minimum value of $\log_2(\frac{1}{65})$. CpG sites with

coverage of less than five reads were removed.

M2A uses a promoter region–based windowed approach, comprising 20 windows of two sizes (250

bp and 2.5 kbp) and a step size equal to the window size, centered on a given TSS. For instance,

$\boldsymbol{W}_{ij} = \{W_{i1}, W_{i2}, W_{i3} \dots, W_{i20}\}$ is the vector of windows where $\boldsymbol{i}$ represents a particular TSS and $\boldsymbol{j}$

represents a particular window corresponding with the $\boldsymbol{i^{th}}$ TSS. This means that W_{i10} and W_{i11}

represent the windows immediately downstream and upstream of the $\boldsymbol{i^{th}}$ TSS. Therefore,

$$W_{i1} = TSS_i - 10^{th} \; window,$$

567    $$W_{i2} = TSS_i - 9^{th} \; window, \dots$$

$$W_{i19} = TSS_i + 9^{th} \; window,$$

$$W_{i20} = TSS_i + 10^{th} \; window$$

571   Each feature was calculated in this manner. The DNA methylation features, including the windowed

572   M-value mean, variance, and sum of squared differences (SSD), and the fraction of the SSD of M-

573   values (FSSD), were calculated and represented by the feature vectors $\boldsymbol{Mave}_i$, $\boldsymbol{Mvar}_i$, $\boldsymbol{Mssd}_i$, and

574    $\boldsymbol{Mfssd}_i$. Therefore, the features for a particular window, denoted as W_{ij}, would be calculated as follows:

$$Mave_{ij} = \frac{1}{n_{ij}} \sum Mval_{ik(j)},$$

576    $$Mssd_{ij} = \sum (Mave_{ij} - Mval_{ik(j)})^2,$$

$$Mvar_{ij} = \frac{1}{n_{ij}} \sum (Mave_{ij} - Mval_{ik(j)})^2,$$

578   and

$$Mfssd_{ij} = \frac{Mssd_{ij}}{\sum (Mave_i - Mval_k)^2},$$

580   where

$$Mave_i = \frac{1}{n_i} \sum Mval_k.$$

Here, i represents the promoter, j represents a specific window for a particular promoter, and

$Mval_k$ represents the Mval for individual CpGs in a region where $Mval_{k(j)}$ is the Mval for an individual

CpG in a specific window. Each feature was interleaved by window size to provide model input wherein

each window contained a number of "channels" equal to the number of features, with the feature array

shape being (*N*, 2, 20, 4), where *N* represents the total number of TSSs in a sample, 2 represents the

number of window sizes (250 bp and 2.5 kbp), 20 represents the number of windows, and 4 represents

the number of features per window. All features were scaled from 0.1 to 1 (using MinMaxScaler with

default values from sklearn version 0.22); in instances where windows overlapped regions without

methylation data, resulting in NaNs (such as chromosomal boundaries, telomeric regions, and

centromeric regions), these feature values were marked as 0.

Calculating histone modification enrichment

The response variable was calculated for each non-overlapping promoter region, a 2000-bp region

centered on each TSS. Histone modification enrichment (HM) for the i^{th} promoter region is calculated

as follows:

$$HM_i = log_2 \frac{\sum_k ChIP_{ik} + \alpha}{\sum_k Input_{ik} + \alpha},$$

where $\sum_k ChIP_{ik}$ represents the sum of either the H3K27ac or the H3K4me3 read signal mapped to the

promoter region at each position, $\sum_k Input_{ik}$ represents the sum of the control read signal mapped to

the promoter region, and α represents the 25[th] percentile of the $\sum_k Input_{ik}$ calculated for a given

sample.

M2A topology

M2A is a machine learning framework that leverages canonical deep-learning strategies, including

convolutional neural network (CNN) and fully connected (FC) layers. Each layer employs a LeakyReLU

(alpha = 0.1) and a kernel constraint by L2-normalization using maxnorm(3). CNN layers are two-

dimensional, with zero padding, a stepsize of (1,1), and a kernel size of (1, 2) to maintain feature space

and prevent convolutions across features from different window sizes. To test the efficacy of this

approach, we compared the performance of a traditional artificial neural network (ANN) consisting of

two FC layers versus three CNN layers in addition to two FC layers. During transfer learning, weights

corresponding to each of the three CNN layers of the six–NBL O-PDX M2A model trained previously

were frozen; only the weights corresponding to the two FC layers were optimized. A summary of all

model topologies and parameters can be found in Additional file 2: Table S16. To train and test each

model topology, we used Keras (v2.2.4) and Tensorflow (v2.1.0) in Python 3.6.5.

Training M2A

The core M2A model (without transfer) training set consisted of six O-PDX samples from the 16-

sample NBL cohort (Additional file 2: Table S17); we trained separate models for H3K27ac and

H3K4me3 HMs with the same WGBS features as the input. After the base models were trained,

transfer learning was employed for three separate datasets, namely the RMS, AML, and EWS

datasets. Each transfer learning model was trained using one sample from the cohort, for a total

of N models, where N equals the number of samples in the cohort. For the RMS and EWS

cohorts, an ensemble approach was used, whereby an averaged prediction from N−1 models

was generated after transfer learning with each sample. The same approach was used with the

AML cohort, except that only samples with $R^2 \geq 0.50$ between FPKM and H3K27ac were used

for transfer learning.

For each training scheme, the same parameters were used, including an 80/20 training/validation

split and a batch size of 128 (Additional file 2: Table S16). All sample input was randomized before

training. The Keras implementation of adadelta (default parameters) minimizing the mean squared error

(MSE) was used to optimize M2A. To prevent overtraining, the EarlyStopping method was employed by

monitoring validation loss for 10 epochs without at least a minimal gain in performance (min_delta =

0.0001) for a maximum of 80 epochs. In no case was the maximum number of epochs reached.

Determining promoter diversity in NBL models

Promoter region–based H3K27ac distributions clearly show a bimodal distribution (see

Additional file 1: Figure S1); therefore, to determine class occupancy (active versus inactive), a

Gaussian mixture model (GaussianMixture from sklearn version 0.22 with n_components = 2)

was applied for each individual sample. To determine the percentage of differentially active

promoters among all active promoters, we used a pairwise comparison approach for all samples.

Cancer consensus genes were downloaded from COSMIC (https://cancer.sanger.ac.uk/census)

(accessed on February 1, 2020). To avoid artificially inflated values from genes with multiple

TSSs, only cancer consensus genes with a single TSS according to GENCODE GRCh37.p13

(release 19) definitions were considered.

Evaluating M2A performance

When determining prediction performance, two primary metrics were considered, namely the R^2 and

the root mean squared error (RMSE). To measure the accuracy of M2A in predicting expressed genes,

we calculated the AUC-ROC by using roc_curve and the precision-recall curve AUC by using

average_precision_score from sklearn 0.22. Paired analyses were tested for statistical significance by

using a Wilcoxon signed-rank test (R v3.4.1). To determine outliers, a median-based method was

implemented using the "outlier" function in the R package GmAMisc [51]. To ensure that low-

mappability regions were not a confounding factor, we used the

wgEncodeCrgMApabilityAlign100mer.bw file downloaded from the UCSC Genome Browser

(http://genome.ucsc.edu/cgi-bin/hgFileUi?db=hg19&g=wgEncodeMapability). For GRCh38.p13

annotations, we used liftOver from UCSC (http://hgdownload.cse.ucsc.edu/admin/exe/linux.x86_64/) to

convert the mappability track to GRCh38. For all performance-related analyses, the average value of

this track within each non-overlapping 2000-bp promoter region was calculated, and promoter regions

with mappability > 0.75 were retained for evaluation.

For genes with multiple promoters, we use the maximum promoter activity to represent the gene

(in analyses of promoter activity vs. gene expression and for identifying genes with differential promoter

activities).

M2A captures the impact of DMRs on promoter region activity

The lists of genes that are differentially expressed in ERMS and ARMS samples and genes with DMRs

were previously reported in reference [42].

Alternative promoter usage analyses

To infer alternative promoter usage that was specific to the RMS subtypes ARMS and ERMS, we first

delineated the "active" vs. "inactive" promoters in a subtype (ERMS or ARMS) by applying a threshold

of $mean(H3K27ac) > 1$ for the average samples in the subtype. Next, primary promoters from multi-

promoter genes were determined by the average H3K27ac level within a specific subtype, and the

promoter with the maximum activity from a given gene was counted as the primary promoter. The

differences in promoter usage between two subtypes was defined as the difference between the activity

sum of the primary promoters and the activity sum of the secondary promoters in the two subtypes.

Analysis of alternative promoter usage in EWS

In the same manner as the RMS subtype analysis, EWS alternative promoter usage was determined in

two patient sample groups: *TP53* mutant and *TP53* wild-type groups (for sample status, see Additional

file 2: Table S17). Candidate genes were identified as active genes (H3K27ac ≥ 1 in at least one group)

with potential differential promoter activities (absolute difference ≥ 1) in the *TP53* wild-type and *TP53*

mutant groups. Candidate genes with alternative promoter usage were identified as genes that used

different active primary promoters in the wild-type and mutant groups and had an average promoter

usage difference of at least 0.5 between groups. Both alternative promoters and differentially active

promoters were considered in prognostic analyses (univariate screening followed by a multivariate

analysis) using Cox proportional hazard models (R 3.4.1). The final model was derived from backward

stepwise selection from a Cox proportional hazards model including *TP53* mutation status and all

potential markers (all genes or promoters with an FDR < 0.5 in the univariate analysis).

M2A code availability

The latest M2A models, feature generation, and prediction pipeline are available for download at

https://github.com/chenlab-sj/M2A. Additionally, source code with detailed instructions for transfer

learning using the M2A model with input samples from other domains is available.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analysed during the current study are publicly available, as detailed in

Additional File 2: Table S15.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships

that could have appeared to influence the work reported in this paper.

Funding

National Cancer Institute of the National Institutes of Health [P30CA021765]; American Lebanese

Syrian Associated Charities (ALSAC). The content is solely the responsibility of the authors and does

not necessarily represent the official views of the National Institutes of Health.

Authors' contributions

JW: Methodology, software, data analysis, result interpretation, writing - original draft, writing - review &

editing. **BX**: Resources, data curation. **DP**: Software, resources. **AT**: Software, resources. **XC**:

Conceptualization, methodology, data analysis, result interpretation, writing - review & editing,

supervision.

Acknowledgements

We thank Keith A. Laycock, PhD, ELS, for editing the manuscript.

References

1.   Davuluri RV, Suzuki Y, Sugano S, Plass C, Huang TH. The functional consequences of
alternative promoter use in mammalian genomes. Trends Genet. 2008;24:167–177.

2.   Demircioğlu D, Cukuroglu E, Kindermans M, Nandi T, Calabrese C, Fonseca NA, et al. A
pan-cancer transcriptome analysis reveals pervasive regulation through alternative
promoters. Cell. 2019;178:1465–1477.e17.

3.   Qamra A, Xing M, Padmanabhan N, Kwok JJT, Zhang S, Xu C, et al. Epigenomic
promoter alterations amplify gene isoform and immunogenic diversity in gastric
adenocarcinoma. Cancer Discov. 2017;7:630–651.

4.   Sotillo E, Barrett DM, Black KL, Bagashev A, Oldridge D, Wu G, et al. Convergence of
acquired mutations and alternative splicing of CD19 enables resistance to CART-19
immunotherapy. Cancer Discov. 2015;5:1282–1295.

5.   Ma X, Liu Y, Liu Y, Alexandrov LB, Edmonson MN, Gawad C, et al. Pan-cancer genome
and transcriptome analyses of 1,699 paediatric leukaemias and solid tumours. Nature.
2018;555:371–376.

6.   Gröbner SN, Worst BC, Weischenfeldt J, Buchhalter I, Kleinheinz K, Rudneva VA, et al.
The landscape of genomic alterations across childhood cancers. Nature. 2018;555:321–
327.

7.   Huether R, Dong L, Chen X, Wu G, Parker M, Wei L, et al. The landscape of somatic
mutations in epigenetic regulators across 1,000 paediatric cancer genomes. Nat
Commun. 2014;5:3630.

8.   Andersson R, Sandelin A. Determinants of enhancer and promoter activities of
regulatory elements. Nat Rev Genet. 2020;21:71–87.

9.   Kelley DZ, Flam EL, Izumchenko E, Danilova LV, Wulf HA, Guo T, et al. Integrated
analysis of whole-genome ChIP-seq and RNA-seq data of primary head and neck tumor
samples associates HPV integration sites with open chromatin marks. Cancer Res.
2017;77:6538–6550.

10.   Kagohara LT, Stein-O'Brien GL, Kelley D, Flam E, Wick HC, Danilova LV, et al.

Epigenetic regulation of gene expression in cancer: techniques, resources and analysis.

Brief Funct Genomics. 2018;17:49–63.

11.   Zhang P, Lehmann BD, Shyr Y, Guo Y. The utilization of formalin fixed-paraffin-

embedded specimens in high throughput genomic studies. Int J Genomics.

2017;2017:1926304.

12.   Moran S, Vizoso M, Martinez-Cardús A, Gomez A, Matías-Guiu X, Chiavenna SM, et al.

Validation of DNA methylation profiling in formalin-fixed paraffin-embedded samples

using the Infinium HumanMethylation450 Microarray. Epigenetics. 2014;9:829–833.

13.   de Ruijter TC, de Hoon JP, Slaats J, de Vries B, Janssen MJ, van Wezel T, et al.

Formalin-fixed, paraffin-embedded (FFPE) tissue epigenomics using Infinium

HumanMethylation450 BeadChip assays. Lab Invest. 2015;95:833–842.

14.   Gu H, Bock C, Mikkelsen TS, Jäger N, Smith ZD, Tomazou E, et al. Genome-scale DNA

methylation mapping of clinical samples at single-nucleotide resolution. Nat Methods.

2010;7:133–136.

15.   Ziller MJ, Gu H, Müller F, Donaghey J, Tsai LT, Kohlbacher O, et al. Charting a dynamic

DNA methylation landscape of the human genome. Nature. 2013;500:477–481.

16.   Charlet J, Duymich CE, Lay FD, Mundbjerg K, Dalsgaard Sørensen K, Liang G, et al.

Bivalent regions of cytosine methylation and H3K27 acetylation suggest an active role

for DNA methylation at enhancers. Mol Cell. 2016;62:422–431.

17.   Onuchic V, Lurie E, Carrero I, Pawliczek P, Patel RY, Rozowsky J, et al. Allele-specific

epigenome maps reveal sequence-dependent stochastic switching at regulatory loci.

Science. 2018;361:eaar3146.

18.   Stadler MB, Murr R, Burger L, Ivanek R, Lienert F, Schöler A, et al. DNA-binding factors

shape the mouse methylome at distal regulatory regions. Nature. 2011;480:490–495.

19.   Sheffield NC, Pierron G, Klughammer J, Datlinger P, Schönegger A, Schuster M, et al.

770   DNA methylation heterogeneity defines a disease spectrum in Ewing sarcoma. Nat Med.

771   2017;23:386–395.

772   20.   Zhu H, Wang G, Qian J. Transcription factors as readers and effectors of DNA

773   methylation. Nat Rev Genet. 2016;17:551–565.

774   21.   Kondo Y. Epigenetic cross-talk between DNA methylation and histone modifications in

775   human cancers. Yonsei Med J. 2009;50:455–463.

776   22.   Rothbart SB, Strahl BD. Interpreting the language of histone and DNA modifications.

777   Biochim Biophys Acta. 2014;1839:627–643.

778   23.   Hashimshony T, Zhang J, Keshet I, Bustin M, Cedar H. The role of DNA methylation in

779   setting up chromatin structure during development. Nat Genet. 2003;34:187–192.

780   24.   Moore LD, Le T, Fan G. DNA methylation and its basic function.

781   Neuropsychopharmacology. 2013, 38:23–38.

782   25.   Jenkinson G, Pujadas E, Goutsias J, Feinberg AP. Potential energy landscapes identify

783   the information-theoretic nature of the epigenome. Nat Genet. 2017;49:719–729.

784   26.   Fortin JP, Hansen KD. Reconstructing A/B compartments as revealed by Hi-C using

785   long-range correlations in epigenetic data. Genome Biol. 2015;16:180.

786   27.   Simmonds P, Loomis E, Curry E. DNA methylation-based chromatin compartments and

787   ChIP-seq profiles reveal transcriptional drivers of prostate carcinogenesis. Genome Med.

788   2017;9:54.

789   28.   Mack SC, Witt H, Piro RM, Gu L, Zuyderduyn S, Stütz AM, et al. Epigenomic alterations

790   define lethal CIMP-positive ependymomas of infancy. Nature. 2014;506:445–450.

791   29.   Teodoridis JM, Hardie C, Brown R. CpG island methylator phenotype (CIMP) in cancer:

792   causes and implications. Cancer Lett. 2008;268:177–186.

30. Capper D, Jones DTW, Sill M, Hovestadt V, Schrimpf D, Sturm D, et al. DNA

methylation-based classification of central nervous system tumours. Nature.

2018;555:469–474.

31. Perez E, Capper D. Invited Review: DNA methylation-based classification of paediatric

brain tumours. Neuropathol Appl Neurobiol. 2020;46:28–47.

32. Strahl BD, Allis CD. The language of covalent histone modifications. Nature.

2000;403:41–45.

33. Musselman CA, Lalonde ME, Côté J, Kutateladze TG. Perceiving the epigenetic

landscape through histone readers. Nat Struct Mol Biol. 2012;19:1218–1227.

34. Spainhour JCG, Lim HS, Yi SV, Qiu P. Correlation patterns between DNA methylation

and gene expression in The Cancer Genome Atlas. Cancer Inform.

2019;18:1176935119828776.

35. Baylin SB. DNA methylation and gene silencing in cancer. Nat. Clin. Pract. Oncol.

2005;2:S4–S11.

36. Jones PA. Functions of DNA methylation: islands, start sites, gene bodies and beyond.

Nat. Rev. Genet. 2012;13:484–492.

37. Lay FD, Liu Y, Kelly TK, Witt H, Farnham PJ, Jones PA, et al. The role of DNA

methylation in directing the functional organization of the cancer epigenome. Genome

Res. 2015;25:467–477.

38. Kagohara LT, Stein-O'Brien GL, Kelley D, Flam E, Wick HC, Danilova LV, et al.

Epigenetic regulation of gene expression in cancer: techniques, resources and analysis.

Brief Funct Genomics. 2018;17:49–63.

39. Kapourani CA, Sanguinetti G. Higher order methylation features for clustering and

prediction in epigenomic studies. Bioinformatics. 2016;32:i405–i412.

40. Zeineldin M, Federico S, Chen X, Fan Y, Xu B, Stewart E, et al. *MYCN* amplification and

ATRX mutations are incompatible in neuroblastoma. Nat Commun. 2020;11:913.

41.   Karlić R, Chung HR, Lasserre J, Vlahovicek K, Vingron M. Histone modification levels

are predictive for gene expression. Proc Natl Acad Sci U S A. 2010;107:2926–2931.

42.   Stewart E, McEvoy J, Wang H, Chen X, Honnell V, Ocarz M, et al. Identification of

therapeutic targets in rhabdomyosarcoma through integrated genomic, epigenomic, and

proteomic analyses. Cancer Cell. 2018;34:411–426.e19.

43.   ENCODE Project Consortium. An integrated encyclopedia of DNA elements in the

human genome. Nature. 2012;489:57–74.

44.   Stunnenberg HG, The International Human Epigenome Consortium, Hirst M. The

International Human Epigenome Consortium: a blueprint for scientific collaboration and

discovery. Cell. 2016;167:1145–1149.

45.   Singh AA, Schuurman K, Nevedomskaya E, Stelloo S, Linder S, Droog M, et al.

Optimized ChIP-seq method facilitates transcription factor profiling in human tumors. Life

Sci Alliance. 2019,2:e201800115.

46.   Seligson DB, Horvath S, Shi T, Yu H, Tze S, Grunstein M, et al. Global histone

modification patterns predict risk of prostate cancer recurrence. Nature. 2005;435:1262–

1266.

47.   Chen X, Stewart E, Shelat AA, Qu C, Bahrami A, Hatley M, et al. Targeting oxidative

stress in embryonal rhabdomyosarcoma. Cancer Cell. 2013;24:710–724.

48.   Stewart E, Shelat A, Bradley C, Chen X, Federico S, Thiagarajan S, et al. Development

and characterization of a human orthotopic neuroblastoma xenograft. Dev Biol.

2015;407:344–355.

49.   Stewart E, Federico SM, Chen X, Shelat AA, Bradley C, Gordon B, et al. Orthotopic

patient-derived xenografts of paediatric solid tumours. Nature. 2017;549:96–100.

50.   Murphy AJ, Chen X, Pinto EM, Williams JS, Clay MR, Pounds SB, et al. Forty-five

patient-derived xenografts capture the clinical and biological heterogeneity of Wilms

tumor. Nat Commun. 2019;10:5806.

51.   Wilcox RR: The normal curve and outlier detection. In: Fundamentals of Modern

846   Statistical Methods: Substantially Improving Power and Accuracy. New York, NY:

847   Springer; 2010. pp 29–45.

848   52.   Golob JL, Kumar RM, Guenther MG, Pabon LM, Pratt GA, Loring JF, et al. Evidence that

849   gene activation and silencing during stem cell differentiation requires a transcriptionally

850   paused intermediate state. PLoS One. 2011;6:e22416.

851   53.   Berdasco M, Esteller M. Clinical epigenetics: seizing opportunities for translation. Nat

852   Rev Genet. 2019;20:109–127.

853   54.   Hinoue T, Weisenberger DJ, Lange CPE, Shen H, Byun HM, Van Den Berg D, et al.

854   Genome-scale analysis of aberrant DNA methylation in colorectal cancer. Genome Res.

855   2011;22:271–282.

856   55.   Cao L, Yu Y, Bilke S, Walker RL, Mayeenuddin LH, Azorsa DO, et al. Genome-wide

857   identification of PAX3-FKHR binding sites in rhabdomyosarcoma reveals candidate

858   target genes important for development and cancer. Cancer Res. 2010;70:6497–6508.

859   56.   Laé M, Ahn E, Mercado G, Chuai S, Edgar M, Pawel B, et al. Global gene expression

860   profiling of PAX-FKHR fusion-positive alveolar and PAX-FKHR fusion-negative

861   embryonal rhabdomyosarcomas. J Pathol. 2007;212:143–151.

862   57.   Marshall AD, Grosveld GC. Alveolar rhabdomyosarcoma—the molecular drivers of

863   PAX3/7-FOXO1-induced tumorigenesis. Skelet Muscle. 2012;2:25.

864   58.   Honda T, Inui M. PDZRN3 regulates differentiation of myoblasts into myotubes through

865   transcriptional and posttranslational control of Id2. J Cell Physiol. 2019;234:2963–2972.

866   59.   Tirode F, Surdez D, Ma X, Parker M, Le Deley MC, Bahrami A, et al. Genomic

867   landscape of Ewing sarcoma defines an aggressive subtype with co-association of

868   *STAG2* and *TP53* mutations. Cancer Discov. 2014;4:1342–1353.

869   60.   Virden RA, Thiele CJ, Liu Z. Characterization of critical domains within the tumor

870   suppressor CASZ1 required for transcriptional regulation and growth suppression. Mol

871   Cell Biol. 2012;32:1518–1528.

872   61.   Wang JL, Yang MY, Xiao S, Sun B, Li YM, Yang LY. Downregulation of castor zinc

873   finger 1 predicts poor prognosis and facilitates hepatocellular carcinoma progression via

874   MAPK/ERK signaling. J Exp Clin Cancer Res. 2018;37:45.

875   62.   Kim B, Jung M, Moon KC. The prognostic significance of protein expression of CASZ1 in

876   clear cell renal cell carcinoma. Dis Markers. 2019;2019:1342161.

877   63.   Shen H, Laird PW. Interplay between the cancer genome and epigenome. Cell.

878   2013;153:38–55.

879   64.   De Craene B, Berx G. Regulatory networks defining EMT during cancer initiation and

880   progression. Nat Rev Cancer. 2013;13:97–110.

881   65.   Roadmap Epigenomics Consortium, Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A,

882   Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang J, et al. Integrative analysis of 111

883   reference human epigenomes. Nature. 2015;518:317–330.

884   66.   Ooi WF, Xing M, Xu C, Yao X, Ramlee MK, Lim MC, et al. Epigenomic profiling of

885   primary gastric adenocarcinoma reveals super-enhancer heterogeneity. Nat Commun.

886   2016;7:12983.

887   67.   Dong X, Greven MC, Kundaje A, Djebali S, Brown JB, Cheng C, et al. Modeling gene

888   expression using chromatin features in various cellular contexts. Genome Biol.

889   2012;13:R53.

890   68.   Singh R, Lanchantin J, Robins G, Qi Y. DeepChrome: deep-learning for predicting gene

891   expression from histone modifications. Bioinformatics. 2016;32:i639–i648.

892   69.   Schmitz U, Pinello N, Jia F, Alasmari S, Ritchie W, Keightley MC, et al. Intron retention

893   enhances gene regulatory complexity in vertebrates. Genome Biol. 2017;18:216.

894   70.   Pennacchio LA, Bickmore W, Dean A, Nobrega MA, Bejerano G. Enhancers: five

895   essential questions. Nat Rev Genet. 2013;14:288–295.

896   71.   Weigel C, Chaisaingmongkol J, Assenov Y, Kuhmann C, Winkler V, Santi I, et al. DNA

897   methylation at an enhancer of the three prime repair exonuclease 2 gene (*TREX2*) is

898   linked to gene expression and survival in laryngeal cancer. Clin Epigenetics. 2019;11:67.

899   72.   Heyn H, Vidal E, Ferreira HJ, Vizoso M, Sayols S, Gomez A, et al. Epigenomic analysis

900   detects aberrant super-enhancer DNA methylation in human cancer. Genome Biol.

901   2016;17:11.

Figure Legends

Figure 1. Evaluation of H3K27ac promoter activity variation between NBL samples.

(a–c) A comparison of H3K27ac promoter activity in **(a)** NBL cell line samples, **(b)** NBL O-PDX

samples, and **(c)** NBL cell line and O-PDX samples, where the promoter region is defined as the

TSS ± 1000 bp. **(d)** Venn diagram of results of NBL cell line promoter activity comparison. **(e)**

Venn diagram of results of O-PDX promoter activity analysis. **(f)** The proportion of sample-

specific active promoters among all active promoters was compared within and between cell line

and O-PDX NBL samples.

Figure 2. M2A feature processing and training workflow.

The M2A framework hinges on the feature processing pipeline **(a)**. First, windowed features

centered around the TSS are calculated from WGBS data for each unique promoter region,

serving as input for model initial training and transfer learning. In this case, matching response

variables to these features were generated for H3K27ac and H3K4me3 for each promoter

region corresponding to a unique TSS. The matching response variables serve as input to the

model topologies depicted in **(b)**, where M2A first extracts high-level features by using a series

of convolutional layers then maps these features to response variables in fully-connected (FC)

layers. Transfer learning with M2A leverages pretrained feature extraction, training only the FC

layers.

Figure 3. M2A performance in NBL and RMS cohorts.

(a, b) Analysis of the performance of M2A in NBL and RMS cohorts with **(a)** H3K4me3 inference

and **(b)** H3K27ac inference, as compared to the replicate consistency from the ENCODE

dataset. Sample KMS-11 was excluded as an apparent outlier $R2=0.016$, RMSE = 1.869 **(c–g)**

Individual examples of median M2A performers in **(c)** NBL cell line H3K27ac inference, **(d)** NBL

cell line H3K4me3 inference, **(e)** RMS H3K27ac inference (pre-transfer), **(f)** RMS H3K4me3

inference, and **(g)** RMS H3K27ac inference (post-transfer). **(h)** A comparison of the

performance of M2A with RMS samples before and after transfer, as measured by RMSE.

Figure 4. M2A recapitulates subtype differences in RMS.

(a) Embryonal and alveolar RMS subtypes are clearly delineated in a tSNE analysis of observed

H3K27ac promoter levels, M2A base model–inferred H3K27ac levels, and M2A post-transfer–

inferred H3K27ac levels. **(b–d)** The subtype-specific genes *GAS2* **(b)** and *NOS1* **(c)** show these

specificities at the DNA methylation, H3K27ac, and H3K4me3 levels; this is faithfully

recapitulated by the M2A H3K27ac inferences for *GAS2* **(d)** and *NOS1* **(e)**.

Figure 5. M2A reveals alternate promoter usage in RMS and EWS.

(a) M2A predicts RMS subtype–specific promoter usage in *PDZRN3*, a known target of the

PAX3/7–FOXO1 fusion protein, which is consistent at the observed values of H3K27ac,

H3K4me3, and RNA-seq. **(b)** A Cox proportional hazards model of alternate promoter usage in

EWS patient samples with and without *TP53* mutations reveals isoforms with potential

prognostic value.

Additional Files

Additional file 1 (file type .PDF, 1.5 MB):

Figure S1. NBL sample H3K27ac promoter distribution.

A comparison of promoter H3K27ac enrichment in *MYCN* amplified (MNA) NBLcell line and O-

PDX samples shows a clear bimodal distribution, delineating "active" and "non-active"

promoters.

Figure S2. M2A prediction generalizability analysis.

(a) The observed promoter H3K27ac levels in samples from M2A training sets and both NBL

and RMS cohort test sets show similar distributions despite the different cancer types. **(b)** The

prediction consistency of M2A outperforms the observed consistency between the most similar

training samples and the test samples from the NBL and RMS cohorts.

Figure S3. M2A with transfer learning outperforms a baseline M2A model of the same

cancer type.

An M2A baseline model trained initially with six NBL O-PDX samples and then with RMS LOO

transfer learning consistently outperforms an M2A baseline model trained with only RMS

training samples.

Figure S4. Signal-to-noise analysis of ENCODE and NBL datasets.

Although the values are highly correlated, sample comparison at the observed H3K27ac

promoter level **(a)** and the H3K4me3 promoter level **(b)** indicates higher RMSE values and

different signal-to-noise profiles.

Figure S5. M2A ENCODE cohort performance.

M2A accurately infers both H3K27ac and H4K4me3 promoter levels in the publicly available

ENCODE dataset.

Figure S6. Analysis of outlier H1-ESC.

(a) When inferring H3K27ac promoter levels, M2A was substantially less accurate with the

ENCODE sample H1-ESC, which is an outlier in the ENCODE cohort. **(b, c)** M2A-predicted

H3K27ac promoter levels **(b)** are more consistent with, and more predictive of, H1-ESC gene

expression than are the actual observed H1-ESC H3K27ac promoter levels **(c)**.

Figure S7. Predicting gene expression in the ENCODE dataset.

(a–i) The indirect ability of M2A to predict gene expression (i.e., expressed vs. not expressed)

on the basis of both M2A-predicted and observed H3K27ac promoter levels for each sample

was determined by comparing the AUCs of the receiver operating characteristic (ROC) curves.

Figure S8. Consistency of gene expression and H3K27ac promoter levels in the AML

cohort.

The consistency, as determined by Pearson's R^2, between the observed values for gene

expression and the H3K27ac promoter levels is remarkably predictive of the performance M2A

in predicting H3K27ac promoter levels in samples from the AML cohort.

Figure S9. M2A accurately determines subtype differences between embryonal and

alveolar RMS.

(a) The promoter activities of single-promoter, differentially expressed genes in the RMS

subtypes ERMS and ARMS are accurately inferred by an M2A base model (trained with six O-

PDX NBL samples). **(b)** The predictive performance of M2A is further boosted by transfer

learning with one RMS sample. **(c, d)** The performance of M2A declines slightly when the model

is applied to all promoters of differentially expressed genes **(c)**, but it recovers when an M2A

model with transfer learning with only one RMS training sample is applied **(d)**.

Figure S10. Kaplan–Meier log-rank analysis by mutation status in EWS.

The prognostic ability of **(a)** *TP53* or **(b)** *STAG2* mutation status in the EWS cohort was

determined by the log-rank test and visualized using the Kaplan–Meier survival curve.

1003

Additional file 2 (file type: .XLSX, 153 KB).

Table S1: H3K27ac active cancer consensus genes in 3 NBL cell lines, and 3 NBL O–PDX

samples.

Table S2: a basic ANN model vs. M2A predictive performance comparison.

Table S3: M2A predictive performance in NBL cell line samples.

Table S4: observed H3K27ac and H3K4me3 ENCODE replicate consistencies.

Table S5: M2A predictive performance in RMS O–PDX samples.

Table S6: M2A RMS transfer model predictive performance in RMS O–PDX samples.

Table S7: M2A predictive performance in ENCODE dataset.

Table S8: M2A predictive performance in AML samples.

Table S9: ERMS vs. ARMS DMRs and associated genes (overexpressed in ERMS).

Table S10: ERMS vs. ARMS DMRs and associated genes (overexpressed in ARMS).

Table S11: M2A alternate promoter usage predictive performance between ARMS and

ERMS samples.

Table S12: M2A predictive performance in EWS samples, before and after transfer.

Table S13: a univariate survival analysis of differential H3K27ac promoter activity

between TP53 mutant and TP53 wild–type EWS tumors.

Table S14: a univariate survival analysis of alternate promoter usage between TP53

mutant and TP53 wild–type EWS tumors.

Table S15: dataset availability.

Table S16: M2A model topologies.

Table S17: sample summary information.

(a)

Windowed feature processing

2500 bp window

250 bp window

features

features

TSS

DNA methylation

Response variables

H3K27ac

H3K4me3

-25000 bp

-2500 bp
+2500 bp

+25000 bp

training samples

Model input

High-level feature extraction

Feature-response mapping

Methyl2Activity topology

Convolution

Convolution

Convolution

FC

FC

Output

Model input

Pretrained convolution layers

Feature-response mapping

Methyl2Activity transfer-learning topology

FC

FC

Output

Response variable processing

$$HM_i = \log_2 \frac{\sum_k ChIP_{ik} + \alpha}{\sum_k Input_{ik} + \alpha}$$

(b)

High-level feature extraction

Feature-response mapping

Model input

Methyl2Activity topology

Convolution

Convolution

Convolution

FC

FC

Output

Pretrained convolution layers

Feature-response mapping

Model input

Methyl2Activity transfer-learning topology

FC

FC

Output

(a)

Observed promoter activity landscape of RMS tumors

Promoter activity landscape of RMS tumors inferred from the baseline model

Promoter activity landscape of RMS tumors inferred from the transfer model

tSNE-2

tSNE-1

ERMS
ARMS
Spindle

(b) SJRHB013757_X1 (ARMS)

WGBS
H3K4me3
H3K27ac

1 kbp

SJRHB013758_X2 (ERMS)

WGBS
H3K4me3
H3K27ac

GAS2

(c) SJRHB013757_X1 (ARMS)

WGBS
H3K4me3
H3K27ac

1 kbp

SJRHB013758_X2 (ERMS)

WGBS
H3K4me3
H3K27ac

NOS1

(d) GAS2

H3K27ac level

ARMS   ERMS
RMS subtype

(e) NOS1-005

H3K27ac level

ARMS   ERMS
RMS subtype

Observed ARMS
Observed ERMS
Predictions from M2A

(a)

25 kbp   SJRHB010468_X1 (ARMS)

H3K4me3

H3K27ac

RNA-seq   *PDZRN3-201*

SJRHB000026_X1 (ERMS)

H3K4me3

H3K27ac

RNA-seq   *PDZRN3-001*

PDZRN3

(b)

	hazard ratio		p-value
TP53_mutation *(N=72)*	3.83 (1.30 - 11.27)		*0.015 **
CASZ1 *(N=72)* (ENST00000496432.2)	0.41 (0.19 - 0.88)		*0.023 **
RET *(N=72)* (ENST00000479913.1)	0.53 (0.29 - 0.97)		*0.039 **
TEX40 *(N=72)* (ENST00000535981.1)	0.39 (0.15 - 0.97)		*0.043 **

 # Events: 31; Global p-value (Log-Rank): < .001
AIC: 218.7; Concordance Index: 0.71

0.2   0.5   1   2   5   10