4 October 2020
Nyeema C. Harris
University of Michigan
1105 N. University Ave.
Ann Arbor, MI. 48109
(734) 647-3765
nyeema@umich.edu
8
9   RH: Gadsden et al. • Michigan ZoomIN
10

Michigan ZoomIN: validating crowd-sourcing to identify mammals from
camera surveys

GABRIEL I. GADSDEN,[1] *1105 N. University Ave., Applied Wildlife Ecology, Ecology and*

Evolutionary Biology, University of Michigan, Ann Arbor MI 48109, USA

RUMAAN MALHOTRA, *1105 N. University Ave., Applied Wildlife Ecology, Ecology and*

Evolutionary Biology, University of Michigan, Ann Arbor MI 48109, USA

JUSTIN SCHELL, *Shapiro Design Lab, University of Michigan, Ann Arbor MI 48109, USA*

TIFFANY CAREY,[2] *1105 N. University Ave., Applied Wildlife Ecology, Ecology and*

Evolutionary Biology, University of Michigan, Ann Arbor MI 48109, USA

NYEEMA C. HARRIS, [3]*1105 N. University Ave., Applied Wildlife Ecology, Ecology and*

Evolutionary Biology, University of Michigan, Ann Arbor MI 48109, USA

ABSTRACT

Camera trap studies have become a popular medium to assess many ecological phenomena

including population dynamics, patterns of biodiversity, and monitoring of endangered species.

In conjunction with the benefit to scientists, camera traps present an unprecedented opportunity

[1] *Current Address: Urban Energy Justice Lab, School For Environment and Sustainability, University of Michigan 440 Church St, Ann Arbor, MI 48109*
[2] *Current Address: National Wildlife Federation, Great Lakes Regional Center, Ann Arbor MI 48109*
[3] *Email:* nyeema@umich.edu

to involve the public in scientific research via image classifications. However, this engagement

strategy comes with a myriad of complications. Volunteers vary in their familiarity with wildlife,

and thus, the accuracy of user-derived classifications may be biased by the commonness or

popularity of species and user-experience. From an extensive multi-site camera trap study across

Michigan U.S.A, images were compiled and identified through a public science platform called

Michigan ZoomIN. We aggregated responses from 15 independent users per image using

multiple consensus methods to assess accuracy by comparing to species identification completed

by wildlife experts. We also evaluated how different factors including consensus algorithms,

study area, wildlife species, user support, and camera type influenced the accuracy of user-

derived classifications. Overall accuracy of user-derived classification was 97%; although,

several canid (e.g., *Canis lupus, Vulpes vulpes*) and mustelid (e.g., *Neovison vison*) species were

repeatedly difficult to identify by users and had lower accuracy. When validating user-derived

classification, we found that study area, consensus method, and user support best explained

accuracy. To continue to overcome stigma associated with data from untrained participants, we

demonstrated their value by showing the accuracy from volunteers was comparable to experts

when classifying North American mammals. Our hierarchical workflow that integrated multiple

consensus methods lead to more image classifications without extensive training and even when

the expertise of the volunteer was unknown. Ultimately, adopting such an approach can harness

broader participation, expedite future camera trap data synthesis, and improve allocation of

resources by scholars to enhance performance of public participants and increase accuracy of

user-derived data.

KEYWORDS accuracy, carnivores, citizen science, community, geographic variation,

engagement, Michigan, mustelid, validation.

Science is a public good. In recent years, an interest in broadening participation in science has

resulted in a wave of public (participatory, community, or citizen) science projects initiated by

scholars (Dickinson et al. 2010, Theobald et al. 2015). The seemingly unlikely partnership

between the public and scientists creates dialogue that can aid in demystifying the scientific

process, facilitate exposure to the natural world, and yield reciprocal benefits (Bonney et al.

2016, Schuttler et al. 2018b). Public participation in ecology is particularly salient due to humans

being a dominant force in natural systems with increasing pressures stemming from land

conversion, urban sprawl, and over-exploitation (Raudsepp-Hearne et al. 2010, Halpern et al.

2015, Young et al. 2016). The public can gather large amounts of data during a person's

everyday activity to monitor biodiversity and environmental change (Trumbull et al. 2000). For

example, someone interested in how urbanization affects species can assist in collecting bird

skyscraper collisions or simply take images with a smartphone and document sightings of

particular organisms (Ries and Oberhauser 2015, Shipley et al. 2018).

Digital technology, including cameras, is emerging as an important tool for participatory

science given the multitude of highly accessible platforms available to share data. As avid users

of technology, engaged youth have the potential to amplify positive effects of participatory

science through intergenerational learning (Arts et al. 2015, Peterson et al. 2019). Hence,

incorporating images garnered from camera surveys are now commonplace for public science

programs (e.g., Schuttler et al. 2018a). Camera surveys can answer a plethora of ecological,

behavioral, and even evolutionary questions across spatial and temporal scales, in a variety of

environments, and on a diversity of species (Burton et al. 2015, McShea et al. 2015, Steenweg et

al. 2017).

Although crowdsourcing has gained popularity, image identification from wildlife

surveys by communities of non-scientists is fraught with challenges (Hochachka et al. 2012).

Researchers not only have to develop the infrastructure to facilitate easy access, but also

construct a useful workflow to aggregate and structure responses into manageable data (Waetjen

and Shilling 2017, Egerer et al. 2019, Parrish et al. 2019). Chief amongst the rising concerns is

that public-generated data are susceptible to biases and inaccuracy of responses that can

compromise usability and hinder broader implementation in future scientific research (Dickinson

et al. 2010, Hochachka et al. 2012, Brown et al. 2018). By design, researchers are often not able

to control for who contributes to citizen science projects, resulting in heterogeneity in user

expertise. Projects also vary in the extensiveness of aids such as sample photos or comparison

charts for similar species that could alter a user's capacity to accurately classify images

(Skarlatidou et al. 2019). Through targeted outreach, engagement, and education strategies, some

projects are able to target specific audiences (e.g., K-12) and thoroughly train volunteers; though,

this is not always feasible and requires significant resources. Varying methods of image

classification, and the implementation of automated and crowd-sourcing aids in classifying

images to discern patterns from large amounts of biological information can be helpful in

reducing processing time (Maes et al. 2015, Norouzzadeh et al. 2018). Standardized guidelines

for processing and evaluating user-derived data will allow for scientists to better evaluate their

utility and leverage public science to tackle large data sets regardless of the research system

(Freitag and Pfeffer 2013, Maes et al. 2015, Egerer et al. 2019, Locke et al. 2019).

Overall, findings have been inconsistent concerning reliability of user-derived data for

scientific research, requiring additional scrutiny based on project workflows and inclusion of

expert validation (Danielsen et al. 2005, Crall et al. 2011). Katrak-Adefowora et al. (2020) found

that training allowed participants even with no biology background to correctly identify images

with comparable accuracy to biology experts. Other studies found that there was high agreement

among classifying individuals, indicating that user expertise was not significant in correct

classifications, but animal size and distinct markings had a greater impact on accuracy (Swanson

et al. 2016, Potter et al. 2019). Still other studies have found that environmental, species-specific

behaviors, and land use affect classification accuracy (Broeckhoven and le Fras Nortier Mouton

2015, Meek et al. 2016). Generally, existing studies on validation of user derived data fail to

explain factors contributing to accuracy, occur outside a North American context, or employ a

single metric for deriving accuracy from aggregated responses. Another consideration that

remains unexplored for camera surveys is whether data generated from public participants are

comparable across geographies. Potential differences in species composition and image quality

from environmental conditions such as weather may influence how the public classifies images,

which may affect accuracy (Meek et al. 2015, Steger et al. 2017).

Here, we used public generated image identifications from an extensive multi-site, multi-

season camera trap survey throughout the state of Michigan. Specifically, we created a public

science online platform called *Michigan ZoomIN (MZI)* for image processing and developed a

science communication strategy to solicit volunteers. We then evaluated user responses by

comparing various consensus methods to obtain a converged species identification as well as

examined factors that influenced the accuracy of users. We hypothesized that common species

will be classified with higher accuracy while physically similar or rare species would have lower

user accuracy and reach consensus using a relaxed consensus method. Using a hierarchical

approach to assess user responses, we hypothesized high agreement would positively correlate

with accuracy for all images regardless of consensus method and species characteristics. We also

hypothesized that camera type would influence user accuracy with images from cameras that

used a flash at night having higher accuracy. We hypothesized that the study area would affect

accuracy, as sampling locations varied in vegetation and environmental conditions that could

distort images and make classifying harder. We contribute three novel insights to research

involving image identification from crowdsourcing by: 1) determining whether hierarchical

consensus criteria can mitigate potential bias from volunteers with varying levels of expertise, 2)

assessing multiple consensus thresholds to compare accuracy among species, and 3) identifying

various factors that can diminish accuracy. Results from our work also have relevance in

minimizing workload in classifying for researchers, as well as providing a workflow for

assessing overall accuracy of user-derived classifications for other camera studies.

STUDY AREA

Throughout 2015—2017, the Applied Wildlife Ecology (AWE) lab at the University of

Michigan implemented camera trap surveys to investigate questions of community ecology

within the carnivore guild of Michigan. We deployed 273 remotely-triggered Reconyx© cameras

(PC 850, PC 900, PC 850C, PC 900C) across three study areas that had distinct environmental

and habitat attributes from north to south: the Huron Mountain Club (HMC; Marquette County),

University of Michigan Biological Station (UMBS; Pellston County), and the Shiawassee

National Wildlife Refuge (SNWR; Saginaw County). The Huron Mountain Club was our most

ecologically diverse site, comprised privately-owned mixed secondary and old growth forest

along the southern coast of Lake Superior. The University of Michigan Biological Station, a

research station, comprised a mix of deciduous and coniferous forest between Douglas and Burt

Lakes in the northern tip of Michigan's Lower Peninsula. The Shiawassee National Wildlife

Refuge, a floodplain of Saginaw County, comprised marsh, bottomland hardwood forest, and

grasslands abutting urban development and agriculture managed by the U.S. Fish & Wildlife

Service. At each site, unbaited cameras were placed 0.5—1.0 m off the ground and affixed to

trees > 0.5m diameter. Camera settings included: high sensitivity, one-second lapse between a

sequence of three pictures in a trigger, and a 15-second quiet period between triggers. Our survey

efforts across 32,542 trap nights resulted in 10,199 of camera trap images that required accurate

species identifications.

METHODS

Description of *Michigan ZoomIN*

We developed a public science website called *Michigan ZoomIN (MZI)* on the

Zooniverse Project Builder platform where we uploaded images and collated user responses over

multiple seasons (Figure 1). From its origins in 2007 with the Galaxy Zoo project, which asked

volunteers to classify shapes of galaxies, the Zooniverse crowdsourcing platform has created and

hosted more than 200 different projects from disciplines as diverse as astronomy, ecology,

physics, history, and more to promote people-powered research (Lintott 2019; Zooniverse.org).

Through both a coordinated outreach program throughout Michigan and the existing Zooniverse

community, 645,000 classifications were completed over one year by 3,950 registered users

(unregistered users also contributed but were not trackable). There were 39 possible

classifications ranging from the focal carnivores in our empirical field research (e.g., coyote

Canis latrans, raccoon *Procyon lotor,* American black bear *Ursus americianus,* red fox *Vulpes*

vulpes), and their prey (e.g., white-tailed deer *Odocoileus virginianus*, squirrel *Sciurus*

carolinensis, opossum *Didelphis virginiana*, cottontail rabbit *Sylvilagus floridanus*, porcupine

Erethizon dorsatum, small mammal). Users were given written descriptions of species as well as

photos to aid in classifying species from survey images (Figure 1 – label 7). We also

incorporated the "looks like" tool in MZI, which allows volunteers to narrow possible

classifications based on color, shape, and body size (Figure 1 – label 5). We did not include

'unknown' as a possible choice for selection to promote more discrimination and an "educated"

guess. Users were also provided a field guide with descriptions of common ecological tenets

such as competition and dietary separation, and additional images and information useful for

classification of species (Figure 1 – label 6). The field guide resource was included to provide

more education to improve understanding and stimulate greater enthusiasm particularly for

younger audiences. Images were retired from viewing once 15 unique users classified the image.

Data Aggregation

To produce a final identification for multiple user responses, we aggregated

classifications into a single image species identification using various consensus approaches

(Figure 2). Images were considered to have reached consensus if they matched the criteria of any

one of four consensus methods: agreement, Kappa, Kappa and evenness, or gold standard. 1)

Agreement: if all 15 users agreed on the species identification for an image. 2) Kappa: A Fleiss

Kappa statistic (modified from Cohen's Kappa, for more than 2 users) above 0.8 designating a

substantial strength of agreement (Landis and Koch 1977). 3) Kappa and Evenness: A Fleiss

Kappa above 0.6 (lower than the recommendations of Landis and Koch (1977)), and evenness

below 0.5. Pielou's evenness is an index that measures the proportion of votes given to a

particular species in comparison to the number of species picked for the image in question. The

scale runs from 0—1, with 0 indicating low evenness and higher accuracy (Pielou 1966,

Swanson et al. 2015). 4) Gold Standard: At least one of the users classifying the image was from

our research team and thus categorized as an expert with a gold standard tag. If the expert was

unsure of classification, they removed the gold tag prior to identification submission. Images that

did not reach any consensus method were classified as Majority and later verified by a project

researcher (Figure 2). The following process allowed us to test a hierarchical approach that

ranged from rigorous (100% user agreement) to more relaxed criteria of user agreement. We

specifically tested Kappa and Kappa and Evenness as metrics that allow progressively lower

levels of user agreement to account for the differing levels of experience inherent to a diverse

group of users, while producing comparable accuracy to stricter consensus criteria (100% user

agreement amongst all 15 users).

Expert Review

To test the efficacy of the consensus algorithm, researchers verified a random sample of

5,000 images to compare against the user-generated identifications. In addition, all images that

had a carnivore identification (with the exception of raccoons) generated by the consensus

algorithm were also reviewed, for a total review dataset of n = 5,734. Beyond the initial random

sample of 5,000 images, additional raccoon images were not included for expert verification due

to the assumption that they did not fit the rarity criterion and were well represented in the

random sample.

Evaluation of Accuracy

As a rough measure of the accuracy for each of the consensus methods used in the

consensus algorithm, we calculated the proportion of correct answers for each method. For those

images that did not meet any of the criteria to produce consensus identification, we calculated the

proportion correctly identified among the 15 user classifications. For hypothesis testing, we first

constructed a generalized linear model to evaluate factors influencing accuracy. Aggregated user

responses from the consensus output were compared to the expert dataset to generate a binary

output of 1(correct) or 0 (incorrect). For comparison against images that were classified by

consensus, images that did not meet a consensus criterion were classified under the consensus

type as majority. Majority represented the most widely chosen species by public users when

consensus was not reached by other methods. We included: consensus methods, species, study

area, camera model, and percent user agreement as fixed effects in the model. Species was used

to better understand if animal similarity or rarity of a species affected the accuracy. We used two

different camera models in our study. Infrared cameras provide black and white images in low

light settings, while white flash cameras produce color images, even at night. Percent user

agreement represented a measure on how well users agreed on a classification and corresponding

difficulty depending on species. We conducted model selection using Akaike information

criterion (AIC). All models < 3 ΔAIC were considered top models and beta coefficients for

variables from top selected models were then used to test the following hypotheses:

1. Species that have a high degree of similarity will be harder to identify, which will

negatively affect accuracy;

2. Species accuracy will be greater for species that are more common and have greater

abundance indexed by trigger activity;

3. Stronger similarity in user agreement will correlate to overall higher accuracy;

4. White flash cameras will yield higher accuracy;

5. Accuracy will vary across study areas with the site closest to human habitation yielding

higher accuracy.

All model construction and selection were conducted using the lme4 package in R (version

3.4.4).

RESULTS

From our public science project, *Michigan ZoomIN* involved 3,950 registered users that were

involved in classifying 10,199 image sequences across three study areas. (HMC: 1,630; UMBS:

2,485; SNWR: 6,084). Each sequence consisted of three images and was reviewed by 15 unique

users. The most common species were raccoon and white-tailed deer with 5,594 (51%) and 2,592

(14%) classifications, respectively. Most images (95%) reached a consensus method to converge

on a specific species identification. The remaining 5% remained unidentifiable even after being

evaluated by a team researcher (Figure 2). Two-thirds of images were classified from complete

agreement among all 15 users. Raccoon, deer, and striped skunk (*Mephitis mephitis*) were the

only species to have higher than 50% of their images reach agreement (Figure 3). Kappa

evenness was the second most common consensus method, comprising 19% of images. Majority,

yielded completion of 10%, and majority plus gold standard comprised < 1% of images.

Groundhog (*Marmota monax*), fisher (*Pekania pennanti*), mink (*Neovison vison*), muskrat

(*Ondatra zibethicus*), and snowshoe hare (*Lepus americanus*) had 100% of their images go to

majority, and all these species represented the fewest captures in the dataset (Figure 3).

We built generalized linear models to evaluate factors that influenced the accuracy of

species classifications. The top model included: consensus method, percent user agreement, and

study area (Table 1). Less strict consensus criteria were only marginally worse except for

majority: Kappa ($\beta = -1.61$; $p = 0.014$), kappa evenness ($\beta = -1.162$; $p = 0.058$), and majority (β

$= -3.355$; $p < 0.0001$). Percent user agreement ($\beta = 5.328$; $p < 0.0001$) was strongly associated

with higher accuracy. Contrary to expectations, the camera model did not influence accuracy. As

previously stated, consensus of 100% user agreement was the most common method in which

images reached the final dataset. While most species were represented in images that reached

consensus, species varied in which consensus method was most effective for identification.

Indeed, the wide geographic coverage of the camera surveys presented the possibility for

differences in accuracy based on study area. With a 0.1 p-value threshold, images from SNWR

were significant in identification accuracy compared to other study areas (β= 5.047; p-value =

0.059). In contrast, images from HMC, the most remote site with the highest species diversity

located in the Upper Peninsula of Michigan, had the lowest accuracy of species classification.

Notably, the public was 99% accurate in identifying false positives, incidental triggers

from wind or vegetation movement. Small mammals were often confused as partial images of

cottontail rabbits, or mistakenly missed and labeled as nothing there. Though many species had

high accuracy (>90%, Figure 4), accuracy was statistically significant for squirrels (β= 2.180; p =

0.041) and porcupines (β= 2.299; p = 0.042) likely because of the ubiquity of squirrels and the

uniqueness of porcupines. Classifying animals incorrectly were common only for particular

species such as mustelids and coyotes. The high false positives were due to similarities to other

species based upon body size, color, and activity patterns. Mustelids had the lowest overall

accuracy (Figure 4); arguably, because the family comprises species with similar anatomical

features (e.g., marten *Martes americana*; long-tailed weasel *Mustela frenata*; fisher; mink) that

expectedly would prove difficult to discern for non-experts.

To underscore the importance of considering multiple consensus thresholds, we evaluated

results for coyotes as a case study, given their dynamic role within the carnivore community and

because they are commonly mistaken for other canid species (Prugh et al. 2009, Colborn et al.

2020). There were 520 classifications of coyotes by users, of which 265 were classified through

one of our consensus criteria. Only 9% passed consensus from complete user agreement, while

still obtaining >90% accuracy. Instead, the kappa statistic was the best consensus method for

identifying this species, comprising 48% of coyote classifications (Figure 3).

We found marginal support for differences in accuracy across species related to study

area and species interactions. All study areas had either 99–100% accuracy for common species

such as deer, raccoon, and striped skunk. However, rare species, as either being infrequently

detected or geographically constrained, had lower accuracy, consistent with our hypothesis. For

example, fisher and the small-bodied muskrat were only detected at HMC and SNWR,

respectively and had accuracies <50%. Gray wolves (*Canis lupus*) are similar in appearance to

coyotes and found only at HMC where they were most commonly misidentified as coyotes.

Likewise, though coyotes occurred in every study area, they had the lowest accuracy at HMC. In

contrast, the badger (*Taxidea taxus*), which has a noticeably distinct body shape and striking

facial pattern, was only captured at UMBS and yet had 75% accuracy.

DISCUSSION

Our study builds on previous research by suggesting common species, those with distinguishing

features, and generally larger species are most accurately identified by the public whereas

smaller bodied species and rarer species are less accurately classified. Similarly, Potter et al.

(2019) found the most distinct mammal species such as short-beaked echidna (*Tachyglossus*

aculeatus) had the highest accuracy from camera surveys in northern Australia, and groups of

closely related species with similar morphology, such as mustelids had the most errors. In an

African ecosystem, Swanson et al. (2015) found that different species of gazelles as well as

weasel-like small mammals were often misidentified from camera data derived from public

users. Such variation in accuracy by species may reflect differences in capturing high quality

images due to camera placement. As the optimal height for most mesocarnivores is 2-3 feet off

the ground (Jacobs and Ausband 2018), smaller or larger bodied species would only be partly

captured, which may create more difficulty for users completing identifications.

We also found accuracy varied by study area; though, what specific attributes in

geography accounted for these differences is unknown. In addition to habitat characteristics and

environmental conditions, the emergent biotic properties such as range shift of mesocarnivores

and proximity to urbanization can influence rarity and behavior, altering both the frequency of

detection and consequently diminishing user accuracy. Although it is difficult to disentangle the

individual components of differences in accuracy between geographies, additional considerations

in sampling design could offset differences in study area. For example, additional consideration

for standardizing sampling efforts by introducing measuring devices for animal size reference,

altering camera height, focusing more on rare species identification in training, and placing

multiple cameras at a single location to capture different angles for a species could help account

for classification differences related to study area characteristics (Hofmeester et al. 2019).

Overall, given the sensitivity of accuracy based upon species and study area underscores the

necessity of developing a flexible workflow to process and aggregate user-derived data from

camera surveys.

In evaluating our consensus hierarchy, we found that lower threshold consensus criteria

such as the third ranking kappa-evenness metric performed nearly as well as 100% agreement.

The high accuracy of images classified by these metrics (kappa, kappa and evenness) suggests

that hierarchical consensus criteria could be useful in mitigating potential bias from volunteers

with varying levels of expertise. By including lower thresholds of kappa and evenness, both of

which were more robust to incorrect classifications when aggregating responses, a reliable final

classification can still be reached even with multiple individual errors. For researchers, this

means that they can be less selective about their volunteer pool and invest less in training and

more in data collection or analysis. Given a reasonable mix of expertise in the volunteer pool,

more difficult species can still attain sufficient kappa and evenness thresholds from agreement

among a smaller number of correct classifications from experienced volunteers. While less

experienced members are still valuable in accurately classifying more common species, which

may comprise a large proportion of the total images.

Our work highlights the contributions and limitations of untrained participants to

accurately classify images of Michigan mammals. We found that most species were eventually

correctly identified by employing multiple consensus methods through a robust hierarchical

workflow. As a result, we are now confident in lowering the classification per image threshold to

below 15 users. Although researchers may find it helpful in future works to first set consensus

criteria based on conservative estimates of user knowledge. Future research is needed to

determine the degree to which demographic attributes, motivations, environmental stewardship

capacity and media content impact classification accuracy and public participation.

MANAGEMENT IMPLICATIONS

We found collaborating with the public was fruitful for investigating mammals of Michigan– a

typical species assemblage found throughout the contiguous United States. We suggest

highlighting distinguishing features of species through interactive field guides and species

comparisons tools to improve accuracy of user-derived data. Our findings underscore that even

with a relaxed criterion of user agreement most species were well represented in the final species

classifications. Using a hierarchical workflow with multiple thresholds for reaching consensus to

classify images will allow managers to harness a broad volunteer community to maximize

workload reduction without compromising accuracy.

ACKNOWLEDGMENTS

First, we recognize implementing our field research of camera surveys was conducted on lands

originally belonging to the People of the Three Fires. We thank the Shiawassee National Wildlife

Refuge, the University of Michigan Biological Station, and the Huron Mountain Club for

providing access to implement camera surveys for fieldwork on their property. We would like to

thank the Applied Wildlife Ecology (AWE) Lab at the University of Michigan for assistance

with validating classification. Our work is not human subjects research requiring IRB review,

though we remain forever indebted to the nearly 4,000 anonymous volunteers that contributed

their time to identify images on *Michigan ZoomIN (MZI)* and the dozens of colleagues that

integrated *MZI* into curricula. Our publication used data generated via the Zooniverse.org

platform, development of which was funded by generous support including a Global Impact

Award from Google and by a grant from the Alfred P. Sloan Foundation. We appreciate

technological support from University of Michigan LSA Information Technology and

Technology Services teams especially A. Roelofs and P. Knoop. We thank G. Kuhnlein,

Michigan News, and the University of Michigan Museum of Natural History for assistance with

dissemination on multi-media platforms to recruit volunteers. We appreciate the funding

provided by the Detroit Zoological Society, the Huron Mountain Wildlife Foundation, and the

University of Michigan Biological Station.

AUTHOR CONTRIBUTIONS:

370 N.C.H. conceived and designed the research. N.C.H and J.S. developed virtual interface for

371 species identification by public and expert confirmation. G.I.G. and T.C. constructed outreach

372 materials and led trainings on *Michigan ZoomIN*. G.I.G. and R.M. analyzed the data. G.I.G. and

373 N.C.H. wrote the manuscript. All authors contributed to fieldwork, image validation, public

374 engagement and editing the manuscript for final approval.

LITERATURE CITED

Arts, K., R. van der Wal, and W. M. Adams. 2015. Digital technology and the conservation of
nature. Ambio 44:661–673.
Bonney, R., T. B. Phillips, H. L. Ballard, and J. W. Enck. 2016. Can citizen science enhance
public understanding of science? Public Understanding of Science 25:2–16.
Broeckhoven, C., and le Fras Nortier Mouton, P. 2015. Some Like It Hot: Camera Traps Unravel
the Effects of Weather Conditions and Predator Presence on the Activity Levels of Two
Lizards. PloS one, 10(9):e0137428.
Brown, G., McAlpine, C., Rhodes, J., Lunney, D., Goldingay, R., Fielding, K., and Vass, L.
2018. Assessing the validity of crowdsourced wildlife observations for conservation
using public participatory mapping methods. Biological Conservation 227:141–151.
Burton, A. C., E. Neilson, D. Moreira, A. Ladle, R. Steenweg, J. T. Fisher, E. Bayne, and S.
Boutin. 2015. REVIEW: Wildlife camera trapping: a review and recommendations for
linking surveys to ecological processes. Journal of Applied Ecology 52:675–685.
Colborn A. S., Kuntze C. C., Gadsden G. I., Harris N. C. 2020. Spatial variation in diet-microbe
associations across populations of a generalist North American carnivore Journal of
Animal Ecology 89:1952-1960.
Crall, A. W., G. J. Newman, T. J. Stohlgren, K. A. Holfelder, J. Graham, and D. M. Waller.
2011. Assessing citizen science data quality: an invasive species case study. Conservation
Letters 4:433–442.
Danielsen, F., N. D. Burgess, and A. Balmford. 2005. Monitoring Matters: Examining the
Potential of Locally-based Approaches. Biodiversity & Conservation 14:2507–2542.
Dickinson, J. L., B. Zuckerberg, and D. N. Bonter. 2010. Citizen Science as an Ecological
Research Tool: Challenges and Benefits. Annual Review of Ecology, Evolution, and
Systematics 41. 41:149–172
Egerer, M., B. B. Lin, and D. Kendal. 2019. Towards better species identification processes
between scientists and community participants. Science of the Total Environment
694:133738.

Freitag, A., and M. J. Pfeffer. 2013. Process, Not Product: Investigating Recommendations for
Improving Citizen Science "Success". PLoS One 8:e64079.
Halpern, B. S., M. Frazier, J. Potapenko, K. S. Casey, K. Koenig, C. Longo, J. S. Lowndes, R. C.
Rockwood, E. R. Selig, K. A. Selkoe, and S. Walbridge. 2015. Spatial and temporal
changes in cumulative human impacts on the world's ocean. Nature Communications
6:7615.
Hochachka, W. M., D. Fink, R. A. Hutchinson, D. Sheldon, W. K. Wong, and S. Kelling. 2012.
Data-intensive science applied to broad-scale citizen science. Trends in Ecology &
Evolution 27:130–137.
Hofmeester, T. R., J. P. G. M. Cromsigt, J. Odden, H. Andrén, J. Kindberg, and J. D. C.
Linnell. 2019. Framing pictures: A conceptual framework to identify and correct for
biases in detection probability of camera traps enabling multi-species comparison.
Ecology and Evolution 9:2320– 2336.
Jacobs, C. E., and D. E. Ausband. 2018. An evaluation of camera trap performance – What are
we missing and does deployment height matter? Remote Sensing in Ecology and
Conservation 4:352–360.
Katrak-Adefowora, R., Blickley, J.L. and Zellmer, A.J. 2020. Just-in-Time Training Improves
Accuracy of Citizen Scientist Wildlife Identifications from Camera Trap Photos. Citizen
Science: Theory and Practice 5:8.
Landis, J. R., and G. G. Koch. 1977. The measurement of observer agreement for categorical
data. Biometrics 33:159–174.
Lintott, C. 2019. The Crowd and the Cosmos: Adventures in the Zooniverse. OUP Oxford
Oxford, United Kingdom.
Locke, C. M., C. M. Anhalt-Depies, S. Frett, J. L. Stenglein, S. Cameron, V. Malleshappa, T.
Peltier, B. Zuckerberg, and P. A. Townsend. 2019. Managing a large citizen science
project to monitor wildlife. Wildlife Society Bulletin 43:4–10.
Maes, D., N. J. B. Isaac, C. A. Harrower, B. Collen, A. J. van Strien, and D. B. Roy. 2015. The
use of opportunistic data for IUCN Red List assessments. Biological Journal of the
Linnean Society 115:690–706.
McShea, W. J., Forrester, T., Costello, R., Zhihai He., and Roland Kays. 2016. Volunteer-run
cameras as distributed sensors for macrosystem mammal research. Landscape Ecology
31:55–66.
Meek, P., Ballard G, and Fleming, P. 2015. The pitfalls of wildlife camera trapping as a survey
tool in Australia. Australian Mammalogy 37:13–22.
Meek, P., Ballard, G., Fleming, P. and Falzon, G. 2016. Are we getting the full picture? Animal
responses to camera traps and implications for predator studies. Ecology and Evolution 6:
3216–3225.
Norouzzadeh, M. S., A. Nguyen, M. Kosmala, A. Swanson, M. S. Palmer, C. Packer, and J.
Clune. 2018. Automatically identifying, counting, and describing wild animals in camera-
trap images with deep learning. Proceedings of the National Academy of Sciences
115:E5716-E5725.
Parrish, J. K., T. Jones, H. K. Burgess, Y. He, L. Fortson, and D. Cavalier. 2019. Hoping for
optimality or designing for inclusion: Persistence, learning, and the social network of
citizen science. Proceedings of the National Academy of Sciences 116:1894–1901.

Peterson, M. N., K. T. Stevenson, and D. F. Lawson. 2019. Reviewing how intergenerational
learning can help conservation biology face its greatest challenge. Biological
Conservation 235:290–294.
Pielou, E. C. 1966. The measurement of diversity in different types of biological collections.
Journal of Theoretical Biology 13:131–144
Potter, L. C., C. J. Brady, and B. P. Murphy. 2019. Accuracy of identifications of mammal
species from camera trap images: A northern Australian case study. Austral Ecology
44:473–483.
Prugh, L. R., C. J. Stoner, C. W. Epps, W. T. Bean, W. J. Ripple, A. S. Laliberte, and J. S.
Brashares. 2009. The rise of the mesopredator. Bioscience 59:779–791.
Raudsepp-Hearne, C., G. D. Peterson, and E. M. Bennett. 2010. Ecosystem service bundles for
analyzing tradeoffs in diverse landscapes. Proceedings of the National Academy of
Sciences 107:5242–5247.
Ries, L., and K. Oberhauser. 2015. A Citizen Army for Science: Quantifying the Contributions
of Citizen Scientists to our Understanding of Monarch Butterfly Biology. Bioscience
65:419–430.
Schuttler, S. G., R. S. Sears, I. Orendain, R. Khot, D. Rubenstein, N. Rubenstein, R. R. Dunn, E.
Baird, K. Kandros, T. O'Brien, and R. Kays. 2018a. Citizen Science in Schools: Students
Collect Valuable Mammal Data for Science, Conservation, and Community Engagement.
Bioscience 69:69–79.
Schuttler, S. G., A. E. Sorensen, R. C. Jordan, C. Cooper, and A. Shwartz. 2018b. Bridging the
nature gap: can citizen science reverse the extinction of experience? Frontiers in Ecology
and the Environment 16:405–411.
Shipley, J. R., J. F. Kelly, and W. F. Frick. 2018. Toward integrating citizen science and radar
data for migrant bird conservation. Remote Sensing in Ecology and Conservation 4:127–
136.
Skarlatidou, A., A. Hamilton, M. Vitos, and M. Haklay. 2019. What do volunteers want from
citizen science technologies? A systematic literature review and best practice guidelines. .
Journal of Science Communication 18(01):A02
Steenweg, R., M. Hebblewhite, R. Kays, J. Ahumada, J. T. Fisher, C. Burton, S. E. Townsend, C.
Carbone, J. M. Rowcliffe, J. Whittington, J. Brodie, J. A. Royle, A. Switalski, A. P.
Clevenger, N. Heim, and L. N. Rich. 2017. Scaling-up camera traps: monitoring the
planet's biodiversity with networks of remote sensors. Frontiers in Ecology and the
Environment 15:26–34.
Steger, C., Butt, B. and Hooten, M.B. 2017. Safari Science: assessing the reliability of citizen
science data for wildlife surveys. Journal of Applied Ecology 54: 2053–2062.
Swanson, A., M. Kosmala, C. Lintott, and C. Packer. 2016. A generalized approach for
producing, quantifying, and validating citizen science data from wildlife images.
Conservation Biology 30:520–531.
Swanson, A., M. Kosmala, C. Lintott, R. Simpson, A. Smith, and C. Packer. 2015. Snapshot
Serengeti, high-frequency annotated camera trap images of 40 mammalian species in an
African savanna. Scientific Data 2:150026.
Theobald, E. J., A. K. Ettinger, H. K. Burgess, L. B. DeBey, N. R. Schmidt, H. E. Froehlich, C.
Wagner, J. HilleRisLambers, J. Tewksbury, M. A. Harsch, and J. K. Parrish. 2015.
Global change and local solutions: Tapping the unrealized potential of citizen science for
biodiversity research. Biological Conservation 181:236–244.

Trumbull, D. J., R. Bonney, D. Bascom, and A. Cabral. 2000. Thinking scientifically during
495   participation in a citizen-science project. Science Education 84:265–275.
496 Waetjen, D.P., Shilling, F.M., 2017. Large extent volunteer roadkill and wildlife observation
497   systems as sources of reliable data. Frontiers in Ecology and Evolution 5:89
498 Young, H. S., D. J. McCauley, M. Galetti, and R. Dirzo. 2016. Patterns, Causes, and
499   Consequences of Anthropocene Defaunation. Annual Review of Ecology, Evolution and
500   Systematics 47:333–358.

502 *Associate Editor:*

Figure Captions

Figure 1. *Michigan ZoomIN* interface for public user classifications of mammals from camera
surveys throughout Michigan. 1) welcome screen with tutorial, 2) sample images active on talk
board from public comments, 3) image for user identification, 4) list of species to select for
classification, 5) filter features to narrow species selections, 6) field guide to provide additional
information, 7) reference photo of user selection to compare to image, and 8) additional
questions to provide more information concerning the image.
531
Figure 2. *Michigan ZoomIN* workflow from user-derived input to consensus assessment of
aggregated responses to expert validation necessary to create a final dataset ready for analysis of
classifying images. Images classified by an expert are tagged with gold standard which
immediately passed consensus and sent to the final data set. All other images went through a
hierarchical process with 15 different users classifying each image. Images that have complete
agreement among users went to the final dataset. If there is not unanimous agreement, different
algorithms were calculated from user responses to determine if they meet the Kappa threshold
and if not the kappa evenness threshold. Images that do not meet any consensus went into an
expert review file, which was then classified by a research team member to determine species.
541
Figure 3. Distribution of images for each species reaching consensus across various criteria.
Agreement resulted when all 15 users agreed on the species identification. Kappa was based on a
Fleiss Kappa statistic above 0.8, while Kappa and evenness used Fleiss Kappa statistic above 0.6
combined with Pielou's evenness below 0.5. Majority represented the most widely chosen
species by public users when consensus not reached by other methods. Gold standard indicated
when an expert contributed to the image classification, but only represented <1% of data.
548
Figure 4. Mammals identified by public volunteers through *Michigan ZoomIN*. Species in green
panel obtained ≥90% accuracy: 1) Domestic cat- *Felis catus,* 2) Human- *Homo sapien,* 3)
Raccoon- *Procyon lotor,* 4) White-tailed deer- *Odocoileus virginianus,* 5) Striped skunk-
Mephitis mephitis, 6) Opossum- *Didelphis virginiana,* 7) Squirrel- *Sciurus carolinensis,* 8)
Bobcat- *Lynx rufus, 9)* Coyote- *Canis latrans,* 10) Porcupine- *Erethizon dorsatum*; species in
yellow panel obtained 80-89% accuracy: 11) Chipmunk- *Tamias striatus*, 12) Black bear- *Ursus*
americanus, 13) Cottontail rabbit- *Sylvilagus floridanus*, 14) Groundhog- *Marmota monax*, 15)
Domestic dog- *Canis lupus familiaris*; and species in the red panel accuracy < 79% accuracy: 16)
Snowshoe hare- *Lepus americanus*, 17) American badger- *Taxidea taxus*, 18) Small mammal-
e.g., *Peromyscus maniculatus*, 19) Marten- *Martes americana*, 20) Fisher- *Pekania pennanti*, 21)
Muskrat- *Ondatra zibethicus*, 22) Weasel- e.g., *Mustela frenata*, 23) Grey wolf- *Canis lupus*,
24) Red fox- *Vulpes vulpes*, 25) Mink- *Neovison vison*. Gray fox- *Urocyon cinereoargenteus* not
shown.

Table 1 Top models presented with ΔAIC < 3 from model selection analysis for accuracy
assessment from comparison between user derived classification and expert validation.

Model	df	logLik	Δ AIC	W
Consensus method + Percent user agreement + Study area	8	-570.335	0.00	0.493
Consensus method + Percent user agreement + Study area + Camera model	9	-570.327	1.99	0.182
Consensus method + Percent user agreement + Species	30	-549.265	2.03	0.179
Consensus method + Percent user agreement + Species + Study Area	32	-547.451	2.43	0.146

596   Figure 1

623   Figure 2

650   Figure 3

677   Figure 4

694   **Summary for online Table of Contents**: Engaging the public proved extremely helpful in

695   processing images of mammals from a large scale camera survey in Michigan, given their high

696   accuracy in species identification. We found that study area, consensus method, and user support

697   best explained accuracy when validating classification from our public science project.