Neuro-computational mechanisms of action-outcome learning under moral
conflict

L. Fornari[1§], K. Ioumpa[1§], A. D. Nostro[1ǂ], N. J. Evans[2ǂ], L. De Angelis[1], R. Paracampo[1], S.
Gallo[1], M. Spezio[3], C. Keysers[1,4,#] & V. Gazzola[1,4,#,*]

[1] *Netherlands Institute for Neuroscience, KNAW, Meibergdreef 47, 1105BA Amsterdam, The Netherlands*
[2] *School of Psychology, University of Queensland, Australia*
[3] *Psychology, Neuroscience, & Data Science, Scripps College, 1030 Columbia Ave, Claremont, CA 91711*
Claremont, CA USA
[4] *Dep. of Psychology, University of Amsterdam, Nieuwe Achtergracht 129-B, 1018 WT Amsterdam, The*
Netherlands

[§,ǂ,#] Equal contribution
Corresponding author: Valeria Gazzola, Meibergdreef 47, 1105BA Amsterdam,
v.gazzola@nin.knaw.nl, +31205661716

Classification: Biological sciences, Psychological and Cognitive Sciences

Keywords: reinforcement learning theory, prosocial behavior, empathy, pain, reward, mirror
neurons, simulation theory

Author contribution: The experiments were conceived by VG and CK with input from all
authors. Funding and project leadership by VG. The fMRI data were collected by KI with the
help of SG; the behavioral data by LF. Pilot data for the preparation of the study were obtained
by RP, AN and SG. FMRI data were analyzed by KI with guidance from VG and CK; the RLT
models were developed and programmed by AN, CK, LD, LF, MS, NE, and VG. AN, CK, KI,
LF, MS, and VG wrote the manuscript with edits and comments from all other authors.

Abstract

Predicting how actions result in conflicting outcomes for self and others is essential for social
functioning. We tested whether Reinforcement Learning Theory captures how participants
learn to choose between symbols that define a moral conflict between financial self-gain and

other-pain. We tested whether choices are better explained by model-free learning (decisions based on combined historical values of past outcomes), or model-based learning (decisions based on the current value of separately expected outcomes) by including trials in which participants know that either self-gain or other-pain will not be delivered. Some participants favored options benefiting themselves, others, preventing other-pain. When removing the favored outcome, participants instantly altered their choices, suggesting model-based learning. Computational modelling confirmed choices were best described by model-based learning in which participants track expected values of self-gain and other-pain separately, with an individual valuation parameter capturing their relative weight. This valuation parameter predicted costly helping in an independent task. The expectations of self-gain and other-pain were also biased: the favoured outcome was associated with more differentiated symbol-outcome probability reports than the less favoured outcome. FMRI helped localize this bias: signals in the pain-observation network covaried with pain prediction errors without linear dependency on individual preferences, while the ventromedial prefrontal cortex contained separable signals covarying with pain prediction errors in ways that did and did not reflected individual preferences.

Introduction

We often have to learn that certain actions lead to favorable outcomes for us, but harm others, while alternative actions are less favorable for us but avoid or mitigate harms to others (1). Much is already known about the brain structures involved in making moral choices when the relevant action-outcome contingencies are well known (2–9), but how we *learn* these contingencies remains poorly understood, especially in situations pitting gains to self against losses for others.

Reinforcement learning theory (RLT) has successfully described how individuals learn to benefit themselves (10, 11) and most recently, how they learn to benefit others (12–15). At the core of reinforcement learning is the notion that we update expected values (EV) of actions via prediction errors (PE) – the differences between actual outcomes and expected values represented in mind.

Ambiguity in morally relevant action-outcome associations raises specific questions with regard to RLT, especially if outcomes for self and others conflict. If actions benefit the self and harm others, are these conflicting outcomes combined into a common valuational

representation as in model-free/habit learning; or do we track separate expectations for benefits

to the self and harm to others, as in model-based/goal-directed learning (16)? In addition,

people differ in how they represent benefits and harms to self (17), and in whether they prefer

to maximize benefits for the self vs. minimizing harms to others (3, 4, 6). How can such

differences be computationally represented using RLT? Would people emphasizing one aspect,

such as the harm to others, already show increased prediction errors and expected value signals

when confronted with this outcome type, or are expectations tracked independently of one's

preferences, such that preferences only play out when decisions are being taken?

To address these questions, we performed two experiments: an Online experiment with

79 and an fMRI experiment with 27 participants (Table 1). In the core task common to both

experiments, participants learned morally relevant action-outcome associations over different

blocks of 10 trials each (Figure 1A, C; Table 1). In each block, participants saw two new

symbols and faced a conflictual situation (Conflict condition): one symbol led to high monetary

gains for the self 80% of the time, and to a painful but tolerable shock to the hand of a

confederate with the same probability. We refer to this symbol as 'lucrative', since it was

associated with higher monetary outcomes. The other symbol led to low monetary gains for

the self 80% of the time, and to non-painful shocks to the confederate with the same probability.

We refer to this symbol as 'pain-reducing'. Importantly, to partially de-correlate

representations of shock and money, the probabilities of high and low monetary reward and

pain and no-pain to others were drawn independently. At the beginning of each block,

participants did not know the associations between symbols and outcomes. Choosing which

symbol best satisfies the moral values that participants act upon in the task thus involves

learning to predict the outcomes associated with each symbol.

In the online experiment, participants performed a number of additional tasks. First, to

explore whether participants learned the symbol-outcome association probabilities for both the

self-money and other-shocks, even if their preferences may emphasize one, we added blocks

in which we crossed the association to remove the conflict. In this No-Conflict condition

(Figure 1B), the symbol that led to high money in 80% of cases also led to low shocks in 80%

of the cases, and the symbol leading to low money in 80% of the cases also led to high shocks

in 80% of the cases. After each of these blocks we then explicitly asked participants to report

the learned associations (Figure 1C). Only if participants learn symbol-outcome association for

the less preferred outcome, will they report different probabilities under the Conflict and

NoConflict for this outcome. Second, to distinguish model-based from model-free learning, we

adapted the devaluation approach pioneered in animals (16). In a classic example of this

approach (18), animals learned to press a lever for an attractive food A. All rats then underwent
taste aversion learning, but for some, food A, the one they worked for, was paired with a
lithium-chloride injection, for others the injection was paired with a different food B. When
both groups could later press the lever again, only those for whom food A had been devalued
reduced their usage of the lever. This showed their behavior was model-based: driven by an
understanding that the lever didn't simply lead to rewarding states, but to food A - a food that
had lost its value and wasn't worth working for anymore (18). Here, we therefore added blocks
(DropOut condition, Figure 1D) in which after 10 trials to learn the symbol-outcome
associations, we informed participants that the self-money or the other-shocks would not be
delivered on the following trials. We then examined the choice on the 11^{th} trial, before
participants witnessed the modified outcome. If participants use a model-based strategy, we
expect them to show different choices depending on which outcome is removed, with choices
expected to change substantially from the 10^{th} to the 11^{th} trial, if the outcome they weigh more
is removed. If participants use a model-free strategy, as has recently been suggested using a
two-stage decision-task(15), we expect them to continue choosing their previously favoured
option. The comparison of participant's choices at the 11^{th} trials between the Conflict and No-
Conflict conditions, further informs us on their ability to form separate representations for
money and shocks.

For the participants in the fMRI experiment, beside the core learning task, we also
performed a different costly helping task previously developed in our group (4), to test whether
parameters we extract from our conflictual learning task can predict choices in a different costly
helping task not requiring learning (Supplementary Material §3).

After describing participants' behavior, we used Hierarchical Bayesian Model
comparisons to test which alternative computational formulations of the RLT models
(Figure 2) better describe participant's choices, and gain insights into how people combine the
two outcomes in their morally relevant learning experience. Except in our random choice
model (M0), in all the learning models we compare, shock and money are additively combined
using an individual weighting factor *wf* ranging from 0 to 1. This *wf* captures the value of the
monetary outcome for self relative to the value of shock to the other and is not unlike the
salience alpha in the Rescorla-Wagner Learning Rule (19, 20). A *wf* closer to zero would
characterize a participant preferring to minimize the harm to the other person, a *wf* closer to
one corresponds to more lucrative choices. In particular, we investigate whether choices are
better predicted (i) by an RLT model that combines money and shock as soon as outcomes are
revealed (M1) or (ii) models that keep separate representations for the two types of outcomes

(M2). M1 instantiates model-free learning in which participant's decisions are based on the
history of past reward values, without representation of the nature of outcomes, while M2
instantiates model-based learning with separable representations of the expected outcomes for
self and others. For M2, we further compared a variant that scales outcomes based on personal
preferences for money vs. shock (M2Out) vs. a variant that initially tracks expectations
independently of personal preferences regarding outcomes, but that introduces weights at the
decision phase of the task (M2Dec). We expect all but M0 to perform comparably well during
the conflict phase of our experiment. We however expect that if participants use model-based
learning under conflict, M2 should outperform M1 during the 11^{th} trial when one outcome is
removed. We had no specific predictions about whether M2Dec or M2Out would predict
choices at the 11^{th} trial more accurately.

Shocks were made visible to the participants through pre-recorded videos showing
facial expressions from the confederate. In the fMRI experiment participants believed these
recordings to be part of a real-time video-feed, while in the Online experiment they were aware
that the videos had been pre-recorded, but believed that after the task we would select 15 trials
and administer the corresponding shocks to another participant in the lab. We used this video-
feedback, instead of the symbolic feedback more often used in neuroeconomic paradigms, to
explore the neural systems activated in situations where the consequences of our actions are
only available from the facial expressions of the people around us, and to remain closer in
design to the paradigms used in nonhuman animal models (21).

Finally, we used the neuroimaging data to inform our understanding of how our
participants update values in our tasks. Given substantial literature on how individuals learn to
maximize gains for the self (22, 23), we focus on how the brain updates values based on shocks
to others. In particular, given the conceptual difference between M2Out and M2Dec, we ask
whether we can distinguish and localize areas in which BOLD signals covary with prediction
errors in ways that do vs. do not depend on *wf*. More specifically, given an extensive literature
that has outlined a network of brain regions recruited while witnessing painful vs non-painful
facial expressions (24–27), and the availability of a multivariate signature to quantify activity
in this network (28), we first ask whether activity in this network covaries with prediction errors
for shocks, and if so, whether it does so in ways that do or do not depend linearly on *wf*. Second,
we perform a more explorative analysis to localize where in the brain the BOLD signal covaries
with pain prediction errors in ways that do or do not significantly depend linearly on *wf*. Recent
reviews identified that the ventromedial prefrontal cortex (vmPFC) has BOLD signals that
covary positively with the current value of multiple outcomes, in particular for chosen options

(22, 29), and outcomes for others appear to also be encoded in this region (30, 31).
Interestingly, when switching between making choices to benefit the self and others, the most
ventral medial prefrontal cortex had BOLD signals covarying with the value that was at the
focus of a particular trial while a more dorsal part covaried with the irrelevant value (32). In
our task we may thus expect the ventral mPFC to have signals that covary with prediction errors
for shocks in ways that depend on *wf* (and at thus at the focus of a particular trial) while more
dorsal mPFC regions may also represent prediction errors for shocks in ways that do not depend
on *wf* (which may become relevant in the future if contingencies change).

176
177 **Figure 1. Learning Task. (A)** Trial structure. Red outline indicates an exemplary choice for this trial. The picture
178 showing the confederate's response for the outcome is a still frame captured from one of the videos used in the
179 online experiment and in which the confederate received a painful shock (see Supplementary Materials §1 for

A   Choice and value updating during the first 10 trials

	Choice in Decision phase	Outcome phase if [shock] is chosen	
M0	categorical_logit([0,0])		
M1	cat_logit(τ*[EV[face], EV[shock]])	PE[shock] = (Out$_M$*wf + Out$_S$*(1-wf)) - EV[shock]	EV[shock] = EV[shock] + λ*PE[shock] EV[face] = EV[face]
M2 Out	cat_logit(τ*[EV[face]$_M$ + EV[face]$_S$, EV[shock]$_M$ + EV[shock]$_S$])	PE[shock]$_M$ = (Out$_M$*wf) - EV[shock]$_M$ PE[shock]$_S$ = (Out$_S$*(1-wf)) - EV[shock]$_S$	EV[shock]$_M$ = EV[shock]$_M$ + λ_M*PE[shock]$_M$ EV[shock]$_S$ = EV[shock]$_S$ + λ^*_SPE[shock]$_S$
M2 Dec	cat_logit(τ*[EV[face]$_M$*wf + EV[face]$_S$*(1-wf), EV[shock]$_M$*wf + EV[shock]$_S$*(1-wf)])	PE[shock]$_M$ = Out$_M$ - EV[shock]$_M$ PE[shock]$_S$ = Out$_S$ - EV[shock]$_S$	EV[face]$_M$ = EV[face]$_M$ EV[face]$_S$ = EV[face]$_S$

B   Choice in Decision phase at the 11th trial

	when money is removed	when shock is removed
M0	categorical_logit([0,0])	categorical_logit([0,0])
M1	cat_logit(τ*[EV[face], EV[shock]])	cat_logit(τ*[EV[face], EV[shock]])
M2 Out Dec	cat_logit(τ*[EV[face]$_S$ + EV[shock]$_S$])	cat_logit(τ*[EV[face]$_M$ + EV[shock]$_M$])

191
192

Figure 2. Tested RLT Model. (A) Model formalization for the symbol chosen in Figure 1a. In all our models,
decisions are based on a categorical logit function (aka softmax), contrasting the expected value across the
alternatives with inverse temperature τ [0,5] such that for a given difference in expected value, participants with
higher τ choose the symbol with higher expected value more often. The learning rate λ [0,1] captures the speed
at which expected values are updated, with higher λ values indicating that expected values are more strongly
updated following surprising outcomes. EV=Expected Value, PE=Prediction Error, Out=Outcome. Subscript
M=money, Subscript S=shocks. Outcomes are coded based on their value, such that high-shock OutS=-1, low-
shock OutS=+1, high-Money OutM=+1, low-money OutM=-1. Trials in which one outcome is withheld during
dropout is coded as Out=0. Note that outcomes are set to the same values (+1 and -1) for money and shock because
we chose the amount of money at stake to be close to the indifference point (see Supplementary Materials §2).
(B) Model formalization for the symbol chosen in Figure 1a during the decision phase at the 11th trial for the
DropOut condition.
205

Results

Participants vary in their preference but most favour one symbol in the first 10 trials of a block

Figure 3A-B shows participants' choices in the Conflict condition (see also Table 1). Using a binomial distribution, with 60 trials for the fMRI experiment (10 trials x 6 blocks), and 120 trials for the Online experiment (10 trials x 12 blocks), we found that in the fMRI experiment about half the participants chose the pain-reducing option above chance, about half were within what would be expected without preference, and just 3 chose the lucrative option above chance. For shorthand, we refer to the choice patterns in these subgroups, divided by the binomial probability, as 'considerate', 'ambiguous' or 'lucrative' preferences. In the Online experiment, participants were evenly subdivided across these three preference subgroups. Averaging the learning curves of participants showing opposite preferences would occlude learning, and Figure 3B thus visualizes the average choices for each group separately. The groups with lucrative and considerate choice preferences show typical learning curves, starting at chance (50% preference for the pain reducing option) on the first trial of each block and then gravitating towards the pain reducing (for considerate) or lucrative (for lucrative preferences) option, with the last four trials showing relatively stable preferences around 80%, as would be expected for an RLT with 80/20 probability. The curve of the group with ambiguous preference, consistently remains around 50% showing no clear learning curve (Figure 3B, orange and yellow). Finally, participant's average choices did not seem to be influenced by how much participants believed in the shock delivery (Supplementary Materials §4), and were in agreement with participant's reported motivation (Supplementary Materials §5).

		Online Exp.	fMRI Exp.
Participants	Total participants	79	27
	Average Age ± SD	25y±7	37y±17
	Females, males	39,40	27,0
	Subj. with Considerate preference	29 (37%)	13 (48%)
	Subj. with Lucrative preference	24 (30%)	3 (11%)
	Subj. with Ambiguous preference	26 (33%)	11 (41%)
	Conflict	4Bx10$_{Conf}$T /B	6Bx10$_{Conf}$T /B
	No-Conflict	4Bx10$_{NoConf}$T /B	n.a.

Conditions in the Learning task	Explicit learning task	8Tx4Questions	n.a.
	Conflict Money Dropout	4Bx(10$_{Conf}$T/B+10$_{Drop}$T/B)	n.a.
	Conflict Shock Dropout	4Bx(10$_{Conf}$T/B+10$_{Drop}$T/B)	n.a.
	No-Conflict Money Dropout	4Bx(10$_{NoConf}$T/B+10$_{Drop}$T/B)	n.a.
	No-Conflict Shock Dropout	4Bx(10$_{NoConf}$T/B+10$_{Drop}$T/B)	n.a.
Helping Task	Helping Blocks	n.a.	30T

230
231
Table 1. Participants and Task-conditions overview. For each experiment, the table reports the total number of
participants included in the behavioural analysis, and how many of those fell within the Considerate, Lucrative
and Ambiguous preference pattern as calculated by the Binomial distribution. Additionally, the table reports the
number of repetition of each condition included in that particular experiment. T=trial; B=block; Conf=Conflict;
NoConf=NoConflict; Drop=Dropout. fMRI data only included the right handed participants, 25 (35y±15SD; 25f),
while behavioural data were collected for the 27 indicated in the table. Note: although age differed between the
fMRI and Online experiment (Mann-Whitney test, BF10=8.022, p<0.001), the choices did not differ significantly
(Kendall's tau(age, %Considerate choices)=0.113, BF$_{10}$=0.543, p=0.096).

240
241
242

Considerate Online · Considerate fMRI · Ambiguous Online · Ambiguous fMRI · Lucrative Online · Lucrative fMRI

Figure 3. Choice allocation in the Conflict condition. (A) Histogram showing the proportion of pain-reducing
choices per participant over the first 10 trials of all available blocks of the online (left, 12 blocks of 10 trials) or
fMRI (right, 6 blocks of 10 trials) experiment. For blocks with DropOut, only the first 10 trials, before DropOut
occurs, are included (black rectangles in Figure 1D). Participants with above chance proportion of pain-reducing
choices according to a binomial (*p*<0.05, considerate preference) are shown in dark (online) or light (fMRI) green,

participant with below chance (*p*<0.05, lucrative preference) are shown in dark (online) or light (fMRI) blue, and

participants within chance level (*p*>0.05, ambiguous preference) are shown in orange (online) or yellow (fMRI).

(B) Choices as a function of trial number within a block, averaged over blocks, separately for the three preference

subgroups, with error bars representing the s.e.m. over participants. Colour-code as in (A). As in (A) all first 10

trials of the Conflict conditions are included, independently of whether they belong to a DropOut or No-DropOut

block. For the comparison of choice allocations across conditions see Figure 3_figure supplement 1.

Figure 3_figure supplement 1. Choice allocation in DropOut and No-DropOut Conflict Online conditions.

(A) Average choices allocation per participant during the first 10 trials of the 8 blocks that will later lead to

DropOut (x-axis) and the 4 conflict No-DropOut blocks (y-axis). Pearson's correlation= 0.89, $t_{(77)}$=17.23, p<2.2e-

16, BF_{10}>1000; mean No-DropOut=0.501±0.025s.e m.; mean DropOut=0.512±0.023s.e m.; Paired sample test

$t_{(78)}$= 1.04, p=0.302; BF_{10}=0.208. Color code represents the subgroup classification including all trials. The dashed

lines represent the critical values for the binomial distribution *p*<0.05 separately for the DropOut and No-DropOut

blocks. Regression line in black with shaded s.e.m. **(B)** As in (A) for the first 10 trials of the 4 DropOut blocks in

which money will later be removed (x-axis), and the 4 DropOut blocks in which shock will later be removed (y-

axis). Pearson's correlation=0.78, p<0.001, BF_{10}>1000; mean DropOut Money=0.52±0.024s.e m.; mean DropOut

Shock=0.503±0.025s.e.m.; Paired sample test $t_{(78)}$=1.21, p=0.23, BF_{10}=0.25

Participants have explicit and separable access to the symbol-outcome associations for money and shock that are biased by preference

To probe participants' explicit learning, in the Online experiment we asked them to report how likely each symbol was associated with high-shock and high-money. Overall, participants tended to report higher probabilities for symbols with higher probability, also capturing the difference between the Conflict and NoConflict condition (comparing the *N* and *U* shapes in Figure 4A). Because choices are thought to be driven by the *difference* in expected value across options, we summarized explicit reports as the difference in reported probability for the favored minus the non-favoured symbol, with larger reported differences in the correct direction providing more evidence for (explicit) learning (Figure 4B). We observed that the reported differences in probability across the symbols was different from zero, and in the expected direction in all cases (All $BF_{10}>3$), showing that as a group, participants with considerate and lucrative preferences also learned something about the association with outcomes that they appear to be less interested in. Second, participants show evidence of noticing the Conflict vs No-Conflict difference, which they would not need to do if they focused exclusively on maximizing money or minimizing shocks: participants with considerate preference (green) reported lower association with high-shock for their preferred (shock-reducing) symbol in the conflict and no-conflict block (both 3rd and 4th violin below zero), and they reported that their favored symbol had lower high-money probability in the Conflict (1st violin) and higher high-money probability in the No-Conflict (2nd violin) than their non-preferred symbol. A similar pattern is true for participants with lucrative preference (blue): they reported that their favorite (money-maximizing) symbol had a higher high-money probability than their non-favored symbol after both Conflict and No-Conflict blocks, and that this symbol was more likely in Conflict and less likely in No-Conflict blocks to lead to high-shock than their non-preferred symbol. Also participants with ambiguous preference could report above chance differences in probability for the two symbols in the correct direction in all cases (Figure 4C).

Interestingly, the magnitude of explicitly reported probability difference was biased in favour of the outcome participants seem to weigh more strongly. When separating participants in preference-groups, we see that those choosing the pain-reducing option above chance show a stronger difference, that was also closer to the actual difference, in reported probability for the shock than money outcomes, and those choosing the lucrative option show a stronger difference, that was also closer to the actual difference, for the money than the shock outcomes

(difference between the filled green violins in Figure 4B for the considerate preference=

18.2 ± 4.45s.e.m, W=377.5, BF_{10}=577.22, p<0.001 and between the filled blue violins for the

lucrative=9.2 ± 3.25s.e.m., W=233, BF_{10}=7.67, p<0.018). Because the thresholds for grouping

participants in these three preference groups is somewhat arbitrary, we also examined whether

over all participants, the preference (calculated as proportion of pain-reducing choices) was

predictive of their bias, and this was the case (Figure 4D, Pearson's r= 0.51; $t_{(77)}$=5.2, p= 1.6 x

10^{-6}).

That ambiguous preference participants could still report above chance probability

estimates (Wilcoxon signed-ranked, all p<0.001, BF_{10}>100; Figure 4C) suggests that their lack

of above chance choice preference was not due to a total lack of symbol-outcome association

learning. A lack of clear preference for one outcome over the other seems also to have played

a role. Indeed, when the conflict was removed (Figure 4E orange), ambiguous preference

participants do demonstrate learning, but choose the favorable option less frequently than the

other two groups (Bayesian one way ANOVA, $F_{(2,76)}$=15.6, p<0.001; $BF_{incl\ group}$=8284.52,

Figure 4F). At the single subject level, in No-Conflict blocks, 54% of the ambiguous preference

subjects fell below the learning threshold determined using a binomial distribution (71/120

correct choices; red line in Figure 4F), while just 7% of those with considerate and 4% of those

with lucrative preference were below this threshold. The ambiguous preference group thus

appears to contain a mix of individuals that do not show significant learning in our task even

without conflict, and individuals that can learn without conflict but fail to display clear

preferences in the presence of conflict.

Figure 4. Participant reports and No-Conflict choices. (A) Participants' reported perceived high-outcome

probability for both symbols (see Fig. 1C) from the Conflict (left) and No-Conflict (right) conditions. The x-axis

specifies which symbol and question were asked: short/tall pile of money=reported high-money probability for

the symbol actually associated with low/high-money; painless/painful face=reported high-shock probability for

the symbol associated with low/high-shock. As a group, most participants correctly assigned higher probabilities

to symbols that had higher probability, and reported a different pattern of probabilities after Conflict and No-

Conflict blocks. Thick black lines = average of reported probabilities; dotted black lines = programmed/expected

probability. **(B)** Difference between reported probabilities for the preferred minus non-preferred symbol

separately for the Considerate (green) and Lucrative (blue) preference group and for the Conflict (filled violins)

and No-Conflict (empty violins) condition. The symbols on the x-axis specify whether the question was to estimate

the probability of leading to high-money (coins) or high-shock (face). Violin plots report median and quartiles.

The BF above a violin represents the result of a Wilcoxon signed rank test of the differences vs. zero; the BF

below a pair of violins represents the result of a Wilcoxon signed rank test comparing the conflict and no-conflict

conditions. Figure 4_figure supplement 1 guides the reader through the computations presented in this panel.

Dotted black lines = programmed probability. **(C)** Same as in (e) for the ambiguous group. **(D)** Correlation

between the participant's report bias and the proportion of pain-reducing choices during the conflict task, with

Bias= (rpS(high shock) - rpS(low shock)) - ((rpM(high money) - rpM(low money)), where rpS/rpM stand for
reported probability of high-Shock/high-Money outcome, and highshock, lowshock, highmoney, lowmoney refer
to the symbols based on their actually most likely outcome. **(E)** Choice in the No-Conflict condition. Note that
the classification in terms of preference is still based on the Conflict choices. **(F)** Proportion of favorable choices
(i.e. choosing the symbol most likely to lead to high-money and low-shock) separately for the three groups.
346

348 **Figure 4_figure supplement 1. Reading key for Figure 4B.** For each of the violin plots in Figure 4B, we here
349 show how the violin plot is computed based on the difference between the two symbols.
350

DropOut trials suggest model-based learning with separate outcome representations biased by preference

Examining the choices on the critical 11th DropOut trial, similar to a classic devaluation procedure, speaks in favour of a model-based learning by confirming that participants have separable representations of self-money and other-shock, and have also learned more about the outcome that drives their decisions than about the other outcome. We find that for the considerate and lucrative groups, which showed strong preferences on the 10th trial, removing the quantity that guided their choice most, led to a highly significant change in their choice allocation (Figure 5A, both $BF_{10}>80$), while removing the quantity that guided their choices less, left their choices unchanged (both $BF_{10}<1/3$).

Figure 5. Participant's choices at the 11th trial. (A) Choices in the Conflict DropOut blocks averaged separately for the preference subgroups as classified using all Conflict trials. For each subgroup, the solid line represents the average of all trials, the dashed line, blocks in which money will be withheld, and the dotted lines, blocks in which shocks will be withheld. BF_{10} values shown between the 10th and 11th trial indicate the result of a Bayesian paired t-test (i.e. a Wilcoxon signed-rank test on the differences) comparing choices in these two trials. BF_{10} value shown above the 10th or the 11th trial, the result of the same test comparing choices on trials where money will be or is withheld vs trials where shocks will be or are withheld. Error bars are the s.e m. across participants. **(B)** Violin plot of choices across participants on the 11th trial of conflict drop-out blocks as a function of preference subgroup and outcome withheld. The boxplot within each violin signifies the 4 quartiles of the data, the black dot represents one outlier.

Dropout trials in NoConflict blocks did not lead to comparable changes in choices
(Supplementary Figure 4; all 11^{th} vs 10^{th} trial $BF_{10}<2.1$), confirming that the change in choices
in the Conflict condition are not the result of a simple rule 'reset choice when my prefered
preferred outcome is removed, but not when my non-preferred outcome is removed', but
considers the probability of outcomes associated with the remaining outcome, which are the
only difference between Conflict and No-Conflict blocks.

Interestingly, removing the guiding outcome in Conflict trials did not lead to a mirror
symmetrical (i.e. ~80%) preference for the other symbol: the average choice allocation
switched from ~80% (considerate) or ~20% (lucrative) pain-reducing choices on the 10^{th} to
just below, or above, ~50%, on the 11^{th}, respectively (Figure 5A). That the first choice after
DropOut is not as polarized as the 10^{th} trial is in line with the bias we observed in the explicit
reports (Figure 4B,D): given that the difference in reported outcome probabilities was less
pronounced for the less-guiding outcome, one would expect less polarized choices in a decision
based on these lesser differences in expected values. For the ambiguous group, there was no
robust evidence that removing either quantities lead to a change of preference towards the
better option for the remaining outcome. It is important to emphasize that the choice on trial
11 occurs before the participant is presented with the single remaining outcome, and thus
probes choices exclusively based on expected values that were learned during conflict trials.
Supplementary Figure 4 also shows choices on trials 12-20, but as these trials do not contain
two potentially conflicting outcomes, but only one outcome, we do not expect our models to
adequately represent participants' reasoning and learning over those subsequence trials.
In summary, choices reveal substantial individual variability in preference, but most
participants show evidence of some form of learning. Explicit reports and their difference
across Conflict and No-Conflict blocks reveal that participants have explicit and separable
representations of outcome probability differences across symbols in terms of money and
shock, although these representations are more differentiated for the outcome that their choices
indicate to bear more weight. Finally, devaluation performed by informing participants that
their preferred outcome will no longer be delivered leads to an immediate switch away from
their previously preferred option during Conflict. Together this supports the notion that choices
in our conflictual conditions are dominated by model-based learning in which participants
represent separable outcomes for money to self and shock to others, with the individual
preferences influencing the magnitude of their respective symbol-outcome association. This

would suggest that M2Out (Figure 2) might be the computational implementation of RLT
learning most suitable to capture our participants' choices in our paradigm.
410

M2Out predicts participant's choices more accurately

Devaluation, as implemented in our conflict Drop-Out blocks, is critical to distinguish
model-based and model-free learning. To directly compare our neurocomputational models
(Figure 2A), we therefore fitted our competing models on the first 10 trials of the Conflict
DropOut blocks, and then examined (i) how well they predicted those first 10 choices using an
approximation of the leave-one-out information criterium (LOOIC(33)) and (ii) how well they
predicted choices on the 11^{th} trial that was not included into the model fitting, when one of the
outcomes is removed. Because the 11^{th} trial was not included in the fit, we can directly quantify
the likelihood of the observed choices on the 11^{th} trial under the different models to assess
which model best predicts behavior under devaluation. *A priori*, we expect the four models to
make very different predictions on the 11^{th} trial. M0, by design, continues to predict random
choices. Because M1 does not have separable representations of money and shock, its
predictions for the 11^{th} trial are simply based on the symbol with highest composite expected
value (EV) based on the first 10 trials, and thus predicts that choice behavior does not change
on the 11^{th} trial. Finally, because the M2 models have separate EV for money and shock, we
programed the model to transform the information that participants receive *before* they perform
the 11^{th} choice (i.e. one outcome will no longer be delivered), into a revised decision criterion
based on the remaining expected value only, without using a *wf* in their choice, as there is no
longer a conflict to resolve (Figure 2B). For M2Out and M2Dec, choice is then dictated for the
11^{th} trial by cat_logit(τ(EV$_R$(symbol1),EV$_R$(symbol2)), where subscript R represents the
remaining quantity: M (money) or S (shock). We thus expect choices not to change much if
the less preferred outcome is removed, and to change significantly, if the preferred outcome is
removed.
Figure 6A illustrates the predictions of the three learning models (M1, M2Out, M2Dec)
as compared with the actual choices for the critical DropOut blocks. For visualization purposes,
we did not include M0 and the ambiguous preference participants in this figure, because they
do not predict or show the learning behavior we are trying to capture. During the initial 10 trials
in which the conflict is present, all three learning models (M1, M2Out, M2Dec) capture the
general shape of the learning curve, and can accommodate individual differences in preference
through the weighting factor *wf*, generating increasingly more pain-reducing choices as the trial

number increases for participants with considerate preferences, and fewer pain-reducing

choices for participants with lucrative preference. Including all participants (including the

ambiguous participants), model comparison over the first 10 trials with the LOOIC confirms

that M1, M2Out and M2Dec perform similarly well, i.e. remain within a standard error of one-

another, and perform better than the random-choice model M0 (Figure 6B, and Supplementary

Figure 5 for results that exclude the ambiguous and non-believers groups). Note that the

information criterion (IC) scale captures how much information is lost when comparing model

predictions with actual choices, and smaller IC thus characterize models that better describe

behavior. A similar pattern is true for the choices in the fMRI experiment (Supplementary

Figure 5c). Next, we used the 11th trial to arbitrate across M1, M2Out and M2Dec. As we saw

in Figure 5, participants' actually responded to devaluation in ways that depended on their

preference. Lucrative preference participants kept their choices unchanged when shocks were

removed, but switched their preference to just above 50% pain reducing choices on average

when money was removed. The converse was true for Considerate preference participants.

When money is removed and only shocks remain (Figure 6B, gray background), all learning

models correctly predict that considerate preference participants do not change their choices,

but only M2Out (purple, and marked by the black arrow-head) accurately predicts that

participants with lucrative preference change their preference to just above 50%. In contrast,

M1 (gray) fails to predict any change for those lucrative participants, and M2Dec (light blue)

overestimates their change. When shocks are removed and only money remains (Figure 6B

yellow background), all learning models correctly predict a lack of change for the lucrative

preference participants, but again, only M2Out correctly predict the magnitude of change in

the considerate participants, that now choose the pain-reducing symbol just under 50% of cases

on average, while M1 fails to predict any change and M2Dec overestimates the change. When

considering all participants' choices (including the ambiguous), Figure 6C shows that M2Out

consistently predicts the choices on the 11th trial better than any of the other models. M2Out

also outperformed the other models when excluding ambiguous participants or those that

signaled some degree of scepticism about whether someone would really be administered

shocks (Supplementary Figure 5).

Figure 6C estimates the performance of the models at the group level, by adding the

log-likelihood of all participants and trials, and plotting this overall likelihood for each of the

4000 samples drawn from the posterior distribution. To compare models at the single subject

level, we extracted the mean log-likelihood of the 4000 samples for each participant and model,

and calculated the likelihood ratio K in favour of M2Out, against each of the other models, as

the ratio of their mean likelihoods. Using the traditional bounds for moderate evidence that

have become standard in Bayesian statistics ($K>3$(34)), in all cases we find more participants

with evidence in favour of M2Out than for the other models (Figure 6D).

That M2Out outperforms M2Dec in particular, is because M2Out scales expected

values based on *wf*. For instance, an extremely considerate participant has *wf*≈0, and the EV_M

thus remains close to zero for both symbols across the first 10 trials, because $OutM$ are

multiplied with this *wf*≈0 to calculate PE_M (Figure 2), and EV_M thus is the sum of many PE_M

close to zero. Accordingly, when shocks are removed, choices are between two symbols

withEV_M≈0, predicting the ~50% preferences observed. In contrast, for M2Dec, such scaling

of EV by *wf* does not occur, and on the 11^{th} trial, participants have EV values for their less-

favored outcome that are as differentiated as those of the favored outcome. Note that if we

simulate a version of M2Dec that would maintain *wf* in the decision-phase of the 11^{th} trial,

M2Dec would make predictions extremely similar to M2Out.

Figure 6. Model Comparison. **(A)** For the Conflict DropOut blocks of the online experiment, the graphs shows the trial-by-trial proportions of pain-reducing and lucrative choices for the first 10 trials of each block for participants with considerate (green) and lucrative (blue) preferences (preference classification is based on all conflict trials also including No-DropOut blocks), together with the predictions made by M1, M2Dec and M2Out. M0 always predicts 0.5 and is thus not shown, and participants with ambiguous preference are also not shown. In all cases, choices and predictions are averaged over all blocks (and samples for the models), and error bars then represent the s.e m. across participants. The models were fitted to the choices in the first 10 trials only.

The inlet, shows the leave-one-out information criterion (LOOIC(33)) of the models over the first 10 trials, with the error bars representing the estimation error. M0 performs poorly, but the three learning models perform similarly. **(B)** White background: participant choices and model predictions for the 10^{th} trial as in (a). Gray and yellow backgrounds: participant choices and model predictions for the 11^{th} trial, with dashed lines connecting the 10^{th} trial to 11^{th} trials in blocks in which money is removed (gray background), and the dotted lines those where shocks are removed (yellow background), separately for the considerate and lucrative preference groups. Note that the 11^{th} trial was not included in the fitting of the models. M2Out (marked by black arrowheads) makes the best predictions for the 11^{th} trial. **(C)** Likelihood of the choices during devaluation at the 11^{th} trial over all

participants (incl. those with ambiguous preference) and including both money and shock Dropout trials, under
all four models. For each of the 4000 posterior draws, we summed the log likelihood of the 11th trial for all
participants and blocks, and the histograms show the distribution of those likelihood over the 4000 posterior
draws. Results were multiplied by -2 to place them on the information scale also used for LOOIC, that quantifies
the information lost by each model, with lower values indicating a model with better predictions, and zero would
indicate perfect predictions without lost information. M2Out clearly outperformed all other models in its ability
to predict out-of-sample choices after devaluation of one of the outcomes. Note that we use LOOIC in (b), because
these first 10 trials were included in the fitting of the model, but log-likelihood in (c) because it is not included in
the model fit. **(D)** The distribution of likelihood ratio (*K*) in favour of M2Out across subjects relative to the
competing models based on the log-likelihood of the 11th trial.
K=exp(log_lik_M2Out)/exp(log_lik_OtherModel). *K*>3 represents subjects with at least modest evidence to have
been more likely to have used M2Out than the alternative, and *K*<⅓, more likely to have used the alternative
model. Note that participants with *K*>3 always outnumber those for *K*<⅓, showing that also at the single subject
level, M2Out is the model most favoured by the data during devaluation.

M2Out can accurately recover the value of wf, and this value predicts both preference in
our task and helping in a different task.

For our winning model M2Out, fitting the model on the first 10 trials of the DropOut
blocks leads to a wide range of *wf* values (Figure 7A) across participants that echoes the wide
distribution of choices (Figure 5). Indeed the *wf* parameter estimates correlate highly with the
proportion of pain-reducing choices (in both experiments, Kendall's Tau$_{(wf,choice)}$<-0.82,
p<0.001, BF$_{10}$>1000, Figure 7B). Parameter recovery further shows that if one simulates
participants with different *wf*, M2Out accurately recovered much of that variance
(r(*wf*$_{simulated}$,*wf*$_{estimated}$)=0.69, p<10^{-6}, BF$_{10}$>10^6 Supplementary Figure 6). For our fMRI
participants, we also performed a Helping task not involving learning that we had previously
developed(4) (Supplementary material §3). In each Helping trial, participants viewed a victim
receive a painful stimulation, and could decide to donate money. If no money was donated, the
victim received a second stimulation of equal intensity. If some money was donated, each €
donated reduced the second stimulation by one point on the 10 point pain scale. We found that
the *wf* value estimated in our learning task in the fMRI experiment could significantly predict
how much money participants on average donated to reduce the victims pain in this different
Helping task (Kendall τ(wf,choice)=-0.47, BF$_{10}$=76, p<0.001, Figure 7C), providing evidence
that *wf* captures a property of the participant that generalises to other situations in which
participants need to resolve a conflict between self-money and other-pain. A Bayesian multiple
linear regression including *wf* together with more traditional trait questionnaires (four subscales
of the IRI [26] as a traditional measure of empathy and MAS [27] as a traditional measure of the

attitude towards money), found strong evidence that including *wf* improves the predictive
power (BF_incl=11.46, Supplementary Table 3), while none of these questionnaires explained
variance in the Helping Task (all BF_incl<⅓, Supplementary Table 3). This establishes the utility
of our task as a predictor of costly helping. The distribution of the other parameters of M2Out
can be found in Supplementary Materials §8.

547
Figure 7. *Wf* Predictive power. (A) Distribution of *wf* parameter estimates for M2Out in the Online (purple, only
including the first 10 trials of the Conflict-DropOut blocks) or fMRI (lavender, including all trials) across
participants. The whisker plots represent the quartiles, dots outliers. **(B)** Relation between *wf* and proportion of
pain reducing choices in the online (left) and fMRI (right) experiment. For the online experiment only the 10 first
trials of DropOut blocks are considered in the model fit. The line and shading represent the regression line and its
standard error, *p* and BF values are for a two-tailed test on the Kendall's tau value. **(C)** Relation between *wf* in
fMRI experiment and donation in the helping experiment.

Neural correlates of prediction errors for shocks can be found in the pain observation
network and the vmPFC.
Participants showed robust brain activity across a wide network of brain regions when
the outcomes of their decisions were revealed (Figure 8, Supplementary Table 5)
To examine whether signals in the affective vicarious pain signature (AVPS) covary
with prediction errors for shocks (PE_S), and whether they do so in a way that correlates with
wf or not, we extracted the *wf*-normalized PE_S parameter estimate image (β_{PES}) from each
participant, and dot-multiplied with the AVPS neural signature. It is critical for the
interpretation of our fMRI analysis that we use *wf*-normalized PE predictors in our fMRI
analysis (as described in Methods: fMRI data analysis: PE *wf*-normalization) to avoid that *wf*
influences results twice: in calculating the parameter estimate image and in calculating the
regression of the parameter estimates and *wf*. The AVPS is known to have larger signals when
viewing painful compared to non-painful facial expressions (28). We coded shock outcomes in

terms of their *value* for a participant and not their intensity, i.e. a non-painful shock had value

+1 and a painful shock of -1. Hence, trials with painful shocks should typically have negative

prediction errors, and β_{PES} should load *negatively* on this neural signature. We found this to be

the case (Figure 8B, $t_{(24)}=-5.55$, $p=1E-5$, $BF_{10}=2090$). Next, we asked whether the strength of

the loading depended on *wf* using a Bayesian correlation, and found evidence of the absence

of such a relationship ($r=-0.058$, $p=0.78$, $BF_{10}=0.257$, Figure 8C). The facial pain observation

network thus does carry signals that negatively covary with prediction errors for shocks in our

paradigm, and these signals do not depend on personal preferences (*wf*). Voxels contributing

substantially to the loading can be found in Figure 8D.

Next, we performed a more explorative voxel-wise linear regression that predicts the

parameter estimate of the PEs modulator using a constant and *wf*. As expected for a region

involved in valuation, we found the vmPFC to have signals covarying positively with PEs (i.e.

higher signals when shocks were lower than expected). A more ventral cluster showed signals

associated with PEs in a way that depended on *wf* (Figure 8E, warm colors and Supplementary

Table 6), while a more dorsal vmPFC cluster showed an association with PEs after removing

variance explained by *wf* (Figure 8E, cold colors and Supplementary Table 7). In addition, we

find that the left somato-motor cortex, including BA4 and 3, also harboured signals covarying

positively with PEs, with a more dorsal location representing it with a significant dependence

on *wf*, and a more ventral region in a way that did not significantly depend on *wf*.

Because PEs and PE$_M$ predictors in our fMRI design are somewhat correlated due to

the outcome structure of our task (average $r=-0.26$, ranging from -0.49 to -0.03 across our

participants), we performed a parameter recovery analysis (See Supplementary Material §11)

that confirmed that if we simulate participants with signals that covary with PEs but not PE$_M$,

we recover significant parameter estimates for PEs but not PE$_M$, and vice versa. We also found

that the PE$_M$ parameter estimate images do not load on the AVPS, and found evidence of

absence of such loading ($t_{(24)}=0.392$, $p=0.7$, $BF_{10}=0.226$, Figure 8B), confirming our ability to

appropriately separate and localize PE$_M$ from PEs signals.

Figure 8. Key fMRI results. **(A)** Results of the contrast Outcomes>0 indicating voxels where BOLD signals during the outcome phase (independently of prediction errors) are increased, p<0.001, k=FWEc=749 voxels. **(B)** Loading of the normalized parameter estimate images (βPE) for PE_S and PE_M on the AVPS(28). Each dot is one participant, the horizontal line is the mean. *p* and BF values reflect a one-sample parametric (because normality was not violated) *t*-test against zero in JASP. **(C)** Loading of the parameter estimate for PE_S (β_{PES}) onto the AVPS as a function of *wf*. *p* and BF values represent the outcomes of a parametric correlation analysis using Pearson's *r*. **(D)** Voxels contributing substantially to the loading of β_{PES} onto AVPS. Because the overall loading is calculated by first multiplying each voxel's β_{PES} value with the value in this voxel of AVPS, and then summing these multiplication over all voxels, here we simply calculated the first step (the voxelwise multiplication of β_{PES} and AVPS), and averaged the result across our participants. To not overcrowd the image, we only show voxels with values above 50 or below -50. **(E)** Warm colors on the central two columns: regression between PE_S and 1-*wf* (p_{unc}<0.001, k=FWEc=162voxels). This identifies voxels with signals that increase more for shock outcomes

that are less intense than expected (i.e. positive PE_S) in participants that place more weights on shocks (high 1-*wf*

values). Cold color in the outer two columns: results of the contrast constant>0 in the same linear regression with

(1-wf) (p_{unc}<0.001, k=FWEc=132voxels). This identifies voxels with signals that increase with increasing PE_S

after the variance explained by (1-*wf*) is removed. All results in (E) were FWE cluster corrected at p<0.05,

following cluster cutting at p_{unc}<0.001, specified using the critical FWE cluster size FWEc. Note that the reverse

contrasts for (e) did not yield results that survive FWE cluster correction. The light gray line helps the viewer to

visually compare the location of the clusters across the two contrasts.

Discussion

Our aim was to investigate how participants learn action-outcome associations during

moral conflict - so called morally salient features (35) - that contrast gains for the self against

pain for others. We had three central questions. First, whether such learning would be model-

free or model-based, and whether they can be modeled using reinforcement learning models.

Second, whether individual preferences for maximizing self-gains vs. minimizing other-pain

would bias learning towards the preferred outcome. Third, if such bias exists, whether and how

the brain would process other-pain exclusively in a biased fashion, or whether it might have

co-existing representations that are biased and unbiased.

Our results indicate that participants' choices appear to be dominated by biased model-

based learning (16). The dominance of model-based learning was apparent (i) from the ability

to provide above chance explicit reports for both outcomes that also captured the difference

between Conflict and No-Conflict blocks and (ii) from the reaction to devaluation by not

altering choices if the preferred outcome was preserved, but by switching preference when the

preferred outcome was removed in the conflict blocks. That models were biased was borne out

from the fact that (i) individuals had more differentiated reports for their favoured outcome-

type (ii) that preferences after devaluation were asymmetric, with preference levels depending

on the weight of the remaining quantity.

Bayesian model-comparison of our models confirmed this conclusion (see

Supplementary Materials §12 for a detailed discussion of the predictions and rationale of each

model). During the first 10 trials of Conflict, all the learning models outperformed the random-

choice model (M0), but made virtually indistinguishable predictions. However, when one

outcome was devalued, M2Out outperformed the other learning models at the group level, and

in the majority of participants. Specifically, M2Out was the only model that captured the

prevalent behavior of unchanged preference, when the less favoured outcome was removed

and weak preference, when the most valued outcome was removed. This was because M2Out

tracked separate self-money and other-shock expectations for each symbol (and is hence
model-based), but did so in ways that depend on individual preference (and is thus what we
call biased). For a minority of participants, we found evidence that unbiased model-based
learning or model-free learning was a better description of choices under devaluation, and
further studies may want to investigate which variables drive such individual differences.

That our participant's learning seemed dominated by a model-based form of learning
contrasts with a recent report that participants choices appear to be dominated by model-free
learning when learning to prevent shocks to others(15). Two main experimental differences
could account for these contrasting findings. Firstly, in Lockwood et al., agents following
model-based learning do not outperform agents following a model-free strategy(15), while in
our DropOut condition, model-based learning allows swifter switching to the most favourable
option. This difference in incentives could have biased our participants towards the arguably
more effortful model-based strategy. Second, in Lockwood et al., participants only had to track
one outcome (shocks for self *or* others), while in our task, two outcomes (self-money and other-
shocks) were always involved. Dual, conflicting outcomes may prioritise more model-based
forms of learning. Overall, this difference emphasizes the need and opportunity to investigate
how situational demands shape the choice participants make between model-based and model-
free learning in morally relevant situations.

Considering the dominance of the biased model-based learning, we next asked how this
biased model-based learning would be implemented in the brain. Given the considerable
literature on learning to choose options that maximize gains for the self (22, 36), we focused
on how shocks to others are processed in the brain under conflict. In principle, there might be
two ways in which the bias in our model-based learning could be implemented in the brain: (1)
the bias occurs so early that a person maximizing self-gain fails to process shocks altogether
(but see Supplementary material §15 for preliminary eye-tracking data) or (2) all participants
initially process shocks in similar ways and modulate these signals with *wf* only in later stages
of processing. In case (1) we would fail to find BOLD signals associated with PEs in a way
that does not depend on *wf* anywhere in the brain, while in case(2) we could expect that earlier
stages of processing (e.g. the network associated with pain observation(24–26, 28)) has signals
associated with PEs that do not depend on *wf*, while later stages of processing associated with
valuation, particularly in the vmPFC(14, 22, 29–31) might have PEs signals that depend on *wf*
and inform choices. The psychological (37) and neuroimaging(38–42) literature cannot
adjudicate these alternatives, as they have shown biases of empathy at both early and late levels
of processing. However, our fMRI data speaks to option 2: the pain observation network, as

quantified using the affective vicarious pain signature(28), had signals that covaried with
prediction errors for shocks with evidence for the absence of a linear dependence on *wf*, while
within the medial prefrontal cortex, we found more dorsal regions with signals that covary with
prediction errors for shocks with magnitudes that did not significantly depend on *wf*, as well as
more ventral signals that did. That the ventral but not the dorsal cluster had valuation signals
that depend on *wf*, i.e. on whether a participant focuses on minimizing other-pain or not,
dovetails with a finding of Nicolle et al. (32) that if participants have to switch from trial to
trail between focusing on making a monetary decision according to their own values or those
of someone else, more ventral mPFC voxels had valuation signals corresponding to the values
of the current focus, while more dorsal clusters had valuation signals corresponding to those of
the alternative focus.

A limitation of our fMRI study is that, unlike prediction errors for shocks and money
that we show to be distinguishable, outcomes and prediction errors were so highly correlated
(shock: mean r= 0.84, ranging 0.75-0.87; money: mean r= 0.84, ranging 0.78-0.86) that we
cannot establish whether signals are truly representing prediction errors rather than outcomes.
While this problem is not unusual (e.g. (12)), a novel version of the paradigm better
dissociating prediction error and outcome will be necessary to explore whether the pain
observation network simply processes outcomes or already integrates outcomes with
expectations as predictive coding theories might predict(43–45). However, whether the pain
observation network processes outcomes or prediction errors, it does so in ways that do not
depend on *wf*, thereby providing evidence against the notion that the bias is so pervasive that
the brain does not at all represent shocks independently of bias. This hybrid representation of
some signals that are biased and others that are not, echoes the fact that in our Online
experiments, even participants with lucrative preference have above chance access to the
association of the symbols to shocks (Figure 4B), while the magnitude in probability difference
does correlate with *wf* (Figure 4D). A promising direction in which our study could be
expanded would involve the use of electro-physiological recordings. In particular, one could
then explore whether signals in the pain observation network covarying with prediction errors
earlier than the vmPFC signals that do depend on *wf*, and whether coherence across these
regions increases during shock observation, to test the idea that vmPFC valuation may be
informed by unbiased activity in the pain observation network.

In addition to these core fMRI findings, we made a couple of additional, less expected
findings in our explorative analysis. First, we failed to find signals robustly associated with
prediction errors for money when applying a whole brain correction for multiple comparisons

(but for a lower uncorrected threshold results see

https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254

Figure8_fMRI/SecondLevel_Regression_PEM_wf). That we had few participants with very

high *wf* (i.e. with highly lucrative preferences) and more participants with low *wf*, in our fMRI

study, could help explain why it was more difficult to find signals that represent the self-money

compared to the other-shocks. Additionally, money was always delivered (i.e. money was

delivered also in the low-money condition), and we had robust activations in regions associated

with monetary rewards when outcomes were revealed. The strength of that trial-independent

signal may have made it more difficult to detect the apparently smaller differences between

trials with higher and lower monetary outcomes. In the future, it might be helpful to either

compare conditions with no monetary reward at all against a condition with monetary reward,

and to recruit a larger number of lucrative participants. Second, we also found regions around

the left central sulcus encoding prediction errors for shocks in ways that depend on *wf* or not,

extending into both SI (area 3b and 1) and MI (area 4). While it is tempting to interpret these

activations as representing some form of embodiment of the somatic pain of others onto the

observer's own somatomotor cortex (4, 27), such conclusions should be made with care

considering the limitations of reverse inference, in particular given the explorative nature of

this whole brain analysis (46). Neuromodulating these foci, akin to what we have done

previously in our helping task (4), and exploring the impact on choices, would be important to

explore their causal contribution to learning and decision making.

Participants varied substantially in their preferences under conflict. In our Online study,

about one third prefered the symbol leading to high monetary gain, another third, the symbol

minimizing the pain to the other, while for the remaining third the number of trials we acquired

did not suffice to identify a significant preference. This pattern was true despite that to

maximize conflict, in the online experiment we asked each participant to adjust the amount of

money on offer for the high money outcome to be equivalent to the cost of the other receiving

the painful shock-intensity we used (Supplementary Material §2). Indeed, participants that had

a lower indifference point, later choose the lucrative option more often, and these individual

differences in choice are thus not a trivial consequence of individual differences in the amount

of money on offer. In our models, we find that an individual parameter, the weighting factor

(*wf*, conceptually similar to the salience factor alpha in the Rescorla-Wagner Learning Model

(19)) is an effective way to capture these individual differences in the relative weight placed

on the money vs shocks. Importantly, *wf* had external validity, in that it predicted how much

money the same participants gave to reduce shocks to the same confederate in a different task,

the Helping Task (4), which does not require learning. We also found that *wf* was better than

our trait measures of empathy and money attitude scales at predicting donation in this Helping

Task. Recent work has supported the view that state empathy is regulated by motives and

context (see (37, 47) for reviews). Our moral conflict task creates financial incentives known

to downregulate empathy (37). It is thus perhaps unsurprising that the IRI, which measures

self-reported trait empathy and which does not probe empathy under such conflictual situations,

fails to predict decisions under conflict, while *wf*, which is estimated for learning during moral

conflict, is better at predicting decisions during moral conflict in the Helping task. This speaks

to the need to develop situated state empathy measures that probe the propensity of participants

to deploy their empathy in specific situations, to complement measures of empathy as a context

free trait (37, 47). Our paradigm might be particularly well suited in phenotyping antisocial

populations and could provide insights into the atypical tuning of their morally relevant

learning and decision-making in the spirit of computational psychiatry (48). This is because

psychopathic offenders show atypical instrumental learning (49) and reduced activity in

empathy related networks while witnessing the pain of others (50) despite typical levels of trait

empathy in self-report questionnaires.

In our fMRI study, most participants (13/27) showed a significant preference for the

pain-reducing option, with only 3/27 showing a significant preference for the lucrative option.

Why this distribution was different from the Online experiment is unclear, but two factors may

play a role. First, in the fMRI experiment, that was conducted on site, participants were under

the impression that all shocks were delivered in real time to the other person, while in the online

experiment, they were led to believe that a subset of the shocks would later be delivered to

another person. This may have made the value of reducing shocks higher in the fMRI than in

the Online experiment. Second, the vicinity of the experimenter in the fMRI experiment may

have increased the motivation to act in a socially desirable way compared to the online

experiment. As there is no significant correlation between our variables of interest (*wf* and

percentage of considerate choices) with age, the difference in age between the experiment is

unlikely to have caused this difference.

Our study has several additional limitations. First, we limited our model comparison to

a number of hypotheses driven by RLT models. We did not test ratio or logarithmic ratio

models of valuational representation in this study. These valuational structures are known to

occur but are less often indicated in modeling gains and losses to the self (17). Future

experiments could be optimized to explore whether such alternative ways to combine these

values may be more appropriate under certain moral conflicts. Second, our participants or at

least some of them, may not use a RLT model at all. Instead, they may use rules such as 'choose
a symbol randomly, and only switch if you encounter *x* unfavorable outcomes in a row'. Future
studies may benefit from studying how well such heuristics may perform. Third, we used *wf*
to address quantitative individual differences. We recognize that *wf* only indicates preferences
during our tasks and that it is risky to interpret the *wf* as suggesting stable moral values, as is
the case for all behavioral studies absent strong evidence regarding stable moral commitments.
Future studies may wish to explore whether different individuals may be best captured using
qualitatively different models and in conjunction with validated evidence of long-term moral
commitments and values. This might be particularly relevant when including participants with
independently demonstrated morally considerate commitments on the one hand, or psychiatric
disorders affecting social functioning on the other. Fourth, our evidence for a biased model-
based approach hinges on drop-out trials in our Online experiment, and our request for
probability reports. We introduced these trials to rigorously reveal the nature of the learning
that participants deploy in conflict situations - and they indeed provided the data necessary to
adjudicate across our models. However, we must also consider the possibility that these very
trials also influenced participants to deploy the model-based learning that provides more
optimal decisions during drop-out and more accurate probability reports. Performing a conflict
task with only a single devaluation trial on a large number of participants may be a way to
exclude this possibility. However, if participants were to have adapted their strategy to
optimally fit the requirements of the task, M2Dec would have been even more adaptive.

799   To summarize, our data sheds light onto the processes at play when adults have to learn
that certain actions lead to favorable outcomes for us but harm others, while alternative actions
are less favorable for us but avoid harm to others. We show that in our task, the choices of a
majority of participants are best described by a biased model-based approach, in which some
brain signals (e.g. in the pain observation network) covary with prediction errors for the harm
to others in ways that do not depend on whether participants prefer to maximize gains for self
or minimize harm to others, while others (e.g. in the vmPFC) do depend on this individual
preference. We foresee that the neurocognitive model we introduce and this task will be
particularly useful, in the future, to understand how neurocomputational processes may differ
in antisocial populations that often fail to make choices that appropriately consider harm to
others in situations in which such harm results from pursuing personal goals.

Materials and Methods

Two independent experiments were performed: an Online study and an fMRI study. Behavior was tested in a larger sample of participants, but this had to be done online, using the Online platform Gorilla (https://gorilla.sc/), due to COVID-19 restrictions in place at the time. Brain activity was measured in a smaller number of participants in our fMRI scanner. Table 1 gives an overview of the number of participants and experimental conditions included in each study.

Participants

In total, 106 healthy volunteers with normal or corrected-to-normal vision, and no history of neurological, psychiatric, or other medical problems, or any contraindication to fMRI (for the fMRI experiment only) were recruited for our experiments (Table 1). Particularly, 27 participants (37y±17SD; 27f) took part in the fMRI, and 79 (25y±7SD, 39f) in the Online experiment, with the two samples being independent. Participants taking part in the Online experiment were significantly younger than the ones who took part in the fMRI study (Mann-Whitney test, BF_{10}=8.022, p<0.001). However, choices (in terms of percentage of considerate choices) and *wf* did not significantly depend on age (Kendall's tau(age, %Considerate choices)=0.113, BF_{10}=0.543, p=0.096; Kendall's Tau(age,wf)=-0.115, BF_{10}=0.561, p=0.091). Participants for the online task were recruited through Prolific (https://prolific.ac/), while those for the fMRI through advertisements of the experiment on social media.

As two of the 27 participants in the fMRI were left handed, and stimuli showed movements of the right hand of the actor, to reduce variability possibly induced by lateralization of the brain responses, these two participants only performed the tasks off-line (i.e. no fMRI data acquired). The studies were approved by the Ethics Committee of the University of Amsterdam, The Netherlands (2017-EXT-8201, 2018-EXT-8864, 2020-EXT-12450). Consent forms for participation and authorization for the publication of images have been obtained.

Online Experiment general procedure

In the Online experiment, participants' main task was the probabilistic reinforcement learning tasks, in which participants had to choose between two new symbols based on learning how much money each symbol could earn them, and whether the symbols lead to a painful shock to another participant (Figure 1). The task included a deceptive cover-story in which

participants were informed, at the very beginning of the study, that while they perform the task

online, a second participant is present in our lab. It was explained that the separation of the two

participants allowed the experiment to be performed under COVID19 restrictions. In reality,

no one was invited to our lab, and this cover story served to create a situation in which

participants took what they thought to be decisions with real implications for self and others,

while at the same time minimizing discomfort to others (no shocks actually delivered to the

other person). Participants then received an general overview of all the tasks they will

encounter during the experiment. The text also informed them that 15 trials would be randomly

selected from the main Learning task, and that the self-money of these trials would be added

to the online participants payment, while the electrical stimulation associated with these trials

would then be delivered on the hand of the other participant in the lab. Participants knew they

would be offered the choice to see (via Zoom), or not, the other person receiving the stimulation

at the end of the task. The experimental session started with a Stress Tolerance questionnaire

assessing participants' susceptibility to stress, which generated the advice to continue or not

with the experiment (Supplementary material §13). No participant received the advice to

withdraw, and the session continued with the Optimization Task, aimed at determining the

amount of money necessary to create a true conflict for a given person between self-gain and

other-shock (this value is referred to as *indifference point*). This was done by asking

participants to choose between different predetermined combinations of money and stimulation

intensity (Supplementary material §2). The choices in this task also count toward the final

bonus and number of stimulation to the other. The experiment then continued with the core

Reinforcement Learning Task (see below). The experiment ended with a debriefing about the

deception, asking how much they believed the cover story (Supplementary material §4) and

with a series of questions to investigate participants' motivations and their feelings toward the

experiment (Supplementary material §14). Participants were aware that they could withdraw

from the experiment at any time.

Reinforcement Learning task (Figure 1-D)

Conflict. Participants had to learn symbol-outcome associations under a moral conflict - i.e. in

a context in which choices that are better for the self (high monetary gain) are usually worse

for the other (painful shock), while those that are worse for the self (low monetary gain) are

usually better for the other (non-painful shock). Each block was composed of 10 trials (Figure

1A) in which participants had to learn the outcomes associated with each of the two presented

symbols. The same symbol pair was presented in all trials of a block, but changed across blocks.
In each block one of the symbols would most likely lead to higher monetary reward for the
participant, and a noxious electrical stimulation to the dorsum of the actor's hands (lucrative
symbol or choice), and the other symbol to lower monetary reward and a non-noxious
stimulation to the actor (pain-reducing symbol or choice). Symbol-outcome associations were
drawn independently for each symbol and outcome, following the schema illustrated in Figure
1B. This procedure helped decorrelate representations of shock and money. While the
monetary reward was presented in the outcome screen as a numerical amount of euros, the
outcome for the other was presented as a video recording of the author A.N. while receiving
the stimulation (Supplementary material §1). Participants could therefore infer whether the
stimulation was or not noxious only based on the facial reaction of the actor in the videos.
Participants were aware that the person in the video and the other participant in the lab were
not the same individual, and that the videos were pre-recorded to illustrate what responses to
the stimulations would look like.
890
No-Conflict. This condition was identical to the Conflict blocks with the exception that the
symbol-outcome associations did not likely introduced a moral conflict because the symbol
that was usually best for the self was also usually best for the other: one symbol was most likely
associated with high monetary reward for the participants and non-noxious stimulation to the
other participant, while the other symbol was usually associated with low monetary reward and
a noxious stimulations. There was therefore a clear incentive to choose for the symbol leading
to the higher reward, as it was also the symbol that would be most beneficial for the other.
898
ConflictMoneyDropout. Each block included two sets of 10 trials. The first 10 trials were
identical to those presented in the Conflict condition. After the 10th trial, the participants were
informed that the money outcome was removed and the block continued with the last 10 trials,
in which participants would only see the videos of the actor receiving the stimulation, and no
money would be rewarded. Participants were informed that the probability of each symbol of
being associated with high or low stimulation would remain the same over the 20 trials.
905
ConflictShockDropout. This condition was identical to the ConflictMoneyDropout with the
exception that the removed outcome was the stimulation to the other. From the 11th to 20th
trials participants therefore only saw the amount of money associated with the chosen symbol
and no consequences were to happen to the other.

No-ConflictMoneyDropout and *No-ConflictShockDropout.* These two conditions were the
same as the ConflictMoneyDropout and ConflictShockDropout, except that the first 10 trials
were the same as those presented in the NoConflict conditions, in which symbol-outcome
associations did not likely introduce a moral conflict.

The position of each symbol on the screen within a pair was randomized across trials. Symbols
composing a pair were kept the same across individuals (i.e. the same two symbols formed the
same pair throughout the experiment), but pairs were randomly distributed across conditions
and participants. Symbol-outcome associations were determined based on the matrix in Figure
1B, and were kept constant across participants. Conditions were presented in a randomized
order within and across participants.

Explicit recall questions
At the end of each of the four Conflict and four NoCoflict blocks only, participants were asked,
for each symbol separately, to recall the probability of that symbol to be associated with higher
monetary reward and with the noxious stimulation using a scale from 0 to 100% (Figure 1C).
The order in which the two symbols were presented and the type of question (association with
high shock or with high reward) was randomized between participants. The slider starting
position was always at 50%.

fMRI Experiment
Participants in the fMRI experiment performed two experimental tasks: a Helping task
as published in Gallo et al. (4), and the Conflict condition of the reinforcement learning task
used in the Online experiment. The experiment consisted of four sessions of the Helping task
and one session of the Conflict task, positioned after the first two sessions of Helping. fMRI
results of the Helping task will be the object of a separate publication. In the current work we
only report the fMRI data associated with the Learning task, and the amount of money
participants were willing to give up to reduce the intensity of the confederate's stimulation in
the Helping task in order to test the predictive power of the weighting factor.
Briefly, the Helping task showed videos of another participant (author S.G.) receiving
noxious stimulations of different intensities (Supplementary material §3). Participants were
then endowed with 6 euros, which they could decide to keep for themselves or to give all or in

part up to proportionally reduce the intensity of the next stimulation. Videos either showed the
facial reaction of the confederate or the hand reaction (no face). Because the online experiment
mainly showed facial reactions, only donations associated with the trials showing the facial
reactions in the Helping task were included in the current manuscript (results would though not
change when including the full donation dataset (Kendall τ(wf,choice)=-0.54, BF_{10}=13.19,
p=0.004). Participants were led to believe that the videos they saw were a life-video-feed of
another participant receiving these shocks in a closeby room, although in reality they were pre-
recorded movies. All participants were presented the same videos, albeit in randomized order,
so that we can compare the average donation of the participants as a willingness to give up
money to reduce the pain of others.

As the participants of the fMRI experiment came in person to the lab, the cover story
was slightly different from that adopted in the online version. For the fMRI we used the same
cover story used and validated in Gallo et al. (4). Each participant was paired with what they
believed to be another participant like them, although in reality it was a confederate, author
S.G. They drew lots to decide who plays the role of the learner (or the donor in the Helping
task) and of the pain-taker. The lots were rigged so that the confederate would always be the
pain-taker. The participant was then taken to the scanning room while the confederate was
brought to an adjacent room, connected through a video camera. Participants were misled to
think that electrical stimulations were delivered to the confederate in real-time, and that what
the participants saw on the monitor was a live feed from the pain-taker's room. In reality, we
presented pre-recorded videos of the confederate's reactions.

In contrast to the Online experiment, in which the high-money amount was selected for
each participant using the Optimization task, in the fMRI Learning task, the high-money
amount offered was fixed at 1.5€ for all participants, and corresponded to the average amount
associated with the indifferent point in the Optimization task (1.53 ± 0.37SD; Supplementary
Material 2). The low-money amount was the same as in the Online experiment, 0.5€.

At the end of the fMRI tasks, participants were debriefed and asked to fill out the
interpersonal reactivity index (IRI) empathy questionnaire (51), and the money attitude scale
(MAS) (52). To assess whether participants believed that the other participant really was
receiving electrical shocks, at the end of the experiment, participants were asked 'Do you think
the experimental setup was realistic enough to believe it' on a scale from 1 (strongly disagree)
to 7 (strongly agree). All participants reported that they at least somewhat agreed with the
statement (i.e. 5 or higher).

The fMRI task was programmed in Presentation (www.neurobs.com), and presented

under Windows 10 on a 32inch BOLD screen from Cambridge Research Systems visible to

participants through a mirror (distance eye to mirror: ~10cm; from mirror to the screen:

~148cm). The timing of the task was adapted to the requirements of fMRI: each trial started

with a jittered fixation cross lasting 3-9 seconds. Then the two symbols appeared and

participants could make their choice without a time restriction. After the button press, there

was a fixation cross raging from 3-9 seconds and the video with a duration of 2 seconds

followed.

Analysis of Behavioral Data

Quantification of choices was performed in RStudio (Version 1.1.453) and Matlab

(https://nl.mathworks.com/). Statistical analyses were then performed using JASP (https://jasp-

stats.org, version 0.11.1), to provide both Bayes factors and p values. Bayes factors were

important because in many cases we are as interested in evidence of absence than in evidence

of the presence of an effect, and p-values cannot quantify evidence for the absence of an effect

(53). For instance, we expect that removing money outcomes in considerate participants should

not alter choices, while removing shocks should. We used traditional bounds of $BF_{10}>3$ to infer

the presence of an effect and $BF_{10}<\frac{1}{3}$ to infer the absence of an effect (34, 53). Two-tailed tests

are indicated by BF_{10},i.e. $p(\text{Data}|H_1)/p(\text{Data}|H_0)$ while one-tailed tests are indicated by BF_{+0}.

Where ANOVAs were used, we report BF_{incl} which reports the probability of the data given a

model including the factor divided by the average probability of the data given the models not

including that factor. Normality was tested using Shapiro-Wilk's. If this test rejected the null-

hypothesis of a normal (or bi-variate normal for correlations) distribution, we used non-

parametric tests such as the Wilcoxon signed rank or Kendall's Tau test, while if the null-

hypothesis was not rejected, we used parametric tests. We always used default priors for

Bayesian statistics as used in JASP.

Computational Modeling of Behavioral Data

Our experiment represents a variation of a classical two armed-bandit task and was

modeled using a reinforcement learning (RL) algorithm with a Rescorla-Wagner updating

rule (19). We compared 4 models explained in Figure 1E. Models were fitted in RStan

(version 2.18.2, http://mc-stan.org/rstan/) using a hierarchical Bayesian approach, i.e. by

estimating the actual posterior distribution through Bayes rule. Our models were adapted

from the R package *hBayesDM* (for "hierarchical Bayesian modeling of Decision-Making

tasks") described in detail in Ahn et al (2017) (54).

Model comparison was also performed in a fully Bayesian way. All the 3 models were

fit just on the conflict condition, and just the first 10 trials of each block (the ones involving

the conflictual choices). To assess the ability of a model to fit data of these initial 10 trials that

were included in the parameter-fitting, we use the leave-one-out informative criterion (LOOIC

(33)), which computes a pointwise log-likelihood of the posterior distribution to calculate the

model evidence, rather than using only point estimates as with the other methods, e.g. of Akaike

information criterion (AIC (55)) and the deviance information criterion (DIC (56)). The

LOOIC is on an information criterion scale: lower values indicate better out-of-sample

prediction accuracy. For the Conflict-Dropout blocks, on the other hand, we aimed to assess

the ability of a model fitted on the first 10 trials of a Conflict-Dropout block to predict behavior

on the devaluation trial (trial 11). The model was then fitted using the first 10 trials of all the

Conflict-Dropout blocks, and no leave-one-out procedure was necessary to assess the

predictive performance on the 11th trial because it was not included in the parameter fitting.

We thus directly used the log likelihoods of the 11th trial, which capture the probability of the

choices on the 11th trial given the model fitted on the preceding 10 trials. Specifically, at the

group level, we added the log likelihood of all the 11th trials of all the participants for each of

the 4000 draws from the posterior distribution, and then examined the distribution of these

values (Fig. 4C). To compare the ability of different models to predict this 11th trial at the

single subject level, we then calculated a likelihood ratio (similar to a Bayes factor) for each

participant: we summed the log-likelihood of the 8 11th trials available for each participant,

and used the mean across the 4000 draws as our point estimate. We then calculated the ratio of

the exponents of the log-likelihoods for the competing models.

As priors on the hyperparameters   and *wf*, we use the recommended Stan method of

using a hidden variable distributed along N(0,1), that is then transformed using the cumulative

normal distribution to map onto a space from 0 to 1. For   , we use the same method but then

multiply the result by 5 to have the function map onto the interval [0,5].

For the Dropout blocks of the Online experiment, in which one of the two outcomes

was removed after the 10th trials, the three models M1, M2Out, M2Dec were modified from

the 11th trial to account for the fact that participants were told which quantity would be

removed. For the M2 models, this was implemented by setting EV=0 for the removed quantity

before decision-making on the 11th trial. In addition, *wf* is modified only to value the remaining

EV (i.e. *wf*=1 if shocks are removed and *wf*=0 if money is removed). For M1, we cannot reset

expectations for a specific quantity (shock or money) and EV has to remain unchanged.

However, the *wf* is adapted (i.e. *wf*=1 if shocks are removed and *wf*=0 if money is removed),

which maximizes what can be learned from the following trials. In all cases, the outcomes for

the missing quantity are always set to zero after Dropout.

MRI Data acquisition

MRI images were acquired with a 3-Tesla Philips Ingenia CX system using a 32-

channel head coil. One T1-weighted structural image (matrix = 240x222; 170 slices; voxel

size = 1x1x1mm) was collected per participant together with an average of 775.83 EPI

volumes ± 23.11 SD (matrix M x P: 80 x 78; 32 transversal slices acquired in ascending

order; TR = 1.7 seconds; TE = 27.6ms; flip angle: 72.90°; voxel size = 3x3x3mm, including a

.349mm slice gap).

fMRI Data preprocessing

MRI data were processed in SPM12 (57). EPI images were slice-time corrected to the

middle slice and realigned to the mean EPI. High quality T1 images were coregistered to the

mean EPI image and segmented. The normalization parameters computed during the

segmentation were used to normalize the gray matter segment (1mmx1mmx1mm) and the EPIs

(2mmx2mmx2mm) to the MNI templates. Finally, EPIs images were smoothed with a 6mm

kernel.

fMRI Data analysis

For the Learning Task, our aim was to identify brain activity that scales with PEs for

shock when the outcome is revealed, and to compare this activity across participants with

different weighting factors. Analyses and the experimental design therefore focused on the

outcome phase. In line with other studies (e.g., (12)), activity during the decision phase will

not be analysed here for two reasons. First, to isolate activity when outcomes are revealed, we

randomized the interval between outcomes and the response screen. As a result, decisions can

have occurred at any time between the last outcome and the next button press, making it

difficult to capture the activity linked to that decision. Second, the button press required for the

response triggers significant brain activity in frontal regions that could have been hard to

dissociate from the valuation processes we are interested in.

To analyse activity during the outcome phase, we ideally would have included

predictors for the monetary outcome (Out_M) and the shock (Out_S) as well as their prediction

errors (PE_M and PE_S). Unfortunately, PE_S and Out_S are too highly correlated (0.741, ranging

0.750 to 0.866 across our 27 participants), and the same is true for PE_M and Out_M (0.749,

ranging 0.781 to 0.862 across our 27 participants), to allow us to include them into a single

design matrix. Indeed, we performed a parameter recovery attempt that simulates voxels with

various mixtures of Out_S, PE_S, Out_M and PE_M across 24 participants with the addition of

noise, and tried to recover them using design matrices with all 4 parametric modulators, and

failed to have appropriate recovery (data not shown). This is not surprising, because for time-

efficiency, here we aimed to focus on the initial learning of the association, including only 10

trials per block, and during the initial learning outcomes with high Out_S will also cause high

PE_S and trials with high Out_M will also cause high PE_M. Our parameter recovery simulations

however confirmed that we can include PE_S and PE_M in a single design matrix, as these are

only weakly correlated (average r=-0.26, ranging from -0.49 to -0.03 across the participants),

and correctly identify voxels simulated to contain PE_S but not PE_M, PE_M and not PE_S or both

(Supplementary Materials §11). The parameter recovery also indicated that we can dissociate

voxels in which participants with lower *wf* show stronger PE_S related signals. This then allows

us to focus on the processing of shocks in the context of learning by identifying voxels with

signals that covary with PE_S in ways that do or do not linearly depend on *wf*, while being aware

that signals that covary with PE_S might in fact have covaried with Out_S.

Our fMRI design matrix thus included the following regressors. The decision regressor

started with the appearance of the two symbols and ended with the button press of the

participant. The outcome regressor was aligned with the presentation of the video and had a

fixed duration of 2 seconds, corresponding to the duration of the stimulus. The decision

regressor had 5 parametric modulators (EV_{S1}, EV_{S2}, EV_{M1}, EV_{M2}, choice) and the outcome

regressor 2 parametric modulators (PE_S and PE_M). The modulators were derived from the

winning M2Out, with EV_{S1}, EV_{S2}, EV_{M1}, EV_{M2} representing the expected values in terms of

shock and money for symbol 1 and symbol 2, and choice had value 1 or 2 based on the chosen

symbol. Symbols '1' or '2' correspond to the lucrative and pain-reducing choice respectively.

PE wf-normalization: Because we are interested in whether PE_S or PE_M representations

depend on *wf* or not, we divided PE_S with (1-*wf*) and PE_M with *wf* before entering them into

the design matrix. As a result, the first PE_S value in the parametric modulator would always be

PE_S=-1 if it was a high-shock outcome or PE_S=+1 for a low-shock outcome, independently of

the participant's *wf* value. If signals covary with PE_S in a way that does not linearly depend on

wf, the parameter estimate across participants (β_{PES}) would violate H_0:β_{PES}=0, but not

H_0:r(β_{PES}=0,wf)=0. If signals covary with PE_S in a way that does linearly depend on *wf*, it

would violate both of these H_0. Note that for outcomes, the coding was +1 for good outcomes

(i.e. high money or low shock) and -1 for bad outcomes (i.e. low money or high shock). EV

and PE follow that polarity.

Results were then analysed in two ways. First, to improve reverse inference, we use a

multivariate signature developed to distinguish the sight of facial expressions of pain vs. neutral

facial expressions that captures the overall activation of the facial pain observation network as

a single scalar value: the affective vicarious pain signature (AVPS(28)). To explore if signals

in this network covary with PE_S, we then simply dot-multiplied the PES parameter estimate

volume for each participant separately with the AVPS, after having brought the AVPS into our

fMRI analysis space using ImageCalc. The result of the dot-multiplication indicates how much

the covariance with PE_S loads on the AVPS. We then brought these values into JASP, and

compared them against zero and correlated them with *wf*. Because the values were normally

distributed, we used parametric analyses. We did the same for PE_M to confirm that our analysis

can separate PE_M and PE_S signals. Second, we performed a similar analysis at the voxel level,

by bringing the parameter estimate images for PE_S into a second level linear regression using

a constant and wf as the two predictors. A t-test on the constant then reveals regions in which

signals covary with PE_S after removing variance explained by *wf*. A t-test on the *wf* parameter

estimate then reveals regions where the signals covary with PE_S in ways that depend on *wf*. To

test if signals covary with 1-*wf*, we simply used a negative contrast in the t-test. Results were

familywise corrected at the cluster level using the established two-step procedure in SPM: (i)

for cluster-cutting we visualized results at $p_{unc} < .001$ k=10, and identified the FWEc minimum

cluster size for family wise error correction from the results table, (ii) we reloaded the results

at $p_{unc} < .001$ k=FWEc, so that all displayed results survive FWE at cluster-level.

Data Availability

All data and analysis code are available at

https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254

Acknowledgements

The work was funded by European Union's Horizon 2020 research and innovation programme

under ERC-StG 'HelpUS' 758703 to VG, Dutch Research Council (NWO) VIDI grant (452-

14-015) to V.G and VICI grant (453-15-009) to C.K. M.S. thanks the John Templeton

Foundation (Grant 21338) for support in developing the ideas regarding self/other valuational

representation.

We thank C. Gavanozi, I. Gembutaite and B. Hoekzema for helping with data acquisition in
the Replication and Outcome-Dropout study. We thank A. Veggerby Lind for helping
recording the stimuli used in the Replication and Outcome-Dropout studies. We thank A.
Gentile for his input and help with developing the learning models. VG, CK, AN, LD, KI, SG,
LF and RP thank P. Lockwood as her work inspired the development of the learning tasks.

Competing interests

The authors declare no competing interests.

1145

1146 **Bibliography**

1147

1.   J. M. Cowell, *et al.*, The development of generosity and moral cognition across five
cultures. *Dev. Sci.* **20** (2017).
2.   G. Hein, G. Silani, K. Preuschoff, C. D. Batson, T. Singer, Neural Responses to Ingroup
and Outgroup Members' Suffering Predict Individual Differences in Costly Helping.
Neuron **68**, 149–160 (2010).
3.   M. J. Crockett, J. Z. Siegel, Z. Kurth-Nelson, P. Dayan, R. J. Dolan, Moral
transgressions corrupt neural representations of value. *Nat. Neurosci.* **20**, 879–885
(2017).
4.   S. Gallo, *et al.*, The causal role of the somatosensory cortex in prosocial behaviour.
eLife **7** (2018).
5.   J. D. Greene, "The cognitive neuroscience of moral judgment and decision making" in
The Cognitive Neurosciences, 5th Ed, (MIT Press, 2014), pp. 1013–1023.
6.   O. FeldmanHall, T. Dalgleish, D. Evans, D. Mobbs, Empathic concern drives costly
altruism. *NeuroImage* **105**, 347–356 (2015).
7.   C. A. Hutcherson, B. Bushong, A. Rangel, A Neurocomputational Model of Altruistic
Choice and Its Implications. *Neuron* **87**, 451–462 (2015).
8.   J. K. Rilling, A. G. Sanfey, The neuroscience of social decision-making. *Annu. Rev.*
Psychol. **62**, 23–48 (2011).
9.   G. Hein, Y. Morishima, S. Leiberg, S. Sul, E. Fehr, The brain's functional network
architecture reveals human motives. *Science* **351**, 1074–1078 (2016).
10. R. S. Sutton, A. G. Barto, *Reinforcement Learning: An Introduction* (The MIT Press,
1998).

11. W. Schultz, Updating dopamine reward signals. *Curr. Opin. Neurobiol.* **23**, 229–238
(2013).

12. P. L. Lockwood, M. A. J. Apps, V. Valton, E. Viding, J. P. Roiser, Neurocomputational
mechanisms of prosocial learning and links to empathy. *Proc. Natl. Acad. Sci. U. S. A.*
113, 9763–8 (2016).

13. M. A. J. Apps, M. F. S. Rushworth, S. W. C. Chang, The Anterior Cingulate Gyrus and
Social Cognition: Tracking the Motivation of Others. *Neuron* **90**, 692–707 (2016).

14. L. L. Lengersdorff, I. C. Wagner, P. L. Lockwood, C. Lamm, When Implicit
Prosociality Trumps Selfishness: The Neural Valuation System Underpins More
Optimal Choices When Learning to Avoid Harm to Others Than to Oneself. *J. Neurosci.*
Off. J. Soc. Neurosci. **40**, 7286–7299 (2020).

15. P. L. Lockwood, M. C. Klein-Flügge, A. Abdurahman, M. J. Crockett, Model-free
decision making is prioritized when learning to avoid harming others. *Proc. Natl. Acad.*
Sci. U. S. A. **117**, 27719–27730 (2020).

16. R. J. Dolan, P. Dayan, Goals and habits in the brain. *Neuron* **80**, 312–325 (2013).

17. D. Talmi, A. Pine, How Costs Influence Decision Values for Mixed Outcomes. *Front.*
Neurosci. **0** (2012).

18. C. Adams, A. Dickinson, Instrumental Responding Following Reinforcer Devaluation.
Q. J. Exp. Psychol. Sect. B-Comp. Physiol. Psychol. **33**, 109–121 (1981).

19. R. A. Rescorla, A. R. Wagner, A theory of Pavlovian conditioning: Variations in the
effectiveness of reinforcement and nonreinforcement. *Class. Cond. II Curr. Res. Theory*
21, 64–99 (1972).

20. R. R. Miller, R. C. Barnet, N. J. Grahame, Assessment of the Rescorla-Wagner model.
117, 363–386 (1995).

21. J. Hernandez-Lallement, *et al.*, Harm to Others Acts as a Negative Reinforcer in Rats.
Curr. Biol. CB **30**, 949-961.e7 (2020).

22. O. Bartra, J. T. McGuire, J. W. Kable, The valuation system: a coordinate-based meta-
analysis of BOLD fMRI experiments examining neural correlates of subjective value.
NeuroImage **76**, 412–427 (2013).

23. M. Watabe-Uchida, N. Eshel, N. Uchida, Neural Circuitry of Reward Prediction Error.
Annu. Rev. Neurosci. **40**, 373–394 (2017).

24. C. Lamm, J. Decety, T. Singer, Meta-analytic evidence for common and distinct neural
networks associated with directly experienced pain and empathy for pain. *NeuroImage*
54, 2492–2502 (2011).

25. I. Timmers, *et al.*, Is Empathy for Pain Unique in Its Neural Correlates? A Meta-
Analysis of Neuroimaging Studies of Empathy. *Front. Behav. Neurosci.* **12** (2018).

26. J. Jauniaux, A. Khatibi, P. Rainville, P. L. Jackson, A meta-analysis of neuroimaging
studies on pain empathy: investigating the role of visual information and observers'
perspective. *Soc. Cogn. Affect. Neurosci.* **14**, 789–813 (2019).

27. C. Keysers, J. H. Kaas, V. Gazzola, Somatosensation in social perception. *Nat. Rev.*
Neurosci. **11**, 417–28 (2010).

28. F. Zhou, *et al.*, Empathic pain evoked by sensory and emotional-communicative cues
share common and process-specific neural representations. *eLife* **9**, e56929 (2020).

29. M. F. S. Rushworth, T. E. J. Behrens, Choice, Uncertainty and Value in Prefrontal and
Cingulate Cortex. *Nat. Neurosci.* **11**, 389-�397 (2008).

30. C. C. Ruff, E. Fehr, The neurobiology of rewards and values in social decision making.
Nat. Rev. Neurosci. **15**, 549–562 (2014).

31. C. J. Burke, P. N. Tobler, M. Baddeley, W. Schultz, Neural mechanisms of
observational learning. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 14431–14436 (2010).

32. A. Nicolle, *et al.*, An agent independent axis for executed and modeled choice in medial
prefrontal cortex. *Neuron* **75**, 1114–1121 (2012).

33. A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-
out cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).

34. H. Jeffreys, *Theory of Probability*, 2nd Ed. (Oxford University Press, 1939).

35. G. D. Reeder, A. E. Monroe, J. B. Pryor, Impressions of Milgram's obedient teachers:
Situational cues inform inferences about motives and traits. **95**, 1–17 (2008).

36. J. Garrison, B. Erdeniz, J. Done, Prediction error in reinforcement learning: a meta-
analysis of neuroimaging studies. *Neurosci. Biobehav. Rev.* **37**, 1297–1310 (2013).

37. J. Zaki, Empathy: a motivated account. *Psychol. Bull.* **140**, 1608–47 (2014).

38. G. Hein, T. Singer, I feel how you feel but not always: the empathic brain and its
modulation. *Curr. Opin. Neurobiol.* **18**, 153–158 (2008).

39. C. Lamm, H. C. Nusbaum, A. N. Meltzoff, J. Decety, What are you feeling? Using
functional magnetic resonance imaging to assess the modulation of sensory and
affective responses during empathy for pain. *PloS One* **2**, e1292 (2007).

40. K. C. Borja Jimenez, *et al.*, Changes in brain activity following the voluntary control of
empathy. *NeuroImage* **216** (2020).

41. F. Cui, A.-R. Abdelgabar, C. Keysers, V. Gazzola, Responsibility modulates pain-
matrix activation elicited by the expressions of others in pain. *NeuroImage* **114**, 371–
378 (2015).

42. E. A. Caspar, K. Ioumpa, C. Keysers, V. Gazzola, Obeying orders reduces vicarious
brain activation towards victims' pain. *NeuroImage* **222**, 117251 (2020).

43. R. M. Thomas, T. De Sanctis, V. Gazzola, C. Keysers, Where and how our brain
represents the temporal structure of observed action. *NeuroImage* **183**, 677–697 (2018).

44. J. M. Kilner, K. J. Friston, C. D. Frith, Predictive coding: an account of the mirror
neuron system. *Cogn. Process.* **8**, 159–66 (2007).

45. C. Keysers, V. Gazzola, Hebbian learning and predictive mirror neurons for actions,
sensations and emotions. *Philos. Trans. R. Soc. B Biol. Sci.* **369**, 20130175–20130175
(2014).

46. R. A. Poldrack, Inferring mental states from neuroimaging data: From reverse inference
to large-scale decoding (2011).

47. C. Keysers, V. Gazzola, Dissociating the ability and propensity for empathy. *Trends*
Cogn. Sci. **18**, 163–6 (2014).

48. P. R. Montague, R. J. Dolan, K. J. Friston, P. Dayan, Computational psychiatry. *Trends*
Cogn. Sci. **16**, 72–80 (2012).

49. R. J. R. Blair, The amygdala and ventromedial prefrontal cortex in morality and
psychopathy. *Trends Cogn. Sci.* **11**, 387–392 (2007).

50. H. Meffert, V. Gazzola, J. A. den Boer, A. A. J. Bartels, C. Keysers, Reduced
spontaneous but relatively normal deliberate vicarious representations in psychopathy.
Brain J. Neurol. **136**, 2550–62 (2013).

51. M. H. Davis, Measuring individual differences in empathy: Evidence for a
multidimensional approach. *J. Pers. Soc. Psychol.* **44**, 113 (1983).

52. K. T. Yamauchi, D. J. Templer, The development of a money attitude scale. *J. Pers.*
Assess. **46**, 522–528 (1982).

53. C. Keysers, V. Gazzola, E.-J. Wagenmakers, Using Bayes factor hypothesis testing in
neuroscience to establish evidence of absence. *Nat. Neurosci.* **23**, 788–799 (2020).

54. W.-Y. Ahn, N. Haines, L. Zhang, Revealing neurocomputational mechanisms of
reinforcement learning and decision-making with the hBayesDM package. *Comput.*
Psychiatry, 24–57 (2017).

55. H. Akaike, "Factor analysis and AIC" in *Selected Papers of Hirotugu Akaike*, (Springer,
1987), pp. 371–386.

56. D. J. Spiegelhalter, N. G. Best, B. P. Carlin, A. V. D. Linde, Bayesian measures of

model complexity and fit. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **64**, 583–639 (2002).

57. J. Ashburner, *et al.*, SPM12 manual. *Wellcome Trust Cent. Neuroimaging Lond. UK*

2464 (2014).

1274

1275

1276

1277

1278

1279

Supplementary material associated to the manuscript

'Neuro-computational mechanisms of action-outcome learning under moral

conflict'

Authored by L. Fornari[1§], K. Ioumpa[1§], A. D. Nostro[*], N. J. Evans[*], L. De Angelis, R.

Paracampo, S. Gallo, M. Spezio, C. Keysers[#] & V. Gazzola[#,*]

[§,*,#] equal contribution

* corresponding author: v.gazzola@nin.knaw.nl

1287

1288 **Table of Content:**

1305

1306

1. Stimuli creation and validation for the Learning task

Videos

Videos were generated in house. The authors SG and AN played the role of the actress during the fMRI and Online experiment respectively. All videos were recorded to start with an initial neutral facial expression, followed by the facial expression in response to the electrical stimulation delivered to the right-hand dorsum. The upper part of the actress' body was clearly visible on a black background. The actresses were encouraged to produce realistic and clear facial expressions in response to the temporally unpredictable stimulations. The intensity of the stimulation was decided with the actresses, and the facial expressions were exaggerated to reflect a clear expression of pain. All original recordings were cut to last 2s and to have the frame showing the beginning of the facial expression in response to the stimulation exactly at 1s. All videos were validated by independent groups of subjects, who were asked to rate the intensity of the pain experienced by the confederate on a scale from 1 to 10, with '1' being 'just a simple touch sensation' and '10' being 'most intense imaginable pain'. As low pain intensity videos we selected the ones with ratings of 1 and 2 and as high pain intensity videos we selected the ones with ratings of 4, 5 and 6 Videos were edited using Adobe Premiere Pro CS6 (Adobe, San Jose, CA, USA).

The videos used for the Learning task in the fMRI experiment were selected from the Hand videos recorded for Gallo et al., 2018 (1) (https://elifesciences.org/articles/32740, see also Supplementary Material§3 below, and https://doi.org/10.7554/eLife.32740.006, https://doi.org/10.7554/eLife.32740.009). Thirteen videos of value 1 and 14 of value 2 were used as low pain stimuli, while 9 videos of value 4, 13 of value 5 and 3 of value 6 were chosen for as high intensity stimuli. A full description of how these videos were created can be found in (1). A similar procedure was then used to generate the video of the Online experiment (for two examples see https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254 Figure1_Task). The main difference was that in the stimuli for the Online experiment both actress's hands were shown (Figure 1A), instead of only the right hand. For the Online experiment, an initial pool of 300 videos showing a painful stimulation and 300 videos showing an innocuous stimulation were recorded. The final pool of stimuli was then selected based on an online validation (survey done through Gorilla, https://gorilla.sc/) performed by 191 participants (aged 18 – 35, 103 females). Participants were recruited through Prolific (https://prolific.ac/). From the initial pool, 140 high intensity and 130 low intensity stimuli

were selected. Since for each condition we needed 140 trials (120 trials with both outcomes

plus 40 trials with money dropout), 10 low videos were repeated twice in each condition. The

exact same videos used in the Conflict blocks were repeated in the No-Conflict blocks, this

allowed us to rule-out any effect due to the difference in the displayed stimuli between

conditions. The repetition did not have any negative impact on participants' performance, since

in the Online experiment participants were aware of the fact that these videos had been pre-

recorded.

The high intensity pain stimuli used in the fMRI Learning task had an average rating of

$4.76 \pm 0.66SD$ and the low intensity stimuli of $1.52 \pm 0.51SD$ (Bayesian Mann-Whitney t-tests

W=675, p<0.001,BF_{10}=2710.5). For the Online study, the high intensity pain stimuli had an

average rating of $5.74 \pm 0.47SD$ and the low intensity stimuli of $1.54 \pm 0.18SD$ (Bayesian

Mann-Whitney t-tests, W=18200, BF_{10}=7.7*10^{11}, p<0.001). There was no difference between

the average rating of the low intensity pain stimuli used in the fMRI and in the Online

experiment (Mann-Whitney t-test W=1750, BF_{10}=0.24, p=0.983). On the other hand, the

difference between the average rating of the high intensity pain stimuli used in the two

experiments was significant (W=502.50, BF_{10}=456.31, p<0.001). Having high intensity pain

stimuli perceived as of lower intensity in the fMRI experiment might a priori predict more

lucrative preferences in the fMRI compared to the Online experiment. In contrast, lucrative

preferences were rarer in the fMRI than in the Online experiment (Table 1), suggesting that

this difference in the high intensity stimuli did not impact behaviour substantially.

1360

Symbols

All the symbols used in the learning task were created using Adobe Illustrator. For the

creation of symbols we used simple geometrical shapes without obvious meaning. Each symbol

was then paired with a second one created using the same shape elements organized in a

different position. All the symbols covered the same area on the screen. Twenty-four pairs of

symbols were used in the test: 12 pairs were used in the Conflict, and 12 in the No-Conflict

condition. The absence of *a priori* preference for a symbols in each pair was assessed in an

independent group of subjects (Bayesian Wilcoxon Signed-Rank test, W=31,, BF_{10}=0.31,

p=0.547).

1370

2. Optimization task

In the Online experiment, before the learning tasks, we determined the amount of monetary reward that would have a subjective value equivalent to the painful shock received by the other participant. This task enabled us to personalize the amount of money participants were later offered as high reward in the learning tasks to create a meaningful conflict. Participants always had to choose between a pain-reducing option combining 0.5€ for them with a low shock to the confederate, and a lucrative option combining a higher amount of money for themselves with a high shock to the confederate. The amount of money offered in the lucrative option varied in steps of 0.25€ across the 5 types of choices (see the table below). Before making the choice, the participant was able to see the intensity of the electrical stimulation for the confederate by playing pre-recorded videos.

	Pain-reducing option	Lucrative option	Nr participants
1	0.5€ + Low Shock	1€ + High Shock	12 (6L, 3A, 3C)
2	0.5€ + Low Shock	1.25€ + High Shock	22 (12L, 8A, 2C)
3	0.5€ + Low Shock	1.5€ + High Shock	13 (3L, 7A, 3C)
4	0.5€ + Low Shock	1.75€ + High Shock	7 (2L, 2A, 3C)
5	0.5€ + Low Shock	2€ + High Shock	25 (1L, 6A, 18C)

Supplementary Table 1. Optimization Task. The table lists the five possible options offered during the optimization task, with the number of participants to which each option was assigned during the Learning Task of the Online experiment. The numbers in brackets indicate the number of participants showing a Lucrative (L), Considerate (C) or Ambiguous (A) preference during the task.

Each of the 5 types of choices were presented 6 times, for a total of 30 decisions. A sigmoid was then fitted to the choice data, and the indifference point was selected based on where the sigmoid crosses the 0.5 pain-reducing proportion. To select the high reward to offer during the learning task, we picked the lowest value for which each participant chose half of the times the low-electrical stimulation option and half the high electrical stimulation option. In case there was not a value for which the number of pain-reducing choices was equal to the number of lucrative choices, we picked the lowest value for which the number of lucrative choices was higher than the number of pain-reducing choices. If a participant had always chosen the pain-reducing option, we picked the highest value (2€) as the high reward for the learning experiment. Conversely, if a participant had always chosen the lucrative option, we picked the lowest value (1€) as the high reward for the learning experiment. Due to an error in programming the task, to 4 participants who had 1€ as real indifference, 1.25€ was offered as

1398 high reward during learning. Another subject whom should have been offered 1€ got offered
1399 2€ as high reward. This could have induced a more lucrative behaviour in this subject, however
1400 this participant was still in the Considerate group, meaning that this didn't have a strong impact
1401 on performance. In the Online experiment, the average indifference point was 1.53 ± 0.37SD,
1402 which matched the 1.5 € given as high reward to every participant in the fMRI experiment.

1403

3. Helping Task

1405 In the fMRI experiment, participants performed the Helping Task presented in Gallo et
1406 al. (1) as main experimental task. Only the behavioral results (i.e. average donation per
1407 participant associated with the Face stimuli) of the Helping Task are included in the current
1408 publication. Briefly, participants performed 60 trials in which they watched a first (pre-
1409 recorded) video of the same confederate as in the Learning Task receiving a painful stimulation.
1410 The intensity of the stimulation could vary between 1 and 6 on a 10 point pain scale, and was
1411 chosen on each trial by the computer program. In each trial participants also receive 6 credits,
1412 and could decide to donate some of these credits to reduce the intensity of the second
1413 stimulation to the confederate. Each credit donated back to the experimenter reduced the next
1414 stimulation by 1 point on the 10 point pain scale. Participants then watched a second video
1415 showing the confederate's response to the second stimulation. At the end of the task,
1416 participants were paid the sum of the amount of money that they had kept for themselves from
1417 all the trials divided by 10. We capture individual differences as the average number of credits
1418 given up per trials ("donation"). Two types of videos were presented. One showing the
1419 confederate receiving an electroshock on the hand and expressing the pain she felt by only
1420 reacting with facial expression. The other showed a belt hitting the dorsum of the confederate
1421 right hand, and the confederate expressing how much pain she felt by a reaction of the hand
1422 alone. The face was not visible in the latter stimuli. Hand and Face videos were presented in
1423 separate sessions.

1424 In total there were 2 sessions of 15 trials each presenting the hand video and 2
1425 presenting the face. The single session with 6 blocks of Learning task was presented at the end,
1426 after the four Helping Task sessions. This was because the fMRI experiment was centered on
1427 the Helping task, and the Learning task was meant as a pilot data, which should have been
1428 followed by a second fMRI data acquisition centered to the Learning, which was however
1429 impossible due to COVID19 restrictions.

1430

1431

1432 **Supplementary Figure 1. Helping task structure.** Top: two screen shots taken from one exemplar video
1433 showing the confederate receiving the slap on her right hand, and from a video showing the confederate receiving
1434 the shock on the hand and manifesting its painfulness through facial reactions. Bottom: trial structure.

1435

4. Participants' belief that someone will get shocks

At the end of the Online study, participants were asked to express the degree of
agreement to the statement: "Before choosing whether or not to see the other person receiving
the electrical stimulations, I believed that someone was really going to receive them" on a scale
from 1 (strongly disagree) to 7 (strongly agree). If they at least somewhat disagreed, they were
additionally asked whether they nevertheless acted as if they believed someone would get
shocks or not. An ANOVA with level of agreement as a nominal factor showed that average
choices were not influenced by how much a participant believed in shock delivery (Main effect
of belief in shocks, $F_{(6,72)}=0.739$, p=0.62, $BF_{incl}=0.109$). However, of the 28 Non-Believers (i.e.
those who responded 'somewhat disagree', 'disagree', or 'strongly disagree'), 24 reported they
nevertheless acted as if someone were to receive the shocks (Supplementary Table 2). This
might explain why people that doubted anyone would get shocks would nevertheless give up
money to reduce shocks. In contrast, the 4 participants reporting they acted as if no one were
to get shocks (red in Supplementary Figure 2) chose pain-reducing options on average
significantly less than the other 75 participants (Independent sample test $t_{(77)}=-2.97$, p=0.004,

BF$_{10}$=8.75). In the fMRI study, all 27 participants reported that they believed that someone was

getting a shock.

1453

	Total Nr of participants	Believers	Non-Believers	
			Acting as if believing	*Non-acting as if believing*
Online participants	79	51	24	4
fMRI participants	27	27	0	0

1454

1455 **Supplementary Table 2. Distribution of participants' beliefs.** The number in each cell is the participants count.

1456 Believers were considered those reporting values from 4 to 7 (neither agree to totally agree), non-believers those

1457 reporting 3 to 1 (somewhat disagree to totally disagree). Nr=number.

1458

Supplementary Figure 2: Effect of belief in the fact that the victim will receive shocks on choice allocation in the Online study. The figure shows the proportion of pain-reducing choices across all the conflict trials of the Online experiment (including No-Dropout and the 10 first trial of Dropout blocks) as a function of the answer provided. Whisker plots represent the quartiles, black dots the outliers, and red dots participants reporting that they did not act as if someone were getting shocks. Participants responding that they at least somewhat disagree are considered Non-Believers for the purpose of Supplementary Figure 5.

1459
1460

5. Participants' motivation

To better understand what drove participants' choices in the Online learning task, at the end of the Online task, participants were asked the question "What motivated your choices?". Five options were proposed to the participant and they could choose to select 1 or more. In the case they selected more than 1, they were asked to select them in order of importance, starting from the most important. The options presented to participants were the following:

A) Gaining as much money as possible

B) Giving a good impression

C) Being in line with my moral beliefs

D) Avoiding to see the other person receiving the shocks

E) Preventing harm to others

Supplementary Figure 3A below reports how often a particular motivation was chosen as the primary motivation in each group.

Supplementary Figure 3. Motivation as a function of preference. (a) for each of the preference groups, the color indicates the proportion of participants that have chosen a particular motivation as their primary motivation, and the white numbers represent the actual number of participants in that cell. **(b)** The value of the pain-avoidance index, calculated as (D+E)/9-A/5, as a function of group preference. The whiskers represent the quartile. This index has the value of +1 if participants select pain avoiding motivations as their first two motivations (i.e. preventing harm to others and avoiding to see the video of the other person receiving the shocks) and do not select 'gaining as much money as possible', and -1 if they select 'gaining as much money as possible' and not the two pain avoiding motivations.

Since there were 5 alternatives, to quantify motivation, values from 0 to 5 were assigned
to A-E according to its position in the list of motivations (0 if it had not been selected, 5 if it
had been indicated as first motivation, 4 as second, 3 as third, 2 as forth and 1 as fifth). To
assess whether the motivation overall was more lucrative or pain-reducing, we then derived a
secondary measure of '*Pain-Avoidance*' that would be negative if the motivation was mainly
'gaining as much money as possible' and positive if the motivation was 'preventing harm to
others' or 'avoiding to see the other person receiving the shocks'. Specifically, this was
calculated as (D+E)/9-A/5. We divided D+E by 9 and A by 5, because a person with maximum
pain reducing motivation could choose D and E as their first motivations and thus obtain
5+4=9, and one maximally lucrative would choose A as their first motivation. We decided on
this contrast of D+E and A, because we found D+E to correlate highly negatively with A (tau=-
0.38, $p<0.001$, $BF_{10}=29232.57$). Supplementary Table 3 below gives an overview of the
average pain avoidance score calculator for the Lucrative, Considerate and Ambiguous group
preferences separately (see also Supplementary Figure 3B).
1498

	Pain-Avoidance score ((D+E)/9-A/5)		
	lucrative	**ambiguous**	**considerate**
Participants	24	26	29
Mean	-0.519	0.120	0.506
Std. Deviation	0.447	0.509	0.440
Shapiro-Wilk	0.903	0.960	0.887
P-value of Shapiro-Wilk	0.025	0.384	0.005
Minimum	-1.000	-1.000	-0.800
Maximum	0.556	1.000	1.000

1499 **Supplementary Table 3: Descriptive statistics for the Pain-Avoidance score.** As can be seen, the score indeed
1500 varies all the way from -1 to 1 amongst our participants, but is not normally distributed, and is thus analysed non-
1501 parametrically.

1502

To test whether participants that chose the pain-reducing options during the task, within
or below chance (i.e. considerate, neutral and lucrative) differed in their pain avoidance score,
we performed a Bayesian ANOVA with group as factor. This analysis revealed a significant
main effect of group (Kruskal Wallis tests, $H_{(2)}=35.18$, $p<0.001$,) and post-hoc Bayesian Mann-

Whitney t-tests confirmed differences between all the 3 groups (Considerate-Ambiguous:
$W=196.50$, $p=0.002$, $BF_{10}=9.31$; Lucrative-Ambiguous: $W=111.50$, $p<0.001$, $BF_{10}=38.14$;
Lucrative-Considerate: $W=47.50$, $p<0.001$, $BF_{10}=397.01$).

Testing each value against zero using a Wilcoxon signed rank test showed that for the
considerate preference group values were significantly above zero ($W=380$, $p<0.001$,
$BF_{10}=714.32$), for the lucrative preference group they were below zero ($W=13.5$, $p<0.001$,
$BF_{10}=748.88$) but for the ambiguous preference group, they were close to zero ($W=227.5$,
$p=0.19$, $BF_{10}=0.48$). This suggests that participants in the Considerate and Lucrative group
could and decided to consciously report motivations that were in line with their choices, while
the Ambiguous group appeared to have less clearly polarized monetary or pain-avoiding
motivations.

1518

6. Choices in the NoConflict Dropout and across all 20 trials

1520

1521

Supplementary Fig. 4. Participant's choices in the DropOut blocks. For both Conflict and No-Conflict blocks,

the figure shows the average ± sem pain-reducing choices as a function of trial separately for the three preference

groups. Note that preference grouping is based on all Conflict trials, including the No-Dropout blocks, while the

figure only shows choices on the Dropout trials. Tables summarize the Wilcoxon signed-rank values and BF_{10} for

key comparisons. BF rather than p values are shown, to specify evidence for ($BF_{10} > 3$) or against ($BF_{10} < 1/3$) the

presence of a difference. Here we also show choices on trials 12-20 after Drop-out. These choices are less relevant

to the main paper, because they capture what is probably a different form of learning, when symbols are only

associated with 1 outcome rather than 2 outcomes that can conflict. NoSh=No-Shock, i.e. the outcome shock has

been removed, and only money remains. NoMon=No-Money, i.e. the outcome Money has been removed, and

only the shock remains.

7. Model comparison during Conflict Dropout excluding certain participants

Supplementary Figure 5 (Related to Fig 4). Model Comparison. (A) Mean LOOIC and the Likelihood of the choices during devaluation at the 11th trial (as inlet Figure 6A, and Figure 6C), but excluding participants with ambiguous preferences. (B) As in (A) but excluding participants that expressed doubts. (C) Mean LOOIC for all fMRI participants.

8. M2Out Parameter Distributions and Recovery

The distribution of the parameters that were estimated for τ, λS and τ, λM can be found in Supplementary Figure 6A,B. We also performed a parameter recovery assessment on the M2Out model to assess the robustness of inferences on the parameter values estimated from this model. Generating values for the parameters wf, λS and λM were all drawn from a uniform distribution between 0 and 1, and τ from an exponential distribution between 0.2 and 5 (i.e., the natural logarithm of τ was drawn from a uniform distribution between -1.61 and 1.61). Parameter sets were sampled using a Latin hypercube sampling design (2, 3) to ensure an adequate coverage of the joint parameter space. The simulated data were generated using the full process of the M2Out model (i.e., simulated trial 1 informs simulated trial 2, etc.), with the amount of data matching the fMRI experiment (i.e., 6 blocks, each with 10 trials). Parameter values were estimated using Differential Evolution Markov Chain Monte Carlo (4), with 12

chains run in parallel for 2000 iterations and the first 1000 iterations discarded as burn-in. The

priors on the *wf*, λS and λM parameters were all truncated normal distribution (between 0 and

1) with mean 0.5 and standard deviation 0.2, while the prior for the τ parameter was a truncated

normal distribution (between 0 and infinity) with mean 1 and standard deviation 3. The

estimates plotted in Figure 6c display the estimated posterior means from each simulated data

set.

In the main manuscript, we use a hierarchical Bayesian model to estimate these parameters

under the assumption that we sampled multiple individuals from the same underlying

population. In particular for λS and λM, it is likely that different participants may gravitate

onto similar λS and λM given the similarity in volatility experienced by them. For the

parameter recovery we perform here, this assumption is not true: we deliberately sampled the

entire space of possible λS and λM using a uniform distribution. Accordingly, for the parameter

recovery, each simulated behavior was analysed individually using a non-hierarchical model.

The priors for this individual implementation were not informed by the hyper-parameters of

the final study, and were simply aimed at informing us about the proportion of all possible

variance in these parameters that can be retrieved from analysing participants one at a time.

The inverse temperature (τ) parameter had a relatively broad distribution across our

participants in our two experiments (Supplementary Figure 6A), and parameter recovery shows

that when simulating participants with different τ, the τ estimates recovered by fitting M2Out

correlated quite highly with the simulated τ values (Kendall's τ (τ simulated, τ estimate)=0.53,

p<0.001, BF$_{10}$>1000). As reported in the main text, the same was true for *wf*. In contrast,

parameter recovery shows that although the learning rates can be significantly recovered by

M2Out, the correlation values are more modest than for *wf* and τ (Kendall's τ (λS$_{simulated}$,

λS$_{estimate}$)=0.25, Kendall's τ (λM$_{simulated}$, λM$_{estimate}$)=0.24, both p<0.001, BF$_{10}$>1000), and the

τ values obtained from our model-fitting should thus be interpreted more tentatively. In

particular, we find that τ is difficult to estimate for the outcome that participants consider less

in their choices: for simulated participants with *wf*<0.1 (i.e. that minimize shocks to others),

Kendall's τ between simulated and estimated values drops to 0.03 for λM but is 0.36 for λS,

while for simulated participants with *wf*>0.9 (i.e. that maximize gains to the self), it is 0.32 for

λM but drops to 0.06 for λS. Accordingly, we did not include these parameter estimates in the

main manuscript due to their reduced robustness.

It might also be noted, that the median learning rates estimated by the hierarchical Bayesian

model for the Online experiment (Supplementary Figure 6B) was close to 0.25, and was very

different from the medial of the parameters recovered by the non-hierarchical model with a

prior with mean 0.5. Adjusting the prior of the parameter recovery to match the hyper-

parameter obtained in the Online experiment fixes this anomaly.

1589

Supplementary Figure 6. (A, B) Distribution of parameter estimates for M2Out in the Online (purple, only

including the first 10 trials of the Conflict-DropOut blocks) or fMRI (lavender, including all trials) across

participants. The whisker plots represent the quartiles, dots outliers. **(C-F)** Parameter Recovery Performance of a

non-hierarchical implementation of M2Out. The figure shows the result of simulating the choices of 4000

participants that use M2Out for their decision, with a range of *wf*, $\boldsymbol{\tau}$, λS and λM values, then estimating the

parameters using M2Out. See §8 for details of the procedure. Note that this was done non-hierarchically to avoid

influence from one simulation on the other, using priors that were not informed by the findings of our actual study.

We performed a Shapiro-Wilk test for bivariate normality that considered for each of the four parameters both the

simulated and estimated values. For *wf*, p=0.048, and we thus report both Pearson's r and Kendall's $\boldsymbol{\tau}$. For the

other three parameters, the Shapiro-Wilk p was always below 0.001, and we thus only report Kendall's $\boldsymbol{\tau}$. The *p*

and BF values always refer to one-tailed tests with H1: $\boldsymbol{\tau}$ >0, as this test does not require bi-variate normality.

Note how for *wf* and $\boldsymbol{\tau}$, the estimated values capture the simulated values fairly well, although the slope of the

regression line is <1. For $\boldsymbol{\tau}$, however, the prior distribution (truncated normal distribution, between 0 and 1, with

mean 0.5 and standard deviation 0.2) appears to have a stronger influence on the estimated values than the

simulated data, which strongly compresses the estimated values towards the peak of the prior and reduces the

coupling between simulated and estimated values. **(G-H)** Adapting the priors for λS and λM to 0.2, closer to the

values observed in the Online study, shifts the recovered parameters to a range closer to what was observed in the

online study.

1608

9. Predicting helping based on *wf* but not IRI or MAS

1609

	Coefficient	Mean	SD	P(incl)	P(incl|data)	BF inclusion
	Intercept	2.643	0.211	1.000	1.000	1.000
	wf_10T_M2 Dec	-1.767	1.075	0.500	0.835	**5.058**
I R I	Fantasizing	-0.004	0.032	0.500	0.291	0.410
	Perspective taking	-0.003	0.029	0.500	0.284	0.397
	Empathic Concern	0.011	0.032	0.500	0.318	0.466
	Personal Distress	-9.326e -4	0.026	0.500	0.282	0.393
M A S	Power-Prestige	-2.308e -4	0.027	0.500	0.289	0.406
	Retention-time	-0.010	0.032	0.500	0.310	0.450
	Distrust	0.021	0.039	0.500	0.386	0.630
	Anxiety	-0.018	0.041	0.500	0.350	0.539

1610

1611 **Supplementary Table 4. Bayesian linear regression posterior summary of coefficients.** The table summarizes

1612 the Bayesian linear regression model comparison for models explaining the donation in the Helping task using

1613 the *wf* of the Learning task, the subscales of the IRI (5) (FS, PT, EC and PD) and MAS (6) (Power-Prestige,

1614 Retention-time, Distrust and Anxiety). The first column indicates the variable under consideration, followed by

1615 the mean and sd of the regression parameter estimate, followed by the prior probability of including each variable

10. Results of the fMRI analysis for Outcome and PEₛ

1622

Cluster size	Voxels in cyto	% Cluster	Hem	Cyto or Anatomical description	% Area	Peak t-value	MNI coordinates		
							x	y	z
Main effect of Outcome (5% FWE correction . t=3.47 . p<.001 . k=749)									
29347	852	2.9	L	Lobule VIIa crusI	28				
	679.1	2.3	L	Lobule VI	36.3				
	522.3	1.8	R	Area 45 IFG (p. Triangularis)	50.6	11.84	50	20	8
	513.3	1.7	R	Lobule VI	28.5				
	495.4	1.7	R	Area hOc4la	55.9				
	487.1	1.7	L	Area hOc4la	57				
	467.3	1.6	R	Area PFm (IPL)	66.3				
	384.4	1.3	L	Area hOc4lp	44.9				
	371.1	1.3	R	Thal: Prefrontal	66.3				
	367	1.3	L	Thal: Prefrontal	58.1				
	336.5	1.1	R	Area hIP3 (IPS)	73.8				
	324.6	1.1	L	Lobule VIIa crusII	19.9				
	320.5	1.1	R	Area FG4	65.5				
	319.6	1.1	R	Lobule VIIa crusI	9.8				
	315.9	1.1	L	Area hOc4v [V4(v)]	43.5				
	311.6	1.1	R	Area 44	51.9				
	290	1	L	Area FG3	35.1	13.59	-42	-52	-20
	289.5	1	R	Area hOc4v [V4(v)]	46.6				
	278.5	0.9	R	Area hOc4lp	49.8				
	260.3	0.9	L	Area FG2 Fusiform Gyrus	51.1	11.83	-42	-68	-18
	258	0.9	R	Area hIP1 (IPS)	89.1				
	243.1	0.8	R	Area hOc1 [V1]	11.8				
	232.1	0.8	R	Area PGa (IPL)	31.3				
	224.8	0.8	R	Area hOc3v [V3v]	26.4				
	221.6	0.8	R	Area FG3 Fusiform Gyrus	33.8	11.24	44	-54	-20
	205	0.7	R	Area FG2	63				
	198.1	0.7	L	Area hOc3v [V3v]	21.4				
	188.3	0.6	R	Area FG1	75.7				
	177.9	0.6	L	Area FG4	30.1				
	166.6	0.6	R	Area hIP2 (IPS)	79.1				
	165	0.6	L	Area FG1	64.7				
	156.9	0.5	L	Lobule IX	25.2				
	154.3	0.5	R	Thal: Temporal	28.2				
	144.9	0.5	R	Area PF (IPL)	21.5				
	134.5	0.5	L	Area hOc1 [V1]	6.6				
	127.8	0.4	R	Thal: Parietal	38.3				
	119.6	0.4	L	Thal: Parietal	37.6				
	103.8	0.4	L	Lobule VI (Verm)	49.6				
	99.6	0.3	L	Thal: Temporal	18.7	14.05	-24	-26	-6
	96.5	0.3	R	Lobule VI (Verm)	41.6				
	92	0.3	L	Lobule VIIIa (Verm)	62.2				
	90.9	0.3	L	Lobule IX (Verm)	101.4				

	90.1	0.3	L	Lobule VIIb	13.3			
	87.8	0.3	R	Lobule VIIIa (Verm)	41.9			
	83	0.3	R	Lobule IX	11.8			
	82.9	0.3	R	Lobule IX (Verm)	79			
	82.3	0.3	R	Area hOc2 [V2]	8.1			
	80.5	0.3	L	Area hOc5 [V5/MT]	100.2			
	78.8	0.3	R	Lobule VIIa crusII	5.6			
	65.8	0.2	R	Area hOc3d [V3d]	12			
	62.9	0.2	R	Amygdala (LB)	29.4			
	61.6	0.2	L	Area hOc3d [V3d]	6.2			
	60.3	0.2	R	Lobule VIIb	9.2			
	58.3	0.2	R	Area hOc5 [V5/MT]	100			
	51	0.2	R	Area Fp1	3			
	50.4	0.2	R	Area 7A (SPL)	6.5			
	49.1	0.2	L	Thal: Visual	54.8			
	42.9	0.1	R	Thal: Visual	103			
	38.4	0.1	R	Area PFcm (IPL)	11.8			
	37.9	0.1	L	Area hOc2 [V2]	4	13.03	-18	-102	0
	36.3	0.1	L	Lobule VIIIa	4.8			
	36	0.1	L	Lobule VIIb (Verm)	117.6			
	35.4	0.1	R	Lobule V	4.4			
	34	0.1	R	Area PGp (IPL)	3.4			
	32.1	0.1	R	Subiculum	8.5			
	30.9	0.1	R	Area PFt (IPL)	7.4			
	30.6	0.1	R	Lobule VIIb (Verm)	93.5			
	30.4	0.1	R	Lobule VIIa crusII (Verm)	53.5			
	28.6	0.1	R	Area 7PC (SPL)	6.3			
	28.1	0.1	L	Area PGa (IPL)	4.4			
	28	0.1	L	Area PFm (IPL)	4.8			
	23.8	0.1	R	Area hOc4d [V3A]	5.6			
	23.1	0.1	L	Lobule VIIa crusII (Verm)	51.8			
	21.6	0.1	R	Lobule VIIIa	3			
	21.4	0.1	R	Area Fo2	1.9			
	18.9	0.1	R	Amygdala (SF)	39.6			
	14.8	0.1	R	Amygdala (CM)	53.9			
			L	Hippocampus		14.20	-20	-30	-6
			L	Fusiform Gyrus		13.59	-42	-52	-20
			R	Insula Lobe		12.72	42	20	-2
			R	IFG (p. Opercularis)		12 .26	46	10	30
			L	Cerebelum (Crus 2)		11.78	-8	-74	-32
2032			R	Superior Medial Gyrus		10.81	4	32	52
			L	Superior Medial Gyrus		8 .68	2	36	40
			R	Posterior-Medial Frontal		7.95	4	22	60
			L	Superior Medial Gyrus		6.43	-6	26	40
			L	ACC		4.21	-10	26	28
1853	368.3	19.9	L	Area hIP3 (IPS)	80.4	11.15	-30	-52	46
	263	14.2	L	Area hIP1 (IPS) Inferior Parietal Lobule	72.2	5.05	-50	-40	46
	104.4	5.6	L	Area hIP2 (IPS)	46.4			
	62.4	3.4	L	Area 7A (SPL) Superior Parietal Lobule	5	4.53	-30	-66	56
	29.1	1.6	L	Area 7P (SPL)	8.4			
	21.4	1.2	L	Area 7M (SPL)	13.9			

Cluster size	Voxels in cyto	% Cluster	Hem	Cyto or Anatomical description	% Area	Peak t-value	x	y	z
	18.8	1	R	Area 7M (SPL)	18.4				
	11.1	0.6	R	Area 7P (SPL) Precuneus	2.4	4.17	8	-76	50
	6.3	0.3	L	Area PFt (IPL)	1.1				
	4.8	0.3	R	Area 7A (SPL)	0.6				
	4.4	0.2	L	Area PF (IPL)	0.8				
	3.1	0.2	L	Area 2	0.6				
	2.6	0.1	L	Area 7PC (SPL)	1.5				
	1.8	0.1	L	Area PFm (IPL)	0.3				
			R	Precuneus		9.27	8	-64	40
			L	Precuneus		5.81	-4	-72	42
			L	Middle Occipital Gyrus		5.63	-28	-66	32
927	110.5	11.9	L	Area 44 IFG (p. Triangularis)		5.89	-50	16	26
	9.5	1	L	Area 45					
			L	Precentral Gyrus		6.96	-40	4	32
			L	IFG (p. Triangularis)		6.80	-40	10	26
			L	Middle Frontal Gyrus		4.93	-44	4	52
			L	Precentral Gyrus		4.92	-42	0	40
			L	IFG (p. Opercularis)		4.13	-48	10	14
749	29	3.9	R	Area 33 ACC	13.4	9.18	6	4	28
	16.5	2.2	L	Area 33	7.8				

1623

Supplementary Table 5. BOLD activity associated with the main effect of Outcome. Only clusters surviving
a 5% FWE correction at the cluster size are reported (t=3.47, p < .001, cluster size 749; resampled voxel size:
2x2x2mm; Supplementary Figure 2). Brain regions are identified using the Anatomy Toolbox[30]. The columns
refer to the size in voxels of each cluster; the number of voxels of that cluster falling within a specific
cytoarchitectonic region; the percentage of voxels in that region; hemisphere; cytoarchitectonic region (if
available) or macro-anatomical description of the region; percentage of cytoarchitectonic region activated by
cluster; peak t-value within a particular region; and MNI coordinates of the peak. If more peaks were identified
within the same cyto-architectonic or anatomical region only the peak with the highest t-value was included in the
table. Peaks falling outside the gray matter are not included in the table. Cyto architectonic description is only
reported when a voxel has a probability over 40% to fall in that area and only for cyto-architectonic areas available
in the anatomy toolbox; anatomical description is otherwise reported. The '*.txt' file generated by the Anatomy
toolbox can be found at: https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254

1636

1637

Cluster size	Voxels in cyto	% Cluster	Hem	Cyto or Anatomical description	% Area	Peak t-value	x	y	z
Linear regression of PE_s and 1-wf (5% FWE correction . t=3.47 . p<.001 . k=162)									
166	63.3	38.1	L	Area 4a Precentral Gyrus	6.8	5.63	-32	-30	64
	31.5	19	L	Area 4p	9.7				
	29.1	17.5	L	Area 3b Postcentral Gyrus	5.2	5.27	-44	-20	54
	0.5	0.3	L	Area 1	0.1				
162	23.6	14.6	R	Area s32 Mid Orbital Gyrus Area Fo2	16	3.80 3.77	6	36	-12
	4.7	2.9	R	R IFG (p. Orbitalis)	0.4		30	28	-18

0.9	0.5	R	Area Fo2	0.2				
0.4	0.2	R	Area s24	0.3				
0.3	0.2	R	Area 33	0.1				
		R	Insula Lobe		4.79	26	24	-12
		R	Rectal Gyrus		4.65	20	18	-12

Supplementary Table 6. BOLD activity covarying with PEs in a way that depends linearly on wf. Results of the linear regression of PEs and 1-*wf*. Note that because the PES values from M2Out have been divided by 1-*wf* prior to entering them into the first level parametric regressor, the PES parameter estimates would no longer depend on 1-wf if PES signals were similarly strong across participants. Only clusters surviving a 5% FWE correction at the cluster size are reported (t=3.47, p < .001, cluster size 162; resampled voxel size: 2x2x2mm; Supplementary Figure 2). Brain regions are identified using the Anatomy Toolbox[30]. The columns refer to the size in voxels of each cluster; the number of voxels of that cluster falling within a specific cytoarchitectonic region; the percentage of voxels in that region; hemisphere; cytoarchitectonic region (if available) or macro-anatomical description of the region; percentage of cytoarchitectonic region activated by cluster; peak t-value within a particular region; and MNI coordinates of the peak. If more peaks were identified within the same cyto-architectonic or anatomical region only the peak with the highest t-value was included in the table. Peaks falling outside the gray matter are not included in the table. Cyto architectonic description is only reported when a voxel has a probability over 40% to fall in that area and only for cyto-architectonic areas available in the anatomy toolbox; anatomical description is otherwise reported. The '*.txt' file generated by the Anatomy toolbox can be found at: https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254

Cluster size	Voxels in cyto	% Cluster	Hem	Cyto or Anatomical description	% Area	Peak t-value	x	y	z
colspan				Linear regression of PE$_S$ (5% FWE correction . t=3.47 . p<.001 . k=132)					
580	39.4	6.8	L	Area s32 Mid Orbital Gyrus	18.8	4.29	-12	34	-12
	31.3	5.4	R	Area Fp1 Middle Frontal Gyrus	1.8	4.82	26	50	4
	16.1	2.8	L	Area Fp2 ACC	2.2	4.47	-4	50	-2
	5	0.9	R	Area 33	2.3				
	3.6	0.6	L	Area Fo2	0.3				
	2.1	0.4	R	Area s32	1.5				
			R	ACC		5.09	10	38	0
			R	Mid Orbital Gyrus		5.04	16	42	-4
			L	Superior Orbital Gyrus		4.25	-20	38	-12
			R	Mid Orbital Gyrus		4.24	2	40	-6
153	7.9	5.1	L	Area Fp1	0.4				
			L	Middle Frontal Gyrus		5.34	-36	52	4
			L	IFG (p. Triangularis)		4.24	-36	36	10
132	9.8	7.4	L	Area 3a	3.4				
	9.8	7.4	L	Area 4p Postcentral Gyrus	3	4.28	-52	-8	26
	9.1	6.9	L	Area 44	1				
	3.3	2.5	L	Area 3b	0.6				
			L	Precentral Gyrus		4.22	-50	-6	22

1655

Supplementary Table 7. BOLD activity covarying positively with PES after removing variance explained

by wf. Results of the constant in the linear regression of PE$_S$ and 1-*wf*, capturing voxels where the parametric

modulator for PES is above zero after removing variance explained by 1-*wf*. Note that because the PE$_S$ values

from M2Out have been divided by 1-wf prior to entering them into the first level parametric regressor, the PE$_S$

parameter estimates used in this analysis would survive this contrast particularly well if all participants had

similarly positive prediction errors. Only clusters surviving a 5% FWE correction at the cluster size are reported

(t=3.47, p < .001, cluster size 132; resampled voxel size: 2x2x2mm; Supplementary Figure 2). Brain regions are

identified using the Anatomy Toolbox[30]. The columns refer to the size in voxels of each cluster; the number of

voxels of that cluster falling within a specific cytoarchitectonic region; the percentage of voxels in that region;

hemisphere; cytoarchitectonic region (if available) or macro-anatomical description of the region; percentage of

cytoarchitectonic region activated by cluster; peak t-value within a particular region; and MNI coordinates of the

peak. If more peaks were identified within the same cyto-architectonic or anatomical region only the peak with

the highest t-value was included in the table. Peaks falling outside the gray matter are not included in the table.

Cyto architectonic description is only reported when a voxel has a probability over 40% to fall in that area and

only for cyto-architectonic areas available in the anatomy toolbox; anatomical description is otherwise reported.

The '*.txt' file generated by the Anatomy toolbox can be found at:

https://osf.io/rk8w4/?view_only=98b193a58aff48dda40b9d3d91ac5254

1673

1674

11. fMRI Parameter Recovery

The average correlation between the time courses of the parametric modulators for PES and PEM was -0.26, ranging from -0.49 to -0.03. Due to this correlation, we explored whether our experimental design and GLM approach can disentangle voxels that represent PES from those representing PEM, and whether they can differentiate voxels linearly dependent on *wf* from those that are not. Our GLM included, during the outcome period, a boxcar for the duration of the movie with two parametric modulators, one for PES and one for PEM. Both have been normalized by dividing them with 1-*wf* and wf respectively. This was done, as described in the Methods and Materials section of the main manuscript, to ensure that PES and PEM predictors become independent of preference and *wf* per se. When used in the GLM, the parameter estimates for these normalised PES values can then be compared across participants to identify if the brains of participants with higher weight on shocks (i.e. larger value for 1-*wf*) show larger signals for a given outcome than participants with lower weight on shocks. Using the original PES values would make that interpretation difficult, because they are already dependent on *wf*.

For this parameter recovery, we ran 1000 simulations. In each, we simulated 25 participants. For each participant, we used their own design matrices (the same used for the actual GLM first level analysis of the fMRI activity after convolution with the haemodynamic response function) to mix signals in each subject using three mixings (i) -1*PES+0*PEM+noise; (ii) 0*PES+1*PEM+noise, and (iii) -1*PES+1*PEM+noise. Noise was a random gaussian set at 1std of the mixed signal. Next, we ran a GLM using the same design matrix, and saved the parameter estimates for PES (we will call βPES) and PEM (we will call βPEM) for each participant. We then perform a t-test for βPES and one for βPEM to see if across the 25 parameter estimates (one per participant) there is evidence against the null hypothesis H0:βPES=0 or H0:βPEM=0. Of course, if PES was mixed into the voxels activity (case i or iii), a significant t-test would be a hit, while a non-significant t-test would be a miss, and the same applies to PEM for case ii. After repeating this procedure 1000 times, we count the proportion of the 1000 simulations where a t-test was significant against H0:βPES=0 or βPEM=0. Additionally, to see how often the analysis falsely detects a dependence on wf although wf was not included in the mixing, we also look at r(*wf*, βPES)=0 and r(*wf*, βPEM)=0. Initially, we use $p<0.05$ as a criterion, to look at the specificity and sensitivity for the case in which we explore responses in the AVPS, which is univariate. We also indicate proportions at $p<0.001$, but this time for a one-tailed test, in parenthesis, to provide results relevant for an

explorative whole brain analysis where the cluster-cutting threshold was set at 0.001. The proportion of significant results was as follows:

H0　　　　　mixing	-1·PES+0·PEM	0·PES+1·PEM	-1·PES+1·PEM
βPES=0	99.8(92.4)	4.4(0)	99.8(90.6)
βPEM=0	5.5(0.1)	99.8(90.8)	99.5(91.9)
r(wf, βPES)=0	5.2(0.7)	5.3(0.1)	5.4(0.4)
r(wf, βPEM)=0	4.5(0.2)	5.1(0.1)	3.8(0.0)

Supplementary Table 8: Percentage of two-tailed t-tests significant at p<0.05 from the 1000 simulations using signals generated without multiplications with wf or (1-wf), and in brackets, the percentage of one tailed $p_{1\text{-tailed}}$<0.001. The top row specifies how the signals were generated before adding 1sd of noise, the leftmost column, the null hypothesis that was tested in the hypothesis testing.

We then repeated the same analysis, but this time multiplying the signals with (1-*wf*) and *wf* as indicated in Supplementary Table 9 to simulate cases of voxels where signal strength depends on preference.

H0　　　　mixing	-1·(1-wf)·PES+0·wf·PEM	0·(1-wf)·PES+1·wf·PEM	-1·(1-wf)·PES+1·wf·PEM
βPES=0	99.3(81.1)	3.5(0.3)	99(80.5)
βPEM=0	3.8(0.2)	98.5(64.8)	98.7(62.3)
r(wf, βPES)=0	75.9(26.4)	5.1(0.0)	74.6(26.5)
r(wf, βPEM)=0	5(0.0)	98.3(77.3)	98.5(79.3)

Supplementary Table 9: Percentage of two-tailed t-tests significant at p<0.05 from the 1000 simulations using signals generated with multiplications with *wf* or (1-*wf*), and in brackets, the percentage of $p_{1\text{-tailed}}$<0.001. The top row specifies how the signals were generated before adding 1sd of noise, the leftmost column, the null hypothesis that was tested in the hypothesis testing.

The above tables explore evidence against the null hypothesis, but for univariate analysis we also ask whether we can actually provide evidence for voxels mixed without a certain factor that the GLM provides evidence for the null hypothesis using Bayesian statistics (8), using a bound of BF_{10}<1/3. Using a Bayesian test, with n=25, we know that |t|<1 provides evidence in

favour of H0: βPES=0 over H1: βPES≠0, and |r|<0.17 for H0: r(wf, βPES)=0 over H1: r(wf, βPES)≠0 (BF$_{10}$<1/3, using default priors in JASP). We thus counted the proportion with evidence in favour of H0 over H1 in all cases using these bounds.

H0 mixing	-1·PES+0·PEM	0·PES+1·PEM	-1·PES+1·PEM
βPES=0	0	69.6	0
βPEM=0	66	0	0
r(wf, βPES)=0	57.1	58.5	59.3
r(wf, βPEM)=0	58.0	56.8	58.4

Supplementary Table 10: Percentage of BF$_{10}$<1/3 from the 1000 simulations using signals generated without multiplications with wf or (1-wf). The top row specifies how the signals were generated before adding 1sd of noise, the leftmost column, the null hypothesis that was tested in the hypothesis testing.

H0 mixing	-1·(1-wf)·PES+0·wf·PEM	0·(1-wf)·PES+1·wf·PEM	-1·(1-wf)·PES+1·wf·PEM
βPES=0	0	70.1	0
βPEM=0	68.1	0.1	0
r(wf, βPES)=0	2.0	57.5	3.6
r(wf, βPEM)=0	57.5	0.0	0.0

Supplementary Table 11: Percentage of BF$_{10}$<1/3 from the 1000 simulations using signals generated with multiplications with wf or (1-wf). The top row specifies how the signals were generated before adding 1sd of noise, the leftmost column, the null hypothesis that was tested in the hypothesis testing.

Summary: In our simulations, with 1sd of noise, we can detect voxels with signals linearly dependent on PES and/or PEM accurately: If we use alpha=0.05, as we would for the AVPS analysis, signals generated by including PES but not PEM are detected as representing PES in ~99% of cases, and only in ~5% of cases as representing PEM, and vice versa for voxels generated to include PEM but not PES. Using Bayesian statistics, we can even provide evidence in favour of the H0 for the former (βPES=0) in ~70% of cases, and the latter H0 (βPEM=0) in 66% of cases. Within our sample size, we thus have power to arrive at

1750   conclusions that match the way we generated the signals in the majority of simulations. Even

1751   when p<0.001 is used, as it would for our exploratory whole brain analysis, power remains

1752   decent.

1753

1754   With regard to linear dependence on wf, we find that for voxels generated with PES but not

1755   PEM signals, if the signal was generated using (1-*wf*) as a multiplier, a significant correlation

1756   is detected in 76% of cases, and when not used in the generation, a significant correlation is

1757   found in 5% of cases, while evidence for H0:r(*wf*, βPES)=0 is found 57% of cases.

1758

1759   12. Psychological description of the difference between our learning
1760   models

1761

All our learning models had in common that they learned by updating expected values for the

two symbols using prediction errors, and additively combined self-money and other-shock. All

of them captured individual variability in preference using a weighting factor (*wf*). What varied

was when the two outcomes were combined. In our model-free version (M1), the outcomes

themselves are combined into a single composite outcome value, which is already biased by

the participants preference (Figure 2). Translated in psychological terms, this model would

capture a model-free learning in which each symbol is associated with a value that captures

how good or bad the outcomes for those symbols have felt in the past. In case of devaluation,

this model-free learning doesn't switch its preference to the symbol that has the highest

expected value on the remaining quantity because it does not have an internal model that

separates expectations for self-money and other-shock. In contrast, in our unbiased model-

based version (M2Dec), participants track expectations separately for self-money and other-

shock - independently of preference - and preference only plays out during the decision-phase.

In psychological terms, this captures an unbiased model-based learning in which participants

separately know how likely each symbol will lead to self-money or other-shock, respectively.

Only when a decision must be taken, will they combine these predictions with the relative value

that self-money and other-pain have for them personally, to come to a decision under conflict

(Figure 2). In this version, the variable 'expected value' represents expectation in the objective

units in which the outcome is coded, independently of whether this particular outcome is more

or less valued by the participant in the conflict situation. Accordingly, in case of devaluation,

the participant can base their decision on accurate predictions for the remaining quantity, and
should have ~80% preference for the symbol with the highest expected value for the remaining
quantity. Finally, in our *biased* model-based version (M2Out), participants also track separate
expectations for both symbols in terms of self-money and other-shock, but they do so in ways
that depend on their individual preference. Specifically, when self-money and other-shock
outcomes are revealed, they are multiplied with their subjective weight (*wf* for self-money and
1-wf for other-shock, Figure 2), and expectations are updated using these weighted values. At
the decision-stage, the comparison between the two symbols is then done based on a simple
sum of these already weighted expected values. Psychologically, this means that people have
separate models for self-money and other-shock, but that the models predict the subjective
value of each choice rather than the objective outcomes they are associated with. As a result,
in case of devaluation, participants will have weaker preferences for the symbol that leads to
more favorable outcomes if the outcome that remains was the outcome they valued less, than
if the outcome that remained was the outcome they valued more.

1796

13. Stress Tolerance short questionnaire (STSQ)

1798 The Stress Tolerance Short Questionnaire is designed to measure the stress tolerance quotient
1799 and the participants' anxiety level.
1800 In the first part participants have to rate each item from 1 (always) to 5 (never), according to
1801 how much of the time the statement is true for them.
1802

	Always	Frequently	Neutral	Sometimes	Never
1. I eat at least one balanced meal a day.	1	2	3	4	5
2. I exercise to the point of perspiration at least twice a week.	1	2	3	4	5
3. I limit myself to less than half a pack of cigarettes a day	1	2	3	4	5

4. I take fewer than five cups of tea a week.	1	2	3	4	5
5. I regularly attend social activities.	1	2	3	4	5
6. I am in good health (including eye-sight, hearing, teeth).	1	2	3	4	5
7. do something for fun at least once a week.	1	2	3	4	5
8. I am able to organize my time effectively.	1	2	3	4	5
9. I drink fewer than three cups of coffee a day.	1	2	3	4	5
10. I take some quite time for myself during the day.	1	2	3	4	5

1803

1804 In the second part participants have to rate each item from 0 (not at all sure) to 3 (nearly every

1805 day), according to how often they have been bothered by the following problems over the last

1806 2 weeks.

1807

	Not at all sure	Several days	Over half the days	Nearly every day
11. Feeling nervous, anxious, or on edge	0	1	2	3
12. Not being able to stop or control worrying	0	1	2	3
13. Becoming easily annoyed or irritable	0	1	2	3
14. Feeling afraid as if something awful might happen	0	1	2	3

1808

If participants obtained a score higher than 50 in the STSQ we would advise them not to

continue with the experiment. This is because during the experiment they would be confronted

with a number of possibly stressful videos, and put in a position in which their own benefit is

detrimental to another individual. However, in case participants obtained a score higher than

50, we still gave them the possibility to continue with the task, after witnessing a sample video

displaying a high intensity electrical stimulation to the confederate. None of our participants

obtained a score higher than 50 in the STSQ.

14. Debriefing Questionnaire

Please indicate how much you agree with the following statements	strongly disagree	disagree	Somewhat disagree	Neither agree or disagree	Somewhat agree	agree	totally agree
1. I felt uncomfortable when I saw the person in the videos getting the painful stimulation							
2. I am concerned about the welfare of the person who will receive the electrical stimulations							
3. I very quickly understood the different outcomes associated with each symbol							
4. Once I discovered the outcome-symbol association, I voluntarily chose the most financially rewarding symbol for myself							
5. I always felt guilty when I saw a video depicting a high intensity electrical stimulation							
6. I felt responsible for my choices							
7. I felt uncomfortable when not knowing the consequences of each symbol							
8. Before the experiment started I felt distressed							
9. During the experiment I felt distressed because of the conflictual situation							
10. I now feel distressed because of the experience							
11. It was easy to understand the task							
12. I was motivated throughout the task							
13. I participated to earn money							
14. I participated to contribute to scientific knowledge							

15. I participated out of curiosity						
16. I felt I could help the person who will receive the electrical stimulations by choosing one symbol instead of the other						
17. I feel that conflicts of this type are part of my daily life						

This questionnaire was administered after the learning task, to assess the motivation of the participants and the distress caused by the experiment. Highlighted in yellow are the questions which we considered more interesting to analyze, in relation to the behaviour in the learning task.

-Question 3: No effect of group on the declared understanding of the task ($F_{(2,76)}=0.53$, p=0.594, $BF_{incl}=0.165$).

-Question 5: effect of group on the sense of guilt when witnessing the high intensity electrical stimulations (Main effect of group on responsibility, $F_{(2,76)}=4.61$, p=0.01, $BF_{incl}=3.77$). Particularly Considerate declared to feel more guilty than both Lucrative (Independent sample t-test $t_{(51)}=2.85$, p=0.01, $BF_{10}=6.93$) and Ambiguous (Independent sample t-test $t_{(53)}=-2.38$, p=0.02, $BF_{10}=2.44$); while there was no evidence of difference between the amount of guilt reported by Lucrative and Ambiguous (Independent sample t-test $t_{(48)}=0.83$, p=0.41, $BF_{10}=0.38$).

-Question 6: absence of evidence of the effect of group on the amount of responsibility perceived during the task (Main effect of group on responsibility, $F_{(2,76)}=2.10$, p=0.130, $BF_{incl}=0.564$)

-Question 12: The three groups do not differ in the amount of motivation ($F_{(2,76)}=0.05$, p=0.95, $BF_{incl}=0.115$)

-Sum of the answers to question 8-9-10: No effect of group on the amount of stress caused by the task ($F_{(2,76)}=0.769$, p=0.47, $BF_{incl}=0.201$).

15. Eye gaze analysis

In an independent group of participants (N=41) we tested whether participants maximizing self-gain would look away from the facial expressions of pain (compared to participants minimizing the pain to others), to reduce their moral conflict. We collected eye gaze data through Prolific.sc. Specifically, we correlated the percentage of time spent by participants fixating the facial expression of pain videos in the Conflict condition, with the percent of considerate choices they made. As the percentage of fixation on the videos was not normally distributed (Shapiro-Wilk= 0.914, p=0.004) we calculated the Kendall's τ. Supplementary figure 7 shows the results of this correlation: despite a tendency in the expected direction, the relation was not significant and the Bayes factor leaned in favour of the null hypotheses. This preliminary data speaks against the idea that participants with more lucrative preference strongly avoided to look at the facial expressions of pain compared to participants with more considerate preference.

Supplementary Figure 7. Correlation between time spent looking at the video and considerate choices in the Conflict condition.

16. References

1. S. Gallo, *et al.*, The causal role of the somatosensory cortex in prosocial behaviour. *eLife* **7** (2018).

2. M. D. Mckay, R. J. Beckman, W. J. Conover, A Comparison of Three Methods for Selecting Values of Input Variables in the Analysis of Output From a Computer Code. *Technometrics* **42**, 55–61 (2000).

3.   N. J. Evans, J. S. Trueblood, W. R. Holmes, A parameter recovery assessment of time-
variant models of decision-making. *Behav. Res. Methods* **52**, 193–206 (2020).

4.   B. M. Turner, P. B. Sederberg, S. D. Brown, M. Steyvers, A Method for Efficiently
Sampling from Distributions with Correlated Dimensions. *Psychol. Methods* **18**, 368–
384 (2013).

5.   M. H. Davis, A. P. Association, A multidimensional approach to individual differences
in empathy. *JSAS Cat. Sel. Doc. Psychol.* (1980).

6.   K. T. Yamauchi, D. J. Templer, The development of a money attitude scale. *J. Pers.*
Assess. **46**, 522–528 (1982).

7.   C. Keysers, V. Gazzola, E.-J. Wagenmakers, Using Bayes factor hypothesis testing in
neuroscience to establish evidence of absence. *Nat. Neurosci.* **23**, 788–799 (2020).

8.   C. Keysers, V. Gazzola, E.-J. Wagenmakers, Author Correction: Using Bayes factor
hypothesis testing in neuroscience to establish evidence of absence (Nature
Neuroscience, (2020), 23, 7, (788-799), 10.1038/s41593-020-0660-4). *Nat. Neurosci.*
(2020) https:/doi.org/10.1038/s41593-020-00710-7.

1878

1879

1880

1881

1882

1883

1884

1885

1886

1887

1888

1889

1890

1891

1892

1893

1894

1895