1 **Assessing the reliability of species distribution projections in climate change research**
2
3

4 Luca Santini [1-2*], Ana Benítez-López [2-3], Mirza Čengić [1], Luigi Maiorano [4], Mark A.J. Huijbregts [1]
5
6 [1] National Research Council, Institute of Research on Terrestrial Ecosystems (CNR-IRET), Via
7 Salaria km 29.300, 00015, Monterotondo (Rome), Italy
8

9 [2] Department of Environmental Science, Institute for Wetland and Water Research, Faculty of
10 Science, Radboud University, P.O. Box 9010, NL-6500 GL, Nijmegen, The Netherlands
11
12
13 [3] Integrative Ecology Group, Estación Biológica de Doñana (EBD-CSIC), Sevilla, Spain
14
[4] Department of Biology and Biotechnologies, Sapienza Università di Roma, Rome, Italy

* corresponding author: luca.santini.eco@gmail.com

Abstract

Forecasting changes in species distribution under future scenarios is one of the most prolific

areas of application for species distribution models (SDMs). However, no consensus yet exists on

the reliability of such models for drawing conclusions on species distribution response to changing

climate. In this study we first provide an overview of common modelling practices in the field by

reviewing the papers published in the last 5 years. Then, we use a virtual species approach and three

commonly applied SDM algorithms (GLM, MaxEnt and Random Forest) to assess the estimated

(cross-validated) and actual predictive performance of models parameterized with different

modelling settings and violations of modelling assumptions.

Our literature review shows that most papers that model species distribution under climate

change rely on single models (65%), include models fitted on small samples (< 50 presence points,

62%), use presence-only data (85%), and binarize models' output to estimate range shift,

contraction or expansion (74%). Our virtual species approach reveals that the estimated predictive

performance (TSS and AUC) tends to be over-optimistic compared to the real predictive

performance. Further, the binarization of predicted probabilities of presence reduces actual models'

predictive ability considerably. Interestingly, sample size is one of the main predictors of real

accuracy, but has little influence on estimated accuracy. Therefore, models based on larger sample

sizes are more reliable than reflected by their estimated performance. Finally, the inclusion of

irrelevant predictors and the violation of modelling assumptions tends to increase estimated

accuracy but decrease real accuracy of model projections, and leads to biased estimates of range contraction and expansion.

Our study calls for extreme caution in the application and interpretation of SDMs in the context of biodiversity conservation and climate change research, especially when modelling a large number of species where species-specific model settings become impracticable.

Keywords

Area Under the Curve (AUC), bias, climate change projections, disequilibrium, geographic extent, sample size, Niche modelling, spurious relationships, True Skill Statistics (TSS).

1. Introduction

Understanding how climate shapes species distribution and how range shifts may be driven by future climatic change is more urgent than ever. In the last thirty years, studies aimed at developing, improving and applying species distribution models (SDMs) have proliferated (Araújo et al. 2019), and forecasting changes in species distribution under future scenarios is one of the most popular areas of application for SDMs today (Thuiller et al. 2011, Schloss et al. 2012, Newbold 2018). In SDM-based climate change forecasting studies, models are trained on current data and used to predict the probability of presence under present and future conditions. Models' predictions are often binarized, allowing one to assess whether a species distribution is expected to shift, contract or expand (Newbold 2018). Although many modelling techniques require presence and absence data, many models are fitted using presence-only data, i.e., contrasting presences with random pseudo-absences, or background points, that represent available conditions (Guillera-Arroita et al. 2015). The predictive performance of these models is commonly assessed by randomly splitting the dataset into training and testing, and fitting the model on the training dataset and validating it on the testing dataset using discrimination metrics such as the True Skill Statistic (TSS) or the Area Under the Curve (AUC). While several authors have warned about the challenges and uncertainties of projecting future species distribution (Dormann 2007, Peterson et al. 2018), only few studies have tested model performance with empirical data, reporting mixed results (Rapacciuolo et al. 2012, Morán-Ordóñez et al. 2017, Sofaer et al. 2018).

The literature on SDMs has grown very quickly and extensively, with papers adhering to different schools of thoughts and supporting the use of one or another technique (see Norberg et al. 2019 for an overview), suggesting different validation measures (e.g. Allouche et al. 2006, Leroy et al. 2018) or approaches (e.g. testing on spatially independent data; Bahn and McGill 2013). Additionally, a number of studies made different conclusions about the minimum number of presence points needed (Stockwell and Peterson 2002, Hernandez et al. 2006, Wisz et al. 2008, van Proosdij et al. 2016), area of sampling of background points (e.g. VanDerWal et al. 2009, Anderson and Raza 2010, Elith et al. 2010, Barve et al. 2011), or choice of environmental predictors and approaches to reduce collinearity (see Fourcade et al. 2018 for an overview). This can make it challenging and disorientating for people that approach the field of SDM for the first time. The existence of modeling software that make the application of these models easier and more accessible to people with limited modelling background (e.g. MaxEnt Phillips et al. 2006), may be counterproductive, as running an SDM in one of these software may appear simpler than it is. This is particularly worrying considering that SDMs are largely used to inform conservation science (Newbold 2018, Manish and Pandit 2019).

Recently, a number of experts have delineated a set of best practices, and shown that many

studies still apply inconsistent approaches that do not adhere to the best standards (Araújo et al.

2019). This generates a self-perpetuating problem, because published papers create a precedent, and

are used to justify modelling choices in new papers. For example, while several authors argued that

models' predictors should be chosen considering the biology of the species (Araújo and Guisan

2006, Austin and Van Niel 2011), it has become a common practice to include all bioclimatic

variables excluding collinear variables using automatic procedures irrespective of species-specific

biological considerations (e.g. Manish and Pandit 2019), increasing the risk of detecting spurious

relationships (Synes and Osborne 2011, Fourcade et al. 2018). Worryingly, it has been shown that

non-biologically relevant predictors can contribute to increase the predictive ability of the models

(Fourcade et al. 2018), so discrimination accuracy metrics may suggest a very good model while the

relationships estimated do not have a biological meaning (Journé et al. 2019, Warren et al. 2020).

Spurious relationships become particularly problematic when the model is projected to new areas or

environmental scenarios (Heikkinen et al. 2012, Bahn and McGill 2013, Yackulic et al. 2013,

Merow et al. 2014). Similarly, methodological papers that suggest less demanding requirements can

become preferred and widely cited references, reinforcing the trend. For example, van Proosdij et

al. (2016) concluded that 14 to 25 observations may be sufficient to run species distribution models.

This is now often cited to justify the use of small sample sizes (e.g. Carlson et al. 2017, Chen et al.

2017) despite previous recommendations suggesting a minimum of 50 points (Stockwell and

Peterson 2002, Hernandez et al. 2006, Wisz et al. 2008).

The extent to which SDMs perform adequately also depends on the degree to which

modelling assumptions are met. SDMs are often fitted on opportunistically collected data that

violate the assumption of random sampling (e.g. Guillera-Arroita et al. 2015). Furthermore, the

present distribution of species is rarely in equilibrium with the environment, meaning that species

only occupy a portion of the fundamental niche, not only because of biotic (e.g. competition or

predation) or dispersal constraints (e.g. physical barriers, limited dispersal abilities; Soberon and

Peterson 2005), but also because they may have recently contracted their distribution due to human

influence (e.g. Varela et al. 2009, Di Marco and Santini 2015, Faurby and Svenning 2015) or

stochastic events. This problem has often been discussed in the literature in relation to the

inferences made (Varela et al. 2009, Maiorano et al. 2013, Martínez-Freiría et al. 2016, Faurby and

Araújo 2018). Yet, methodological papers aimed at assessing optimal settings to run species

distribution models typically assume ideal conditions (e.g. van Proosdij et al. 2016).

Models used for future projections to inform conservation need to adhere to even higher

standards than those used for present predictions. In fact, while a model used for predicting current

distribution can still provide meaningful predictions even though the inferred relationships are

wrong (Fourcade et al. 2018, Warren et al. 2020), relationships need to be realistic in order to make

meaningful predictions to different conditions. However, one can only validate models on present

data and assume they perform equally well for future predictions. While a number of studies have

discussed and tested the influence of multiple sources of uncertainty on the predictive accuracy of

SDM predictions under present conditions (e.g. Wenger and Olden 2012, Vale et al. 2014, Fourcade

et al. 2018, Fernandes et al. 2019), to our knowledge, no study has yet tested the reliability of both

present and future predictions while considering the effects of different modelling settings and

several violations in model assumptions simultaneously.

In this study we first provide an overview of common practices in the field by reviewing the

papers published in the last 5 years. We focused on the sample size used, choice and selection of

environmental predictors, types of models employed, the sampling approach of background (or

pseudo-absence) points, and the method used for binarization of model outputs. Then, we employed

a virtual species approach (Zurell et al. 2010, Meynard et al. 2019) to assess the contribution of

different modelling settings and violation of assumptions to the predictive accuracy and projected

responses to climate change of SDMs for three commonly applied model algorithms (GLM,

MaxEnt and RandomForest). Our approach allows validating model predictions against the virtual

"reality", therefore estimating true model predictive accuracy. We generated 50 virtual species

distributions, fitted SDMs under different conditions, and assessed the discrimination ability of

present and future model predictions against the real distribution. We also compared this predictive

ability with that estimated using a cross-validation approach, which is the most common way of

assessing model discrimination accuracy in most SDM studies. We systematically assessed the

combined effect of 1) the number of presence points (i.e. sample size), 2) the geographic extent

over which background points are drawn, 3) the number of biologically relevant (i.e. true niche

axes) and irrelevant predictors (i.e. spurious correlates), 4) the species prevalence (proportion of

study area occupied by the species), 5) the sample prevalence (proportion of presences over

background points), 6) the proportion of niche filling (the degree to which the species is at

equilibrium with the environment), 7) and the spatial bias in presence points. We then assessed

model predictions using two common discrimination metrics: the AUC and the TSS.

146
147   **2 Methods**

148
149   <u>2.1 Literature review</u>

150   We conducted a literature review on common practices in SDM papers that projected models to a

151   different time period (past or future). We queried Web of Science and focused on papers published

152   in the last 5 years (2015-2019) to reflect the most recent trends in the field. We randomly selected

50 papers per year for a total of 250 papers. From each paper we extracted the following information: sample size, occurrence data type (e.g. presence-only vs. presence-absence), models used, the variable selection criteria, and whether probabilistic output were binarized or not. A detailed description of the literature search and data extraction is presented in Appendix S1.

2.2 Environmental variables

We obtained 19 bioclimatic variables from CHELSA (http://chelsa-climate.org; Karger et al. 2017) for the present and for the future (year 2050) RCP 8.5 taking the median over all the General Circulation Models (GCM). We also downloaded the human footprint index for 2009 from https://wcshumanfootprint.org/ (Venter et al. 2016).

2.3 Virtual species

We generated 50 virtual species using the 'virtualspecies' R package (Leroy et al. 2016). For each virtual species, we first generated a random extent between 3 and 10 decimal degrees centered around a random location in the globe (Fig. 1). We then selected 6 random bioclimatic variables and sampled their values within the extent using 100 random points. We used the mean and standard deviation estimated for the 6 bioclimatic variables to generate the niche tolerance for the virtual species.

We then projected the niche within the study area for present and future conditions and binarized the distribution into presence/absence using a threshold sampled randomly between the 0.2 to 0.8 quantiles of the suitability values in the study area. Note that the virtual species can potentially be present outside the original study area in environmentally analogous conditions, but we assume that the species is either limited by dispersal, absent because of biotic interactions, or its presence outside the study area is simply unknown to the modeller.

2.4 Scenario settings

For each virtual species, we fitted species distribution models using different cross-combinations of the settings presented in Table 1. To assess the influence of sample size, we sampled random presence points (10, 25, 50, 100, 250, 500 and 1000 points) within the distribution area of the species. Then, to assess the influence of the geographic extent, we fitted a minimum convex polygon (MCP) around presence areas to generated a buffer expressed as percentage of the MCP, which delimited the geographic extent within which the background points were sampled (0%, 100%, 500%, 5,000%, 50,000%; the latter often resulting in the entire continent). We set the number of background points depending on the number of presence points and the level of sample

prevalences. We used three sample prevalence: 0.01, 0.1 and 1. Not all background points,

however, could always be sampled depending on the selected geographic extent (i.e. insufficient

number of cells), leading to variable sample prevalence values.

In each model, we used a total number of predictor variables between 3 and 12. To assess the

influence of biologically relevant and irrelevant predictors of species presence, we sampled none, 3,

or 6 relevant bioclimatic predictors (those describing the true species niche), and none, 3 or 6

irrelevant bioclimatic predictors (not describing the niche) from the other 13 bioclimatic variables

(Table 1). Combinations yielding 0 predictor variables were not considered. We tested collinearity

using a stepwise VIF selection and only retained variables with VIF<3 (Zuur et al. 2010), so the

final number of biologically relevant or irrelevant predictors could be different from multiples of 3.

We also considered violations of two important assumptions underlying SDMs that are common in

real study cases: non-equilibrium with the environment (niche filling) and non-random sampling of

presence points that results in a bias along an environmental gradient. Decreasing proportions of

niche filling were simulated by only sampling presence points below a given quantile (0.33, 0.66, 1)

of the human footprint index values within the study area (Table 1). This mimics a scenario where a

species is potentially present (given climatic conditions) and yet absent because of human impact.

Note that species may be in disequilibrium with the environment for different reasons (e.g. biotic

interactions, dispersal limitations) but the result would be similar. For simplicity we restrict our

analyses to the case where species are not an equilibrium because of human impact.

Sampling bias was simulated by randomly sampling one of the biologically relevant

bioclimatic predictors used in the distribution model and sampling presences only below a given

quantile (0.33, 0.66, 1) of the distribution of environmental values (Table 1). This represents the

situation where the species has only been observed under certain conditions (i.e. sampling bias

correlates with environmental gradients, "environmental bias"), therefore potentially biasing the

estimation of the species niche. When no biologically relevant variable was included, an irrelevant

predictor was selected instead. The full set of combinations of settings in Table 1 corresponded to

7560 models; to reduce the computational effort we sampled 500 model settings from the

multidimensional space using a conditional Latin hypercube approach (Minasny and McBratney

2006), which ensured that the subset of models is representative of the real variability occurring in

the original 7560 models.

2.5 Model fitting and validation

We used this synthetic dataset to fit three distribution model algorithms: MaxEnt (using a 'cloglog'

transformation), Generalized Linear Model (GLM) and Random Forest. For each, we run a repeated

split sample cross-validation by splitting the dataset into training (80%) and testing datasets (20%)

10 times. We estimated model discrimination accuracy with the Area Under the Curve (AUC) and

the True Skill Statistic (TSS) (Lawson et al. 2014). We binarized the predictions into presence-

absence by using the threshold that maximized TSS. Finally, we validated the model predictions for

the present, future, and areas of contraction and expansion against the virtual reality using the same

discrimination metrics. We matched the predicted probabilities with the true presences and absences

of the virtual species to estimate the true AUC, and the predicted presences and absences from the

binarized model with the true presences and absences of the virtual species to estimate the true TSS

using the threshold that maximized TSS on the testing dataset. By doing this, we were able to both

1) estimate model discrimination accuracy mimicking a typical ecological modeller, and 2) quantify

the real model discrimination accuracy by comparing the model to the virtual reality.

234
<u>2.6 Evaluation of model settings</u>

As a post-processing step, we used a Random Forest regression to estimate the influence of

different modelling settings and confounding factors on the discrimination accuracy of the three

distribution model algorithms. We fitted a Random Forest with 1,000 trees to each species using all

discrimination performance metrics (TSS and AUC, both estimated and true for the present and the

future) and estimated changes in distribution (% of range contraction and expansion) as dependent

variable, and the values of each treatment (number of presences, sample prevalence, species

prevalence, number of relevant predictors, number of irrelevant predictors; % buffer, degree of bias

in sampling points, niche filling proportion) as independent variables. The 'mtry' parameter in the

random forest model was set to the number of predictors divided by 3 (Breiman 2001).

We then estimated the relative importance by permutation and partial response curves for

each predictor per species, and then averaged all relative importance estimates and partial response

curves per variable across all species. The estimated relative importance values were transformed to

percentages (rescaled to 100) for interpretability. Confidence intervals for both variable importance

and partial response curves were estimated from the standard error of the mean across species

models.

251
<u>2.7 R packages</u>

All analyses were computed in R v. 3.5.3 (R Core Team 2018) using the packages

'virtualspecies' (Leroy et al. 2016), 'raster' (Hijmans and van Etten 2014), 'PresenceAbsence'

(Freeman and Moisen 2015), 'dismo' (Hijmans et al. 2017), 'rgeos' (Bivand and Rundel 2013),

'pROC' (Robin et al. 2013), 'usdm' (Naimi 2013) and 'GISTools' (Brunsdon and Chen 2014) for

generating virtual species and fitting species distribution models, and 'clhs' for the conditional

Latin hypercube sampling (Roudier 2011). We used the R package 'ranger' for fitting Random

Forest models (Wright and Ziegler 2017) and 'pdp' for estimating the partial response curves

(Greenwell, Brandon 2019). The codes used for the analyses of this paper are available as part of

the supplementary materials.

263
3. Results

3.1 Common practices in SDMs

Among 250 papers revised, 92 included correlative species distribution models projected to

different times, and therefore were deemed relevant for our scopes (Table S1). Based on our sample

and using a bootstrapping approach, we estimated that the total number of papers published

between 2015 and 2019 that matched this criterion is 1194-1665 (95CI), indicating that we sampled

approximately between 5.5 and 7.7% of the total (Appendix S1).

Most of the papers inspected included models fitted on relatively small sample sizes (N < 50; Fig.

2a), with only 18.4% including minimum samples larger than 50 and 16.1% not reporting the

sample size used. More than 50% of the papers included all bioclimatic variables with no biological

justification (Fig. 2b). Among these papers, in ~50% of the cases the authors reduced the number of

variables using automatized approaches based on correlations or best fit to the data. A smaller

number of studies selected variables a priori, some of which did not provide a justification for this

choice (7.8%). Most of the studies used a single model (Fig. 2c,d), with MaxEnt being the most

common algorithm used (78.3%), followed by linear models (GLM = 30.4%; GAM = 26.1%) and

machine learning models (RF = 27.2%; GBM = 20.7%) (Fig. 2c). The majority of studies did not

include real absences but used pseudo-absences, background data, or presence-only methods (i.e.

climatic envelopes; 84.8%; Fig. 2e). A large proportion of papers using pseudo-absences or

background points did not report the area of sampling (48.7%), while others used a variety of

different approaches, the most common being sampling randomly across the pre-defined study area

(Fig. 2f). Finally, most studies binarized the continuous probability outputs based on discrimination

metrics (e.g. max TSS or equal sensitivity and specificity; 73.9%), almost one quarter of the studies

reported the continuous output (22.8%), and a small percent (3.3%) categorized probabilities into

multiple arbitrary categories (e.g. 0.3, 0.6 and > 0.6; Fig. 2g).

3.2 Reliability of climate change predictions

The three algorithms showed a consistent pattern across the two scenarios and accuracy metrics.

The predictive accuracy of models' predictions estimated by cross-validation was consistently

above the acceptable performance thresholds (AUC=0.7, TSS=0.7; Landis and Koch 1977, Swets

1988), and higher than the true predictive accuracy for present, future predictions, and contraction

and expansion areas (Fig. 3). However, the accuracy of binary predictions (TSS) was substantially

lower than that measured for continuous predictive outputs (AUC), suggesting that the binarization

of relative probabilities of presence decreases models' predictive ability considerably (Fig. 3).

Under optimal modelling settings (e.g. large sample size, relevant predictors, no violation of

assumptions regarding niche filling and unbiased sampling), models performed relatively well

according to AUC (Fig. S1a), but poorly when considering binary outputs (Fig. S1c). On the

contrary, under poor modelling settings and conditions (small sample size, irrelevant predictors,

violation of the main assumptions), the estimated predictive abilities remained high, but the true

predictive abilities dropped considerably, especially when predictions were binarized into presence-

absence (Fig. S1b,d). TSS and AUC were highly correlated, but while high TSS always

corresponded to high AUC, the opposite was not always true (Fig. S2).

3.3 Determinants of estimated predictive ability

The importance and effect of different factors on the estimated predictive accuracy by cross-

validation was qualitatively similar when using TSS or AUC (Fig. S2-S8). The most important

predictors of estimated predictive accuracy were species prevalence, the environmental gradient

sampled (inverse of environmental bias), and the geographic extent of background point sampling

(Fig. 4). Additionally, sample prevalence was important for Random Forest, and the number of

presence points for GLM (Fig. 4). Predictive accuracy decreased with increasing species prevalence

and decreasing environmental gradient sampled (i.e. increased with environmental bias), and

increased with increasing geographic extent sampled (Fig. S3-S8). The number of presence points

had a positive effect when fitting GLM, but was only weakly positive in MaxEnt models, and had

no effect when using Random Forests. Sample prevalence a had positive effect in Random Forests,

and weakly negative in the other two models. The number of relevant and irrelevant predictors had

a weak but positive effect regardless of the model (Fig. S3-S8).

3.4 Determinants of true predictive ability

The true predictive accuracy (i.e. measured against the virtual reality) of the models for the present

was mostly affected by sample prevalence, the number of presences, and the environmental gradient

sampled. The geographic extent was also important when fitting MaxEnt models, and sample

prevalence in the case of Random Forests (Fig. 4). Both the number of presence points and the

environmental gradient sampled had a positive influence on predictive accuracy, and geographic

extent and sample prevalence had a positive influence in MaxEnt and Random Forests, respectively

(Fig. S3-S8).

The number of biologically relevant and irrelevant predictors, and niche filling, became more
influential for the predictive accuracy of models projected into the future, with the number of
relevant predictors and niche filling increasing predictive performance, and the number of irrelevant
predictors decreasing predictive performance (Fig. S3-S8).
Species with high prevalence were more likely to expand and less likely to contract the
range. However, a number of additional factors contributed to these estimates (Fig. S9-S12), such as
the number biologically relevant and irrelevant predictors, showing a positive effect on contraction
and expansion estimates in GLM and MaxEnt, and a negative effect in random forest (Fig. S10-
S12). Violation of equilibrium and random sampling assumption also contributed to increase range
contraction and expansion estimates (Fig. S10-S12).
339
340
4. Discussion
In this paper we report on common practices in SDM and use this information to assess the effects
of these practices on the predictive accuracy of SDMs, and thus, on the reliability of future climate-
induced range shifts. Our literature review points out that a large part of papers that model species
distribution under climate change rely on single models (typically MaxEnt), include models fitted
on very small samples, use presence-only data, and typically binarize models' output to measure
range shift, contraction or expansion. Consistently with previous analyses (Araújo et al. 2019), it
also highlighted how poor modelling practices are common in the literature, especially in relation to
the use of very small samples, lack of ecological considerations in the selection of model predictors,
and non-reporting of fundamental information on background sample selection and study area
(Zurell et al. 2020).
When exploring the influence of these practices on the predictive accuracy using a virtual species
approach, we found out that the estimated discrimination capacity by TSS and AUC does not reflect
the actual predictive ability of SDMs, and tends to be over-optimistic compared to the real model
performance when predicted under present conditions, and especially when projected to future
(different) conditions. The ability of models to discriminate presences from absences as measured
by the TSS is particularly low, even under optimal model settings, good ecological knowledge of
the species climatic requirements, and modelling assumptions are fully met. The extent to which
predictions are reliable depends on a number of model parameters (e.g. number of presence points),
actual proportion of species distribution within the geographic extent (species prevalence) and our
degree of knowledge of the species ecology (predictor variables included in the model). Under
optimal settings and a good ecological knowledge of the species climatic requirements, future
predictions show low discrimination ability, whereas under non-optimal settings, predictions may

not be better than random. Ultimately, our results suggest that irrespective of the estimated

performance, we may be unable to make meaningful future predictions for many species, and even

when we can, binarization of models' outputs should be avoided. Based on our results, we elaborate

in the following paragraphs on guidelines and recommendations for good modelling practices when

fitting SDMs.

4.1 Aim for large sample sizes

An important determinant of predictive accuracy that is often undervalued is sample size.

Previous studies suggested a minimum of 50 points (Stockwell and Peterson 2002, Hernandez et al.

2006, Wisz et al. 2008), and van Proosdij et al. (2016) suggested even fewer were needed.

However, these studies assessed the number of points needed under optimal conditions where the

modeller uses biologically relevant environmental predictors, points are sampled randomly, and

species are in equilibrium with the environment; or used real species (therefore estimating accuracy

with testing data, e.g. Wisz et al. 2008). Our results show that while sample size has a little

influence on the estimated (cross-validated) accuracy, it is one of the most important predictors of

true accuracy. The relationship with sample size is asymptotic, and tends to stabilize around 200-

500 points. We must stress, however, that no magic number exists, and these values are contingent

on the settings in our simulation (e.g. the number of predictor variables used in the models).

Because we are rarely aware if the predictor variables are directly linked to species ecology, or if

the species is in equilibrium with the environment or presence points are biased, one should always

aim for the largest possible sample. This may be impracticable for many species, that are either

poorly known, or narrow ranged. In the absence of biological information on e.g. species' thermal

tolerance, it is hard to say, however, if species that are narrow ranged are specialist of specific

climate conditions, or are in disequilibrium with the environment. This second case likely would

result in an under-estimation of niche tolerance and over-prediction of range contraction under

climate change (Araújo and Pearson 2005, Martínez-Freiría et al. 2016, Faurby and Araújo 2018).

In these cases, alternative conservation assessments should be considered when possible. Projecting

SDMs trained on insufficient samples does not improve our knowledge in any meaningful way and

may actually be detrimental.

4.2 Behold sample prevalence, not the absolute number of background points

Many SDM studies using presence-only data sample a large number of background points or

pseudo-absences (e.g. 10,000), often citing Barbet-Massin et al. (2012) or Phillips and Dudík (2008)

as supporting reference. However, Barbet-Massin et al. did not test MaxEnt, and showed important

differences between algorithms. In turn, Phillips and Dudík (2008) tested MaxEnt but they report

their results for many species with different numbers of presence points. Hence, the positive

relationship between AUC and the number of background points they found should be interpreted

carefully as it is mediated by sample prevalence. A recent study concluded that the number of

background points depends on the modelling technique used (Liu et al. 2019), with accuracy in

MaxEnt stabilizing above a few hundreds of background points, and large numbers being only

relevant for common species with small samples of training presences. Our results show that GLM

and MaxEnt work best when sample prevalence is very low, supporting the practice of sampling a

large number of background points or pseudo-absences compared to the number of presences.

However, matching the findings by Barbet-Massin et al. (2012), we found that Random Forest

models perform best with high sample prevalence. This reinforces the notion that no rule of thumb

exists and settings should be model- and sample-specific, which is often ignored in ensemble

forecasting approaches that fit all models on the same dataset (e.g. Avalos and Hernández 2015,

Sales et al. 2017).

<u>4.3 Choose predictors carefully</u>

The number and quality of predictors does not seem to have a clear effect on estimated accuracy, if

any, increasing the number irrespective of the true underlying relationship, tends to deceitfully

increase estimated performance, and increase or decrease the estimated range contraction and

expansion. Choosing biologically meaningful predictors may not be particularly problematic when

predicting to present conditions (Fourcade et al. 2018), but it becomes a serious issue when the

model is transferred in space or time (Wenger and Olden 2012). Here we considered an optimistic

scenario where only 6 climatic variables influence species distributions. In reality, there might be

many biologically relevant variables that determine or influence the distribution of a species, but

our results suggest that when model is projected under different conditions is better to aim for few

variables for which we have clear biological expectations than many variables with unclear effects

on the species' distribution (Araújo and Guisan 2006, Austin and Van Niel 2011).

<u>4.4 Geographic extents should accommodate the purpose of the study</u>

Previous studies suggest sampling background points in areas that are potentially accessible to

the species (e.g. biome or continent) (Araújo et al. 2019) or considering the historical biogeography

of the species (Barve et al. 2011, Merow et al. 2013, Cooper and Soberón 2018). This is meant to

allow a fair comparison between what is used and what is available. Sampling over large areas tend

to inflate estimated predictive accuracy, whereas the effect on true predictive accuracy of present

and future predictions is inconsistent across metrics (positive for AUC and negative or flat for TSS)

and models. This suggests that the most appropriate geographic area for sampling background

points varies across species and it should be tailored to the objective of the study. Setting a

biologically meaningful sampling area requires a deep knowledge of species ecology (e.g. dispersal

distance, physical and biotic barriers) and biogeography (e.g. historical distribution), which is

unavailable for most species, an important future avenue of research may be delineating rules of

thumbs that tend to improve accuracy.

<u>4.5 Noise is inevitable</u>

An important driver of the variation in model performance is species prevalence (Leroy et al. 2018).

Our results concur with previous studies showing that generalist species are harder to predict than

specialist species (Evangelista et al. 2008). However, "generalist" and "specialist" are relative terms

in the context of species distribution models, as they are defined based on the geographic extent

being sampled. Species prevalence over the geographic extent is something we are unaware of in

real study cases, and will always be an unknown factor that affects our predictive ability (Leroy et

al. 2018). In this sense, we should aim to optimize other model settings that can be controlled for,

such as the choice of predictors, the sample prevalence or having a biologically plausible

geographic extent.

<u>4.6 Violation of modelling assumptions provides a false sense of accuracy</u>

Species distribution models rely on the assumptions of random sampling and species equilibrium

with the environment. Worryingly, our results show that when these two assumptions are not met,

the estimated accuracy by cross-validation can be inflated, therefore giving the false impression that

the model performs well. The extensive use of citizen science data in SDMs make models

particularly prone to sample bias, with points more often collected in areas highly accessible to

humans (Bean et al. 2012), or in countries that upload their data to platforms like GBIF more

consistently (Meyer et al. 2016). Bias can be controlled via a number of techniques, such as

including covariates that act as a proxy for the bias (Warton et al. 2013), manipulating background

points (Ranc et al. 2016, Vollering et al. 2019), thinning presence points across the geographic

(Veloz 2009) or the environmental space (de Oliveira et al. 2014), or weighting data points (Elith et

al. 2010). While sampling bias can be sometimes obvious when we compare our sample to the

known approximate distribution of the species (e.g. by using IUCN range maps, or atlases), niche

filling is harder to evaluate, as we only have good knowledge of the historical biogeography of a

relatively small number of species. In many cases, the current distribution of species may result in a

circular reasoning, where small ranges may suggest narrow climatic tolerance while the species

only persists in a given geographic area for different reasons, e.g. because of anthropogenic impact

(Di Marco and Santini 2016). Multiple studies have discussed the under-estimation of the niche due

to historical range contractions (Varela et al., 2009; Maiorano et al., 2013; Martínez-Freiría et al.,

2016; Faurby & Araújo, 2018), and demonstrated these may largely influence our future projections

(Martínez-Freiría et al. 2016, Faurby and Araújo 2018). A possibility to alleviate this effect is using

a multi-temporal approach (or time-calibrated models) by including historical data associated with

the corresponding temporal climatic variables in the model training (Nogués-Bravo 2009, Maiorano

et al. 2013). Yet, historical records are rarely available, so we should expect that the niche always

tends to be under-estimated by an unknown extent compared to the true species potential, and

climatic projections may therefore tend to be pessimistic on average about future species occurrence

(Martínez-Freiría et al. 2016, Faurby and Araújo 2018).

<u>4.7 Binarization</u>

Accuracy metrics can fool us easily, and should not be used acritically to assess the reliability of a

model, especially considering that they can provide higher estimates in sub-optimal conditions as

we have shown here (Fig. S1). Some of the problems discussed above arise from the binarization of

probabilistic model outputs into suitable and unsuitable areas (e.g. to determine the area of range

contraction or expansion) based on a threshold. In fact, the true AUC tends to perform better than

true TSS (Fig. S1, S2), and the estimated AUC has similar values to those of the true AUC under

optimal conditions, whereas the estimated TSS is consistently higher than the true TSS, thus

overestimating accuracy. Studies using MaxEnt typically only show AUC values (standard output

of the software), even though model predictions are binarized. Here we show that while AUC is, as

expected, highly correlated with TSS, high AUC can correspond to low TSS (Fig. S2). The problem

arises from the fact that even when the model performs well, the treshold that maximizes

discriminatory ability on the training dataset may not discriminate well true presence/absence,

especially under different environmental conditions. Additionally, classical cross-validation is

performed by using a split-sample approach, but a better and more informative option is to cross-

validate on spatially independent samples (Bahn and McGill 2013, Roberts et al. 2017).

Additionally, other performance metrics focusing on probabilities (e.g. Boyce index) should be

considered when possible. Several authors have argued that binarization should be entirely avoided

unless it is clearly justified by the model application's objective (Guillera-Arroita et al. 2015). Our

results support this recommendation, and actually indicate that binary outputs should never be

considered or used to quantify changes in distribution areas. Alternative approaches to summarize

the results should be considered, such as looking at trends in predicted probabilities per areas.

<u>4.8 Additional sources of uncertainty to be considered</u>

In this study we evaluated the sensitivity of SDM predictions to a number of modelling settings and

common violations of SDM assumptions. However, there are additional factors that we did not

consider that can further contribute to making predictions less reliable. These include the spatial

accuracy of data points in relation to the resolution used (Graham et al. 2008), the taxonomic

accuracy of the data points (i.e., species confused with others, especially from citizen science data),

and the ambiguous taxonomy of the species that may lead to merging data for different species, or

viceversa not considering part of the distribution of a species (Araújo et al. 2019). Furthermore,

species distribution models assume that the species niche is static, thereby ignoring intraspecific

variation and local adaptations across populations (Pearman et al. 2010, Valladares et al. 2014).

This can be particularly problematic in climate change studies, as populations can adapt to climate

change (Hoffmann and Sgró 2011), and different populations can hold diverse degrees of adaptation

potential (Razgour et al. 2019).

4.9 Concluding remarks

Estimating the distribution of a species is a non-trivial task, as it requires a careful consideration of

the biology of the species and its historical biogeography. Uncertainty is expected to be particularly

high in studies modelling hundreds or thousands of species (Warren et al. 2013, 2018, Visconti et

al. 2016, Newbold 2018, Thuiller et al. 2019), where species-specific considerations on the

geographic extent or variables to include become impracticable, and normally the same geographic

extent for sampling background points or set of variables is used. These studies are powerful for

communicating important messages at the level of geographic areas (e.g., biomes) and entire

communities, but need to be interpreted with extreme caution, and are ill-suited for drawing

inferences at the level of species.

Our study indicates that our ability to predict future species distribution is low under on average,

and can be low to the point of not being meaningful when conditions are far from optimal,

especially when models' predictions are binarized. Hence, SDM based climate change forecasting

must adhere to the highest standards, must be clearly described (Zurell et al. 2020), and the

estimated accuracy of models should be interpreted with extreme care, as well as the results,

especially in relation to the quantification of range shifts, contraction and expansion, and the

identification of areas that will be lost or gained. These considerations are also valid (and perhaps

more problematic considering the wide temporal window and static niche assumption) in the case of

hind-casting to paleoclimates, which is now common in studies focused on refugia and

phylogeography (e.g. Svenning et al. 2011). Future research may focus on developing novel

approaches to improve, synthesize and communicate SDM projections.

References

Allouche, O. et al. 2006. Assessing the accuracy of species distribution models: Prevalence, kappa and the true skill statistic (TSS). - J. Appl. Ecol. in press.

Anderson, R. P. and Raza, A. 2010. The effect of the extent of the study region on GIS models of species geographic distributions and estimates of niche evolution: Preliminary tests with montane rodents (genus Nephelomys) in Venezuela. - J. Biogeogr. 37: 1378–1393.

Araújo, M. B. and Pearson, R. G. 2005. Equilibrium of species' distributions with climate. - Ecography (Cop.). 28: 693–695.

Araújo, M. B. and Guisan, A. 2006. Five (or so) challenges for species distribution modelling. - J. Biogeogr. 33: 1677–1688.

Araújo, M. B. et al. 2019. Standards for distribution models in biodiversity assessments. - Sci. Adv. 5: eaat4858.

Austin, M. P. and Van Niel, K. P. 2011. Improving species distribution models for climate change studies: Variable selection and scale. - J. Biogeogr. 38: 1–8.

Avalos, V. del R. and Hernández, J. 2015. Projected distribution shifts and protected area coverage of range-restricted Andean birds under climate change. - Glob. Ecol. Conserv. 4: 459–469.

Bahn, V. and McGill, B. J. 2013. Testing the predictive performance of distribution models. - Oikos 122: 321–331.

Barbet-Massin, M. et al. 2012. Selecting pseudo-absences for species distribution models: How, where and how many? - Methods Ecol. Evol. 3: 327–338.

Barve, N. et al. 2011. The crucial role of the accessible area in ecological niche modeling and species distribution modeling. - Ecol. Modell. 222: 1810–1819.

Bean, W. T. et al. 2012. The effects of small sample size and sample bias on threshold selection and accuracy assessment of species distribution models. - Ecography (Cop.). 35: 250–258.

Bivand, R. and Rundel, C. 2013. rgeos: Interface to Geometry Engine - Open Source (GEOS). R package version 0.3-2. - R Packag. version 0.1-8: 61.

Breiman, L. 2001. Machine Learning. - Mach. Learn. 45: 5–32.

Brunsdon, C. and Chen, H. 2014. GISTools: some further GIS capabilities for R. - R Packag. version 0.7-4. https//CRAN.R-project.org/package=GISTools. in press.

Carlson, C. J. et al. 2017. Parasite biodiversity faces extinction and redistribution in a changing climate. - Sci. Adv. 3: e1602422.

Chen, Y. et al. 2017. Assessing the effectiveness of China's protected areas to conserve current and future amphibian diversity. - Divers. Distrib. 23: 146–157.

Cooper, J. C. and Soberón, J. 2018. Creating individual accessible area hypotheses improves stacked species distribution model performance. - Glob. Ecol. Biogeogr. 27: 156–165.

de Oliveira, G. et al. 2014. Evaluating, partitioning, and mapping the spatial autocorrelation
component in ecological niche modeling: A new approach based on environmentally
equidistant records. - Ecography (Cop.). 37: 637–647.

Di Marco, M. and Santini, L. 2015. Human pressures predict species' geographic range size better
than biological traits. - Glob. Chang. Biol. 21: 2169–2178.

Di Marco, M. and Santini, L. 2016. Climatic tolerance or geographic breadth: what are we
measuring? - Glob. Chang. Biol. 22: 972–973.

Dormann, C. F. 2007. Promising the future? Global change projections of species distributions. -
Basic Appl. Ecol. 8: 387–397.

Elith, J. et al. 2010. The art of modelling range-shifting species. - Methods Ecol. Evol. 1: 330–342.

Evangelista, P. H. et al. 2008. Modelling invasion for a habitat generalist and a specialist plant
species. - Divers. Distrib. 14: 808–817.

Faurby, S. and Svenning, J. C. 2015. Historic and prehistoric human-driven extinctions have
reshaped global mammal diversity patterns. - Divers. Distrib. 21: 1155–1166.

Faurby, S. and Araújo, M. B. 2018. Anthropogenic range contractions bias species climate change
forecasts. - Nat. Clim. Chang. 8: 252.

Fernandes, R. F. et al. 2019. Effects of simulated observation errors on the performance of species
distribution models. - Divers. Distrib. 25: 400–413.

Fourcade, Y. et al. 2018. Paintings predict the distribution of species, or the challenge of selecting
environmental predictors and evaluation statistics. - Glob. Ecol. Biogeogr. 27: 245–256.

Freeman, E. A. and Moisen, G. 2015. PresenceAbsence : An R Package for Presence Absence
Analysis. - J. Stat. Softw. 23: 1–31.

Graham, C. H. et al. 2008. The influence of spatial errors in species occurrence data used in
distribution models. - J. Appl. Ecol. 45: 239–247.

Greenwell, Brandon, M. 2019. pdp: An R Package for Constructing Partial Dependence Plots. - R J.
9: 421–436.

Guillera-Arroita, G. et al. 2015. Is my species distribution model fit for purpose? Matching data and
models to applications. - Glob. Ecol. Biogeogr. 24: 276–292.

Heikkinen, R. K. et al. 2012. Does the interpolation accuracy of species distribution models come at
the expense of transferability? - Ecography (Cop.). 35: 276–288.

Hernandez, P. A. et al. 2006. The effect of sample size and species characteristics on performance
of different species distribution modeling methods. - Ecography (Cop.). 29: 773–785.

Hijmans, R. J. and van Etten, J. 2014. raster: Geographic data analysis and modeling. - R Packag.
version 2.3-12. http//CRAN.R-project.org/package=raster in press.

Hijmans, R. J. et al. 2017. Package ' dismo .' - R Packag. version 1.1-4. https//CRAN.R-

project.org/package=dismo 9: 1–68.

Hoffmann, A. A. and Sgró, C. M. 2011. Climate change and evolutionary adaptation. - Nature 470:
479–485.

Journé, V. et al. 2019. Correlative climatic niche models predict real and virtual species
distributions equally well. - Ecology in press.

Karger, D. N. et al. 2017. Climatologies at high resolution for the earth's land surface areas. - Sci.
Data 4: 170122.

Landis, J. R. and Koch, G. G. 1977. The Measurement of Observer Agreement for Categorical
Data. - Biometrics 33: 159–174.

Lawson, C. R. et al. 2014. Prevalence, thresholds and the performance of presence-absence models.
- Methods Ecol. Evol. 5: 54–64.

Leroy, B. et al. 2016. virtualspecies, an R package to generate virtual species distributions. -
Ecography (Cop.). 39: 599–607.

Leroy, B. et al. 2018. Without quality presence–absence data, discrimination metrics such as TSS
can be misleading measures of model performance. - J. Biogeogr. 45: 1994–2002.

Liu, C. et al. 2019. The effect of sample size on the accuracy of species distribution models:
considering both presences and pseudo-absences or background sites. - Ecography (Cop.). 42:
535–548.

Maiorano, L. et al. 2013. Building the niche through time: Using 13,000 years of data to predict the
effects of climate change on three tree species in Europe. - Glob. Ecol. Biogeogr. 22: 302–317.

Manish, K. and Pandit, M. K. 2019. Identifying conservation priorities for plant species in the
Himalaya in current and future climates: A case study from Sikkim Himalaya, India. - Biol.
Conserv. 233: 176–184.

Martínez-Freiría, F. et al. 2016. Contemporary niche contraction affects climate change predictions
for elephants and giraffes. - Divers. Distrib. 22: 432–444.

Merow, C. et al. 2013. A practical guide to MaxEnt for modeling species' distributions: What it
does, and why inputs and settings matter. - Ecography (Cop.). 36: 1058–1069.

Merow, C. et al. 2014. What do we gain from simplicity versus complexity in species distribution
models? - Ecography (Cop.). 37: 1267–1281.

Meyer, C. et al. 2016. Multidimensional biases, gaps and uncertainties in global plant occurrence
information. - Ecol. Lett. 19: 992–1006.

Meynard, C. N. et al. 2019. Testing methods in species distribution modelling using virtual species:
what have we learnt and what are we missing? - Ecography (Cop.). in press.

Minasny, B. and McBratney, A. B. 2006. A conditioned Latin hypercube method for sampling in
the presence of ancillary information. - Comput. Geosci. 32: 1378–1388.

Morán-Ordóñez, A. et al. 2017. Evaluating 318 continental-scale species distribution models over a
60-year prediction horizon: what factors influence the reliability of predictions? - Glob. Ecol.
Biogeogr. 26: 371–384.

Naimi, B. 2013. usdm: Uncertainty analysis for species distribution models. - R Packag. version
1.1-18. https//cran.r-project.org/web/packages/usdm/index.html: 1–15.

Newbold, T. 2018. Future effects of climate and land-use change on terrestrial vertebrate
community diversity under different scenarios. - Proc. R. Soc. B Biol. Sci. 285: 20180792.

Nogués-Bravo, D. 2009. Predicting the past distribution of species climatic niches. - Glob. Ecol.
Biogeogr. 18: 521–531.

Norberg, A. et al. 2019. A comprehensive evaluation of predictive performance of 33 species
distribution models at species and community levels. - Ecol. Monogr. in press.

Pearman, P. B. et al. 2010. Within-taxon niche structure: Niche conservatism, divergence and
predicted effects of climate change. - Ecography (Cop.). 33: 990–1003.

Peterson, A. T. et al. 2018. Major challenges for correlational ecological niche model projections to
future climate conditions. - Ann. N. Y. Acad. Sci. 1429: 66–77.

Phillips, S. J. and Dudík, M. 2008. Modeling of species distributions with Maxent: New extensions
and a comprehensive evaluation. - Ecography (Cop.). 31: 161–175.

Phillips, S. J. et al. 2006. Maximum entropy modeling of species geographic distributions. - Ecol.
Modell. 190: 231–259.

R Core Team 2018. R: A language and environment for statistical computing. - R Found. Stat.
Comput. Vienna, Austria. https//www.R-project.org/ in press.

Ranc, N. et al. 2016. Performance tradeoffs in target-group bias correction for species distribution
models. - Ecography (Cop.). 40: 1076–1087.

Rapacciuolo, G. et al. 2012. Climatic associations of British species distributions show good
transferability in time but low predictive accuracy for range change. - PLoS One 7: e40212.

Razgour, O. et al. 2019. Considering adaptive genetic variation in climate change vulnerability
assessment reduces species range loss projections. - Proc. Natl. Acad. Sci. U. S. A. 116:
10418–10423.

Roberts, D. R. et al. 2017. Cross-validation strategies for data with temporal, spatial, hierarchical, or
phylogenetic structure. - Ecography (Cop.). 40: 913–929.

Robin, X. et al. 2013. pROC: an open-source package for R and S+ to analyze and compare ROC
curves. - BMC Bioinformatics 12: 77.

Roudier, P. 2011. clhs: a R package for conditioned Latin hypercube sampling. in press.

Sales, L. P. et al. 2017. Model uncertainties do not affect observed patterns of species richness in
the Amazon. - PLoS One in press.

Schloss, C. A. et al. 2012. Dispersal will limit ability of mammals to track climate change in the
Western Hemisphere. - Proc. Natl. Acad. Sci. USA 109: 8606–8611.

Soberon, J. and Peterson, A. T. 2005. Interpretation of Models of Fundamental Ecological Niches
and Species' Distributional Areas. - Biodivers. Informatics 2: 1–10.

Sofaer, H. R. et al. 2018. Misleading prioritizations from modelling range shifts under climate
change. - Glob. Ecol. Biogeogr. 27: 658–666.

Stockwell, D. R. B. and Peterson, A. T. 2002. Effects of sample size on accuracy of species
distribution models. - Ecol. Modell. 148: 1–13.

Svenning, J. C. et al. 2011. Applications of species distribution modeling to paleobiology. - Quat.
Sci. Rev. 30: 2930–2947.

Swets, J. A. 1988. Measuring the accuracy of diagnostic systems. - Science (80-.). 240: 1285–1293.

Synes, N. W. and Osborne, P. E. 2011. Choice of predictor variables as a source of uncertainty in
continental-scale species distribution modelling under climate change. - Glob. Ecol. Biogeogr.
20: 904–914.

Thuiller, W. et al. 2011. Consequences of climate change on the tree of life in Europe. - Nature 470:
531–534.

Thuiller, W. et al. 2019. Uncertainty in ensembles of global biodiversity scenarios. - Nat. Commun.
10: 1446.

Vale, C. G. et al. 2014. Predicting species distribution at range margins: Testing the effects of study
area extent, resolution and threshold selection in the Sahara-Sahel transition zone. - Divers.
Distrib. 20: 20–33.

Valladares, F. et al. 2014. The effects of phenotypic plasticity and local adaptation on forecasts of
species range shifts under climate change. - Ecol. Lett. 17: 1351–1364.

van Proosdij, A. S. J. et al. 2016. Minimum required number of specimen records to develop
accurate species distribution models. - Ecography (Cop.). 39: 542–552.

VanDerWal, J. et al. 2009. Abundance and the Environmental Niche: Environmental Suitability
Estimated from Niche Models Predicts the Upper Limit of Local Abundance. - Am. Nat. 174:
282–291.

Varela, S. et al. 2009. Is current climatic equilibrium a guarantee for the transferability of
distribution model predictions? A case study of the spotted hyena. - J. Biogeogr. 36: 1645–
1655.

Veloz, S. D. 2009. Spatially autocorrelated sampling falsely inflates measures of accuracy for
presence-only niche models. - J. Biogeogr. 36: 2290–2299.

Venter, O. et al. 2016. Sixteen years of change in the global terrestrial human footprint and
implications for biodiversity conservation. - Nat. Commun. 7: 11.

Visconti, P. et al. 2016. Projecting global biodiversity indicators under future development
scenarios. - Conserv. Lett. 9: 5–13.

Vollering, J. et al. 2019. Bunching up the background betters bias in species distribution models. -
Ecography (Cop.). in press.

Warren, R. et al. 2013. Quantifying the benefit of early climate change mitigation in avoiding
biodiversity loss. - Nat. Clim. Chang. 3: 678.

Warren, R. et al. 2018. The projected effect on insects, vertebrates, and plants of limiting global
warming to 1.5°C rather than 2°C. - Science (80-.). 360: 791–795.

Warren, D. L. et al. 2020. Evaluating presence-only species distribution models with discrimination
accuracy is uninformative for many applications. - J. Biogeogr. 47: 167–180.

Warton, D. I. et al. 2013. Model-based control of observer bias for the analysis of presence-only
data in ecology. - PLoS One 8: e79168.

Wenger, S. J. and Olden, J. D. 2012. Assessing transferability of ecological models: an
underappreciated aspect of statistical validation. - Methods Ecol. Evol. 3: 260–267.

Wisz, M. S. et al. 2008. Effects of sample size on the performance of species distribution models. -
Divers. Distrib. 14: 763–773.

Wright, M. N. and Ziegler, A. 2017. ranger : A Fast Implementation of Random Forests for High
Dimensional Data in C++ and R. - J. Stat. Softw. 77: 1–17.

Yackulic, C. B. et al. 2013. Presence-only modelling using MAXENT: when can we trust the
inferences? - Methods Ecol. Evol. 4: 236–243.

Zurell, D. et al. 2010. The virtual ecologist approach: Simulating data and observers. - Oikos 119:
622–635.

Zurell, D. et al. 2020. A standard protocol for reporting species distribution models. - Ecography
(Cop.). in press.

Zuur, A. F. et al. 2010. A protocol for data exploration to avoid common statistical problems. -
Methods Ecol. Evol. 1: 3–14.

Table 1. Summary of treatments considered for fitting the species distribution models on virtual
species.
747

Treatment	Values
Presences	10, 25, 50, 100, 250, 500, 1000
Sample Prevalence	0.01, 0.1, 1
Buffer (%)	0, 100, 500, 5000, 50000
Bias (%)	33, 66, 100
Niche filling (%)	33, 66, 100
Relevant predictors	0, 3, 6
Irrelevant predictors	0, 3, 6

Fig. 1. Modelling steps taken to generate virtual species and fit and project the species distribution
 model.

Fig. 2. Summary of the literature review. (a) Number of presences used in the models (minimum among species if multiple species were modelled); PtP = Polygons converted to presence points; NR = sample size not reported; (b) Variable selection approach. all ClmVr = all climatic variables were considered; sel. ClmVr = a subset of climatic variable was considered; Automatic selection = collinear variables were excluded using automatized approaches based on correlations, variance inflation factors, or best fit to the data; Informed selection = collinear variables were excluded based on expert opinion; No selection = Collinear variables were not excluded, or no collinearity was not found or reported; No justification = no justification provided for the rationale underlying the subset of variables chosen; (c) Percentages of studies using MaxEnt, generalized linear models (GLM), generalized additive models (GAM), random forests (RF), generalized boosted trees (GBM), multivariate adaptive regression splines (MARS), classification trees (CTA), artificial neural networks (ANN) and flexible discriminant analysis (FDA). Other models used in a minority of instances are not reported here (see Table S1); (d) Percentage of studies using one or multiple models, or ensemble modelling approach; (e) Pseudo-absences or background points sampling approach; Bias = sampling that mimics sampling bias; Buffer = random sampling within a buffer around presence points; Distance-weighted = Sampling with higher intensity near (-) or far from (+) presence points; Globally = Random sampling globally; Outside climate envelope = Beyond climatic conditions observed for presence points; Study area = random sampling within a pre-defined study area; (f) Percentage of studies using presences only, presences + background points / pseudo-absences, and those using presences and real absences; (g) Percentage of studies binarizing the probabilities into suitable/unsuitable, or in multiple arbitrary categories, or not applying any form of binarization.

775 **Fig. 3.** AUC (a) and TSS (b) of the models fitted. Estimated = Estimated through internal cross-validation; Present and Future = True value validated
776 against virtual reality for present and future; Contraction and Expansion = True value validated against virtual reality for predicted contraction and
777 expansion areas; Dashed line = null expectation (no better than random); Dotted line = Value typically considered as "good" performance thresholds.
778 The box edges are the 25th and 75th percentiles of the distribution, and whiskers 1.5 the inter-quartile range.
779
780
781
782

Fig. 4. Relative variable importance of different settings and conditions on the TSS (a, b, c) and AUC (d, e, f) estimated by cross-validation (a, d), and measured against virtual reality for the present (b, e) and future predictions (c, f). Relative importance values are rescaled to 100 for each species. Bars represent the mean over all virtual species and error bars the standard error around the mean.