CD38 dependent NAD$^+$ depletion contributes to oligodendrocyte loss and inhibition of

myelin regeneration

Authors

Monica R. Langley[1], Chan-Il Choi[1], Thais R. Peclat[3], Whitney Simon[1], Hyesook Yoon[1,2], Laurel

Kleppe[1], Claudia C.S. Chini[3,4], Eduardo N. Chini[3,4], and Isobel A. Scarisbrick[1,2,*]

Affiliations

[1] Department of Physical Medicine & Rehabilitation, Rehabilitation Medicine Research Center,

Mayo Clinic, Rochester, MN 55905, USA

[2] Department of Physiology and Biomedical Engineering, Mayo Clinic, Rochester, MN 55905,

USA

[3] Robert & Arlene Kogod Center on Aging, Mayo Clinic, Rochester, MN 55905, USA

[4] Department of Anesthesiology, Mayo Clinic, Rochester, MN 55905, USA

[*] Corresponding author. Tel: 507-774-0124; E-mail:scarisbrick.isobel@mayo.edu

Abstract

Western-style diets cause disruptions in myelinating cells and astrocytes within the CNS. We

identified increased CD38 expression in the mouse spinal cord following chronic high fat

consumption or focal demyelinating injury. CD38-catalytically inactive mice are significantly

protected from high fat-induced NAD^+ depletion, oligodendrocyte loss, oxidative damage, and

astrogliosis. 78c, a CD38 inhibitor, increased NAD^+ and attenuated neuroinflammatory changes

in astrocytes induced by saturated fat. Conditioned media from saturated fat-treated astrocytes

impaired oligodendrocyte differentiation pointing to indirect mechanisms of

oligodendrogliopathy. Combined saturated fat and lysolecithin demyelination in cerebellar slices

resulted in additional deficits in myelin proteins that were mitigated by concomitant 78c

treatment. Importantly, oral 78c increased counts of oligodendrocytes and remyelinated axons

after focal demyelination. Our findings suggest that high fat diet impairs oligodendrocyte

survival and differentiation through astrocyte-linked mechanisms mediated by the NAD^+ase

CD38, and highlight the use of CD38 inhibitors as potential therapeutic candidates to improve

myelin regeneration.

Keywords

CD38, Glia, high fat diet, NAD^+, remyelination

Introduction

Obesity is a key risk factor for developing multiple sclerosis (MS) (Gianfrancesco, Acuna et al., 2014), which further increases risk when interacting with other known genetic, lifestyle, and environmental risk factors (Olsson, Barcellos et al., 2017). Moreover, recent studies suggest that obesity and related metabolic comorbidities also influence disease progression and disability burden in MS patients (Manouchehrinia, Hedstrom et al., 2018, Marrie, Rudick et al., 2010, Tettey, Simpson et al., 2014). Although current investigations point to obesity-induced white matter abnormalities and loss of oligodendrocytes within the central nervous system (CNS) following chronic consumption of high fat diets (HFD) in rodent models, how this may influence MS pathophysiology is largely unknown (Graham, Grabowska et al., 2019, Langley, Yoon et al., 2019, Yoon, Kleven et al., 2016).

Astrocyte activation occurs following chronic HFD feeding in the CNS and exerts oligotoxic effects (Buckman, Thompson et al., 2015, Buckman, Thompson et al., 2013, Clarke, Liddelow et al., 2018, Liddelow, Guttenplan et al., 2017, Wanrooy, Kumar et al., 2018). On the other hand, building and maintenance of myelin membranes requires oligodendrocyte fatty acid synthesis. In addition, oligodendrocytes can utilize lipids from the diet or synthesized by astrocytes, at least under normal conditions (Camargo, Goudriaan et al., 2017, Dimas, Montani et al., 2019). The potential to target astrocytes to improve outcomes in MS is already suggested since some currently prescribed MS disease modifying therapies do impact astrocytes (Ponath, Park et al., 2018). Prior studies regarding the role of HFDs in MS animal models have utilized immune-mediated models such as experimental autoimmune encephalomyelitis (EAE) and demonstrate exacerbation of peripheral immune responses or neuroinflammation with HFD consumption (Hasan, Seo et al., 2017, Herrmann, Barth et al., 2016, Ji, Wu et al., 2019,

Timmermans, Bogie et al., 2014). Also, recent findings suggest that insulin sensitizing agents

such as metformin and pioglitazone provide benefit to MS patients with metabolic syndrome

(Negrotto, Farez et al., 2016). There are few if any studies regarding the effects of HFD on the

capacity for remyelination. Since oligodendrocyte function is intimately linked to neighboring

glial cells and neuronal activity, in addition to microenvironmental factors, the most promising

interventions will likely involve a combinatorial approach (Alizadeh, Dyck et al., 2015, Alizadeh

& Karimi-Abdolrezaee, 2016, Ohtomo, Iwata et al., 2018).

CD38 is an NAD^+-degrading enzyme expressed by astrocytes and was recently shown to

be upregulated in MS animal models (Herrmann et al., 2016, Roboon, Hattori et al., 2019,

Zhang, Chen et al., 2014, Zhang, Sloan et al., 2016). CD38 is the main NAD^+ase in many

mammalian tissues and regulates tissue and cellular NAD^+ homeostasis as well as NAD^+

dependent enzymes such as PARPs and Sirtuins (Aksoy, Escande et al., 2006, Aksoy, White et

al., 2006). Global CD38 knockout confers protection against demyelination, T cell responses,

and EAE disease severity as well as HFD-induced metabolic dysfunction in rodent models

(Barbosa, Soares et al., 2007, Chiang, Harrington et al., 2015, Herrmann et al., 2016, Roboon et

al., 2019). Here we test the hypotheses that HFD consumption promotes oligodendrocyte loss in

a CD38-dependent manner and that CD38 inhibition will enhance myelin regeneration. First we

established increased CD38 expression following HFD consumption. Next, we demonstrated that

CD38 catalytically inactive mice fed either a regular or HFD were protected from HFD-mediated

loss of oligodendrocytes with concomitant increases in NAD^+ levels and reduced

neuroinflammation and oxidative damage. *In vitro* approaches identified oligodendrocyte direct

and astrocyte-linked indirect mechanisms by which excessive fatty acid impairs myelinating cell

function. Finally, the value of a small molecule inhibitor of CD38, 78c, was then identified as a

novel protective strategy to enhance myelin regeneration in a model of focal demyelinating

injury. Together these findings highlight a novel indirect pathway of myelin injury involving

pathogenic changes in astrocytes. Moreover, these changes can be attenuated by inhibition of

CD38 in the context of chronic HFD consumption and targeted to improve myelin regeneration

in the CNS.

Results

Astrocytes increase CD38 in the spinal cord following chronic HFD or demyelination

Our recent studies revealed spinal cord changes involving ER stress, mitochondrial dysfunction, and oxidative stress in the spinal cords of mice chronically consuming a HFD and the direct potential of saturated fat palmitate (PA) to inhibit differentiation of primary murine oligodendrocytes from their progenitors or neural stem cells (Langley et al., 2019, Yoon et al., 2016). Although saturated fat inhibited markers of differentiation *in vitro*, it was not oligotoxic, suggesting a more complex mechanism at play *in vivo* (Langley et al., 2019). Considering the emerging role of astrocytes in the pathology of various neurological disorders (Liddelow et al., 2017, Ponath et al., 2018, Yoon, Choi et al., 2020) and in the CNS following chronic HFD (Buckman et al., 2015, Ogrodnik, Zhu et al., 2019, Wanrooy et al., 2018), along with their known essential metabolic and functional links with oligodendrocytes (Domingues, Portugal et al., 2016, Kiray, Lindsay et al., 2016), we decided to address the possibility of indirect effects of HFD mediated by astrocytes (Fig 1A).

NAD^+ is a key mediator of cellular energy status and is involved in mitochondrial function and quality control processes primarily through the activity of sirtuins that require NAD^+ as a cofactor (Camacho-Pereira, Tarrago et al., 2016, Canto, Menzies et al., 2015, Long, Klimova et al., 2017)(Foolad, Khodagholi et al., 2019). Exercise is known to increase NAD^+ levels in various tissues and protected against "Western-style diet"-induced deficits in myelin and myelinating cells in two recent studies (Chiang et al., 2015, Graham et al., 2019, Yoon et al., 2016). Our recent untargeted metabolomics profiling of spinal cord tissue identified NAD^+ metabolism as one of the top pathways impacted by HFD consumption (Langley et al., 2019), which prompted us to quantify spinal cord NAD^+ levels using a cycling assay. Spinal cord NAD^+

levels from HFD-fed mice were less than half that of RD-fed mice (Fig 1B). A heatmap with

HFD-induced gene expression changes in NAD$^+$-related enzymes was also derived from this

papers sequencing results (Figure 1- figure supplement 1a)(Langley et al., 2019).

Since CD38 is the main NAD$^+$-glycohydrolase in many mammalian tissues (Camacho-

Pereira et al., 2016, Chini, Chini et al., 2018), we next quantified CD38 expression in spinal cord

astrocytes to establish a potential link. Quantification by qPCR indicated higher RNA expression

of CD38 in HFD-fed mice compared to regular diet-fed controls (Fig 1C). Consistent with

findings demonstrating CD38 is primarily localized to astrocytes within the CNS (Hattori, Kaji et

al., 2017, Kou, Banerjee et al., 2009, Roboon et al., 2019, Zhang et al., 2014, Zhang et al., 2016),

increases in CD38 immunoreactivity (Fig 1D-G) were observed in GFAP+ astrocytes following

either HFD (Fig 1D,F) or lysolecithin-induced demyelination (Fig 1E,G). Using multiplex *in*

situ hybridization, CD38 RNA levels were also higher in ALDH1L1+ astrocytes of HFD-fed

mice (Figure 1- supplement 1b). Similar to the HFD spinal cords, primary murine astrocytes

cultured with PA (100 µM, 24 h) showed increased GFAP (Fig 1H) and CD38 (Fig 1I) RNA

expression and depleted NAD$^+$ levels (Fig 1J). Collectively, these data suggest that HFD can

induce astrocytosis in the spinal cord or primary cultures, which increases expression of CD38

and subsequently depletes NAD$^+$ levels.

CD38 catalytically inactive (CD38ci) mice are resistant to HFD-induced oligodendrocyte

loss

To determine the potential therapeutic significance of CD38 upregulation to HFD-

induced oligodendrocyte loss, we collected spinal cords from wildtype (WT) and CD38-

catalytically inactive (CD38ci) mice fed a regular diet (RD) or HFD for 30 wk beginning at 50

wk of age for histological analysis (Fig 2A). CD38ci mice have the CD38 gene knocked out and

reinserted with a mutation at E230 of the CD38 gene, rendering the CD38-NAD$^+$ase catalytic

site inactive while preserving overall expression levels of CD38 (Tarrago, Chini et al., 2018). As

seen in our previous studies (Langley et al., 2019, Yoon et al., 2016), there was a significant

reduction in the number of Olig2+ oligodendrocyte progenitors (Fig 2B) as well as GST3+

mature oligodendrocytes (Fig 2C) in the HFD-fed wildtype mice compared to regular diet-fed

mice. However, CD38ci mice fed HFD did not have reductions in oligodendrocyte lineage cell

counts compared to their regular diet genotype controls.   Across spinal cord regions, no

significant changes in MBP due to genotype or diet group were found (Fig 2D). Overall, these

data suggest that oligodendrocyte lineage cell survival is preserved in CD38ci mice following

chronic HFD consumption.

Diminished neuroinflammatory responses and oxidative damage in CD38ci mice following

chronic HFD

Neuroinflammation, including complex interplay between microglia and astrocytes, is

now appreciated to contribute to oligotoxicity and MS pathophysiology (Domingues et al., 2016,

Liddelow et al., 2017). To explore the potential impact of blocking CD38 catalytic activity on

CNS neuroinflammation following chronic HFD consumption, we next performed

immunohistochemical analysis on the spinal cords from wildtype and CD38ci mice fed regular

or HFD for markers of microglia and astrocytes. Quantification of the thresholded area

considered positive for ionized calcium-binding adapter molecule 1 (IBA1) demonstrated

decreased microglia (Fig 3A) in the ventral grey matter of spinal cords from CD38ci mice

compared to wildtype mice overall. In addition, HFD-fed CD38ci mice showed significantly

reduced IBA1 immunoreactivity in the ventral grey and white matter of the spinal cord compared

to wildtype mice fed a HFD. Considering that astrocytes have endogenous expression of CD38

as well as elevated CD38 expression following HFD or demyelination (Roboon et al., 2019,

Zhang et al., 2014, Zhang et al., 2016) (Fig 1), we also investigated astrocyte reactivity in these

mice. In the ventral spinal cord, quantification of astrocyte marker glial fibrillary acidic protein

(GFAP, Fig 3B) demonstrated significantly reduced positive area in the white matter of CD38ci

mice compared to wildtype mice. Wildtype HFD-fed mice showed significantly more astrocyte

reactivity compared to either regular diet or HFD-fed CD38ci mice.

Given our previous data suggesting increased oxidative damage markers in the CNS

following chronic consumption of HFDs (Langley et al., 2019, Yoon et al., 2016) and the

relevance of oxidative stress in glia of MS lesions (Haider, Fischer et al., 2011, Lassmann & van

Horssen, 2016, Miller, Dziedzic et al., 2019), we next sought to compare the extent of oxidative

damage in the spinal cords of wildtype and CD38ci mice fed regular or HFD. More abundant

lipid peroxidation marker 4-hydroxynonenal (4HNE) was present in the HFD-fed wildtype mice

than regular diet-fed wildtype mice in the spinal cord dorsal column. Importantly, a significant

reduction in 4HNE was found in the HFD-CD38ci mice compared to HFD-wildtype mice in the

dorsal column and ventral spinal cord (grey and white matter). To confirm that preservation of

NAD^+ levels was achieved in the CD38ci mice, we next measured NAD^+ levels on spinal cord

tissue from the same cohort of mice. As expected based on our data in Fig 1, NAD^+ levels were

significantly reduced following chronic HFD consumption in wildtype mice (Fig 3D). Moreover,

in CD38ci mice, which had diminished CD38 activity (Fig 3E), NAD^+ levels were significantly

higher overall and in the CD38ci-HFD mice compared to the wildtype-HFD mice. Next, our

histological findings were further substantiated by observed gene expression differences in

SOD2 (Fig 3F) and pro-repair astrocyte markers genes BDNF and S100a10, as determined by

qRT-PCR (Fig 3G-H). Overall, the observed reductions in neuroinflammation and oxidative

stress makers and increased NAD$^+$ levels in the CD38ci mice likely provide a more suitable

microenvironment for oligodendrocyte survival, differentiation, and repair following chronic

HFD consumption.

Astrocytes cultured in high saturated fat conditions increase CD38 and pro-inflammatory

cytokine expression and inhibit oligodendrocyte differentiation

Astrocytes release various growth factors, cytokines, chemokines, extracellular matrix

proteins, glutamate, and reactive oxygen species that heavily influence oligodendrocyte

proliferation and differentiation (Alizadeh & Karimi-Abdolrezaee, 2016, Domingues et al., 2016,

Kiray et al., 2016, Lassmann & van Horssen, 2016). Therefore, we next investigated the effects

of culturing oligodendrocytes in conditioned media obtained from PA-treated astrocytes with or

without 78c co-treatment, as depicted in Fig 4A. 78c is a thiazoloquin(az)olin(on)e CD38

inhibitor that increases NAD$^+$ levels in various tissues and cell types and is orally bioavailable

(Boslett, Helal et al., 2018, Haffner, Becherer et al., 2015, Tarrago et al., 2018). As expected,

astrocyte-conditioned medium (ACM) from PA-treated astrocytes significantly reduced

oligodendrocyte PLP and MBP expression. However, this negative impact on myelin protein

production was partially rescued by co-treatment with the small molecule CD38 inhibitor 78c (3

μM) (Haffner et al., 2015, Tarrago et al., 2018). Likewise, addition of exogenous NAD$^+$ (50

μM) to the PA-treated astrocyte cultures (Fig 4B) significantly increased PLP expression, while

ACM-treated oligodendrocyte numbers remained unchanged across treatment groups. Along

with other research that indicates the role of NAD$^+$ in reducing astrocyte proinflammatory and

oxidative stress responses (Harlan, Pehar et al., 2016, Roboon et al., 2019, Wilhelm & Hirrlinger,

2012), these data indicate the importance of maintaining proper astrocyte NAD$^+$ levels in order

to promote oligodendrocyte differentiation.

To address whether the possible NAD^+-driven astrocytic changes indirectly mediate the

observed effects on oligodendrocyte, we next used primary murine astrocyte cultures to assess

NAD^+ levels and transcriptional changes. Addition of 78c (3µM) was effective to restore NAD^+

levels in PA-treated astrocytes when quantified using an NAD^+ cycling assay (Fig 4C). However,

astrocytes derived from CD38 knockout mice did not have depleted NAD^+ levels in response to

PA. Next, we determined whether CD38 regulates astrocyte pro-inflammatory responses. After

treating astrocyte cultures with PA and/or 78c, we used qRT-PCR and a multiplex bioassay for

determining cytokine RNA transcripts and release into culture supernatant, respectively. We

observed significant increases in IL-1β and IL-6 expression (Fig 4D) in astrocytes stimulated

with PA and 78c significantly attenuated PA-induced RNA expression of IL-1β, IL-6, TNFα, and

reactive astrocyte marker H2D1. Additionally, the mitochondrial antioxidant SOD2 was

significantly increased in astrocytes treated with 78c or both PA and 78c. Significant increases in

secretion of cytokines triggered by PA were observed using a Bioplex assay including CXCL1

(KC), MIP1α, RANTES, IL-6, MIP1β, and MCP1 (Fig 4E). Notably, co-treatment with 78c

successfully mitigated PA-stimulated secretion of CXCL1, MIP1α, RANTES, IL-6, MIP1β, and

MCP1. Thus, inflammatory and oxidative stress-related changes in cultures astrocytes may help

to explain reduced differentiation potential of the oligodendrocyte progenitors in the presence of

PA-treated astrocyte conditioned medium.

NAD^+ and NMN restore oligodendrocyte differentiation in PA-treated cultures

Our recent findings demonstrated that PA can directly impair oligodendrocyte differentiation

from neural stem cells or oligodendrocyte progenitors (Langley et al., 2019), and others have

indicated that depletions in cellular NAD^+ can prevent differentiation of stem cells to mature

oligodendrocytes (Stein & Imai, 2014). To examine the effects of replenishing NAD^+ levels on

oligodendrocytes, we next supplemented cultures with exogenous NAD^+ (50 µM) and monitored

PLP expression (Fig 5A). Co-treatment with PA and NAD^+ significantly enhanced PLP

expression (Fig 5 B) compared to exclusive PA treatment. However, treatment of

oligodendrocytes with CD38 inhibitor 78c did not attenuate this effect (Figure 2- figure

supplement 1).

Interestingly, CD38+ cells can deplete NAD^+ and metabolites available for low-CD38 expressing

cells (Chini, Hogan et al., 2019, Tarrago et al., 2018), such as oligodendrocytes (Hattori et al.,

2017, Roboon et al., 2019, Zhang et al., 2014, Zhang et al., 2016). We found that direct effects of

saturated fat treatment on oligodendrocytes include mitochondria function and morphology

changes (Langley et al., 2019) while other mechanisms, potentially including NAD^+-depleting

mechanisms (Penberthy & Tsunoda, 2009, Penke, Schuster et al., 2017, Stein & Imai, 2014),

could also play a part. We hypothesized that the pro-myelinating effects of increasing NAD^+ are

SIRT1-dependent, so we determined if the beneficial effects of NAD^+ in the presence of PA

could be blocked with a SIRT1 inhibitor Ex-527. Similar to adding exogenous NAD^+, treatment

with the NAD^+ precursor NMN (100 µM) also restored markers of oligodendrocyte

differentiation in PA-treated cultures (Fig 5C). However, in the presence of Ex-527 (5 µM), the

increases in PLP and MBP protein in response to NMN were suppressed. At the doses used, PA,

NMN, and Ex-527 did not significantly alter the number of oligodendrocytes. Hence, the

impaired ability of oligodendrocytes cultured with excess saturated fat to differentiate is at least

partially regulated by NAD^+-dependent sirtuin activity.

78c prevents lysolecithin- and PA-mediated myelin protein and oligodendrocyte loss

Since we observed protective effects of genetic CD38 inhibition on oligodendrocyte

survival following HFD (Fig 2), as well as increased CD38 expression in a demyelination model

(Fig 1), we next sought to explore benefits conferred by pharmacologic inhibition using 78c in a

model of lysolecithin-induced *ex vivo* demyelination. Murine cerebellar slices from postnatal day

7-9 day pups were demyelinated using lysolecithin (LL, 0.5 mg/ml, 18 h), followed by a period

of one week to allow for repair (Fig 6A). Confocal imaging of immunofluorescence staining

demonstrated reductions in MBP after LL that was significantly attenuated by 78c co-treatment

(Fig 6 B).

Finally, to evaluate the therapeutic potential of 78c in a model of high fat exposure

combined with demyelination, organotypic cerebellar slices were demyelinated as described

above and cultured for 7 d in the presence or absence of PA and/or 78c. All cerebellar slices

showed fewer Olig2+ cells and less MBP immunoreactivity following LL insult (Fig 6 C). After

7 d of recovery, MBP levels were successfully restored, although Olig2+ cells remained

significantly depleted in the 7 d LL group. Inclusion of PA in the media further diminished

Olig2+ cells in the LL slices. Importantly, 78c and PA co-treatment significantly increased the

percentage of Olig2+ cells and MBP expression in the demyelinated cerebellar slices. However,

neither palmitate nor 78c alone, without lysolecithin, for two weeks significantly altered MBP

expression (Figure 6- figure supplement 1). These findings demonstrate the capability of 78c to

restore myelin protein expression even in a combined saturated fat- and demyelinating agent-

induced condition.

CD38 inhibition with 78c improves myelin repair following lysolecithin-induced focal

demyelination of the ventral spinal cord

To determine whether pharmacologic inhibition of CD38 NAD$^+$ase activity affects the

potential for remyelination *in vivo*, we administered 78c to mice in each diet group (regular or

high fat) and used histological approaches to investigate numbers of oligodendrocyte lineage

cells and remyelinated axons in the lysolecithin model of demyelination. Six-week old male

wildtype mice were given either a regular or HFD for 5 weeks, then split into additional groups:

receiving the same diet with or without 78c (600ppm) (Fig 7A). All mice were subjected to a

focal demyelinating injury to the ventral spinal cord by microinjection of 1% lysolecithin. At the

study endpoint, HFD mice had gained significantly more weight than their RD counterparts,

irrespective of genotype, while 78c treatment slightly decreased yet did not have a significant

effect on body weights (Figure 7-figure supplement 1a). Relative to wild type mice, the 78c-

treated mice in the regular and high-fat groups consumed significantly less chow (Figure 7-figure

supplement 1b). Based on the average food intake and 78c concentration (600ppm) in the diets, it

can be approximated that mice consumed 16-19.5 mg 78c daily through oral consumption.

Hematoxylin and eosin (H&E) staining indicates that lesion area (Figure 7-figure supplement

1C) and neurofilament (a marker of axon integrity, Figure 7- figure supplement 1D) remain

unchanged by drug or diet.

After two weeks of recovery, lesions were collected from perfused mice and used for

subsequent counting of remyelinated axons and myelinating cells. Overall, lysolecithin mice fed

78c along with either a RD or HFD showed significantly greater numbers of remyelinated axons

compared to the lysolecithin mice fed a RD or HFD without 78c (Fig 7B). In particular, more

than double the number of remyelinated axons was present in the 78c-regular diet mice over

regular diet alone. Further supporting a role in myelin repair, MBP immunoreactivity in the total

lesion or the lesion rim was higher in the 78c-treated mice irrespective of diet (Fig 7C).

Importantly, numbers of Olig2+ (Fig 7D) and GST3+ (Fig 7E) cells were significantly higher in

the combined 78c group (which includes LL-RD-78c and LL-HFD-78c mice) versus the

combined group without 78c treatment (which includes LL-RD and LL-HFD). Moreover, the

number of Olig2+ cells in the lesion rim and GST3+ cells in the rim or total lesion were

significantly higher in the LL-RD-78c versus the LL-RD group. In addition to the demyelinated

lesion site, 78c also attenuated HFD-induced oligodendrocyte loss on the uninjected side of the

ventral spinal cord (Figure 7-figure supplement 2). These findings suggest that, *in vivo,* 78c helps

to preserve or replenish myelinating cells and their progenitors, leading to increased markers of

myelin repair. Furthermore, these results provide a strong rationale for targeting CD38 and

overall NAD$^+$ metabolism as a new strategy to attenuate oligodendrocyte loss and aid in myelin

regeneration.

78c promotes pro-repair astrocytes after lysolecithin-induced demyelination

Next, we chose to explore the role of 78c on neuroinflammation markers in the spinal cord

following demyelination. Histological quantification of threshold-positive staining for microglia

marker IBA1 (Fig 8A) indicated a significant interaction between diet and drug treatment on the

lesion rim, yet no significant effect of diet or drug alone. Although there is a trend of decrease

for both IBA1 and GFAP in the 78c treated group, no statistically significant attenuation of

astrocyte marker GFAP was observed (Fig 8B). In both the RD- and HFD-fed mice, there was,

however, a significant increase in S100a10 immunoreactivity with 78c treatment (Fig 8C),

considered a marker for pro-repair astrocytes. There was also a slight decrease in C3d staining, a

marker of reactive astrocytes, although this did not reach statistical significance (Fig 8D).

Overall, these data suggest that 78c modulates astrocyte responses elicited by demyelinating

injury, skewing them towards a phenotype that enhances oligodendrocyte survival and

differentiation and myelin repair (Fig 9).

Discussion

Diet-induced obesity increases the risk for developing MS, especially when combined with other known risk factors (Gianfrancesco et al., 2014, Mokry, Ross et al., 2016, Olsson et al., 2017). The results of the current study suggest that metabolic disturbances, triggered by consumption of a diet high in fat, promote oligodendrogliopathy and impair myelin regeneration through astrocyte-linked indirect NAD^+-dependent mechanisms. We demonstrate that restoring NAD^+ levels via genetic inactivation of CD38 can overcome these effects. Moreover, we show that therapeutic inactivation of CD38 can enhance myelin regeneration. Together these findings point to a new metabolic targeting strategy positioned to improve disease course in MS and other conditions in which the integrity of myelin is a key concern.

Neuroinflammation and astrogliosis are predominant pathological mediators of neurodegenerative conditions, including MS, and astrogliosis is linked to HFD-induced CNS pathology in rodent models (Buckman et al., 2015, Buckman et al., 2013, Liddelow et al., 2017, Mizunoya, Ohnuki et al., 2013, Ogrodnik et al., 2019, Ponath, Ramanan et al., 2017, Thaler, Yi et al., 2012, Wanrooy et al., 2018). In spinal cord astrocytes of HFD mice, we observed increases in CD38 expression and an overall depletion of NAD^+. This suggests that CD38, the major constitutively active NAD^+-degrading enzyme, is induced in astrocytes with chronic exposure to a HFD. These findings are consistent with outcomes in studies of neuroinflammation that indicate increased expression of CD38 in the brain of individuals with MS or human immunodeficiency virus encephalitis and in rodent experimental models of Alzheimer's disease (AD), sepsis associated brain injury, and toxin-induced brain convulsions (Blacher, Dadali et al., 2015, Ginwala, McTish et al., 2016, Kou et al., 2009, Machado-Santos, Saji et al., 2018, Mamik, Banerjee et al., 2011, Peng, Wang et al., 2018, Zou, He et al., 2017). In the current study, we

found that HFD or focal lysolecithin-mediated demyelinating injury each drove increased CD38

expression in astrocytes, while genetic or pharmacological CD38 inhibition after HFD or

demyelination, respectively, attenuated astrocyte activation.

Upstream activators of CD38 expression include cytokines, lipopolysaccharide (LPS),

25-hydroxycholesterol (25-HC), glutamate, and factors secreted by senescent cells (Bruzzone,

Verderio et al., 2004, Chini, Hogan et al., 2019, Chini et al., 2018, Mamik et al., 2011).

Separately, astrocytosis and CD38 are reported to increase with normal aging, and growing

evidence supports a role for HFD-induced neuroinflammation in promoting premature

"inflammaging" (Camacho-Pereira et al., 2016, Chini et al., 2019, Clarke et al., 2018, Denver,

Gault et al., 2018, Escande, Nin et al., 2013, Freeman, Haley-Zitlin et al., 2014). Likely related

to the older age of mice in the current study (~80 wks), HFD did not increase IBA1 or GFAP

levels. In palmitate-stimulated astrocytes, downstream inflammatory indicators of TLR4

activation, including IL-6 and TNFα were increased. Pointing to a therapeutic strategy, this

response was blunted when NAD$^+$ levels were restored by co-treatment with 78c. Also, studies

reporting the role of CD38 in neuroinflammation indicate CD38 inhibition or knockout decreases

cytokine secretion in LPS-stimulated microglia, macrophages, and monocytes (Amici, Young et

al., 2018, Iqbal & Zaidi, 2007, Matalonga, Glaria et al., 2017, Roboon et al., 2019, Wang, Liu et

al., 2017). Therefore, improving NAD$^+$ levels by targeting CD38 shows therapeutic promise

across multiple neurological and non-neurological disease types, and here we establish its

significance to oligodendrocyte heath and regeneration with strong linkage to its effects on

astrocyte metabolism.

Astrocytes and microglia secrete a variety of chemokines, cytokines and growth factors

that regulate myelin injury and regeneration (Alizadeh & Karimi-Abdolrezaee, 2016, Domingues

et al., 2016, Kiray et al., 2016). Of cytokines increased in astrocytes after palmitate-exposure,

CCL3/MIP1α, MIP1β, and RANTES also increase in active demyelinating MS lesions and fluids

(Boven, Montagne et al., 2000, Janssen, Rickert et al., 2016, Mori, Nistico et al., 2016). These

factors are thought to enhance disease progression by recruiting peripheral immune cells and by

activating resident glia (Boven et al., 2000, Mori et al., 2016, Tejera-Alhambra, Casrouge et al.,

2015). Opposing effects are seen for CXCL1/KC on myelination and glia activation (Clarner,

Janssen et al., 2015, Karim, Kim et al., 2018), while combined CCL2/MCP1 and CCL3/MIP1α

absence attenuates demyelination, oligodendrocyte loss, and astrogliosis in the cuprizone model

(Janssen et al., 2016). Recently, knockout of CD38 or supplementation with NAD^+ was shown to

suppress activation of astrocytes in culture (Roboon et al., 2019). Here we showed that saturated

fat- induced reductions in NAD^+ result in proinflammatory responses in astrocytes which trigger

impaired differentiation and survival in oligodendrocyte lineage cells. Moreover, we demonstrate

that 78c mitigates the dysmyelinating effects of astrocyte secreted factors.

In addition to cytokines such as IL-1β and TNFα, reactive oxygen species can inhibit

oligodendrocyte survival (Liddelow et al., 2017, Valentin-Torres, Savarin et al., 2016).

Oxidative damage can also activate inflammasome signaling in peripheral immune cells and glia,

including astrocytes (Lassmann & van Horssen, 2016, Liao, Lai et al., 2018, Moyon, Dubessy et

al., 2015, Sarkar, Malovic et al., 2017). Complementing this, we provide new RNA evidence for

more SOD2 and less IL1β in high fat-78c co-treated astrocyte cultures compared to treatment

with high fat alone. We also found more SOD2, BDNF and S100a10 in the CD38ci mouse spinal

cord pointing to a shift towards pro-repair astrocyte properties (Liddelow et al., 2017). Blocking

downstream CD38 signaling also reduced markers of oxidative damage in a rodent toxin-induced

convulsion model (Zou et al., 2017). Our findings suggest that the beneficial effects of CD38

inhibition include protection against astrocyte pro-inflammatory factors and reactive oxygen

species known to negatively impact oligodendrocyte renewal and maturation.

The effects of CD38 activity and NAD$^+$ levels may depend heavily on cell type and

disease context, with findings here supporting a critical protective role of lowering astrocyte

CD38 activity for protection of myelinating cells and for myelin regeneration in the adult CNS.

In addition, CD38+ cells regulate NAD$^+$ levels in neighboring CD38- cells by influencing NAD$^+$

precursor availability (Chini et al., 2019, Tarrago et al., 2018), and this may also be the case for

astrocytes and oligodendrocytes in our cell culture studies of high fat and experimental

demyelination. Consistent with this model, 78c had no direct effect on oligodendrocyte

differentiation in the context of palmitate *in vitro*. In a recent study, palmitate inhibited NAMPT

activity, depleted cellular NAD$^+$ levels, and resulted in apoptosis in neurons at a higher dose than

studied here (Penke, Schuster et al., 2017). Another report indicates NAMPT inhibition prevents

differentiation of stem cells to oligodendrocyte lineage cells (Stein & Imai, 2014). These studies

align well with the direct inhibitory effect of palmitate on oligodendrocyte differentiation

observed here. PARP deficiency has been shown to either increase severity of disease or

attenuate clinical signs in animal models of MS (Selvaraj et al, 2009). Thus, complex alterations

in NAD$^+$ biosynthesis, salvage, and degradation pathways may contribute to the effects of

palmitate on oligodendrocytes that we document can be rescued by NAD$^+$ replenishment.

Several recent studies have explored supplementing NAD$^+$ precursors or flavonoids

known to inhibit NAD$^+$ase activity to improve myelin repair, neuroprotection, and to decrease

immune cellular responses. For example, an NAD$^+$/BDNF/TrkB pathway is proposed to be

involved in nicotinamide-mediated protection, including improved remyelination in a stroke

model and precursor supplementation increased BDNF in a Huntington's disease model

(Hathorn, Snyder-Keller et al., 2011, Wang, Zhang et al., 2017). Supporting a possible NAD^+-

BDNF pathway for remyelination, we observed increased BDNF expression in the spinal cord of

CD38ci mice and improved remyelination with oral 78c administration. Although direct effects

on myelinating cells were not examined, mice with CD38-knockout, given the CD38 inhibitor

apigenin, or supplemented with NAD^+ each showed reduced EAE disease severity (Ginwala et

al., 2016, Herrmann et al., 2016, Wang, Zhao et al., 2016). NAD^+ levels can impact

oligodendrocyte lineage cell proliferation and survival via cyclins and E2f1, while differentiation

is reported to be regulated redundantly by SIRT1 and SIRT2 (Jablonska, Gierdalski et al., 2016,

Ji, Doucette et al., 2011, Penberthy & Tsunoda, 2009, Stein & Imai, 2014). Results here

demonstrate that the ability of NAD^+ supplementation to promote oligodendrocyte differentiation

in vitro is indeed at least partially reliant on SIRT1 function. Restoring NAD^+ levels was also

beneficial in the context of palmitate or high fat-induced inflammation and oxidative stress via

SIRT1 in liver, heart, and muscle (Escande, Chini et al., 2010, Li, Dou et al., 2015, Prola, Pires

Da Silva et al., 2017, Sadeghi, Seyyed Ebrahimi et al., 2017). Improvements in expression of

myelin proteins in the spinal cord of mice consuming a high fat high sucrose diet in conjunction

with coordinate wheel running, were also linked to increases in SIRT1 expression (Yoon et al.,

2016). Collectively, these findings are particularly encouraging since effective modulation of

disease progression will likely include a combination of therapeutic strategies encompassing

anti-inflammatory, myelin regeneration, and neuroprotective interventions.

HFD consumption did not further impair markers of myelin integrity in our *in vivo* model

of myelin injury and repair as it did in *ex vivo* remyelinating cerebellar slice cultures. Although

treatment with 78c did improve myelin regeneration in mice consuming a regular diet, little

impact was seen in mice consuming a HFD, suggesting the need to explore additional factors in

the context of HFD-demyelination that contribute impaired remyelination. Alternately, a higher

dose or even a combination of drugs (to target various aspects of astrocyte responses) may be

required to improve remyelination in the context of HFD. Although a preference for pro-repair

versus pro-inflammatory phenotype was indicated following 78c treatment in the lysolecithin

model of focal demyelination studied, a complete astrocyte-specific transcriptomic profile in the

HFD mice with demyelination will be needed to elucidate specific pathogenic astrocyte

populations. This approach would additionally identify novel genes in the astrocyte

transcriptome that could be targeted to mitigate the indirect oligotoxic effects we observe.

Genetic CD38 inhibition attenuated HFD-induced oligodendrocyte loss and dietary

supplementation with 78c increased counts of oligodendrocytes and remyelinated axons

following focal demyelinating injury. These findings suggest that increases in astrocyte CD38,

and the associated depletion of NAD^+ levels, may contribute indirectly to impairments in

oligodendrocyte health and myelin regeneration via inflammation and oxidative stress. Tarrago

et al. (2018) recently established 78c as a specific, reversible, and non-competitive inhibitor of

CD38 NAD^+ase activity improves age-related metabolic function (Hattori et al., 2017).

Altogether, these findings suggest that genetic or pharmacologic CD38 inhibition provides

therapeutic benefits in HFD-induced metabolic dysfunction and downstream signaling events in

the CNS, in addition to the already described peripheral benefits on age-related metabolic

dysfunction. Importantly, inhibition of CD38 with 78c can improve myelin regeneration in the

adult CNS including reducing the pro-inflammatory profile while increasing the pro-repair

profile of astrocytes. Together these findings point to the potential translational value of targeting

CD38 to improve outcomes in MS and other demyelinating conditions.

A few current MS disease modifying therapies (i.e. glatiramer acetate, fingolimod,

dimethyl fumarate) affect CD38+ cell counts or inhibit astrocyte activation, but these roles have

not been directly linked to therapeutic benefits (Bruck, Popescu et al., 2012, Choi, Gardell et al.,

2011, Herrmann et al., 2016, Pavelek, Vysata et al., 2018, Ponath et al., 2018, van de Donk,

Richardson et al., 2018, Wilms, Sievers et al., 2010). For instance, fingolimod treatment

decreased CD38 expression in a rat EAE model (Herrmann et al., 2016), and was shown to

promote beneficial astrocyte responses in culture (Hoffmann, Hofereiter et al., 2015).

Monoclonal antibodies targeting CD38 have already met with success in clinical trials for

conditions outside the CNS (Chini et al., 2018, Deckert, Wetzel et al., 2014, Lokhorst, Plesner et

al., 2015, Martin, Baz et al., 2017), including Daratumumab which is FDA approved for use in

multiple myeloma. Accordingly, the research findings we present should inspire further research

into ways to mitigate the negative consequences of HFD on oligodendrocytes alone and in the

context of white matter injury which may include dietary intake of NAD^+ precursors,

administration of CD38 inhibitors, or exercise-related rehabilitation strategies.

Materials and Methods

Key Resources Table

Reagent or Resource	**Source**	**Additional information**
Antibodies		
Mouse anti-CD38	Novus	1g7f4 (1:500)
Rabbit anti-GFAP	Dako	Z0334 (1:5000)
mouse anti-Olig2	Millipore	MABN50(1:200)
rabbit anti-Olig2	Millipore	Ab9610; AB_10141047 (1:500)
Rabbit anti-GST3	Abcam	Ab153949 (1:1000)
rat anti-Myelin basic protein (MBP)	Millipore	MAB386; AB_94975 (1:750)
Rabbit anti-PLP	Abcam	A28486 (1:750)
Rabbit anti-IBA1	Wako	019-19741(1:3000)
rabbit anti-4-hydroxynonenol	Abcam	ab46545 (1:500)
chicken anti-GFAP	Abcam	ab4647 (1:2000)
rabbit anti-Neurofilament	Sigma Aldrich	N4142; AB_477272 (1:1300)
Goat anti-C3d	R&D Systems	AF2655 (1:500)
Rabbit anti-S100a10	Proteintech	11250-1-AP (1:7500)
biotinylated goat anti-rat secondary	Jackson	112-066-072 (1:100)
biotinylated donkey anti-rabbit secondary	Jackson	711-065-152 (1:100)
biotinylated donkey anti-mouse secondary	Jackson	715-066-151 (1:100)
goat-anti-rabbit AF488 secondary	Jackson	111-545-047 (1:100)
goat anti-rat Cy3 secondary	Jackson	112-166-072 (1:100)
goat anti-chicken AF647 secondary	Jackson	703-606-155 (1:100)
goat anti-rabbit AF488 secondary	Jackson	111-545-041 (1:100)
donkey anti-rabbit Cy3 secondary	Jackson	711-166-152 (1:100)
goat anti-rabbit AF647 secondary	Jackson	111-606-144 (1:100)
goat anti-rat AF647 secondary	Jackson	112-606-143 (1:100)
Oligonucleotides		

(*primers/probes*)		
BDNF	Applied Biosystems	Mm04230607_s1
SOD2	Applied Biosystems	Mm01313000_m1; NM_013671.3;
S100a10	Integrated DNA Technologies	Mm.PT.58.6571055
CD38	Applied Biosystems	Mm01220906_m1
OLIG2	Applied Biosystems	NM_016967; Mm.PT.58.42319010
MBP	Integrated DNA Technologies	NM_001025251; CCAGTAGTCCATTTCTTCAAGAACAT/GCCGATTTATAGTCGGAAGCTC
PLP	Integrated DNA Technologies	NM_011123.2; TCTTTGGCGACTACAAGACCAC/CACAAACTTGTCGGGATGTCCTA
GFAP	Integrated DNA Technologies	NM_010277.2; GCAGATGAAGCCACCCTGG/GAGGTCTGGCTTGGCCAC
IL1β	Integrated DNA Technologies	Mm.PT.51.17212823
IL6	Applied Biosystems	NM_031168.1; Mm00446190_m1
H2D1	Integrated DNA Technologies	Mm.PT.58.42136026.g
TNFα	Applied Biosystems	Mm00443258_m1
Rn18s	Applied Biosystems	NR_003278.3; Mm03928990_g1
GAPDH	Applied biosystems	Mm99999915_g1
RNAscope Probe - Mm-Aldh1l1-C2	Advanced Cell Diagnostics	405891-C2
RNAscope Probe - Mm-CD38	Advanced Cell Diagnostics	513061
Chemicals		
78c CD38 inhibitor	Millipore	Cat#538763
Nicotinamide adenine dinucleotide	Sigma Aldrich	Cat#10127965001
Nicotinamide 1,N6-	Sigma Aldrich	Cat#41628

ethenoadenine dinucleotide		
flavin mononucleotide	Sigma Aldrich	Cat#F8399
Alcohol dehydrogenase	Sigma Aldrich	Cat#A3263
Diaphorase	Sigma Aldrich	Cat#D5540
resazurin	Sigma Aldrich	Cat#119303
palmitate	Sigma Aldrich	Cat#P0500
DAPI	ThermoFisher	Cat#D1306; AB_2629482
lysolecithin	Sigma Aldrich	Cat#L-4129
Methyl green	Sigma Aldrich	67060
Commercial kits		
RNA scope Duplex Manual kit	Advanced Cell Diagnostics	Cat#322430
Bio-Plex Mouse Cytokine 23-plex Assay	Bio-Rad	Cat#m60009rdpd
Software		
GraphPad Prism 7	GraphPad Software	https://www.graphpad.com/scientificsoftware/prism
CellSens	Olympus	https://www.olympus-lifescience.com/en/software/cellsens/
Zen	Zeiss	https://www.zeiss.com/microscopy/us/products/microscope-software/zen.html
Fuji ImageJ	NIH	https://imagej.net/Fiji

Animal husbandry and diets

All experimental diets and procedures were approved by The Mayo Clinic Institutional

Animal Care and Use Committee (IACUC) and performed in accordance with institutional and

regulatory guidelines. Mice were provided *ad libitum* access to diets along with tap water. For

initial characterization of CD38 expression and NAD^+ levels following a high-fat diet, male

C57BL6/J mice were obtained from Jackson Labs (Bar Harbor, ME) and were randomly

assigned to feeding on either a regular diet (RD, 10 % kcal fat, D12450K, Research Diets, New

Brunswick, NJ) or a high-fat diet (HFD, 60% kcal fat, D12492, Research Diets, New Brunswick,

NJ) beginning at 10 wk of age, as previously described (Langley et al., 2019).

For studies comparing wildtype and CD38ci mice fed RD and HFDs, mice were bred

onsite after initial generation at TransViragen (Chapel Hill, NC) via recombineering technology,

as previously described (Partida-Sanchez, Cockayne et al., 2001, Tarrago et al., 2018). Male

mice received a regular (10.4% kcal fat, Teklad TD000357) or HFD (42% kcal fat, Teklad

TD88137) beginning at 50 wk of age for 30 wk. Diets used for lysolecithin demyelination studies

were these same diets, with or without 600 ppm 78c added, and were started one week prior to

the spinal cord microinjection procedure. Food intake and body weights were monitored weekly.

Lysolecithin microinjection into spinal cord

For focal demyelination of the ventral white matter (laminectomy at thoracic 10^{th}

vertebra; mediolateral 0.3, dorsoventral -1.4 mm relative to the dorsal surface) of the murine

spinal cord, 8 wk old male mice were injected with 2 µl of a saline solution containing 1%

lysophosphatidyl choline with 0.01% Evan's Blue (Sigma L4129) at 0.2 µl/min using a 30-40

µm glass micropitppette and stereotaxic microinjection device (Stoelting, Inc., Wood Dale, IL).

For anesthesia, 100 mg/kg ketamine and 10 mg/kg xylazine were administered

intraperitoneally prior to surgery, while postoperative care consisted of an antibiotic (Baytril 50

mg/kg/day, i.p.), an analgesic (buprinex 0.05 mg/kg, twice daily, subcutaneous) and saline

(0.2ml/d, intraperitoneal) for three days. After a two-week recovery period, mice were deeply

anesthetized (ketamine 150mg/kg and xylazine 15 mg/kg, i.p.) and transcardially perfused with

4% paraformaldehyde (Fisher T353-500) for collection of spinal cord tissue. Spinal cord

segments were microdissected, with the rostral 2 mm segment processed for paraffin-embedded

blocks and the caudal segment osmicated (1% osmium tetroxide, 0223B, Polysciences) and

embedded in Araldite. To identify myelin sheaths, Araldite blocks were cut to 1 µm sections,

stained with p-phenylendiamine (Sigma P6001), imaged with 60x objective (Olympus BX51

microscope), and stitched together (Adobe Photoshop) to comprise the entire lesion. Number of

remyelinated axons per square millimeter of lesion was reported after blinded counting in ImageJ

(Yoon et al., 2020). Immunohistochemical analysis of focal demyelinated lesions utilized overall

neuropathological changes (hematoxylin and eosin) along with myelin loss to determine the

lesion core, while the lesion rim was defined as also encompassing neuroinflammatory markers

(GFAP, IBA1). Lesion areas between 50,000 - 180,000 μm2 were included in the analysis.

Immunohistochemistry

Spinal cords were fixed in 4% paraformaldehyde, processed, paraffin-embedded, and cut

to 6 μm for all immunoperoxidase and immunofluorescence procedures, as described previously

(Langley et al., 2019, Yoon et al., 2020). When necessary, heat-induced antigen retrieval was

achieved through incubation with sodium citrate (10 mM, pH 6.0) in a steamer basket. Sections

were blocked (20 % normal goat serum), then incubated in primary antibodies described in the

key resources table.

The following day, incubation with appropriate secondary antibodies (biotinylated or

fluorochrome-conjugated, Jackson, 1:200) for immunoperoxidase or immunofluorescence

staining was performed. After incubating in an avidin peroxidase solution (Vectastain Elite ABC

kit, PK-6100, Vector Laboratories), color development was achieved by incubation with 3,3-

diaminobenzidine (DAB, Sigma D5637) and counterstaining with methyl green (Sigma, 67060).

Slides were dehydrated, coverslipped with DPX (Sigma), and images acquired using and

Olympus BX51 microscope with DP72 camera. Immunofluroescence slides were counterstained

with 4',6'-diamindino-2-phenylindole (DAPI), mounted with fluoromount, and imaged with an

Olympus BX51 microscope and captured with Olympus XM10 camera. Cell and axon counting,

as well as thresholding of images, was performed in ImageJ (NIH) without knowledge of

treatment groups using cropped regions of interest (Adobe Photoshop) and averaged from two

sections per animal. Rolled or damaged tissue sections were excluded from histological

quantification.

in situ **hybridization using multiplex RNAscope:**

For visualizing RNA expression of CD38 in tissue sections, slides containing RD- or

HFD-fed mouse corpus callosum were processed for RNAscope *in situ* as previously described

(Choi, Yoon et al., 2018, Wang, Flanagan et al., 2012). In short, slides were deparaffinized, and

incubated with hydrogen peroxide for 10 min at room temperature (RT), target retrieval solution

at 100 °C for 30 min, and protease plus solution at 40°C for 30 min. Next, probes (Mm-CD38-

C1, 513061 and Mm-Aldh1l1-C2, 405891, ACD) were incubated at 40 °C for 2 h and sections

were stored in 5x saline sodium citrate (SSC) buffer overnight. The following day, probes were

hybridized and chromogenic development steps were performed in freshly prepared solutions

following manufacturer instructions as previously described (322430, ACD, Newark, CA)1.

Briefly, six amplification steps and washes were performed prior to AP-based Fast Red color

development for 10 min at RT. Then, four additional amplification steps and HRP-based Green

chromogen development was achieved by incubating for 10 min at RT. Stained slides were

counterstained with dilute hematoxylin and allowed to dry for 20 min at 60 °C prior to being

cooled and mounted with Vectamount (H5000, Vector Labs, Burlingame, CA). All probes and

RNAscope 2.5 HD Multiplex reagents were purchased from Advanced Cell Diagnostics

(322430, ACD, Newark, CA).

RNA isolation and qRT-PCR

To determine gene expression from spinal cord or primary murine cells, total RNA was

extracted using RNA Stat60 (Tel-Test; Friendswood, TX) per manufacturer's instructions and

used for qRT-PCR using BioRad iTaq Universal Probes or primers and SYBR Green One-step

kits. Starting quantity for quantification of each gene's expression was normalized to Rn18s or

GAPDH. Detailed primer and probe information are annotated in the key resources table.

Mixed glial preparations and primary cell cultures

Primary mixed glial cultures were prepared from P0-3 mice, as previously described

(Langley et al., 2019, Yoon et al., 2020). In brief, dissected cerebral cortices were dissociated

and grown in DMEM (Fisher 11960-044) with 1 mM sodium pyruvate (Corning 11360070), 20

mM HEPES (Gibco 15630-080), 100 U/ml PenStrep (Life Technologies 15140122), 10% fetal

bovine serum (FBS, Atlanta 337 Biologicals S11150), and 5 µg/ml insulin. Following 10-12

days in culture, sequential immunopanning was performed to separate astrocyte, microglia, and

oligodendrocyte progenitor cells by agitating at 225 rpm on an orbital shaker (Innova 2000).

After removing microglia via differential adhesion, purified oligodendrocyte progenitors were

plated onto PLL-coated glass coverslips and cultured in differentiation medium (Neurobasal A

media (Life Technologies, 10888022) with N2 (17502048), B27 (17504044), PenStrep, sodium

pyruvate, 2 mM glutamax (35050-061), 40 ng/ml T3, and 5% BSA (Sigma A4503)) for two

days. Treatments with palmitate (PA, 100 µM) NAD^+ (50 µM), NMN (100 µM), EX527 (5 µM)

and/or 78c (3 µM) were added and cultures were allowed to differentiate for 72 hr. Positive

staining for myelin proteins was compared to control levels by immunofluorescence staining and

threshold quantification in ImageJ.

In experiments characterizing astrocyte responses and effects of astrocyte-conditioned

media on oligodendrocytes, astrocytes were plated at a density of 450,000 cells/well on PLL-

coated 6-well plates and grown for 2-3 days in serum-free Neurobasal A media containing 1%

N2, 2% B27, P/S, 1mM sodium pyruvate, 0.45% glucose, 5% BSA, and 50µM β-

mercaptoethanol before treating with or without palmitate (PA, 100 µM) and 78c (3 µM).

Astrocytes were collected for an NAD^+ cycling assay or RNA isolation. Supernatants from

treated astrocytes were collected 24 h post-treatment and concentrated using Vivaspin sample

concentrators (10 and 100 kDa molecular weight cutoffs, Sigma) and stored frozen as 10x

protein concentrate until use for treating OLCs (Liddelow et al., 2017, Yoon et al., 2020).

Conditioned medium from astrocytes (ACM) was brought back to 1x concentration in

oligodendrocyte differentiation medium and placed on oligodendrocytes to differentiate as

described above.

NAD^+ cycling and CD38 activity assays

To determine intracellular NAD^+ levels, one well of a 6-well plate of astrocytes, or

approximately 20 mg of spinal cord tissue, was homogenized in 300 µl of 10% trichloroacetic

acid. Samples were extracted using an organic solvent mixture (1,1,2-Trichloro-1,2,2-

trifluoroethane and triocylamine), diluted, and measured along with a standard curve, as

previously described (Tarrago et al., 2018). Briefly, a cycling reagent (0.76% ethanol, 4 µM

FMN, 27.2 U/ml alcohol dehydrogenase, 0.44 U/ml diaphorase, and 8 µM resazurin) was

prepared in phosphate buffer (100 mM NaH_2PO_4/Na_2HPO_4, pH 8.0) and added to an equal

volume of pre-diluted cell or tissue extract. Plates were measured every 30 seconds for 1 h at

544nm using a SpectraMax plate reader. Raw values were normalized to protein quantification

(Bradford reagent, Biorad) and expressed as nmols/mg protein. For measuring CD38 NAD^+ase

activity, spinal cord tissue was homogenized in NETN buffer (100 mM NaCl, 1 mM EDTA, 20

mM Tris-HCl, and 0.5% Nonidet P-40) supplemented with 5 mM NaF and protease and

phosphatase inhibitors (Tarrago et al., 2018). After centrifugation (12000 rpm, 10 min),

supernatant volumes were adjusted to protein concentrations and reaction mix containing 50 µM

1,N6-Etheno NAD$^+$ in 0.25 M sucrose buffer was added at an equal volume just prior to reading

the plate at 300 excitation and 410 emission every minute for 1 hr on a Spark Magellan plate

reader.

Bioplex assay for cytokines

Astrocyte supernatants from cell treatments described above were used for determination

of secreted cytokine levels using a Bio-Plex Mouse cytokine 23-plex assay (BioRad

m60009rdpd) per manufacturer instructions, as previously described (Langley, Ghosh et al.,

2017). Briefly, post-treatment medium was collected and incubated with primary antibody-

conjugated magnetic beads for 30 min. Wells were washed thrice and then incubated for 30 min

with detection antibodies. After washing, samples were incubated for 10 min with a

streptavidin-phycoerythrin solution, and then washed again prior to resuspension and reading

with a BioPlex 200 (BioRad) reader and Bio-Plex Manager software.

Organotypic cerebellar slice culture and demyelination

Organotypic cerebellar slice cultures were cultured and demyelinated using previously

described methods (Birgbauer, Rao et al., 2004, Hill, Medved et al., 2014, Zhang, Jarjour et al.,

2011). Cerebella were dissected from postnatal day 7-9 pups, embedded in low-melting point

agar (Invitrogen 16520050), and 350 µm slices collected into artificial CSF solution using a

McIlwain tissue chopper (Mickle Laboratory Engineering Co; Guildford, UK). Four slices were

transferred and grown per each Millicell insert (Sigma PIHP03050) in 50% MEM, 22.5% HBSS,

24% HI Horse Serum, 1.5% Glucose, 1% GlutaMax, 1% P/S medium.

Demyelination was induced after 7 d in culture by treating with 0.5 mg/ml

lysophosphatidycholine (LL, Sigma) for 18 h. Organotypic slices were placed back in LL-free

medium and collected for immunohistochemistry and confocal imaging after 24 h

(demyelination) or 7 d (myelin regeneration) with or without addition of PA (100 μM) and/or

78c (3 μM) added to the growth media. Confocal images were acquired at 20x magnification

from 5 regions randomly selected per slice using a Zeiss LSM 780 and Zen Software in the

Mayo Clinic Microscopy and Cell Analysis Core. Images were thresholded for each stain, and

immunoreactivity or counts were quantified in ImageJ.

Statistical analysis

Data were graphed with bars representing mean ± SEM with sample size for each

experiment indicated in the figure legends. To determine sample size, power analysis using the

fpower macro in SAS was used to determine animal numbers for each endpoint based on 80%

power, $\alpha=0.05$, and 4 treatment groups. Differences and SD used to determine δ were used from

analysis of previous similarly-designed studies in our laboratory.   No power analysis or other

statistical method was used to compute sample size for cell culture experiments. Cell culture

studies were repeated twice and each biological replicate per group was isolated from different

pups. Assays were run in technical duplicate or triplicate.

No statistical outliers were removed, but the ROUT test in GraphPad Prism was used to

check for outliers. Comparisons between two groups were analyzed by Student's t-test, whereas

one-way or two-way ANOVAs with Bonferroni post-test were applied for multiple-comparisons.

Analysis was performed in GraphPad Prism 7.0 software with p-values ≤ 0.05 considered

significant.

Acknowledgements

78c-containing chow for these experiments was generously provided by Calico Life

Sciences LLC. The diagrams within figures were created using elements from the Biomedical-

PPT-Toolkit-Suite (Motifolio).

Author Contributions

IAS and MRL conceived and designed the study. MRL, CC, and TRPM conducted *in*

vivo experiments and necropsies. Embedding and histological staining was performed by MRL

and LK. Image acquisition and quantifications were performed by MRL under guidance from

CC. MRL completed *in vitro* assays and qRT-PCR. WS and HY assisted with glial preparations

and organotypic slice cultures. TRP, CCSC, and ENC provided expertise in NAD$^+$ quantification

and experimental design as well as CD38 mice. MRL and IAS wrote the manuscript and all

authors reviewed and edited the manuscript.

Conflict of Interest

ENC holds a patent on the use of CD38 inhibitors for metabolic diseases. The other authors

have no conflicts of interest to disclose.

References

References

Aksoy P, Escande C, White TA, Thompson M, Soares S, Benech JC, Chini EN (2006) Regulation

of SIRT 1 mediated NAD dependent deacetylation: a novel role for the multifunctional

enzyme CD38. *Biochem Biophys Res Commun* 349: 353-9

Aksoy P, White TA, Thompson M, Chini EN (2006) Regulation of intracellular levels of NAD: a

novel role for CD38. *Biochem Biophys Res Commun* 345: 1386-92

Alizadeh A, Dyck SM, Karimi-Abdolrezaee S (2015) Myelin damage and repair in pathologic

CNS: challenges and prospects. *Front Mol Neurosci* 8: 35

Alizadeh A, Karimi-Abdolrezaee S (2016) Microenvironmental regulation of oligodendrocyte

replacement and remyelination in spinal cord injury. *J Physiol* 594: 3539-52

Amici SA, Young NA, Narvaez-Miranda J, Jablonski KA, Arcos J, Rosas L, Papenfuss TL,

Torrelles JB, Jarjour WN, Guerau-de-Arellano M (2018) CD38 Is Robustly Induced in

Human Macrophages and Monocytes in Inflammatory Conditions. *Front Immunol* 9: 1593

Barbosa MT, Soares SM, Novak CM, Sinclair D, Levine JA, Aksoy P, Chini EN (2007) The

enzyme CD38 (a NAD glycohydrolase, EC 3.2.2.5) is necessary for the development of

diet-induced obesity. *FASEB J* 21: 3629-39

Birgbauer E, Rao TS, Webb M (2004) Lysolecithin induces demyelination in vitro in a cerebellar

slice culture system. *J Neurosci Res* 78: 157-66

Blacher E, Dadali T, Bespalko A, Haupenthal VJ, Grimm MO, Hartmann T, Lund FE, Stein R,

Levy A (2015) Alzheimer's disease pathology is attenuated in a CD38-deficient mouse

model. *Ann Neurol* 78: 88-103

Boslett J, Helal M, Chini E, Zweier JL (2018) Genetic deletion of CD38 confers post-ischemic

myocardial protection through preserved pyridine nucleotides. *J Mol Cell Cardiol* 118: 81-

94

Boven LA, Montagne L, Nottet HS, De Groot CJ (2000) Macrophage inflammatory protein-1alpha

(MIP-1alpha), MIP-1beta, and RANTES mRNA semiquantification and protein expression

in active demyelinating multiple sclerosis (MS) lesions. *Clin Exp Immunol* 122: 257-63

Bruck W, Popescu B, Lucchinetti CF, Markovic-Plese S, Gold R, Thal DR, Metz I (2012)

Neuromyelitis optica lesions may inform multiple sclerosis heterogeneity debate. *Ann*

Neurol 72: 385-94

Bruzzone S, Verderio C, Schenk U, Fedele E, Zocchi E, Matteoli M, De Flora A (2004) Glutamate-

mediated overexpression of CD38 in astrocytes cultured with neurones. *J Neurochem* 89:

264-72

Buckman LB, Thompson MM, Lippert RN, Blackwell TS, Yull FE, Ellacott KL (2015) Evidence

for a novel functional role of astrocytes in the acute homeostatic response to high-fat diet

intake in mice. *Mol Metab* 4: 58-63

Buckman LB, Thompson MM, Moreno HN, Ellacott KL (2013) Regional astrogliosis in the mouse

hypothalamus in response to obesity. *J Comp Neurol* 521: 1322-33

Camacho-Pereira J, Tarrago MG, Chini CCS, Nin V, Escande C, Warner GM, Puranik AS, Schoon

RA, Reid JM, Galina A, Chini EN (2016) CD38 Dictates Age-Related NAD Decline and

Mitochondrial Dysfunction through an SIRT3-Dependent Mechanism. *Cell Metab* 23:

1127-1139

Camargo N, Goudriaan A, van Deijk AF, Otte WM, Brouwers JF, Lodder H, Gutmann DH, Nave

KA, Dijkhuizen RM, Mansvelder HD, Chrast R, Smit AB, Verheijen MHG (2017)

Oligodendroglial myelination requires astrocyte-derived lipids. *PLoS Biol* 15: e1002605

Canto C, Menzies KJ, Auwerx J (2015) NAD(+) Metabolism and the Control of Energy

Homeostasis: A Balancing Act between Mitochondria and the Nucleus. *Cell Metab* 22: 31-

53

Chiang SH, Harrington WW, Luo G, Milliken NO, Ulrich JC, Chen J, Rajpal DK, Qian Y,

Carpenter T, Murray R, Geske RS, Stimpson SA, Kramer HF, Haffner CD, Becherer JD,

Preugschat F, Billin AN (2015) Genetic Ablation of CD38 Protects against Western Diet-

Induced Exercise Intolerance and Metabolic Inflexibility. *PLoS One* 10: e0134927

Chini C, Hogan KA, Warner GM, Tarrago MG, Peclat TR, Tchkonia T, Kirkland JL, Chini E

(2019) The NADase CD38 is induced by factors secreted from senescent cells providing a

potential link between senescence and age-related cellular NAD(+) decline. *Biochem*

Biophys Res Commun 513: 486-493

Chini EN, Chini CCS, Espindola Netto JM, de Oliveira GC, van Schooten W (2018) The

Pharmacology of CD38/NADase: An Emerging Target in Cancer and Diseases of Aging.

Trends Pharmacol Sci 39: 424-436

Choi JW, Gardell SE, Herr DR, Rivera R, Lee CW, Noguchi K, Teo ST, Yung YC, Lu M,

Kennedy G, Chun J (2011) FTY720 (fingolimod) efficacy in an animal model of multiple

sclerosis requires astrocyte sphingosine 1-phosphate receptor 1 (S1P1) modulation. *Proc*

Natl Acad Sci U S A 108: 751-6

Clarke LE, Liddelow SA, Chakraborty C, Munch AE, Heiman M, Barres BA (2018) Normal aging

induces A1-like astrocyte reactivity. *Proc Natl Acad Sci U S A* 115: E1896-E1905

Clarner T, Janssen K, Nellessen L, Stangel M, Skripuletz T, Krauspe B, Hess FM, Denecke B,

Beutner C, Linnartz-Gerlach B, Neumann H, Vallieres L, Amor S, Ohl K, Tenbrock K,

Beyer C, Kipp M (2015) CXCL10 triggers early microglial activation in the cuprizone

model. *J Immunol* 194: 3400-13

Deckert J, Wetzel MC, Bartle LM, Skaletskaya A, Goldmacher VS, Vallee F, Zhou-Liu Q, Ferrari

P, Pouzieux S, Lahoute C, Dumontet C, Plesa A, Chiron M, Lejeune P, Chittenden T, Park

PU, Blanc V (2014) SAR650984, a novel humanized CD38-targeting antibody,

demonstrates potent antitumor activity in models of multiple myeloma and other CD38+

hematologic malignancies. *Clin Cancer Res* 20: 4574-83

Denver P, Gault VA, McClean PL (2018) Sustained high-fat diet modulates inflammation, insulin

signalling and cognition in mice and a modified xenin peptide ameliorates neuropathology

in a chronic high-fat model. *Diabetes Obes Metab* 20: 1166-1175

Dimas P, Montani L, Pereira JA, Moreno D, Trotzmuller M, Gerber J, Semenkovich CF, Kofeler

HC, Suter U (2019) CNS myelination and remyelination depend on fatty acid synthesis by

oligodendrocytes. *Elife* 8

Domingues HS, Portugal CC, Socodato R, Relvas JB (2016) Oligodendrocyte, Astrocyte, and

Microglia Crosstalk in Myelin Development, Damage, and Repair. *Front Cell Dev Biol* 4:

71

Escande C, Chini CC, Nin V, Dykhouse KM, Novak CM, Levine J, van Deursen J, Gores GJ, Chen

J, Lou Z, Chini EN (2010) Deleted in breast cancer-1 regulates SIRT1 activity and

contributes to high-fat diet-induced liver steatosis in mice. *J Clin Invest* 120: 545-58

Escande C, Nin V, Price NL, Capellini V, Gomes AP, Barbosa MT, O'Neil L, White TA, Sinclair

DA, Chini EN (2013) Flavonoid apigenin is an inhibitor of the NAD+ ase CD38:

implications for cellular NAD+ metabolism, protein acetylation, and treatment of metabolic

syndrome. *Diabetes* 62: 1084-93

Foolad F, Khodagholi F, Javan M (2019) Sirtuins in Multiple Sclerosis: The crossroad of

neurodegeneration, autoimmunity and metabolism. *Mult Scler Relat Disord* 34: 47-58

Freeman LR, Haley-Zitlin V, Rosenberger DS, Granholm AC (2014) Damaging effects of a high-

fat diet to the brain and cognition: a review of proposed mechanisms. *Nutr Neurosci* 17:

241-51

Gianfrancesco MA, Acuna B, Shen L, Briggs FB, Quach H, Bellesis KH, Bernstein A, Hedstrom

AK, Kockum I, Alfredsson L, Olsson T, Schaefer C, Barcellos LF (2014) Obesity during

childhood and adolescence increases susceptibility to multiple sclerosis after accounting for

established genetic and environmental risk factors. *Obes Res Clin Pract* 8: e435-47

Ginwala R, McTish E, Raman C, Singh N, Nagarkatti M, Nagarkatti P, Sagar D, Jain P, Khan ZK

(2016) Apigenin, a Natural Flavonoid, Attenuates EAE Severity Through the Modulation of

Dendritic Cell and Other Immune Cell Functions. *J Neuroimmune Pharmacol* 11: 36-47

Graham LC, Grabowska WA, Chun Y, Risacher SL, Philip VM, Saykin AJ, Alzheimer's Disease

Neuroimaging I, Sukoff Rizzo SJ, Howell GR (2019) Exercise prevents obesity-induced

cognitive decline and white matter damage in mice. *Neurobiol Aging* 80: 154-172

Haffner CD, Becherer JD, Boros EE, Cadilla R, Carpenter T, Cowan D, Deaton DN, Guo Y,

Harrington W, Henke BR, Jeune MR, Kaldor I, Milliken N, Petrov KG, Preugschat F,

Schulte C, Shearer BG, Shearer T, Smalley TL, Jr., Stewart EL et al. (2015) Discovery,

Synthesis, and Biological Evaluation of Thiazoloquin(az)olin(on)es as Potent CD38

Inhibitors. *J Med Chem* 58: 3548-71

Haider L, Fischer MT, Frischer JM, Bauer J, Hoftberger R, Botond G, Esterbauer H, Binder CJ,

Witztum JL, Lassmann H (2011) Oxidative damage in multiple sclerosis lesions. *Brain* 134:

1914-24

Harlan BA, Pehar M, Sharma DR, Beeson G, Beeson CC, Vargas MR (2016) Enhancing NAD+

Salvage Pathway Reverts the Toxicity of Primary Astrocytes Expressing Amyotrophic

Lateral Sclerosis-linked Mutant Superoxide Dismutase 1 (SOD1). *J Biol Chem* 291: 10836-

46

Hasan M, Seo JE, Rahaman KA, Min H, Kim KH, Park JH, Sung C, Son J, Kang MJ, Jung BH,

Park WS, Kwon OS (2017) Novel genes in brain tissues of EAE-induced normal and obese

mice: Upregulation of metal ion-binding protein genes in obese-EAE mice. *Neuroscience*

343: 322-336

Hathorn T, Snyder-Keller A, Messer A (2011) Nicotinamide improves motor deficits and

upregulates PGC-1alpha and BDNF gene expression in a mouse model of Huntington's

disease. *Neurobiol Dis* 41: 43-50

Hattori T, Kaji M, Ishii H, Jureepon R, Takarada-Iemata M, Minh Ta H, Manh Le T, Konno A,

Hirai H, Shiraishi Y, Ozaki N, Yamamoto Y, Okamoto H, Yokoyama S, Higashida H,

Kitao Y, Hori O (2017) CD38 positively regulates postnatal development of astrocytes cell-

autonomously and oligodendrocytes non-cell-autonomously. *Glia* 65: 974-989

Herrmann MM, Barth S, Greve B, Schumann KM, Bartels A, Weissert R (2016) Identification of

gene expression patterns crucially involved in experimental autoimmune encephalomyelitis

and multiple sclerosis. *Dis Model Mech* 9: 1211-1220

Hill RA, Medved J, Patel KD, Nishiyama A (2014) Organotypic slice cultures to study

oligodendrocyte dynamics and myelination. *J Vis Exp*: e51835

Hoffmann FS, Hofereiter J, Rubsamen H, Melms J, Schwarz S, Faber H, Weber P, Putz B, Loleit

V, Weber F, Hohlfeld R, Meinl E, Krumbholz M (2015) Fingolimod induces

neuroprotective factors in human astrocytes. *J Neuroinflammation* 12: 184

Iqbal J, Zaidi M (2007) CD38 is required for priming by TNF-alpha: a mechanism for extracellular

coordination of cell fate. *Am J Physiol Renal Physiol* 292: F1283-90

Jablonska B, Gierdalski M, Chew LJ, Hawley T, Catron M, Lichauco A, Cabrera-Luque J, Yuen T,

Rowitch D, Gallo V (2016) Sirt1 regulates glial progenitor proliferation and regeneration in

white matter after neonatal brain injury. *Nat Commun* 7: 13866

Janssen K, Rickert M, Clarner T, Beyer C, Kipp M (2016) Absence of CCL2 and CCL3

Ameliorates Central Nervous System Grey Matter But Not White Matter Demyelination in

the Presence of an Intact Blood-Brain Barrier. *Mol Neurobiol* 53: 1551-1564

Ji S, Doucette JR, Nazarali AJ (2011) Sirt2 is a novel in vivo downstream target of Nkx2.2 and

enhances oligodendroglial cell differentiation. *J Mol Cell Biol* 3: 351-9

Ji Z, Wu S, Xu Y, Qi J, Su X, Shen L (2019) Obesity Promotes EAE Through IL-6 and CCL-2-

Mediated T Cells Infiltration. *Front Immunol* 10: 1881

Karim H, Kim SH, Lapato AS, Yasui N, Katzenellenbogen JA, Tiwari-Woodruff SK (2018)

Increase in chemokine CXCL1 by ERbeta ligand treatment is a key mediator in promoting

axon myelination. *Proc Natl Acad Sci U S A* 115: 6291-6296

Kiray H, Lindsay SL, Hosseinzadeh S, Barnett SC (2016) The multifaceted role of astrocytes in

regulating myelination. *Exp Neurol* 283: 541-9

Kou W, Banerjee S, Eudy J, Smith LM, Persidsky R, Borgmann K, Wu L, Sakhuja N, Deshpande

MS, Walseth TF, Ghorpade A (2009) CD38 regulation in activated astrocytes: implications

for neuroinflammation and HIV-1 brain infection. *J Neurosci Res* 87: 2326-39

Langley M, Ghosh A, Charli A, Sarkar S, Ay M, Luo J, Zielonka J, Brenza T, Bennett B, Jin H,

Ghaisas S, Schlichtmann B, Kim D, Anantharam V, Kanthasamy A, Narasimhan B,

Kalyanaraman B, Kanthasamy AG (2017) Mito-Apocynin Prevents Mitochondrial

Dysfunction, Microglial Activation, Oxidative Damage, and Progressive

Neurodegeneration in MitoPark Transgenic Mice. *Antioxid Redox Signal* 27: 1048-1066

Langley MR, Yoon H, Kim HN, Choi CI, Simon W, Kleppe L, Lanza IR, LeBrasseur NK,

Matveyenko A, Scarisbrick IA (2019) High fat diet consumption results in mitochondrial

dysfunction, oxidative stress, and oligodendrocyte loss in the central nervous system.

Biochim Biophys Acta Mol Basis Dis: 165630

Lassmann H, van Horssen J (2016) Oxidative stress and its impact on neurons and glia in multiple

sclerosis lesions. *Biochim Biophys Acta* 1862: 506-10

Li J, Dou X, Li S, Zhang X, Zeng Y, Song Z (2015) Nicotinamide ameliorates palmitate-induced

ER stress in hepatocytes via cAMP/PKA/CREB pathway-dependent Sirt1 upregulation.

Biochim Biophys Acta 1853: 2929-36

Liao PC, Lai MH, Hsu KP, Kuo YH, Chen J, Tsai MC, Li CX, Yin XJ, Jeyashoke N, Chao LK

(2018) Identification of beta-Sitosterol as in Vitro Anti-Inflammatory Constituent in

Moringa oleifera. *J Agric Food Chem* 66: 10748-10759

Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, Bennett ML,

Munch AE, Chung WS, Peterson TC, Wilton DK, Frouin A, Napier BA, Panicker N,

Kumar M, Buckwalter MS, Rowitch DH, Dawson VL, Dawson TM, Stevens B et al. (2017)

Neurotoxic reactive astrocytes are induced by activated microglia. *Nature* 541: 481-487

Lokhorst HM, Plesner T, Laubach JP, Nahi H, Gimsing P, Hansson M, Minnema MC, Lassen U,

Krejcik J, Palumbo A, van de Donk NW, Ahmadi T, Khan I, Uhlar CM, Wang J, Sasser

AK, Losic N, Lisby S, Basse L, Brun N et al. (2015) Targeting CD38 with Daratumumab Monotherapy in Multiple Myeloma. *N Engl J Med* 373: 1207-19

Long A, Klimova N, Kristian T (2017) Mitochondrial NUDIX hydrolases: A metabolic link between NAD catabolism, GTP and mitochondrial dynamics. *Neurochem Int* 109: 193-201

Machado-Santos J, Saji E, Troscher AR, Paunovic M, Liblau R, Gabriely G, Bien CG, Bauer J, Lassmann H (2018) The compartmentalized inflammatory response in the multiple sclerosis brain is composed of tissue-resident CD8+ T lymphocytes and B cells. *Brain* 141: 2066-2082

Mamik MK, Banerjee S, Walseth TF, Hirte R, Tang L, Borgmann K, Ghorpade A (2011) HIV-1 and IL-1beta regulate astrocytic CD38 through mitogen-activated protein kinases and nuclear factor-kappaB signaling mechanisms. *J Neuroinflammation* 8: 145

Manouchehrinia A, Hedstrom AK, Alfredsson L, Olsson T, Hillert J, Ramanujam R (2018) Association of Pre-Disease Body Mass Index With Multiple Sclerosis Prognosis. *Front Neurol* 9: 232

Marrie RA, Rudick R, Horwitz R, Cutter G, Tyry T, Campagnolo D, Vollmer T (2010) Vascular comorbidity is associated with more rapid disability progression in multiple sclerosis. *Neurology* 74: 1041-7

Martin T, Baz R, Benson DM, Lendvai N, Wolf J, Munster P, Lesokhin AM, Wack C, Charpentier E, Campana F, Vij R (2017) A phase 1b study of isatuximab plus lenalidomide and dexamethasone for relapsed/refractory multiple myeloma. *Blood* 129: 3294-3303

Matalonga J, Glaria E, Bresque M, Escande C, Carbo JM, Kiefer K, Vicente R, Leon TE, Beceiro S, Pascual-Garcia M, Serret J, Sanjurjo L, Moron-Ros S, Riera A, Paytubi S, Juarez A, Sotillo F, Lindbom L, Caelles C, Sarrias MR et al. (2017) The Nuclear Receptor LXR

Limits Bacterial Infection of Host Macrophages through a Mechanism that Impacts Cellular

NAD Metabolism. *Cell Rep* 18: 1241-1255

Miller ED, Dziedzic A, Saluk-Bijak J, Bijak M (2019) A Review of Various Antioxidant

Compounds and their Potential Utility as Complementary Therapy in Multiple Sclerosis.

Nutrients 11

Mizunoya W, Ohnuki K, Baba K, Miyahara H, Shimizu N, Tabata K, Kino T, Sato Y, Tatsumi R,

Ikeuchi Y (2013) Effect of dietary fat type on anxiety-like and depression-like behavior in

mice. *Springerplus* 2: 165

Mokry LE, Ross S, Timpson NJ, Sawcer S, Davey Smith G, Richards JB (2016) Obesity and

Multiple Sclerosis: A Mendelian Randomization Study. *PLoS Med* 13: e1002053

Mori F, Nistico R, Nicoletti CG, Zagaglia S, Mandolesi G, Piccinin S, Martino G, Finardi A,

Rossini PM, Marfia GA, Furlan R, Centonze D (2016) RANTES correlates with

inflammatory activity and synaptic excitability in multiple sclerosis. *Mult Scler* 22: 1405-

1412

Moyon S, Dubessy AL, Aigrot MS, Trotter M, Huang JK, Dauphinot L, Potier MC, Kerninon C,

Melik Parsadaniantz S, Franklin RJ, Lubetzki C (2015) Demyelination causes adult CNS

progenitors to revert to an immature state and express immune cues that support their

migration. *J Neurosci* 35: 4-20

Negrotto L, Farez MF, Correale J (2016) Immunologic Effects of Metformin and Pioglitazone

Treatment on Metabolic Syndrome and Multiple Sclerosis. *JAMA Neurol* 73: 520-8

Ogrodnik M, Zhu Y, Langhi LGP, Tchkonia T, Kruger P, Fielder E, Victorelli S, Ruswhandi RA,

Giorgadze N, Pirtskhalava T, Podgorni O, Enikolopov G, Johnson KO, Xu M, Inman C,

Palmer AK, Schafer M, Weigl M, Ikeno Y, Burns TC et al. (2019) Obesity-Induced

Cellular Senescence Drives Anxiety and Impairs Neurogenesis. *Cell Metab*

Ohtomo R, Iwata A, Arai K (2018) Molecular Mechanisms of Oligodendrocyte Regeneration in

White Matter-Related Diseases. *Int J Mol Sci* 19

Olsson T, Barcellos LF, Alfredsson L (2017) Interactions between genetic, lifestyle and

environmental risk factors for multiple sclerosis. *Nat Rev Neurol* 13: 25-36

Partida-Sanchez S, Cockayne DA, Monard S, Jacobson EL, Oppenheimer N, Garvy B, Kusser K,

Goodrich S, Howard M, Harmsen A, Randall TD, Lund FE (2001) Cyclic ADP-ribose

production by CD38 regulates intracellular calcium release, extracellular calcium influx and

chemotaxis in neutrophils and is required for bacterial clearance in vivo. *Nat Med* 7: 1209-

16

Pavelek Z, Vysata O, Sobisek L, Klimova B, Andrys C, Vokurkova D, Mazurova R, Stourac P,

Valis M (2018) Lymphocyte populations and their change during five-year glatiramer

acetate treatment. *Neurol Neurochir Pol* 52: 587-592

Penberthy WT, Tsunoda I (2009) The importance of NAD in multiple sclerosis. *Curr Pharm Des*

15: 64-99

Peng QY, Wang YM, Chen CX, Zou Y, Zhang LN, Deng SY, Ai YH (2018) Inhibiting the

CD38/cADPR pathway protected rats against sepsis associated brain injury. *Brain Res*

1678: 56-63

Penke M, Schuster S, Gorski T, Gebhardt R, Kiess W, Garten A (2017) Oleate ameliorates

palmitate-induced reduction of NAMPT activity and NAD levels in primary human

hepatocytes and hepatocarcinoma cells. *Lipids Health Dis* 16: 191

Ponath G, Park C, Pitt D (2018) The Role of Astrocytes in Multiple Sclerosis. *Front Immunol* 9:

217

Ponath G, Ramanan S, Mubarak M, Housley W, Lee S, Sahinkaya FR, Vortmeyer A, Raine CS,

Pitt D (2017) Myelin phagocytosis by astrocytes after myelin damage promotes lesion

pathology. *Brain* 140: 399-413

Prola A, Pires Da Silva J, Guilbert A, Lecru L, Piquereau J, Ribeiro M, Mateo P, Gressette M,

Fortin D, Boursier C, Gallerne C, Caillard A, Samuel JL, Francois H, Sinclair DA, Eid P,

Ventura-Clapier R, Garnier A, Lemaire C (2017) SIRT1 protects the heart from ER stress-

induced cell death through eIF2alpha deacetylation. *Cell Death Differ* 24: 343-356

Roboon J, Hattori T, Ishii H, Takarada-Iemata M, Le TM, Shiraishi Y, Ozaki N, Yamamoto Y,

Sugawara A, Okamoto H, Higashida H, Kitao Y, Hori O (2019) Deletion of CD38

Suppresses Glial Activation and Neuroinflammation in a Mouse Model of Demyelination.

Front Cell Neurosci 13: 258

Sadeghi A, Seyyed Ebrahimi SS, Golestani A, Meshkani R (2017) Resveratrol Ameliorates

Palmitate-Induced Inflammation in Skeletal Muscle Cells by Attenuating Oxidative Stress

and JNK/NF-kappaB Pathway in a SIRT1-Independent Mechanism. *J Cell Biochem* 118:

2654-2663

Sarkar S, Malovic E, Harishchandra DS, Ghaisas S, Panicker N, Charli A, Palanisamy BN, Rokad

D, Jin H, Anantharam V, Kanthasamy A, Kanthasamy AG (2017) Mitochondrial

impairment in microglia amplifies NLRP3 inflammasome proinflammatory signaling in cell

culture and animal models of Parkinson's disease. *NPJ Parkinsons Dis* 3: 30

Stein LR, Imai S (2014) Specific ablation of Nampt in adult neural stem cells recapitulates their

functional defects during aging. *EMBO J* 33: 1321-40

Tarrago MG, Chini CCS, Kanamori KS, Warner GM, Caride A, de Oliveira GC, Rud M, Samani

A, Hein KZ, Huang R, Jurk D, Cho DS, Boslett JJ, Miller JD, Zweier JL, Passos JF, Doles

JD, Becherer DJ, Chini EN (2018) A Potent and Specific CD38 Inhibitor Ameliorates Age-

Related Metabolic Dysfunction by Reversing Tissue NAD(+) Decline. *Cell Metab* 27:

1081-1095 e10

Tejera-Alhambra M, Casrouge A, de Andres C, Seyfferth A, Ramos-Medina R, Alonso B, Vega J,

Fernandez-Paredes L, Albert ML, Sanchez-Ramon S (2015) Plasma biomarkers

discriminate clinical forms of multiple sclerosis. *PLoS One* 10: e0128952

Tettey P, Simpson S, Jr., Taylor B, Blizzard L, Ponsonby AL, Dwyer T, Kostner K, van der Mei I

(2014) An adverse lipid profile is associated with disability and progression in disability, in

people with MS. *Mult Scler* 20: 1737-44

Thaler JP, Yi CX, Schur EA, Guyenet SJ, Hwang BH, Dietrich MO, Zhao X, Sarruf DA, Izgur V,

Maravilla KR, Nguyen HT, Fischer JD, Matsen ME, Wisse BE, Morton GJ, Horvath TL,

Baskin DG, Tschop MH, Schwartz MW (2012) Obesity is associated with hypothalamic

injury in rodents and humans. *J Clin Invest* 122: 153-62

Timmermans S, Bogie JF, Vanmierlo T, Lutjohann D, Stinissen P, Hellings N, Hendriks JJ (2014)

High fat diet exacerbates neuroinflammation in an animal model of multiple sclerosis by

activation of the Renin Angiotensin system. *J Neuroimmune Pharmacol* 9: 209-17

Valentin-Torres A, Savarin C, Hinton DR, Phares TW, Bergmann CC, Stohlman SA (2016)

Sustained TNF production by central nervous system infiltrating macrophages promotes

progressive autoimmune encephalomyelitis. *J Neuroinflammation* 13: 46

van de Donk N, Richardson PG, Malavasi F (2018) CD38 antibodies in multiple myeloma: back to

the future. *Blood* 131: 13-29

Wang C, Zhang Y, Ding J, Zhao Z, Qian C, Luan Y, Teng GJ (2017) Nicotinamide Administration

Improves Remyelination after Stroke. *Neural Plast* 2017: 7019803

Wang J, Zhao C, Kong P, Sun H, Sun Z, Bian G, Sun Y, Guo L (2016) Treatment with NAD(+)

inhibited experimental autoimmune encephalomyelitis by activating AMPK/SIRT1

signaling pathway and modulating Th1/Th17 immune responses in mice. *Int*

Immunopharmacol 39: 287-94

Wang YM, Liu ZY, Ai YH, Zhang LN, Zou Y, Peng QY (2017) Blocking the CD38/cADPR

pathway plays a double-edged role in LPS stimulated microglia. *Neuroscience* 361: 34-42

Wanrooy BJ, Kumar KP, Wen SW, Qin CX, Ritchie RH, Wong CHY (2018) Distinct contributions

of hyperglycemia and high-fat feeding in metabolic syndrome-induced neuroinflammation.

J Neuroinflammation 15: 293

Wilhelm F, Hirrlinger J (2012) Multifunctional roles of NAD(+) and NADH in astrocytes.

Neurochem Res 37: 2317-25

Wilms H, Sievers J, Rickert U, Rostami-Yazdi M, Mrowietz U, Lucius R (2010) Dimethylfumarate

inhibits microglial and astrocytic inflammation by suppressing the synthesis of nitric oxide,

IL-1beta, TNF-alpha and IL-6 in an in-vitro model of brain inflammation. *J*

Neuroinflammation 7: 30

Yoon H, Choi CI, Triplet EM, Langley MR, Kleppe LS, Kim HN, Simon WL, Scarisbrick IA

(2020) Blocking the Thrombin Receptor Promotes Repair of Demyelinated Lesions in the

Adult Brain. *J Neurosci* 40: 1483-1500

Yoon H, Kleven A, Paulsen A, Kleppe L, Wu J, Ying Z, Gomez-Pinilla F, Scarisbrick IA (2016)

Interplay between exercise and dietary fat modulates myelinogenesis in the central nervous

system. *Biochim Biophys Acta* 1862: 545-55

Zhang H, Jarjour AA, Boyd A, Williams A (2011) Central nervous system remyelination in

culture--a tool for multiple sclerosis research. *Exp Neurol* 230: 138-48

Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, Phatnani HP, Guarnieri P,

Caneda C, Ruderisch N, Deng S, Liddelow SA, Zhang C, Daneman R, Maniatis T, Barres

BA, Wu JQ (2014) An RNA-sequencing transcriptome and splicing database of glia,

neurons, and vascular cells of the cerebral cortex. *J Neurosci* 34: 11929-47

Zhang Y, Sloan SA, Clarke LE, Caneda C, Plaza CA, Blumenthal PD, Vogel H, Steinberg GK,

Edwards MS, Li G, Duncan JA, 3rd, Cheshier SH, Shuer LM, Chang EF, Grant GA,

Gephart MG, Barres BA (2016) Purification and Characterization of Progenitor and Mature

Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse. *Neuron*

89: 37-53

Zou Y, He X, Peng QY, Guo QL (2017) Inhibition of CD38/Cyclic ADP-ribose Pathway Protects

Rats against Ropivacaine-induced Convulsion. *Chin Med J (Engl)* 130: 2354-2360

Figures and Figure Legends

<u>**Figure 1.**</u> **CD38 expression in astrocytes of chronic HFD-fed mice and in a model of**

demyelination.

A Schematic representation of previously shown direct effects of high fat diet (HFD) on

oligodendrocyte lineage cells including apoptosis and impaired differentiation with potential

astrocyte-mediated indirect effects, represented by dotted lines, that comprise the hypothesis to

be tested.

B Spinal cord tissue of regular diet (RD) and high fat diet (HFD) mice showing a depletion of

NAD^+ after 12 wk (n=5 mice/group) as determined by cycling assay.

C Increased RNA expression of CD38 (n=4), assessed by qPCR, in spinal cord tissue of HFD-

fed mice when compared to RD-fed mice.

D- G Immunofluorescence staining with double-labeling (*F, G*) and corresponding quantification

(*D, E)* indicates increased CD38 (green) immunoreactivity in astrocytes (GFAP, red) within the

spinal cords of HFD-fed mice (*D, F,* n=7) or in the lysolecithin (LL) model of experimental

demyelination (*E, G,* n=4)versus the uninjected (UI) side. Broken white line (*G*) designates

lesion boundary. Scale bar = 50 μm (*F*) and 100μm (*G*).

H-J High fat consumption was modeled in primary murine astrocyte cultures by including

saturated fat palmitate in the media (PA, 100 μM, 24 h). Increases in RNA expression of GFAP

(*H,* n=4) and CD38 (**I,** n=6) were observed in PA-containing astrocyte cultures. *J*, Depleted

NAD^+ levels were found in PA-treated astrocytes when assessed using an NAD^+ cycling assay,

n=7-8. Bar graphs represent the mean +/- SEM. Asterisks indicate levels of statistical

significance as determined by Student's t-test, where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

1015

1016   The following figure supplement is available for figure1:

1017   **Figure 1- Figure Supplement 1**. CD38 RNA expression in mouse spinal cord tissue.

1018   **Figure 1- Source Data 1.**

1019

1020

Figure 2. CD38-catalytically inactive (CD38ci) mice are resistant to HFD-induced

oligodendrocyte loss.

Histological analysis from wildtype (WT) and CD38-catalytically inactive mice (CD38ci) fed a

regular diet (RD) or high fat diet (HFD) for 30 wk beginning at 50 wk of age. Red and blue

outlined regions depict regions imaged and quantified, including dorsal column and ventral white

matter (broken gray line) and ventral grey matter (broken black line).

A Schematic depicting experimental design for diets administration.

B-C Olig2+ oligodendrocyte lineage cells (**B**) and GST3+ mature oligodendrocytes (**C**) were

significantly depleted in HFD-fed WT mice, yet CD38ci mice fed a HFD had no reduction in

oligodendrocyte numbers. When compared to WT-HFD mice, CD38ci-HFD mice had

significantly more oligodendrocytes remaining in the dorsal column (representative images

pictured; ###, $p < 0.001$). Scale bar = 50 µm.

D Representative images of immunohistochemical staining of the dorsal column and

quantification of the dorsal column and ventral grey matter and white matter of the spinal cords

confirming no significant changes in MBP immunoreactivity across genotype or diet groups.

Scale bar= 100 µm.

Bar graphs represent mean +/- SEM. Two-way ANOVA followed by Bonferroni post-hoc test

(n=7-10 mice/group). Asterisks indicate levels of statistical significance when compared to

control, where *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

1040

1041   The following figure supplement is available for figure 2:

1042   **Figure 2- Figure Supplement 1**. Effect of direct CD38 inhibition on oligodendrocyte

1043   progenitors.

1044   **Figure 2- Source Data 1.**

1045

1048

Figure 3. Diminished neuroinflammatory responses in CD38 catalytically inactive (CD38ci)

mice following chronic HFD.

Histological analysis from wildtype (WT) and CD38-catalytically inactive mice (CD38ci) fed a

regular diet (RD) or high fat diet (HFD) for 30 wk beginning at 50 wk of age. Scale bar= 50 µm.

and n=7-10 mice/group.

A Representative images of the ventral white matter and immunohistochemical analysis of spinal

cord regions demonstrates decreased microgliosis (IBA1) in CD38ci mice. When compared to

WT-HFD mice, CD38ci-HFD mice had significantly less IBA1+ immunoreactivity in the spinal

cord ventral gray and white matter.

B Representative images of the ventral white matter and corresponding quantification confirm

reduced astrogliosis (GFAP) in the spinal cords of CD38 catalytically inactive (CD38ci) mice

overall when compared to wildtype (WT) mice. Decreased GFAP+ immunoreactivity in the

ventral white matter when compared to HFD mice.

C Representative images of the dorsal column and quantification of the spinal cords confirming

significant changes in 4HNE immunoreactivity, an oxidative stress marker, between genotypes

for HFD mice.

D NAD^+ levels as determined by cycling assay indicate reductions in WT-HFD mice compared

to WT-RD and increased NAD^+ levels in the CD38ci mice versus WT.

E CD38 activity levels based on etheno-NAD^+ assay show impaired activity in CD38ci mice

versus WTs and higher activity in WT-HFD mice than WT-RD (n=5/group).

F-H qRT-PCRs data indicting differences in antioxidant (F) and pro-repair astrocyte marker

genes (G-H, BDNF and S100a10, respectively) in the CD38ci mice versus controls.

Bar graphs represent mean +/- SEM. Two-way ANOVA with Bonferroni post-test. Asterisks

indicate levels of statistical significance, where *, p<0.05 and **, p<0.01. When compared to

WT-HFD mice, #, p<0.05.

The following figure supplement is available for figure 3:

Figure 3- Source Data 1.

1076

1077

Figure 4. Astrocytes cultured in high saturated fat conditions increase CD38 and pro-

inflammatory cytokine expression and inhibit oligodendrocyte differentiation.

A-D Schematic depicting how astrocyte conditioned media (ACM) treatments on primary

murine oligodendrocytes were performed (***A***). Conditioned media obtained from PA-treated

murine astrocytes (ACM (PA)) reduced protein expression of mature oligodendrocyte markers

PLP (***B, C***) and MBP (***B, D***), while PA-78c-ACM-treated OLCs have enhanced PLP and MBP

expression.

E-H Representational diagram (**E**) of astrocyte treatments with PA and NAD^+ to collect ACM for

OPC treatments. The addition of NAD^+ restored PLP levels (***F, G***) when compared to ACM-PA,

but had no effect on Olig2 (***F, H***). Scale bar = 100µm.

I-K High fat consumption was modeled in primary murine astrocyte cultures by including

saturated fat palmitate in the media (PA, 100 µM, 24 h). ***I***, Addition of a small molecule CD38

inhibitor (78c, 3µM) was effective to restore NAD^+ levels in PA-treated astrocytes when assessed

using an NAD^+ cycling assay. Astrocytes derived from CD38 knockout mice did not show

depleted NAD^+ levels in response to PA. RNA levels by qPCR (***J***) of proinflammatory cytokines,

antioxidant enzyme, and an astrocyte activation marker in PA and 78c treated astrocytes and

cytokine levels (***K***) detected in astrocyte conditioned media (ACM) by Bioplex assay.

Bar graphs represent the mean +/- SEM. Asterisks indicate levels of statistical significance when

compared to control as determined by Student's t-test (***A***, ***B***) or one-way ANOVA with

Bonferroni post-test (***C-L***), where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. # indicates $p<0.05$ when

compared to ACM-PA group by one-way ANOVA with Bonferroni post-test.

1099

1100   The following figure supplement is available for figure 4:

1101   **Figure 4- Source Data 1.**

1102

<u>Figure 5.</u> NAD$^+$ replenishment attenuates PA-induced deficits in oligodendrocyte

differentiation in a SIRT1-dependent manner.

A-C Representational diagram (**A**) of experimental treatments in primary murine

oligodendrocyte lineage cells (OLCs). Adding exogenous NAD^+ (50 µM) to oligodendrocyte

cultures significantly restores PLP protein expression (**B, C**) in PA-exposed OLCs.

D-G Treatment with NAD^+ precursor nicotinamide mononucleotide (NMN,100 µM) also

enhances expression of mature oligodendrocyte markers in PA-treated cultures, although not

when SIRT1 inhibitor Ex-527 (5 µM) is also included in the media, as determined by

immunostaining quantification for PLP (**D, E**) and MBP (**D, F)** per DAPI+ cell (**G**). Scale bar=

100 µm.

Graphical results represented as the mean ± SEM (n=4-8/group). Asterisks indicate levels of

statistical significance when compared to designated group as determined by one-way ANOVA

with Bonferroni post-test, where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$. # indicates $p<0.05$ and ##

represents $p<0.001$when compared to the ACM-PA group.

1117

1118   The following figure supplement is available for figure 5:

1119   **Figure 5- Source Data 1.**

1120

1121

Figure 6. CD38 inhibition protects from saturated fat and lysolecithin-induced

demyelination in cerebellar slices

A-C Representational diagram (**A**) of *ex vivo* remyelination experiments where murine

organotypic cerebellar slices were demyelinated with lysolecithin (LL, 0.5 mg/ml, overnight)

followed by subsequent culture in control media for 1 wk. Confocal microscopic imaging of

slices indicate reduced MBP (red, *B*) expression when normalized to neurofilament (NF, green).

LL-demyelinated slices cultured with 78c had significantly increased MBP expression when

normalized to NF staining and expressed as a percentage of the control group (**C**).

D-F Slices cultured in media containing both LL-palmitic acid (PA, 100 µM) and 78c, a CD38

inhibitor, 3 µM) had significant improvements in Olig2 (***D,E***) and MBP (***D, F***) expression

compared to the LL-PA alone group (#, $p<0.05$).

Bar graphs represent the mean ± SEM (n=4-8/group); *, $p<0.05$, **, $p<0.01$, and ***, $p<0.001$

when compared to control by one-way ANOVA with Bonferroni post-test.

1135

1136   The following figure supplement is available for Figure 6:

1137   **Figure Supplement 1**. Cerebellar slice culture treatment controls.

1138   **Figure 6- Source Data 1.**

1139

1140

Figure 7. CD38 inhibition with 78c improves myelin repair following lysolecithin-induced

focal demyelination to the ventral spinal cord.

Eight week old male C57Bl6 mice were fed a regular diet (RD) or high fat diet (HFD) for 5 wks,

then split into two additional groups: one receiving the same diet and the other half receiving the

respective diet with 78c added (600 ppm). One week later, a focal demyelinating lesion to the

ventral spinal cord was induced by lysolecithin (LL, 1% with Evans Blue dye) and mice were

allowed to recover for two weeks. Lesion boundary regions were determined by limiting to the

area of continuous myelin loss (core) or by including the entire inflammatory lesion (rim) based

on immunohistochemical analyses.

A Schematic depicting experimental design and lesion boundaries.

B Representative images and corresponding analysis illustrating the mean number of

remyelinated axons after focal lysolecithin-mediated demyelination of the ventral spinal cord

was increased in 78c-treated mice at 14 days post injury when compared to LL only mice (**,

$p < 0.01$, two-way ANOVA with Bonferroni post-test) as assessed by paraphenylenediamine

(PPD)-stained semi-thin sections of araldite-embedded spinal cord tissue. Scale bar= 20 μm.

C Immunohistochemical staining of the total lesion and lesion rim identified significant changes

in MBP immunoreactivity in 78c treated animals.

D-E Olig2+ oligodendrocyte lineage cells (OLCs, *D*) and GST3+ mature oligodendrocytes (*E*)

were significantly higher in 78c-RD mice when compared to RD controls.

Bar graphs represent mean +/- SEM. Two-way ANOVA (n=7-10 mice/group). Asterisks indicate

levels of statistical significance, where *, $p < 0.05$ and **, $p < 0.01$.

1162

1163   The following figure supplements are available for Figure 7:

1164   **Figure 7- Figure supplement 1.** Lysolecithin-induced demyelination model characterization.

1165   **Figure 7- Figure supplement 2.** 78c attenuates HFD-induced oligodendrocyte loss.

1166   **Figure 7- Source Data 1.**

1167

1168

Figure 8. 78c promotes pro-repair astrocytes after lysolecithin-induced demyelination.

Histological analysis of markers of neuroinflammation in animals fed a regular diet (RD) or high

fat diet (HFD) with or without 78c (CD38 inhibitor, 600 ppm) and injected with lysolecithin

(LL) followed by a 2-week recovery period.

A, B IBA1 (**A**, microgliosis) and GFAP (**B**, astrogliosis) were not significantly affected by either

diet or 78c alone, although a significant interaction for drug and diet was detected for IBA1

immunoreactivity in the lesion rim. Scale bar= 100 microns.

C-D, Changes in and pro-repair (S100a10, brown) pro-inflammatory (C3d, red) astrocyte (GFAP,

green) markers in adjacent sections within the LL lesion area indicates a shift toward pro-repair

in the 78c-treated mice. Scale bar = 100 μm.

Bar graphs represent mean +/- SEM. Two-way ANOVA (n=5-8 mice/group). Asterisks indicate

levels of statistical significance, where **, $p<0.01$ when compared to LL-RD and #, $p<0.05$

when compared to LL-HFD group by Bonferroni post-test.

1182

1183   The following figure supplement is available for figure 8:

1184   **Figure 8- Source Data 1.**

1185

1186

<u>Figure 9.</u> Schematic representation of proposed hypothesis.

Resting astrocytes typically help to provide a supportive environment for oligodendrocyte

differentiation and repair, when necessary. High fat diet (HFD) and lysolecithin (LL) increase

CD38 expression in reactive astrocytes, thereby depleting NAD$^+$ levels. Reductions in NAD$^+$

result in oxidative stress and inflammation which triggers impaired differentiation and survival in

oligodendrocyte lineage cells. 78c inhibits CD38 to attenuate the described detrimental effects of

astrocytes on oligodendrocytes.

1194

1195   **Supplemental Figures and Figure Legends**

1196

1197   **Figure 1- Figure Supplement 1. CD38 RNA expression in mouse spinal cord tissue.**

1198   **A** Heat map of HFD-induced changes in RNA expression of NAD$^+$-related genes acquired from

1199   previously published dataset (Langley et al, 2019).

1200   **B** RNA scope *in situ* hybridization identified more CD38+ puncta in ALDH1+ (astrocyte marker)

1201   cells following HFD versus RD.

Figure 2- Figure Supplement 1. Effect of direct CD38 inhibition on oligodendrocyte progenitors.

A-D Immunofluorescent staining of primary murine oligodendrocyte progenitors cultured in differentiation media indicates reduced PLP (B) and MBP (C) expression per cell (D) in the presence of palmitate (PA, 100 µm) that was not attenuated by direct 78c (3 µM) treatment. Graphical results represented as the mean ± SEM (n=3/group). Asterisks indicate levels of statistical significance when compared to designated group as determined by one-way ANOVA with Bonferroni post-test, where *, $p<0.05$; **, $p<0.01$.

1212

1213 **Figure 6- Figure Supplement 1. Cerebellar slice culture treatment controls.**

1214 **A-B** Confocal imaging of cerebellar murine slice cultures after two weeks shows no changes in

1215 MBP expression with 78c (3 µm) or palmitate (PA, 100 µm) treatments.

1216 Bar graphs represent the mean ± SEM (n=3-4/group); Data were analyzed by one-way ANOVA

1217 with Bonferroni post-test.

1218

Figure 7- Figure Supplement 1. Lysolecithin-induced demyelination model

characterization.

Eight week old male C57Bl6 mice were fed a regular diet (RD) or high fat diet (HFD) for 5 wks,

and then split into two additional groups: one receiving the same diet and the other half receiving

the respective diet with 78c added (600 ppm). One week later, a focal demyelinating lesion to the

ventral spinal cord was induced by lysolecithin (LL, 1% with Evans Blue dye) and mice were

allowed to recover for two weeks. Lesion boundary regions were determined by limiting to the

area of continuous myelin loss (core) or by including the entire inflammatory lesion (rim) based

on immunohistochemical analyses.

A Body weights were not significantly affected by 78c, but were by high fat diet (HFD)

consumption when compared to regular diet (RD).

B Food intake was significantly altered by HFD or 78c consumption.

C H&E staining shows the lesion area (C, entire boundary outlined by dotted line and

determined by consideration of all stains performed). No significant differences between RD and

HFD groups were noted.

D Neurofilament, (a marker of axon integrity) as determined by immunohistochemical

quantification, was unchanged by either drug or diet.

E Quantification of the core only region from histological analyses indicates no significant

changes.

Bar graphs represent mean +/- SEM. Two-way ANOVA (n=7-10 mice/group). Asterisks indicate

levels of statistical significance, where *, $p<0.05$ and **, $p<0.01$.

1241

Figure 7- Figure Supplement 2. 78c attenuates HFD-induced oligodendrocyte loss.

1242

A Histological analysis of the uninjected (UI) ventral side of the spinal cord from mice fed a

regular diet (RD) or high fat diet (HFD) for 5 wks, and then split into two additional groups: one

receiving the same diet and the other half receiving the respective diet with 78c added (600

ppm). Olig2+ oligodendrocyte lineage cells are depleted in HFD-fed mice compared to RD, yet

mice fed HFD with 78c added had significantly more oligodendrocytes than HFD alone mice (#,

$p<0.05$).

Bar graphs represent mean +/- SEM. Two-way ANOVA followed by Bonferroni post-hoc test

(n=6-9 mice/group). Asterisks indicate levels of statistical significance when compared to

control, where **, $p<0.01$; ***, $p<0.001$.

1252

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 1- supplement 1

A

	RD1	RD2	RD3	HFD1	HFD2	HFD3	HFD4	
NAD biosynthesis or salvage								Qprt
								Nmrk1
								Nmrk2
								Nampt
								Nmnat1
								Nmnat2
								Nmnat3
								Nadsyn1
								ACMSD
NAD consuming								Sirt1
								Sirt2
								Sirt3
								Sirt4
								Sirt5
								Sirt6
								Sirt7
								CD38
								PARP1
								PARP2
								PARP3
								PARP4
								PARP6
								PARP8
								PARP9
								PARP10
								PARP11
								PARP12
								PARP14
								PARP16

Relative

Row Minimum — Row Maximum

B

Figure 2- supplement 1

A

	Control	78c	PA	PA + 78c
PLP				
MBP				
DAPI				
Merge				

B

PLP

PLP/ DAPI

Control 78c PA PA + 78c

C

MBP

MBP/Dapi

Control 78c PA PA + 78c

D

Dapi count

# of cells/ field

Control 78c PA PA + 78c

Figure 6- supplement 1

Olig2

Figure 7- supplement 2

A

RD

HFD

RD+78c

HFD+78c

Ventral

Olig2$^+$ cells/mm^2

***

###

UI   UI +78c

■ RD
■ HFD

Ventral GM

Olig2$^+$ cells/mm^2

**

#

UI   UI +78c

■ RD
■ HFD

Ventral WM

Olig2$^+$ cells/mm^2

***

#

UI   UI +78c

■ RD
■ HFD