Critical role for astrocyte NAD$^+$ glycohydrolase in myelin injury and regeneration

Monica R. Langley[1], Chan-II Choi[1], Thais R. Peclat[2], Yong Guo[3], Whitney Simon[1], Hyesook Yoon[1,4], Laurel Kleppe[1], Claudia F. Lucchinetti[3], Claudia C.S. Chini[2,5], Eduardo N. Chini[2,5], and Isobel A. Scarisbrick[1,4,*]

Affiliations
[1] Department of Physical Medicine & Rehabilitation, Rehabilitation Medicine Research Center, Mayo Clinic, Rochester, MN 55905, USA
[2] Robert & Arlene Kogod Center on Aging, Mayo Clinic, Rochester, MN 55905, USA
[3] Department of Neurology, College of Medicine, Mayo Clinic, Rochester, MN, 55905, USA.
[4] Department of Physiology and Biomedical Engineering, Mayo Clinic, Rochester, MN 55905, USA
[5] Department of Anesthesiology, Mayo Clinic, Rochester, MN 55905, USA

[*] Corresponding author. Tel: 507-774-0124; E-mail:scarisbrick.isobel@mayo.edu

Conflict of Interest
ENC holds a patent on the use of CD38 inhibitors for metabolic diseases. The other authors declare no competing financial interests.

Abstract

Western-style diets cause disruptions in myelinating cells and astrocytes within the mouse CNS. CD38 has increased expression in the cuprizone and EAE demyelination models and is the main NAD^+ depleting enzyme in CNS tissue. Altered NAD^+ metabolism has been linked to both high fat consumption and Multiple Sclerosis (MS). We identified increased CD38 expression in the male mouse spinal cord following chronic high fat consumption or focal lysolecithin-induced demyelinating injury as well as in reactive astrocytes within an active MS lesion. CD38-catalytically inactive mice are significantly protected from high fat-induced NAD^+ depletion, oligodendrocyte loss, oxidative damage, and astrogliosis. 78c, a CD38 inhibitor, increased NAD^+ and attenuated neuroinflammatory changes in astrocytes induced by saturated fat. Conditioned media from saturated fat-treated astrocytes impaired oligodendrocyte differentiation pointing to indirect mechanisms of oligodendrogliopathy. Combined saturated fat and lysolecithin demyelination in cerebellar slices resulted in additional deficits in myelin proteins that were mitigated by concomitant 78c treatment. Importantly, oral 78c increased counts of oligodendrocytes and remyelinated axons after focal demyelination. Our findings suggest high fat diet impairs oligodendrocyte survival and differentiation through astrocyte-linked mechanisms mediated by the NAD^+ase CD38, and highlight the use of CD38 inhibitors as potential therapeutic candidates to improve myelin regeneration.

Introduction

Introduction

Obesity is a key risk factor for developing multiple sclerosis (MS)(1, 2). Moreover,

recent studies suggest that obesity and related metabolic comorbidities also influence

disease progression and disability burden in individuals affected by MS (3). Although

obesity-induced white matter abnormalities and loss of oligodendrocytes within the

central nervous system (CNS) following chronic consumption of high fat diets (HFD) has

been found in rodents, how diet influences white matter integrity in humans and

interacts with MS pathophysiology is largely unknown (4-6).

Astrocyte activation occurs following chronic HFD feeding in the CNS (7, 8) and

can exert oligotoxic effects (9). Building and maintenance of myelin membranes

requires oligodendrocyte fatty acid synthesis, yet oligodendrocytes can utilize lipids from

the diet or synthesized by astrocytes (10). The potential to target astrocytes to improve

outcomes in MS is already suggested since some currently prescribed MS disease

modifying therapies impact astrocytes (11). Prior studies regarding the role of HFDs in

demyelination models have utilized immune-mediated models such as experimental

autoimmune encephalomyelitis (EAE) and demonstrate exacerbation of peripheral

immune responses or neuroinflammation with HFD consumption (12-15). There are few,

if any, studies regarding the effects of HFD on the capacity for remyelination. Since

oligodendrocyte function is intimately linked to neighboring glial cells and neuronal

activity, in addition to microenvironmental factors, the most promising interventions will

likely involve a combinatorial approach.

CD38 is an NAD^+-degrading enzyme expressed by astrocytes and was recently

shown to be upregulated in peripheral immune- and toxin-mediated demyelination

models (15-18). CD38 is the main NAD$^+$ase in many mammalian tissues and regulates

tissue and cellular NAD$^+$ homeostasis as well as NAD$^+$ dependent enzymes such as

PARPs and Sirtuins (19, 20). Global CD38 knockout confers protection against

demyelination, T cell responses, and EAE disease severity as well as HFD-induced

metabolic dysfunction in rodent models (15, 18, 21, 22). Here we test the hypotheses

that HFD consumption promotes oligodendrocyte loss in a CD38-dependent manner

and that CD38 inhibition will enhance myelin regeneration. First, we established

increased CD38 expression following HFD consumption. Our previous studies indicate

that high fat diets lead to a reduction in oligodendrocytes and their progenitors within the

mouse CNS (4, 6, 8). Next, we demonstrated that CD38 catalytically inactive mice fed a

HFD were protected from HFD-mediated loss of oligodendrocytes with concomitant

increases in NAD$^+$ levels and reduced neuroinflammation and oxidative damage. *In vitro*

approaches identified oligodendrocyte direct and astrocyte-linked indirect mechanisms

by which excessive fatty acid impairs myelinating cell function. Finally, the value of a

small molecule inhibitor of CD38, 78c, was then identified as a novel protective strategy

to enhance myelin regeneration in a model of focal demyelinating injury. Together these

findings highlight a novel indirect pathway of myelin injury involving pathogenic changes

in astrocytes. Moreover, these changes can be attenuated by inhibition of CD38 in the

context of chronic HFD consumption and targeted to improve myelin regeneration in the

CNS.

Results

Astrocytes increase CD38 in the spinal cord following chronic HFD or demyelination

Our recent studies revealed spinal cord changes involving ER stress, mitochondrial dysfunction, and oxidative stress in the spinal cords of mice chronically consuming a HFD and the direct potential of saturated fat palmitate (PA) to inhibit differentiation of primary murine oligodendrocytes from their progenitors or neural stem cells (4, 6). Although saturated fat inhibited markers of differentiation *in vitro*, it was not oligotoxic, suggesting a more complex mechanism at play *in vivo* (4). Considering the emerging role of astrocytes in the pathology of various neurological disorders (9, 11, 23) and in the CNS following chronic HFD (7, 8, 24), along with their known essential metabolic and functional links with oligodendrocytes (25, 26), we decided to address the possibility of indirect effects of HFD mediated by astrocytes (Figure 1A).

NAD^+ is a key mediator of cellular energy status and is involved in mitochondrial function and quality control processes primarily through the activity of sirtuins that require NAD^+ as a cofactor (19, 20, 27). Exercise is known to increase NAD^+ levels in various tissues (22), and protect against "Western-style diet"-induced deficits in myelin and myelinating cells in two recent studies (5, 6). Our recent untargeted metabolomics profiling of spinal cord tissue identified NAD^+ metabolism as one of the top pathways impacted by HFD consumption (4), which prompted us to quantify spinal cord NAD^+ levels using a cycling assay. Spinal cord NAD^+ levels from HFD-fed mice were less than half that of RD-fed mice (p=0.029, t=2.639, df=8; Figure 1B).

Since CD38 is the main NAD$^+$-glycohydrolase in many mammalian tissues(19,

20, 27), we next quantified CD38 expression in spinal cord astrocytes to establish a

potential link. Quantification by qPCR indicated higher RNA expression of CD38 in

HFD-fed mice compared to regular diet-fed controls (p=0.001, t=5.626, df=6; Figure 1

C). Consistent with findings demonstrating CD38 is primarily localized to astrocytes

within the CNS (16-18, 28), increases in CD38 immunoreactivity were observed in

GFAP$^+$ astrocytes following HFD (p=0.038, t=1.937, df=12; Figure 1D). The HFD-

mediated effects mirror the increased CD38 expression following lysolecithin-induced

demyelination shown here (p=0.004, t=5.105, df=6; Figure 1E) and in EAE (15) and

cuprizone (18) associated demyelination studies. Similar to the HFD spinal cords,

primary murine astrocytes cultured with PA (100 µM, 24 h) showed increased GFAP

(p=0.046, t=2.507, df=6; Figure 1F) and CD38 (p=0.005, t=3.576, df=10; Figure 1G)

RNA expression and depleted NAD$^+$ levels (p=0.011, t=2.962, df=13; Figure 1H).

Collectively, these data suggest that HFD can induce astrocytosis in the spinal cord or

cultures, which increases expression of CD38 and subsequently depletes NAD$^+$ levels.

Microscopic imaging of human brain sections demonstrated extensive CD38

immunoreactivity in white matter astrocytes within control brain tissue and increased

CD38 in some hypertrophic reactive astrocytes within MS lesion tissue with active

demyelinating activity (Figure 1I). The expression and significance of CD38

immunoreactivity among various astrocyte subtypes and disease phenotypes needs to

be further characterized, but is likely not specific to MS since GFAP+CD38+ cells were

found to be increased in HIV-1-associated encephalopathy (HIVE) brain tissue sections

(29).

CD38 catalytically inactive (CD38ci) mice are resistant to HFD-induced

oligodendrocyte loss

To determine the potential therapeutic significance of CD38 upregulation to HFD-

induced oligodendrocyte loss, we collected spinal cords from wildtype (WT) and CD38-

catalytically inactive (CD38ci) mice fed a regular diet (RD) or HFD for 30 wk beginning

at 50 wk of age for histological analysis (Figure 2A). CD38ci mice have the CD38 gene

knocked out and reinserted with a mutation at E230 of the CD38 gene, rendering the

CD38-NAD$^+$ase catalytic site inactive, while preserving overall expression levels of

CD38 (30). To confirm that preservation of NAD$^+$ levels was achieved in the CD38ci

mice, we next measured NAD$^+$ levels on spinal cord tissue from the same cohort of

mice. As expected based on our data in Figure 1, NAD$^+$ levels were significantly

reduced following chronic HFD consumption in wildtype mice ($F_{(1, 24)}$=1.785, p=0.004;

Figure 2B). Moreover, in CD38ci mice, which had diminished CD38 activity ($F_{(1,}$

$_{16)}$=75.13, p<0.0001; Figure 2C), NAD$^+$ levels were significantly higher overall ($F_{(1,}$

$_{24)}$=33.90, p<0.0001) and in the CD38ci-HFD mice compared to the wildtype-HFD mice

(p<0.0001). As seen in our previous studies (4, 6), there was a significant reduction in

the number of PDGFRα$^+$ ($F_{(1, 16)}$= 16.61, p=0.0003 dorsal column and $F_{(1, 16)}$= 17.77,

p=0.0007 ventral; Figure 2E) and Olig2$^+$ oligodendrocyte progenitors ($F_{(1, 26)}$= 9.066,

p=0.006 dorsal column and $F_{(1, 26)}$= 24.90, p<0.0001 ventral; Figure 2 E) as well as

GST3$^+$ mature oligodendrocytes ($F_{(1, 22)}$= 14.53, p=0.001 dorsal column and $F_{(1, 25)}$=

10.23, p=0.004 ventral; Figure 2F) in the HFD-fed wildtype mice compared to regular

diet-fed mice. However, CD38ci mice fed HFD did not have reductions in

oligodendrocyte lineage cell counts compared to their regular diet genotype controls

(p=0.682), yet had significantly higher PDGFRα$^+$ (p=0.0397 dorsal column and

p=0.0198 ventral white matter) and GST3$^+$ (p=0.0001 dorsal column) cell counts versus

WT-HFD mice. The improved cell numbers are likely due to decreased apoptosis rather

than proliferative changes, as evidenced by decreased cleaved-caspase-3 staining ($F_{(1,}$

$_{16)}$= 5.48, p=0.0224 ventral; Figure 2E) in the CD38ci-HFD versus WT-HFD group and

no significant difference in Ki67. Across spinal cord regions, no significant changes in

MBP due to genotype or diet group were found ($F_{(1, 28)}$= 0.002, p= 0.966, and $F_{(1, 28)}$=

0.683, p=0.415, respectively, Figure 2D). Overall, these data suggest that

oligodendrocyte lineage cell survival is preserved in CD38ci mice following chronic HFD

consumption.

Diminished neuroinflammatory responses and oxidative damage in CD38ci mice

following chronic HFD

Neuroinflammation, including complex interplay between microglia and

astrocytes, is now appreciated to contribute to oligotoxicity and MS pathophysiology (9,

25). To explore the potential impact of blocking CD38 catalytic activity on CNS

neuroinflammation following chronic HFD consumption, we next performed

immunohistochemical analysis on the spinal cords from wildtype and CD38ci mice fed

regular or HFD for markers of microglia and astrocytes. Quantification of the

thresholded area considered positive for ionized calcium-binding adapter molecule 1

(IBA1) demonstrated decreased microglia numbers or activity ($F_{(1, 21)}$= 5.079, p=0.030;

Figure 3A) in the ventral grey matter of spinal cords from CD38ci mice compared to

wildtype mice overall. In addition, HFD-fed CD38ci mice showed significantly reduced

IBA1 immunoreactivity in the ventral grey and white matter of the spinal cord compared

to wildtype mice fed a HFD. Although HFD did not significantly increase IBA1-

immunoreactivity in this study, this is consistent with a recent Western-style diet study

from our laboratory and older mice may already have residually higher levels (8, 31).

Considering that astrocytes have endogenous expression of CD38 as well as elevated

CD38 expression following HFD or demyelination (16-18), we also investigated

astrocyte reactivity in these mice. In the ventral spinal cord, quantification of astrocyte

marker glial fibrillary acidic protein (GFAP, $F_{(1, 28)}$= 12.66, p=0.0014; Figure 3B)

demonstrated significantly reduced positive area in the white matter of CD38ci mice

compared to wildtype mice. Wildtype HFD-fed mice showed significantly more astrocyte

reactivity compared to either regular diet (p=0.023) or HFD-fed CD38ci mice (p=0.043).

190   Given our previous data suggesting increased oxidative damage markers in the

CNS following chronic consumption of HFDs (4, 6) and the significance of oxidative

stress in glia of MS lesions (32), we next sought to compare the extent of oxidative

damage in the spinal cords of wildtype and CD38ci mice fed regular or HFD. More

abundant lipid peroxidation marker 4-hydroxynonenal (4HNE) was present in the HFD-

fed wildtype mice than regular diet-fed wildtype mice in the spinal cord dorsal column

($F_{(1, 28)}$= 3.358, p=0.019, Figure 3C). Importantly, a significant reduction in 4HNE was

found in the HFD-CD38ci mice compared to HFD-wildtype mice in the dorsal column

and ventral spinal cord (grey and white matter). Superoxide dismutase 2 (SOD2) is a

mitochondrial antioxidant that can protect from high fat-induced changes in oxidative

stress and metabolism (33), while brain-derived neurotrophic factor (BDNF) is

permissive to myelination (23) and S100a10 is a recently defined pro-repair astrocyte

maker (9). Next, our histological findings were further substantiated by observed gene

expression increases in SOD2 ($F_{(1, 8)}$= 7.721, p=0.024 ; Figure 3D) and pro-repair

astrocyte markers genes BDNF($F_{(1, 8)}$= 5.685, p= 0.044; Figure 3E) and S100a10 ($F_{(1,}$

$_{8)}$= 24.73, p= 0.0011; Figure 3 F) in the CD38ci mice, as determined by qRT-PCR.

Overall, the observed reductions in neuroinflammation and oxidative stress markers and

increased NAD$^+$ levels in the CD38ci mice likely provide a more suitable

microenvironment for oligodendrocyte survival, differentiation, and repair following

chronic HFD consumption.

CD38 inhibition with 78c improves myelin repair following lysolecithin-induced

focal demyelination of the ventral spinal cord

78c is a thiazoloquin(az)olin(on)e CD38 inhibitor that increases NAD$^+$ levels in

various tissues and cell types and is orally bioavailable (30, 34). To determine whether

pharmacologic inhibition of CD38 affects the potential for remyelination, we

administered 78c to mice in each diet group (regular or high fat) and used histological

approaches to investigate numbers of oligodendrocyte lineage cells and remyelinated

axons in the lysolecithin model of demyelination. Six-week old male wildtype mice were

given either a regular or HFD for 5 weeks, then split into additional groups: receiving the

same diet with or without 78c (600ppm) (Figure 4A). All mice were subjected to a focal

demyelinating injury to the ventral spinal cord by microinjection of 1% lysolecithin.

After two weeks of recovery, lesions were collected from perfused mice and used

for subsequent counting of remyelinated axons and myelinating cells. At the study

endpoint, HFD mice had gained significantly more weight than their RD counterparts

($F_{(1, 36)}$=51.91, p<0.0001, Supplemental Figure 1A), irrespective of genotype. While 78c

treatment did not have a significant effect on body weights, mice given 78c consumed

less food (Supplemental Figure 1B). Average lesion area (0.107 mm^2 RD, 0.092 mm^2

HFD, 0.132 mm^2 RD-78c, and 0.117 mm^2 HFD-78c, as determined by hematoxylin and

eosin, Supplemental Figure 1C) was not significantly altered by either HFD ($F_{(1, 24)}$=

2.642, p=0.117) or 78c ($F_{(1, 24)}$=0.886, p=0.356). Neurofilament immunoreactivity (a

marker of axon integrity, Supplemental Figure 1D), was 18.09% RD, 15.76% HFD,

18.29% RD-78c, and 18.71% of the lesion area on average, and also remained

unchanged by drug ($F_{(1, 24)}$= 0.717, p=0.405) or diet ($F_{(1, 24)}$= 0.261, p=0.614). Overall,

lysolecithin mice fed 78c along with either a RD or HFD showed significantly greater

numbers of remyelinated axons compared to the lysolecithin mice fed a RD or HFD

without 78c ($F_{(1, 25)}$= 10.11, p= 0.004; Figure 4B). In particular, more than double the

number of remyelinated axons was present in the 78c-regular diet mice over regular diet

alone (p=0.0013). Further supporting a role of CD38 inhibition in myelin repair, MBP

immunoreactivity in the total lesion or the lesion rim was higher in the 78c-treated mice

irrespective of diet ($F_{(1, 24)}$= 5.71, p=0.025; Figure 4C). Importantly, numbers of Olig2$^+$

($F_{(1, 23)}$ = 9.508, p= 0.005; Figure 4D) and GST3$^+$ ($F_{(1, 24)}$= 6.172, p=0.020; Figure 4E)

cells were significantly higher in the combined 78c group (which includes LL-RD-78c

and LL-HFD-78c mice) versus the combined group without 78c treatment (which

includes LL-RD and LL-HFD). Moreover, the number of Olig2$^+$ cells in the lesion rim

(p=0.049) and GST3$^+$ cells in the rim (p= 0.044) or total lesion (p= 0.032) were

significantly higher in the LL-RD-78c versus the LL-RD group. These findings suggest

that, *in vivo,* 78c helps to preserve or replenish myelinating cells and their progenitors,

leading to increased markers of myelin repair. Furthermore, these results provide a

strong rationale for targeting CD38 and overall NAD$^+$ metabolism as a new strategy to

attenuate oligodendrocyte loss and aid in myelin regeneration.

78c promotes pro-repair astrocytes after lysolecithin-induced demyelination

Next, we chose to explore the role of 78c on neuroinflammation markers in the

spinal cord following demyelination. Histological quantification of threshold-positive

staining for microglia marker IBA1 (Figure 5A) indicated a significant interaction

between diet and drug treatment on the lesion rim ($F_{(1, 24)}$= 6.936, p=0.015), yet no

significant effect of diet ($F_{(1, 24)}$= 2.160, p=0.154) or drug ($F_{(1, 24)}$= 0.2549, p=0.618)

alone. Although there is a trend of decrease for both IBA1 and GFAP in the 78c treated

group, no statistically significant attenuation of astrocyte marker GFAP was observed

(Figure 5 B). In both the RD- and HFD-fed mice, there was, however, a significant

increase in S100a10 immunoreactivity with 78c treatment ($F_{(1, 22)}$= 20.77, RD: p=0.001,

HFD: p=0.045; Figure 5C), considered a marker for pro-repair astrocytes. There was

also a slight decrease in C3d staining, a marker of reactive astrocytes, with 78c

treatment, although this did not reach statistical significance ($F_{(1, 19)}$= 3.681, p=0.069;

Figure 5D).

Astrocytes cultured in high saturated fat conditions increase CD38 and pro-

inflammatory cytokine expression and inhibit oligodendrocyte differentiation

269   Astrocytes release various growth factors, cytokines, chemokines, extracellular

270 matrix proteins, glutamate, and reactive oxygen species that heavily influence

271 oligodendrocyte proliferation and differentiation (25, 26, 32, 35). Therefore, we next

272 investigated the effects of culturing oligodendrocytes in conditioned media obtained

273 from PA-treated astrocytes with or without 78c co-treatment, as depicted in Figure 6A.

274 As expected, astrocyte-conditioned medium (ACM) from PA-treated astrocytes

275 significantly reduced oligodendrocyte PLP ($F_{(2, 17)}$= 9.396, p=0.006) and MBP

276 expression ($F_{(3, 8)}$= 8.155, p=0.003). However, this negative impact on myelin protein

277 production was partially rescued by co-treatment with the small molecule CD38 inhibitor

278 78c (3 µM) (p=0.002, PLP; and p=0.040, MBP) (30, 34). Likewise, addition of

279 exogenous NAD^+ (50 µM) to the PA-treated astrocyte cultures ($F_{(2, 9)}$= 8.502, p=0.008;

280 Figure 6 B) significantly increased PLP expression, while ACM-treated oligodendrocyte

281 numbers remained unchanged across treatment groups ($F_{(2, 9)}$= 1.153, p=0.358). Along

282 with other research that indicates the role of NAD^+ in reducing astrocyte

283 proinflammatory and oxidative stress responses (18, 36), these data indicate the

284 importance of maintaining proper astrocyte NAD^+ levels in order to promote

285 oligodendrocyte differentiation.

286   To address whether the possible NAD^+-driven astrocytic changes indirectly

287 mediate the observed effects on oligodendrocyte, we next used primary murine

288 astrocyte cultures to assess NAD^+ levels and transcriptional changes. Addition of 78c

289 (3µM) was effective to restore NAD^+ levels in PA-treated astrocytes when quantified

290 using an NAD^+ cycling assay ($F_{(4, 23)}$= 6.113; p=0.004; Figure 6C). However, astrocytes

291 derived from CD38 knockout mice did not have depleted NAD^+ levels in response to PA.

Next, we determined whether CD38 regulates astrocyte pro-inflammatory responses.

After treating astrocyte cultures with PA and/or 78c, we used qRT-PCR and a multiplex

bioassay for determining cytokine RNA transcripts and release into culture supernatant,

respectively. We observed significant increases in IL-1β and IL-6 expression ($F_{(3, 8)}$=

8.427, p=0.007, IL-1 β; and $F_{(3, 8)}$= 11.83 , p=0.007, IL-6; Figure 6D) in astrocytes

stimulated with PA and 78c significantly attenuated PA-induced RNA expression of IL-

1β (p=0.007), IL-6 (p=0.008), TNFα (p=0.017), and reactive astrocyte marker H2D1

(p=0.028). Additionally, the mitochondrial antioxidant SOD2 was significantly increased

in astrocytes treated with 78c (p=0.022) or both PA and 78c (p=0.035). Significant

increases in secretion of cytokines triggered by PA were observed using a Bioplex

assay including CXCL1 (KC) ($F_{(3, 8)}$ = 9.296, p= 0.014), MIP1α ($F_{(3, 8)}$= 549.8,

p<0.0001), RANTES ($F_{(3, 8)}$= 418.5, p<0.0001), IL-6 ($F_{(3, 8)}$= 18.00, p=0.001), MIP1β

($F_{(3, 8)}$= 533.5, p<0.0001), and MCP1 ($F_{(3, 8)}$= 66.96, p<0.0001)(Figure 6 E). Notably, co-

treatment with 78c successfully mitigated PA-stimulated secretion of CXCL1 (p=0.006),

MIP1α (p<0.0001), RANTES (p<0.0001), IL-6 (p=0.001), MIP1β (p<0.0001), and MCP1

(p<0.0001). Thus, inflammatory and oxidative stress-related changes in cultures

astrocytes may help to explain reduced differentiation potential of the oligodendrocyte

progenitors in the presence of PA-treated astrocyte conditioned medium.

NAD$^+$ and NMN restore oligodendrocyte differentiation in PA-treated cultures

Our recent findings demonstrated that PA can directly impair oligodendrocyte

differentiation from neural stem cells or oligodendrocyte progenitors (4), and others

have indicated that depletions in cellular NAD$^+$ can prevent differentiation of stem cells

to mature oligodendrocytes (37). To examine the effects of replenishing NAD$^+$ levels on

oligodendrocytes, we next supplemented cultures with exogenous NAD$^+$ (50 µM) and

monitored PLP expression (Figure 7A). Co-treatment with PA and NAD$^+$ significantly

enhanced PLP expression ($F_{(2, 16)}$=3.458, p=0.024; Figure 7B) compared to exclusive

PA treatment. However, treatment of oligodendrocytes with CD38 inhibitor 78c did not

attenuate this effect (data not shown).

321   Interestingly, CD38$^+$ cells can deplete NAD$^+$ and metabolites available for low-

CD38 expressing cells (30, 38), such as oligodendrocytes (16-18, 28). We found that

direct effects of saturated fat treatment on oligodendrocytes include mitochondria

function and morphology changes (4) while other mechanisms, potentially including

NAD$^+$-depleting mechanisms (37, 39, 40), could also play a part. SIRT1 regulates

mitochondria function, was decreased following HFD in spinal cord tissue (4), and is a

NAD$^+$-dependent deacetylase.   We hypothesized that the pro-myelinating effects of

increasing NAD$^+$ are SIRT1-dependent, so we determined if the beneficial effects of

NAD$^+$ in the presence of PA could be blocked with a SIRT1 inhibitor Ex-527. Similar to

adding exogenous NAD$^+$, treatment with the NAD$^+$ precursor NMN (100 µM) also

restored markers of oligodendrocyte differentiation in PA-treated cultures ($F_{(3, 25)}$=8.449,

p=0.011; Figure 7C). However, in the presence of Ex-527 (5 µM), the increases in PLP

and MBP protein in response to NMN were suppressed (p=0.003). At the doses used,

PA, NMN, and Ex-527 did not significantly alter the number of oligodendrocytes. Hence,

the impaired ability of oligodendrocytes cultured with excess saturated fat to

differentiate is at least partially regulated by NAD$^+$-dependent sirtuin activity (Figure 7D).

78c prevents lysolecithin- and PA-mediated myelin protein and oligodendrocyte loss

Since we observed protective effects of genetic CD38 inhibition on oligodendrocyte survival following HFD, as well as increased CD38 expression in a spinal cord demyelination model, we next sought to explore benefits conferred by pharmacologic inhibition using 78c in a brain model of lysolecithin-induced *ex vivo* demyelination. Murine cerebellar slices from postnatal pups were demyelinated using lysolecithin (LL, 0.5 mg/ml, 18 h), followed by a period of one week to allow for repair (Figure 8 A). Confocal imaging of immunofluorescence staining demonstrated reductions in MBP after LL that was significantly attenuated by 78c co-treatment ($F_{(2, 25)}$=4.578, p=0.020; Figure 8B).

Finally, to evaluate the therapeutic potential of 78c in a brain model of high fat exposure combined with demyelination, organotypic cerebellar slices were demyelinated as described above and cultured for 7 d in the presence or absence of PA and/or 78c. Cerebellar slices showed fewer Olig2$^+$ cells ($F_{(6, 21)}$=4.964, p=0.003) with LL insult. Inclusion of PA in the media further diminished Olig2+ cells in the LL slices. After 7 d of recovery, Olig2$^+$ cells remained significantly depleted in the 7 d LL group. There was also less MBP immunoreactivity following LL ($F_{(6, 21)}$=2.899, p=0.020; Figure 8 C), however MBP levels were successfully restored after 7 d recovery. Importantly, 78c and PA co-treatment significantly increased the percentage of Olig2$^+$ cells (p=0.049) and MBP expression (p=0.023) in the demyelinated cerebellar slices. However, neither palmitate nor 78c alone, without lysolecithin, for two weeks significantly altered MBP expression (PA, p=0.668, and 78c, p=0.566 versus control). These findings demonstrate

361   the capability of 78c to restore myelin protein expression even in a combined saturated

362   fat- and demyelinating agent-induced condition. Overall, these data suggest that 78c

363   modulates astrocyte responses elicited by demyelinating injury or high fat diet in the

364   CNS, skewing them towards a phenotype that enhances oligodendrocyte survival and

365   differentiation and myelin repair (Figure 9).

Discussion

Relevance of Diet and NAD⁺ homeostasis to Multiple Sclerosis

Myelin disturbances and oligodendrocyte loss can leave axons vulnerable

leading to permanent neurological deficits. The results of the current study suggest that

metabolic disturbances, triggered by consumption of a diet high in fat, promote

oligodendrogliopathy and impair myelin regeneration through astrocyte-linked indirect

NAD^+-dependent mechanisms. Diet-induced obesity increases the risk for developing

MS, especially when combined with other known risk factors (1, 2). We demonstrate

that restoring NAD^+ levels via genetic inactivation of CD38 can overcome these effects.

Moreover, we show that therapeutic inactivation of CD38 can enhance myelin

regeneration. Together these findings point to a new metabolic targeting strategy

positioned to improve disease course in MS and other conditions in which the integrity

of myelin is a key concern.

Neuroinflammation and CD38

Neuroinflammation and astrogliosis are predominant pathological mediators of

neurodegenerative conditions, including MS (9, 41), and astrogliosis is linked to HFD-

induced CNS pathology in rodents (8, 24). In spinal cord astrocytes of HFD mice, we

observed increased CD38 expression and an overall depletion of NAD^+. These findings

are consistent with our human immunostaining data and other reports of increased

CD38 in the brain of an individual with MS (42) and in experimental models of

neurological injury and disease (21). Known upstream activators of CD38 expression

include cytokines, lipopolysaccharide (LPS), 25-hydroxycholesterol (25-HC), glutamate,

and factors secreted by senescent cells (38, 43-45). In the current study, we found that

HFD or focal lysolecithin-mediated demyelinating injury each drove increased CD38

expression in astrocytes, while genetic or pharmacological CD38 inhibition after HFD or

demyelination, respectively, attenuated astrocyte activation markers. Confirming the

human relevance of this effect, we observed CD38 in hypertrophic reactive astrocytes

within an active demyelinating lesion from an individual with MS.

Astrocytes and microglia secrete a variety of chemokines, cytokines and growth

factors that regulate myelin injury and regeneration (25, 35). In palmitate-stimulated

astrocytes, IL-6 and other cytokines were increased. CCL3/MIP1α, MIP1β, and

RANTES also increase in active demyelinating MS lesions and fluids (46, 47). Pointing

to a therapeutic strategy, this response was blunted when NAD^+ levels were restored by

co-treatment with 78c. Other studies indicate that CD38 inhibition or knockout

decreases cytokine secretion in LPS-stimulated microglia, macrophages, and

monocytes (48). Recently, knockout of CD38 or supplementation with NAD^+ was shown

to suppress activation of astrocytes in culture (18). Therefore, improving NAD^+ levels by

targeting CD38 shows therapeutic promise across multiple neurological and non-

neurological disease types (21), and here we establish its significance to

oligodendrocyte heath and regeneration with strong linkage to its effects on astrocyte

metabolism.

Oxidative stress and oligodendrocyte differentiation

Reactive oxygen species can also inhibit oligodendrocyte survival (32).

Complementing this, we demonstrate more SOD2 and less IL1β in high fat-78c co-

treated astrocyte cultures compared to treatment with high fat alone. We also found

more SOD2, BDNF and S100a10 in the CD38ci mouse spinal cord pointing to a shift

towards pro-repair astrocyte properties (9). Blocking downstream CD38 signaling also

reduced markers of oxidative damage in a convulsion model (49). Our findings suggest

that the beneficial effects of CD38 inhibition include protection against astrocyte pro-

inflammatory factors and reactive oxygen species known to negatively impact

oligodendrocyte renewal and maturation.

The effects of CD38 activity and NAD^+ levels may depend heavily on cell type

and disease context, with findings here supporting a critical protective role of lowering

astrocyte CD38 activity for protection of myelinating cells and for myelin regeneration in

the adult CNS. Future studies could further solidify this finding once astrocyte-specific

conditional knockouts for CD38 become available. CD38-positive cells can regulate

NAD^+ levels in neighboring CD38-negative cells by influencing NAD^+ precursor

availability (30, 38), which may be the case for astrocytes and oligodendrocytes in our

studies. Consistent with this model, 78c had no direct effect on oligodendrocyte

differentiation in the context of palmitate *in vitro*. Palmitate can inhibit nicotinamide

phosphoribosyltransperase (NAMPT) activity (40), a rate-limiting enzyme in the NAD^+

salvage pathway, as well which prevents differentiation of stem cells to oligodendrocyte

lineage cells (37) and potentially explains the direct inhibitory effect of palmitate on

oligodendrocyte differentiation observed here. Thus, complex alterations in NAD^+

biosynthesis, salvage, and degradation pathways may contribute to the effects of

palmitate on oligodendrocytes that we document can be rescued by NAD^+

replenishment.

NAD^+-boosting strategies for neural repair

Several recent studies have explored supplementing NAD$^+$ precursors or

flavonoids known to inhibit NAD$^+$ase activity to improve myelin repair, neuroprotection,

and to decrease immune cellular responses. For example, an NAD$^+$/BDNF/TrkB

pathway is proposed to be involved in nicotinamide-mediated protection, including

improved remyelination in a stroke model (50) and precursor supplementation increased

BDNF in a Huntington's disease model (51). Supporting a possible NAD$^+$-BDNF

pathway for remyelination, we observed increased BDNF expression in the spinal cord

of CD38ci mice and improved remyelination with oral 78c administration. Although direct

effects on myelinating cells were not examined, mice with CD38-knockout, given the

CD38 inhibitor apigenin, or supplemented with NAD$^+$ each showed reduced EAE

disease severity (15, 52, 53). NAD$^+$ levels can impact oligodendrocyte lineage cell

proliferation and survival via cyclins and E2f1, while differentiation is reported to be

regulated redundantly by SIRT1 and SIRT2 (37, 54, 55). Results here demonstrate that

the ability of NAD$^+$ supplementation to promote oligodendrocyte differentiation *in vitro* is

indeed at least partially reliant on SIRT1 function. Restoring NAD$^+$ levels was also

beneficial in the context of palmitate or high fat-induced inflammation and oxidative

stress via SIRT1 in liver, heart, and muscle (56-58). Improvements in expression of

myelin proteins in the spinal cord of mice consuming a high fat high sucrose diet in

conjunction with coordinate wheel running, were also linked to increases in SIRT1

expression (6) . Collectively, these findings are particularly encouraging since effective

modulation of disease progression will likely include a combination of therapeutic

strategies encompassing anti-inflammatory, myelin regeneration, and neuroprotective

interventions.

Study considerations

HFD consumption did not further impair markers of myelin integrity in our *in vivo*

model of myelin injury and repair as it did in *ex vivo* remyelinating cerebellar slice

cultures. Although treatment with 78c did improve myelin regeneration in mice

consuming a regular diet, little impact was seen in mice consuming a HFD, suggesting

the need to explore additional factors in the context of HFD-demyelination that

contribute to impairments. Alternately, a higher dose or combination of drugs (to target

various aspects of astrocyte responses) may be required to improve remyelination in

the context of HFD. Although a preference for pro-repair versus pro-inflammatory

phenotype was indicated following 78c treatment in the lysolecithin model of focal

demyelination studied, a complete astrocyte-specific transcriptomic profile in the HFD

mice with demyelination will be needed to identify novel genes in the astrocyte

transcriptome that could be targeted to mitigate the indirect oligotoxic effects we

observe.

Conclusions

Our data demonstrates an essential role of NAD^+ levels in maintaining CNS white

matter integrity, which can be impaired by focal demyelination or high fat diet

consumption. Further, we have shown that inhibition of the NADase CD38 attenuates

oligodendrocyte loss and promotes myelin repair. Genetic CD38 inhibition attenuated

HFD-induced oligodendrocyte loss and dietary supplementation with 78c increased

counts of oligodendrocytes and remyelinated axons following focal demyelinating injury.

These findings suggest that increases in astrocyte CD38, and the associated depletion

of NAD^+ levels, may contribute indirectly to impairments in oligodendrocyte health and

myelin regeneration via inflammation and oxidative stress. Tarrago, et al. (30) recently

established 78c as a specific, reversible, and non-competitive inhibitor of CD38

NAD^+ase activity improves age-related metabolic function (28). Altogether, these

findings suggest that genetic or pharmacologic CD38 inhibition provides therapeutic

benefits in HFD-induced metabolic dysfunction and downstream signaling events in the

CNS, in addition to the already described peripheral benefits on age-related metabolic

dysfunction. Importantly, inhibition of CD38 with 78c can improve myelin regeneration in

the adult CNS including reducing the pro-inflammatory profile while increasing the pro-

repair profile of astrocytes. Together these findings point to the potential translational

value of targeting CD38 to improve outcomes in MS and other demyelinating conditions.

A few current MS disease modifying therapies (i.e. glatiramer acetate, fingolimod,

dimethyl fumarate) affect CD38[+] cell counts or inhibit astrocyte activation, but these

roles have not been directly linked to therapeutic benefits (11, 15, 59-62). For instance,

fingolimod treatment decreased CD38 expression in a rat EAE model (15), and was

shown to promote beneficial astrocyte responses in culture (63). Monoclonal antibodies

targeting CD38 have already met with success in clinical trials for conditions outside the

CNS (43, 64-66), including Daratumumab which is FDA approved for use in multiple

myeloma. Accordingly, the research findings we present should inspire further research

into ways to mitigate the negative consequences of HFD on oligodendrocytes alone and

in the context of white matter injury which may include dietary intake of NAD^+

precursors, administration of CD38 inhibitors, or exercise-related rehabilitation

strategies.

Materials and Methods

See Table 1 for key reagents.

Animal husbandry and diets

All experimental diets and procedures were approved by The Mayo Clinic

Institutional Animal Care and Use Committee (IACUC) and performed in accordance

with institutional and regulatory guidelines. Mice were provided *ad libitum* access to

diets along with tap water. For initial characterization of CD38 expression and NAD$^+$

levels following a high-fat diet, male C57BL6/J mice were obtained from Jackson Labs

(Bar Harbor, ME) and were randomly assigned to feeding on either a regular diet (RD,

10 % kcal fat, D12450K, Research Diets, New Brunswick, NJ) or a high-fat diet (HFD,

60% kcal fat, D12492, Research Diets, New Brunswick, NJ) beginning at 10 wk of age,

as previously described (4).

For studies comparing wildtype and CD38ci mice fed RD and HFDs, mice were

bred onsite after initial generation at TransViragen (Chapel Hill, NC) via recombineering

technology, as previously described (30, 67). Male mice received a regular (10.4% kcal

fat, Teklad TD000357) or HFD (42% kcal fat, Teklad TD88137) beginning at 50 wk of

age for 30 wk. Diets used for lysolecithin demyelination studies were these same diets,

with or without 600 ppm 78c added (Calico Life Sciences, LLC), and were started one

week prior to the spinal cord microinjection procedure. Food intake and body weights

were monitored weekly.

Lysolecithin microinjection into spinal cord

For focal demyelination of the ventral white matter (laminectomy at thoracic 10[th]

vertebra; mediolateral 0.3, dorsoventral -1.4 mm relative to the dorsal surface) of the

murine spinal cord, 12 wk old male mice were injected with 2 µl of a saline solution

containing 1% lysophosphatidyl choline with 0.01% Evan's Blue (Sigma L4129) at 0.2

µl/min using a 30-40 µm glass micropipette and stereotaxic microinjection device

(Stoelting, Inc., Wood Dale, IL). Male mice were selected for lysolecithin injection

procedures to reduce variability due to progesterone-mediated enhancement of myelin

repair and modulation of CNS neuroinflammatory responses (68). Young adult mice

have previously yielded reliable, reproducible lesions and good survival in this model

from studies in our laboratory and others (23, 69). The spinal cord has large white

matter tracts that are more frequently myelinated axons versus corpus callosum,

allowing for more confident quantification of remyelinated axons due to challenges of

determining thin myelin sheaths or simply unmyelinated axons from demyelinated axons

(23, 69, 70).

For anesthesia, 100 mg/kg ketamine and 10 mg/kg xylazine were administered

intraperitoneally prior to surgery, while postoperative care consisted of an antibiotic

(Baytril 50 mg/kg/day, i.p.), an analgesic (Buprinex 0.05 mg/kg, twice daily,

subcutaneous) and saline (0.2ml/d, intraperitoneal) for three days. After a two-week

recovery period, mice were deeply anesthetized (ketamine 150mg/kg and xylazine 15

mg/kg, i.p.) and transcardially perfused with 4% paraformaldehyde (Fisher T353-500)

for collection of spinal cord tissue. Spinal cord segments were microdissected, with the

rostral 2 mm segment processed for paraffin-embedded blocks and the caudal segment

osmicated (1% osmium tetroxide, 0223B, Polysciences) and embedded in Araldite. To

identify myelin sheaths, Araldite blocks were cut to 1 µm sections, stained with p-

phenylenediamine (Sigma P6001), imaged with 60x objective (Olympus BX51

microscope), and stitched together (Adobe Photoshop) to comprise the entire lesion

within the stained tissue section (23, 70). Number of remyelinated axons per square

millimeter of lesion was reported after blinded counting in ImageJ (23, 70).

Immunohistochemical analysis of focal demyelinated lesions utilized overall

neuropathological changes (hematoxylin and eosin) along with myelin loss (based on

MBP staining) to determine the lesion core, while the lesion rim was defined as the

region encompassing the entire border of neuroinflammatory markers (GFAP, IBA1)

(70). Lesion areas between 50,000 - 180,000 µm2 were included in the analysis based

on our previous studies using this model (23, 70). Analyses were performed with

experimenter blinded to treatment group by assigning a random animal number.

Immunohistochemistry

Spinal cords were fixed in 4% paraformaldehyde, processed, paraffin-embedded,

and cut to 6 µm for all immunoperoxidase and immunofluorescence procedures, as

described previously (4, 23). When necessary (all antibodies except GFAP and MBP),

heat-induced antigen retrieval was achieved through incubation with sodium citrate (10

mM, pH 6.0) in a steamer basket. Sections were blocked (20 % v/v normal goat serum

in PBS), then incubated in primary antibodies described in the key resources table.

The following day, incubation with secondary antibodies (biotinylated or

fluorochrome-conjugated, Jackson, 1:200) for immunoperoxidase or

immunofluorescence staining was performed. After incubating in an avidin peroxidase

solution (Vectastain Elite ABC kit, PK-6100, Vector Laboratories), color development

was achieved by incubation with 3,3-diaminobenzidine (DAB, Sigma D5637) and

counterstaining with methyl green (Sigma, 67060). Slides were dehydrated, cover

slipped with DPX (Sigma), and images acquired using and Olympus BX51 microscope

with DP72 camera. Immunofluorescence slides were counterstained with 4',6'-

diamindino-2-phenylindole (DAPI), mounted with fluoromount, and imaged with an

Olympus BX51 microscope and captured with Olympus XM10 camera. Cell and axon

counting, as well as thresholding of images, was performed in ImageJ (NIH) without

knowledge of treatment groups using cropped regions of interest (Allen Spinal Cord

Atlas, Adobe Photoshop) and averaged from two sections per animal (60 μm apart).

Formalin-fixed paraffin-embedded 5 μm thick sections from confirmed MS and

normal CNS control autopsy cases were immunostained for CD38 (1:2000, LifeSpan

BioSciences , LS-C817841) . Steamed antigen retrieval was performed with DAKO high

pH target retrieval solution (K8004, Dako) (71).

RNA isolation and qRT-PCR

To determine gene expression from spinal cord or primary murine cells, total

RNA was extracted using RNA Stat60 (Tel-Test; Friendswood, TX) per manufacturer's

instructions and used for qRT-PCR using BioRad iTaq Universal Probes or primers and

SYBR Green One-step kits. Starting quantity for quantification of each gene's

expression was normalized to Rn18s or GAPDH. Detailed primer and probe information

are annotated in the key resources Table.

Mixed glial preparations and primary cell cultures

Primary mixed glial cultures were prepared from P0-3 mice, as previously described (4, 23). In brief, dissected cerebral cortices were dissociated and grown in DMEM (Fisher 11960-044) with 1 mM sodium pyruvate (Corning 11360070), 20 mM HEPES (Gibco 15630-080), 100 U/ml PenStrep (Life Technologies 15140122), 10% fetal bovine serum (FBS, Atlanta 337 Biologicals S11150), and 5 µg/ml insulin. Following 10-12 days in culture, sequential immunopanning was performed to separate astrocyte, microglia, and oligodendrocyte progenitor cells by agitating at 225 rpm on an orbital shaker (Innova 2000). Purity of the cell types cultured was within the ranges reported previously by our group (72, 73). After removing microglia via differential adhesion, purified oligodendrocyte progenitors were plated onto PLL-coated glass coverslips and cultured in differentiation medium (Neurobasal A media (Life Technologies, 10888022) with N2 (Gibco 17502048), B27 (Gibco 17504044), PenStrep, sodium pyruvate, 2 mM glutamax (Gibco 35050-061), 40 ng/ml T3, and 5% BSA (Sigma A4503)) for two days. Treatments with palmitate (PA, 100 µM) NAD$^+$ (50 µM), NMN (100 µM), EX527 (5 µM) and/or 78c (3 µM) were added and cultures were allowed to differentiate for 72 hr. Doses used in cell culture experiments were based on previous studies (30, 74) and preliminary dose-response experiments (data not shown). Positive staining for myelin proteins was compared to control levels by immunofluorescence staining and threshold quantification in ImageJ. Briefly, a positive threshold for each stain was decided, then applied to each image (5 per coverslip) to determine percentage of field positive. Mean values were graphed as a percentage of the control group.

In experiments characterizing astrocyte responses and effects of astrocyte-

conditioned media on oligodendrocytes, astrocytes were plated at a density of 450,000

cells/well on PLL-coated 6-well plates and grown for 2-3 days in serum-free Neurobasal

A media containing 1% N2, 2% B27, P/S, 1mM sodium pyruvate, 0.45% glucose, 5%

BSA, and 50μM β-mercaptoethanol before treating with or without palmitate (PA, 100

μM) and 78c (3 μM). Astrocytes were collected for an NAD$^+$ cycling assay or RNA

isolation. Supernatants from treated astrocytes were collected 24 h post-treatment and

concentrated using Vivaspin sample concentrators (10 and 100 kDa molecular weight

cutoffs, Sigma) and stored frozen as 10x protein concentrate until use for treating OLCs

(9, 23). Conditioned medium from astrocytes (ACM) was brought back to 1x

concentration in oligodendrocyte differentiation medium and placed on oligodendrocytes

to differentiate as described above.

NAD$^+$ cycling and CD38 activity assays

To determine intracellular NAD$^+$ levels, one well of a 6-well plate of astrocytes, or

approximately 20 mg of spinal cord tissue, was homogenized in 300 μl of 10%

trichloroacetic acid in water. Samples were extracted using an organic solvent mixture

(1,1,2-Trichloro-1,2,2-trifluoroethane and triocylamine), diluted, and measured along

with a standard curve, as previously described (30). Briefly, a cycling reagent (0.76%

ethanol, 4 μM FMN, 27.2 U/ml alcohol dehydrogenase, 0.44 U/ml diaphorase, and 8 μM

resazurin (Sigma)) was prepared in phosphate buffer (100 mM NaH$_2$PO$_4$/Na$_2$HPO$_4$, pH

8.0) and added to an equal volume of pre-diluted cell or tissue extract. Plates were

measured every 30 seconds for 1 h at 544 nm using a SpectraMax plate reader. Raw

values were compared to a standard curve, then divided by protein quantification

(Bradford reagent, Biorad) to be expressed as nmols/mg protein. For measuring CD38

NAD$^+$ase activity, spinal cord tissue was homogenized in NETN buffer (100 mM NaCl,

1 mM EDTA, 20 mM Tris-HCl, and 0.5% Nonidet P-40 (Sigma)) supplemented with 5

mM NaF (Sigma) and protease and phosphatase inhibitors (Thermo Fisher) (30). After

centrifugation (12000 rpm, 10 min), supernatant volumes were adjusted to protein

concentrations and reaction mix containing 50 µM 1,N6-Etheno NAD$^+$ in 0.25 M

sucrose buffer was added at an equal volume just prior to reading the plate at 300 nm

excitation and 410 nm emission every minute for 1 hr on a Spark Magellan plate reader.

Bioplex assay for cytokines

Astrocyte supernatants from cell treatments described above were used for

determination of secreted cytokine levels using a Bio-Plex Mouse cytokine 23-plex

assay (BioRad m60009rdpd) per manufacturer instructions. Briefly, post-treatment

medium was collected and incubated with primary antibody-conjugated magnetic beads

for 30 min. Wells were washed thrice and then incubated for 30 min with detection

antibodies. After washing, samples were incubated for 10 min with a streptavidin-

phycoerythrin solution, and then washed again prior to resuspension and reading

fluorescence with a BioPlex 200 (BioRad) reader and Bio-Plex Manager software.

Organotypic cerebellar slice culture and demyelination

Organotypic cerebellar slice cultures were cultured and demyelinated using

previously described methods (8). Cerebella were dissected from postnatal day 7-9

pups, embedded in low-melting point agar (Invitrogen 16520050), and 350 µm slices

collected into artificial CSF solution using a McIlwain tissue chopper (Mickle Laboratory

Engineering Co; Guildford, UK). Four slices were transferred and grown per each

Millicell insert (Sigma PIHP03050) in 50% MEM, 22.5% Hank's Balanced Salt Solution

(HBSS), 24% heat-inactivated (HI) Horse Serum, 1.5% Glucose, 1% GlutaMax, 1% P/S

medium.

Demyelination was induced after 7 d in culture by treating with 0.5 mg/ml

lysophosphatidycholine (LL, Sigma) for 18 h. Organotypic slices were placed back in

LL-free medium and collected for immunohistochemistry and confocal imaging after 24

h (demyelination) or 7 d (myelin regeneration) with or without addition of PA (100 µM)

and/or 78c (3 µM) added to the growth media. Confocal images were acquired at 20x

magnification from 5 regions randomly selected per slice using a Zeiss LSM 780 and

Zen Software in the Mayo Clinic Microscopy and Cell Analysis Core. Images were

thresholded for each stain, and stain-positive immunoreactivity area or cell counts per

total field area were quantified in ImageJ.

Statistics

Data were graphed with bars representing mean ± SEM with sample size for

each experiment indicated in the figure legends. To determine sample size, power

analysis using the fpower macro in SAS was used to determine animal numbers for

each endpoint based on 80% power, $\alpha=0.05$, and 4 treatment groups. Differences and

SD used to determine δ were used from analysis of previous similarly-designed studies

in our laboratory.   Cell culture studies used a combination of male and female pups,

were repeated twice, and assays were run in technical duplicate or triplicate. No

statistical outliers were removed, but the ROUT test in GraphPad Prism was used to

check for outliers and data normality was assessed using the Shapiro-Wilk test.

Comparisons between two groups were analyzed by Student's t-test, whereas one-way

or two-way ANOVAs with Bonferroni post-test were applied for multiple-comparisons.

Analysis was performed in GraphPad Prism 7.0 software with p-values ≤ 0.05

considered significant.

Study Approval

All animal experiments were approved by the IACUC and the human pathology

study was approved by the Institutional Review Board of Mayo Clinic, Rochester, MN

(IRB 2067-99).

Author Contributions

IAS and MRL conceived and designed the study. MRL, CC, and TRPM

conducted *in vivo* experiments and necropsies. Embedding and histological staining of

mouse tissues was performed by MRL and LK. Image acquisition and quantifications

were performed by MRL under guidance from CC. YG and CL performed and analyzed

human histological staining. MRL completed *in vitro* assays and qRT-PCR. WS and HY

assisted with glial preparations and organotypic slice cultures. TRP, CCSC, and ENC

provided expertise in NAD^+ quantification and experimental design as well as CD38

mice. MRL and IAS wrote the manuscript and all authors reviewed and edited the

manuscript.

Acknowledgements

The research was supported by the Mayo Clinic Center for Multiple Sclerosis and

Autoimmune Neurology, the Eugene and Marcia Applebaum Foundation, and the Mayo

Clinic Center for Biomedical Discovery. Portions of this work were supported by the NIH

R01NS052741-10, NIA R01AG058812, the National Multiple Sclerosis Society (G-1508-

05951, RG-1901-33209 and FG-1908-34819) and the Minnesota State Spinal Cord

Injury and Traumatic Brain Injury Research Program. 78c-containing chow was

generously provided by Calico Life Sciences LLC. Diagrams were created using

elements from the Biomedical-PPT-Toolkit-Suite (Motifolio).

References

References

1.   Gianfrancesco MA, Acuna B, Shen L, Briggs FB, Quach H, Bellesis KH, et al.

Obesity during childhood and adolescence increases susceptibility to multiple

sclerosis after accounting for established genetic and environmental risk factors.

Obes Res Clin Pract. 2014;8(5):e435-47.

2.   Olsson T, Barcellos LF, and Alfredsson L. Interactions between genetic, lifestyle

and environmental risk factors for multiple sclerosis. *Nat Rev Neurol.*

2017;13(1):25-36.

3.   Castro K, Ntranos A, Amatruda M, Petracca M, Kosa P, Chen EY, et al. Body

Mass Index in Multiple Sclerosis modulates ceramide-induced DNA methylation

and disease course. *EBioMedicine.* 2019;43:392-410.

4.   Langley MR, Yoon H, Kim HN, Choi CI, Simon W, Kleppe L, et al. High fat diet

consumption results in mitochondrial dysfunction, oxidative stress, and

oligodendrocyte loss in the central nervous system. *Biochim Biophys Acta Mol*

Basis Dis. 2019:165630.

5.   Graham LC, Grabowska WA, Chun Y, Risacher SL, Philip VM, Saykin AJ, et al.

Exercise prevents obesity-induced cognitive decline and white matter damage in

mice. *Neurobiol Aging.* 2019;80:154-72.

6.   Yoon H, Kleven A, Paulsen A, Kleppe L, Wu J, Ying Z, et al. Interplay between

exercise and dietary fat modulates myelinogenesis in the central nervous system.

Biochim Biophys Acta. 2016;1862(4):545-55.

7.   Buckman LB, Thompson MM, Lippert RN, Blackwell TS, Yull FE, and Ellacott KL.

Evidence for a novel functional role of astrocytes in the acute homeostatic

response to high-fat diet intake in mice. *Mol Metab.* 2015;4(1):58-63.

8.   Kim HN, Langley MR, Simon WL, Yoon H, Kleppe L, Lanza IR, et al. A Western

diet impairs CNS energy homeostasis and recovery after spinal cord injury: Link

to astrocyte metabolism. *Neurobiol Dis.* 2020;141:104934.

9.   Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, et

al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature.*

2017;541(7638):481-7.

10. Dimas P, Montani L, Pereira JA, Moreno D, Trotzmuller M, Gerber J, et al. CNS

myelination and remyelination depend on fatty acid synthesis by

oligodendrocytes. *Elife.* 2019;8.

11. Ponath G, Park C, and Pitt D. The Role of Astrocytes in Multiple Sclerosis. *Front*

Immunol. 2018;9:217.

12. Hasan M, Seo JE, Rahaman KA, Min H, Kim KH, Park JH, et al. Novel genes in

brain tissues of EAE-induced normal and obese mice: Upregulation of metal ion-

binding protein genes in obese-EAE mice. *Neuroscience.* 2017;343:322-36.

13. Timmermans S, Bogie JF, Vanmierlo T, Lutjohann D, Stinissen P, Hellings N, et

al. High fat diet exacerbates neuroinflammation in an animal model of multiple

sclerosis by activation of the Renin Angiotensin system. *J Neuroimmune*

Pharmacol. 2014;9(2):209-17.

14. Ji Z, Wu S, Xu Y, Qi J, Su X, and Shen L. Obesity Promotes EAE Through IL-6

and CCL-2-Mediated T Cells Infiltration. *Front Immunol.* 2019;10:1881.

15.   Herrmann MM, Barth S, Greve B, Schumann KM, Bartels A, and Weissert R.

Identification of gene expression patterns crucially involved in experimental

autoimmune encephalomyelitis and multiple sclerosis. *Dis Model Mech.*

2016;9(10):1211-20.

16.   Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, et al. An

RNA-sequencing transcriptome and splicing database of glia, neurons, and

vascular cells of the cerebral cortex. *J Neurosci.* 2014;34(36):11929-47.

17.   Zhang Y, Sloan SA, Clarke LE, Caneda C, Plaza CA, Blumenthal PD, et al.

Purification and Characterization of Progenitor and Mature Human Astrocytes

Reveals Transcriptional and Functional Differences with Mouse. *Neuron.*

2016;89(1):37-53.

18.   Roboon J, Hattori T, Ishii H, Takarada-Iemata M, Le TM, Shiraishi Y, et al.

Deletion of CD38 Suppresses Glial Activation and Neuroinflammation in a Mouse

Model of Demyelination. *Front Cell Neurosci.* 2019;13:258.

19.   Aksoy P, White TA, Thompson M, and Chini EN. Regulation of intracellular levels

of NAD: a novel role for CD38. *Biochem Biophys Res Commun.*

2006;345(4):1386-92.

20.   Aksoy P, Escande C, White TA, Thompson M, Soares S, Benech JC, et al.

Regulation of SIRT 1 mediated NAD dependent deacetylation: a novel role for

the multifunctional enzyme CD38. *Biochem Biophys Res Commun.*

2006;349(1):353-9.

21.   Guerreiro S, Privat AL, Bressac L, and Toulorge D. CD38 in Neurodegeneration

and Neuroinflammation. *Cells.* 2020;9(2).

22.   Chiang SH, Harrington WW, Luo G, Milliken NO, Ulrich JC, Chen J, et al. Genetic

Ablation of CD38 Protects against Western Diet-Induced Exercise Intolerance

and Metabolic Inflexibility. *PLoS One.* 2015;10(8):e0134927.

23.   Yoon H, Choi CI, Triplet EM, Langley MR, Kleppe LS, Kim HN, et al. Blocking the

Thrombin Receptor Promotes Repair of Demyelinated Lesions in the Adult Brain.

J Neurosci. 2020;40(7):1483-500.

24.   Ogrodnik M, Zhu Y, Langhi LGP, Tchkonia T, Kruger P, Fielder E, et al. Obesity-

Induced Cellular Senescence Drives Anxiety and Impairs Neurogenesis. *Cell*

Metab. 2019.

25.   Domingues HS, Portugal CC, Socodato R, and Relvas JB. Oligodendrocyte,

Astrocyte, and Microglia Crosstalk in Myelin Development, Damage, and Repair.

Front Cell Dev Biol. 2016;4:71.

26.   Kiray H, Lindsay SL, Hosseinzadeh S, and Barnett SC. The multifaceted role of

astrocytes in regulating myelination. *Exp Neurol.* 2016;283(Pt B):541-9.

27.   Camacho-Pereira J, Tarrago MG, Chini CCS, Nin V, Escande C, Warner GM, et

al. CD38 Dictates Age-Related NAD Decline and Mitochondrial Dysfunction

through an SIRT3-Dependent Mechanism. *Cell Metab.* 2016;23(6):1127-39.

28.   Hattori T, Kaji M, Ishii H, Jureepon R, Takarada-Iemata M, Minh Ta H, et al.

CD38 positively regulates postnatal development of astrocytes cell-autonomously

and oligodendrocytes non-cell-autonomously. *Glia.* 2017;65(6):974-89.

29.   Kou W, Banerjee S, Eudy J, Smith LM, Persidsky R, Borgmann K, et al. CD38

regulation in activated astrocytes: implications for neuroinflammation and HIV-1

brain infection. *J Neurosci Res.* 2009;87(10):2326-39.

30.   Tarrago MG, Chini CCS, Kanamori KS, Warner GM, Caride A, de Oliveira GC, et

al. A Potent and Specific CD38 Inhibitor Ameliorates Age-Related Metabolic

Dysfunction by Reversing Tissue NAD(+) Decline. *Cell Metab.* 2018;27(5):1081-

95 e10.

31.   Clarke LE, Liddelow SA, Chakraborty C, Munch AE, Heiman M, and Barres BA.

Normal aging induces A1-like astrocyte reactivity. *Proc Natl Acad Sci U S A.*

2018;115(8):E1896-E905.

32.   Lassmann H, and van Horssen J. Oxidative stress and its impact on neurons and

glia in multiple sclerosis lesions. *Biochim Biophys Acta.* 2016;1862(3):506-10.

33.   Liu Y, Guo Y, Huang W, Deng KY, Qian Y, and Xin HB. 17beta-Estradiol

Promotes Apoptosis in Airway Smooth Muscle Cells Through CD38/SIRT1/p53

Pathway. *Front Endocrinol (Lausanne).* 2018;9:770.

34.   Haffner CD, Becherer JD, Boros EE, Cadilla R, Carpenter T, Cowan D, et al.

Discovery, Synthesis, and Biological Evaluation of Thiazoloquin(az)olin(on)es as

Potent CD38 Inhibitors. *J Med Chem.* 2015;58(8):3548-71.

35.   Alizadeh A, and Karimi-Abdolrezaee S. Microenvironmental regulation of

oligodendrocyte replacement and remyelination in spinal cord injury. *J Physiol.*

2016;594(13):3539-52.

36.   Harlan BA, Pehar M, Sharma DR, Beeson G, Beeson CC, and Vargas MR.

Enhancing NAD+ Salvage Pathway Reverts the Toxicity of Primary Astrocytes

Expressing Amyotrophic Lateral Sclerosis-linked Mutant Superoxide Dismutase 1

(SOD1). *J Biol Chem.* 2016;291(20):10836-46.

37.   Stein LR, and Imai S. Specific ablation of Nampt in adult neural stem cells

recapitulates their functional defects during aging. *EMBO J.* 2014;33(12):1321-

40.

38.   Chini C, Hogan KA, Warner GM, Tarrago MG, Peclat TR, Tchkonia T, et al. The

NADase CD38 is induced by factors secreted from senescent cells providing a

potential link between senescence and age-related cellular NAD(+) decline.

Biochem Biophys Res Commun. 2019;513(2):486-93.

39.   Penberthy WT, and Tsunoda I. The importance of NAD in multiple sclerosis. *Curr*

Pharm Des. 2009;15(1):64-99.

40.   Penke M, Schuster S, Gorski T, Gebhardt R, Kiess W, and Garten A. Oleate

ameliorates palmitate-induced reduction of NAMPT activity and NAD levels in

primary human hepatocytes and hepatocarcinoma cells. *Lipids Health Dis.*

2017;16(1):191.

41.   Ponath G, Ramanan S, Mubarak M, Housley W, Lee S, Sahinkaya FR, et al.

Myelin phagocytosis by astrocytes after myelin damage promotes lesion

pathology. *Brain.* 2017;140(2):399-413.

42.   Machado-Santos J, Saji E, Troscher AR, Paunovic M, Liblau R, Gabriely G, et al.

The compartmentalized inflammatory response in the multiple sclerosis brain is

composed of tissue-resident CD8+ T lymphocytes and B cells. *Brain.*

2018;141(7):2066-82.

43.   Chini EN, Chini CCS, Espindola Netto JM, de Oliveira GC, and van Schooten W.

The Pharmacology of CD38/NADase: An Emerging Target in Cancer and

Diseases of Aging. *Trends Pharmacol Sci.* 2018;39(4):424-36.

44.   Bruzzone S, Verderio C, Schenk U, Fedele E, Zocchi E, Matteoli M, et al.

859   Glutamate-mediated overexpression of CD38 in astrocytes cultured with

860   neurones. *J Neurochem.* 2004;89(1):264-72.

861   45.   Mamik MK, Banerjee S, Walseth TF, Hirte R, Tang L, Borgmann K, et al. HIV-1

862   and IL-1beta regulate astrocytic CD38 through mitogen-activated protein kinases

863   and nuclear factor-kappaB signaling mechanisms. *J Neuroinflammation.*

864   2011;8:145.

865   46.   Boven LA, Montagne L, Nottet HS, and De Groot CJ. Macrophage inflammatory

866   protein-1alpha (MIP-1alpha), MIP-1beta, and RANTES mRNA semiquantification

867   and protein expression in active demyelinating multiple sclerosis (MS) lesions.

868   *Clin Exp Immunol.* 2000;122(2):257-63.

869   47.   Mori F, Nistico R, Nicoletti CG, Zagaglia S, Mandolesi G, Piccinin S, et al.

870   RANTES correlates with inflammatory activity and synaptic excitability in multiple

871   sclerosis. *Mult Scler.* 2016;22(11):1405-12.

872   48.   Amici SA, Young NA, Narvaez-Miranda J, Jablonski KA, Arcos J, Rosas L, et al.

873   CD38 Is Robustly Induced in Human Macrophages and Monocytes in

874   Inflammatory Conditions. *Front Immunol.* 2018;9:1593.

875   49.   Zou Y, He X, Peng QY, and Guo QL. Inhibition of CD38/Cyclic ADP-ribose

876   Pathway Protects Rats against Ropivacaine-induced Convulsion. *Chin Med J*

877   *(Engl).* 2017;130(19):2354-60.

878   50.   Wang C, Zhang Y, Ding J, Zhao Z, Qian C, Luan Y, et al. Nicotinamide

879   Administration Improves Remyelination after Stroke. *Neural Plast.*

880   2017;2017:7019803.

51. Hathorn T, Snyder-Keller A, and Messer A. Nicotinamide improves motor deficits

and upregulates PGC-1alpha and BDNF gene expression in a mouse model of

Huntington's disease. *Neurobiol Dis.* 2011;41(1):43-50.

52. Ginwala R, McTish E, Raman C, Singh N, Nagarkatti M, Nagarkatti P, et al.

Apigenin, a Natural Flavonoid, Attenuates EAE Severity Through the Modulation

of Dendritic Cell and Other Immune Cell Functions. *J Neuroimmune Pharmacol.*

2016;11(1):36-47.

53. Wang J, Zhao C, Kong P, Sun H, Sun Z, Bian G, et al. Treatment with NAD(+)

inhibited experimental autoimmune encephalomyelitis by activating AMPK/SIRT1

signaling pathway and modulating Th1/Th17 immune responses in mice. *Int*

Immunopharmacol. 2016;39:287-94.

54. Ji S, Doucette JR, and Nazarali AJ. Sirt2 is a novel in vivo downstream target of

Nkx2.2 and enhances oligodendroglial cell differentiation. *J Mol Cell Biol.*

2011;3(6):351-9.

55. Jablonska B, Gierdalski M, Chew LJ, Hawley T, Catron M, Lichauco A, et al. Sirt1

regulates glial progenitor proliferation and regeneration in white matter after

neonatal brain injury. *Nat Commun.* 2016;7:13866.

56. Li J, Dou X, Li S, Zhang X, Zeng Y, and Song Z. Nicotinamide ameliorates

palmitate-induced ER stress in hepatocytes via cAMP/PKA/CREB pathway-

dependent Sirt1 upregulation. *Biochim Biophys Acta.* 2015;1853(11 Pt A):2929-

36.

57.   Prola A, Pires Da Silva J, Guilbert A, Lecru L, Piquereau J, Ribeiro M, et al.

903   SIRT1 protects the heart from ER stress-induced cell death through eIF2alpha

904   deacetylation. *Cell Death Differ.* 2017;24(2):343-56.

905   58.   Sadeghi A, Seyyed Ebrahimi SS, Golestani A, and Meshkani R. Resveratrol

906   Ameliorates Palmitate-Induced Inflammation in Skeletal Muscle Cells by

907   Attenuating Oxidative Stress and JNK/NF-kappaB Pathway in a SIRT1-

908   Independent Mechanism. *J Cell Biochem.* 2017;118(9):2654-63.

909   59.   Wilms H, Sievers J, Rickert U, Rostami-Yazdi M, Mrowietz U, and Lucius R.

910   Dimethylfumarate inhibits microglial and astrocytic inflammation by suppressing

911   the synthesis of nitric oxide, IL-1beta, TNF-alpha and IL-6 in an in-vitro model of

912   brain inflammation. *J Neuroinflammation.* 2010;7:30.

913   60.   Choi JW, Gardell SE, Herr DR, Rivera R, Lee CW, Noguchi K, et al. FTY720

914   (fingolimod) efficacy in an animal model of multiple sclerosis requires astrocyte

915   sphingosine 1-phosphate receptor 1 (S1P1) modulation. *Proc Natl Acad Sci U S*

916   *A.* 2011;108(2):751-6.

917   61.   van de Donk N, Richardson PG, and Malavasi F. CD38 antibodies in multiple

918   myeloma: back to the future. *Blood.* 2018;131(1):13-29.

919   62.   Pavelek Z, Vysata O, Sobisek L, Klimova B, Andrys C, Vokurkova D, et al.

920   Lymphocyte populations and their change during five-year glatiramer acetate

921   treatment. *Neurol Neurochir Pol.* 2018;52(5):587-92.

922   63.   Hoffmann FS, Hofereiter J, Rubsamen H, Melms J, Schwarz S, Faber H, et al.

923   Fingolimod induces neuroprotective factors in human astrocytes. *J*

924   *Neuroinflammation.* 2015;12:184.

64.   Deckert J, Wetzel MC, Bartle LM, Skaletskaya A, Goldmacher VS, Vallee F, et al.

SAR650984, a novel humanized CD38-targeting antibody, demonstrates potent

antitumor activity in models of multiple myeloma and other CD38+ hematologic

malignancies. *Clin Cancer Res.* 2014;20(17):4574-83.

65.   Lokhorst HM, Plesner T, Laubach JP, Nahi H, Gimsing P, Hansson M, et al.

Targeting CD38 with Daratumumab Monotherapy in Multiple Myeloma. *N Engl J*

Med. 2015;373(13):1207-19.

66.   Martin T, Baz R, Benson DM, Lendvai N, Wolf J, Munster P, et al. A phase 1b

study of isatuximab plus lenalidomide and dexamethasone for relapsed/refractory

multiple myeloma. *Blood.* 2017;129(25):3294-303.

67.   Partida-Sanchez S, Cockayne DA, Monard S, Jacobson EL, Oppenheimer N,

Garvy B, et al. Cyclic ADP-ribose production by CD38 regulates intracellular

calcium release, extracellular calcium influx and chemotaxis in neutrophils and is

required for bacterial clearance in vivo. *Nat Med.* 2001;7(11):1209-16.

68.   Ghoumari AM, Abi Ghanem C, Asbelaoui N, Schumacher M, and Hussain R.

Roles of Progesterone, Testosterone and Their Nuclear Receptors in Central

Nervous System Myelination and Remyelination. *Int J Mol Sci.* 2020;21(9).

69.   Keough MB, Jensen SK, and Yong VW. Experimental demyelination and

remyelination of murine spinal cord by focal injection of lysolecithin. *J Vis Exp.*

2015(97).

70.   Pavelko KD, van Engelen BG, and Rodriguez M. Acceleration in the rate of CNS

remyelination in lysolecithin-induced demyelination. *J Neurosci.* 1998;18(7):2498-

505.

71.   Chuang TY, Guo Y, Seki SM, Rosen AM, Johanson DM, Mandell JW, et al. LRP1

expression in microglia is protective during CNS autoimmunity. *Acta Neuropathol*

Commun. 2016;4(1):68.

72.   Burda JE, Radulovic M, Yoon H, and Scarisbrick IA. Critical role for PAR1 in

kallikrein 6-mediated oligodendrogliopathy. *Glia.* 2013;61(9):1456-70.

73.   Radulovic M, Yoon H, Wu J, Mustafa K, and Scarisbrick IA. Targeting the

thrombin receptor modulates inflammation and astrogliosis to improve recovery

after spinal cord injury. *Neurobiol Dis.* 2016;93:226-42.

74.   Valle C, Salvatori I, Gerbino V, Rossi S, Palamiuc L, Rene F, et al. Tissue-

specific deregulation of selected HDACs characterizes ALS progression in

mouse models: pharmacological characterization of SIRT1 and SIRT2 pathways.

Cell Death Dis. 2014;5:e1296.

Figures and Tables

Figure 1. CD38 expression in astrocytes of chronic HFD-fed mice and in a model of demyelination.

A Schematic representation of previously shown direct effects of high fat diet (HFD) on

oligodendrocyte lineage cells including apoptosis and impaired differentiation with

potential astrocyte-mediated indirect effects, represented by dotted lines, that comprise

the hypothesis to be tested.

B Spinal cord tissue of regular diet (RD) and high fat diet (HFD) mice showing a

depletion of NAD^+ after 12 wk (n=5 mice/group) as determined by cycling assay.

C Increased RNA expression of CD38 (n=4), assessed by qPCR, in spinal cord tissue of

HFD-fed mice when compared to RD-fed mice.

D- E Immunofluorescence staining with double-labeling and corresponding

quantification of immunoreactivity and cell counts indicates increased CD38 (green)

immunoreactivity in astrocytes (GFAP, red) within the spinal cords of HFD-fed mice (**D**,

n=7) or in the lysolecithin (LL) model of experimental demyelination (**E**, n=4)versus the

uninjected (UI) side. Broken white line designates lesion boundary. Scale bar = 50 µm

(**D**) and 100µm (**E**).

F-H High fat consumption was modeled in primary murine astrocyte cultures by

including saturated fat palmitate in the media (PA, 100 µM, 24 h). Increases in RNA

expression of GFAP (**F,** n=4) and CD38 (**G,** n=6) were observed in PA-containing

astrocyte cultures. **H**, Depleted NAD^+ levels were found in PA-treated astrocytes when

assessed using an NAD^+ cycling assay, n=7-8.

I Representative microscopic images of CD38 staining in control brain tissue and an MS

lesion with active demyelinating activity. Scale bar = 50 µm.

Bar graphs represent the mean +/- SEM. Asterisks indicate levels of statistical

significance as determined by Student's t-test, where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

Figure 2. **CD38-catalytically inactive (CD38ci) mice are resistant to HFD-induced**

oligodendrocyte loss.

Histological analysis from wildtype (WT) and CD38-catalytically inactive mice (CD38ci)

fed a regular diet (RD) or high fat diet (HFD) for 30 wk beginning at 50 wk of age.

A Schematic depicting experimental design.

B NAD^+ levels as determined by cycling assay indicate reductions in WT-HFD mice

compared to WT-RD and increased NAD^+ levels in the CD38ci mice versus WT.

C CD38 activity levels by etheno-NAD^+ assay show impaired activity in CD38ci mice

versus WTs and higher activity in WT-HFD mice than WT-RD (n=5/group).

D, Outlined regions depict regions imaged, including dorsal column (broken gray line)

and ventral white matter (broken black line).

E PDGFRα double immunofluorescent staining with Ki67 or cleaved-caspase-3 shows a

depletion of oligodendrocyte progenitor cells with HFD that was significantly attenuated

in CD38ci mice versus wildtype. Increased cleaved-caspase-3 and PDGFRα co-labeled

cells were found in HFD versus RD WT mice, while fewer cleaved-caspase-3 and

PDGFRα co-labeled cells we observed in the CD38ci HFD than WT HFD ventral WM.

F-G $Olig2^+$ oligodendrocyte lineage cells (**F**) and $GST3^+$ mature oligodendrocytes (**G**)

were significantly depleted in HFD-fed WT mice, yet CD38ci mice fed a HFD had no

reduction in oligodendrocyte numbers. When compared to WT-HFD mice, CD38ci-HFD

mice had significantly more oligodendrocytes remaining in the dorsal column

(representative images pictured). Scale bar = 50 µm. Gray lines indicate genotype

means, while gray broken lines indicate comparison of overall genotype effect.

H Representative images of immunohistochemical staining confirming no significant

changes in MBP immunoreactivity across genotype or diet groups. Scale bar= 100 µm.

Bar graphs represent mean +/- SEM. Two-way ANOVA followed by Bonferroni post-hoc

test (n=7-10 mice/group). Asterisks indicate levels of statistical significance when

compared to control, where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

1019

<u>Figure 3.</u> Diminished neuroinflammatory responses in CD38 catalytically inactive

(CD38ci) mice following chronic HFD.

Histological analysis from wildtype (WT) and CD38-catalytically inactive mice (CD38ci)

fed a regular diet (RD) or high fat diet (HFD) for 30 wk beginning at 50 wk of age. Scale

bar= 50 μm. and n=7-10 mice/group.

A Representative images of the ventral white matter and immunohistochemical analysis

of spinal cord regions demonstrates decreased microgliosis (IBA1) in CD38ci mice.

When compared to WT-HFD mice, CD38ci-HFD mice had significantly less IBA1$^+$

immunoreactivity in the spinal cord ventral gray and white matter.

B Representative images of the ventral white matter and corresponding quantification

confirm reduced astrogliosis (GFAP) in the spinal cords of CD38 catalytically inactive

(CD38ci) mice overall when compared to wildtype (WT) mice. Decreased GFAP$^+$

immunoreactivity in the ventral white matter when compared to HFD mice.

C Representative images of the dorsal column and quantification of the spinal cords

confirming significant changes in 4HNE immunoreactivity, an oxidative stress marker,

between genotypes for HFD mice.

D-F qRT-PCRs data indicting differences in antioxidant (F) and pro-repair astrocyte

marker genes (G-H, BDNF and S100a10, respectively) in the CD38ci mice versus

controls. Bar graphs represent mean +/- SEM. Two-way ANOVA with Bonferroni post-

test. Asterisks indicate levels of statistical significance, where *, $p<0.05$ and **, $p<0.01$.

Gray lines on graphs indicate genotype means, while gray broken lines with asterisks

indicate a significant effect of genotype overall.

1042

1043

Figure 4. CD38 inhibition with 78c improves myelin repair following lysolecithin-

induced focal demyelination to the ventral spinal cord.

Eight week old male C57Bl6 mice were fed a regular diet (RD) or high fat diet (HFD) for

5 wks, then split into two additional groups: one receiving the same diet and the other

half receiving the respective diet with 78c added (600 ppm). One week later, a focal

demyelinating lesion to the ventral spinal cord was induced by lysolecithin (LL, 1% with

Evans Blue dye) and mice were allowed to recover for two weeks. Lesion boundary

regions were determined by limiting to the area of continuous myelin loss (core) or by

including the entire inflammatory lesion (rim) based on immunohistochemical analyses.

A Schematic depicting experimental design and lesion boundaries.

B Representative images and corresponding analysis illustrating the mean number of

remyelinated axons after focal lysolecithin-mediated demyelination of the ventral spinal

cord was increased in 78c-treated mice at 14 days post injury when compared to LL

only mice (**, $p<0.01$, two-way ANOVA with Bonferroni post-test) as assessed by

paraphenylenediamine (PPD)-stained semi-thin sections of araldite-embedded spinal

cord tissue. Scale bar= 20 µm.

C Immunohistochemical staining of the total lesion and lesion rim identified significant

changes in MBP immunoreactivity in 78c treated animals.

D-E Olig2$^+$ oligodendrocyte lineage cells (OLCs, **D**) and GST3$^+$ mature oligodendrocytes

(**E**) were significantly higher in 78c-RD mice when compared to RD controls.

Bar graphs represent mean +/- SEM. Two-way ANOVA (n=7-10 mice/group). *, $p<0.05$

and **, $p<0.01$. Gray lines on graphs indicate genotype means, while gray broken lines

indicate a significant effect of genotype overall.

1067

1068

<u>Figure 5.</u> 78c promotes pro-repair astrocytes after lysolecithin-induced

demyelination.

Histological analysis of markers of neuroinflammation in animals fed a regular diet (RD)

or high fat diet (HFD) with or without 78c (CD38 inhibitor, 600 ppm) and injected with

lysolecithin (LL) followed by a 2-week recovery period.

A, B IBA1 (**A**, microgliosis) and GFAP (**B**, astrogliosis) were not significantly affected by

either diet or 78c alone, although a significant interaction for drug and diet was detected

for IBA1 immunoreactivity in the lesion rim. Scale bar= 100 microns.

C-D, Changes in and pro-repair (S100a10, brown) pro-inflammatory (C3d, red)

astrocyte (GFAP, green) markers in adjacent sections within the LL lesion area indicates

a shift toward pro-repair in the 78c-treated mice. Scale bar = 100 μm.

Bar graphs represent mean +/- SEM. Two-way ANOVA (n=5-8 mice/group). Asterisks

indicate levels of statistical significance, where **, $p < 0.01$ when compared to LL-RD by

Bonferroni post-test. Gray lines on graphs indicate genotype means, while gray broken

lines with asterisks indicate a significant effect of genotype overall.

1084

1085

<u>**Figure 6.**</u> **Astrocytes cultured in high saturated fat conditions increase CD38 and**

pro-inflammatory cytokine expression and inhibit oligodendrocyte differentiation.

A Schematic depicting how astrocyte conditioned media (ACM) treatments on primary

murine oligodendrocytes were performed. Conditioned media obtained from PA-treated

murine astrocytes (ACM (PA)) reduced protein expression of mature oligodendrocyte

markers PLP and MBP, while PA-78c-ACM-treated OLCs have enhanced PLP and MBP

expression.

B Representational diagram of astrocyte treatments with PA and NAD^+ to collect ACM

for OPC treatments. The addition of NAD^+ restored PLP levels when compared to ACM-

PA, but had no effect on Olig2 . Scale bar = 100µm.

C-E High fat consumption was modeled in primary murine astrocyte cultures by

including saturated fat palmitate in the media (PA, 100 µM, 24 h). **C**, Addition of a small

molecule CD38 inhibitor (78c, 3µM) was effective to restore NAD^+ levels in PA-treated

astrocytes when assessed using an NAD^+ cycling assay. Astrocytes derived from CD38

knockout mice did not show depleted NAD^+ levels in response to PA. RNA levels by

qPCR (**D**) of proinflammatory cytokines, antioxidant enzyme, and an astrocyte activation

marker in PA and 78c treated astrocytes and cytokine levels (**E***)* detected in astrocyte

conditioned media (ACM) by Bioplex assay.

Bar graphs represent the mean +/- SEM. Asterisks indicate levels of statistical

significance when compared to control as determined by or one-way ANOVA with

Bonferroni post-test , where *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$ versus control.

1107

1108

Figure 7. NAD⁺ replenishment attenuates PA-induced deficits in oligodendrocyte

differentiation in a SIRT1-dependent manner.

A Representational diagram of experimental treatments in primary murine

oligodendrocyte lineage cells (OLCs). Adding exogenous NAD⁺ (50 µM) to

oligodendrocyte cultures significantly restores PLP protein expression (**B**) in PA-

exposed OLCs.

C Treatment with NAD$^+$ precursor nicotinamide mononucleotide (NMN,100 μM) also

enhances expression of mature oligodendrocyte markers in PA-treated cultures,

although not when SIRT1 inhibitor Ex-527 (5 μM) is also included in the media, as

determined by immunostaining quantification for PLP and MBP per DAPI$^+$ cell . Scale

bar= 100 μm.

D Schematic summarizing direct and indirect effects of palmitate on oligodendrocytes

characterized *in vitro*.

Graphical results represented as the mean ± SEM (n=4-8/group). Asterisks indicate

levels of statistical significance when compared to designated group as determined by

one-way ANOVA with Bonferroni post-test, where *, $p<0.05$; **, $p<0.01$; ***, $p<0.001$.

1125

1126

<u>**Figure 8.**</u> **CD38 inhibition protects from saturated fat and lysolecithin-induced**

demyelination in cerebellar slices

A-B Representational diagram (**A**) of *ex vivo* remyelination experiments where murine

organotypic cerebellar slices were demyelinated with lysolecithin (LL, 0.5 mg/ml,

overnight) followed by subsequent culture in control media for 1 wk. Confocal

microscopic imaging of slices indicate reduced MBP (red, **B**) expression when

normalized to neurofilament (NF, green). LL-demyelinated slices cultured with 78c had

significantly increased MBP expression when normalized to NF staining and expressed

as a percentage of the control group.

C Slices cultured in media containing both LL-palmitic acid (PA, 100 μM) and 78c, a

CD38 inhibitor, 3 μM) had significant improvements in Olig2 and MBP expression

compared to the LL-PA alone group.

Bar graphs represent the mean ± SEM (n=4-8/group); *, $p < 0.05$, **, $p < 0.01$, and ***,

$p < 0.001$ when compared to control by one-way ANOVA with Bonferroni post-test.

1141

1142

Figure 9. **Schematic representation of proposed hypothesis.**

Resting astrocytes typically help to provide a supportive environment for

oligodendrocyte differentiation and repair, when necessary. High fat diet (HFD) and

lysolecithin (LL) increase CD38 expression in reactive astrocytes, thereby depleting

NAD$^+$ levels. Reductions in NAD$^+$ result in oxidative stress and inflammation which

triggers impaired differentiation and survival in oligodendrocyte lineage cells. 78c

inhibits CD38 to attenuate the described detrimental effects of astrocytes on

oligodendrocytes.

Reagent or Resource	Source	Additional information
Antibodies		
Mouse anti-CD38	Novus	1g7f4 (1:500)
Rabbit anti-GFAP	Dako	Z0334 (1:5000)
mouse anti-Olig2	Millipore	MABN50(1:200)
rabbit anti-Olig2	Millipore	Ab9610; AB_10141047 (1:500)
Rabbit anti-GST3	Abcam	Ab153949 (1:1000)
rat anti-Myelin basic protein (MBP)	Millipore	MAB386; AB_94975 (1:750)
Rabbit anti-PLP	Abcam	A28486 (1:750)
Rabbit anti-IBA1	Wako	019-19741(1:3000)
rabbit anti-4-hydroxynonenol	Abcam	ab46545 (1:500)
chicken anti-GFAP	Abcam	ab4647 (1:2000)
Goat anti-C3d	R&D Systems	AF2655 (1:500)
Rabbit anti-S100a10	Proteintech	11250-1-AP (1:7500)
biotinylated goat anti-rat	Jackson	112-066-072 (1:100)
biotinylated donkey anti-rabbit	Jackson	711-065-152 (1:100)
biotinylated donkey anti-mouse	Jackson	715-066-151 (1:100)
goat-anti-rabbit AF488	Jackson	111-545-047 (1:100)
goat anti-rat Cy3	Jackson	112-166-072 (1:100)
goat anti-chicken AF647	Jackson	703-606-155 (1:100)
Oligonucleotides		
BDNF	Applied Biosystems	Mm04230607_s1
SOD2	Applied Biosystems	Mm01313000_m1; NM_013671.3;
S100a10	IDT	Mm.PT.58.6571055
CD38	Applied Biosystems	Mm01220906_m1
OLIG2	Applied Biosystems	NM_016967; Mm.PT.58.42319010
GFAP	IDT	NM_010277.2; GCAGATGAAGCCACCCTGG/ GAGGTCTGGCTTGGCCAC
IL1β	IDT	Mm.PT.51.17212823
IL6	Applied Biosystems	NM_031168.1;Mm00446190_m1
H2D1	IDT	Mm.PT.58.42136026.g
TNFα	Applied Biosystems	Mm00443258_m1
Rn18s	Applied	NR_003278.3; Mm03928990_g1

	Biosystems	
GAPDH	Applied biosystems	Mm99999915_g1
Chemicals		
78c CD38 inhibitor	Millipore	Cat#538763
Nicotinamide adenine dinucleotide	Sigma	Cat#10127965001
Nicotinamide 1,N6-ethenoadenine dinucleotide	Sigma	Cat#41628
flavin mononucleotide	Sigma	Cat#F8399
Alcohol dehydrogenase	Sigma	Cat#A3263
Diaphorase	Sigma	Cat#D5540
resazurin	Sigma	Cat#119303
palmitate	Sigma	Cat#P0500
DAPI	ThermoFisher	Cat#D1306; AB_2629482
lysolecithin	Sigma	Cat#L-4129
Methyl green	Sigma	67060
Commercial kits		
Bio-Plex Mouse Cytokine 23-plex Assay	Bio-Rad	Cat#m60009rdpd
Software		
GraphPad Prism 7	GraphPad	https://www.graphpad.com/scientificsoftware/prism
CellSens	Olympus	https://www.olympus-lifescience.com/en/software/cellsens/
Zen	Zeiss	https://www.zeiss.com/microscopy/us/products/microscope-software/zen.html
Fiji ImageJ	NIH	https://imagej.net/Fiji

1151

1152 **<u>Table 1.</u> Key Resources Table**