Intraspecific diversity loss in a predator species alters prey community structure and

ecosystem functions

Allan Raffard[1,2], Julien Cucherousset[3], José M. Montoya[1], Murielle Richard[1], Samson Acoca-

Pidolle[1], Camille Poésy[1], Alexandre Garreau[1], Frédéric Santoul[2] & Simon Blanchet[1].

[1] CNRS, Université Toulouse III Paul Sabatier, Station d'Écologie Théorique et Expérimentale du

CNRS à Moulis, UMR-5321, F-09200 Moulis, France.

[2] EcoLab, Université de Toulouse, CNRS, INPT, UPS, Toulouse, France

[3] CNRS, Université Toulouse III Paul Sabatier, UMR-5174 EDB (Laboratoire Evolution & Diversité

Biologique), 118 route de Narbonne, F-31062, Toulouse, France.

Corresponding author:

Allan Raffard

Station d'Ecologie Théorique et Experimentale du CNRS à Moulis.

UMR-5321(CNRS-UPS)

2 route du CNRS, F-09200 Moulis, France

Phone: (+33) 5.61.55.67.47

E-mail: allan.raffard@sete.cnrs.fr; allan.raffard@uclouvain.be

Authors' contributions: AR, JC and SB conceived the study. AR, MR, SA-P, CP and AG

carried out the experiment. AR performed the statistical analyses. AR, JC, JMM, FS and SB

interpreted and discussed the results. AR and SB wrote the article, and all authors made

corrections.

Keywords: ecosystem functioning, community, genotype, functional traits, intraspecific

diversity, trophic cascade, predator, food chain

Abstract

Loss in intraspecific diversity can alter ecosystem functions, but the underlying mechanisms

are still elusive, and intraspecific biodiversity-ecosystem function relationships (iBEF) have

been restrained to primary producers. Here, we manipulated genetic and functional richness of

a fish consumer (*Phoxinus phoxinus*), to test whether iBEF relationships exist in consumer

species, and whether they are more likely sustained by genetic or functional richness. We

found that both genotypic and functional richness affected ecosystem functioning, either

independently or in interaction. Loss in genotypic richness reduced benthic invertebrate

diversity consistently across functional richness treatments, whereas it reduced zooplankton

diversity only when functional richness was high. Finally, both losses in genotypic and

functional richness altered essential functions (e.g. decomposition) through trophic cascades.

We concluded that iBEF relationships lead to substantial top-down effects on entire food

chains. The loss of genotypic richness impacted ecological properties as much as the loss of

functional richness, probably because it sustains "cryptic" functional diversity.

Introduction

Human disturbances associated with global change are increasingly altering worldwide patterns of species diversity, as well the functions and services provided by ecosystems (Parmesan 2006; Naeem *et al.* 2009; Isbell *et al.* 2017). Nonetheless changes observed at the species and ecosystem levels are always preceded by changes in phenotypic and genotypic composition within plant and animal populations (Clements & Ozgul 2016; Clements *et al.* 2017; Baruah *et al.* 2019). Accordingly, extremely rapid changes in intraspecific diversity are currently occurring worldwide (Darimont *et al.* 2009; Mimura *et al.* 2016; Miraldo *et al.* 2016; Leigh *et al.* 2019). Changes in intraspecific diversity can affect species turnover and composition (Vellend 2005; Vellend & Geber 2005), as well as ecosystem functioning (Des Roches *et al.* 2018; Raffard *et al.* 2019c). For instance, the loss of genotypes within primary producers can reduce ecosystem process rates and species diversity (Crutsinger *et al.* 2006; Johnson *et al.* 2006; Fridley & Grime 2010), suggesting the existence of biodiversity-ecosystem functioning relationships at the intraspecific level (iBEF, Hughes *et al.* 2008; Koricheva & Hayes 2018; Raffard *et al.* 2019c), in addition to the more widely studied BEFs at the interspecific level.

The relationships between intraspecific diversity and ecosystem functioning are challenging to study and this is primarily because of the confusion between genetic and functional trait diversity. iBEF studies have initially manipulated genotypic richness within experimental populations (e.g. Crutsinger *et al.* 2006; Hughes & Stachowicz 2009; Abbott *et al.* 2017). Genetic variability is expected to encapsulate a large proportion of trait variability, and thus higher genotypic richness should maintain higher functional diversity (Hughes et al. 2008). Although this approach allows deciphering the general effects of intraspecific diversity on ecosystem functioning, it does not provide mechanisms since ecological interactions are supported by functional traits that are -partly- genetically encoded. As a consequence, parallel

studies have directly manipulated functional trait richness within populations, which enabled

to determine a direct mechanistic linkage between functional richness and community

structure and ecosystem functioning (Violle *et al.* 2007; Matthews *et al.* 2011). For instance,

individual body mass (and traits covarying with body mass) has strong ecological effects

because of the associated functional differences and resource use complementarity among

individuals (Woodward *et al.* 2005; Hildrew *et al.* 2007; Rudolf *et al.* 2014; Harmon *et al.*

2009; Rudolf & Rasmussen 2013a). Focusing on specific traits, such as body mass, might

conversely blur the ecological effects of "cryptic" trait variation (i.e. unmeasured functional

traits such as metabolic rate or feeding behaviour and activity, Brown *et al.* 2004; Wolf &

Weissing 2012) that is likely supported by genotypic richness. We therefore argue that

manipulating simultaneously genotypic richness and the diversity of key functional traits such

as body mass should allow assessing whether cryptic trait diversity is ecologically important,

and could provide a better mechanistic understanding of iBEF relationships.

The ecological effects of biodiversity changes can be particularly strong when the later

occur at high trophic levels such as secondary consumers or predators (Duffy 2002, 2003;

Griffin *et al.* 2013). Changes in the diversity and abundance of predatory species can trigger

important effects in functions supported by lower trophic levels, especially in regulating the

abundance of the lower trophic levels and indirectly ecosystem functioning, such as biomass

production (Shurin *et al.* 2002; Barnes *et al.* 2018). High predator species richness can

sometimes favour resource use complementarity and decrease prey abundance (Cardinale *et*

al. 2006; Griffin *et al.* 2013). In some cases, however, increasing predator richness increases

prey abundance through mechanisms such as predator interference (Sih *et al.* 1998; O'Connor

& Bruno 2009), and can modify multiple ecosystem functions along food webs (Antiqueira *et*

al. 2018; Barnes *et al.* 2018). These mechanisms have all been investigated at the interspecific

level, but they may also apply *within* a predatory species (intraspecific diversity), and we can

expect relationships –either positive or negative- between predator intraspecific diversity and

the structure of prey communities, that could subsequently cascade down on ecosystem

functions such as decomposition rate or primary production (Duffy, 2002; Bruno &

O'Connor, 2005). However, it is still difficult to forecast how loss in intraspecific diversity in

consumer species could affect ecosystem functioning, because iBEF studies have primarily

focused on primary producers (Koricheva & Hayes 2018; Raffard *et al.* 2019c). This is

despite the fact that human activities strongly affect predator and consumer populations, for

instance through harvest or fisheries activities, which may alter ecosystem functioning

through intraspecific changes in traits and genotypes (Hauser *et al.* 2002; Estes *et al.* 2011;

Palkovacs *et al.* 2012, 2018).

In this study, we investigated whether a loss in genotypic and functional diversity

within a consumer species at the top of a three-level trophic chain could mediate top-down

effects on key ecosystem functions. In a nine-months pond mesocosm experiment, we

simultaneously manipulated *genotypic richness* (number of genetic entities) and the *functional*

richness (variance in individuals body mass) of experimental populations of a freshwater fish,

the European minnow (*Phoxinus phoxinus*); a common and abundant species with important

ecosystems effects (Miró *et al.* 2018; Raffard *et al.* 2019b). We predicted that increased

functional richness should affect ecological functions; if functional richness captures the

entire functional differences among genotypes, then increasing genotypic richness should not

impact ecological functions further (Fig. 1a). Alternatively, if functional richness does not

capture all the functional differences among genotypes, then increasing genotypic richness

should increase functional diversity and affect ecological functions (Fig. 1b). The ecological

effects of genotypic and functional richness might display different shape following different

conceptual predictions to which we compared our experimental findings (Fig. 1): (i)

"additive" effects when loss in genotypic richness affects ecological functions regardless of

changes in functional richness, (ii) "enhancing" effects when loss in genotypic richness affect

ecological functions solely at higher levels of functional richness, or (iii) "compensatory"

effects when high genotypic richness compensates for the loss of functional richness,

maintaining higher ecological functions at lower levels of functional richness. Finally, we

investigated the mechanistic basis of the ecological effects of genotypic and functional

richness. In particular, we expected that genotypic and functional richness affect directly

community structure through trophic interactions, and indirectly ecosystem functions at lower

trophic levels through trophic cascades. Hence, we specifically tested whether the loss in

genotypic and functional richness directly affected population performance and community

structure, and indirectly ecosystem functioning mediated by changes in community structure

of benthic and pelagic food web.

Material and methods

Model species

The European minnow (*P. phoxinus*) is a common species occurring across Western Europe,

living in relatively cold water (summer water temperature generally lower than 24°C)

including mountains lakes, small rivers at intermediate altitude and mountain streams (Keith

et al. 2011). It is a small-bodied cyprinid fish species (<12 cm long, 4-8 cm long as an adult

in general) playing the role of secondary consumer in most ecosystems (Raffard *et al.* 2020),

with a generalist diet composed of small invertebrates, algae or zooplankton (Frost 1943;

Collin & Fumagalli 2011). Populations of European minnows differed in their genetic and

phenotypic richness (Fourtune *et al.* 2018), and previous works revealed that genetically and

phenotypically unique populations differently affect prey community abundance and

ecosystem functions (Raffard *et al.* 2019b).

We selected ten populations from a large river basin in southern France (the Garonne

catchment) based on *a priori* knowledge to maximise both genetic and functional

differentiations among populations (Fourtune *et al.* 2018; Raffard *et al.* 2019a) (Fig. S1).

Specifically, the ten selected populations displayed a high level of genetic differentiation

(mean pairwise Fst = 0.133, range = 0.029-0.320) and greatly varied in the number of alleles

they harbour (from 5.470 alleles in average for the less diverse population to 10.176 alleles

for the most diverse population) according to data measured using seventeen microsatellite

markers (from Fourtune *et al.* 2018; Prunier *et al.* 2019; Raffard *et al.* 2019a). Moreover, we

selected five populations mainly composed of small-bodied adults (mean body mass ±

standard error (SE) = 1.03 g ± 0.02) and five populations mainly composed of large-bodied

adults (mean body mass ± SE = 3.06 g ± 0.07) (Fig. S2a). These differences in adult body

mass are due to selective pressures from the local environment such as mean water

temperature; the higher the mean water temperature, the smallest the adult body mass due to

increased metabolic rate and an accelerated pace-of-life (Raffard *et al.* 2019a). Finally,

populations are geographically distant one from each other, and they will hereafter be each

considered as functionally and genetically unique entity, and intraspecific richness will be

manipulated by varying the number of populations in independent mesocosms (see details

below).

In September 2017, individuals were collected in each river by electrofishing a 200-m

section and visually selecting 30-50 individuals reflecting the size range of adults. Fish were

maintained at similar density in outdoor tanks for 3 weeks before the start of the experiment.

Fish were fed ad libitum with frozen Chironomidae during this period.

Mesocosm experiment

In October 2016, 24 outdoor mesocosms were filled with 900 L of water, and 3 cm of gravel.

Nutrients were added to the mesocosms using 10 mL of solution containing nitrogen and

phosphorus (ratio N:P:K = 3.3:1.1:5.8). Each mesocosm was then inoculated twice (October

2016 and May 2017) with 200 mL of a concentrated solution of phytoplankton and 200 mL of

concentrated solution of zooplankton collected from a unique lake nearby (Lamartine Lake,

France 43°30'21.5"N, 1°20'32.7"E). In May 2017, we introduced three adult pond-snails

(Lymnaeidae) and ten adult isopods (Asellidae) in each mesocosms. They were let uncovered

to allow natural colonization by other invertebrates and community assemblage until the start

of the experiment that occurred approximately 6 months later.

In October 2017, eight fish were weighted and introduced in each mesocosm, which

were assigned to one of four treatments according to a full-factorial design with *genotypic*

richness (two levels, *high* and *low* genotypic richness) and *functional* richness (two levels,

high and *low* functional richness) as the main factors. Genotypic richness was manipulated by

introducing individuals sourced from either two (*low genotypic richness*, four individuals

from each population) or four (*high genotypic richness*, two individuals from each

population) out of the ten populations displaying significant genetic differentiation (Fourtune

et al. 2018; Raffard *et al.* 2019a). These two levels of genotypic richness were selected as it

has previously been shown in a meta-analysis that the effect of intraspecific richness on

ecosystem functioning increases linearly until approximately four genotypes and then reaches

a plateau beyond that (Raffard *et al.* 2019c). Since we aimed at testing the effect of richness

while minimizing the effects of population identity, each replicate of each treatment of

richness contained a different (unique) assemblage of populations. Functional richness

consisted in manipulating the body mass of the individuals present in the mesocosms; hence,

experimental populations contained individuals sourced either from large or small populations

(i.e., one functional entity, *low functional richness*)*, or from both small and large populations

(i.e., two functional entities, *high functional richness*, see Table S1 and Fig. S2 for details on

the different experimental populations). It is noteworthy that populations containing only

either small or large individuals were actually more functionally redundant than populations

containing large and small individuals. Experimental populations hence varied simultaneously

according to their genotypic and functional richness, leading to four treatments.

The experiment lasted 30 weeks (210 days) after fish were introduced. Mesocosms

were checked daily for mortality, which was rarely observed over the course of the

experiment. At the end of the experiment, we measured several ecological parameters to

assess effects of genotypic and functional richness on population performance, community

structure and ecosystem functioning:

- *Population performance*: All tanks were emptied and we recaptured all remaining

fish to estimate mortality rate ($1 - \frac{remaining\ fish}{introduced\ fish}$; mean per tank $\pm$ standard error (SE) =

0.22 $\pm$ 0.01). Fish were weighted to assess fish biomass production (biomass production =

averaged final weight - averaged initial weight) of each experimental population during the

experiment. Since individuals with small initial body mass intrinsically displayed a higher

growth rate than large individuals, a standardised biomass production index was calculated as

the residuals of the relationship between biomass production and initial fish size.

- *Community structure*: Zooplankton community structure was quantified by filtering

5 L of water through a 200 μm sieve. Samples were conserved in a 70% ethanol solution and

subsequently identified to the order or family levels, including Cyclopoida, Calanoida,

Daphniidae, Chydoridae and Bosminidae. The diversity of zooplankton was calculated as the

Simpson's diversity (*D-zoo*) representing the probability that two randomly chosen

individuals belong to different clades. *D-zoo* was calculated as $1 - \frac{\sum N_i * (N_i - 1)}{N_{tot} * (N_{tot} - 1)}$, where N_{tot}

was the total number of sampled individuals, and N_i the number of sampled individuals for

each group (Simpson 1949; Lande 1996). Zooplankton abundance was quantified as the total

number of individuals for all taxa pooled at the tank level.

Benthic invertebrates were collected from the mesh bags used to measure

decomposition rates (*see below*), conserved in a 70% ethanol solution, and identified as

Isopoda, Diptera, Gastropoda, Ephemeroptera, Plecoptera, Odonata, Copepoda, Cladocera,

and Ostracoda. The diversity of benthic invertebrates was calculated as the Simpson's

diversity (*D-inv*). The abundance of benthic invertebrates was quantified as the total number

of individuals for all taxa pooled at the tank level.

- *Ecosystem functioning*: Decomposition rate was measured by quantifying the mass

loss of black poplar (*Populus nigra*, a dominant riparian tree in southern France) abscised

leaves (Alp *et al.* 2016). On the 7[th] of March 2018 (19 weeks after the introduction of fish), 4

g of air-dried leaves were put in each mesocosm within a coarse mesh (1 x 1 cm) bag. At the

end of the experiment, the remaining leaf material was removed from the mesocosms, rinsed

with tap water, oven dried at 60°C for three days and weighed to the nearest 0.001 g to assess

the loss of biomass. The decomposition rate was calculated as $k = -\frac{ln\,(x)}{t}$ (Alp *et al.* 2016),

where *X* is the proportion of litter remaining and *t* is the elapsed time in days.

Pelagic algae stock was measured as the chlorophyll-a concentration ($\mu g.L^{-1}$) in the

water column using a multiparametric probe (OTT, Hydrolab DS5®). Five measurements

were taken in each mesocosm and averaged for subsequent analyses. Since phytoplankton are

the main primary producers in these mesocosms (Downing & Leibold 2002), pelagic algae

stock can be considered as a proxy for the biomass of primary producers.

Statistical analyses

Prior to analysis, the pelagic algae stock (i.e. the chlorophyll-a concentration), zooplankton

and benthic invertebrate abundances were log-transformed to reach normality of models'

residuals. After analyses of outliers using Cook's distance, we removed one mesocosm from

the final analyses that displayed outliers for several of the measured variables (Fig. S3).

Analyses were hence run on 23 replicates.

We first assessed the ecological effects of genotypic and functional richness using

mixed effects linear models (LMMs). We ran one model for each ecological parameter as a

dependent variable, and the genotypic richness, functional richness, and the resulting two-

term interaction (that was removed when non-significant) as explanatory variables. Fish

mortality rate during the experiment was included as a fixed effect to control for a potential

effect of final density on ecological processes. To control for the disposition of the tanks

during the experiment, the position of tanks was added as a random term. To compare the

relative strength of a loss in genotypic and functional richness on ecological parameters we

calculated effect sizes. Hedges' g were computed as follow: $g = \frac{X_2 - X_1}{s_{pooled}}$, where X_1 and X_2 are

means of treatments (for each genotypic or functional richness treatment separately) measured

for each ecological parameters respectively, and s_{pooled} is the pooled standard deviation

computed as $s_{pooled} = \sqrt{\frac{(n_2-1)s_2^2+(n_1-1)s_1^2}{n_1+n_2-2}}$, where n is sample size, and s^2 is variance

(Nakagawa & Cuthill 2007). An effect size was hence measured for each variable response (n

= 7, and each treatment) independently. These individual effect sizes were then averaged for

genotypic (g_G) and functional richness (g_S) respectively so as to get an absolute mean effect

sizes that were compared visually based on 95% confidence intervals (CI).

Secondly, we ran confirmatory path-analysis (Shipley 2009) to assess the direct and

indirect links among intraspecific diversity (genotypic and functional richness), prey

(invertebrates and zooplankton) community structure and ecosystem functioning. We

expected that intraspecific diversity in European minnows affects directly invertebrate

community (both diversity and abundance) through predation, and indirectly ecosystem

functions (decomposition rate, algae stock) through trophic cascades. Specifically, benthic

invertebrates consume leaf litter, affecting decomposition rate (Gessner *et al.* 1999, 2010),

while zooplankton forages on phytoplankton, affecting pelagic algae stock (brown and green

trophic food chain, respectively). Because genotypic and functional richness likely interact to

shape processes (see Fig. 1), we initially included in all models the interaction term between

genotypic and functional richness on community structure. As above, interaction terms were

removed when not significant. Specifically, we computed two path analysis: one for the

benthic food chain (fish-invertebrates-decomposition), and one for the pelagic food chain

(fish-zooplankton-algae stock) (Fig. S4). The fits of the structural models were assessed using

the C statistic (see independent claims in Table S2), that follows a χ^2 distribution and whose

associated *p*-value indicates whether the models adequately fit the data or not. Alternative

models including direct effects of fish intraspecific diversity on community components and

ecosystem functions were performed (Fig. S4) to test whether genotypic and functional

richness affected ecosystem functions through unmeasured parameters (e.g., other community

parameters, nutrient recycling). This approach allows testing whether the predicted indirect

effects are the most likely effects that explain the data. Finally, the structural models were

further compared to simplified models that did not include the effects of genotypic and

functional richness on community parameters (see Fig. S4) using Akaike Information Criteria

(AIC, Cardon *et al.* 2011; Shipley 2013). This later test allows assessing whether the indirect

effects of genotypic and functional richness lead to a better explanation of the data, and

whether trophic cascades were important or not.

All statistical analysis were run using R software (R Core Team 2013), LMM were

run using the R-package lme4 (Bates *et al.* 2014), and confirmatory path-analyses were run

using the R-package ggm (Marchetti *et al.* 2020).

Results

Both genotypic and functional richness of experimental populations significantly

affected several ecological processes (Table 1). At the population level, we found that the

interaction between genotypic and functional richness tended (p = 0.057) to alter fish biomass

production (i.e., population performance) of experimental populations (Table 1, Fig. 2a).

Specifically, the less diversified experimental populations (low genotypic richness and low

functional richness) displayed a lower biomass production than all other treatments (Fig. 2a).

At the community level, the diversity of benthic invertebrates was significantly higher in the

high genotypic richness treatment (mean *D-inv* ± SE = 0.64 ± 0.04) than in the low genotypic

richness treatment (*D-inv* ± SE = 0.53 ± 0.03), and this pattern was repeatable across

functional richness treatments (Fig. 2d, Table 1). Irrespective of the genotypic richness

treatment, mesocosms containing fish populations with low functional richness had higher

diversity of benthic invertebrates than mesocosms containing fish populations with high

functional diversity (Fig. 2d, Table 1). In contrast, the diversity of zooplankton (*D-zoo*) was

affected by the interaction between genotypic and functional richness (Fig. 2e, Table 1).

Zooplankton diversity was strikingly higher in populations with both high genotypic richness

and high functional richness (Fig. 2e, Table 1), suggesting that, in this case, the effect of

genotypic richness depended on the functional richness of experimental fish populations.

Benthic invertebrates and zooplankton abundances were consistently (i.e. across genotypic

richness treatment) enhanced when increasing functional richness (Fig. 2f, g, and Table 1).

Genotypic richness of populations did not affect benthic invertebrates and zooplankton

abundances (Fig. 2f, g, and Table 1). A weak but significant negative effect of functional

richness on the pelagic algae stock was detected (Table 1, Fig. 2c) and there was no

significant effects genotypic and functional richness on decomposition rate (Table 2). Overall,

genotypic and functional richness induced ecological effects of similar intensity (mean $|g_G|$ =

0.356, CIs = 0.043-0.669, and mean $|g_S|$ = 0.564, CIs = 0.247-0.882, see Fig. S5 for details).

Confirmatory path analyses revealed that effects of intraspecific diversity on

ecosystem functioning were primarily mediated by trophic cascades, through changes in

community structure (Fig. 3). First, genotypic and functional richness affected the diversity

and abundance of benthic invertebrates (Fig. 3), which subsequently and positively affected

the decomposition rate ($p < 0.001$, Fig. 4a, b and Table S3). Second, functional richness

positively affected abundance of zooplankton, leading to a decrease in pelagic algae stock (p

< 0.001, Fig. 4c, d, Table S3). Overall, the confirmatory path-analysis confirmed the key role

of intraspecific diversity for both food chains, and its indirect effects for ecosystem functions

(Fig. 3 and Table 2). Indeed, models including genotypic and functional richness reproduced

adequately the causal pathways (Table 2), and their AICs were better than that of both

alternative and simplified models for each ecosystem function (Table 2).

Discussion

The present study demonstrates that loosing intraspecific diversity in a secondary consumer

species has substantial top-down consequences for community structure and ecosystem

functioning in food webs. We showed that both losses in functional (mass) and genotypic

richness sustained these iBEF relationships, suggesting that differences in mass among

individuals (an important functional trait for food web dynamics, Woodward *et al.* 2005) did

not capture the entire functional space, and thus that genotypic richness encapsulates

important and "cryptic" (unmeasured) functional diversity. The loss of genotypes within

consumer populations can affect both the community structure and the abundance of lower

trophic levels, as well as ecosystem functions with a similar strength than the loss of

functional entities (i.e. body mass). Specifically, we found that diversity loss (genotypic and

functional) within populations indirectly affects primary producer biomass and organic matter

recycling, two ecosystem functions forming the bases of food chains. This suggests that

intraspecific diversity is a key but understudied facet of biodiversity as it indirectly sustains

BEF relationships, even when changes in intraspecific diversity occur at the top of the food

chains.

Our study suggests that genotypic richness can support non-negligible cryptic

functional diversity. Cryptic functional traits, such as physiological and behavioural traits, can

affect interactions in food webs and ecological functions. As such, behavioural variation

within populations is widespread in nature, and can be genetically encoded (Sih *et al.* 2004;

Anreiter & Sokolowski 2019). For instance, personality traits (e.g., activity or boldness) can

determine foraging activity and diet, ultimately affecting trophic interactions and ecosystem

functioning in food webs (Wolf & Weissing 2012; Toscano *et al.* 2016). Additionally,

metabolic and stoichiometric traits, can also strongly differ within species, and can be

functionally important in shaping energetic needs both quantitatively and qualitatively (i.e.,

specific elemental needs) (Brown *et al.* 2004; Leal, *et al.* 2016; Rosenblatt & Schmitz 2016).

Hence, fish with different genotypes may share obvious functional traits (such as body mass)

but may subtly differ in other cryptic functional traits, making them unique ecologically

(Loreau 2001). Although such "ecological dissimilarity-despite-morphological similarity" has

rarely been demonstrated within species, there is now ample evidence that cryptic species

(species being morphologically and phylogenetically similar) can actually be ecologically

unique regarding their influence on ecosystems (e.g. Fišer *et al.* 2015).

Interestingly, this cryptic diversity can interact with functional richness in various

ways (i.e., additive, enhancing, or compensatory) depending on the ecological parameter

considered (i.e., benthic invertebrate diversity, zooplankton diversity, and fish biomass

production, respectively). For instance, we found that genotypic richness can compensate for

the loss of functional diversity for population performance (i.e., increase fish biomass

production), with high genotypic richness maintaining high biomass production even when

the distribution of body mass in the population is limited. We speculate that genotypic

richness can increase niche variation among individuals, thereby limiting the loss of biomass

production when decreasing functional richness (Duffy *et al.* 2001). These findings echo and

expand on studies at the interspecific level demonstrating that phylogenetic diversity among

species explains variance in ecosystem functions that is not explained either by species or

functional diversity, suggesting that phylogenetic diversity sustains unmeasured functional

differences among species (Cadotte *et al.* 2012; Mouquet *et al.* 2012; Le Bagousse-Pinguet *et*

al. 2019). We argue that iBEF relationships are sustained by differences in functional traits

among individuals that can be directly measured (e.g. mass) and/or indirectly captured

through quantification of the genetic pool of individuals composing populations.

Accordingly, we showed that genotypic and functional richness can affect

independently (i.e., additively) and consistently community structure, demonstrating that

multiple facets of intraspecific diversity can regulate lower trophic levels. First, genotypic

richness increased benthic invertebrate diversity. This effect was very robust, as it was

repeatable and consistent across functional richness treatments. Genotypic richness probably

enhanced resource partitioning, allowing individuals to forage on a more diverse array of

resources, regulating the abundance of each taxonomic group and leading to a higher diversity

(Duffy, 2002; Johnson *et al.*, 2006; Hughes et al. 2008). Second, functional richness led to

increased prey abundance that was repeatable across both communities of benthic

invertebrates and zooplankton. Previous studies at the inter- and intraspecific levels provide

non-consistent predictions, since an increase in consumer diversity can affect either positively

or negatively resource abundance (Sih *et al.* 1998; Griffin *et al.* 2013; Rudolf & Rasmussen

2013; Antiqueira *et al.* 2018; Barnes *et al.* 2018). Our results indicate that functionally rich

populations did not consume fewer resources than functionally poor populations since they

had higher biomass production. This suggests that the increase in prey abundance was not due

to an increase in inter-individual competition in functionally rich populations. At the opposite,

flexible exploitation of resources might occur in functional rich populations because

European minnows are omnivorous (Frost 1943; Collin & Fumagalli 2011), their diet

probably included some periphyton, decreasing the predation risk on animal resources, and

hence increasing invertebrate abundance. Because the top-down control of intraspecific

diversity on community structure is likely driven by trophic mechanisms, quantifying

individual diet (e.g. using gut content, eDNA or stable isotope analysis) and its temporal

dynamic in such experiments would allow making more accurate predictions regarding

trophic niche partitioning.

Our results further revealed how changes in top consumers' genotypic and functional

richness percolate through the food web and alter ecosystem functions at the base of the food

chains. Such trophic cascade induced by intraspecific diversity could be driven by a

"classical" (i.e. interspecific) BEF relationship between benthic invertebrate diversity and

decomposition rate (Hooper *et al.* 2005; Gessner *et al.* 2010). Specifically, fish genotypic

richness increased benthic invertebrate diversity that accelerated litter decomposition rate.

The higher decomposition rate of organic matter is likely produced by higher consumption

efficiency through trophic complementarity among clades of invertebrates in diverse

community (Gessner *et al.* 2010). Invertebrate community with a high diversity probably

harboured a high functional diversity (Cadotte *et al.* 2011), and focusing on the functional

type of invertebrates might allow a more precise understanding of this link. These results echo

those reported at the community level and those manipulating richness within primary

producer species (Downing & Leibold 2002; Crutsinger *et al.* 2006), while implying a

modification of top-down effects by intraspecific diversity in consumers. Biodiversity within

consumer species is largely impacted by anthropogenic activities (Miraldo *et al.* 2016),

involving the loss of non-measured functional diversity, which might further lead to

underestimated cascading impacts on ecosystem functioning.

In conclusion, we demonstrated that both genetic and functional richness within

consumer populations are important facets of biological diversity, inducing effects on prey

community structure and trophic cascades mediating ecosystem functions. These results are

consistent with previous synthetic works (Koricheva & Hayes 2018; Raffard *et al.* 2019c),

reinforcing the call for considering changes of intraspecific diversity in consumer species as

an important ecological predictor. Importantly, genotypic richness can sustain important

cryptic functional diversity, and future investigations should aim at developing a general

framework from genes to ecosystems to better understand the global links existing among the

multiple facets of biodiversity and ecosystem functioning, and ultimately ecosystem services.

Acknowledgements

We thank Kéoni Saint-Pe and Jérôme G. Prunier for their help for fish sampling. AR is

financially supported by a Doctoral scholarship from the Université Fédérale de Toulouse.

Electrofishing was performed under the authorization of local authorities (Arrêtés

Préfectoraux from the Direction Départementale des Territoires of French departments

Ariège, Haute-Garonne, Tarn, Aveyron, Tarn-et-Garonne, Lot). This work was undertaken at

SETE, which is part of the "Laboratoire d'Excellence" (LABEX) entitled TULIP (ANR-10-

LABX-41).

Conflict of interest disclosure

The authors of this article declare that they have no financial conflict of interest with the

content of this article. Simon Blanchet and José M. Montoya are one of the PCI Ecology

recommenders.

Data availability statement: Data are available on Figshare 10.6084/m9.figshare.12459065
440

441 References

Abbott, J.M., Grosberg, R.K., Williams, S.L. & Stachowicz, J.J. (2017). Multiple dimensions
of intraspecific diversity affect biomass of eelgrass and its associated community.
Ecology, 98, 3152–3164.
Alp, M., Cucherousset, J., Buoro, M. & Lecerf, A. (2016). Phenological response of a key
ecosystem function to biological invasion. *Ecol. Lett.*, 19, 519–527.
Anreiter, I. & Sokolowski, M.B. (2019). The *foraging* gene and its behavioral effects:
pleiotropy and plasticity. *Annu. Rev. Genet.*, 53, 373–392.
Antiqueira, P.A.P., Petchey, O.L. & Romero, G.Q. (2018). Warming and top predator loss
drive ecosystem multifunctionality. *Ecol. Lett.*, 21, 72–82.
Barnes, A.D., Jochum, M., Lefcheck, J.S., Eisenhauer, N., Scherber, C., O'Connor, M.I., *et
al.* (2018). Energy flux: the link between multitrophic biodiversity and ecosystem
functioning. *Trends Ecol. Evol.*, 33, 186–197.
Baruah, G., Clements, C.F., Guillaume, F. & Ozgul, A. (2019). When do shifts in trait
dynamics precede population declines? *Am. Nat.*, 193, 633–644.
Bates, D., Maechler, M, Bolker, B & Walker, S. (2014). lme4: Linear mixed-effects models
using Eigen and S4. R package version 1.1-7.
Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). Toward a
metabolic theory of ecology. *Ecology*, 85, 1771–1789.
Bruno, J.F. & O'Connor, M.I. (2005). Cascading effects of predator diversity and omnivory in
a marine food web. *Ecol. Lett.*, 8, 1048–1056.
Cadotte, M.W., Carscadden, K. & Mirotchnick, N. (2011). Beyond species: functional
diversity and the maintenance of ecological processes and services. *J. Appl. Ecol.*, 48,
1079–1087.
Cadotte, M.W., Dinnage, R. & Tilman, D. (2012). Phylogenetic diversity promotes ecosystem
stability. *Ecology*, 93, S223–S233.
Cardinale, B.J., Srivastava, D.S., Emmett Duffy, J., Wright, J.P., Downing, A.L., Sankaran,
M., *et al.* (2006). Effects of biodiversity on the functioning of trophic groups and
ecosystems. *Nature*, 443, 989–992.
Cardon, M., Loot, G., Grenouillet, G. & Blanchet, S. (2011). Host characteristics and
environmental factors differentially drive the burden and pathogenicity of an
ectoparasite: a multilevel causal analysis: Causalities in host-parasite interactions. *J.
Anim. Ecol.*, 80, 657–667.
Clements, C.F., Blanchard, J.L., Nash, K.L., Hindell, M.A. & Ozgul, A. (2017). Body size
shifts and early warning signals precede the historic collapse of whale stocks. *Nat.
Ecol. Evol.*, 1, 0188.

Clements, C.F. & Ozgul, A. (2016). Including trait-based early warning signals helps predict
population collapse. *Nat. Commun.*, 7, 10984.
Collin, H. & Fumagalli, L. (2011). Evidence for morphological and adaptive genetic
divergence between lake and stream habitats in European minnows (*Phoxinus*
phoxinus , Cyprinidae). *Mol. Ecol.*, 20, 4490–4502.
Crutsinger, G.M., Collins, M.D., Fordyce, J.A., Gompert, Z., Nice, C.C. & Sanders, N.J.
(2006). Plant genotypic diversity predicts community structure and governs an
ecosystem process. *Science*, 313, 966–968.
Darimont, C.T., Carlson, S.M., Kinnison, M.T., Paquet, P.C., Reimchen, T.E. & Wilmers,
C.C. (2009). Human predators outpace other agents of trait change in the wild. *Proc.*
Natl. Acad. Sci., 106, 952–954.
Des Roches, S., Post, D.M., Turley, N.E., Bailey, J.K., Hendry, A.P., Kinnison, M.T., *et al.*
(2018). The ecological importance of intraspecific variation. *Nat. Ecol. Evol.*, 2, 57–
64.
Downing, A.L. & Leibold, M.A. (2002). Ecosystem consequences of species richness and
composition in pond food web. *Nature*, 416, 837–-841.
Duffy, J.E. (2002). Biodiversity and ecosystem function: the consumer connection. *Oikos*, 99,
201–219.
Duffy, J.E. (2003). Biodiversity loss, trophic skew and ecosystem functioning. *Ecol. Lett.*, 6,
680–687.
Duffy, J.E., Macdonald, K.S., Rhode, J.M. & Parker, J.D. (2001). Grazer diversity, functional
redundancy, and productivity in seagrass beds: an experimental test. *Ecology*, 82,
2417– 2434.
Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., *et al.* (2011).
Trophic Downgrading of Planet Earth. *Science*, 333, 301–306.
Fišer, Ž., Altermatt, F., Zakšek, V., Knapič, T. & Fišer, C. (2015). Morphologically Cryptic
Amphipod Species Are "Ecological Clones" at Regional but Not at Local Scale: A
Case Study of Four Niphargus Species. *PLOS ONE*, 10, e0134384.
Fourtune, L., Prunier, J.G., Mathieu-Begne, E., Canto, N., Veyssiere, C., Loot, G., *et al.*
(2018). Intraspecific genetic and phenotypic diversity: parallel processes and
correlated patterns?
Fridley, J.D. & Grime, J.P. (2010). Community and ecosystem effects of intraspecific genetic
diversity in grassland microcosms of varying species diversity. *Ecology*, 91, 2272–
2283.
Frost, W.E. (1943). The natural history of the minnow, *Phoxinus phoxinus. J. Anim. Ecol.*, 12,
139.
Gessner, M.O., Chauvet, E. & Dobson, M. (1999). A perspective on leaf litter breakdown in
streams. *Oikos*, 85, 377–384.
Gessner, M.O., Swan, C.M., Dang, C.K., McKie, B.G., Bardgett, R.D., Wall, D.H., *et al.*
(2010). Diversity meets decomposition. *Trends Ecol. Evol.*, 25, 372–380.
Griffin, J.N., Byrnes, J.E.K. & Cardinale, B.J. (2013). Effects of predator richness on prey
suppression: a meta-analysis. *Ecology*, 94, 2180–2187.
Hauser, L., Adcock, G.J., Smith, P.J., Bernal Ramirez, J.H. & Carvalho, G.R. (2002). Loss of
microsatellite diversity and low effective population size in an overexploited
population of New Zealand snapper (Pagrus auratus). *Proc. Natl. Acad. Sci.*, 99,
11742–11747.
Hildrew, A.G., Raffaelli, D. & Edmonds-Browns, R. (2007). *Body size: the structure and*
function of aquatic ecosystems. Cambridge University Press, Cambridge, UK.

Hooper, D.U., Chapin, F.S., Ewel, J.J., Hector, A., Inchausti, P., Lavorel, S., *et al.* (2005).
Effects of biodiversity on ecosystem functioning: a consensus of current knowledge.
Ecol. Monogr., 75, 3–35.
Hughes, A.R., Inouye, B.D., Johnson, M.T.J., Underwood, N. & Vellend, M. (2008).
Ecological consequences of genetic diversity. *Ecol. Lett.*, 11, 609–623.
Hughes, A.R. & Stachowicz, J.J. (2009). Ecological impacts of genotypic diversity in the
clonal seagrass Zostera marina. *Ecology*, 90, 1412–1419.
Isbell, F., Gonzalez, A., Loreau, M., Cowles, J., Díaz, S., Hector, A., *et al.* (2017). Linking
the influence and dependence of people on biodiversity across scales. *Nature*, 546,
65–72.
Johnson, M.T.J., Lajeunesse, M.J. & Agrawal, A.A. (2006). Additive and interactive effects
of plant genotypic diversity on arthropod communities and plant fitness. *Ecol. Lett.*, 9,
24–34.
Keith, P., Persat, H., Feunteun, E. & Allardi, J. (2011). *Les poissons d'eau douce de France.*
Muséum National d'Histoire Naturelle and Publications Biotope. Paris and Mèze.
Koricheva, J. & Hayes, D. (2018). The relative importance of plant intraspecific diversity in
structuring arthropod communities: A meta-analysis. *Funct. Ecol.*, 32, 1704–1717.
Lande, R. (1996). Statistics and partitioning of species diversity, and similarity among
multiple communities. *Oikos*, 76, 5.
Le Bagousse-Pinguet, Y., Soliveres, S., Gross, N., Torices, R., Berdugo, M. & Maestre, F.T.
(2019). Phylogenetic, functional, and taxonomic richness have both positive and
negative effects on ecosystem multifunctionality. *Proc. Natl. Acad. Sci.*, 116, 8419–
8424.
Leal, M.C., Seehausen, O. & Matthews, B. (2016). The ecology and evolution of
stoichiometric phenotypes. *Trends Ecol. Evol.*
Leigh, D.M., Hendry, A.P., Vázquez-Domínguez, E. & Friesen, V.L. (2019). Estimated six
per cent loss of genetic variation in wild populations since the industrial revolution.
Evol. Appl., eva.12810.
Loreau, M. (2001). Biodiversity and Ecosystem Functioning: Current Knowledge and Future
Challenges. *Science*, 294, 804–808.
Marchetti, G.M., Drton, M. & Sadeghi, K. (2020). ggm: functions for graphical Markov
models. Version 2.5.
Matthews, B., Narwani, A., Hausch, S., Nonaka, E., Peter, H., Yamamichi, M., *et al.* (2011).
Toward an integration of evolutionary biology and ecosystem science. *Ecol. Lett.*, 14,
690–701.
Mimura, M., Yahara, T., Faith, D.P., Vázquez-Domínguez, E., Colautti, R.I., Araki, H., *et al.*
(2016). Understanding and monitoring the consequences of human impacts on
intraspecific variation. *Evol. Appl.*, 10, 121–139.
Miraldo, A., Li, S., Borregaard, M.K., Florez-Rodriguez, A., Gopalakrishnan, S., Rizvanovic,
M., *et al.* (2016). An Anthropocene map of genetic diversity. *Science*, 353, 1532–
1535.
Miró, A., Sabás, I. & Ventura, M. (2018). Large negative effect of non-native trout and
minnows on Pyrenean lake amphibians. *Biol. Conserv.*, 218, 144–153.
Mouquet, N., Devictor, V., Meynard, C.N., Munoz, F., Bersier, L.-F., Chave, J., *et al.* (2012).
Ecophylogenetics: advances and perspectives. *Biol. Rev.*, 87, 769–785.
Naeem, S., Bunker, D.E., Hector, A., Loreau, M. & Perrings, C. (2009). *Biodiversity,*
Ecosystem Functioning, and Human Wellbeing. Oxford University Press.
Nakagawa, S. & Cuthill, I.C. (2007). Effect size, confidence interval and statistical
significance: a practical guide for biologists. *Biol. Rev.*, 82, 591–605.

O'Connor, M.I. & Bruno, J.F. (2009). Predator richness has no effect in a diverse marine food
web. *J. Anim. Ecol.*, 78, 732–740.
Palkovacs, E.P., Kinnison, M.T., Correa, C., Dalton, C.M. & Hendry, A.P. (2012). Fates
beyond traits: ecological consequences of human-induced trait change. *Evol. Appl.*, 5,
183–191.
Palkovacs, E.P., Moritsch, M.M., Contolini, G.M. & Pelletier, F. (2018). Ecology of harvest-
driven trait changes and implications for ecosystem management. *Front. Ecol.*
Environ., 16, 20–28.
Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annu.*
Rev. Ecol. Evol. Syst., 37, 637–669.
Prunier, J.G., Chevalier, M., Raffard, A., Loot, G., Poulet, N. & Blanchet, S. (2019).
Contemporary loss of genetic diversity in wild fish populations reduces biomass
stability over time. *bioRxiv*.
R Core Team. (2013). R: a language and environment for statistical computing.
Raffard, A., Cucherousset, J., Prunier, J.G., Loot, G., Santoul, F. & Blanchet, S. (2019a).
Variability of functional traits and their syndromes in a freshwater fish species (
Phoxinus phoxinus): The role of adaptive and nonadaptive processes. *Ecol. Evol.*, 9,
2833–2846.
Raffard, A., Cucherousset, J., Santoul, F., Di Gesu, L. & Blanchet, S. (2019b). Intraspecific
variation alters ecosystem and next-generation performance as much as temperature.
bioRxiv.
Raffard, A., Santoul, F., Blanchet, S. & Cucherousset, J. (2020). Linking intraspecific
variability in trophic and functional niches along an environmental gradient. *Freshw.*
Biol., fwb.13508.
Raffard, A., Santoul, F., Cucherousset, J. & Blanchet, S. (2019c). The community and
ecosystem consequences of intraspecific diversity: a meta-analysis. *Biol. Rev.*, 94,
648–661.
Rosenblatt, A.E. & Schmitz, O.J. (2016). Climate change, nutrition, and bottom-up and top-
down food web processes. *Trends Ecol. Evol.*, 31, 965–975.
Rudolf, V.H.W. & Rasmussen, N.L. (2013). Population structure determines functional
differences among species and ecosystem processes. *Nat. Commun.*, 4.
Rudolf, V.H.W., Rasmussen, N.L., Dibble, C.J. & Van Allen, B.G. (2014). Resolving the
roles of body size and species identity in driving functional diversity. *Proc. R. Soc. B*
Biol. Sci., 281, 20133203.
Shipley, B. (2009). Confirmatory path analysis in a generalized multilevel context. *Ecology*,
90, 363–368.
Shipley, B. (2013). The AIC model selection method applied to path analytic models
compared using ad-separation test. *Ecology*, 94, 560–564.
Shurin, J.B., Borer, E.T., Seabloom, E.W., Anderson, K., Blanchette, C.A., Broitman, B., *et*
al. (2002). A cross-ecosystem comparison of the strength of trophic cascades. *Ecol.*
Lett., 5, 785–791.
Sih, A., Bell, A. & Johnson, J.C. (2004). Behavioral syndromes: an ecological and
evolutionary overview. *Trends Ecol. Evol.*, 19, 372–378.
Sih, A., Englund, G. & Wooster, D. (1998). Emergent impacts of multiple predators on prey.
Trends Ecol. Evol., 13, 350–355.
Simpson, E.H. (1949). Measurement of diversity. *Nature*, 163, 688.
Toscano, B.J., Gownaris, N.J., Heerhartz, S.M. & Monaco, C.J. (2016). Personality, foraging
behavior and specialization: integrating behavioral and food web ecology at the
individual level. *Oecologia*, 182, 55–69.

Vellend, M. (2005). Species diversity and genetic diversity: parallel processes and correlated
patterns. *Am. Nat.*, 166, 199–215.
Vellend, M. & Geber, M.A. (2005). Connections between species diversity and genetic
diversity. *Ecol. Lett.*, 8, 767–781.
Violle, C., Navas, M.-L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., *et al.* (2007). Let
the concept of trait be functional! *Oikos*, 116, 882–892.
Wolf, M. & Weissing, F.J. (2012). Animal personalities: consequences for ecology and
evolution. *Trends Ecol. Evol.*, 27, 452–461.
Woodward, G., Ebenman, B., Emmerson, M., Montoya, J., Olesen, J., Valido, A., *et al.*
(2005). Body size in ecological networks. *Trends Ecol. Evol.*, 20, 402–409.

633
634

Figure captions

Fig. 1. Predicted ecological effects of genotypic richness in relation with functional richness. Functional richness is assumed to positively affect ecological functions, whereas genotypic richness might have different effects depending on it support cryptic functional diversity. First, if functional richness does capture the essential of the functional differences among genotypes, then genotypic richness would have no ecological effects (a). Second, if genotypic richness supports cryptic functional diversity, the effects could be (b) additive, (c) enhancing, or (d) compensatory.

Fig. 2. Effects of genotypic and functional richness on populations biomass production index (a), decomposition rate (b), algae stock (c), diversity of benthic invertebrates (e), diversity of zooplankton (f), abundance of benthic invertebrates (g), and abundance of zooplankton (h). Error bars represent ± 1 SE.

Fig. 3. Causal pathways between genotypic richness, functional richness, community structure, and ecosystem functioning. Two models have been run separately on decomposition rate (a), and pelagic algae stock (b). Only significant links are drawn, blue arrows represent positive links, yellow arrows negative links, and they are weighted according to the *p*-values extracted from the model.

Fig. 4. Effects of the diversity (a) and the abundance (b) of benthic invertebrates on decomposition rate; and effects of the diversity and the abundance of zooplankton on pelagic algae stocks (c and d, respectively). Points are partial residuals extracted from models

658   described in the statistical analysis section (see also Table S3). Straight lines represent the

659   slope predicted by the models (see statistical analyses) when significant ($\alpha < 0.05$), and grey

660   shadows represent 95% CIs.

(a) No effect   (b) Additive effect   (c) Enhancing effect   (d) Compensatory effect

HG = High genotypic richness

LG = Low genotypic richness

663 **Fig. 1.**

665
666 **Fig. 2.**

668
669 **Fig. 3**
670

671
672
673　**Fig. 4.**

674 **Table 1.** Results of the mixed effect linear models quantifying the relationships between

675 genotypic richness, functional richness and ecological parameters. Significant -and near

676 significant- *p*-values are displayed in bold. Interaction terms were removed from the model

677 when not significant.

Response	Effect	*p*-value	χ^2, d.f.
Fish biomass production	Genotypic richness	0.301	(1.071, 1)
	Functional richness	0.634	(0.227, 1)
	Genotypic richness* Functional richness	**0.057**	(3.621, 1)
	Mortality	**0.023**	(5.196, 1)
Benthic invertebrates diversity	**Genotypic richness**	**0.007**	(7.152, 1)
	Functional richness	**0.002**	(9.484, 1)
	Mortality	0.212	(1.551, 1)
Zooplankton diversity	**Genotypic richness**	**0.052**	(3.756, 1)
	Functional richness	**0.004**	(8.163, 1)
	Genotypic richness* Functional richness	**0.038**	(4.287, 1)
	Mortality	0.851	(0.035, 1)
Benthic invertebrates abundance	Genotypic richness	0.300	(1.072, 1)
	Functional richness	**0.031**	(4.667, 1)
	Mortality	0.238	(1.392, 1)
Zooplankton abundance	Genotypic richness	0.885	(0.021, 1)
	Functional richness	**0.004**	(8.154, 1)
	Mortality	0.220	(1.503, 1)
Decomposition rate	Genotypic richness	0.140	(2.169, 1)
	Functional richness	0.690	(0.158, 1)
	Mortality	0.893	(0.018, 1)
Algae stock	Genotypic richness	0.834	(0.044, 1)
	Functional richness	**0.053**	(3.742, 1)
	Mortality	0.412	(0.673, 1)

679
680

681   **Table 2.** Model fits of the two confirmatory path-analyses explaining decomposition rate and

682   algae stock variation. C-statistic (*C*), degree of freedom (*df*), and *p*-value are given for full

683   models as indication.

Variable	*C*	df	*p*-value	AIC full model	AIC alternative model	AIC simplified model
Decomposition rate	8.555	6	0.200	26.555	30.555	53.042
Algae stock	6.745	10	0.749	26.746	28.367	49.679