Title

Responsiveness variability during anaesthesia relates to inherent differences in brain structure

and function of the fronto-parietal networks

Authors

Feng Deng[a], Nicola Taylor[a], Adrian M. Owen[bd], Rhodri Cusack[a], Lorina Naci[ac*]

Affiliations

[a] Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,

Dublin, Ireland.

[b]Brain and Mind Institute, Western University, London, Canada.

[c]Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

[d] Department of Physiology and Pharmacology and Department of Psychology, Western

University, London, Canada.

[*]**Corresponding author:**

Lorina Naci

School of Psychology

Trinity College Institute of Neuroscience

Global Brain Health Institute

Trinity College Dublin

Dublin, Ireland

Telephone: +353 (0)87 688 5642

Email: nacil@tcd.ie

Abstract

Anaesthesia combined with functional neuroimaging provides a powerful approach for

understanding the brain mechanisms of consciousness. Although propofol is used

ubiquitously in clinical interventions that reversibly suppress consciousness, it shows large

inter-individual variability, and the brain bases of this variability remain poorly understood.

We asked whether three networks key to conscious cognition — the dorsal attention (DAN),

executive control (ECN), and default mode (DMN) — underlie responsiveness variability

under anaesthesia. Healthy participants (N=17) were moderately anaesthetized during

narrative understanding and resting state conditions inside the Magnetic Resonance Imaging

scanner. A target detection task measured behavioural responsiveness. An independent

behavioural study (N=25) qualified the attention demands of narrative understanding. 30% of

participants were unaffected in their response times, thus thwarting a key aim of anaesthesia

— the suppression of behavioural responsiveness. Individuals with stronger functional

connectivity within the DAN and ECN, between them, and to the DMN, and with larger grey

matter volume in frontal regions were more resilient to anaesthesia. For the first time, we

show that responsiveness variability during propofol anaesthesia relates to inherent

differences in brain structure and function of the fronto-parietal networks, which can be

predicted prior to sedation. Results highlight novel markers for improving awareness

monitoring during clinical anaesthesia.

Keywords

Consciousness; Frontal cortex; Fronto-parietal networks; Neuroimaging; Propofol

1. Introduction

Understanding how the human brain gives rise to consciousness remains a grand challenge

for modern neuroscience. A first step to understanding consciousness is to define what it is

not. To this end, anaesthesia combined with functional neuroimaging provides a powerful

approach for studying the brain mechanisms that change as consciousness fades (Vatansever

et al, 2020; Pal et al., 2020; Varley et al., 2020; Luppi et al., 2019; Demertzi et al., 2019;

Mashour and Hudetz 2018; Naci et al., 2018; MacDonald et al., 2015; Sarasso et al., 2015;

Stamatakis et al., 2010), and, conversely, that are necessary for realizing human

consciousness. Anaesthesia has been used for over 150 years to reversibly abolish

consciousness in clinical medicine, but its effect can vary substantially between individuals.

At moderate dosages, the intended suppression of behavioural responsiveness is highly

variable (Chennu et al., 2016; Bola et al., 2019), and at deep anaesthesia dosages, in rare

cases (0.1-0.2%; Mashour and Avidan 2015; Sandin et al., 2000), individuals retain conscious

awareness, also known as 'unintended intraoperative awareness' (Pandit et al., 2017; Sanders

et al., 2017; Sandin et a., 2000; Mashour and Avidan 2015). A much higher percentage of

patients presumed to be unconscious during general anaesthesia (22%; Leslie et al., 2007)

may have subjective experiences, such as dreaming. The brain bases of this considerable

inter-individual effect variability remain poorly understood.

A key and mostly overlooked question is what these individual differences (Searle and

Hopkins 2009; Palanca et al., 2009) can reveal about the unravelling of conscious cognition

as consciousness falters during anaesthesia. Propofol is the most common anaesthetic agent

in clinical interventions that require the reversible suppression of consciousness. Two recent

electroencephalography (EEG) studies that measured variable behavioural responsiveness

during mild and moderate propofol anaesthesia reported that the participants' varying levels

of responsiveness were differentiated by alpha band connectivity during wakefulness

(Chennu et al., 2016), and changing patterns of EEG signal diversity from wakefulness to

moderate sedation (Bola et al., 2019). However, the brain regional- or network-bases for this

substantial individual variability (Searle and Hopkins 2009; Palanca et al., 2009), and its

potential link to cognitive function, remain unknown. Propofol-induced sedation produces

selective metabolic impairment and reduces brain activity bilaterally in frontal and parietal

associative regions (Witon et al., 2020; Boly et al., 2008; Plourde et al., 2006; Alkire 2005;

Baars et al., 2003; Fiset et al., 1999; for a review see MacDonald et al., 2015; but see Pal et

al., 2020). Therefore, three brain networks with frontal and parietal lobe distribution, the

dorsal attention network (DAN), executive control network (ECN), and default mode

network (DMN) are primary candidate sources of individual variability under propofol

sedation.

The DAN and ECN, distributed laterally across frontal and parietal lobes, are key to

orchestrating stimulus-driven and goal-directed cognition. Their roles include orienting and

modulating attention to the saliency of incoming stimuli, monitoring and analysing

information from the environment and integrating it with internally generated goals, as well

as planning and adapting new behavioural schemas to take account of this information

(Duncan 2010; Elliott 2003; Corbetta and Schulman 2002; Kroger et al., 2002; Shallice

1988). The DMN extends partially in lateral and medial frontal and parietal lobes and is

thought to mediate internally-oriented cognition, e.g., autobiographical memory, imagination,

and thinking about the self (Andrews-Hanna et al. 2010; Schneider et al., 2008; Buckner et

al., 2008; Beer 2007; D'Argembeau et al., 2005; Wicker et al., 2003; Gusnard et al., 2001).

Recent studies show that it is also involved in external environment monitoring (Spreng et al.

2014; Buckner et al. 2008; Hahn et al., 2007), including encoding of scenes, episodes or

contexts (Vatansever et al., 2017), or shifts in contextually relevant information (Smith et al.,

2018). Recent studies that investigate state-related manipulations (conscious vs unconscious)

show that by, contrast to other brain networks, the DAN, ECN and DMN are selectively

impaired during loss of consciousness, not only under anaesthesia but also during severe

brain injury (Luppi et al., 2019; Naci et al., 2018). Thus, they provide strong evidence that

these three networks are part of a common brain mechanism disrupted during loss of

consciousness across vastly different conditions.

It is well established that the DAN/ECN and the DMN display an antagonistic relationship

(Huang et al., 2020; Fox et al., 2005; Fransson et al., 2005) during spontaneous thought and

goal-directed cognition. During conditions that engage external attention, the reduction of

functional activity in the DMN (Greicius et al., 2003, Raichle et al., 2001, Shulman et al.

1997), is concomitant with an increase of activity in the DAN and ECN (Seeley et al. 2007;

Dosenbach et al. 2007; Fox et al., 2005; Sridharan et al. 2008). The anticorrelation between

these is related to individual differences in performance variability (Kelly 2008), and directly

supports sustained attention (Kucyi et al., 2020). More broadly, it has been suggested to

facilitate ongoing switches between internally- and externally biased modes of attention

(Honey et al., 2017; Buckner et al., 2013) that contribute to consciousness (Carhart-Harris

and Friston 2010). Conversely, this antagonistic relationship breaks down in state-related

manipulations of consciousness, such as during sleep (Tagliazucchi et al., 2013), anaesthesia

(Bonhomme at el., 2012; Boveroux et al., 2010), and severe brain injury (Haugg et al., 2018;

Boly et al., 2009). Despite this accumulating evidence for the key and interrelated roles of the

DAN, ECN, and DMN in supporting conscious cognition (Huang et al., 2020; Demertzi et

al., 2013; Vanhaudenhuyse et al., 2011), and their primacy as target sites of propofol-induced

sedation, their roles in individual differences under anaesthesia have not been previously

investigated.

To address this gap, in two studies we tested whether variability or impairments in functional

connectivity within and between the DAN, ECN, and DMN underlie individual differences in

responsiveness during propofol anaesthesia. To directly investigate their individual and joint

roles, in a first functional Magnetic Resonance Imaging (fMRI) study we scanned healthy

participants (N=17) during wakefulness and during the administration of propofol at dosages

of 'moderate anaesthesia,' expressly aimed at engendering individual differences. Variability

in behavioural responsiveness in each state was assessed with an auditory target detection

task prior to scanning. To test whether any individual differences in responsiveness were

related to differences in perceptual or high-level attention processes that were invisible to the

clinical sedation scale (Ramsay et al., 1974), participants were scanned during an active

listening condition comprised of a brief (5 minutes) engaging auditory narrative, and during a

control condition of spontaneous thought, i.e., the resting state. Listening to plot-driven

narratives is naturally engaging, requires minimal behavioural collaboration from

participants, and therefore, is highly suitable for testing perceptual or high-level attention

processes independently of behavioural output or eye opening (Naci et al., 2018; 2017),

which are impaired in moderate anaesthesia. In a second behavioural study, we assessed the

high-level attention demands of narrative understanding in an independent behavioural group

(N=25), and related them to the brain activity of participants who underwent scanning.

Finally, to test whether individual differences under anaesthesia related to inherent brain

features independent from the propofol sedation, we investigated the functional connectivity

between and within the three networks during wakefulness, as well as grey matter volume

differences across participants.

2. Methods

2.1. Participants

Study 1. Healthy participants for the fMRI anaesthesia study (N=17; 18-40 years; 13 males) were tested in the MRI scanner at the Robarts Research Institute, Western University, in Canada. Ethical approval was obtained from the Health Sciences Research Ethics Board and Psychology Research Ethics Board of Western University. *Study 2*. An independent group of healthy participants (N=25; 18-40 years; 7 males) were tested behaviourally at the Global Brain health Institute at Trinity College Dublin, in Ireland. Ethical approval was obtained from the School of Psychology Research Ethics Board, Trinity College Dublin. All healthy participants were right-handed, native English speakers, had self-reported normal hearing and no history of neurological disorders. Informed consent was obtained for each participant prior to the experiment.

2.2. Stimuli and Design

Study 1

The study comprised a structural scan at the start, followed by two experimental sessions, one during wakefulness and the other during moderate anaesthesia, each with the same design. At the start of each session, a clinical assessment of sedation (Ramsay, 1974) was conducted to confirm wakefulness or moderate anaesthesia. Subsequently, a behavioural task, and two fMRI acquisitions, a free listening to an auditory narrative and a resting state scan, were conducted in each session. The narrative was the same in both sessions. The data analysed here was collected as part of a larger study, and the wakefulness data has previously been reported in relation to deep anaesthesia (Naci et al. 2018). See below for full protocol details.

Behavioural testing. *Auditory target detection task*. Before commencing the scanning for

either session, participants were asked to perform a computerized auditory target detection

task (50 trials), which aimed to assess individual responsiveness differences during moderate

anaesthesia. fMRI was not acquired during this task. Sound was presented with the

Sensimetrics (S14; www.sens.com) headphones. A volume level deemed comfortable by

each individual was determined prior to testing and used for the duration of the experiment.

Participants were instructed to press a button with their index finger as soon as they heard an

auditory beep (1200 Hz, 100 ms duration, 10 ms ramps) and to keep their eyes on the fixation

cross on the screen. Participant had up to 3000 ms to make a response. Once a response was

made, a pause of 1500 ms occurred prior to the presentation of the next stimulus. The

stimulus onset asynchrony (SOA) = min [reaction time (RT), 3000ms] + 1500 ms. The SOA

was jittered by virtue of variable RTs between trials. To ensure the participant did not lose

contact with the response box during moderate anaesthesia, the response box was attached

securely to their dominant hand. Individual and group RTs were used for behavioural

analyses.

MRI testing. Inside the MRI scanner, participants underwent two functional scans during

wakefulness and moderate anaesthesia. A plot-driven auditory narrative (5 minutes) was

presented over MRI compatible noise cancelation headphones (Sensimetrics, S14;

www.sens.com). Participants were asked to simply listen with eyes closed. The narrative

comprised a highly engaging auditory excerpt from the movie 'Taken' depicting dramatic

events, where a young girl travelling abroad without her family is kidnapped while speaking

on the phone to her father. The same narrative was presented during wakefulness and

moderate anaesthesia. A similar eyes-closed, resting state condition (8 minutes) was also

acquired in each state for comparison with the narrative condition.

Sedation Procedure. The level of sedation was measured with the Ramsay clinical sedation

scale for each participant by 3 independent assessors (two anaesthesiologists and one

anaesthesia nurse) in person inside the scanner room. Before entering the fMRI scanner, a

20G I.V. cannula was inserted into a vein on the dorsum of the non-dominant hand of the

participants. The propofol infusion system was connected to the cannula prior to the first

scanning session. No propofol was administered during the wakeful session. Participants

were fully wakeful, alert and communicated appropriately (Ramsay 1) and wakefulness (eye-

opening) was monitored with an infrared camera placed inside the scanner. At the

commencement of the moderate anaesthesia session, intravenous propofol was administered

with a Baxter AS 50 (Singapore). An effect-site/plasma steering algorithm was used

combined with the computer-controlled infusion pump to achieve step-wise increments in the

sedative effect of Propofol. This infusion pump was manually adjusted to achieve the desired

levels of sedation, guided by targeted concentrations of Propofol, as predicted by the TIVA

Trainer (the European Society for Intravenous Anaesthesia, eurosiva.eu) pharmacokinetic

simulation program. The pharmacokinetic model provided target-controlled infusion by

adjusting infusion rates of Propofol over time to achieve and maintain the target blood

concentrations as specified by the Marsh 3 compartment algorithm for each participant, as

incorporated in the TIVA Trainer software (Marsh et al., 1991). In accordance with the

Canadian Anaesthesia Society (CAS) guidelines, non-invasive blood pressure (NIBP), heart

rate, oxygen saturation (SpO2) and end-tidal carbon dioxide (ETCO2) were monitored

continuously through the use of a dedicated MR compatible anaesthesia monitor. Complete

resuscitation equipment was present throughout the testing.

Propofol infusion commenced with a target effect-site concentration of 0.6 μg/ml and oxygen

was titrated to maintain SpO2 above 96%. Throughout sedation, participants remained

capable of spontaneous cardiovascular function and ventilation. Supplemental oxygen was

administered via nasal cannulae to ensure adequate levels of oxygen at all times. If the

Ramsay level was lower than 3 for moderate sedation, the concentration was slowly

increased by increments of 0.3 μg/ml with repeated assessments of responsiveness between

increments to obtain a Ramsay score of 3. During administration of propofol, participants

generally became calm and slowed in their response to verbal communication. Once

participants stopped engaging in spontaneous conversation, and speech became sluggish, they

were classified as being at Ramsey level 3. Nevertheless, when asked via loud verbal

communication, participants agreed to perform the auditory target detection task as would be

expected at Ramsey level 3. A brief recall task was adopted from the Mini Mental State

Exam (Folstein et al., 1975), as a behavioural test complimentary to the anaesthesiologists'

assessments of whether sedation commensurate with Ramsay level 3 had been achieved. At

the start of each session (wakefulness/moderate anaesthesia sessions), the researcher named

three different sets of unrelated objects clearly and slowly and asked the participant to name

each of them. The participant was instructed to remember the words in order to be able to

repeat them in a short while. For 10 minutes following the word presentation, the participant

was allowed to rest and performed no other task/was not exposed to any experimental stimuli.

Subsequently, the participant was asked to repeat the words. Two different lists (a. Ball-

Flag-Tree; b. Flower-Egg-Rope), counterbalanced across participants, were used to avoid

familiarity effects between the wakefulness and moderate anaesthesia states. As expected,

during the wakefulness session, all participants correctly repeated all three words, whereas

during the moderate anaesthesia session, performance was varied with participants sluggishly

repeating one or more of the words, consistent with Ramsay level 3 sedation. During

moderate anaesthesia (Ramsey 3), the mean estimated effect-site propofol concentration, as

provided by the TIVA Trainer software, was 1.99 (1.59- 2.39) μg/ml and the mean estimated

plasma propofol concentration was 2.02 (1.56- 2.48) μg/ml.

Study 2

To provide a subjective measure of attention during the same story, an independent

behavioural group of participants that did not undergo scanning, rated how "suspenseful" this

story was every 2 seconds, from 'least' (1) to 'most suspenseful' (9). The audio excerpt was

divided into 156 clips, each 2 seconds long to match the repetition time (TR, 2 sec) used in

the independent participant group from Study 1 (who underwent MRI scanning at the Robarts

Research Institute, at Western University, in Canada). Participants heard the stimuli through

over-ear headphones connected to the stimulus presentation computer and used the keyboard

to record responses, in a sound-isolated laboratory at the Global Brain Health Institute, at

Trinity College Dublin, in Ireland. Participants had up to 3000 ms to make a response, at

which point the next sequential clip began immediately. At the end of the experiment,

participants indicated via a feedback questionnaire that the interruptions did not disrupt the

coherence of the story's plot and their perception of suspense throughout.

2.3. Data Analyses

Study 1

Analysis of behavioural data. As responses beyond 3000ms were not monitored, for the

purposes of behavioural analyses, we considered participants who did not respond within

3000ms during anaesthesia as very delayed in their response time rather as missing data

points. Participants were awake and responsive but very sluggish, consistent with moderate

anaesthesia/Ramsay level 3 designation, and, therefore, it is very likely that responses were

made outside the 3000ms window. Individual-level differences in reaction times (RTs) of the

target detection task between wakefulness and moderate anaesthesia sessions were assessed

by independent samples t-tests. The rate of RT change from baseline was calculated as

follows: [(RT during anaesthesia - RT during wakefulness)/RT during wakefulness]*100.

The hit rate was calculated as the number of responses divided by the number of trials for

wakefulness and moderate anaesthesia separately. The Mahalanobis distance was computed

to detect outliers in our multivariate dataset (Hadi 1992), comprised of the rate of RT change

from baseline and hit rate during anaesthesia. Participants were considered outliers if their

robust Mahalanobis distance from the rest of the distribution was significant at $p < 0.01$.

MRI Data Acquisition. Functional images were obtained on a 3T Siemens Prisma system,

with a 32-channel head coil. The high-resolution brain structural images were acquired using

a T1-weighted 3D MPRAGE sequence with the following parameters, voxel size: 1 x 1 x 1

mm, TA=5 minutes and 38 seconds, echo time (TE) = 4.25ms, matrix size=240x256x192,

flip angle (FA) = 9 degrees. The functional echo-planar images (EPI) images were obtained

with following parameters, 33 slices, voxel size: 3 x 3 x 3, inter-slice gap of 25%, repetition

time (TR) = 2000ms, TE=30ms, matrix size=64x64, FA=75 degrees. The audio narrative and

resting state had 155 and 256 scans, respectively.

MRI preprocessing: 1) Functional data. Standard preprocessing procedures and data analyses

were performed with SPM8 (Statistical Parametric Mapping; Wellcome Institute of Cognitive

Neurology, http://www.fil.ion.ucl.ac.uk/spm/software/spm8/) and the AA pipeline software

(Cusack et al., 2015). In the preprocessing pipeline, we performed slice timing correction,

motion correction, registration to structural images, normalization to a template brain, and

smoothing. The data were smoothed with a Gaussian smoothing kernel of 10mm FWHM

(Peigneux et al. 2006). Spatial normalization was performed using SPM8's segment-and-

normalize procedure, whereby the T1 structural was segmented into grey and white matter

and normalized to a segmented MNI-152 template. These normalization parameters were

then applied to all EPIs. The time series in each voxel was high pass-filtered with a cutoff of

1/128 Hz to remove low-frequency noise, and scaled to a grand mean of 100 across voxels

and scans in each session. Prior to analyses, the first five scans of each session were

discarded to achieve T1 equilibrium and to allow participants to adjust to the noise of the

scanner. To avoid the formation of artificial anti-correlations (Murphy et al. 2009; Anderson

et al., 2011), we performed no global signal regression. 2) Structural data. The brain

structural images were processed using FreeSurfer package

(https://surfer.nmr.mgh.harvard.edu/), a well-documented automated program which is

widely used to perform surface-based morphometric analysis (Dale et al. 1999). The

processing steps include: 1) removing the non-brain tissue; 2) transforming the skull-

stripping brain volume to Talairach-like space; 3) segmenting brain tissues to GM, WM,

cerebrospinal fluid (CSF); 4) performing intensity normalization to remove the effect of bias

field; 5) building a surface tessellation to generate a triangular cortical mesh consisting of

about 300,000 vertices in the whole brain surface; 6) correcting topological deficits of

cortical surface; and 7) deforming brain surface to generate optimized models of GM/WM

and GM/CSF boundaries.

Analyses of fMRI data. Group-level correlational analyses explored, for each voxel, the cross-

subject synchronization of brain activity by measuring the correlation of each participant's

time-course with the mean time-course of all other participants (Naci et al., 2018; 2017;

2014). Further, to investigate whether the behavioural variability under anaesthesia related to

perceptual or higher-order processing differences among participants, we performed two

mixed data-driven and model-based analyses. First, we extracted the sound envelope of the

auditory narrative via the Matlab *MIR*toolbox

(http://www.jyu.fi/hum/laitokset/musiikki/en/research/coe/materials/mirtoolbox) and built a

generalised linear model (GLM) by using statistical parametric mapping to derive auditory

characteristic-related brain activation for each individual. Subsequently, the z-scored average

suspense ratings of the narrative obtained by the independent group of participants in study 2

(see SI) were used as a regressor in the fMRI data of individuals who underwent propofol

anaesthesia in study 1, to measure the neural correlates of perceptual or high-level attention

processes during the narrative condition. The regressors were generated by convolving

boxcar functions with the canonical hemodynamic response function (Friston et al., 1989).

Also included in the general linear model were nuisance variables, comprising the movement

parameters in the three directions of motion and three degrees of rotation, as well as a

constant (all-ones vector) that served to regress out the session mean. Fixed-effect analyses

were performed in each subject, corrected for temporal auto-correlation using an AR

(1)+white noise model. Linear contrasts were used to obtain subject-specific estimates for

each effect of interest. Linear contrast coefficients for each participant were entered into the

second level random-effects analysis. Clusters or voxels that survived at $p < 0.05$ threshold,

corrected for multiple comparisons with the family-wise error (FWE) were considered

statistically significant. For thresholding at the cluster level, an uncorrected cluster forming

voxel threshold of $p < 0.001$ was used.

Functional connectivity (FC) within and between the DAN, ECN, and DMN was assessed by

computing the Pearson correlation of the fMRI time courses between 19 regions of interests

(ROIs; spherical, 10mm diameter) (from Raichle 2011) constituting these different brain

networks, as identified by resting state studies (Table 1). This parcellation method is

theoretically-driven based on meta-analyses of these three particular networks, and,

furthermore, helps to relate our current findings to our previous findings based on the same

parcellation method (Naci et al., 2018; Haugg et al., 2018). Permutation (1000 times) tests

were used to explore FC difference between conditions and were false discovery rate (FDR)

corrected for multiple comparisons. All of the analyses were conducted using Fisher z-

transformed correlation (Pearson r). Glass's delta was used to compute the effect size of the

comparison between wakeful and moderate anaesthesia states, because of the different

standard deviations of the two states, while Hedges' g was used to compute the effect size of

the comparison between fast and slow participants because of the different sample sizes for

the two groups.

Table 1. Overview of selected regions of interests for the three functional networks

Network/Region	ROI	MNI coordinates		
		x	y	z
Default Mode Network	Posterior cingulate/precuneus	0	-52	27
	Medial prefrontal	-1	54	27
	Left lateral parietal	-46	-66	30
	Right lateral parietal	49	-63	33
	Left inferior temporal	-61	-24	-9
	Right inferior temporal	58	-24	-9
Dorsal Attention Network	Left frontal eye field	-29	-9	54
	Right frontal eye field	29	-9	54
	Left posterior IPS	-26	-66	48
	Right posterior IPS	26	-66	48
	Left anterior IPS	-44	-39	45
	Right anterior IPS	41	-39	45
	Left MT	-50	-66	-6
	Right MT	53	-63	-6
Executive Control Network	Dorsal medial PFC	0	24	46
	Left anterior PFC	-44	45	0
	Right anterior PFC	44	45	0
	Left superior parietal	-50	-51	45
	Right superior parietal	50	-51	45

357   Abbreviations: IPS: Intraparietal sulcus, MT: Middle temporal area, PFC: Prefrontal cortex

Grey matter volume analyses. Vertex-wise grey matter volume (GMV) was computed

(Winkler et al., 2018) and smoothed with a 10-mm full-width at half maximum Gaussian

kernel to perform the statistics. Monte Carlo simulation cluster analysis and a cluster-wise

threshold of $p < 0.05$ was adopted for multiple comparisons correction (Hagler et al., 2006).

Then, averaged GMV values for each significant cluster, for each individual participant, was

extracted to perform permutation tests (1000 times) that tested for clusters/regions showing

significant GMV differences between fast and slow participants. The FDR method was used

to correct for multiple comparisons.

Study 2

To determine how similar suspense ratings were across the group, the inter-subject

correlation of suspense ratings was computed as the average of the Pearson correlations of

each participant's data with the mean data from the rest of the group To account for the

non-normalized distribution of correlation values (Fisher 1915), all statistical analyses were

performed on z-transformed correlation values, using Fisher's r-to-z transformation. For

visualization purposes, we re-transformed these z-values in correlation values.

3. Results

3.1 The effect of moderate anaesthesia on behavioural responsiveness

Despite the same Ramsay 3 clinical score, independently determined by three assessors, we

observed significant heterogeneity in the RTs of the auditory detection task (Figure 1A).

5/17 participants were not delayed significantly relative to their wakeful responses, 9/17 were

significantly delayed and had fragmented responses (showing 2–40% missing trials), and

3/17 failed to make any responses within the 3000ms time window (Figure 1A–B), despite

agreeing to have understood the task instructions and to make responses via the button box

affixed to their dominant hand. This large individual variability was at odds with the propofol

infusion rates titrated for each participant based on the pharmacokinetic model adjusted for

demographic variables (Marsh et al., 1991), to maintain stable target blood concentrations

consistent with Ramsay 3 sedation level (Ramsay, 1974).

Based on the rate of RT change from the wakeful baseline and hit rates during anaesthesia

(Figure 1B), multivariate data outlier detection showed that 5 participants had significantly

different robust Mahalanobis distance from the rest (Figure 1C). On this basis, individuals

were divided into two groups for further analyses: 'fast' participants (FPs; N=12), and 'slow'

participants (SPs; N=5). Individual responsiveness variability was not explained by

demographic differences, as suggested by no differences in age (between-samples t-test, $t =$

1.3, $p = 0.2$) and gender (Fisher's exact test, odds ratio $= 1.3$, $p = 1$) between FPs and SPs.

Behavioural Responses

Figure 1. Behavioural responses. (A) Reaction times and (B) hit rates during the auditory

target detection task in the wakeful (blue) and moderate anaesthesia (orange) states. (C)

Multivariate outlier detection based on the robust Mahalanobis distance measure. Green/blue

dotted lines depict the significance threshold of $p<0.05/p<0.01$. For further analyses,

participants that showed robust Mahalanobis distance significantly different from the rest at

$p<0.01$ (above the blue line) were labelled 'slow' participants (N=5), whereas the rest (below

the blue line) were labelled 'fast' participants (N=12). The horizontal axis represents each

individual (S1-S17=Subject1-Subject17).

Rather, the brain bases of this variability may be related to three other factors. First, the

individual variability could be related to underlying differences in perceptual or high-level

attention processes, which may have been invisible to the behavioural examination during the

Ramsay assessment. Second, responsiveness differences may be due to inherent connectivity

differences prior to sedation, and/or alterations of connectivity due to sedation, within and

between the DAN, ECN and DMN networks. Third, individual responsiveness differences

may be related to inherent structural brain differences, which may, in turn, also manifest as

differences in functional connectivity. In the following analyses, we explored each of these

factors in turn.

3.2 Perceptual and high-level attention processes and behavioural responsiveness during

moderate anaesthesia

We investigated whether each individual's sensory-driven auditory perceptual processes, and

high-level attention processes during the narrative condition related to their response times

during the target detection task. First, we used SPM to model the relationship between the

narrative's perceptual properties captured by the sound envelope, and changes in brain

activity over time. Then, each individual's brain activity was related to their RTs in the target

detection task. The sound envelope predicted brain activity in bilateral auditory cortex and

right inferior frontal gyrus during wakefulness (Figure 2A), which as expected was

dramatically reduced in moderate anaesthesia, to one small subthreshold cluster in left

auditory cortex (p=0.05 FWE corrected) (Figure 2B). We observed no correlation between

the participants' reaction times in either wakefulness or moderate anaesthesia and the extent

of their individual activations in auditory regions (Figure 2E–F).

Figure 2. Perceptual and high-level attention processes during the narrative. At the group

level, the sound envelope of the narrative predicted significant (p<0.05; FWE corrected)

clusters in the (A) auditory cortex and left inferior frontal gyrus during wakefulness, and (B)

left auditory cortex in moderate anaesthesia [did not reach statistical significance (p=0.05

FWE corrected)]. (C)/(D) At the group level, the suspense ratings of the audio narrative

predicted significant (p<0.05; FWE corrected) clusters in the (C) auditory attention and

salience networks during wakefulness, and (D) auditory attention network during moderate

anaesthesia. Red arrows indicate the location of the coronal views with respect to the

anterior-posterior dimension. (E)–(H) Panels display the number of voxels in primary

auditory regions, auditory attention regions, and salience networks of individual participants

that were predicted by the sound envelope and suspense ratings, relative to each individual's

reaction time in the target detection task. At the individual level, the sound envelope

predicted significant activations in the auditory regions in (E) 14/17 of participants during

wakefulness, and (F) 10/17 of participants during moderate anaesthesia. At the individual

level, the suspense ratings predicted significant activations in the auditory attention and

salience network regions in (G) 17/17 of participants during the wakefulness and (H) 14/17

of participants during moderate anaesthesia. There was no correlation between the perceptual

or higher-order processes during the narrative and reaction times, during the target detection

task, either in wakefulness (E)/(G) or in moderate anaesthesia (F)/(H). Abbreviations: AAN:

Auditory Attention Network; SN: Salience Network.

Second, to investigate differential recruitment of high-level attention processes during the

narrative, we used a previously established qualitative measure of the listeners' sustained

attention throughout the narrative — the perception of suspense on a moment-by-moment

basis (Naci et al., 2014; 2017). We measured the perception of suspense in Study 2, in an

independent group from those who underwent scanning. Suspense ratings throughout the

narrative were highly and significantly correlated across individuals (r=0.90; SE=0.07)

[t(24)=20.56, p<0.0001], confirming the similar high-level attention processes, and

understanding of the narrative across different individuals. We used SPM to model the

relationship between this qualitative measure of the narrative's ongoing high-level attention

demands and changes in brain activity over time, during the wakeful and moderate

anaesthesia states. At the group level, during wakefulness, suspense ratings significantly

(p<0.05; FWE corrected) predicted activity in regions of the auditory attention network

(Tobyne et al., 2018; Michalka et al., 2015; Naci et al., 2013) and the salience network

(Seeley 2019; Menon 2015) (Figure 2C), with clips rated as 'highly suspenseful' predicting

stronger activity in these networks (Table 2). By contrast, during moderate anaesthesia, one

small activation cluster was observed (Figure 2D).

Table 2. Coordinates for activation clusters associated with the perception of suspense during

the narrative.

Cluster Index	Cluster Peak Location	Peak MNI			Cluster-wise p-value
		x	y	z	
1	Cingulate Sulcus	-2	20	36	<0.0001
2	Right Superior Temporal Gyrus	58	-36	14	<0.0001
3	Right Anterior Insular Gyrus	40	24	8	<0.0001
4	Posterior Cingulate Gyrus	14	-44	14	<0.0001
5	Left Anterior Insular Gyrus	-36	14	-8	<0.0001
6	Thalamus	4	-20	10	<0.005

As previously established (Naci et al. 2014; 2017; 2018), the similarity of the narrative's

suspense ratings enabled model-based predictions of the underlying brain activity, from the

group data, in a leave-one-out fashion to individual participants. Using this approach, we

related the activity in suspense-driven activation clusters for each individual to their

respective RTs during the target detection task. No correlation was observed between the

participants' activation in the auditory attention and salience networks regions to their RTs

(Figure 2 G–H). Taken together, these results strongly suggested that differences in

behavioural responsiveness under anaesthesia were not driven by differences in perceptual or

high-level attention processes.

3.3 Network connectivity and behavioural responsiveness during moderate anaesthesia

Rather, alterations in the connectivity within and between the DAN, ECN, and DMN during

moderate anaesthesia (Figure S1, Supplemental Information) may drive these individual

differences (Figure 3). In the narrative condition, during wakefulness, functional connectivity

within the DAN ($p < 0.005$), within the ECN ($p < 0.005$), between the DMN and DAN ($p <$

0.05), between DMN and ECN ($p < 0.05$), and between DAN and ECN ($p < 0.05$)

differentiated the two response groups, with significantly higher FC for FPs relative to SPs

(Figure 3B). During moderate anaesthesia, functional connectivity within the DMN ($p <$

0.05), within the ECN ($p < 0.05$), and between DAN and ECN ($p < 0.05$) differentiated FPs

from SPs, with significantly higher FC for FPs relative to SPs (Figure 3B). In the resting state

condition, during wakefulness, no differences were observed between FP and SP in any of

connectivity types. During moderate anaesthesia, functional connectivity within and between

the DAN, ECN, and DMN differentiated the two groups (DMN, $p < 0.05$; DAN, $p < 0.05$;

ECN, $p < 0.05$; DMN-DAN, $p < 0.005$; DMN-ECN, $p < 0.005$; DAN-ECN, $p < 0.005$), with

significantly higher FC for FPs relative to SPs (Figure 3B). All the above were permutation

tests FDR corrected.

Figure 3. Functional connectivity differences between fast and slow participants. (A) FC

matrices for the fast and slow participants during wakeful and moderate anaesthesia states in

narrative and resting state conditions separately. The three black and white patterned

rectangles on the left hand side represent, from top to bottom, the DMN, DAN and ECN. The

colour bar indicates z-transformed Pearson correlation strength (low-high: blue-red). Each

cell in the matrix represents FC between each node of the three networks. (B) Comparisons

of averaged FC within and between the three brain networks, between fast and slow

participants for each condition and state separately. * = p<0.05 FDR corrected for multiple

comparisons. Abbreviations: DMN, default mode network; DAN, dorsal attention network;

ECN, executive control network.

A 2x2x2 ANOVA explored the effects of *condition* (narrative, resting state), *state*

(wakefulness, moderate anaesthesia) and *response group* (FPs, SPs), and their interactions.

We found a main effect of response group (F (1,15) = 19.1, p < 0.005) that was driven by

higher connectivity for FPs, and no effects on condition or state. However, effect size

analyses showed higher difference between FPs and SPs in the narrative condition during

wakefulness (Hedges' g = 3.1) over all other conditions, including relative to moderate

anaesthesia (Hedges' g = 1.1), and relative to the resting state condition during wakefulness

(Hedges' g = 0.9), and moderate anaesthesia (Hedges' g = 2.3).

3.4. Structural brain differences and behavioural responsiveness during moderate anaesthesia

Finally, we tested whether individual differences in behavioural responsiveness were related

to structural brain differences, which may also underlie functional connectivity differences.

To this end, we performed whole-brain vertex-based comparisons of grey matter volume

between FPs and SPs. We found that FPs had uniformly significantly higher grey matter

volume relative to the SPs, in two frontal regions, the left superior and dorsolateral frontal

cortex (L SFC), including the presupplementary motor area (pre-SMA), and the left rostral

middle frontal cortex (L rMFC) (Figure 4A, Table 3). The averaged GMV for each of these

two regions were higher for FPs relative to SPs (L SFC, $p < 0.0005$, L rMFC, $p < 0.0005$;

FDR corrected, Figure 4B). The opposite effect, i.e., higher grey matter for SP than FP, was

not observed anywhere in the brain. The L SFC and L rMFC overlapped with the functional

ROIs that defined the frontal nodes of the executive control network (Table 1), further

supporting the functional connectivity results above (Figure 3). Together they strongly

suggested that inherent brain structural and functional features of the fronto-parietal

networks, especially the ECN, underlies individual differences in behavioural responsiveness

under moderate anaesthesia.

Figure 4. Grey matter volume differences between fast and slow participants. (A) Vertex-

wise analysis. Two clusters showed significant between-group difference in GMV (p < 0.05,

Monte Carlo simulation corrected). Colour bar indicates the p-value (-lg(p)). (B) Averaged

GMV was extracted for each of the two clusters, and between-group comparisons of the

averaged GMV were performed by using permutation tests. *** = p<0.0005 FDR corrected

for multiple comparisons. Abbreviations: L SFC, left superior frontal cortex; L rMFC, left

rostral middle frontal cortex.

Table 3. Regions showing significant grey matter volume difference between fast and slow

participants.

Index	Peak location	Vertex number	Size (mm^2)	Peak MNI coordinates			Cluster-wise p-value
				x	y	z	
1	L_SFC	3910	2217.98	-8	29	52	<0.0001
2	L rMFC	1542	937.67	-39	26	21	0.04

P values represent the cluster-wise corrected p for the peak vertex. Abbreviations: L SFC, left

superior frontal cortex; L rMFC, left rostral middle frontal cortex.

This was further confirmed by a direct comparison of the FC differences between FPs and

SPs in the ROIs covering the frontal (dorsal medial PFC, left and right anterior PFC) and

parietal (left and right superior parietal cortex) aspects of the ECN separately. A 2x2x2

ANOVA with factors *state* (wakefulness, moderate anaesthesia), *region* (frontal ECN,

parietal ECN), and *response group* (FPs, SPs) showed a main effect of region [F (1,15) =

11.9, p < 0.005], driven by higher connectivity in the parietal aspect of the ECN, and a main

effect of group [$F(1,15) = 8.2, p < 0.05$)], driven by higher connectivity in FPs relative to

SPs. Direction comparisons between FPs and SPs in frontal and parietal aspects of ECN,

showed that FC within the frontal, but not parietal, aspect of the ECN distinguished FPs and

SPs during wakefulness and during moderate anaesthesia ($p < 0.05$, FDR corrected) (Figure

5).

572 **Figure 5**. Differences of functional connectivity in frontal and parietal aspects of executive

573 control network between fast and slow participants, in wakeful and moderate anaesthesia

574 states in narrative condition. * = p <0.05 FDR corrected for multiple comparisons.

4. Discussion

Although anaesthesia has been used for over 150 years to reversibly abolish consciousness in clinical medicine, the brain bases of its considerable inter-individual effect variability remain poorly understood. To address this gap, we asked whether the connectivity within and between three networks that are primary sites of propofol-induced sedation and key to conscious cognition — the DAN, ECN and DMN — underlie individual responsiveness differences under anaesthesia. Consistent with previous studies (Chennu et al., 2016; Bola et al., 2019), we observed substantial individual differences in reaction times and hit rates under moderate anaesthesia. These were not related to differences in perceptual or high-level attention processes. Rather, the inherent level of functional connectivity during the narrative condition prior to sedation, within the DAN and ECN, between them, and to the DMN differentiated the participants' responsiveness level. Furthermore, grey matter volume in the frontal cortex aspect of the ECN was significantly related to behavioural responsiveness during sedation. Participants whose reaction times showed little change from the wakeful baseline had significantly higher grey matter volume in these regions than participants who showed big changes. Therefore, for the first time, we show that responsiveness variability during propofol anaesthesia relates to differences in inherent brain structural and functional features of the fronto-parietal networks, which can be predicted prior to sedation. Additionally, we found that during moderate anaesthesia, the connectivity within and between the DAN, ECN and DMN differentiated fast and slow participants, suggesting that perturbations by anaesthesia further exacerbate inherent inter-individual functional connectivity differences.

The differential impact of moderate anaesthesia on cognitive process can be inferred in a post hoc manner, based on the perceptual and cognitive processes recruited by the target detection

601 task and the brain networks and regions that showed functional and/or structural differences

602 between the participants. The task involved a hierarchy of perceptual and cognitive

603 processes. These included (a) auditory perception, (b) understanding of linguistic instruction,

604 (c) short-term memory to remember instructions for the duration of the experiment, (d)

605 attention to auditory stimuli, (e) attention to intention to make a response, (f) cognitive

606 control over other, competing and distracting mental processes, and the (g) execution of a

607 motor response, when prompted by the auditory stimuli. The clinical assessment determined

608 similar basic language communication (b), and short-term memory function (c) across

609 participants during moderate anaesthesia. Furthermore, our results showed no relationship

610 between behavioural responsiveness and sensory/auditory (a), or higher-order, including

611 linguistic processes that relied on sustained attention (b, d), across participants. Therefore,

612 differences in responsiveness were likely underpinned by differences in complex mental

613 faculties, such as attention to intention (e), cognitive control (f) and action execution (g).

615 Consistent with this inference, our results demonstrated that FC within and between the DAN

616 and ECN differentiated the participants' level of responsiveness, suggesting that moderate

617 anaesthesia impacts attention to intention and goal and action execution, subserved by the

618 DAN and ECN (Marek and Dosenbach 2018; Dosenbach et al., 2007; Ernst and Paulus

619 2005). Furthermore, our structural imaging results demonstrated that grey matter volume

620 differences in the frontal cortex aspect of the ECN, including the DLPFC and pre-SMA,

621 underscore individual responsiveness differences. In addition to their role in action selection

622 and execution (Lee et al., 1999; Boly et al., 2007), the DLPFC and pre-SMA represent

623 higher-order attention, i.e., attention to intention. Lau et al (2004) investigated the brain

624 regions that were preferentially involved in attending to the intention to move, relative to the

625 actual movement, and found that brain activity in pre-SMA, as well as its functional

connectivity with right DPFC represented the intention to move. The pre-SMA has also been

implicated in sustained cognitive control (Nachev et al., 2005), another process that is key to

successful task performance. Therefore, a post-hoc interpretation of our findings is that

moderate anaesthesia impacts attention to intention, sustained cognitive control, and action

execution, and that individuals with smaller grey matter volume in frontal regions and weaker

functional connectivity within and between the DAN and ECN, may have a vulnerability for

stronger suppression of behavioural responsiveness than those with higher values for these

features. Additionally, the finding DMN connectivity during moderate anaesthesia

differentiated fast and slow participants suggests that individual differences in responsiveness

may also reflect the effect of sedation on self-referential processes (Andrews-Hanna et al.

2010; Schneider et al., 2008; Buckner et al., 2008; Beer 2007; D'Argembeau et al., 2005;

Wicker et al., 2003; Gusnard et al., 2001), and context processing (Smith et al., 2018;

Vatansever et al., 2017), which are supported by the DMN.

As the suppression of behavioural responsiveness is a key aim of anaesthesia, the converse

finding was also highly relevant. Surprisingly, we found that 30% of participants showed no

delay in reaction times under moderate anaesthesia relative to wakefulness, and critically,

these exhibited inherent brain differences to participants who were significantly delayed. This

result is highly relevant to the prediction of responsiveness variability under anaesthesia

(Chennu et al., 2016; Palanca et al 2009) and monitoring depth-of-anaesthesia for the

detection of unintended intraoperative awareness. Although rare (0.1-0.2%, Mashour and

Avidan 2015; Sandin et al., 2000), unintended awareness can be very traumatic and lead to

negative long-term health outcomes, such as post-traumatic stress disorder (up to 70%), as

well as clinical depression or phobias (Pandit et al., 2017; 2014; Mashour and Avidan 2015).

Given its rarity, unintended intraoperative awareness does not lend itself to direct

investigation in the relatively small groups of typical research studies with vulnerable

populations, including the present study, where individuals were deeply anaesthetized

without intubation in a research MRI setting.

Our results suggest that individuals with larger grey matter volume in frontal regions and

stronger functional connectivity within DAN and ECN, between them and to the DMN, may

require higher doses of propofol to become non-responsive to the same extent as individuals

with weaker connectivity in these networks and smaller grey matter volume in frontal cortex.

If replicated in a clinical context, these findings will provide novel markers that may help to

improve the accuracy of awareness monitoring during clinical anaesthesia. We note that

future clinical studies ought to measure the clearance of anaesthetic agents by each individual

directly for, potentially, a more precise assessment of the metabolic variation across

participants, than that provided by the anaesthesia delivery instrument used here. It is also

worth noting that propofol was used here due to its wide prominence in clinical interventions,

and future studies that employ the same paradigm across different agents will determine

whether the brain-behaviour relationships we report here generalize to other anaesthetic

agents.

Finally, we observed higher power for detecting individual responsiveness differences under

anaesthesia in the wakeful narrative condition, than in the resting state. This may be due to

higher signal to noise during the narrative condition. Previous studies have shown that the

active attention engagement during well-crafted narratives leads to reduced movement

(Centeno et al., 2016) and less sleep in scanner (Centeno et al., 2016, Vanderwal et al., 2015),

resulting in higher signal-to-noise relative to resting state (rs) fMRI (Wang et al., 2017).

Moreover, it may be due to higher power to detect functional roles of brain networks during

naturalistic stimuli relative to rsfMRI. Rs fMRI is entirely unconstrained, and thus it is

difficult to separate signal due to cognitive processes from extraneous sources, including

motion, cardiac and respiratory physiological noise, arterial CO_2 concentration, blood

pressure, and vasomotion (Murphy et al., 2013; Birn et al., 2006). By contrast, as we have

previously established, naturalistic narratives drive diverging responses across brain

networks, highlighting their functionally distinct roles, and thus may be more sensitive for

investigating how their differing cognitive roles are impacted by anaesthesia than rsfMRI

(Naci et al., 2018; Haugg et al., 2018; Naci et al., 2014). Our results are consistent with a

growing recognition of the importance of naturalistic stimuli for studying cognitive processes

and understanding the neural basis of real-world functioning (Sonkusare et al., 2019).

Funding

This work was supported by the Provost PhD Award Scheme from Trinity College Dublin, an

L'Oreal for Women In Science International Rising Talent Award, and the Welcome Trust

Institutional Strategic Support grant to L.N and by the Canadian Institute for Advanced

Research (CIFAR), the Canadian Institutes for Health Research (CIHR), the Natural Sciences

and Engineering Research Council of Canada (NSERC), and a Canada Excellence Research

Chairs award (Grant No. 215063) to AMO and by an ERC Advanced Grant (FOUNDCOG)

(Grant No. 787981) to R.C..

Conflict of Interest

The authors declare no conflict of interest.

Data Availability Statement

In study 1, reaction time and fMRI data were collected in 2015, under ethics from Western

University, Canada. This stipulated that data would be shared only with the research team

directly involved in this study. Therefore, individual data cannot be freely shared due to

ethics protocol regulation. However, we have uploaded group maps to NeuroVault

(https://identifiers.org/neurovault.collection:9576).

In study 2, suspense ratings data were collected under ethics from Trinity College Dublin,

Ireland, which stipulated that anonymized data can be shared with the scientific community.

Therefore, the behavioral data and code will be shared upon contacting the authors. The

sharing practices are in line with the guidelines from the funding body.

References

Alkire, M. T., & Miller, J. (2005). General anesthesia and the neural correlates of

consciousness. *Progress in brain research*, 150, 229-597.

Anderson, J. S., Druzgal, T. J., Lopez-Larson, M., Jeong, E. K., Desai, K., & Yurgelun-Todd,

D. (2011). Network anticorrelations, global regression, and phase-shifted soft tissue

correction. *Human brain mapping*, *32*(6), 919-934.

Andrews-Hanna, J. R., Reidler, J. S., Huang, C., & Buckner, R. L. (2010). Evidence for the

default network's role in spontaneous cognition. *Journal of neurophysiology*, *104*(1), 322-

335.

Baars, B. J., Ramsøy, T. Z., & Laureys, S. (2003). Brain, conscious experience and the

observing self. *Trends in neurosciences*, 26(12), 671-675.

Beer, J. S. (2007). The default self: feeling good or being right?. *Trends in cognitive*

sciences, *11*(5), 187-189.

Birn, R. M., Diamond, J. B., Smith, M. A., & Bandettini, P. A. (2006). Separating

respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in

fMRI. *Neuroimage*, 31(4), 1536-1548.

Bola, M., Orlowski, P., Plomecka, M., & Marchewka, A. (2019). EEG signal diversity during

propofol sedation: an increase in sedated but responsive, a decrease in sedated and

unresponsive subjects. *bioRxiv*, 444281.

Boly, M., Balteau, E., Schnakers, C., Degueldre, C., Moonen, G., Luxen, A., ... & Laureys, S.

(2007). Baseline brain activity fluctuations predict somatosensory perception in

humans. *Proceedings of the National Academy of Sciences*, 104(29), 12187-12192.

Boly, Mélanie, Christophe Phillips, Luaba Tshibanda, Audrey Vanhaudenhuyse, M. Schabus,

Thien Thanh Dang-Vu, Gustave Moonen, Roland Hustinx, Pierre Maquet, and Steven

Laureys. "Intrinsic brain activity in altered states of consciousness: how conscious is the

734 default mode of brain function?." *Annals of the New York Academy of Sciences* 1129 (2008):

735 119.

736 Boly, M., Tshibanda, L., Vanhaudenhuyse, A., Noirhomme, Q., Schnakers, C., Ledoux, D., ...

737 & Luxen, A. (2009). Functional connectivity in the default network during resting state is

738 preserved in a vegetative but not in a brain dead patient. *Human brain mapping*, 30(8), 2393-

739 2400.

740 Bonhomme, V. L. G., Boveroux, P., Brichant, J. F., Laureys, S., & Boly, M. (2012). Neural

741 correlates of consciousness during general anesthesia using functional magnetic resonance

742 imaging (fMRI). *Archives italiennes de biologie*, 150(2/3), 155-163.

743 Boveroux, P., Vanhaudenhuyse, A., Bruno, M. A., Noirhomme, Q., Lauwick, S., Luxen, A.,

744 ... & Brichant, J. F. (2010). Breakdown of within-and between-network resting state

745 functional magnetic resonance imaging connectivity during propofol-induced loss of

746 consciousness. *Anesthesiology: The Journal of the American Society of*

747 *Anesthesiologists*, *113*(5), 1038-1053.

748 Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network:

749 anatomy, function, and relevance to disease.

750 Buckner, R. L., Krienen, F. M., & Yeo, B. T. (2013). Opportunities and limitations of

751 intrinsic functional connectivity MRI. *Nature neuroscience*, 16(7), 832.

752 Carhart-Harris, R. L., & Friston, K. J. (2010). The default-mode, ego-functions and free-

753 energy: a neurobiological account of Freudian ideas. *Brain*, *133*(4), 1265-1283.

754 Centeno, M., Tierney, T. M., Perani, S., Shamshiri, E. A., StPier, K., Wilkinson, C., ... &

755 Pressler, R. M. (2016). Optimising EEG-fMRI for localisation of focal epilepsy in

756 children. *PloS one*, 11(2).

Chennu, S., O'Connor, S., Adapa, R., Menon, D. K., & Bekinschtein, T. A. (2016). Brain

connectivity dissociates responsiveness from drug exposure during propofol-induced

transitions of consciousness. *PLoS computational biology*, *12*(1), e1004669.

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven

attention in the brain. *Nature reviews neuroscience*, *3*(3), 201-215.

Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., &

Peelle, J. E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and parallel

processing using Matlab and XML. *Frontiers in neuroinformatics*, *8*, 90.

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I.

Segmentation and surface reconstruction. *Neuroimage*, *9*(2), 179-194.

D'Argembeau, A., Collette, F., Van der Linden, M., Laureys, S., Del Fiore, G., Degueldre, C.,

... & Salmon, E. (2005). Self-referential reflective activity and its relationship with rest: a

PET study. *Neuroimage*, *25*(2), 616-624.

Davis, M. H. et al. Dissociating speech perception and comprehension at reduced levels of

awareness. *Proc. Natl. Acad. Sci.* USA 104, 16032–16037 (2007).

Demertzi, A., Soddu, A., & Laureys, S. (2013). Consciousness supporting networks. *Current*

Opinion in Neurobiology, *23*(2), 239-244.

Demertzi, A., Tagliazucchi, E., Dehaene, S., Deco, G., Barttfeld, P., Raimondo, F., ... &

Schiff, N. D. (2019). Human consciousness is supported by dynamic complex patterns of

brain signal coordination. *Science advances*, 5(2), eaat7603.

Dosenbach, N. U., Fair, D. A., Miezin, F. M., Cohen, A. L., Wenger, K. K., Dosenbach, R.

A., ... & Schlaggar, B. L. (2007). Distinct brain networks for adaptive and stable task control

in humans. *Proceedings of the National Academy of Sciences*, *104*(26), 11073-11078.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs

for intelligent behaviour. *Trends in cognitive sciences*, *14*(4), 172-179.

Elliott, R. (2003). Executive functions and their disorders: Imaging in clinical

neuroscience. *British medical bulletin*, 65(1), 49-59.

Ernst, M., & Paulus, M. P. (2005). Neurobiology of decision making: a selective review from

a neurocognitive and clinical perspective. *Biological psychiatry*, *58*(8), 597-604.

Fiset, P., Paus, T., Daloze, T., Plourde, G., Meuret, P., Bonhomme, V., ... & Evans, A. C.

(1999). Brain mechanisms of propofol-induced loss of consciousness in humans: a positron

emission tomographic study. Journal of Neuroscience, 19(13), 5506-5513.

Fisher, R. A. (1915). Frequency distribution of the values of the correlation coefficient in

samples from an indefinitely large population. Biometrika, 10(4), 507-521.

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state": a practical

method for grading the cognitive state of patients for the clinician. *Journal of psychiatric*

research, *12*(3), 189-198.

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E.

(2005). The human brain is intrinsically organized into dynamic, anticorrelated functional

networks. *Proceedings of the National Academy of Sciences*, 102(27), 9673-9678.

Fransson, P. (2005). Spontaneous low-frequency BOLD signal fluctuations: An fMRI

investigation of the resting-state default mode of brain function hypothesis. *Human brain*

mapping, 26(1), 15-29.

Friston K.J., Fletcher P., Josephs O., Holmes A., Rugg M.D., Turner R. (1998). Event-

Related fMRI: Characterizing Differential Responses. NeuroImage, Vol 7, 1,30-40.

Greicius, M. D., Krasnow, B., Reiss, A. L., & Menon, V. (2003). Functional connectivity in

the resting brain: a network analysis of the default mode hypothesis. *Proceedings of the*

National Academy of Sciences, *100*(1), 253-258.

Gusnard, D. A., Akbudak, E., Shulman, G. L., & Raichle, M. E. (2001). Medial prefrontal

cortex and self-referential mental activity: relation to a default mode of brain

function. *Proceedings of the National Academy of Sciences*, *98*(7), 4259-4264.

Hadi, A.S., (1992), 'Identifying multiple outliers in multivariate data', Journal of the Royal

Statistical Society. Series B (Methodological), Vol. 54, No. 3, pp. 761-771.

Hagler Jr, D. J., Saygin, A. P., & Sereno, M. I. (2006). Smoothing and cluster thresholding

for cortical surface-based group analysis of fMRI data. Neuroimage, 33(4), 1093-1103.

Hahn, B., Ross, T. J., & Stein, E. A. (2007). Cingulate activation increases dynamically with

response speed under stimulus unpredictability. *Cerebral cortex*, 17(7), 1664-1671.

Haugg, A., Cusack, R., Gonzalez-Lara, L. E., Sorger, B., Owen, A. M., & Naci, L. (2018).

Do patients thought to lack consciousness retain the capacity for internal as well as external

awareness?. *Frontiers in neurology*, *9*, 492.

Hauk, O., Johnsrude, I., & Pulvermüller, F. (2004). Somatotopic representation of action

words in human motor and premotor cortex. *Neuron*, *41*(2), 301-307.

Honey, C. J., Newman, E. L., & Schapiro, A. C. (2017). Switching between internal and

external modes: a multiscale learning principle. *Network Neuroscience*, 1(4), 339-356.

Huang, Z., Zhang, J., Wu, J., Mashour, G. A., & Hudetz, A. G. (2020). Temporal circuit of

macroscale dynamic brain activity supports human consciousness. *Science advances*, *6*(11),

eaaz0087.

Kelly, A. C., Uddin, L. Q., Biswal, B. B., Castellanos, F. X., & Milham, M. P. (2008).

Competition between functional brain networks mediates behavioral

variability. *Neuroimage*, 39(1), 527-537.

Kroger, J. K., Sabb, F. W., Fales, C. L., Bookheimer, S. Y., Cohen, M. S., & Holyoak, K. J.

(2002). Recruitment of anterior dorsolateral prefrontal cortex in human reasoning: a

parametric study of relational complexity. *Cerebral cortex*, *12*(5), 477-485.

Kucyi, A., Daitch, A., Raccah, O., Zhao, B., Zhang, C., Esterman, M., ... & Parvizi, J. (2020).

Electrophysiological dynamics of antagonistic brain networks reflect attentional

fluctuations. *Nature Communications*, 11(1), 1-14.

Långsjö, J. W., Alkire, M. T., Kaskinoro, K., Hayama, H., Maksimow, A., Kaisti, K. K., ... &

Scheinin, H. (2012). Returning from oblivion: imaging the neural core of consciousness.

Journal of Neuroscience, 32(14), 4935-4943.

Lau, H. C., Rogers, R. D., Haggard, P., & Passingham, R. E. (2004). Attention to

intention. *science*, *303*(5661), 1208-1210.

Lee, K. M., Chang, K. H., & Roh, J. K. (1999). Subregions within the supplementary motor

area activated at different stages of movement preparation and execution. *Neuroimage*, *9*(1),

117-123.

Leslie, K., Skrzypek, H., Paech, M. J., Kurowski, I., & Whybrow, T. (2007). Dreaming

during Anesthesia and Anesthetic Depth in Elective Surgery PatientsA Prospective Cohort

Study. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 106(1), 33-

42.

Luppi, A., Craig, M., Pappas, I., Finoia, P., Williams, G., Allanson, J., ... & Stamatakis, E.

(2019). Consciousness-specific dynamic interactions of brain integration and functional

diversity.

MacDonald, A. A., Naci, L., MacDonald, P. A., & Owen, A. M. (2015). Anaesthesia and

neuroimaging: investigating the neural correlates of unconsciousness. *Trends in Cognitive*

Sciences, *19*(2), 100-107.

Marek, S., & Dosenbach, N. U. (2018). The frontoparietal network: function,

electrophysiology, and importance of individual precision mapping. *Dialogues in clinical*

neuroscience, *20*(2), 133.

Marsh, B. M. W. M. N., White, M., Morton, N., & Kenny, G. N. C. (1991). Pharmacokinetic

model driven infusion of propofol in children. *BJA: British Journal of Anaesthesia*, *67*(1),

41-48.

Mashour, G. A., & Avidan, M. S. (2015). Intraoperative awareness: controversies and non-

controversies. *British journal of anaesthesia*, *115*(suppl_1), i20-i26.

Mashour, G. A., & Hudetz, A. G. (2018). Neural correlates of unconsciousness in large-scale

brain networks. *Trends in neurosciences*, 41(3), 150-160.

Menon, V. (2015). Salience network. In A. W. Toga (Ed.), Brain mapping: An encyclopedic

reference (Vol. 2, pp. 597–611). London: Academic Press: Elsevier.

Michalka, S. W., Kong, L., Rosen, M. L., Shinn-Cunningham, B. G., & Somers, D. C.

(2015). Short-term memory for space and time flexibly recruit complementary sensory-biased

frontal lobe attention networks. Neuron, 87(4), 882-892.

Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The

impact of global signal regression on resting state correlations: are anti-correlated networks

introduced?. *Neuroimage*, *44*(3), 893-905.

Murphy, K., Birn, R. M., & Bandettini, P. A. (2013). Resting-state fMRI confounds and

cleanup. *Neuroimage*, 80, 349-359.

Naci, L., Cusack, R., Jia, V. Z., & Owen, A. M. (2013). The brain's silent messenger: using

selective attention to decode human thought for brain-based communication. *Journal of*

Neuroscience, *33*(22), 9385-9393.

Naci, L., Cusack, R., Anello, M., & Owen, A. M. (2014). A common neural code for similar

conscious experiences in different individuals. *Proceedings of the National Academy of*

Sciences, 111(39), 14277-14282.

Naci, L., Sinai, L., & Owen, A. M. (2017). Detecting and interpreting conscious experiences

in behaviorally non-responsive patients. *NeuroImage*, *145*, 304-313.

Naci, L., Haugg, A., MacDonald, A., Anello, M., Houldin, E., Naqshbandi, S., ... & Owen, A.

M. (2018). Functional diversity of brain networks supports consciousness and verbal

intelligence. *Scientific reports*, *8*(1), 13259.

Nachev, P., Rees, G., Parton, A., Kennard, C., & Husain, M. (2005). Volition and conflict in

human medial frontal cortex. *Current Biology*, 15(2), 122-128.

Pal, D., Li, D., Dean, J. G., Brito, M. A., Liu, T., Fryzel, A. M., ... & Mashour, G. A. (2020).

Level of consciousness is dissociable from electroencephalographic measures of cortical

connectivity, slow oscillations, and complexity. *Journal of Neuroscience*, *40*(3), 605-618.

Palanca, B. J. A., Mashour, G. A., & Avidan, M. S. (2009). Processed electroencephalogram

in depth of anesthesia monitoring. *Current Opinion in Anesthesiology*, *22*(5), 553-559.

Pandit, J. J., Andrade, J., Bogod, D. G., Hitchman, J. M., Jonker, W. R., Lucas, N., ... & Paul,

R. G. (2014). 5th National Audit Project (NAP5) on accidental awareness during general

anaesthesia: summary of main findings and risk factors. *British journal of*

anaesthesia, *113*(4), 549-559.

Pandit JJ, Cook TM; The NAP5 Steering Panel. NAP5. Accidental Awareness During

General Anesthesia in the United Kingdom and Ireland: Report and Findings. *London: The*

Royal College of Anesthetists and Association of Anesthetists of Great Britain and Ireland;

2014. Available at: http://www.nationalauditprojects.org.uk/NAP5_home. Accessed on May

3, 2017.

Peigneux, P., Orban, P., Balteau, E., Degueldre, C., Luxen, A., Laureys, S., & Maquet, P.

(2006). Offline persistence of memory-related cerebral activity during active

wakefulness. *PLoS biology*, *4*(4).

Plourde, G., Belin, P., Chartrand, D., Fiset, P., Backman, S. B., Xie, G., & Zatorre, R. J.

(2006). Cortical processing of complex auditory stimuli during alterations of consciousness

with the general anesthetic propofol. *Anesthesiology: The Journal of the American Society of*

Anesthesiologists, 104(3), 448-457.

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman,

G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of*

Sciences, 98(2), 676-682.

Raichle, M. E. (2011). The restless brain. *Brain connectivity*, *1*(1), 3-12.

Ramsay, M. A. E., Savege, T. M., Simpson, B. R. J., & Goodwin, R. (1974). Controlled

sedation with alphaxalone-alphadolone. *Br med J*, *2*(5920), 656-659.

Sanders, R. D., Gaskell, A., Raz, A., Winders, J., Stevanovic, A., Rossaint, R., ... & Meier, S.

(2017). Incidence of Connected Consciousness after Tracheal IntubationA Prospective,

International, Multicenter Cohort Study of the Isolated Forearm Technique. *Anesthesiology:*

The Journal of the American Society of Anesthesiologists, *126*(2), 214-222.

Sandin, R. H., Enlund, G., Samuelsson, P., & Lennmarken, C. (2000). Awareness during

anaesthesia: a prospective case study. *The Lancet*, *355*(9205), 707-711.

Sarasso, S., Boly, M., Napolitani, M., Gosseries, O., Charland-Verville, V., Casarotto, S., ...

& Rex, S. (2015). Consciousness and complexity during unresponsiveness induced by

propofol, xenon, and ketamine. *Current Biology*, 25(23), 3099-3105.

Seeley, W. W. (2019). The salience network: a neural system for perceiving and responding

to homeostatic demands. Journal of Neuroscience, 39(50), 9878-9882.

Schneider, F., Bermpohl, F., Heinzel, A., Rotte, M., Walter, M., Tempelmann, C., ... &

Northoff, G. (2008). The resting brain and our self: self-relatedness modulates resting state

neural activity in cortical midline structures. *Neuroscience*, 157(1), 120-131.

Searle, R., & Hopkins, P. M. (2009). Pharmacogenomic variability and anaesthesia. *British*

journal of anaesthesia, *103*(1), 14-25.

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., ... &

Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing

and executive control. *Journal of Neuroscience*, 27(9), 2349-2356.

Shallice, T. (1988). *From neuropsychology to mental structure*. Cambridge University Press.

Shulman, G. L., Fiez, J. A., Corbetta, M., Buckner, R. L., Miezin, F. M., Raichle, M. E., &

Petersen, S. E. (1997). Common blood flow changes across visual tasks: II. Decreases in

cerebral cortex. *Journal of cognitive neuroscience*, 9(5), 648-663.

Smith, V., Mitchell, D. J., & Duncan, J. (2018). Role of the default mode network in

cognitive transitions. *Cerebral Cortex*, 28(10), 3685-3696.

Sonkusare, S., Breakspear, M., & Guo, C. (2019). Naturalistic Stimuli in Neuroscience:

Critically Acclaimed. *Trends in cognitive sciences*.

Spreng, R. N., DuPre, E., Selarka, D., Garcia, J., Gojkovic, S., Mildner, J., ... & Turner, G. R.

(2014). Goal-congruent default network activity facilitates cognitive control. *Journal of*

Neuroscience, 34(42), 14108-14114.

Sridharan, D., Levitin, D. J., & Menon, V. (2008). A critical role for the right fronto-insular

cortex in switching between central-executive and default-mode networks. *Proceedings of the*

National Academy of Sciences, 105(34), 12569-12574.

Stamatakis, E. A., Adapa, R. M., Absalom, A. R., & Menon, D. K. (2010). Changes in resting

neural connectivity during propofol sedation. *PloS one*, 5(12).

Tagliazucchi, E., von Wegner, F., Morzelewski, A., Brodbeck, V., Jahnke, K., & Laufs, H.

(2013). Breakdown of long-range temporal dependence in default mode and attention

networks during deep sleep. *Proceedings of the National Academy of Sciences*, 110(38),

15419-15424.

Tobyne, S. M., Somers, D. C., Brissenden, J. A., Michalka, S. W., Noyce, A. L., & Osher, D.

E. (2018). Prediction of individualized task activation in sensory modality-selective frontal

cortex with 'connectome fingerprinting'. Neuroimage, 183, 173-185.

Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C., & Castellanos, F. X. (2015). Inscapes: A

movie paradigm to improve compliance in functional magnetic resonance

imaging. *Neuroimage*, 122, 222-232.

Vanhaudenhuyse, A., Demertzi, A., Schabus, M., Noirhomme, Q., Bredart, S., Boly, M., ... &

Laureys, S. (2011). Two distinct neuronal networks mediate the awareness of environment

and of self. *Journal of cognitive neuroscience*, 23(3), 570-578.

Varley, T. F., Luppi, A. I., Pappas, I., Naci, L., Adapa, R., Owen, A. M., ... & Stamatakis, E.

A. (2020). consciousness & Brain functional complexity in propofol Anaesthesia. *Scientific*

reports, *10*(1), 1-13.

Vatansever, D., Manktelow, A. E., Sahakian, B. J., Menon, D. K., & Stamatakis, E. A.

(2017). Angular default mode network connectivity across working memory load. *Human*

brain mapping, *38*(1), 41-52.

Vatansever, D., Schröter, M., Adapa, R. M., Bullmore, E. T., Menon, D. K., & Stamatakis, E.

A. (2020). Reorganisation of Brain Hubs across Altered States of consciousness. *Scientific*

Reports, 10(1), 1-11.

Wang, J., Ren, Y., Hu, X., Nguyen, V. T., Guo, L., Han, J., & Guo, C. C. (2017). Test–retest

reliability of functional connectivity networks during naturalistic fMRI paradigms. *Human*

brain mapping, 38(4), 2226-2241.

Wicker, B., Ruby, P., Royet, J. P., & Fonlupt, P. (2003). A relation between rest and the self

in the brain?. *Brain Research Reviews*, *43*(2), 224-230.

Winkler, A. M., Greve, D. N., Bjuland, K. J., Nichols, T. E., Sabuncu, M. R., Håberg, A. K.,

... & Rimol, L. M. (2018). Joint analysis of cortical area and thickness as a replacement for

the analysis of the volume of the cerebral cortex. Cerebral cortex, 28(2), 738-749.

Witon, A., Shirazibehehsti, A., Aviles, A., Adapa, R., Menon, D., Chennu, S., ... & Bowman,

H. (2020). Sedation modulates fronto-temporal predictive coding circuits and the double

surprise acceleration effect. *bioRxiv*.