1 **Title**
2

The brain-bases of responsiveness variability under moderate anaesthesia

Authors

Feng Deng[a], Nicola Taylor[a], Adrian M. Owen[b,d], Rhodri Cusack[a], Lorina Naci[a,c*]

[a] Trinity College Institute of Neuroscience, School of Psychology, Trinity College Dublin,

Dublin, Ireland.

[b] Brain and Mind Institute, Western University, London, Canada.

[c] Global Brain Health Institute, Trinity College Dublin, Dublin, Ireland

[d] Department of Physiology and Pharmacology and Department of Psychology, Western

University, London, Canada.

[*] **Corresponding author:**

Lorina Naci

School of Psychology

Trinity College Institute of Neuroscience

Global Brain Health Institute

Trinity College Dublin

Dublin, Ireland

Telephone: +353 (0)87 688 5642

Email: nacil@tcd.ie

Abstract

Anaesthesia combined with functional neuroimaging provides a powerful approach for understanding the brain mechanisms that change as consciousness fades. Although propofol is used ubiquitously in clinical interventions that reversibly suppress consciousness, its effect varies substantially between individuals, and the brain bases of this variability remain poorly understood. We asked whether three networks that are primary sites of propofol-induced sedation and key to conscious cognition — the dorsal attention (DAN), executive control (ECN), and default mode (DMN) network — underlie responsiveness variability under anaesthesia. Healthy participants (N=17) underwent propofol sedation inside the fMRI scanner at dosages of 'moderate' anaesthesia, and behavioural responsiveness was measured with a target detection task. To assess information processing, participants were scanned during an active engagement condition comprised of a suspenseful auditory narrative, in addition to the resting state. A behavioural investigation in a second group of non-anesthetized participants (N=25) qualified the attention demands of narrative understanding, which we then related to the brain activity of participants who underwent sedation. 30% of participants showed no delay in reaction times relative to wakefulness, whereas the others, showed significantly delayed and fragmented responses, or full omission of responses. These responsiveness differences did not relate to information processing differences. Rather, only the functional connectivity within the ECN during wakefulness differentiated the participants' responsiveness level, with significantly stronger connectivity in the fast relative to slow responders. Consistent with this finding, fast responders had significantly higher grey matter volume in the frontal cortex aspect of the ECN. For the first time, these results show that responsiveness variability during propofol anaesthesia relates to inherent differences in brain function and structure within the executive control network, which can be predicted prior to sedation. These results shed light on the brain bases of responsiveness differences and highlight novel markers that may help to improve the accuracy of awareness monitoring during clinical anaesthesia.

Introduction

Understanding how the human brain gives rise to consciousness remains a grand challenge for modern neuroscience. A first step to understanding consciousness is to define what it is not. To this end, anaesthesia combined with functional neuroimaging provides a powerful approach for studying the brain mechanisms that change as consciousness fades (Vatansever et al, 2020; Pal et al., 2020; Varley et al., 2020; Luppi et al., 2019; Demertzi et al., 2019; Mashour and Hudetz 2018; Naci et al., 2018; MacDonald et al., 2015; Sarasso et al., 2015; Stamatakis et al., 2010), and, conversely, that are necessary for realizing human consciousness. Anaesthesia has been used for over 150 years to reversibly abolish consciousness in clinical medicine, but its effect can vary substantially between individuals. At moderate dosages, the intended suppression of behavioural responsiveness is highly variable (Chennu et al., 2016; Bola et al., 2019), and at deep anaesthesia dosages, in rare cases (0.1-0.2%; Mashour and Avidan 2015; Sandin et al., 2000), individuals retain conscious awareness, also known as 'unintended intraoperative awareness' (Pandit et al., 2017; Sanders et al., 2017; Sandin et a., 2000; Mashour and Avidan 2015). A much higher percentage of patients presumed to be unconscious during general anaesthesia (22%; Leslie et al., 2007) may have subjective experiences, such as dreaming. The brain-bases of this considerable inter-individual effect variability remain poorly understood.

A key and mostly overlooked question is what these individual differences (Searle and Hopkins 2009; Palanca et al., 2009) can reveal about the unravelling of conscious cognition as consciousness falters during anaesthesia. Propofol is the most common anaesthetic agent in clinical interventions that require the reversible suppression of consciousness. Two recent electroencephalography (EEG) studies that measured variable behavioural responsiveness during mild and moderate propofol anaesthesia reported that the participants' varying levels of responsiveness were differentiated by alpha band connectivity during wakefulness (Chennu et al., 2016), and changing patterns of EEG signal diversity from wakefulness to moderate sedation (Bola et al., 2019). However, the brain regional- or network-bases for this substantial individual variability (Searle and Hopkins 2009; Palanca et al., 2009), and its potential link to cognitive function, remain unknown. Propofol-induced sedation produces selective metabolic impairment and reduces brain activity bilaterally in frontal and parietal associative regions (Witon et al., 2020; Boly et al., 2008; Plourde et al., 2006; Alkire 2005; Baars et al., 2003; Fiset et al., 1999; for a review see MacDonald et al., 2015; but see Pal et

al., 2020). Therefore, three brain networks with frontal and parietal lobe distribution, the

dorsal attention network (DAN), executive control network (ECN), and default mode

network (DMN) are primary candidate sources of individual variability under propofol

sedation.

The DAN and ECN, distributed laterally across frontal and parietal lobes, are key to

orchestrating stimulus-driven and goal-directed cognition. Their roles include, orienting and

modulating attention to the saliency of incoming stimuli, monitoring and analysing

information from the environment and integrating it with internally generated goals, as well

as planning and adapting new behavioural schemas to take account of this information

(Duncan 2010; Elliott 2003; Corbetta and Schulman 2002; Kroger et al., 2002; Shallice

1988). The DMN extends partially in lateral and medial frontal and parietal lobes and is

thought to mediate internally-oriented cognition, e.g., autobiographical memory, imagination,

and thinking about the self (Andrews-Hanna et al. 2010; Schneider et al., 2008; Buckner et

al., 2008; Beer 2007; D'Argembeau et al., 2005; Wicker et al., 2003; Gusnard et al., 2001).

Recent studies show that it is also involved in external environment monitoring (Spreng et al.

2014; Buckner et al. 2008; Hahn et al., 2007), including encoding of scenes, episodes or

contexts (Vatansever et al., 2017), or shifts in contextually relevant information (Smith et al.,

2018). Recent studies that investigate state-related manipulations (conscious vs unconscious)

show that by, contrast to other brain networks, the DAN, ECN and DMN are selectively

impaired during loss of consciousness, not only under anaesthesia but also during severe

brain injury (Luppi et al., 2019; Naci et al., 2018). Thus, they provide strong evidence that

these three networks are part of a common brain mechanism disrupted during loss of

consciousness across vastly different conditions.

It is well established that the DAN/ECN and the DMN display an antagonistic relationship

(Huang et al., 2020;Fox et al., 2005; Fransson et al., 2005) during spontaneous thought and

goal-directed cognition. During conditions that engage external-attention, the reduction of

functional activity in the DMN (Greicius et al., 2003, Raichle et al., 2001, Shulman et al.

1997), is concomitant with an increase of activity in the DAN and ECN (Seeley et al. 2007;

Dosenbach et al. 2007; Fox et al., 2005; Sridharan et al. 2008). The anticorrelation between

these is related to individual differences in performance variability (Kelly 2008), and directly

supports sustained attention (Kucyi et al., 2020). More broadly, it has been suggested to

facilitate ongoing switches between internally- and externally biased modes of attention

(Honey et al., 2017; Buckner et al., 2013) that contribute to consciousness (Carhart-Harris

and Friston 2010). Conversely, this antagonistic relationship breaks down in state-related

manipulations of consciousness, such as during sleep (Tagliazucchi et al., 2013), anaesthesia

(Bonhomme at el., 2012; Boveroux et al., 2010), and severe brain injury (Haugg et al., 2018;

Boly et al., 2009). Despite this accumulating evidence for the key and interrelated roles of the

DAN, ECN, and DMN in supporting conscious cognition (Huang et al., 2020; Demertzi et

al., 2013; Vanhaudenhuyse et al., 2011), and their primacy as target sites of propofol-induced

sedation, their roles in individual differences under anaesthesia have not been previously

investigated.

To address this gap, in two studies we tested whether variability or impairments in functional

connectivity within and between the DAN, ECN, and DMN underlie individual differences in

responsiveness during propofol anaesthesia. To directly investigate their individual and joint

roles, in a first fMRI study we scanned healthy volunteers (N=17) during wakefulness and

during the administration of propofol at dosages of 'moderate anaesthesia,' expressly aimed

at engendering individual differences. Prior to the wakefulness and moderate anaesthesia

scans, participants were asked to perform a verbal recall memory test that assessed basic

memory function. Variability in behavioural responsiveness in each state was assessed with

an auditory target detection task prior to scanning. To test whether any individual differences

in responsiveness were related to differences in information processing that were invisible to

the clinical sedation scale (Ramsay et al., 1974), participants were scanned during an active

information processing condition comprised of a brief (5 minutes) engaging auditory

narrative, and during a control condition of spontaneous thought, i.e., the resting state.

Listening to plot-driven narratives is naturally engaging, requires minimal behavioural

collaboration from volunteers, and therefore, is highly suitable for testing information

processing independently of behavioural output or eye opening (Naci et al., 2018; 2017),

which are impaired in moderate anaesthesia. In a second behavioural study, we assessed the

high-level attention demands of narrative understanding in an independent behavioural group

(N=25), and related them to the brain activity of participants who underwent scanning.

Finally, to test whether individual differences under anaesthesia related to inherent brain

features independent from the propofol sedation, we investigated the functional connectivity

between and within the three networks during wakefulness, as well as grey matter volume

differences across participants.

Methods

Participants

Study 1. Healthy participants for the fMRI scanning anaesthesia study (N=17; 18-40 years; 13
males) were tested at the Robarts Research Institute, Western University, in Canada. Ethical
approval was obtained from the Health Sciences Research Ethics Board and Psychology
Research Ethics Board of Western University. Data from the wakefulness condition was
previously reported in Naci et al., 2018.

Study 2. The independent group of healthy participants (N=25; 18-40 years; 7 males) for the
behavioural suspense rating study were tested at the Global Brain health Institute at Trinity
College Dublin, in Ireland. Ethical approval was obtained from the School of Psychology
Research Ethics Board, Trinity College Dublin.

All healthy participants were right-handed, native English speakers, and had no history of
neurological disorders. Both studies were performed in accordance with the relevant
guidelines and regulations set out by the research ethics boards. Informed consent was
obtained for each participant prior to the experiment.

Stimuli and Design

Study 1

Behavioural testing. *Memory recall task*. The brief recall from the Mini Mental State Exam
(Folstein et al., 1975) was adopted to test memory function under sedation. The researcher
named three unrelated objects clearly and slowly and asked the volunteer to name each of
them. The volunteer was instructed to remember the words in order to be able to repeat them
in a short while. After 10 minutes, the volunteer was asked the repeat the words. Two
different lists (a. Ball- Flag-Tree; b. Flower-Egg-Rope) were used to avoid familiarity effects
between the wakeful and moderate anaesthesia states. Their presentation was
counterbalanced across participants.

Auditory target detection task. Before the wakeful and moderate anaesthesia fMRI sessions,

participants were asked to perform a computerized auditory target detection task (4 minutes),

which aimed to assess individual differences during moderate anaesthesia. Participants were

instructed via headphones to press a button with their index finger as soon as they heard an

auditory beep and to keep their eyes on the fixation cross on the screen. The response box

was tapped to the hand to ensure the participant did not lose contact during anaesthesia. The

averaged reaction time (RT) for each participant was computed for further analyses.

MRI testing. Inside the MRI scanner, participants underwent two functional scans during

wakefulness and moderate anaesthesia. A plot-driven audio narrative (5 minutes) was

presented over MRI compatible noise cancelation headphones (Sensimetrics, S14;

www.sens.com), and participants were asked to simply listen with eyes closed. The story

comprised a highly engaging auditory excerpt from the movie 'Taken' depicting dramatic

events, where a young girl travelling abroad without her family is kidnapped while speaking

on the phone to her father. A similar eyes-closed, resting state condition (8 minutes) was also

acquired in each state for comparison with the narrative condition.

Sedation Procedure. The level of sedation was measured with the Ramsay clinical sedation

scale for each participant by 3 independent assessors (two anaesthesiologists and one

anaesthesia nurse) in person inside the scanner room. Before entering the fMRI scanner, a

20G I.V. cannula was inserted into a vein on the dorsum of the non-dominant hand of the

participants. The propofol infusion system was connected to the cannula prior to the first

scanning session. No propofol was administered during the wakeful session. Participants

were fully wakeful, alert and communicated appropriately (Ramsay 1) and wakefulness (eye-

opening) was monitored with an infrared camera placed inside the scanner. At the

commencement of the moderate anaesthesia session, intravenous propofol was administered

with a Baxter AS 50 (Singapore). An effect-site/plasma steering algorithm was used

combined with the computer-controlled infusion pump to achieve step-wise increments in the

sedative effect of Propofol. This infusion pump was manually adjusted to achieve the desired

levels of sedation, guided by targeted concentrations of Propofol, as predicted by the TIVA

Trainer (the European Society for Intravenous Anaesthesia, eurosiva.eu) pharmacokinetic

simulation program. The pharmacokinetic model provided target-controlled infusion by

adjusting infusion rates of Propofol over time to achieve and maintain the target blood

concentrations as specified by the Marsh 3 compartment algorithm for each participant, as

incorporated in the TIVA Trainer software (Marsh et al., 1991). In accordance with the

Canadian Anaesthesia Society (CAS) guidelines, non-invasive blood pressure (NIBP), heart

rate, oxygen saturation (SpO2) and end-tidal carbon dioxide (ETCO2) were monitored

continuously through the use of a dedicated MR compatible anaesthesia monitor. Complete

resuscitation equipment was present throughout the testing.

Propofol infusion commenced with a target effect-site concentration of 0.6 μg/ml and oxygen

was titrated to maintain SpO2 above 96%. Throughout sedation, participants remained

capable of spontaneous cardiovascular function and ventilation. Supplemental oxygen was

administered via nasal cannulae to ensure adequate levels of oxygen at all times. If the

Ramsay level was lower than 3 for moderate sedation, the concentration was slowly

increased by increments of 0.3 μg/ml with repeated assessments of responsiveness between

increments to obtain a Ramsay score of 3. During administration of propofol, participants

generally became calm and slowed in their response to verbal communication. Once

participants stopped engaging in spontaneous conversation, and speech became sluggish, they

were classified as being at Ramsey level 3. Nevertheless, when asked via loud verbal

communication participants agreed to perform the auditory target detection and memory

recall tasks, as would be expected at Ramsey level 3. During moderate anaesthesia (Ramsey

3), the mean estimated effect-site propofol concentration was 1.99 (1.59- 2.39) μg/ml and the

mean estimated plasma propofol concentration was 2.02 (1.56- 2.48) μg/ml.

Study 2

To provide a subjective measure of attention during the story, participants rated how

"suspenseful" it was every 2 seconds, from 'least' (1) to 'most suspenseful' (9). Participants

heard the stimulus through over-ear headphones in sound-isolated rooms at the Global Brain

Health Institute at Trinity College Dublin. Laboratory computers were used to present the

stimulus, cue and collect responses. The audio excerpt was divided into 156 clips, each 2

seconds long to match the repetition time (TR, 2 sec) used in independent participant group

who underwent MRI scanning. Participants had up to 3 seconds to make a response, at which

point the next sequential clip began immediately. At the end of experiment, participants

indicated via a feedback questionnaire that the interruptions did not disrupt the coherence of

the story's plot and their perception of suspense throughout.

Data Analyses

Study 1

MRI Data Acquisition. Functional images were obtained on a 3T Siemens Prisma system,
with a 32-channel head coil. The high-resolution brain structural images were acquired using
a T1-weighted 3D MPRAGE sequence with following parameters, voxel size: 1 x 1 x 1 mm,
TA=5 minutes and 38 seconds, echo time (TE) = 4.25ms, matrix size=240x256x192, flip
angle (FA) = 9 degrees. The functional echo-planar images (EPI) images were obtained with
following parameters, 33 slices, voxel size: 3 x 3 x 3, inter-slice gap of 25%, repetition time
(TR) = 2000ms, TE=30ms, matrix size=64x64, FA=75 degrees. The audio story and resting
state had 155 and 256 scans, respectively. A volume level deemed comfortable by each
individual was used for the duration of the scanning.

MRI preprocessing: 1) Functional data. Standard preprocessing procedures and data analyses
were performed with SPM8 (Wellcome Institute of Cognitive Neurology,
http://www.fil.ion.ucl.ac.uk/spm/software/spm8/) and the AA pipeline software (Cusack et
al., 2015). In the preprocessing pipeline, we performed slice timing correction, motion
correction, registration to structural images, normalization to a template brain, and
smoothing. The data were smoothed with a Gaussian smoothing kernel of 10mm FWHM
(Peigneux et al. 2006). Spatial normalization was performed using SPM8's segment-and-
normalize procedure, whereby the T1 structural was segmented into grey and white matter
and normalized to a segmented MNI-152 template. These normalization parameters were
then applied to all EPIs. The time series in each voxel was high pass-filtered with a cutoff of
1/128 Hz to remove low-frequency noise, and scaled to a grand mean of 100 across voxels
and scans in each session. The preprocessed data were analyzed in using the general linear
model. Prior to analyses, the first five scans of each session were discarded to achieve T1
equilibrium and to allow participants to adjust to the noise of the scanner. To avoid the
formation of artificial anti-correlations (Murphy et al. 2009; Anderson et al., 2011), we
performed no global signal regression. 2) Structural data. The brain structural images were
processed using FreeSurfer package (https://surfer.nmr.mgh.harvard.edu/), a well-
documented automated program which is widely used to perform surface-based
morphometric analysis (Dale et al. 1999). The processing steps include: 1) removing the non-
brain tissue; 2) transforming the skull-stripping brain volume to Talairach-like space; 3)

segmenting brain tissues to GM, WM, cerebrospinal fluid (CSF); 4) performing intensity

normalization to remove the effect of bias field; 5) building a surface tessellation to generate

a triangular cortical mesh consisting of about 300,000 vertices in the whole brain surface; 6)

correcting topological deficits of cortical surface; and 7) deforming brain surface to generate

optimized models of GM/WM and GM/CSF boundaries.

Analyses of fMRI data. To investigate whether the behavioural variability under anaesthesia

related to perceptual or higher-order processing differences among participants, we

performed two mixed data-driven and model-based analyses. First, we extracted the sound

envelope of the auditory narrative via the Matlab *MIR*toolbox

(http://www.jyu.fi/hum/laitokset/musiikki/en/research/coe/materials/mirtoolbox) and built a

generalised linear model (GLM) by using statistical parametric mapping to derive auditory

characteristic-related brain activation for each individual. Subsequently, the z-scored average

suspense ratings of the narrative obtained by the independent group of participants in study 2

were used as a regressor in the fMRI data of individuals who underwent propofol anaesthesia

in study 1, to measure the neural correlates of information processing during the narrative

condition. The regressors were generated by convolving boxcar functions with the canonical

hemodynamic response function. Also included in the general linear model were nuisance

variables, comprising the movement parameters in the three directions of motion and three

degrees of rotation, as well as the mean of each session. Fixed-effect analyses were

performed in each subject, corrected for temporal auto-correlation using an AR (1)+white

noise model. Linear contrasts were used to obtain subject-specific estimates for each effect of

interest. Linear contrast coefficients for each participant were entered into the second level

random-effects analysis. Clusters or voxels that survived at $p<0.05$ threshold, corrected for

multiple comparisons with the family wise error (FWE) were considered statistically

significant.

Functional connectivity (FC) within and between the DAN, ECN, and DMN was assessed by

computing the Pearson correlation of the fMRI time courses between 19 regions of interests

(ROIs) (from Raichle 2011) constituting these different brain networks, as identified by

resting state studies (Table 1). Permutation (1000 times) tests were used to explore FC

difference between conditions and were FDR corrected for multiple comparisons. All of the

analyses were conducted using Fisher z-transformed correlation (Pearson *r*). Pearson

correlation cannot directly imply causal relations between neural regions, but it is a simple

FC measure with minimal assumptions. It is adequate for our purpose because the time-

course and spatial extent of fronto-parietal and default mode networks encompassed a vast

swath of hierarchical processing cascade and, thus, many regions of cause-effect space were

triggered by the stimulus. FC within and between these networks, indicated by Pearson

correlation, reflected their interactions over several minutes and the resulting computations

on the information content of the narrative condition, or of spontaneous thought during the

resting state condition. Glass's delta was used to compute the effect size of the comparison

between wakeful and moderate anaesthesia states, because of the different standard

deviations of the two states, while Hedges' g was used to compute the effect size of the

comparison between fast and slow responders because of the different sample sizes for the

two groups.

Grey matter volume analyses. Grey Matter Volume (GMV) was computed as the volume of

a truncated tetrahedron. The reconstructed surface was smoothed with a 10-mm full-width at

half maximum (FWHM) Gaussian kernel to perform the statistics. Based on group-level

comparison, we detected the clusters showing significant GWM differences at the group-

level. Monte Carlo simulation cluster analysis and a cluster-wise threshold of $p < 0.05$ was

adopted for multiple comparisons correction. In this way, we identified the regions with

significant group differences in GMV. We then extracted averaged GMV of each significant

cluster for individual participants and performed permutation tests (1000 times) to identify

the group difference between fast and slow responders.

Study 2

Analyses of suspense ratings. A group-averaged set of suspense ratings were calculated by

averaging participants' ratings at each time-point. To determine how similar suspense ratings

were across the group, the inter-subject correlation of suspense ratings was computed as the

average of the Pearson correlations of each participant's data with the mean data from the rest

of the group. To compute the average correlation value, each correlation coefficient was

normalized using Fisher's r-to-z transformation and the average of these values was

computed and then back-transformed into a correlation coefficient.

Results

The effect of moderate anaesthesia on brain network connectivity

Initially, we investigated how moderate propofol anaesthesia perturbed the patterns of functional connectivity (FC) relative to wakefulness across all the participants (Figure 1). For the narrative condition, a 2x2 ANOVA that explored the main effects of *state* (wakefulness, moderate anaesthesia) and *network connectivity* (DMN, DAN, ECN, DMN–DAN, DMN–ECN, DAN–ECN) showed significant main effects of state ($F(1, 16) = 5.3$, $p < 0.05$), network connectivity ($F(2.9, 46.7) = 28.1$, $p < 0.0001$), and a significant interaction effect of state by network connectivity ($F(5, 80) = 8.2$, $p < 0.0001$). We investigated further this interaction effect by testing the effect of sedation state on each network connectivity. We found significantly higher DMN–DAN (permutation test, $p < 0.01$, FDR corrected), as well as DMN–ECN (permutation test, $p < 0.05$, FDR corrected) connectivity in moderate anaesthesia relative to wakefulness, during the narrative condition (Figure 1B).

For the resting state condition, a 2x2 ANOVA that explored the main effects of *state* (wakefulness, moderate anaesthesia) and *network connectivity* (DMN, DAN, ECN, DMN–DAN, DMN–ECN, DAN–ECN) showed a significant main effect of network connectivity ($F(2, 32.3) = 18.9$, $p < 0.0001$) and a significant interaction effect of state by network connectivity ($F(2.4, 38.8) = 4.3$, $p < 0.05$). However, we found no significant differences between wakefulness and moderate anaesthesia for each network connectivity (Figure 1B).

A direct comparison between the narrative and resting state conditions, with a 2x2 ANOVA [*condition* (narrative, resting state) x *network connectivity* (6 levels)] on the FC difference values (moderate anaesthesia - wakefulness), showed a significant main effect of network connectivity ($F(5, 80) = 9.9$, $p < 0.0001$), and no main effect of condition or interaction effect. Although the 2x2 ANOVA did not reveal any interactions, effect size analyses showed higher effect sizes of moderate anaesthesia in the narrative relative to the resting state condition for both DMN–DAN (Glass' Delta = 1.20 vs 0.52) and DMN–ECN (Glass' Delta = 1.02 vs 0.46). In summary, these results suggested that the antagonistic relationship between the DMN and the DAN/ECN and was reduced during moderate anaesthesia, with a stronger and significant result in the narrative condition relative to the resting state.

Figure 1. about here please

The effect of moderate anaesthesia on behavioural responsiveness

Despite the same Ramsay 3 clinical score, independently determined by three assessors, we

observed significant heterogeneity in the reaction time of the auditory detection task (Figure

2). 5/17 participants were not delayed significantly relative to their non-sedated responses,

9/17 were significantly delayed and had fragmented responses (showing 2–40% missing

trials), and 3/17 failed to make any responses (full omission of responses), despite agreeing to

have understood the task instructions and to make responses via the button box affixed to

their dominant hand. This large individual variability was at odds with the propofol infusion

rates titrated for each participant based on the pharmacokinetic model adjusted for

demographic variables (Marsh et al., 1991), to maintain stable target blood concentrations

consistent with Ramsay 3 sedation level (Ramsay, 1974). Additionally, all participants

successfully completed the brief delayed recall memory task.

To further investigate these individual differences in responsiveness, individuals were

divided into two groups, fast responders (FRs; N=11), and slow responders (SRs; N=6) based

on the mean split of reaction time (Figure 2). Individual responsiveness variability was not

explained by demographic differences, as suggested by no differences in age (between-

samples t-test, $t = -0.42$, $p = 0.677$) and gender (Fisher's exact test, odds ratio = 0.47, $p =$

0.584) between FRs and SRs.

382   Figure 2. about here please

Rather, the brain-bases of this variability may be related to three other factors. First, the

individual variability could be related to underlying fine-grained perceptual and/or higher-

order information processing differences between participants, which may have been

invisible to the behavioural examination during the Ramsay assessment. For example,

participants who made no responses during moderate anaesthesia, may have appeared to

understand the task instructions based on their limited verbal responses, despite genuine lack

of understanding, which may have led to failure of button-press responses. Second,

independently of the extent of information processing in each individual, responsiveness

differences may be driven by connectivity alterations within and between the DAN, ECN and

DMN networks, which would suggest impairments in other cognitive processes. Third,

individual responsiveness differences may be related to structural brain differences, which

may, in turn, manifest as differences in information processing and/or functional

connectivity. In the following analyses, we explored each of these factors in turn.

Information processing and behavioural responsiveness during moderate anaesthesia

To measure information processing during the auditory narrative, we used a previously

established method (Naci et al., 2018; 2017; 2014), where we demonstrated that the extent of

stimulus-driven cross-subject correlation provides a measure of regional stimulus-driven

information processing. During wakefulness, we observed widespread and significant

(p<0.05; FWE cor) cross-subject correlation between healthy participants, bilaterally in

sensory-driven (primary and association) auditory cortex, visual cortex, as well as higher-

order frontal and parietal cortical regions (Figure 3A). By contrast, during moderate

anaesthesia, significant (p<0.05; FWE cor) cross-subject correlation was observed only in

primary and secondary auditory cortex and three small clusters (the left post central gyrus,

left paracentral lobule and right precentral gyrus) of premotor and motor cortex that have

been implicated in language comprehension (Hauk et al., 2004) (Figure 3B). As expected,

and consistent with previous studies (Naci et al., 2018; Davis et al., 2007), these results

suggested reduced information processing during moderate anaesthesia relative to

wakefulness.

Figure 3. about here please

We delved deeper into these effects to investigate whether each individual's sensory-driven

(auditory) perceptual processes and high-order attention processes during the story condition,

related to their response times in the target detection task. First, we used SPM to model the

relationship between the story's sound envelope and changes in brain activity over time. At

the group level, during wakefulness, the sound envelope predicted brain activity in bilateral

primary and secondary auditory cortex and right inferior frontal gyrus (Figure 3C). This

activity was dramatically reduced in moderate anaesthesia to one small cluster in left primary auditory cortex (superior temporal gyrus) that did not reach statistical significance (p=0.05 FWE cor) (Figure 3D). At the individual level, the sound envelope predicted significant activations in the auditory regions in 14/17 participants (3/17 showing subthreshold activations) during wakefulness, and in 10/17 during moderate anaesthesia. We observed no correlation between the participants' reaction times in either wakefulness or moderate anaesthesia and the extent of their individual activations in auditory regions (Figure 3 G–H).

Second, to investigate differential recruitment of high-order attention processes during the story, we used a previously established qualitative measure of the listeners' sustained attention throughout the narrative — the perception of suspense on a moment-by-moment basis (Naci et al., 2014). Beyond basic physical properties captured in the sound envelope, such as the amplitude and tone of the musical soundtrack, 'suspense,' in dramatic narratives such as the one used in this study, arises through ongoing sustained attention, as the current features of the movie (e.g., a young girl is alone in a foreign country, far away from her family) are compared to stored knowledge of the world (e.g., human traffickers pray on vulnerable targets), what has happened previously in the story (e.g., the girl's close friend has been kidnapped), and what may happen in future (e.g., she may be kidnapped).

Again, we used SPM to model the relationship between this qualitative measure of the story's ongoing high-level attention demands and changes in brain activity over time, measured in the independent group of participants, who had listened to the story without a secondary task in the fMRI scanner, during the wakeful and moderate anaesthesia states. During wakefulness, suspense ratings significantly (p<0.05; FWE cor) predicted activity in regions of the auditory attention network (the left superior temporal gyrus, left and right inferior frontal cortex, the inferior precentral gyrus/sulcus) (Tobyne et al., 2018; Michalka et al., 2015; Naci et al., 2013) and the salience network (the left/right anterior cingulate, insula, inferior frontal, and the thalamus) (Seeley 2019; Menon 2015), with clips rated as 'highly suspenseful' predicting stronger activity in this networks (Table 2) (Figure 3E). As expected, during moderate anaesthesia, the activation predicted by suspense ratings was reduced to one cluster extending in right superior temporal and marginal gyri (Figure 3F).

Table 2 about here please

In the behavioural group, the suspense ratings throughout the narrative showed a very robust

(r=0.90; SE=0.07)) and significant inter-subject correlation (t(24)=20.56, p=9.6e-17),

confirming the common understanding of the story across different individuals listening to

this narrative. In previous work, we have shown that a common neural code that is reliably

present in individual participants underlies common understanding of engaging audio-visual

narratives (Naci et al., 2017; 2014). Therefore, the similarity of the story's suspense ratings

enabled model-based predictions of the underlying brain activity that could be applied to

individual participants. The suspense ratings predicted significant activations in the auditory

attention network and salience network regions in 17/17 of participants during wakefulness

and 14/17 during moderate anaesthesia. We observed no correlation between the participants'

reaction time in wakefulness or moderate anaesthesia and the extent of their individual

activation in the auditory attention and salience networks regions (Figure 3 I–J).

In summary, these results suggested that differences in behavioural responsiveness under

anaesthesia, were neither driven by differences in auditory perceptual processes, nor in

sustained attention processes and the ability to integrate complex and dynamic information

that unveils over time.

Network connectivity and behavioural responsiveness during moderate anaesthesia

Rather, alterations in the connectivity within and between the DAN, ECN, and DMN during

moderate anaesthesia may drive these individual differences. We observed that functional

connectivity within the ECN was the only FC type that differentiated the two response

groups, with significantly higher FC for FRs relative to SRs, in the narrative condition during

wakefulness (permutation test, $p < 0.005$, FDR corrected) (Figure 4). No FC differences

between FR and SR were observed within the DAN or DMN, or between any of the three

network pairs.

477   Figure 4. about here please

To further investigate the within–ECN FC differences between FRs and SRs, we performed a

2x2x2 ANOVA that explored the main effects of *condition* (narrative, resting state), *state*

(wakefulness, moderate anaesthesia) and *response group* (FRs, SRs), as well as any

interaction among them. We found a main effect of response group (F (1,15) = 7.40, $p <$

0.05) that was driven by higher connectivity for FRs, and no effects on condition or state.

Although the 2x2x2 ANOVA did not reveal any interactions, effect size analyses showed a

higher difference between FRs and SRs in the narrative condition during wakefulness

(Hedges' g = 1.58) relative to moderate anaesthesia (Hedges' g = 1.10), and relative to the

resting state condition during wakefulness (Hedges' g = 0.68), and moderate anaesthesia

(Hedges' g = 0.96). In summary, the within–ECN functional connectivity during the narrative

condition in the wakeful state most strongly related to the individual differences in

behavioural responsiveness observed during moderate anaesthesia.

Structural brain differences and behavioural responsiveness during moderate anaesthesia

Finally, we tested whether individual differences in behavioural responsiveness were related

to structural brain differences, which may also underlie functional connectivity differences.

To this end, we performed whole-brain vertex-based comparisons of grey matter volume

between FRs and SRs. We found that FRs had uniformly significantly higher grey matter

volume relative to the SRs, in five regions bilaterally distributed: the left and right superior

and dorsolateral frontal cortex (L SFC, R SFC), including the presupplementary motor area

(pre-SMA), left and right rostral middle frontal cortex (L rMFC, R rMFC) and the right

precentral cortex (R Precentral) (Figure 5 and Table 3). These regions overlapped with the

three functional ROIs that defined the frontal nodes of the executive control network (Table

1). The opposite effect —higher grey matter for SR than FR—was not observed anywhere in

the brain. These results lend support to the functional connectivity results above, and together

they strongly suggest that connectivity within the ECN, and especially the frontal aspect of

this network, underlies individual differences in behavioural responsiveness under moderate

anaesthesia.

Figure 5. about here please

Table 3 about here please

This was further confirmed by a direct comparison of the FC differences between FRs and

SRs in the ROIs covering the frontal (dorsal medial PFC, left and right anterior PFC) and

parietal (left and right superior parietal cortex) aspects of the ECN separately. A 2x2x2

ANOVA with factors *state* (wakefulness, moderate anaesthesia), *region* (frontal ECN,

parietal ECN), and *response group* (FRs, SRs) showed a main effect of region [F (1,15) =

11.791, $p < 0.01$], driven by higher connectivity in the parietal aspect of the ECN, and three-

way interactions (state x region x response group) [F (1,15) = 8.358, $p < 0.05$). During

wakefulness, connectivity within the frontal or the parietal aspects of the ECN,

independently, could distinguish FRs and SRs (permutation test, $p < 0.05$ for frontal aspect;

permutation test, $p < 0.05$ for parietal aspect, FDR corrected). By contrast, in the moderate

anaesthesia, only FC in the frontal aspect could significantly distinguish the FRs and SRs

(permutation test, $p < 0.05$, FDR corrected) (Figure 6).

Figure 6. about here please

Discussion

Although anaesthesia has been used for over 150 years to reversibly abolish consciousness in
clinical medicine, the brain-bases of its considerable inter-individual effect variability remain
poorly understood. To address this gap, we asked whether the connectivity within and
between three networks that are primary sites of propofol-induced sedation and key to
conscious cognition — the DAN, ECN and DMN — underlie individual responsiveness
differences under anaesthesia. Consistent with previous studies (Chennu et al., 2016; Bola et
al., 2019), we observed substantial individual differences in reaction times under moderate
anaesthesia. These were not related to differences in perceptual or higher-order information
processing. Rather, only the functional connectivity within the ECN during the wakeful
narrative condition differentiated the participants' responsiveness level, with significantly
stronger ECN connectivity in the fast, relative to slow responders. These results were further
supported by our finding that fast responders had significantly higher grey matter volume in
the frontal cortex aspect of the executive control network. Taken together, the functional and
structural imaging results, provided very strong evidence that individual differences in
responsiveness under moderate anaesthesia related to inherent differences in brain function
and structure within the executive control network, which can be predicted prior to sedation.

Consistent with previous studies, we found that the antagonistic relationship between the
DAN/ECN and the DMN was reduced during moderate anaesthesia (Bonhomme et al., 2012;
Boveroux et al., 2010), but the connectivity between the DAN/ECN and DMN, or within the
DMN network showed no relationship to individual responsiveness variability. In addition to
orchestrating internally-oriented cognition (Andrews-Hanna et al. 2010; Schneider et al.,
2008; Buckner et al., 2008; Beer 2007; D'Argembeau et al., 2005; Wicker et al., 2003;
Gusnard et al., 2001), the DMN was recently found to play a role in broad associative
processes (Smith et al., 2018;Vatansever et al., 2017). The absence of a relationship between
the DMN connectivity and responsiveness differences is fitting given that the target detection
task did not elicit rich episodic information or any contextual shifts. Furthermore, it suggests
that individual differences in responsiveness were not driven by differences in internally-
oriented, or self-referential cognitive processes.

The differential impact of moderate anaesthesia on cognitive process can be inferred in a post
hoc manner, based on the perceptual and cognitive processes recruited by the target detection

task and the brain networks and regions that showed functional and/or structural differences
between the participants. The task involved a hierarchy of perceptual and cognitive
processes. These included (a) auditory perception, (b) understanding of linguistic instruction,
(c) short-term memory to remember instructions for the duration of the experiment, (d)
attention to auditory stimuli, (e) attention to intention to make a response, (f) cognitive
control over other, competing and distracting mental processes, and the (g) execution of a
motor response, when prompted by the auditory stimuli. The Ramsay clinical assessment
determined similar basic language communication (b), and we found short-term memory
function (c) across participants during moderate anaesthesia. Furthermore, our results showed
no relationship between behavioural responsiveness and sensory/auditory (a), or higher-order,
including linguistic information-processing that relied on sustained attention (b, d), across
participants. Therefore, differences in responsiveness were likely underpinned by differences
in complex mental faculties, such as attention to intention, cognitive control and action
execution.

Our functional connectivity results demonstrated that FC within the ECN differentiated the
participants' level of responsiveness, suggesting that moderate anaesthesia impacts processes
of goal and action execution subserved by the ECN (Marek and Dosenbach 2018; Dosenbach
et al., 2007; Ernst and Paulus 2005). Furthermore, our structural imaging results
demonstrated that grey matter volume differences in the frontal cortex aspect of the ECN,
including the dorsolateral prefrontal cortex (DLPFC) and pre-supplementary motor area (pre-
SMA), underscore individual responsiveness differences. In addition to their role in action
selection and execution (Lee et al., 1999; Boly et al., 2007), the DLPFC and pre-SMA
represent higher-order attention, i.e., attention to intention. Lau et al (2004) investigated the
brain regions that were preferentially involved in attending to the intention to move, relative
to the actual movement, and found that brain activity in pre-SMA, as well as its functional
connectivity with right DPFC represented the intention to move. The pre-SMA has also been
implicated in sustained cognitive control (Nachev et al., 2005), another process that is key to
successful task performance. In summary, a post-hoc interpretation of our findings would be
that moderate anaesthesia impacts attention to intention, sustained cognitive control, and
action execution, and that individuals with smaller grey matter volume in frontal regions and
weaker functional connectivity within the ECN may have a vulnerability for stronger
suppression of behavioural responsiveness than those with higher values for these features.

As the suppression of behavioural responsiveness is one of the key aims of anaesthesia, the

converse finding was also highly relevant. Surprisingly, we found that 30% of participants

showed no delay in reaction times under moderate anaesthesia relative to wakefulness, and

critically, these exhibited inherent brain differences to participants who were significantly

delayed. This result is highly relevant to the prediction of responsiveness variability under

anaesthesia (Chennu et al., 2016; Palanca et al 2009) and monitoring depth-of-anaesthesia for

the detection of unintended intraoperative awareness. Although rare (0.1-0.2%, Mashour and

Avidan 2015; Sandin et al., 2000), unintended awareness can be very traumatic and lead to

negative long-term health outcomes, such as post-traumatic stress disorder (up to 70%), as

well as clinical depression or phobias (Pandit et al., 2017; 2014; Mashour and Avidan 2015).

Given its rarity, unintended intraoperative awareness does not lend itself to direct

investigation in the relatively small groups of typical research studies with vulnerable

populations, including the present study, where individuals were deeply anaesthetized

without intubation in a research MRI setting.

Our results suggest that individuals with larger grey matter volume in frontal regions and

stronger functional connectivity within the ECN may require higher doses of propofol to

become non-responsive, to the same extent as individuals with smaller grey matter volume

and weaker connectivity in these regions. If replicated in a clinical context, these findings

will provide novel markers that may help to improve the accuracy of awareness monitoring

during clinical anaesthesia. It is worth noting that propofol was used here due to its wide

prominence in clinical interventions, and future studies that employ the same paradigm across

different agents will determine whether the brain-behaviour relationships we report here

generalize to other anaesthetic agents.

Finally, we observed higher power for detecting individual responsiveness differences under

anaesthesia from ECN functional connectivity in the wakeful narrative condition, than in the

resting state. This may be due to higher signal to noise during the narrative condition.

Previous studies have shown that the active attention engagement during well-crafted

narratives leads to reduced movement (Centeno et al., 2016) and less sleep in scanner

(Centeno et al., 2016, Vanderwal et al., 2015), resulting in higher signal-to-noise relative to

resting state (rs) fMRI (Wang et al., 2017). Moreover, it may be due to higher power to detect

functional roles of brain networks during naturalistic stimuli relative to rsfMRI. Rs fMRI is

entirely unconstrained, and thus it is difficult to separate signal due to cognitive processes

from extraneous sources, including motion, cardiac and respiratory physiological noise,

arterial CO_2 concentration, blood pressure, and vasomotion (Murphy et al., 2013; Birn et al.,

2006). By contrast, as we have previously established, naturalistic narratives drive diverging

responses across brain networks, highlighting their functionally distinct roles, and thus may

be more sensitive for investigating how their differing cognitive roles are impacted by

anaesthesia than rsfMRI (Naci et al., 2018; Haugg et al., 2018; Naci et al., 2014). Our results

are consistent with a growing recognition of the importance of naturalistic stimuli for

studying cognitive processes and understanding the neural basis of real-world functioning

(Sonkusare et al., 2019).

 Figures and Legends

628
629

Figure 1. Functional connectivity (FC) in wakeful and moderate anaesthesia states for the

narrative and resting state conditions. (A) Group-averaged FC connectivity matrices during

wakeful and moderate anaesthesia states in the narrative and resting state conditions across

all participants. The color bar indicates connectivity (measured as z-transformed Pearson

correlation; low-high, blue-red). Each cell in the FC matrix represents FC between each node

(brain region) of the three networks. (B) Comparisons of FC averaged across all nodes for

each network, within and between the three brain networks, during the wakeful and moderate

anaesthesia states, in the narrative and resting state conditions separately. FDR correction

was applied for multiple comparison comparisons. Abbreviations: DMN, default mode

network; DAN, dorsal attention network; ECN, executive control network.

Figure 2. Reaction time during an auditory target detection task in the wakeful and moderate

sedation states. Participants were divided into two groups by applying mean split method.

Blue (orange) bars represent reaction time for each individual during wakeful (moderate)

sedated state. Horizontal axis represents each individual (S1-S17=Subject1-Subject17) and

vertical axis represents averaged reaction time across 50 trails.

Figure 3. Perceptual and higher-order information processing during the audio story. (a) During wakefulness, the audio story elicited significant (p<0.05; FWE cor) inter-subject correlation across the brain, including frontal and parietal cortex, thought to support high-level information integration over time.(b) During moderate anaesthesia, significant (p<0.05; FWE cor) inter-subject correlation was limited to the primary and secondary auditory cortex and three small clusters in regions of premotor and motor cortex that are involved in language comprehension. (c)/(d) Clusters predicted by the sound envelope of the audio story in the (c) primary and secondary auditory cortex and left inferior frontal gyrus during wakefulness, and (d) left primary auditory cortex in moderate anaesthesia, that did not reach statistical significance (p=0.05 FWE). (e)/(f) Clusters predicted by the suspense ratings of the audio story in regions of the auditory attention network and salience network during wakefulness

and moderate anaesthesia. (G)/(H) The sound envelope predicted significant activations in the

auditory regions in 14/17 and 10/17 of participants during the wakefulness and moderate anaesthesia

states, respectively. (I)/(J) The suspense ratings predicted significant activations in the auditory

regions in 17/17 and 14/17 of participants during the wakefulness and moderate anaesthesia states,

respectively. (G)–(J) There was no correlation between the perceptual or higher-order

information processing during the narrative and reaction times during the target detection

task, either in wakefulness (G)/(H) or in moderate anaesthesia (H)/(J).

Figure 4. Functional connectivity differences between fast and slow responders. (A) FC maps for fast and slow responders during wakeful and moderate anaesthesia states in narrative and resting state conditions separately. Color bar indicates connectivity (z-transformed Pearson correlation; low-high: blue-red). Each cell in the FC map represents FC between each node (brain region) of the three networks. FC within-ECN is highlighted by the magenta square box in each FC map. (B) Comparisons of averaged FC within and between the three brain networks, between fast and slow responders for each condition and state separately. FDR correction was applied for multiple comparisons. Abbreviations: DMN, default mode network; DAN, dorsal attention network; ECN, executive control network.

Figure 5. Vertex-wise analysis of grey matter volume between fast and slow responders. (A) Five regions showed significant between-group difference in grey matter volume (GMV, $p <$ 0.05, Monte Carlo simulation corrected). Color bar indicates the p-value (-lg(p)). (B) Group comparisons of averaged GMV for each of the five clusters between fast and slow responders. Abbreviations: L SFC, left superior frontal cortex; R SFC, right superior frontal cortex; L rMFC, left rostral middle frontal cortex; R rMFC, right rostral middle frontal cortex. R Precentral, right precentral.

Figure 6. Differences of functional connectivity in frontal and parietal aspects of ECN between fast and slow responders in both wakeful and moderate anaesthesia states in narrative condition. Abbreviations: ECN, executive control network.

Table 1. Overview of selected regions of interests for three functional networks.

Network/Region	ROI	MNI coordinates		
		x	y	z
Default Mode Network	Posterior cingulate/precuneus	0	-52	27
	Medial prefrontal	-1	54	27
	Left lateral parietal	-46	-66	30
	Right lateral parietal	49	-63	33
	Left inferior temporal	-61	-24	-9
	Right inferior temporal	58	-24	-9
Dorsal Attention Network	Left frontal eye field	-29	-9	54
	Right frontal eye field	29	-9	54
	Left posterior IPS	-26	-66	48
	Right posterior IPS	26	-66	48
	Left anterior IPS	-44	-39	45
	Right anterior IPS	41	-39	45
	Left MT	-50	-66	-6
	Right MT	53	-63	-6
Executive Control Network	Dorsal medial PFC	0	24	46
	Left anterior PFC	-44	45	0
	Right anterior PFC	44	45	0
	Left superior parietal	-50	-51	45
	Right superior parietal	50	-51	45

699   Abbreviations: IPS: Intraparietal sulcus, MT: Middle temporal area, PFC: Prefrontal cortex

701   Table 2. Coordinates for activation cluster produced by the story's suspense ratings.

Cluster Index	Cluster Peak Location	Peak Talairach coordinates			Cluster-wise p-value
		x	y	z	
1	Cingulate Sulcus	-2	20	36	<0.000
2	Right Superior Temporal Gyrus	58	-36	14	<0.000
3	Right Anterior Insular Gyrus	40	24	8	<0.000
4	Posterior Cingulate Gyrus	14	-44	14	<0.000
5	Left Anterior Insular Gyrus	-36	14	-8	<0.000
6	Thalamus	4	-20	10	<0.005

704   Table 3. Regions showing significant grey matter volume difference between slow

705   responders and fast responders

Index	Peak location	Vertex number	Size (mm2)	Peak Talairach coordinates			Cluster-wise
				x	y	z	
1	L_SFC	6163	3338.75	-6.70	29.30	48.20	1.00E-04
2	L rMFC	3048	1954.78	-38.20	25.40	22.00	1.00E-04
3	R Precentral	3257	1830.76	37.70	-1.90	40.10	1.00E-04
4	R rMFC	1592	927.63	37.80	26.80	28.50	0.0256
5	R SFC	1718	904.29	7.00	46.80	34.60	0.0304

Notes: Peak coordinates were presented in the Talairach-like space. Abbreviations: L SFC,

left superior frontal cortex; R SFC, right superior frontal cortex; L rMFC, left rostral middle

frontal cortex; R rMFC, right rostral middle frontal cortex. R Precentral, right precentral.

Acknowledgements

F.D. was funded by the Provost PhD Award Scheme from Trinity College Dublin, to L.N.
L.N. was also funded by an L'Oreal for Women In Science International Rising Talent
Award, and the Welcome Trust Institutional Strategic Support grant. AMO is a CIFAR
Fellow and is supported by the Canadian Institutes for Health Research (CIHR) and the
Natural Sciences and Engineering Research Council of Canada (NSERC). Data acquisition
(fMRI) was supported by a Canada Excellence Research Chairs award to AMO (Grant No.
215063)

718
719

720 **Conflict of Interest**
721 The authors declare no conflict of interest.
722

723 **Author contributions**
724 Study Conceptualization, L.N., N.T.; Data Acquisition, L.N. Methodology, R.C., F.D., L.N.;
725 Formal Analyses, F.D., N.T.; Writing, F.D., L.N.; Funding and resource acquisition, A.M.O.,
726 L.N. Project administration, L.N.; Supervision, L.N.
727
728

References

Alkire, M. T., & Miller, J. (2005). General anesthesia and the neural correlates of consciousness. *Progress in brain research*, 150, 229-597.

Anderson, J. S., Druzgal, T. J., Lopez-Larson, M., Jeong, E. K., Desai, K., & Yurgelun-Todd, D. (2011). Network anticorrelations, global regression, and phase-shifted soft tissue correction. *Human brain mapping*, *32*(6), 919-934.

Andrews-Hanna, J. R., Reidler, J. S., Huang, C., & Buckner, R. L. (2010). Evidence for the default network's role in spontaneous cognition. *Journal of neurophysiology*, *104*(1), 322-335.

Baars, B. J., Ramsøy, T. Z., & Laureys, S. (2003). Brain, conscious experience and the observing self. *Trends in neurosciences*, 26(12), 671-675.

Beer, J. S. (2007). The default self: feeling good or being right?. *Trends in cognitive sciences*, *11*(5), 187-189.

Birn, R. M., Diamond, J. B., Smith, M. A., & Bandettini, P. A. (2006). Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. *Neuroimage*, 31(4), 1536-1548.

Bola, M., Orlowski, P., Plomecka, M., & Marchewka, A. (2019). EEG signal diversity during propofol sedation: an increase in sedated but responsive, a decrease in sedated and unresponsive subjects. *bioRxiv*, 444281.

Boly, M., Balteau, E., Schnakers, C., Degueldre, C., Moonen, G., Luxen, A., ... & Laureys, S. (2007). Baseline brain activity fluctuations predict somatosensory perception in humans. *Proceedings of the National Academy of Sciences*, 104(29), 12187-12192.

Boly, Mélanie, Christophe Phillips, Luaba Tshibanda, Audrey Vanhaudenhuyse, M. Schabus, Thien Thanh Dang-Vu, Gustave Moonen, Roland Hustinx, Pierre Maquet, and Steven Laureys. "Intrinsic brain activity in altered states of consciousness: how conscious is the default mode of brain function?." *Annals of the New York Academy of Sciences* 1129 (2008): 119.

Boly, M., Tshibanda, L., Vanhaudenhuyse, A., Noirhomme, Q., Schnakers, C., Ledoux, D., ... & Luxen, A. (2009). Functional connectivity in the default network during resting state is preserved in a vegetative but not in a brain dead patient. *Human brain mapping*, 30(8), 2393-2400.

Bonhomme, V. L. G., Boveroux, P., Brichant, J. F., Laureys, S., & Boly, M. (2012). Neural correlates of consciousness during general anesthesia using functional magnetic resonance imaging (fMRI). *Archives italiennes de biologie*, 150(2/3), 155-163.

Boveroux, P., Vanhaudenhuyse, A., Bruno, M. A., Noirhomme, Q., Lauwick, S., Luxen, A., ... & Brichant, J. F. (2010). Breakdown of within-and between-network resting state functional magnetic resonance imaging connectivity during propofol-induced loss of consciousness. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, *113*(5), 1038-1053.

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: anatomy, function, and relevance to disease.

Buckner, R. L., Krienen, F. M., & Yeo, B. T. (2013). Opportunities and limitations of intrinsic functional connectivity MRI. *Nature neuroscience*, 16(7), 832.

Carhart-Harris, R. L., & Friston, K. J. (2010). The default-mode, ego-functions and free-
energy: a neurobiological account of Freudian ideas. *Brain*, *133*(4), 1265-1283.

Centeno, M., Tierney, T. M., Perani, S., Shamshiri, E. A., StPier, K., Wilkinson, C., ... &
Pressler, R. M. (2016). Optimising EEG-fMRI for localisation of focal epilepsy in
children. *PloS one*, 11(2).

Chennu, S., O'Connor, S., Adapa, R., Menon, D. K., & Bekinschtein, T. A. (2016). Brain
connectivity dissociates responsiveness from drug exposure during propofol-induced
transitions of consciousness. *PLoS computational biology*, *12*(1), e1004669.

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven
attention in the brain. *Nature reviews neuroscience*, *3*(3), 201-215.

Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C., &
Peelle, J. E. (2015). Automatic analysis (aa): efficient neuroimaging workflows and parallel
processing using Matlab and XML. *Frontiers in neuroinformatics*, *8*, 90.

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis: I.
Segmentation and surface reconstruction. *Neuroimage*, *9*(2), 179-194.

D'Argembeau, A., Collette, F., Van der Linden, M., Laureys, S., Del Fiore, G., Degueldre, C.,
... & Salmon, E. (2005). Self-referential reflective activity and its relationship with rest: a
PET study. *Neuroimage*, *25*(2), 616-624.

Davis, M. H. et al. Dissociating speech perception and comprehension at reduced levels of
awareness. *Proc. Natl. Acad. Sci*. USA 104, 16032–16037 (2007).

Demertzi, A., Soddu, A., & Laureys, S. (2013). Consciousness supporting networks. *Current
Opinion in Neurobiology*, *23*(2), 239-244.

Demertzi, A., Tagliazucchi, E., Dehaene, S., Deco, G., Barttfeld, P., Raimondo, F., ... &
Schiff, N. D. (2019). Human consciousness is supported by dynamic complex patterns of
brain signal coordination. *Science advances*, 5(2), eaat7603.

Dosenbach, N. U., Fair, D. A., Miezin, F. M., Cohen, A. L., Wenger, K. K., Dosenbach, R.
A., ... & Schlaggar, B. L. (2007). Distinct brain networks for adaptive and stable task control
in humans. *Proceedings of the National Academy of Sciences*, *104*(26), 11073-11078.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs
for intelligent behaviour. *Trends in cognitive sciences*, *14*(4), 172-179.

Elliott, R. (2003). Executive functions and their disorders: Imaging in clinical
neuroscience. *British medical bulletin*, 65(1), 49-59.

Ernst, M., & Paulus, M. P. (2005). Neurobiology of decision making: a selective review from
a neurocognitive and clinical perspective. *Biological psychiatry*, *58*(8), 597-604.

Fiset, P., Paus, T., Daloze, T., Plourde, G., Meuret, P., Bonhomme, V., ... & Evans, A. C.
(1999). Brain mechanisms of propofol-induced loss of consciousness in humans: a positron
emission tomographic study. Journal of Neuroscience, 19(13), 5506-5513.

Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state": a practical
method for grading the cognitive state of patients for the clinician. *Journal of psychiatric
research*, *12*(3), 189-198.

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E.
(2005). The human brain is intrinsically organized into dynamic, anticorrelated functional
networks. *Proceedings of the National Academy of Sciences*, 102(27), 9673-9678.

Fransson, P. (2005). Spontaneous low-frequency BOLD signal fluctuations: An fMRI
investigation of the resting-state default mode of brain function hypothesis. *Human brain*
mapping, 26(1), 15-29.

Greicius, M. D., Krasnow, B., Reiss, A. L., & Menon, V. (2003). Functional connectivity in
the resting brain: a network analysis of the default mode hypothesis. *Proceedings of the*
National Academy of Sciences, *100*(1), 253-258.

Gusnard, D. A., Akbudak, E., Shulman, G. L., & Raichle, M. E. (2001). Medial prefrontal
cortex and self-referential mental activity: relation to a default mode of brain
function. *Proceedings of the National Academy of Sciences*, *98*(7), 4259-4264.

Hahn, B., Ross, T. J., & Stein, E. A. (2007). Cingulate activation increases dynamically with
response speed under stimulus unpredictability. *Cerebral cortex*, 17(7), 1664-1671.

Haugg, A., Cusack, R., Gonzalez-Lara, L. E., Sorger, B., Owen, A. M., & Naci, L. (2018).
Do patients thought to lack consciousness retain the capacity for internal as well as external
awareness?. *Frontiers in neurology*, *9*, 492.

Hauk, O., Johnsrude, I., & Pulvermüller, F. (2004). Somatotopic representation of action
words in human motor and premotor cortex. *Neuron*, *41*(2), 301-307.

Honey, C. J., Newman, E. L., & Schapiro, A. C. (2017). Switching between internal and
external modes: a multiscale learning principle. *Network Neuroscience*, 1(4), 339-356.

Huang, Z., Zhang, J., Wu, J., Mashour, G. A., & Hudetz, A. G. (2020). Temporal circuit of
macroscale dynamic brain activity supports human consciousness. *Science advances*, *6*(11),
eaaz0087.

Kelly, A. C., Uddin, L. Q., Biswal, B. B., Castellanos, F. X., & Milham, M. P. (2008).
Competition between functional brain networks mediates behavioral
variability. *Neuroimage*, 39(1), 527-537.

Kroger, J. K., Sabb, F. W., Fales, C. L., Bookheimer, S. Y., Cohen, M. S., & Holyoak, K. J.
(2002). Recruitment of anterior dorsolateral prefrontal cortex in human reasoning: a
parametric study of relational complexity. *Cerebral cortex*, *12*(5), 477-485.

Kucyi, A., Daitch, A., Raccah, O., Zhao, B., Zhang, C., Esterman, M., ... & Parvizi, J. (2020).
Electrophysiological dynamics of antagonistic brain networks reflect attentional
fluctuations. *Nature Communications*, 11(1), 1-14.

Lau, H. C., Rogers, R. D., Haggard, P., & Passingham, R. E. (2004). Attention to
intention. *science*, *303*(5661), 1208-1210.

Lee, K. M., Chang, K. H., & Roh, J. K. (1999). Subregions within the supplementary motor
area activated at different stages of movement preparation and execution. *Neuroimage*, *9*(1),
117-123.

Leslie, K., Skrzypek, H., Paech, M. J., Kurowski, I., & Whybrow, T. (2007). Dreaming
during Anesthesia and Anesthetic Depth in Elective Surgery PatientsA Prospective Cohort
Study. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 106(1), 33-
42.

Luppi, A., Craig, M., Pappas, I., Finoia, P., Williams, G., Allanson, J., ... & Stamatakis, E.
(2019). Consciousness-specific dynamic interactions of brain integration and functional
diversity.

MacDonald, A. A., Naci, L., MacDonald, P. A., & Owen, A. M. (2015). Anaesthesia and
neuroimaging: investigating the neural correlates of unconsciousness. *Trends in Cognitive*
Sciences, *19*(2), 100-107.

Marek, S., & Dosenbach, N. U. (2018). The frontoparietal network: function,
electrophysiology, and importance of individual precision mapping. *Dialogues in clinical*
neuroscience, *20*(2), 133.

Marsh, B. M. W. M. N., White, M., Morton, N., & Kenny, G. N. C. (1991). Pharmacokinetic
model driven infusion of propofol in children. *BJA: British Journal of Anaesthesia*, *67*(1),
41-48.

Mashour, G. A., & Avidan, M. S. (2015). Intraoperative awareness: controversies and non-
controversies. *British journal of anaesthesia*, *115*(suppl_1), i20-i26.

Mashour, G. A., & Hudetz, A. G. (2018). Neural correlates of unconsciousness in large-scale
brain networks. *Trends in neurosciences*, 41(3), 150-160.

Menon, V. (2015). Salience network. In A. W. Toga (Ed.), Brain mapping: An encyclopedic
reference (Vol. 2, pp. 597–611). London: Academic Press: Elsevier.

Michalka, S. W., Kong, L., Rosen, M. L., Shinn-Cunningham, B. G., & Somers, D. C.
(2015). Short-term memory for space and time flexibly recruit complementary sensory-biased
frontal lobe attention networks. Neuron, 87(4), 882-892.

Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The
impact of global signal regression on resting state correlations: are anti-correlated networks
introduced?. *Neuroimage*, *44*(3), 893-905.

Murphy, K., Birn, R. M., & Bandettini, P. A. (2013). Resting-state fMRI confounds and
cleanup. *Neuroimage*, 80, 349-359.

Naci, L., Cusack, R., Jia, V. Z., & Owen, A. M. (2013). The brain's silent messenger: using
selective attention to decode human thought for brain-based communication. *Journal of*
Neuroscience, *33*(22), 9385-9393.

Naci, L., Cusack, R., Anello, M., & Owen, A. M. (2014). A common neural code for similar
conscious experiences in different individuals. *Proceedings of the National Academy of*
Sciences, 111(39), 14277-14282.

Naci, L., Sinai, L., & Owen, A. M. (2017). Detecting and interpreting conscious experiences
in behaviorally non-responsive patients. *NeuroImage*, *145*, 304-313.

Naci, L., Haugg, A., MacDonald, A., Anello, M., Houldin, E., Naqshbandi, S., ... & Owen, A.
M. (2018). Functional diversity of brain networks supports consciousness and verbal
intelligence. *Scientific reports*, *8*(1), 13259.

Nachev, P., Rees, G., Parton, A., Kennard, C., & Husain, M. (2005). Volition and conflict in
human medial frontal cortex. *Current Biology*, 15(2), 122-128.

Pal, D., Li, D., Dean, J. G., Brito, M. A., Liu, T., Fryzel, A. M., ... & Mashour, G. A. (2020).
Level of consciousness is dissociable from electroencephalographic measures of cortical
connectivity, slow oscillations, and complexity. *Journal of Neuroscience*, *40*(3), 605-618.

Palanca, B. J. A., Mashour, G. A., & Avidan, M. S. (2009). Processed electroencephalogram
in depth of anesthesia monitoring. *Current Opinion in Anesthesiology*, *22*(5), 553-559.

Pandit, J. J., Andrade, J., Bogod, D. G., Hitchman, J. M., Jonker, W. R., Lucas, N., ... & Paul,
R. G. (2014). 5th National Audit Project (NAP5) on accidental awareness during general

anaesthesia: summary of main findings and risk factors. *British journal of anaesthesia*, *113*(4), 549-559.

Pandit JJ, Cook TM; The NAP5 Steering Panel. NAP5. Accidental Awareness During General Anesthesia in the United Kingdom and Ireland: Report and Findings. *London: The Royal College of Anesthetists and Association of Anesthetists of Great Britain and Ireland*; 2014. Available at: http://www.nationalauditprojects.org.uk/NAP5_home. Accessed on May 3, 2017.

Peigneux, P., Orban, P., Balteau, E., Degueldre, C., Luxen, A., Laureys, S., & Maquet, P. (2006). Offline persistence of memory-related cerebral activity during active wakefulness. *PLoS biology*, *4*(4).

Plourde, G., Belin, P., Chartrand, D., Fiset, P., Backman, S. B., Xie, G., & Zatorre, R. J. (2006). Cortical processing of complex auditory stimuli during alterations of consciousness with the general anesthetic propofol. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, 104(3), 448-457.

Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences*, 98(2), 676-682.

Raichle, M. E. (2011). The restless brain. *Brain connectivity*, *1*(1), 3-12.

Ramsay, M. A. E., Savege, T. M., Simpson, B. R. J., & Goodwin, R. (1974). Controlled sedation with alphaxalone-alphadolone. *Br med J*, *2*(5920), 656-659.

Sanders, R. D., Gaskell, A., Raz, A., Winders, J., Stevanovic, A., Rossaint, R., ... & Meier, S. (2017). Incidence of Connected Consciousness after Tracheal IntubationA Prospective, International, Multicenter Cohort Study of the Isolated Forearm Technique. *Anesthesiology: The Journal of the American Society of Anesthesiologists*, *126*(2), 214-222.

Sandin, R. H., Enlund, G., Samuelsson, P., & Lennmarken, C. (2000). Awareness during anaesthesia: a prospective case study. *The Lancet*, *355*(9205), 707-711.

Sarasso, S., Boly, M., Napolitani, M., Gosseries, O., Charland-Verville, V., Casarotto, S., ... & Rex, S. (2015). Consciousness and complexity during unresponsiveness induced by propofol, xenon, and ketamine. *Current Biology*, 25(23), 3099-3105.

Seeley, W. W. (2019). The salience network: a neural system for perceiving and responding to homeostatic demands. Journal of Neuroscience, 39(50), 9878-9882.

Schneider, F., Bermpohl, F., Heinzel, A., Rotte, M., Walter, M., Tempelmann, C., ... & Northoff, G. (2008). The resting brain and our self: self-relatedness modulates resting state neural activity in cortical midline structures. *Neuroscience*, 157(1), 120-131.

Searle, R., & Hopkins, P. M. (2009). Pharmacogenomic variability and anaesthesia. *British journal of anaesthesia*, *103*(1), 14-25.

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., ... & Greicius, M. D. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *Journal of Neuroscience*, *27*(9), 2349-2356.

Shallice, T. (1988). *From neuropsychology to mental structure*. Cambridge University Press.

Shulman, G. L., Fiez, J. A., Corbetta, M., Buckner, R. L., Miezin, F. M., Raichle, M. E., & Petersen, S. E. (1997). Common blood flow changes across visual tasks: II. Decreases in cerebral cortex. *Journal of cognitive neuroscience*, *9*(5), 648-663.

Smith, V., Mitchell, D. J., & Duncan, J. (2018). Role of the default mode network in
cognitive transitions. *Cerebral Cortex*, 28(10), 3685-3696.

Sonkusare, S., Breakspear, M., & Guo, C. (2019). Naturalistic Stimuli in Neuroscience:
Critically Acclaimed. *Trends in cognitive sciences*.

Spreng, R. N., DuPre, E., Selarka, D., Garcia, J., Gojkovic, S., Mildner, J., ... & Turner, G. R.
(2014). Goal-congruent default network activity facilitates cognitive control. *Journal of*
Neuroscience, 34(42), 14108-14114.

Sridharan, D., Levitin, D. J., & Menon, V. (2008). A critical role for the right fronto-insular
cortex in switching between central-executive and default-mode networks. *Proceedings of the*
National Academy of Sciences, 105(34), 12569-12574.

Stamatakis, E. A., Adapa, R. M., Absalom, A. R., & Menon, D. K. (2010). Changes in resting
neural connectivity during propofol sedation. *PloS one*, *5*(12).

Tagliazucchi, E., von Wegner, F., Morzelewski, A., Brodbeck, V., Jahnke, K., & Laufs, H.
(2013). Breakdown of long-range temporal dependence in default mode and attention
networks during deep sleep. *Proceedings of the National Academy of Sciences*, 110(38),
15419-15424.

Tobyne, S. M., Somers, D. C., Brissenden, J. A., Michalka, S. W., Noyce, A. L., & Osher, D.
E. (2018). Prediction of individualized task activation in sensory modality-selective frontal
cortex with 'connectome fingerprinting'. Neuroimage, 183, 173-185.

Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C., & Castellanos, F. X. (2015). Inscapes: A
movie paradigm to improve compliance in functional magnetic resonance
imaging. *Neuroimage*, 122, 222-232.

Vanhaudenhuyse, A., Demertzi, A., Schabus, M., Noirhomme, Q., Bredart, S., Boly, M., ... &
Laureys, S. (2011). Two distinct neuronal networks mediate the awareness of environment
and of self. *Journal of cognitive neuroscience*, 23(3), 570-578.

Varley, T. F., Luppi, A. I., Pappas, I., Naci, L., Adapa, R., Owen, A. M., ... & Stamatakis, E.
A. (2020). consciousness & Brain functional complexity in propofol Anaesthesia. *Scientific*
reports, *10*(1), 1-13.

Vatansever, D., Manktelow, A. E., Sahakian, B. J., Menon, D. K., & Stamatakis, E. A.
(2017). Angular default mode network connectivity across working memory load. *Human*
brain mapping, *38*(1), 41-52.

Vatansever, D., Schröter, M., Adapa, R. M., Bullmore, E. T., Menon, D. K., & Stamatakis, E.
A. (2020). Reorganisation of Brain Hubs across Altered States of consciousness. *Scientific*
Reports, 10(1), 1-11.

Wang, J., Ren, Y., Hu, X., Nguyen, V. T., Guo, L., Han, J., & Guo, C. C. (2017). Test–retest
reliability of functional connectivity networks during naturalistic fMRI paradigms. *Human*
brain mapping, 38(4), 2226-2241.

Wicker, B., Ruby, P., Royet, J. P., & Fonlupt, P. (2003). A relation between rest and the self
in the brain?. *Brain Research Reviews*, *43*(2), 224-230.

Witon, A., Shirazibehehsti, A., Aviles, A., Adapa, R., Menon, D., Chennu, S., ... & Bowman,
H. (2020). Sedation modulates fronto-temporal predictive coding circuits and the double
surprise acceleration effect. *bioRxiv*.