Emotional Metacognition:

Stimulus Valence Modulates Cardiac Arousal and Metamemory

Nicolas Legrand[1*], Sebastian Scott Engen[1*], Camile Maria Costa Correa[1], Nanna Kildahl Mathiasen[1], Niia Nikolova[1], Francesca Fardo[1,2], Micah Allen[1,3,4]

[1] Center of Functionally Integrative Neuroscience, Aarhus University Hospital, Denmark
[2] Danish Pain Research Centre, Aarhus University Hospital, Denmark
[3] Aarhus Institute of Advanced Studies, Aarhus University, Denmark
[4] Department of Psychiatry, Cambridge University, Cambridge, UK

* Equal contribution.

Corresponding author: nicolas.legrand@cfin.au.dk

Keywords: *emotion, arousal, heart rate variability, metacognition, memory*

Abstract

Emotion alters how we feel, see, and experience the world. In the domain of memory, the emotional valence and arousal of memorised stimuli can modulate both the acuity and content of episodic recall. However, no experiment has investigated whether arousal and valence also influence metacognition for memory (i.e., the process of self-monitoring memories). In a pre-registered study, we applied a novel psychophysiological design together with computational models of metacognition to assess the influence of stimulus valence and arousal on the sensitivity, bias, and efficiency of metamemory. To estimate the role of physiological arousal in mediating these effects, we recorded cardiac measures through pulse oximetry. We found that negative valence substantially decreased both memory performance and subjective confidence, in particular for low arousal words. Simultaneously, we found that emotional valence modulated both heart rate and heart-rate variability (HRV) during recognition memory. Exploratory trial-level analyses further revealed that subjective confidence was encoded in instantaneous heart-rate fluctuations and that this relationship was also modulated by emotional valence. Our results demonstrate that recognition memory and metacognition are influenced by the emotional valence of encoded items and that this correlation is in part related to cardiac activity.

Introduction

The cognitive ability to monitor our thoughts, memories and perceptual experiences is an important part of learning, development and communication (Fleming et al., 2012; Heyes et al., 2020; Shea et al., 2014). Metacognition refers to this executive capacity to control lower-order representations by assessing the fidelity and strength of these signals and updating a model of the probability that one is making correct judgements (Yeung & Summerfield, 2012). More specifically, metacognition for memory or metamemory refers to the ability to assess the accuracy and precision of the recollection of previous experiences (Baird et al., 2013; McCurdy

et al., 2013; Ye et al., 2018). Memories are fragile internals signals that are prone to substantive
decay over time (Davis & Zhong, 2017; Otgaar et al., 2019). Episodic recall can be biased by
emotion and arousal both in the context of encoding (Yonelinas & Ritchey, 2015) and during
active recall (Ochsner, 2000). The metacognition of memory recall can, in turn, be influenced
by the level of details and the "feeling-of-knowing" associated with an episode (Chua et al.,
2014; Reggev et al., 2011). In the context of witness testimony for example, if the suspect had
an unremarkable face or the memory was hazy, then a witness may report lower confidence in
their recollection. A reliable witness should therefore not only recall events as experienced in
detail but also accurately assess the fidelity or confidence associated with those memories. As
little is currently known about the ability to self-monitor memory for emotional stimuli, we
conducted a confirmatory, pre-registered investigation of emotional metamemory.

In controlled laboratory settings, classic metacognition experiments often require
participants to view a stimulus, make a decision (e.g., whether the stimulus is known or
unknown), and report their confidence in this judgement. Healthy individuals typically display
reasonably accurate metacognitive insight and achieve a high correlation between confidence
and accuracy, even in the absence of external feedback. Metacognition tasks have been applied
to investigate a variety of cognitive domains including visual perception (Allen et al., 2016;
Fleming et al., 2015), memory (Baird et al., 2013; Fleming et al., 2014; McCurdy et al., 2013),
and value-based decision-making (De Martino et al., 2013). However, even with these simple
lab-based tasks, participants exhibit substantive interindividual differences in metacognitive
ability, and a variety of manipulations can reliably dissociate confidence and accuracy by
biasing subjective confidence reports (Fleming et al., 2015; Rollwage et al., 2020).

Though little is known about how emotion influences metacognition, previous
investigations of memory and emotion highlight stimulus valence and arousal as likely sources
of bias. For example, words rated as highly arousing or positively valenced are recognized and
detected faster (Kever et al., 2019), and the emotional content of valenced words, either positive
or negative, can increase the learner's confidence in subsequent accurate recall (Tauber &
Dunlosky, 2012). Similarly, physiological arousal can promote the awareness of concepts
congruent with the bodily state (Kever et al., 2015, 2017), and the level of arousal at encoding
can enhance subsequent memory performance (Anderson, 2005; Cahill & McGaugh, 1998) via
greater amygdala activity (Kensinger & Corkin, 2004). Accordingly, flashbulb memories (i.e.,
vivid and detailed memories encoded under arousing conditions) are recalled more easily and
with less decay under specific circumstances (Shields et al., 2017; Yonelinas & Ritchey, 2015).
This line of evidence suggests that emotional content, especially those of a highly arousing or
negative nature, could bias the salience of the memory signal during recall. This can ultimately
result in overconfidence which, in the context of testimony, could bias the individual when
estimating the accuracy of his or her recall.

Additionally, a core aspect of emotion is that it often coincides with and is triggered by
changes in internal bodily states (James, 1884), which is expressed by indices of autonomic
activity such as cardiac or respiratory frequency (Kreibig, 2010; Valenza et al., 2014). Heart
rate, for example, can be altered both when perceiving emotional stimuli and during their
encoding and recollection (Abercrombie et al., 2008; Critchley et al., 2005; Legrand et al.,
2020). These bodily changes exert a substantial effect on the mapping between confidence and
decision accuracy, which can ultimately also bias metacognition. For example, both
experimental and pharmacological modulations of arousal have been shown to bias

metacognitive insight, modulating confidence for error trails in a visual task (Allen et al., 2016;
Hauser et al., 2017).
Here, we hypothesized that both the valence and arousal of an encoded stimulus might
modulate the accuracy of memory itself, and investigated whether healthy individuals are
aware of such emotional effects on their recognition accuracy. To test this hypothesis, we
conducted a pre-registered experiment in which participants memorised lists of words grouped
by their levels of valence and arousal. Although most metamemory research has relied on
"feeling of knowing" self-report measures, these can be subject to substantive biases, i.e. such
as conflating self-report bias with metacognitive sensitivity or being confounded by overall
accuracy level (Fleming & Lau, 2014). To overcome these issues, we adapted a signal-theoretic
modelling approach to estimate recognition metamemory for emotional versus unemotional
words. If arousal primarily biases memory, for example by increasing the salience of encoded
items, we would expect to observe a positive main effect of item arousal on both accuracy and
metacognitive confidence. Conversely, if emotion primarily biased metacognition through a
valence-specific 'anchoring' effect, we would expect to observe a full interaction of stimulus
arousal and valence on both measures. As a third alternative, if metacognition were robust to
emotional biases, we would expect to observe the effects of stimulus valence and arousal on
accuracy and response speed, but not on confidence or metacognition. To complement these
analyses, we further recorded cardiac measures of physiological arousal through pulse
oximetry, to assess their mediating effect on the association between confidence and accuracy.
104

Materials and methods

Pre-registration and Open Materials

To improve our control of type-I and type-II error rates, as well as the overall reproducibility
of the study, the trial was pre-registered before any data collection using the standard Open
Science Foundation template. Detailed information regarding power analysis, sample size
considerations, experimental and trial design, planned analyses and other key points can be
found at the following URL: (https://osf.io/9awtb). In what follows, Confirmatory Analyses
and Results refer to planned analyses detailed in the pre-registration, whereas Exploratory
Analyses and Results refer to post-hoc exploratory analyses conducted following contact with
the data. Additionally, in the case of any minor deviation from the pre-registration, these are
documented on a case by case basis.

Participants

Thirty-five participants (26 females) between the ages of 18 and 26 (M = 21, SD = 1.9) were
recruited through local advertisements and took part in the experiment at Aarhus University
Hospital, Denmark. From the total sample of 35 participants, a sub-set of 30 participants passed
the pre-registered exclusion criteria and were analyzed further. All participants had normal or
corrected to normal vision, were fluent in English and provided informed written consent
before the experiment. The procedures were conducted following the Declaration of Helsinki
and with approval from the Danish Neuroscience Centre's (DNC) Institutional Review Board

(IRB). Participants received monetary compensation of 100 DKK per hour. The estimated total
duration of the test session was 1,5 hours (150 DKK). Participants also completed a post-test
stimulus validation measure in which they provided valence and arousal ratings for all stimuli
for an additional 50 DKK. All 35 participants completed the follow-up rating experiment.
128
Procedure
The experimental procedure included one laboratory and one at-home survey session on two
different days with one week in between. In the laboratory session, participants completed a
word recognition metamemory task designed to assess the effects of valence and arousal on
verbal recognition memory and metacognition. In the survey session, participants rated their
subjective feelings of valence and arousal evoked by the words used during the laboratory
session.
At the beginning of the laboratory session, participants were briefed on the nature of the
investigation, were provided task instructions and completed a brief training session of the
metamemory task. The training included an example learning phase of 50 neutral and
unarousing words, followed by an example testing phase of 10 trials with confidence ratings
(see Metamemory Task and Stimuli).
During the metamemory task, heart rate was monitored using a Nonin 3012LP Xpod USB
pulse   oximeter   together   with   a   Nonin   8000SM   'soft-clip'   fingertip   sensor
(https://www.nonin.com/) attached to the left index finger.

Word selection

Stimuli consisted of 1200 English words selected from the Affective Norms for English Words
(ANEW) database based on valence and arousal ratings measured among a population of
American students (Bradley & Lang, 1999). Although ANEW is not validated in the Danish
population, previous standardization of the database in Dutch, Spanish and Italian populations
(Montefinese et al., 2014; Moors et al., 2013; Redondo et al., 2007) showed good consistency
across both American and European samples. In order to estimate the reliability of these ratings
among our sample, we conducted an at-home valence and arousal rating task after the
behavioural test. We created 4 distinct subgroups of 300 word stimuli, according to a 2 by 2
factorial design, where the factors corresponded to valence (positive vs. negative) and arousal
(low vs. high). To this aim, we used the tertile of the valence and arousal distribution to exclude
words with intermediate ratings, whose valence and arousal might be ambiguous (see Fig. 2a).

Metamemory Task

Participants completed a word recognition metamemory task adapted from a previous study
(McCurdy et al., 2013) to test the influence of emotional valence and arousal on memory and
metacognition. The task included 12 blocks, each consisting of a learning phase (Fig. 1a) and
a testing phase (Fig. 1b). In the learning phase, participants viewed a list of 50 English words
for durations of 30, 60 or 90 seconds. The words were presented on the screen in the form of a
table containing five columns with ten rows of words each, and the participants were instructed
to memorize as many words as possible. The list of words in each learning phase corresponded
to a unique combination of the following factorial conditions: Valence (positive, negative),
Arousal (low, high). Participants were notified when 10 seconds of learning time was left by

the display of a small warning at the bottom of the screen. During the testing phase, participants
completed 50 trials designed to measure recognition memory and metamemory. On each trial,
two-word stimuli were presented to the left and right of a fixation cross. The word pair
consisted of a "target" and a "distractor", corresponding to words that were present or absent
in the previous learning phase, respectively. Target and distractor words were matched by
valence and arousal, and their position was randomized across trials. Participants were
instructed to press either the left or the right arrow key to indicate which of the two words they
recognized from the memorised list. This procedure corresponds to a two-alternative forced-
choice task (2AFC) design, which provides optimal conditions for estimating and comparing
metacognition scores across tasks (Lee et al., 2018). Following the button press, participants
provided a subjective confidence rating from 1 ("not confident at all/guessing") to 7 ("very
confident"). Both button presses and confidence ratings had a maximum time-limit of 3s. If
participants had slower responses, a brief message (i.e., "too slow!") was displayed on the
screen and the trial was marked as missed.
The blocks were presented in a pseudo-randomized order to ensure that high arousal blocks
were systematically interleaved with low-arousal blocks. The block order and the selection of
target vs. distraction lists were counterbalanced across participants.
183

A. Experimental design

B. Post-experimental survey

Figure 1: **A.** Experimental design. The metamemory task contained 12 experimental blocks, each consisting of a learning phase and a testing phase for the 50 words, in a factorial design separated by each Valence and Arousal condition. To limit habituation effects, block orders were counter-balanced in a pseudo-randomized order such that each high arousal block was interceded by a low arousal condition. **B.** Post-experimental survey. To validate our stimulus categories with respect to the original ANEW ratings, participants completed a short subject visual analogue scale rating of valence and arousal for the 1200 words used in the main task (600 target and 600 distractors) in an at-home experiment. This was done in a web-based version of the original procedure used in the original ANEW survey.

Valence and Arousal Rating Task

To validate the arousal and valence stimulus categories in our Danish sample, participants completed an at-home valence and arousal subjective rating task. They were instructed to provide valence and arousal ratings of their subjective experience associated with each word presented in the metamemory task. The ratings were collected using a 9-point visual numerical scale in a web-based version of the original ANEW survey protocol (Bradley & Lang, 1999). Our version was implemented using Pavlovia (https://pavlovia.org), an online platform for running PsychoPy experiments (Peirce et al., 2019). Each word was presented twice, once for valence and once for arousal, and the 9-point scales were complemented with pictures of the original drawings of the Self-Assessment Manikin (Bradley & Lang, 1994), as in the original ANEW survey (Bradley & Lang, 1999). Participants rated a total of 1200 words, self-pacing through all rating trials. We compared the ratings provided by the participants in this study with the normative ANEW ratings using a Spearman rank correlation test (see Fig. 2 c & d). After inspecting histograms of participant responses, we excluded one participant, who only ever pressed the same key (rating 5); this exclusion criterion was not noted in the pre-registered protocol. Overall, stimulus ratings in our sample corresponded very well to the original ANEW ratings, $\rho = [0.93-0.78]$, albeit with lower overall consistency for the arousal vs. valence dimension (Fig. 2).

Figure 2: Stimulus selection and rating validation. Two rating procedures were used to select and validate word stimuli; the original ANEW normed ratings and the PAVLOVIA at-home ratings completed by our participants in the post-experimental survey. Each word was rated on a 9-point scale (1-9) for valence and arousal separately. **A & B**. We selected the words used in the metamemory task by removing items from the central tertile in the arousal and valence rating distributions (panel A). The blue and red dots represent words with negative and positive valence, respectively. The light and dark points represent low and high arousal, respectively. The densities represent the distribution for positive and negative valence (red and blue), and arousal (light and dark). **C & D.** We compared the independent rating provided by the ANEW database to the actual ratings provided by the participants, PAVLOVIA, after the main procedure. Both ratings of valence and arousal showed reasonably high consistency, ρ = [0.93-0.78]. See online article for colour figures. The black dots represent each word in the datasets and the red line shows the identity line.

Signal Theoretic Metacognition Modelling

Here we applied a signal-theoretic computational model to describe participant behaviour on the metamemory task (Fleming, 2017; Maniscalco & Lau, 2012). This approach delineates overall behaviour into 'type-I' and 'type-II' measures, corresponding to a basic discrimination performance versus metacognitive levels of performance (Galvin et al., 2003). The type-I performance was quantified using reaction times (RTs) and the signal-detection theoretic (SDT) measures of d-prime (D') and criterion (c) (Macmillan & Creelman, 2004). Type-II

performance (i.e., metacognition) was assessed by the SDT measures of Meta-d' and Meta-
ratio (M-ratio) (Fleming & Lau, 2014). Briefly, this approach models metacognitive "hits"
(e.g., high/low confidence for correct/error trials, respectively) and "misses" (e.g., high/low
confidence for error/correct trials, respectively). All SDT-based measures were estimated at
the subject level (Fleming, 2017). This model has been extensively described and validated
previously (Fleming, 2017; Mazancieux et al., 2020; Morales et al., 2018), here we recount the
approach in the context of the present study to aid interpretation.
D-prime or d' is a measure of a participant's sensitivity to detect previously studied words
during the learning phase, independently from subjective response biases. Instead, criterion or
c' encodes the participant's response bias, that is, the overall tendency to prefer one response
over the other (e.g. if a participant chose the word presented to the left of the fixation point
more often than the alternative). Together with measures of reaction time, d' and c' are metrics
of "first-order" or "type-I" task performance. In contrast, Meta-d is an estimate of the
sensitivity of subjective confidence ratings to type-I performance (i.e., the probability to be
highly confident when correct, or uncertain when incorrect). Meta-d' is, therefore, a measure
of insight, or how well one can consciously discriminate their own type-I performance (Lau &
Rosenthal, 2011). However, metacognitive sensitivity is also a function of the overall
perceptual signal, and as such is substantively influenced by differences in d'. To control for
this effect, the 'M-ratio' (Meta-d' divided by d') is estimated as a measure of metacognitive
efficiency, denoting how a subject's metacognitive sensitivity over- or underperforms what can
be expected given their type-I sensitivity (Fleming & Lau, 2014). Finally, average confidence
on each condition denotes participants "meta-criterion" or "meta-c", or their overall level of
metacognitive bias denoting the tendency to be confident or uncertain irrespective of accuracy.
Meta d' and meta-c are metrics of "second-order" or "type-II" performance.
258

Confirmatory Analyses

Metamemory task

All data were pre-processed according to the protocols established in our pre-registration.
Accordingly, we excluded all trials with reaction times (RT) faster than 100 ms, greater than 3
standard deviations from the median RT, and missing data (absence of response or because the
response button was pressed too early or too late). Due to an unforeseen technical error, an
absence of response in some trials contaminated the following trial, resulting in negative
response times. These trials were also automatically rejected. This procedure resulted in the
exclusion of 3.49% (±3.68) of the trials. Finally, outliers in task performance for each of the
conditions were detected based on reaction time, d', and confidence distributions. We also
excluded participants showing any extreme value using Tukey's boxplots. Based on these
criteria, 5 participants were excluded from all behavioural analyses. These preprocessing steps
are also extensively described in the interactive Jupyter notebooks made available on the OSF
repository: https://osf.io/9awtb.
The preprocessing of the behavioural data was carried out using custom R scripts, using R
Studio (1.2.5019), the R software (R 3.6.1), and Python scripts using Python 3.7.6. The
Bayesian and frequentist statistical models were implemented using the JASP software

(https://jasp-stats.org/) version 0.12.2 and the R package (AFEX 0.27-2). All Type-1 and Type-
2 SDT measures (d', criterion, meta-d', m-ratio, and mc) were derived from the hierarchical
meta-cognition model (HMM) (Fleming, 2017) implemented in R
(https://github.com/metacoglab/HMeta-d), run on the individual level to enable frequentist
analysis of the resultant parameters.

Heart Rate Monitoring

We monitored instantaneous heart rate variability using a Nonin 3012LP Xpod USB pulse
oximeter together with a Nonin 8000SM 'soft-clip' fingertip sensor (https://www.nonin.com/).
Pulse oximeters indirectly measure peripheral blood oxygen saturation. The abrupt cyclic
increase of oxygenation reflects blood pulse following cardiac contraction. Here, we used the
pulse-to-pulse intervals to estimate the instantaneous heart rate. Oxygenation saturation level
was continuously recorded at a 75 Hz sampling rate. The preprocessing of the pulse oximetry
recording was carried out using Python scripts (Python version 3.7.6) and version 0.1.1 of the
Systole Python package (Legrand & Allen, 2020). Statistical analyses were carried out using
the Pingouin Python package (Vallat, 2018) and MNE Python (Gramfort, 2013). PPG signals
were first upsampled to 1000 Hz and clipping artefacts were corrected using spline
interpolation following recent recommendations (van Gent et al., 2019). The signal was then
squared for peak enhancement and normalized using the mean + standard deviation using a
rolling window (window size: 0.75 seconds). All positive peaks were labelled as systolic
(minimum distance: 0.2 seconds). We then detected ectopic, long, short, missed and extra beats
using adaptive thresholds over the successive beats-to-beats interval (Lipponen & Tarvainen,
2019), as implemented in Systole (Legrand & Allen, 2020). The code implementing these steps
can be found in the Jupyter notebooks made available on the OSF repository:
https://osf.io/9awtb.
Instantaneous Pulse Rate. All pulses labelled as missed or extra beats were corrected by
adding or removing beats, respectively. We then interpolated the instantaneous heart rate at 75
Hz to a continuous recording using the previous values and divided it into epochs (from -1
second pre-trial to 6 seconds after the word presentation). All the epochs that contained, or
were adjacent to, an interval that was labelled as long, short or (pseudo-)ectopic beats were
automatically rejected, resulting in an average rejection rate of 18.22% (±11.49%). The
instantaneous heart rate was then averaged across trials for each condition and downsampled
to 5 Hz for subsequent analyses.
Linear regression. In an exploratory analysis, we used the instantaneous pulse rate as a
predictor of confidence over time to track the relationship between cardiac frequency
modulation and metamemory. We extracted the data following the same procedure, this time
using 1s before the trial start as a baseline and using the initial sampling rate (75 Hz) to facilitate
cluster-based statistical tests. Cluster-based permutation testing was performed using the
permutation_cluster_test() and the *permutation_cluster_1samp_test()* functions from the MNE
Python package (Gramfort, 2013). This enabled us to assess significant point-to-point
deviations from zero in encoded responses while controlling for multiple comparisons.
Pulse Rate Variability. Besides the analysis of the instantaneous pulse rate, we also
performed pulse rate variability analyses. Although targeting a different physiological signal
as compared to a classic electrocardiogram (ECG), the varying length of pulse cycle provides

a sufficiently accurate estimation of the underlying heart rate variability (HRV) when used at
rest for healthy young participants (Schäfer & Vagedes, 2013). Here, we extracted the systolic
peak intervals using the method presented above. Intervals labelled as missed or extra beats
were corrected by adding or removing beats, respectively. Additionally, intervals that were
labelled as short, long, or (pseudo-)ectopic beats were corrected using linear interpolation.
Following our specification in the pre-registration, we reported heart rate variability metrics in
the time (RMSSD, pnn50) and frequency domain (normalized and non-normalized high and
low-frequency power), as well as non-linear indexes (SD1 and SD2). These indexes reflect
changes in beat-to-beat intervals and measure sympathetic and parasympathetic influences on
the heart (Shaffer et al., 2014; Shaffer & Ginsberg, 2017). We inspected the resulting time
series and rejected noisy and unreliable segments (2 segments were rejected in total). The
values of each metric were then averaged across learning time (30, 60 or 90 seconds), and the
summary variables were entered into a 2 by 2 repeated-measures ANOVA with factors stimulus
arousal (arousing vs. unarousing) and valence (positive vs. negative).

Results

Behavioural Results

Overview

Following our pre-registration, the behavioural analyses focused on two levels of performance
during the metamemory task: type-I variables corresponding to the discrimination ability, and
type-II variables describing metacognition. To assess memory performance, we analyzed
decision accuracy, discrimination sensitivity (d'), bias (c), and response time (RT). To assess
metacognition, we analyzed average confidence (i.e., metacognitive bias), metacognitive
sensitivity (Meta-d'), and metacognitive efficiency (M-ratio, Meta-d'/d'). All signal theoretic
measures (d', c, Meta-d', and M-ratio) were estimated using a unified Bayesian approach, as
described previously (Fleming, 2017). All posthoc tests were corrected for multiple
comparisons using the Holm procedure. Here, we reported only the key details of the significant
effects; full ANOVA tables and associated statistics for all analyses can be found in our JASP
notebooks located online at the following URL: https://osf.io/pefnr/.

Recognition Memory (Type-I)

First, we examined the influence of emotional Valence and Arousal on decision accuracy, in a
two-way repeated measures ANOVA, collapsing across block and learning time conditions.
We found a significant main effect of Valence ($F_{(1,29)} = 9.887$, $\eta_p^2 = 0.254$, $p = 0.004$), as well
as a significant interaction between Valence and Arousal ($F_{(1,29)} = 7.779$, $\eta_p^2 = 0.212$, $p = .009$),
as positive words were recognized more accurately than negative ones under low arousal
conditions ($T_{(29)} = 4.20$, $p_{Holm} < 0.001$). Similarly, for sensitivity (d'), we found a significant
effect of Valence ($F_{(1,29)} = 11.34$, $\eta_p^2 = 0.28$, $p = 0.002$), as negative words decreased d', as well
as an interaction between Valence and Arousal ($F_{(1,29)} = 7.34$, $\eta_p^2 = 0.20$, $p = 0.011$), as positive
words were recognized more sensitively than negative ones under low arousal conditions
($T_{(29)} = 4.304$, $p_{Holm} < 0.001$). When analyzing the median response time, we observed a
significant effect of Valence ($F_{(1,29)} = 0.55$, $\eta_p^2 = 0.16$, $p = .025$), an effect of Arousal ($F_{(1,29)} =$

6.94, $\eta_p2 = 0.19$, p = .013) and an interaction between these two factors ($F_{(1,29)} = 7.56$, $\eta_p2 = 0.20$, p = .010), revealing that participants responded faster to positive valence under the low compared to the high arousal condition ($T_{(29)}=3.80$, $p_{Holm} = .002$). Analysis of response criterion revealed no significant main effects or interactions, and no other significant effects were found (all ps > .05).

Metacognition (Type-II)

We then performed a second level of analysis on the metacognition data, comprising average confidence, meta-d and M-ratio. First, we performed a Valence × Arousal repeated measures ANOVA on the average confidence. This procedure showed a strong effect of Valence ($F_{(1,29)}$ = 14.98, $\eta_p2= 0.34$, p < 0.001), as participants were more confident for positive valenced words. No other effects or interactions were significant. Participants were also more sensitive to their performance (Meta-d') when responding to positive valenced words (main effect of Valence $F_{(1,29)} = 11.28$, $\eta_p2 = 0.28$, p = 0.002). Concerning the M-ratio (i.e., Meta-d'/d'), which measures metacognitive efficiency, we found no main effect or interactions (all ps > 0.05). Following our pre-registered protocol, we followed up this analysis with a Bayesian ANOVA (Rouder et al., 2012) implemented in JASP (version 0.12.2) (JASP Team, 2020), to assess the strength of evidence for the null effect. This analysis compares the evidence for nested models of increasing complexity; e.g., comparing a null model to those with only main effects of valence or arousal, or a full model with main effects and interaction terms. This revealed strong relative evidence for the null overall model (including subject offsets), BF_{Model} = 7.28; the next best model was one with a main effect of Valence whose relative BF_{Model} = 0.74, i.e. inconclusive evidence. This analysis suggests that under the default JASP priors, it is very unlikely that Valence, Arousal, or their interaction exerted any effect on metacognitive efficiency.

Fig. 3. Behavioural results showing factorial main effects and interactions on discrimination and metacognitive performance. Modified raincloud plots (Allen et al., 2019) illustrating behavioural results of discrimination measures of accuracy (**A**), criterion (**B**) and reaction time (**C**) as well as metacognitive measures of confidence (**D**), Meta-d' (**E**) and M-ratio (**F**). Repeated measures ANOVA (Valence × Arousal) was carried out for each

condition separately. The upper panel shows that a significant main effect of emotional valence was observed as negative valenced words reduced accuracy (**A**) and slowed down reaction times (**C**). Similarly, the lower panel shows a main effect of valence for both Confidence and Meta-d' are impaired by negative valence. (*** p<0.001, ** p<0.01, * p<0.05).

Encoding duration

The effects of the encoding phase duration on both type I and type II task performances were not included in the pre-registration and so were assessed in an exploratory post hoc analysis. The results are described in the Supplementary Materials.

Physiological results

Pulse rate variability

First, we analyzed the effect of valence and arousal on the heart rate frequency during the experimental blocks. We averaged the estimated beats per minute (Mean BPM) across the different learning times (30, 60 and 90 seconds) and submitted it to a two-way repeated measure ANOVA (Valence × Arousal). Results showed a main effect of Valence ($F_{(1,29)} = 10.852$, $\eta_p2 = 0.272$, $p = 0.003$), meaning lower BPM for negative valenced words, but no other main or interaction effects.

For the low and high-frequency peak analysis, the peak high-frequency (HF peak) revealed an interaction between valence and arousal ($F_{(1,29)} = 14.50$, $\eta_p2 = 0.33$, $p < 0.001$) such that high-frequency cardiac oscillations were suppressed by negative emotional valence under low but not high arousal ($T_{(29)} = 3.36$, $p_{Holm} = .007$).

Concerning the Root Mean Square of the Successive Differences (RMSSD, not shown in Fig. 4), we found a main effect of valence ($F_{(1,29)} = 6.74$, $\eta_p2 = 0.19$, $p = .015$) i.e. negative valence increased RMSSD, but no other main effect or interaction (all ps < .05).

When considering the proportion of successive beat-to-beat intervals deviating by more than 50 ms (pnn50) we observed an effect of Valence ($F_{(1,29)} = 24.17$, $\eta_p2 = 0.45$, $p < 0.001$), as well as an interaction between Valence and Arousal ($F_{(1,29)} = 4.54$, $\eta_p2 = 0.13$, $p = .042$). Under high arousal the positive valence suppressed pnn50 while negative valence increased it ($t_{(29)} = -4.98$, $p_{Holm} < 0.001$).

Finally, we also analyzed the effect of these factors on the non-linear metrics of heart rate variability SD1 and SD2. The SD2 metric revealed an effect of Valence ($F_{(1,29)} = 35.20$, $\eta_p2 = 0.55$, $p < 0.001$), so that negative valence increased SD2 heart rate variability, but we found no other main effects or interactions. Concerning SD1, we found no significant effects (all ps > .05). These results are illustrated in **Fig. 4**; here we reported the main significant effects, however full analyses details and results tables can be found in the HRV JASP notebook located on the Github repository for this study.

427
Fig. 4: Modified raincloud plots illustrating results of pulse rate variability (PRV) analyses. PRV indices were
calculated separately for each 50 trial block and averaged by condition. Mean BPM (**A**), Pnn50 (**B**), High-
frequency peak (**C**), SD2 (**D**). Repeated measures ANOVA (Valence × Arousal) was then carried out for each
variable separately. A significant main effect of emotional valence was observed for mean BPM, as negative
valence decreased cardiac activity frequency, as well as for the pnn50 and the non-linear SD2 metric. We did not
observe a main effect of Arousal, but an interaction with valence was found for the high-frequency peak, such that
high-frequency cardiac oscillations were reduced by negative emotional valence under low but not high arousal.
No other significant effects were found. (*** p<0.001, ** p<0.01, * p<0.05). See *Methods* and *PRV Results* for
more details.
437

Event-related analysis

Next, we analyzed the time-locked instantaneous pulse rate fluctuation following word
presentation. **Figure 5a** shows the evoked cardiac frequency fluctuation following the display
of the two words on the screen. Following the specification of the pre-registered report, we
analyzed the average of this fluctuation across time (**Figure 5b**). Here, we observed no effect
of Valence, ($F_{(1,28)} = 0.441$, $\eta_p2 = 0.015$, $p = 0.511$), Arousal ($F_{(1,28)} = 0.003$, $\eta_p2 < 0.001$, $p = 0.954$) or an interaction between these two factors ($F_{(1,28)} = 0.044$, $\eta_p2 = 0.001$, $p = .833$).

Linear regression

In an additional exploratory analysis, we also tested the possible interaction between the
instantaneous pulse rate modulation observed during decision and metacognition and the
subjective report provided by the participant. For each participant and condition separately, we
used the reported confidence C and the instantaneous pulse rate BPM at each time point s of
the trial t to fit a linear regression of the form:
451

452 $$BPM_{s,t} = \alpha_s + \beta_s \times C_t + \varepsilon$$

All variables were normalized and beta-values for each explanatory regressor and participant were extracted for statistical analysis. First, to test for a difference in beta values from 0 across time across all conditions, i.e. the average effect of trial by trial confidence reports on fluctuations in evoked heart-rate, replicating previous analysis linking these variables (Allen et al., 2016). We averaged between conditions assessed significance via non-parametric cluster-level t-test. Results show a significant cluster (3.94-5.65 seconds after stimulus presentation, p = 0.001). As our HRV and behavioural results emphasized an effect of stimulus valence on both metacognitive behaviour and cardiac activity, we then compared the association between confidence and instantaneous cardiac activity between different valence and arousal conditions. When comparing positive and negative valence conditions (averaging across arousal levels) we found a significant early cluster (1.20-2.96 seconds after stimulus presentation, p=0.047), suggesting that stimulus valence modulates the correlation between evoked heart-rate and confidence. Finally, we repeated this comparison for high vs. low arousal conditions, collapsing stimulus valence. This analysis found no significant clusters. See Fig. 5 for illustration of these results.

Fig. 5: **Modulation of the cardiac activity at the trial level and its relation with reported subjective confidence. A.** Evoked pulse rate activity shows that the overall experimental procedure modulated the instantaneous cardiac frequency over time through an early acceleration component after the trial start (0-2 s) and a later deceleration component (2-6 s). This pattern is consistent with an orientation reflex, suggesting early brief integration and later sensory or memory processing. Interestingly, these two components are also time-locked with the decision and metacognition average response time. Here, we did not observe any difference between the

experimental conditions. **B.** We averaged the instantaneous pulse rate in the window of interest (0-5 s) and confirmed this absence of effect and an overall diminution of cardiac frequency after the trial start. **C**. Beta values over time of the linear regression (Confidence ~ BPM) for positive and negative valence trials separately. The confidence level was associated with the instantaneous cardiac frequency during the late time window corresponding to the metacognition decision. **D.** Beta values over time of the linear regression (Confidence ~ BPM) for high and low arousal trials separately. Using the same approach, contrasting for High and Low level of arousal. Significance assessed using a cluster-level statistical permutation test (alpha=0.05). Shaded areas and error bar show the 68% CI. Significant clusters are shown by a shaded red path for condition contrast, and grey path for null tests. See online article for colour figures.

Discussion

In this study, we investigated the influence of emotion on word memory and metamemory through a combination of experimental psychology, cognitive modelling, and physiology. We adapted a recognition memory paradigm (McCurdy et al., 2013) where participants memorised lists of words varying in arousal and valence. Stimulus valence exerted a consistent influence on recognition performance, metacognitive confidence, and physiological activity. For reaction time and accuracy, this effect was greatest for low vs. high arousal words, suggesting that the influence of stimulus valence on memory depends in part on arousal. On the physiological level, we observed an association between the subjective confidence reported by the participant and the evoked pulse rate, which was also modulated by the word valence. Our results evidence that although recognition memory is impaired for negative emotional stimuli, participants can accurately monitor and report this uncertainty. Further, monitoring the effect of emotion on memory may depend in part on integrating the associated changes in cardiac signals.

The notion that emotional events are better recalled than neutral ones has been extensively discussed (Yonelinas & Ritchey, 2015), but the psychophysiological mechanisms mediating this effect are still unclear. In some cases, emotional influences on memory have been assumed to be almost entirely ascribed to arousal, which is presumed to control the cognitive and behavioural importance of stimuli (Mather, 2007; Mather & Sutherland, 2009), while other works attributed this improvement primarily to valence (Adelman & Estes, 2013; Kensinger, 2009). Previous investigations in the perceptual domain have documented that arousing stimuli "boost" the signal-to-noise ratio of visual motion, as reflected in both models of ballistic evidence accumulation, and subjective confidence (Allen et al., 2016; Lufityanto et al., 2016). However, in our study, the effect of arousal was generally muted or dependent on stimulus valence.

Whereas on type-I task performance, stimulus valence interacted with arousal, for metacognitive type-II variables we observed only a pronounced main effect of valence with no arousal effect or interaction. In general, participant confidence reports closely matched the overall effect of stimulus emotion on performance; negative valence decreased sensitivity, increased reaction times, and decreased confidence. The robust evidence we observed for the absence of effect on metacognitive-efficiency (M-ratio) further underlines this finding; the strong null Bayes factor here demonstrates that shifts in subjective confidence were well reflected by the magnitude of any changes in type-I sensitivity, indicating that subjects make optimal use of the available memory signal during metacognitive judgements, irrespective of any conditional valence or an arousal effect. This finding suggests that, although memory is

degraded under negative emotional contexts, participants can accurately account for this in their subjective confidence.

One possible explanation for the absence of arousal effect is found in our validation rating study (see Figure 2); while the valence dimension was well preserved between the ANEW database and the ratings by our participants after the task, the consistency of arousal ratings was slightly reduced. This limits the extent to which we can infer actual stimulus arousal in our data, and it may be that the stimuli were simply not sufficiently distinct or triggering for a Danish sample. Indeed, in this study arousing versus nonarousing word stimuli did not evoke a significant difference in physiological arousal response. Future studies could benefit from both a larger corpus of validated words, a more general sample of English-speaking participants, and multiple modalities of memorised stimuli which may better preserve arousal-based dimensions. Here, it should also be noted that the use of words instead of images or film is also a potential confounding factor due to the lack of vividness and complexity of the mental imagery elicited by one single item. While previous works evidenced that words classified as highly arousing and with extreme positive or negative valence are associated with better memory recall (Buchanan et al., 2006; Madan et al., 2017, 2019), other dimensions like semantic properties or the functional use of the object such as animacy are more relevant psycholinguistic dimensions to predict free recall (Madan, 2020). Again, controlling for this dimension in future studies could help to refine the influence of emotion on metamemory beyond the dimensions of valence and arousal.

In a related line of research, several investigations have linked physiological activity (e.g., as indexed by pupil dilation or cardiac acceleration) to subjective confidence and metacognition. According to influential predictive-coding accounts of metacognition (Allen et al., 2016; Meyniel et al., 2015; Moulin & Souchay, 2015), confidence reflects the width of a posterior decision variable, such that fluctuations in arousal bias the gain or precision of this distribution. Here, we examined both trial level evoked changes in cardiovascular activity and summary measures of pulse rate variability separately for each condition. When examining instantaneous heart rate variation, we observed a robust sinusoidal pattern that remained stable across conditions, similar to an orientation reflex triggered by trial onset (see Figure 5). The shape and intensity of this cardiac deceleration can reflect several cognitive processes like attention orienting, emotion processing or inhibitory control (Abercrombie et al., 2008; Critchley et al., 2005; Legrand et al., 2020). Replicating previous findings (Allen et al., 2016), we observed a robust association between trial-by-trial fluctuations in subjective confidence during this late interval, with the strength of this association being modulated by stimulus valence during the early, decision-evoked period. These results suggest that at least some variance in the monitoring of emotional inputs on metamemory could arise from monitoring the associated physiological changes. This result corroborates the notion that memory retrieval is an embodied process (Garfinkel et al., 2013), which has implications both for their conservation, the accuracy of their recall, but also their control in the case of distressing emotional memories (Gagnepain et al., 2017; Legrand et al., 2020).

Whereas no overall modulation of instantaneous heart-rate was seen for stimulus valence or arousal, here we observed substantive, robust modulations of heart rate variability (HRV) when subjects recalled negatively valenced stimuli across multiple time, frequency and non-linear indices. HRV (i.e the amount of change across time of the interbeat intervals) can reflect the influence of higher cognitive processes on cardiac frequency through the parasympathetic

nervous system (Smith et al., 2017; Thayer & Lane, 2009). Across the different range of HRV
indices we examined, two showed a strong valence main effect (i.e., Mean BPM & SD2),
whereas others (i.e., high-frequency peak and pnn50) showed a robust interaction between
these factors. Although disentangling what underlies these different effects is far from trivial,
it is interesting to note the dissociation between these effects, and similarity to those observed
for our type-I and type-II metamemory measures. One intriguing possibility is that the high-
frequency variability indexed by the former two measures may be a more direct input for
metacognitive monitoring than the others, as these showed a similar pattern of exaggerated
valence effect with no effect of arousal. One means to probe this hypothesis is to correlate
individual differences in the modulation of confidence by valence with each HRV metric;
however, our study is underpowered for individual differences analyses (Schönbrodt &
Perugini, 2013), leaving it as an intriguing avenue for future research.
Several important limitations should be considered when interpreting our HRV effects. As
HRV is here calculated by collapsing across each 50 trial block, the modulations observed
therein are necessarily a mixture of multiple cognitive states and perceptual inputs; future
studies could benefit from disentangling the encoding, perceptual, and retrieval stages to better
account for these stages of the decision process. Additionally, here we assessed heart rate
variability through pulse oximetry recording. Pulse oximetry recordings are used as an
alternative to the electrocardiogram (ECG) by several clinical and non-clinical studies
(Quintana, Elstad, et al., 2016). The sampling rate of our device (75 Hz) is not optimal when
compared to recommended standards for electrocardiogram (ECG) recording and HRV
measurement (Quintana, Alvares, et al., 2016), which could limit our ability to detect true
effects, particularly in the lower frequency range. Previous reports, however, have shown a
strong consistency between the estimated pulse rate variability and the heart rate variability as
measured through ECG (Lu et al., 2009; Schäfer & Vagedes, 2013). Similarly, we did not
measure or control respiratory cycles during this study, which robustly modulate HRV
measures, in particular in the lower frequency. Collectively, while our results nicely
demonstrate that stimulus emotional content modulates high-frequency indices of
cardiovascular arousal, future studies in this area are likely to benefit from a combination of
more nuanced experimental design and a more sophisticated recording set-up.
594

Conclusion

This pre-registered study sheds light on the biasing effects of valence on metamemory with
possible physiological correlates of these effects. Negatively valenced stimuli globally
decreased both memory performance and metacognition, supporting a role for emotions in
guiding confidence and memory performance. While arousal has often been described as a
possible mechanism of the beneficial effect of emotion on memory, we only found limited
evidence for performance improvement under highly arousing conditions. In line with these
main effects, we found that stimulus valence modulated the overall pulse rate variability and
the association between instantaneous heart-rate and subjective confidence. Collectively, these
results suggest that although negative stimuli do exert a degrading influence on recognition
memory, participants are largely able to account for this effect in their subjective confidence,
perhaps by monitoring physiological states.

Acknowledgements
NL, CMCC, NKM, NN, and MA are supported by a Lundbeckfonden Fellowship (under Grant
[R272-2017-4345]), and the AIAS-COFUND II fellowship programme that is supported by the
Marie Skłodowska-Curie actions under the European Union's Horizon 2020 (under Grant
[754513]), and the Aarhus University Research Foundation.

ORCIDS:

N. L. : 0000-0002-2651-4929

S. S. E. : 0000-0002-5348-6417

C. M. C. C. : 0000-0001-7663-5828

N. K. M. : 0000-0001-6580-5289

N. N. : 0000-0003-1226-1139

F. F. : 0000-0002-9974-6261

M. A. : 0000-0001-9399-4179

Disclosure statement. No financial interest or benefit that has arisen from the direct
applications of this research.

Data availability statement.

The project pre-registration can be found at the following link: https://osf.io/9awtb

Data deposition and supplemental online material.

All behavioural and physiological data can be found at: https://osf.io/pefnr/

634   **References**

Abercrombie, H., Chambers, A., Greischar, L., & Monticelli, R. (2008). Orienting, emotion,

and memory: Phasic and tonic variation in heart rate predicts memory for emotional

pictures in men. *Neurobiol Learn Mem*, *90*, 644–650.

Adelman, J. S., & Estes, Z. (2013). Emotion and memory: A recognition advantage for positive

and negative words independent of arousal. *Cognition*, *129*(3), 530–535.

https://doi.org/10.1016/j.cognition.2013.08.014

Allen, M., Frank, D., Schwarzkopf, D. S., Fardo, F., Winston, J. S., Hauser, T. U., & Rees, G.

(2016). Unexpected arousal modulates the influence of sensory noise on confidence.

ELife, *5*. https://doi.org/10.7554/elife.18103

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019). Raincloud plots:

A multi-platform tool for robust data visualization. *Wellcome Open Research*, *4*, 63.

https://doi.org/10.12688/wellcomeopenres.15191.1

Anderson, A. K. (2005). Affective Influences on the Attentional Dynamics Supporting

Awareness. *Journal of Experimental Psychology: General*, *134*(2), 258–281.

https://doi.org/10.1037/0096-3445.134.2.258

Baird, B., Smallwood, J., Gorgolewski, K. J., & Margulies, D. S. (2013). Medial and Lateral

Networks in Anterior Prefrontal Cortex Support Metacognitive Ability for Memory and

Perception. *Journal of Neuroscience*, *33*(42), 16657–16665.

https://doi.org/10.1523/JNEUROSCI.0786-13.2013

Bradley, M. M., & Lang, P. J. (1994). Measuring emotion: The self-assessment manikin and

the semantic differential. *Journal of Behavior Therapy and Experimental Psychiatry*,

25(1), 49–59. https://doi.org/10.1016/0005-7916(94)90063-9

Bradley, M. M., & Lang, P. J. (1999). *Affective Norms for English Words (ANEW): Instruction*

Manual and Affective Ratings.

Buchanan, T. W., Etzel, J. A., Adolphs, R., & Tranel, D. (2006). The influence of autonomic

arousal and semantic relatedness on memory for emotional words. *International*

Journal   of   Psychophysiology,   *61*(1),   26–33.

https://doi.org/10.1016/j.ijpsycho.2005.10.022

Cahill, L., & McGaugh, J. L. (1998). Mechanisms of emotional arousal and lasting declarative

memory. *Trends in Neurosciences*, *21*(7), 294–299. https://doi.org/10.1016/S0166-

2236(97)01214-9

Chua, E. F., Pergolizzi, D., & Weintraub, R. R. (2014). The Cognitive Neuroscience of

Metamemory Monitoring: Understanding Metamemory Processes, Subjective Levels

Expressed, and Metacognitive Accuracy. In S. M. Fleming & C. D. Frith (Eds.), *The*

Cognitive   Neuroscience   of   Metacognition   (pp.   267–291).   Springer.

https://doi.org/10.1007/978-3-642-45190-4_12

Critchley, H. D., Rotshtein, P., Nagai, Y., O\textquotesingleDoherty, J., Mathias, C. J., &

Dolan, R. J. (2005). Activity in the human brain predicting differential heart rate

responses   to   emotional   facial   expressions.   *NeuroImage*,   *24*(3),   751–762.

https://doi.org/10.1016/j.neuroimage.2004.10.013

Davis, R. L., & Zhong, Y. (2017). The Biology of Forgetting—A Perspective. *Neuron*, *95*(3),

490–503. https://doi.org/10.1016/j.neuron.2017.05.039

De Martino, B., Fleming, S. M., Garrett, N., & Dolan, R. J. (2013). Confidence in value-based

choice. *Nature Neuroscience*, *16*(1), 105–110. https://doi.org/10.1038/nn.3279

Fleming, S. M. (2017). HMeta-d: Hierarchical Bayesian estimation of metacognitive efficiency

from   confidence   ratings.   *Neuroscience   of   Consciousness*,   *2017*(1).

https://doi.org/10.1093/nc/nix007

Fleming, S. M., Dolan, R. J., & Frith, C. D. (2012). Metacognition: Computation, biology and

function. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

367(1594), 1280–1286. https://doi.org/10.1098/rstb.2012.0021

Fleming, S. M., & Lau, H. C. (2014). How to measure metacognition. *Frontiers in Human*

Neuroscience, *8*. https://doi.org/10.3389/fnhum.2014.00443

Fleming, S. M., Maniscalco, B., Ko, Y., Amendi, N., Ro, T., & Lau, H. (2015). Action-Specific

Disruption of Perceptual Confidence. *Psychological Science*, *26*(1), 89–98.

https://doi.org/10.1177/0956797614557697

Fleming, S. M., Ryu, J., Golfinos, J. G., & Blackmon, K. E. (2014). Domain-specific

impairment in metacognitive accuracy following anterior prefrontal lesions. *Brain*,

137(10), 2811–2822. https://doi.org/10.1093/brain/awu221

Gagnepain, P., Hulbert, J., & Anderson, M. C. (2017). Parallel Regulation of Memory and

Emotion Supports the Suppression of Intrusive Memories. *The Journal of*

Neuroscience, *37*(27), 6423–6441. https://doi.org/10.1523/JNEUROSCI.2732-16.2017

Galvin, S. J., Podd, J. V., Drga, V., & Whitmore, J. (2003). Type 2 tasks in the theory of signal

detectability: Discrimination between correct and incorrect decisions. *Psychonomic*

Bulletin & Review, *10*(4), 843–876. https://doi.org/10.3758/BF03196546

Garfinkel, S. N., Barrett, A. B., Minati, L., Dolan, R. J., Seth, A. K., & Critchley, H. D. (2013).

What the heart forgets: Cardiac timing influences memory for words and is modulated

by metacognition and interoceptive sensitivity: Cardiac timing and memory.

Psychophysiology, *50*(6), 505–512. https://doi.org/10.1111/psyp.12039

Gramfort, A. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in*

Neuroscience, *7*. https://doi.org/10.3389/fnins.2013.00267

Hauser, T. U., Allen, M., Purg, N., Moutoussis, M., Rees, G., & Dolan, R. J. (2017).

Noradrenaline blockade specifically enhances metacognitive performance. *ELife*, *6*,

e24901. https://doi.org/10.7554/eLife.24901

Heyes, C., Bang, D., Shea, N., Frith, C. D., & Fleming, S. M. (2020). Knowing Ourselves

Together: The Cultural Origins of Metacognition. *Trends in Cognitive Sciences*, *24*(5),

349–362. https://doi.org/10.1016/j.tics.2020.02.007

James,   W.   (1884).   What   is   an   emotion?   *Mind*,   *9*(34),   188–205.

https://doi.org/10.1093/mind/os-ix.34.188

JASP Team. (2020). *JASP (Version 0.12.2)[Computer software]*. https://jasp-stats.org/

Kensinger, E. A. (2009). Remembering the Details: Effects of Emotion. *Emotion Review*, *1*(2),

99–113. https://doi.org/10.1177/1754073908100432

Kensinger, E. A., & Corkin, S. (2004). Two routes to emotional memory: Distinct neural

processes for valence and arousal. *Proceedings of the National Academy of Sciences*,

101(9), 3310–3315. https://doi.org/10.1073/pnas.0306408101

Kever, A., Grynberg, D., Eeckhout, C., Mermillod, M., Fantini, C., & Vermeulen, N. (2015).

The body language: The spontaneous influence of congruent bodily arousal on the

awareness of emotional words. *Journal of Experimental Psychology: Human*

Perception and Performance, *41*(3), 582–589. https://doi.org/10.1037/xhp0000055

Kever, A., Grynberg, D., Szmalec, A., Smalle, E., & Vermeulen, N. (2019). "Passion" versus

"patience": The effects of valence and arousal on constructive word recognition.

Cognition   and   Emotion,   *33*(6),   1302–1309.

https://doi.org/10.1080/02699931.2018.1561419

Kever, A., Grynberg, D., & Vermeulen, N. (2017). Congruent bodily arousal promotes the

constructive recognition of emotional words. *Consciousness and Cognition*, *53*, 81–88.

https://doi.org/10.1016/j.concog.2017.06.007

Kreibig, S. D. (2010). Autonomic nervous system activity in emotion: A review. *Biological*

Psychology, *84*(3), 394–421. https://doi.org/10.1016/j.biopsycho.2010.03.010

Lau, H., & Rosenthal, D. (2011). Empirical support for higher-order theories of conscious

awareness.   *Trends   in   Cognitive   Sciences*,   *15*(8),   365–373.

https://doi.org/10.1016/j.tics.2011.05.009

Lee, A. L. F., Ruby, E., Giles, N., & Lau, H. (2018). Cross-Domain Association in
Metacognitive Efficiency Depends on First-Order Task Types. *Frontiers in*
Psychology, *9*, 2464. https://doi.org/10.3389/fpsyg.2018.02464

Legrand, N., & Allen, M. (2020). *Systole: V 0.1.2—September 2020* (0.1.2) [Computer
software]. Zenodo. https://doi.org/10.5281/ZENODO.3607912

Legrand, N., Etard, O., Vandevelde, A., Pierre, M., Viader, F., Clochon, P., Doidy, F.,
Peschanski, D., Eustache, F., & Gagnepain, P. (2020). Long-term modulation of cardiac
activity induced by inhibitory control over emotional memories. *Scientific Reports*,
10(1), 15008. https://doi.org/10.1038/s41598-020-71858-2

Lipponen, J. A., & Tarvainen, M. P. (2019). A robust algorithm for heart rate variability time
series artefact correction using novel beat classification. *Journal of Medical*
Engineering & Technology, *43*(3), 173–181.
https://doi.org/10.1080/03091902.2019.1640306

Lu, G., Yang, F., Taylor, J. A., & Stein, J. F. (2009). A comparison of photoplethysmography
and ECG recording to analyse heart rate variability in healthy subjects. *Journal of*
Medical Engineering & Technology, *33*(8), 634–641.
https://doi.org/10.3109/03091900903150998

Lufityanto, G., Donkin, C., & Pearson, J. (2016). Measuring Intuition: Nonconscious
Emotional Information Boosts Decision Accuracy and Confidence. *Psychological*
Science, *27*(5), 622–634. https://doi.org/10.1177/0956797616629403

Macmillan, N. A., & Creelman, C. D. (2004). *Detection Theory: A User's Guide* (2nd ed.).
Psychology Press. https://doi.org/10.4324/9781410611147

Madan, C. R. (2020). Exploring word memorability: How well do different word properties
explain item free-recall probability? *Psychonomic Bulletin & Review*.
https://doi.org/10.3758/s13423-020-01820-w

Madan, C. R., Scott, S. M. E., & Kensinger, E. A. (2019). Positive emotion enhances

association-memory. *Emotion*, *19*(4), 733–740. https://doi.org/10.1037/emo0000465

Madan, C. R., Shafer, A. T., Chan, M., & Singhal, A. (2017). Shock and awe: Distinct effects

of taboo words on lexical decision and free recall. *Quarterly Journal of Experimental*

Psychology (2006), *70*(4), 793–810. https://doi.org/10.1080/17470218.2016.1167925

Maniscalco, B., & Lau, H. (2012). A signal detection theoretic approach for estimating

metacognitive sensitivity from confidence ratings. *Consciousness and Cognition*,

21(1), 422–430. https://doi.org/10.1016/j.concog.2011.09.021

Mather, M. (2007). Emotional Arousal and Memory Binding: An Object-Based Framework.

Perspectives on Psychological Science, *2*(1), 33–52. https://doi.org/10.1111/j.1745-

6916.2007.00028.x

Mather, M., & Sutherland, M. (2009). Disentangling the Effects of Arousal and Valence on

Memory for Intrinsic Details. *Emotion Review*, *1*(2), 118–119.

https://doi.org/10.1177/1754073908100435

Mazancieux, A., Fleming, S. M., Souchay, C., & Moulin, C. J. A. (2020). Is there a G factor

for metacognition? Correlations in retrospective metacognitive sensitivity across tasks.

Journal of Experimental Psychology: General. https://doi.org/10.1037/xge0000746

McCurdy, L. Y., Maniscalco, B., Metcalfe, J., Liu, K. Y., de Lange, F. P., & Lau, H. (2013).

Anatomical Coupling between Distinct Metacognitive Systems for Memory and Visual

Perception. *Journal of Neuroscience*, *33*(5), 1897–1906.

https://doi.org/10.1523/JNEUROSCI.1890-12.2013

Meyniel, F., Sigman, M., & Mainen, Z. F. (2015). Confidence as Bayesian Probability: From

Neural Origins to Behavior. *Neuron*, *88*(1), 78–92.

https://doi.org/10.1016/j.neuron.2015.09.039

Montefinese, M., Ambrosini, E., Fairfield, B., & Mammarella, N. (2014). The adaptation of

the Affective Norms for English Words (ANEW) for Italian. *Behavior Research*

Methods, *46*(3), 887–903. https://doi.org/10.3758/s13428-013-0405-3

Moors, A., De Houwer, J., Hermans, D., Wanmaker, S., van Schie, K., Van Harmelen, A.-L.,

De Schryver, M., De Winne, J., & Brysbaert, M. (2013). Norms of valence, arousal,

dominance, and age of acquisition for 4,300 Dutch words. *Behavior Research Methods*,

45(1), 169–177. https://doi.org/10.3758/s13428-012-0243-8

Morales, J., Lau, H., & Fleming, S. M. (2018). Domain-General and Domain-Specific Patterns

of Activity Supporting Metacognition in Human Prefrontal Cortex. *The Journal of*

Neuroscience, *38*(14), 3534–3546. https://doi.org/10.1523/JNEUROSCI.2360-17.2018

Moulin, C., & Souchay, C. (2015). An active inference and epistemic value view of

metacognition. *Cognitive Neuroscience*, *6*(4), 221–222.

https://doi.org/10.1080/17588928.2015.1051015

Ochsner, K. N. (2000). Are affective events richly recollected or simply familiar? The

experience and process of recognizing feelings past. *Journal of Experimental*

Psychology: General, *129*(2), 242–261. https://doi.org/10.1037/0096-3445.129.2.242

Otgaar, H., Howe, M. L., Patihis, L., Merckelbach, H., Lynn, S. J., Lilienfeld, S. O., & Loftus,

E. F. (2019). The Return of the Repressed: The Persistent and Problematic Claims of

Long-Forgotten Trauma. *Perspectives on Psychological Science*, *14*(6), 1072–1095.

https://doi.org/10.1177/1745691619862306

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E.,

& Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior*

Research Methods, *51*(1), 195–203. https://doi.org/10.3758/s13428-018-01193-y

Quintana, D. S., Alvares, G. A., & Heathers, J. A. J. (2016). Guidelines for Reporting Articles

on Psychiatry and Heart rate variability (GRAPH): Recommendations to advance

research communication. *Translational Psychiatry*, *6*(5), e803–e803. https://doi.org/10.1038/tp.2016.73

Quintana, D. S., Elstad, M., Kaufmann, T., Brandt, C. L., Haatveit, B., Haram, M., Nerhus, M., Westlye, L. T., & Andreassen, O. A. (2016). Resting-state high-frequency heart rate variability is related to respiratory frequency in individuals with severe mental illness but not healthy controls. *Scientific Reports*, *6*(1), 37212. https://doi.org/10.1038/srep37212

Redondo, J., Fraga, I., Padrón, I., & Comesaña, M. (2007). The Spanish adaptation of ANEW (Affective Norms for English Words). *Behavior Research Methods*, *39*(3), 600–605. https://doi.org/10.3758/BF03193031

Reggev, N., Zuckerman, M., & Maril, A. (2011). Are all judgments created equal?: An fMRI study of semantic and episodic metamemory predictions. *Neuropsychologia*, *49*(5), 1332–1342. https://doi.org/10.1016/j.neuropsychologia.2011.01.013

Rollwage, M., Loosen, A., Hauser, T. U., Moran, R., Dolan, R. J., & Fleming, S. M. (2020). Confidence drives a neural confirmation bias. *Nature Communications*, *11*(1), 2634. https://doi.org/10.1038/s41467-020-16278-6

Rouder, J. N., Morey, R. D., Speckman, P. L., & Province, J. M. (2012). Default Bayes factors for ANOVA designs. *Journal of Mathematical Psychology*, *56*(5), 356–374. https://doi.org/10.1016/j.jmp.2012.08.001

Schäfer, A., & Vagedes, J. (2013). How accurate is pulse rate variability as an estimate of heart rate variability? *International Journal of Cardiology*, *166*(1), 15–29. https://doi.org/10.1016/j.ijcard.2012.03.119

Schönbrodt, F. D., & Perugini, M. (2013). At what sample size do correlations stabilize? *Journal of Research in Personality*, *47*(5), 609–612. https://doi.org/10.1016/j.jrp.2013.05.009

Shaffer, F., & Ginsberg, J. P. (2017). An Overview of Heart Rate Variability Metrics and

Norms. *Frontiers in Public Health*, *5*, 258. https://doi.org/10.3389/fpubh.2017.00258

Shaffer, F., McCraty, R., & Zerr, C. L. (2014). A healthy heart is not a metronome: An

integrative review of the heart's anatomy and heart rate variability. *Frontiers in*

Psychology, *5*. https://doi.org/10.3389/fpsyg.2014.01040

Shea, N., Boldt, A., Bang, D., Yeung, N., Heyes, C., & Frith, C. D. (2014). Supra-personal

cognitive control and metacognition. *Trends in Cognitive Sciences*, *18*(4), 186–193.

https://doi.org/10.1016/j.tics.2014.01.006

Shields, G. S., Sazma, M. A., McCullough, A. M., & Yonelinas, A. P. (2017). The effects of

acute stress on episodic memory: A meta-analysis and integrative review.

Psychological Bulletin, *143*(6), 636–675. https://doi.org/10.1037/bul0000100

Smith, R., Thayer, J. F., Khalsa, S. S., & Lane, R. D. (2017). The hierarchical basis of

neurovisceral integration. *Neuroscience & Biobehavioral Reviews*, *75*, 274–296.

https://doi.org/10.1016/j.neubiorev.2017.02.003

Tauber, S. K., & Dunlosky, J. (2012). Can older adults accurately judge their learning of

emotional information? *Psychology and Aging*, *27*(4), 924–933.

https://doi.org/10.1037/a0028447

Thayer, J. F., & Lane, R. D. (2009). Claude Bernard and the heart-brain connection: Further

elaboration of a model of neurovisceral integration. *Neuroscience and Biobehavioral*

Reviews, *33*(2), 81–88. https://doi.org/10.1016/j.neubiorev.2008.08.004

Valenza, G., Citi, L., Lanatá, A., Scilingo, E. P., & Barbieri, R. (2014). Revealing Real-Time

Emotional Responses: A Personalized Assessment based on Heartbeat Dynamics.

Scientific Reports, *4*(1). https://doi.org/10.1038/srep04998

Vallat, R. (2018). Pingouin: Statistics in Python. *Journal of Open Source Software*, *3*(31),

1026. https://doi.org/10.21105/joss.01026

van Gent, P., Farah, H., van Nes, N., & van Arem, B. (2019). Analysing Noisy Driver

Physiology Real-Time Using Off-the-Shelf Sensors: Heart Rate Analysis Software

from the Taking the Fast Lane Project. *Journal of Open Research Software*, *7*, 32.

https://doi.org/10.5334/jors.241

Ye, Q., Zou, F., Lau, H., Hu, Y., & Kwok, S. C. (2018). Causal Evidence for Mnemonic

Metacognition in Human Precuneus. *The Journal of Neuroscience*, *38*(28), 6379–6387.

https://doi.org/10.1523/JNEUROSCI.0660-18.2018

Yeung, N., & Summerfield, C. (2012). Metacognition in human decision-making: Confidence

and error monitoring. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, *367*(1594), 1310–1321. https://doi.org/10.1098/rstb.2011.0416

Yonelinas, A. P., & Ritchey, M. (2015). The slow forgetting of emotional episodic memories:

An emotional binding account. *Trends in Cognitive Sciences*, *19*(5), 259–267.

https://doi.org/10.1016/j.tics.2015.02.009