Comparative metabolic profile of maize genotypes reveals the tolerance mechanism associated under combined stresses.

Suphia Rafique[-1]

Department of Biotechnology, Faculty of Chemical and Life Sciences, Jamia Hamdard, New Delhi-110062. India

[-1] Corresponding Author email: suphia123@gmail.com

Abstract:

Drought, heat, high temperature, waterlogging, low-N stress, and salinity are the major environmental constraints that limit plant productivity. In tropical regions maize grown during the summer rainy season, and often faces irregular rains patterns, which causes drought, and waterlogging simultaneously along with low-N stress and thus affects crop growth and development. The two maize genotypes CML49 and CML100 were subjected to combination of abiotic stresses concurrently (drought and low-N / waterlogging and low-N). Metabolic profiling of leaf and roots of two genotypes was completed using GC-MS technique. The aim of study to reveal the differential response of metabolites in two maize genotypes under combination of stresses and to understand the tolerance mechanism. The results of un-targeted metabolites analysis show, the accumulated metabolites of tolerant genotype (CML49) in response to combined abiotic stresses were related to defense, antioxidants, signaling and some metabolites indirectly involved in nitrogen restoration of the maize plant. Alternatively, some metabolites of sensitive genotype (CML100) were regulated in response to defense, while other metabolites were involved in membrane disruption and also as the signaling antagonist. Therefore, the present study provides insight into the molecular mechanisms of tolerance under various stresses of maize plants, that governed on the regulation of cell wall remodeling, maintain metabolic homeostasis, defense against the pathogen, proper signaling, and restore growth under stress conditions.

Keywords: Combine stresses; GCMS; Metabolomics; drought; low-N stress; waterlogging stress; and Tolerant. Vs Sensitive

Abbrevations:

GC-MS- Gas chromatography mass spectrometry

Low-N- Low nitrogen stress

DAS- Days after sowing

PCA-Principal component analysis

.

1. Introduction

Crop faces various stresses in their life span including abiotic and biotic. Several abiotic stresses that affect crop growth, and development are drought, waterlogging, low nutrient availability, salinity, high temperature and heat. The predicted changes in the global climate has accentuate in the frequency and intensity of abiotic factors, most of the abiotic stresses occur in combination, for instant, in tropical area's terminal drought and waterlogging occurs simultaneously during summer-rainy season. Although on many occasions the stresses are for shorter duration and relatively mild in nature, but have a significant impact on plant growth and development. Therefore, co-occurrence of abiotic stresses are more destructive to crops than their separate occurrence at different crop growth stages (**Mittler 2006; Prasad et al 2011**). Besides, the composite responses of plants due to various concurrent stresses rely on signals, these varied and contrast signals may interact with each other to enhanced or obstruct one another (**Vile et al. 2012; Suzuki et al. 2014**). Maize is the most important cereal, grown at the wide geographical ranges of latitude and longitude. In South Asia, particularly in tropical and subtropical environment maize is largely grown during summer-rainy season, in marginal areas which often faces extreme water availability such as waterlogging or water scarcity in a form of terminal drought. Correspondingly, co-occurrence of both the stresses often headed to the depletion in nitrogen content, thus causing low-N stress in the soil. However, studies on combined stresses by **Rizhsky et al (2004)** revealed that the response of plant to combined stresses is unique and cannot be extrapolated from the response of plant to each individual stress. Though, plants belonging to the same genus showed different molecular response in combined stresses (**Aprile et al 2013**). Therefore, the nature of the interaction is an important aspect to fully identify the influences of combined abiotic stresses on crop plants. Among omics approach is metabolomics that has been neglected in crop improvement programs (**Kumar et al 2017**). It has not be fully explored due to its complexity. Though, metabolites plays important role in plant metabolism that influences the plant biomass and architecture (**Turner et al 2016**). Metabolomics study enable to identify wide variety of metabolites that gives the comprehensive view of the cellular process and physiological condition that display in a particular stress condition. Besides, the metabolites assign the functional gene and impact of the particular gene on the metabolic pathway, and that provides information of regulation as well interruptions of linked pathways (**Wen et al 2015**). Further, metabolomics study enable to improve breeding resources through selecting the remarkable traits (**Zivy et al 2015**). Therefore, the aim of metabolomics profiling is to compare the combined stress response in two maize inbred that differed in their stress adaptability and to unrevealed the complex molecular mechanism of tolerance at metabolomics levels in different stress conditions applied simultaneously.

2. Materials and Methods

Plant Material and Growth conditions

In the present work two maize inbred (CML49 & CML100) were selected that showed different adaptability to various combine stress treatments on this basis we have categorized them as 'tolerant and sensitive.' The selected two inbred (CML49 and CML100) plants with 60 pots each were grown in natural conditions in greenhouse up to 30 days. 30 DAS they were subjected to various stress treatment, the plants were grown in low nitrogen, only 25% of the recommended dose of urea was applied to the treated pots. First 30 pots of each inbred were exposed to a drought stress for 10 days, while 30 control pots were supplied with full nutrient and water. After re-watering for two days normally, the same plants were exposed to waterlogging stress for up to 7 days, to maintain the water level 2-3cm above the soil surface of the pots plants were watered day and night. The roots samples from stressed and control replicates plants of each inbred (CML49 & CML100) were kept on the last day of combined stresses for extracting metabolites. Samples were quickly frozen in liquid nitrogen after removing from the plant and then kept at -80^0 C for further analysis.

Sample Preparations and Metabolites extraction

The leaves and roots samples from three independent biological replicates each from control and treated maize plants (CML49 & CML100) was used for metabolite profiling. Harvested plant samples immediately frozen in liquid Nitrogen in a plastic bag (resistant to liquid N_2). Frozen samples were stored at -80^0C. The leaves and roots were homogenized separately with liquid Nitrogen in a mortar and pestle into fine powder. One gram powder of each sample was extracted three times with 5 ml of methanol at room temperature and sonicated for 15 min and centrifuged. The extract was concentrated to 1 ml by Speed Vac. The supernatant filter through Whatman filter paper (No. 4), extracts collected and stored at 4 ^{0}C for further use. The prepared extracts of both leaves and roots of the control and treated plants were subjected to gas chromatography–mass spectroscopy (GC–MS) analysis. Samples were analyzed using GC-MS for the analysis of untargeted metabolites. For derivatization, methoxylamine in pyridine (10μl) and *N*-methyl-*N*-trimethylsilyltrifuoroacetamide MSTFA (90μl) were added.

Gas Chromatography and Mass spectrometry

The GC-MS analysis was performed with a GC-MS QP-2010 ULTRA equipped with an auto sampler (AOC-20i + s) from Shimadzu (Japan), using Equity-5 column, 30.0 m × 0.25 um × 0.25 mm for separation, and helium was used as a carrier gas at a constant flow rate of 1.0 mL/min. The 1μL sample was injected using 1/100 split-mode injections at a temperature of 260°C. The oven temperature program was initially set at 100°C and held for 2 min. The temperature was gradually increased to 250°C at a rate of 10°C/min, and 300°C at 15°C/min. Ions were generated by a 70ev electron beam at an ionization current of 2.0mA. Total ion chromatogram spectra were recorded in the mass range of 40–900 m/z at the rate of 2.5 spectra s^{-1}. For metabolite identification and annotation, peaks were matched against customized reference spectrum databases including the National Institute of Standards and Technology NIST and the Wiley Registry, internal libraries and

3. Results

Comparison of metabolite levels of two inbred in combined abiotic stress condition

Metabolomics profiling using GC-MS was conducted of tolerant (CML49) and sensitive (CML100)
genotypes under various combined stresses. Total 132 compounds were identified, that includes,
fatty acids, (long, medium and short chains), fatty acid derivatives, fatty aldehydes, fatty alcohols,
carboxylic acids, terpenoids, glucosinolates, heterocyclic compounds, halohydrogens, alkaloids,
phenolic acids, steroids and vitamins. Univariate analysis of two genotypes (CML49 and CML100)
was performed by fold change analysis, t-test and volcano plots. Fold change showed the
comparison between the control group metabolites with that of treated group metabolites and
important features selected by t-test with threshold ($p<0.05$) level, the results was plotted in log 2
scale. Thus, same fold change have the same distance from zero base line **(Table 1).**

Table 1. Metabolic profile of leaf and roots of tolerant and sensitive maize plants after exposure to combined abiotic stresses.

TOLERANT GENOTYPE (CML49)			SENSITIVE GENOTYPE (CML100)		
Metabolites	Class	Fold Change	Metabolites	Class	Fold Change
Linalool	Monoterpenoid	-3.161*	Carvon	Monoterpenoid	3.885**
Erythrodiol	Triterpenoid	-3.155*	Squalene	Steroid	-2.019**
Tocopherol	Steroid	-2.959**	Heptyl hexyl ether	Ether lipid	-1.999**
			Tocopherol Acetate	Steroid	-1.87**
Squalene	Steroid	2.820**	1-Hexadecen-3-ol	Acyclic diterpenoid	1.004*
Oleic acid	Saturated FA	-2.693**	Steraldehyde	Fatty aldehyde	1.235*
Undecen	Alkene	-2.508*	Citronellyl acetate	F alcohol ester	-1.404**
Octanoic acid	Saturated FA	-2.487*	1-Iodononane	Liner alkane	-1.394*
3-Decen-3-one	Fatty Alcohol	-2.384**	Docosane	Straight chain alkane	-1.189*
Root Metabolites			**Roots Metabolites**		
Furan carboxylic	Heterocyclic acid	-2.590**	NAKIJIQUINONE B	Hydroquinone lipids	-1.423*
			Furan-2-carboxylic acid	Carboxylic acid	-1.408*
2-Bromopentane	Halohydrocarbo	-2.416**	Caryophyllene	Sesquiterpenoid	-1.254*

The tolerant and (CML49) and sensitive (CML100) genotypes were exposed to drought,

waterlogging and low-N stresses. Strikingly, in both genotypes an overlap of antioxidant compounds

like α-tocopherol, squalene, β-sitosterol and isophytol was spotted in leaves, but fold change was

higher of tolerant genotype **(Fig 1).** However, sensitive genotype leaves shows higher accumulation

of saturated and unsaturated fatty acids whereas, mostly, fatty alcohols, fatty anhydride and higher

alkanes level was presented in greater amount **(Fig 2)**.

Fig 1. Leaf metabolic profile of CML49 and CML100 after exposure to combined abiotic stresses

Fig 1 shows the percent accumulation and overlapping of anti-oxidative and various other metabolites in leaves of both genotypes.

135
136
137

Fig 2. Roots metabolic profile of CML49 and CML100 after exposure to combined abiotic stresses.

Fig 2. shows the percent accumulation and overlapping of different metabolites in roots of both genotypes

Hence, on analyzing the metabolomics profile of two different genotypes suggested that the response of maize plants to various stresses applied simultaneously was shared as well as also distinctive. The roots metabolite profile of tolerant genotype display large variance in accumulated metabolites most of them are halohydrocarbons, heterocyclic compounds, organic cations and saturated and unsaturated fatty acids, carboxylic acids, esters compared to roots of sensitive genotype which have less diversity in metabolite. The aim of principal component analysis (PCA) methods was to reduce multivariate data complexity. In this method, patterns were identified, and data expressed in ways that highlight similarities and differences between samples. In our study, PCA was applied to metabolites concentration data of control and treated leaves and roots of tolerant and in sensitive plants. Hence, to verify the difference between the metabolic profiles of the control and treated tissues statistically and to identify the main metabolites responsible for the differences. A score scatter plots of tolerant leaf showed the good separation of metabolites between treated and control plants. In PC1 the variance is greater (70.1%) and PC2 showed variance of (19.3%). Similar trend of leaf metabolite separation was observed of sensitive genotype. The treated plants exhibited higher variances in PC1 (91.2%) and control plants variance is less PC2 (5.2%) **Fig 3a, b.**

Fig 3a. Score plot of tolerant leaf

156
157
158

Fig 3 b. Score plot of Sensitive leaf

However in roots the trend was opposite of both genotypes, the score plot for PC1 (94.8%) and PC2

(4%) demonstrate clear separation between control root metabolites as compared to treated root

metabolites it may be presence of many saturated and unsaturated fatty acids and halo-hydrocarbon

in the samples. Similarly, sensitive (CML100) genotype PC1 (94%) and PC2 (5.3%) **Fig 4a, b**,

control root metabolites were well separated and treated roots metabolites patterns have less

variance in separation because presence of similar metabolites like, 2-Furan carboxylic acid and

Furan anhydride. Comparative metabolomics analyses of two combined stresses response have

highlighted a number of metabolites involved in diverse metabolic pathways. The Pathway analysis

of the tolerant leaf and roots showed biosynthesis of unsaturated fatty acids, fatty acid, Steroid,

Sulfur metabolism, fatty acid elongation in mitochondria, Selenoamino acid metabolism, Pyruvate

metabolism, Glycolysis and gluconeogenesis metabolism. However, the pathway impact is highest

of sulfur metabolism.

Fig 4a. Score plot of tolerant roots

175
176

Fig 4 b. score plot of sensitive roots

Correlation studies of metabolites and Hierarchical clustering

The leaf and roots metabolites of tolerant and sensitive genotype shows contrasting pattern.

However, the leaf metabolites of the tolerant genotypes differed with root metabolites. Although,

most of the metabolites tolerant genotype of leaf have higher concentration, and some metabolites

of control and treated leaf have medium and low concentrations, on the other hand roots metabolites

have high and low concentrations only though the metabolites are correlated **Fig 5a, b.** Whereas,

leaf and roots metabolites of sensitive genotype categorized into low and high concentrations **Fig**

6 a, b

Fig 5a. Correlation study of tolerant leaf (CML49).

190
191
192

Fig 5 b.Correlation study of tolerant (CML49 Roots.

193
194

Fig 6a. Correlation analysis CML100L

195
196

Fig 6b. Correlation analysis roots CML100

Hierarchical cluster analysis of tolerant leaf metabolites shows two major class control and treated. Some metabolites (Tocopherol acetate, 9-Decen-3-one, Oleic acid, Linalool, Erythidol, 1-Undecen, Squalene, Octanoic acid) have higher concentration in treated plants. Whereas, (stigmasterol, 2-Dodecyloxirane, Lignoceric acid, 2,6-Octadien, gamma Dodeca;actron, Linolelaidic acid, 1-Hexadecen-3-ol, 1,16 dichlorohexadecane, 1-Pentanol) control leaf metabolites were cluster together. Some metabolites like, N-Pentadecane, Palmitic acid, 2-Ethylnonane, Trimethyl 2,2,4 hexane, 1-Chlorononane and Dodecane, Linolenic acid shows overlapping. Therefore, **Fig 7** provides a change in metabolite profiles in response to various stress treatment and categorized in various clusters.

Fig 7. Heat map of the metabolites detected in roots of tolerant (CML49) genotype samples. A color coded matrix represents the concentration of the metabolites in the three replicates of each plant. Prior to statistical analysis, the values have been log2 transformed and normalized

210   In contrast, tolerant roots metabolites also categorized into two class control and treated. The treated

211   group contains 19 metabolites in response to various abiotic stresses (combined) and were higher

212   concentration. On the other hand, 23 metabolites with low concentration. Whereas, in control group

213   the opposite trend was observed **Fig 8.**

Fig 8. Heat map of the metabolites detected in leaf of tolerant (CML49) genotype samples. A color coded matrix represents the concentration of the metabolites in the three replicates of each plant. Subsequently before statistical analysis, the values have been log2 transformed and normalized

Most of the metabolites in both the class (control and treated) were belongs to single cluster. The sensitive genotype leaves and roots metabolites also have two class (control and treated) only 12 metabolites of leaf (treated group) shows higher concentration and 26 metabolites with low concentration. These two group metabolites shows three large clusters and other small sub clusters **Fig 9**. In contrast, the treated group contains 15 metabolites with high concentration range (0.5 to 1.5) and 15 metabolites with low concentration range (-0.5 to -1.5) **Fig 10** Most of the metabolite of treated and control group belongs to single large cluster. The variation in metabolites was less observed under various stresses in sensitive genotype.

Fig 9. Heat map of the metabolites detected in leaves of sensitive (CML100) genotype samples. A color coded matrix represents the concentration of the metabolites in the three replicates of each plant. Subsequently before statistical analysis, the values have been log2 transformed and normalized

224
225

Fig 10. Heat map of the metabolites detected in roots of sensitive (CML100) genotype samples. A color coded matrix represents the concentration of the metabolites in the three replicates of each plant. Subsequently before statistical analysis, the values have been log2 transformed and normalized

Data analysis

The GC-MS data was analyzed using MetaboAnalyst (www.metaboanalyst.ca) web server. Peak areas of identified metabolites in triplicate in a form of comma separated excel file (.CVS format) were uploaded in this software. The software, check the data integrity of the uploaded files automatically. MetaboAnalyst replaces zero values with a small positive value (the half of the minimum positive number detected in the data). Data filtering, and data normalization was done by pooling controlled groups. The internal data structure was transformed to a table with each row represents a tissue sample (root, leaf) from tolerant genotype and sensitive genotypes and each column representing a feature (a metabolite). The data structure in this format, we performed a row-wise data normalization. The purpose of Row-wise normalization was to normalize each sample (row) so that they are comparable to each other. In Metaboanalyst, correlation analysis shows the correlation between different features. Further, hierarchical clustering analysis was done with the squared Euclidean distance and the average linkage method (between groups).The clustering result in the form of a heatmap. The values of the metabolites mean abundances were used to cluster the metabolomic data.

4. Discussion

Role of tolerant genotype metabolites in multiple stresses

In the present work untargeted metabolites were quantified by GCMS technique and differentially regulated metabolites of tolerant (CML49) and sensitive (CML100) were compared with their respective controls. The major observation was differences in metabolite levels in treated versus untreated plants of both genotype, though less metabolites differed in treated leaves and roots of sensitive genotype comparative tolerant one respectively, these variations caused by simultaneous exposure of genotypes to various stresses and it depend on plants response also. The leaves metabolites of tolerant genotype shows fold increase in terpenoids linalool (monoterpen) and erythrodiol (triterpein) the main role of isoprenoid family have been associated in protection against oxidative and other abiotic stresses, especially volatile isoprenoids (Monoterpen) protective in photosynthetic processes during thermal and oxidative stresses (**Claudia Vikers et al. 2009**). Its role in membrane stabilization (**Sharkey and Singsaas 1995**) a hypothesis supports terpenes behave directly as an antioxidant, scavenges ROS by reactions through the conjugated double bond system (**Loretu and Velikova 2001; Affeck and Yakir 2002**). Similarly, Erythrodiol is a pentacyclic triterpenes, leaf surface contains triterpenoids as constituents of waxes being involved in different roles such as maintenance of leaves structure, provide water, permeability, and plant insect interactions (**Bauer et al 2004; Mintz-Oron et al 2008; Wilson et al 2008**). Correspondingly, non-volatile isoprenoid like Tocopherol acts as lipid soluble redox buffer and important scavenger of singlet oxygen species (**Foyer et al 2005**). Also function as an antioxidants, in transgenic tobacco leaves its deficiency led to an enhancement in the lipid peroxidation (**Abbasi et al 2009**). Majority of the metabolites accumulated in roots of tolerant genotype were belongs to saturated and unsaturated fatty acids, FAs and their derivatives directly involved in providing resistance against the bacterial (C16) and fungal pathogen (18:2 and 18:3) hence responsible for plants basal immunity **Kachroo and Kachroo (2009)**. Moreover, indirect and direct role of 18:0, 18:1, 18:2, and 18:3 plant defense signaling. In parsley, fungal infection rapidly induces the transcription of the 18:3-synthesizing omega-3 FA (**Kirsch et al 1997**). Significant role of Oleic acid as a stimulator of the signaling enzyme phospholipase D (PLD) that function in anti-death cell (**Zhang et al. 2003**). However, free fatty acids not only regulate abiotic stress tolerance, but also involve defenses against wound-induced responses and insect/herbivore feedings (**Upchurch 2008**). In our studies, most of the metabolites of tolerant were involved in Phosphotidiyal and sphingolipid metabolism. Further work of **Yozo Okazaki and Kazuki Saito (2014)** confirms that membrane lipids has specific signal mediator and sensor of various biotic and abiotic stresses. Hence activating defense reactions at transcriptional level. As the lipids are stress mitigators, presence of polyunsaturated fatty acids that reduced cell damage expose to stress by scavenging reactive oxygen species. Therefore reducing the intensity of environmental stresses (**Mene-Saffrane et al 2009**). Also, there are reports on

membrane lipid remodeling in response to various stresses like, freezing, dehydration and nutrition depletion **(Moellering et al 2010; Gasulla et al 2013; Okazaki et al 2013b)**, and this remodeling contributes to the uphold membrane integrity and membrane- bound protein functions. While, Benzoic acid related to salicylic acid, (phenolic phytohormone) involve as an endogenous signal mediating plant defense responses against pathogens. Besides, it influence responses to abiotic stresses and related to aspects of plant growth and development **(Vlot et al 2009; Rivas-San Vicente and Plasencia 2011)**. The presence of haloalkanes (2-Bromopentene, ethyl isocynates) heterocyclic compound (2-Nonyl-1,3-dioxalane), hetrocyclic aromatic compounds (N-Methyl-2-Iodopyrrol), Methyl ester (Thioacetic acid) and Methyl ketone (3-Nitropropane) all these compound processes strong, unpleasant and odor may involve in repelling herbivory insects, pest and bacteria, fungi, nematodes **(Lamberth et al 2012; Francis et al 2013).** The role of low molecular weight heterocyclic compounds derivatives of pyrimidine, pyrazole and isoflavones as a stimulators of seed germination and growth of maize seedlings **(Tsygankova et al 2016).** Likewise, N-methyl-Δ-[1]Pyrrolinium is a cation combined with nicotinic acid to forms the alkaloid (S)-nicotine in roots of tobacco plant **(Katoh et al. 2005)** that inhibits a wide variety of herbivores **(Rosenthal G A 1986)**. *Trichoderma harzianum*, used as biocontrol agent, it produced Nonanoic acid (NA) that was very inhibitory to spore germination and mycelial growth of two cacao pathogens, Crinipellis perniciosa Stahel and Moniliophthora roreri Cif **(Aneja et al 2005).** The antibiotic effect of acetic acid on some bacteria, including *Salmonella typhimurium*, *Escherichia coli* O157, and *Listeria monocytogenes* **(Delaquis et al 1999).** The soluble oxalic acid is a strong acid, besides being toxic and lethal to grazing animals **(Von Burg 1994)**. It showed inhibitory effects on sucking insects' planthopper and aphids **(Yoshihara et al 1980; Massonie 1980).** Therefore we concluded that most of the roots metabolites were involved in providing stability to membranes and plant defense.

Role of sensitive genotypes metabolites in multiple stresses

In sensitive leaf and root shows the accumulation of various metabolites, belongs to different classes, like terpenoids, Steroids, Ether lipids, Hydroquinone lipids, fatty aldehyde, fatty alcohol ester, fatty acyl conjugates, cyclic anhydrides, carboxylate, and linear alkanes. The leaves shows the emission of main terpenoids limnonen-oxide (carvon) and citronell. They are allelopathic in nature thus restricting seed germination and growth of nearby plants **(Maffei and Gertsch 2011).** Moreover, plants emitted volatiles, also perceive or recognize them in inter- and intra-plant communications, so that plants under herbivore attack can alert neighbouring plant species, and priming their chemical defenses **(Heil 2014).** The below ground emission of volatiles is an efficient trait: in maize roots the sesquiterpene (E)-β-caryophyllene is necessary to attract entomopathogenic nematodes to roots damaged by the ferocious maize pest Diabrotica virgifera. Maize varieties that lack this signal have been shown to be far more vulnerable to maize pest **(Rasmann et al 2005; Rasmann and Turlings 2007).** However, there were some reports, that indicate due to lipophilic

nature of monoterpenes the physical arrangement of phospholipids in the membranes were distorted and thus affect membrane functioning **(Zunino and Zygadlo 2004)**. Further, **Kaur et al (2010)** showed that citronellol generates ROS that induced lipid peroxidation, leakage of membrane and altered the membrane structure and permeability in wheat. Markedly as reported **Takshak and Agarwal (2015)** in UV-B treated leaf of *W somnifera* accumulated higher amount of phyto-constituents like Isophytol, trans-squalene, nonacosane and these plants were effective in scavenging free radicals. Furan fatty acids and their role in plants have been reviewed in detail **(Mowlong et al 2014)**. F-acids are mainly bound to phospholipids, they substitute for PUFAs and their strong capability (F-acids) to serve as radical scavengers suggests that plants use them to defend against oxidative stress **(Spiteller 2005).** Moreover, the insecticidal activity of some 2-carboxylbenzofurans and their coumarin precursors against sweet potato weevil, *Cylas formicarius elegantulus* first demonstrated by **Williams et al (1994)** and later insecticidal properties of Benzofuran 2-carboxylic acid derivatives was reported **Jackson et al (1998).** Water deficit led to produced various compounds in plants **Carmo-Silva et al (2009)** observe that in C4 grasses the amount of 5-HNV(2-amino-5- hydroxypentanoic acid (5-hydroxynorvaline, 5-HNV) in leaves of, C. dactylon and Z. japonica, increased under water deficit condition. However, any benefits from this unusual non-protein amino acid for drought resistance should be further explored to see that if it plays an important role in stress defensive mechanisms.

5 Conclusion

The metabolomics study of leaf and roots of tolerant plants were mostly involved in the biosynthesis of fatty acid metabolism, biosynthesis of saturating and unsaturated fatty acids, sulfur metabolism, thereby, regulated signaling, re-fix soil nitrate metabolism and defense against the wide variety of pathogens. Therefore, enhancing the tolerant mechanism of the genotype. In contrast, sensitive plant metabolites were mostly involved in leaking or disrupting the membranes that disrupt signaling between various organelles and also for various stresses, defense against the particular pathogens. In present roots metabolomics study we can say, the metabolites somehow help to survive the sensitive genotype in multiple stress conditions but most of them are the causes of susceptibility. On the basis of the above observation, we can conclude that tolerant genotype plants by remodeling of the cell wall, maintain metabolic homeostasis, and proper signaling thus able to tolerate multiple stress conditions. Therefore, the study provides a comprehensive analysis of tolerance mechanism.

"The authors have declared no conflict of interest".

Acknowledgments/ Grant received: I acknowledge the grant received from DST Ministry of Science, Government of India, under the WOS-A scheme.

Reference No: SR/WOS-A/LS-284/2013

REFERNCES:

Abbasi A. R, Saur A, Hennig P, Tschiersch H, Hajirezaei M, Hofius D, et al. (2009). Tocopherol deficiency in transgenic tobacco (*Nicotianatabacum* L.) plants leads to accelerated senescence. Plant Cell Environ 2009; 32: 144–157.

Affek, H.P. & Yakir, D. Protection by isoprene against singlet oxygen in leaves. Plant Physiol 2002; 129: 269–277.

Ai T N, Naing A H, Yun B-W, Lim S H and Kim C K. Overexpression of RsMYB1 Enhances Anthocyanin Accumulation and Heavy Metal Stress Tolerance in Transgenic Petunia. Front. Plant Sci 2018; 9: 1388.

Alfonso Albacete, Elena Cantero-Navarro, Dominik K, Großkinsky Cintia L, Arias María Encarnación, Balibrea Roque, Bru Lena Fragner, Michel E Ghanem, María de la Cruz, González Jose A, Hernández Cristina, MartínezAndújar Eric, van der Graaff ,Wolfram Weckwerth, Günther Zellnig, Francisco Pérez-Alfocea, and Thomas Roitsch. Ectopic overexpression of the cell wall invertase gene CIN1 leads to dehydration avoidance in tomato. J Exp Botany 2015; 66 (3): 863–878

Alvarez, C., Garcia, I., Moreno, I., Perez-Perez, M.E., Crespo, J.L., Romero, L.C., and Gotor, C. Cysteine-generated sulfide in the cytosol negatively regulates autophagy and modulates the transcriptional profile in *Arabidopsis*. Plant Cell 2012b; 24: 4621–4634.

Aprile, A., Havlickova, L., Panna, R., Marè, C., Borrelli, G. M., Marone, D., et al. Different stress responsive strategies to drought and heat in two durum wheat cultivars with contrasting water use efficiency.BMC Genomics 2013; 14:821.

Bae M S, Cho E J, Choi E Y, and Park O K. Analysis of the Arabidopsis nuclear proteome and its response to cold stress. Plant Journal 2003; 36: 652–663.

Barbrook AC, Howe CJ, Purton S. Why are plastid genomes retained in non-photosynthetic organisms? Trends in Plant Science 2006; 11: 101–108.

Bauer, S., Schulte, E., and Thier, H.-P. (2004). Composition of the surface wax from tomatoes I.
Identification of the components by GC/MS. European Food Research and Technology 2004; 219:

Bevan M. et al 1998. Analysis of 1.9 Mb of contiguous sequence from chromosome 4 of *Arabidopsis*
thaliana. Nature 1998; 391: 485-488.

Bermúdez, M.A., Galmés, J., Moreno, I., Mullineaux, P.M., Gotor, C., and Romero, L.C.
Photosynthetic adaptation to length of day is dependent on S-sulfocysteine synthase activity in the
thylakoid. Plant Physiol 2012; 160, 274–288.

Bradford M M. A rapid and sensitive method for the quantitation of microgram quantities of protein
utilizing the principle of protein-dye binding. Anal Biochemistry 1976; 721 (2): 248-254.

Bui, Q. T., and Grandbastien, M. -A. (2012)."LT Rretrotransposons as controlling elements of
genome response to stress? ,"in *Plant Transposable Elements: Impact on Genome Structure and*
Function Topics in Current Genetics, eds M.-A. Grandbastien and J.M .Casacuberta
(Berlin;Heidelberg:Springer),273–296.

Cantoni G L. Biological methylation: selected aspects. Annu Rev Biochem 1975; 44: 435-451.

Ana E. Carmo-Silva, Alfred J. Keys, Michael H. Beale, Jane L. Ward, John M. Baker, Nathaniel D.
Hawkins, Maria Celeste Arrabaça, Martin A.J. Parry. Drought stress increases the production of 5-
hydroxynorvaline in two C4 grasses. Phytochemistry 2009; 70: 664–671.

Cecilia Gotor, Irene García, José L. Crespo & Luis C. Romero. Sulfide as a signaling molecule in
autophagy, Autophagy 2013; 9: (4), 609-611.

Delaquis PJ, Sholberg PL, Stanich K (1999) Disinfection of mung bean seed with gaseous acetic
acid. J Food Prot 1999; 62: 953–957.

Deserah D. Strand, Nicholas Fisher, and David M. Kramer. The higher plant plastid NAD (P)H
dehydrogenase-like complex (NDH) is a high efficiency proton pump that increases ATP production
by cyclic electron flow. J Biol Chem 2017; 292(28): 11850–11860.

Down's M T. An improved hydrazine reduction method for the automated determination of low
nitrate levels in freshwater. Water Res 1978; 12: 673-675.

Dubald M, Barakate A, Mandaron P, Mache R. The ubiquitous presence of exopolygalacturonase
in maize suggests a fundamental cellular function for this enzyme. Plant J 1993; 4: 781–791.

Ebbs S, J Bushey, S Poston, D Kosma, M Samiotakis, D Dzombak. Transport and metabolism of
free cyanide and iron cyanide complexes by willow. Plant Cell Environ 2003; 26: 1467–1478.

Essemine J, Qu M, Mi H and Zhu X-G. Response of Chloroplast NAD (P) H Dehydrogenase-
Mediated Cyclic Electron Flow to a Shortage or Lack in Ferredoxin-Quinone Oxidoreductase-
Dependent Pathway in Rice Following Short-Term Heat Stress. Front. Plant Sci 2016; 7:383.

Evans H.J., and Nason, A. Pyridine nucleotide-nitrate reductase from extracts of higher plants. Plant
Physiol 1953; 28: 233–254. doi:10.1104/pp.28.2.233.

Erice, G., S. Louahlia, J.J. Irigoyen, M. Sanchez-Diaz, J.C. Avice. 2010. Biomass partitioning,
morphology and water status of four alfalfa genotypes submitted to progressive drought and
subsequent recovery. Journal of Plant Physiology 167(2):114-120.

Foyer C H, and Noctor G. (2005) Oxidant and antioxidant signaling in plants: a re-evaluation of the
concept of oxidative stress in a physiological context. Plant Cell Environ 2005; 29: 1056–1107.

Garcia, I., Castellano, J.M., Vioque, B., Solano, R., Gotor, C., and Romero, L.C. Mitochondrial
{beta}-cyanoalanine synthase is essential for root hair formation in *Arabidopsis thaliana*. Plant Cell
2010; 22: 3268–3279.

GarcõÂa-Romera I, Fry S C. Upper limits for endogenous oligogalacturonides and free galacturonic
acids in rose cell-suspension cultures: Implications for the action of exo- and endo-
polygalacturonases in vivo. J Plant Physiol 1997; 150: 241±246.

Grandbastein M A. Activation of plant retrotransposons under stress conditions. Trends in Plant
Sciences 1998; 3: 5 181-187.

Marie-Ang`ele Grandbastien, H´el`ene Lucas, Jean-Benoˆıt More, Corinne Mhiri, Samantha
Vernhettes & Josep M. Casacuberta . The expression of the tobacco Tnt1 retrotransposon is linked
to plant defense responses. Genetica 1997; 100: 241–252.

Gotor, C., and Romero, L.C. S-sulfocysteine synthase function in sensing chloroplast redox status.
Plant Signal. Behav 2013; 8: e23313.
467
468   Grover H L, Nair T V R, and Abrol Y P. Nitrogen metabolism of the upper three leaf blades of
469   wheat at different soil Nitrogen levels. Physiol Plant 1978; 42: 287-292.
470
471   Guillas, I., Guellim, A., Reze, N. and Baudouin, E. Long chain base changes triggered by a short
472   exposure of Arabidopsis to low temperature are altered by AHb1 non-symbiotic haemoglobin
473   overexpression. Plant Physiol Biochem 2013; 63: 191–195.
474
475   Guo W, Zhao J, Li X, Qin L, Yan X, Liao H. A soybean b expansin gene GmEXPB2 intrinsically
476   involved in root system architecture responses to abiotic stresses. Plant J 2011; 66(3):541–552.
477
478   Hageman, R.H., and Hucklesby, D.P. Nitrate reductase from higher plants. Meth. Enzymol 1971;
479   23: 491–503.
480
481   Hammani K, and Barkan A. An mTERF domain protein functions in group II intron splicing in
482   maize chloroplasts. Nucleic Acids Research 2014; 42 (8): 5033–5042.
483
484   Han Y, Chen Y, Yin S, Zhang M, Wang W. Over-expression of TaEXPB23, a wheat expansin gene,
485   improves oxidative stress tolerance in transgenic tobacco plants. J Plant Physiol 2015; 173:62–71.
486
487   Heil M. Herbivore-induced plant volatiles: targets, perception and unanswered question. New
488   Phytologist 2014; 204: 297–306.
489
490   Hu, Xiuli; Liuji Wu, Feiyun Zhao, Dayong Zhang, NanaLi1 Guohui, Zhu Chaohao Li and Wei
491   Wang. Phosphoproteomic analysis of the response of maize leaves to drought heat and their
492   combination stress. Front Plant Sci 2015; 6: 1-21.
493
494   Iftekhar Alam, Dong-Gi Lee, Kyung-Hee Kim, Choong-Hoon park, Shamima Akhtar Sharmin,
495   Hyoshin Lee Ki-Won OH Byung-Wook Yun and Byung-Hyun. Proteome analysis of soybean roots
496   under waterlogging stress at an early vegetative stage. Journal of Bioscience 2010; 35: 1 49–62.
497
498   Ismond, K. P., Dolferus, R., dePauw, M., Dennis, E. S., and Good, A. G. Enhanced low oxygen
499   survival in Arabidopsis through increased metabolic flux in the fermentative pathway. Plant Physiol
500   2013; 132: 1292–1302.
501

[80] Ismond

Aardra Kachroo and Pradeep Kachroo.2009. Fatty acids- Derived signals in plant defense. Annu Rev. Phytopatho l7: 153–76.

Kaur K, Gupta AK, Kaur N. Effect of water deficit on carbohydrate status and enzymes of carbohydrate metabolism in seedlings of wheat cultivars. Indian Journal of Biochemistry and Biophysics 2007; 44: 223-230..

Kaur, Shalinder Shivani Ranab, Harminder Pal Singha,, Daizy Rani Batishb, and Ravinder Kumar Kohlib Citronellol Disrupts Membrane Integrity by Inducing Free Radical Generation. Z. Naturforsch 2011; 66 c: 260 – 266.

Kevin Francis, Crystal Smitherman, Shirley F. Nishino, Jim C. Spain, Giovanni Gadda. The Biochemistry of the Metabolic Poison Propionate 3-Nitronate and Its Conjugate Acid, 3-Nitropropionate. International Union of Biochemistry and Molecular Biology 2013; 65: (9): 759–768.

S. Kimura, K. Sakaguchi, DNA repair in plants, Chem. Rev. 2006; 106: 753–766.

Kirsch C, Takamiya-Wik M, Reinold S, Hahlbrock K, Somssich IE. Rapid, transient, and highly localized induction of plastidial omega-3 fatty acid desaturase mRNA at fungal infection sites in *Petroselinum crispum*. Proc. Natl. Acad. Sci. USA 1997; 94:2079–84.

Klára Kosová, Pavel Vítámvás, Milan O Urban, Ilja T Prášil and Jenny Renaut. Plant Abiotic Stress Proteomics The Major Factors Determining Alterations in Cellular Proteome. Frontiers in Plant Science 2018; 122: 9 1-22–322.

Komatsu S Deschamps, Thibaut Susumu Hiraga, Mikio Kato Mitsuru, Chiba Akiko Hashiguchi, Makoto Tougou Satoshi, Shimamura Hiroshi Yasue. Characterization of a novel flooding stress-responsive alcohol dehydrogenase expressed in soybean roots. Plant Molecular Biology 2011; 77: 309.

Kun Zhang, Huiting Cui,   Shihao Cao, Li Yan Mingna Li1, Yan Sun. 2019.Overexpression of *CrCOMT* from *Carex rigescens* increases salt stress and modulates melatonin synthesis in *Arabidopsis thaliana.* Plant Cell Reports 223-228.

538 Kwasniewski M, Szarejko I. Molecular cloning and characterization of beta-expansin gene related
539 to root hair formation in barley. Plant Physiol 2006; 141(3):1149–1158.
540

541 Jackson Y A, Mark F. Williams, Lawrence A. D. Williams, Kisha Morgan & Flona A. Redway.
542 Insecticidal Properties of Benzofuran-2-carboxylic Acid Derivatives. Pestic. Sci 1998; 53: 241-244.
543

544 Jander G, Joshi V. Recent progress in deciphering the biosynthesis of aspartate-derived amino acids
545 in plants. Mol Plant 2010; 3:54–65.
546

547 Lang, Nguyen Thi; Nguyen Quang Cao Binh, Chau Thanh Nha, Bui Chi Buu. A candidate gene
548 response to drought stress condition in rice (Oryza sativa L.). Omonrice 2010; 17: 105-113.
549

550 Lee, S.; Lee, E.J.; Yang, E.J.; Lee, J.E.; Park, A.R.; Song, W.H.; Park, O.K Proteomic
551 identification of annexins, calcium-dependent membrane binding proteins that mediate osmotic
552 stress and abscisic acid signal transduction in Arabidopsis. Plant Cell 2004; 16: 1378–1391.
553

554 Lee, Y. and Kende, H. Expression of a-expansin and expansin-like genes in deepwater rice. Plant
555 Physiol 2002. **130**: 1396–1405.
556

557 Li AX, Han YY, Wang X, Chen YH, Zhao MR, Zhou S-M, Wang W (2015a) Root-specific
558 expression of wheat expansin gene TaEXPB23 enhances root growth and water stress tolerance in
559 tobacco. Environ Exp Bot 2015; 110:73–84.
560

561 Lianwei Peng, Hiroshi Yamamoto, Toshiharu Shikanai. Structure and biogenesis of the chloroplast
562 NAD(P)H dehydrogenase complex. Biochimica et Biophysica Acta 2011; 1807: 945–953.
563

564 Lisch D. How important are transposons for plant evolution? Nat. Rev. Genet 2012; 14: 49–61.
565

566 Liu H, Ma Y, Chen NA, Guo S, Liu H, Guo X, Chong K, Xu, Y. Overexpression of stress-inducible
567 OsBURP16, the β subunit of polygalacturonase 1, decreases pectin content and cell adhesion and
568 increases abiotic stress sensitivity in rice. Plant Cell and Environment 2013; 37: 1144-1158.
569

570 Loreto, F., Forster, A., Durr, M., Csiky, O. & Seufert, G. On the monoterpene emission under heat
571 stress and on the increased thermotolerance of leaves of Quercusilex L. fumigated with selected
572 monoterpenes. Plant Cell Environ 1998; 21: 101–107.
573

574   Loreto, F and Velikova, V. Isoprene produced by leaves protects the photosynthetic apparatus
575   against ozone damage, quenches ozone products, and reduces lipid peroxidation of cellular
576   membranes. Plant Physiol 2001; 127: 1781–1787.
577

578   Clemens Lamberth, Stephan Trah, Sebastian Wendeborn, Raphael Dumeunier, Mikael Courbot,
579   Jeremy Godwin, Peter Schneiter. Synthesis and fungicidal activity of tubulin polymerisation
580   promoters. Part 2: Pyridazines. Bioorganic & Medicinal Chemistry 2012; 20: 2803–2810.
581

582   Lunn J E. Sucrose-phosphatase gene families in plants. Gene 2003; 303: 187–196.
583

584   Luis C. Romero, M. Ángeles Aroca, Ana M. Laureano-Marín, Inmaculada Moreno, Irene García,
585   and Cecilia Gotor. Cysteine and Cysteine-Related signaling Pathways in *Arabidopsis thaliana*.
586   Molecular Plant 2014; 7 (4): 264-276.
587

588   McClintock, B. The significance of responses of the genome to challenge. Science 1984*;* 226:
589   792–80.
590

591   Madhu Anejaa, Thomas J. Gianfagnaa, Prakash K. Hebbarb. Trichoderma harzianum produces
592   nonanoic acid, an inhibitor of spore germination and mycelial growth of two cacao pathogens.
593   Physiological and Molecular Plant Pathology 2005; 67: 304–307.
594

595   Ana Maldonado M, Sira Echevarría-Zomeño, Sylvia Jean-Baptiste,Martha Hernández, Jesús Jorrín-
596   Novo V. Evaluation of three different protocols of protein extraction for rabidopsis thaliana leaf
597   proteome analysis by two dimensional electrophoresis. J. Proteomics 2008; 71(4): 461-47.
598

599   Mao,H.,Wang,H.,Liu,S.,Li,Z.,Yang,X.,Yan, J et al. A transposable elementina NAC gene is
600   associated with drought tolerance in maize seedlings. Nat.Commun 2015; 6: 8326.
601

602   Massonie G (1980) Breeding of a biotype of Myzus persicae Sulzer on a synthetic medium. V.
603   Influence of oxalic and gentisic acids on the nutritive value of a synthetic medium. Ann Nutr
604   Aliment 1980; 34: 139–146.
605

606   Mawlong Ibandalin, M. S. Sujith Kumar, Dhiraj Singh. Furan fatty acids: their role in plant
607   systems. Phytochem Rev 2014; 1-7.
608

M`ene-Saffran´e L, Dubugnon L, Ch´etelat A, Stolz S, Gouhier-Darimont C, Farmer EE.
Nonenzymatic oxidation of trienoic fatty acids contributes to reactive oxygen species management
in *Arabidopsis*. J Biol Chem 2009;. 284:1702–708.
Mintz-Oron, S., Mandel, T., Rogachev, I., Feldberg, L., Lotan, O., Yativ, M., Aharoni, A.(2008).
Gene expression and metabolism in tomato fruit surface tissues. Plant Physiology 2008; 147: 823–
851.
615
616 Mittler, R. Abiotic stress, the field environment and stress combination. Trends Plant Sci 2006; 11:
617 15–19.
618
619 Moellering, E.R., Muthan, B. and Benning, C. Freezing tolerance in plants requires lipid remodeling
620 at the outer chloroplast membrane. Science 2010; 330: 226–228.
621
622 Narendra Tuteja, Parvaiz Ahmad, Brahma B. Panda, Renu Tuteja (2009) Genotoxic stress in plants:
623 Shedding light on DNA damage, repair and DNA repair helicases. Mutation Research 2009; 681:
624 134–149.
625
626 Okazaki, Y., Otsuki, H., Narisawa, T., Kobayashi, M., Sawai, S., Kamide, Y., Kusano, M., Aoki,
627 T., Hirai, M.Y. and Saito, K. A new class of plant lipid is essential for protection against phosphorus
628 depletion. Nat. Commun 2013b; 4: 1510.
629
630 Prasad, P. V. V., Pisipati, S. R., Momcilovic, I., and Ristic, Z. Independent and combined effects of
631 high temperature and drought stress during grain filling on plant yield and chloroplast EF-Tu
632 expression in spring wheat. J. Agron. Crop Sci 2011; 197: 430–441.
633
634 Piotrowski, M. Primary or secondary? Versatile nitrilases in plant metabolism. Phytochemistry
635 2008; 69: 2655–2667.
636
637 Ralph Reimholz, Peter Geigenberger, Mark Stitt. Sucrose-phosphate synthase is regulated via
638 metabolites phosphorylation in potato tubers, in a manner analogous to the enzyme in leaves. Planta
639 1994; 192: 480-488.
640
641 Rasmann S. et al. Recruitment of entomo-pathogenic nematodes by insect-damaged maize roots.
642 Nature 2005; 434: 732–737.
643

Rasmann, S., Turlings, T.C.J. Simultaneous feeding by aboveground and belowground herbivores
attenuates plant-mediated attraction of their respective natural enemies. Ecology Letters 2007; 10:
926–936.

Reinhard Korbinian Proels and Ralph Huckelhoven. Cell-wall invertases, key enzymes in the
modulation of plant metabolism during defence responses. *Molecular Plant pathology* 2014; 15(8):
858–864.

Rivas-San Vicente,M. ,and Plasencia,J. Salicylic acid beyond defence: its role in plant growth and
development. J Exp.Bot 2011; 62: 3321–3338.

Rizhsk, L., Liang, H., Shuman, J., Shulaev, V., Davletova, S., and Mittler, R. (2004).When defense
pathways collide. The response of Arabidopsis to a combination of drought and heat stress. Plant
Physiol 2004; 134: 1683–1696.

Robles P, Micol JL, Quesada V. Arabidopsis MDA1, a Nuclear-Encoded Protein, Functions in
Chloroplast Development and Abiotic Stress Responses. PLoS ONE 2012; 7(8): e42924.

Roitsch T, Balibrea ME, Hofmann M, Proels R, Sinha A K. Extracellular invertase: key metabolic
enzyme and PR protein. J Exp Bot 2003; 54: 513–524.

Romero, L.C., Garcia, I., and Gotor, C. L-Cysteine Desulfhydrase 1 modulates the generation of the
signaling molecule sulfide in plant cytosol. Plant Signal Behav 2013; 8: e24007

Rosenthal G. A. Scientific American 1986; 254: 76.

Rufty T W, Huber S C, Volk R J. Alternations in leaf carbohydrate metabolism in response to
nitrogen stress. Plant Physiol 1988; 88:725–730.

Salekdeh GH, Siopongco J, Wade LJ, Ghareyazie B, Bennett J. Proteomic analysis of rice leaves
during drought stress and recovery. Proteomics 2002; 2: 1131–1145.

Siegien I and R Bogatek. Cyanide action in plants: from toxic to regulatory. Acta Physiol Plant
2006; 28: 483–497.

Saori Ogawa and Shiro Mitsuya. *S*-methylmethionine is involved in the salinity tolerance of *Arabidopsis thaliana* plants at germination and early growth stages. Physiologia Plantarum 2012; 144: 13–19. 2012.

Sanchez-Perez R, Jorgensen K, Olsen C E, Dicenta F, Moller B L. Bitterness in almonds. Plant Physiol 2008; 146(3):1040–1052.

Shapiro J. A 21st century view of evolution: genome system architecture, repetitive DNA, and natural genetic engineering. Gene 2005; 345: 91–100.

Sharkey, T D and Singsaas, E L. Why plants emit isoprene. Nature 1995; 374: 769.

Solís-Guzmán, Marí.Gloria., Argüello-Astorga, G., López-Bucio, José., Ruiz-Herrera, Leó.Francisco., López-Meza, J.E., Sánchez-Calderón, L., Carreón-Abud, Yazmí., Martínez-Trujillo, M., *Arabidopsis thaliana* sucrose phosphate synthase (*sps*) genes are expressed differentially in organs and tissues, and their transcription is regulated by osmotic stress, *Gene Expression Patterns* 2017;

Spiteller G. Furan fatty acids: occurrence, synthesis, and reactions. Are furan fatty acids responsible for the cardioprotective effects of a fish diet? Lipids 2005; 40: 755–771.

Suzuki, N., Rivero, R. M., Shulaev, V., Blumwald, E., and Mittler, R. Abiotic and biotic stress combinations. New Phytol 2014; 203: 32–43.

Swidzinski JA, et al. A custom microarray analysis of gene expression during programmed cell death in Arabidopsis thaliana. Plant J 2002; 30: 431-446.

Tahir, J., Watanabe, M., Jing, H.C., Hunter, D.A., Tohge, T., Nunes-Nesi, A., Brotman, Y., Fernie, A.R., Hoefgen, R., and Dijkwel, P.P. Activation of R-mediated innate immunity and disease susceptibility is affected by mutations in a cytosolic O-acetylserine (thiol) lyase in *Arabidopsis*. Plant J 2013; 73: 118–130.

Takshak Swabha; S. B. Agrawal. Alterations in metabolite profile and free radical scavenging activities of Withania somnifera leaf and root extracts under supplemental ultraviolet-B radiation. Acta Physiol Plant 2015; 37:260.

Taylor N L, et al. Analysis of the rice mitochondrial carrier family reveals anaerobic accumulation
of a basic amino acid carrier involved in arginine metabolism during seed germination. Plant
Physiology 2010; 154: 691-704.

Tsygankova Victoria, Yaroslav Andrusevich, Olexandra Shtompel, Artem Hurenko, Roman
Solomyannyj, Galyna Mrug, Mikhaylo Frasinyuk, Volodymyr Brovarets. Stimulating effect of five
and six-membered heterocyclic compounds on seed germination and vegetative growth of maize
(*Zea mays* L.) International Journal of Biology Research 2016; 1 (4): 01-14

Turner, M. F., Heuberger, A. L., Kirkwood, J. S., Collins, C. C., Wolfrum, E. J., Broeckling, C. D.,
et al. Non-targeted metabolomics in diverse Sorghum breeding lines indicates primary and
secondary metabolite profiles are associated with plant biomass accumulation and photosynthesis.
Front. Plant Sci 2016; 7: 953.

Upchurch R G. Fatty acid unsaturation, mobilization, and regulation in the response of plants to
stress. Biotechnol. Lett 2008; 30:967–77.

Urbonavieius J, Qian Q, Durand J M B, Hagervall T G, and Bjork G R.. Improvement of reading
frame maintenance is a common function for several tRNA modification. EMBO J 2001; 20: 4863-
4873.

Velikova, V. and Loreto, F. On the relationship between isoprene emission and thermotolerance in
Phragmites australis leaves exposed to high temperatures and during the recovery from a heat stress.
Plant Cell Environ 2005; 28: 318–327.

Vile, D., Pervent, M., Belluau, M., Vasseur F.,Bresson, J., Muller, B.,et al. Arabidopsis growth
under prolonged high temperature and water deficit: independent or interactive effects? Plant Cell
Environ *2*012; 35: 702–718.

Vlot, A.C., Dempsey,D.M.A., and Klessig,D.F. Salicylic acid, a multi- faceted hormone to combat
disease. Annu.Rev.Phytopathol 2009; 47: 177–206.

Von Burg R. Toxicology update. J Appl Toxicol 1994; 14: 233–237.

Wen, W., Li, K., Alseekh, S., Omranian, N., Zhao, L., Zhou, Y., et al. Genetic determinants of the
network of primary metabolism and their relationships to plant performance in a maize recombinant
inbred line population. Plant Cell 2015; 27: 1839–1856.

Wataru Yamori, Naoki Sakata, Yuji Suzuki, Toshiharu Shikanai and Amane Makino. Cyclic electron flow around photosystem I via chloroplast NAD (P)H dehydrogenase (NDH) complex performs a significant physiological role during photosynthesis and plant growth at low temperature in rice. The Plant Journal 2011; 68: 966–976.

Williams, L. A. D., Anderson, M. J. and Jackson, Y. A., Insecticidal activity of synthetic 2-carboxylbenzofurans and their coumarin precursors. Pestic Sci 1994; 42: 167-71.

Wilson, Shan, H., W. K., Phillips, D. R., Bartel, B., and Matsuda, S. P. T. Trinorlupeol: A major nonsterol triterpenoid in arabidopsis. Organic Letters 2008; 10: 1897–1900.

Wirtz, M., and Hell, R. Functional analysis of the cysteine synthase protein complex from plants: structural, biochemical and regulatory properties. J. Plant Physiol 2006; 163: 273–286.

Winkel-Shirley B. Flavonoid biosynthesis, A colorful model for genetics, biochemistry, cell biology and biotechnology. Plant Physiol 2001; 26:485-93.

Won S, Choi S, Kumari S, Cho M, Lee SH, Cho H. Root hair specific EXPANSIN B genes have been selected for Graminaceae root hairs. Mol Cells 2010. 30(4):369–376.

Xing SC, Li F, Guo QF, Liu DR, Zhao XX, Wang W. The involvement of an expansin geneTaEXPB23 from wheat in regulating plant cell growth. Biol Plant 2009; 53(3):429–434.

Yan J X, Wait R Berkelman, T Harry R A, et al. Electrophoresis 2000; 21: 3666-3672.

Yaeno T, Matsuda O, Iba K. Role of chloroplast trienoic fatty acids in plant disease defense responses. Plant J 2004; 40: 931–41.

Yip W K, Yang S F. Cyanide metabolism in relation to ethylene production in plant tissues. Plant Physiol 1988; 88(2):473–476.

Yoshihara T, Sogawa K, Pathak MD, Juliano BO, Sakamura S. Oxalic acid as a sucking inhibitor of the brown planthopper in rice (Delphacidae, Homoptera). Entomol Exp Appl 1980; 27: 149–155.

Yozo Okazaki and Kazuki Saito. Roles of lipids as signaling molecules and mitigators during stress response in plants. The Plant Journal 2014; 79: 584–596.

788   Zhao Y, Cai M, Zhang X, Li Y, Zhang J, et al. Genome-Wide Identification, Evolution and
789   Expression Analysis of mTERF Gene Family in Maize. PLoS ONE 2014; 9(4): e94126.
790

791   Zhao D, Reddy K R, Kakani V G, Reddy V R. Nitrogen deficiency effects on plant growth, leaf
792   photosynthesis, and hyperspectral reflectance properties of sorghum. Eur J Agron 2005; 22: 391–
793   403.
794

795   Zhao, M.R., Han, Y.Y., Feng, Y.N., Li, F. and Wang, W. Expansins are involved in cell growth
796   mediated by abscisic acid and indole-3-acetic acid under drought stress in wheat. Plant Cell Rep
797   2012. 31: 671–685.
798

799   Zhang W, Wang C, Qin C, Wood T, Olafsdottir G, Welti R, and Wang X. The oleate-stimulated
800   phospholipase D, PLD delta, and phosphatidic acid decrease H2O2-induced cell death in
801   Arabidopsis. Plant Cell 2003, 15: 2285–2295.
802

803   Zivy, M., Wienkoop, S., Renaut, J., Pinheiro, C., Goulas, E., and Carpentier, S. The quest for tolerant
804   varieties: the importance of integrating "omics" techniques to phenotyping. Front. Plant Sci 2015;
805   6: 448.
806

807   Zou H, Wenwen Y, Zang G, Kang Z, Zhang Z, Huang J, Wang G. OsEXPB2, a b-expansin gene, is
808   involved in rice root system architecture. Mol Breed 2015; 35(1):
809

810   Zunino M. P. and Zygadlo J. A. Effect of monoterpenes on lipid peroxidation in maize. Planta 2004;
811   219: 303 – 309.