Factors influencing riverine utilization patterns in two

sympatric macaques

Yosuke Otani[1*], Henry Bernard[2], Anna Wong[2], Joseph Tangah[3], Augustine Tuuga[4], Goro

Hanya[5], Ikki Matsuda[3, 6, 7, 8]

[1]*Center for the Study of Co* Design, Osaka University, Toyonaka, Osaka 560-0043,*

Japan

[2]*Institute for Tropical Biology & Conservation, University of Malaysia Sabah, Sabah,*

Malaysia

[3]*Forest Research Centre, Sabah Forestry Department, Sandakan, Sabah*

[4]*Sabah Wildlife Department, Kota Kinabalu, Sabah, Malaysia*

[5]*Primate Research Institute, Kyoto University, Inuyama, Aichi, Japan*

[6]*Chubu University Academy of Emerging Sciences, Kasugai, Aichi, Japan*

[7]*Wildlife Research Center of Kyoto University, Kyoto, Japan*

[8]*Japan Monkey Centre, Inuyama, Japan*

[*]*Corresponding author, email: otani.primate.res@gmail.com (Tel: +81-6-6210-8250)*

Short title: Riverine utilization patterns of two sympatric macaques

Abstract

Many species of terrestrial animals, including primates, live in varied association with the aquatic (e.g., riverine or coastal) environment. However, the benefits that each species receive from the aquatic environment are thought to vary depending on their social and ecological characteristics, and thus, elucidating those benefits to each species is important for understanding the principles of wild animal behaviour. In the present study, to gain a more complete picture of aquatic environment use, including social and ecological factors in primates, factors affecting riverine habitat utilization of two macaque species *(Macaca nemestrina* and *M. fascicularis)* were identified and qualitative comparisons were made with sympatric proboscis monkeys *(Nasalis larvatus)*, which have different social and ecological characteristics. Temporal variation in sighting frequency of macaques at the riverbanks was positively related to the fruit availability of a dominant riparian plant species and negatively related to the river water level which affects the extent of predation pressure. Riverine utilization of macaques was greatly influenced by distribution and abundance of food (especially fruit) resources, possibly in association with predation pressure. Additionally, qualitative ecological comparisons with sympatric proboscis monkeys suggest that the drivers of riverine utilization depend on the feeding niches of the species, and different anti-predator strategies resulting from their differing social structures.

Keywords

Anti-predator strategy, aquatic environment, foraging strategy, landscape of fear, southern pig-tailed macaque

Introduction

Animals often live in forests characterized as mosaic habitats comprising both terrestrial and aquatic (e.g., riverine or coastal) environments while varying their associations with the aquatic environment. In addition to direct relationships such as obtaining food from the aquatic environment, e.g., bears eating salmon [1]; racoons eating mussels [2], the aquatic environment has an indirect impact on terrestrial animals by bringing about environmental heterogeneity. Differences in terrestrial flora, which are due to varying water abundance and light intensity at the border between aquatic and terrestrial environments, affect foraging behaviour of animals that use plants as a food source. In addition, such boundaries restrict the distribution and movement patterns like a river-barrier [3] and thus are constraints on habitat utilization. Aquatic environments are, therefore, undoubtedly important for terrestrial animals across many taxonomic groups and geographic areas.

Primates are a primary example of terrestrial animals that rely on the aquatic environment. Kempf [4] presented a comprehensive review of primate aquatic behaviours with the conclusions that the use of aquatic resources and the aquatic environment affect various aspects of primate life, including feeding, traveling, predation avoidance, and thermoregulation, although it has been noted that the represented data are inadequate to draw strong conclusions. Reports of aquatic-related primate ecology have increased in recent years, strongly supporting such proposed broader trends with special reference to primates inhabiting flooded habitats [5]. One of the notable flooded habitats is riverine forest, providing a relatively constant availability of fruits, combined with greater plant diversity and higher leaf quality when compared with dryland forest, which is due to the frequent supply of nutrient-rich soils in riverine forests that are exposed to seasonal

flooding [6,7]. Further, in contrast to dryland forest, riverine forest potentially has a better

light environment and more gap-specialists that have leaves containing higher protein

content [8]—one of the important factors influencing primate abundance [9] and dietary

choice e.g., [10,11]. The distinctive food conditions caused by the presence of such large-

scale riparian areas in forest habitat are likely to have a significant impact on primate

behavioural ecology, and thus riverine habitat is an ideal forest type for research

contributing to a fundamental understanding of primate behavioural ecology, including

how the aquatic environment affects primate distribution and ranging behaviours.

Furthermore, studying habitats in riparian environments also plays an

important role in understanding the landscape of fear [12,13]. Animals' responses to

predation risk vary over time and space; for example, they can alter their behaviour,

such as by changing time allocation patterns depending on the level of fear [14].

Boundaries between aquatic and terrestrial environments, e.g., riverbanks, can be

advantageous or disadvantageous in anti-predator strategies, because they provide

animals with a physically heterogeneous environment. In riverine habitats, river-edge

trees in exposed places are preferred by primates for roosting to avoid predation,

because this allows detection of approaching predators [15-18]. For example, proboscis

monkeys (*Nasalis larvatus*) often choose sleeping sites in river-edge trees in areas with

narrow river widths; this provides good escape routes from terrestrial predators such as

clouded leopards (*Neofelis diardi*) [19-21], which generally show a nocturnal activity

pattern [22]. Conversely, on flooded days when water levels are extremely high, proboscis

monkeys remain in inland forest because of reduced predation threat, as terrestrial

predators are prevented from foraging on the forest floor [23]. Therefore, riverine forests

have temporal and spatial environmental heterogeneity, and revealing the behavioural

changes of animals in response to such heterogeneity can provide insight into

behavioural adaptation in response to fear of predators. Consequently, studying primates

that live in riverine habitats is ideal for elucidating their resource exploitation patterns

relative to anti-predator strategies; these are central components to understanding

various primate habitat adaptations.

Long-tailed macaques (*Macaca fascicularis*) and southern pig-tailed macaques

(*M. nemestrina*, hereafter pig-tailed macaques, unless otherwise noted) are widely

distributed throughout the Sundaic region of Southeastern Asia, often coexisting in a

broad variety of habitats [24]. Long-tailed macaques have often been reported sleeping in

river-edge trees [16,25,26], and van Schaik, et al. [16] indicate that multiple ecological factors

(food availability, temperature and predation risk) influence their riverine utilization.

Conversely, ecological data for pig-tailed macaques, including the northern species

(*Macaca leonina*) and southern pig-tailed species, are less complete [27-30], and their

aquatic-related behaviours are rarely reported. Rodman [31] reported that microhabitat

segregation occurs between the two species and that, unlike long-tailed macaques, pig-

tailed macaques generally do not use riverine areas. Conversely, Albert, et al. [18] noted

that the preference of northern pig-tailed macaques to locate their sleeping sites in river-

edge trees was a part of their predator avoidance strategy. In addition to predation

pressure, as in long-tailed macaques, food availability may affect riverine utilization in

pig-tailed macaques. By determining the effects of food availability on riverine forest

use in macaque habitat, it is possible to evaluate whether the ranging behaviour of pig-

tailed macaques is affected by food availability in the same way as that of long-tailed

macaques.

We examined the riverine habitat utilization patterns of sympatric long-tailed

and pig-tailed macaques inhabiting a secondary riverine forest on the island of Borneo.

At this site, it has been reported that sympatric primates, including the two study

macaque species, proboscis monkeys and other species, prefer to utilize river-edge trees

for night-time sleeping, although the frequency of riverine usage is different among

these sympatric primates [32]. A previous study on proboscis monkeys at the site

suggested that riverine preference is related to high predation pressure, but not river-

edge dietary choice [17,23]. However, in theory, predation risk varies with group size and

body weight [33-35]. The two species of macaques in this study may have different anti-

predator strategies from proboscis monkeys, as the macaques live in groups of multiple

males and females and have larger group sizes than proboscis monkeys, which live in

groups consisting of one male and multiple females [35,36] and have significantly different

body weight than the two macaque species, i.e., proboscis monkey: male, 25kg and

female, 14 kg; long-tailed macaque: male, 6 kg and female, 4 kg; pig-tailed-macaque,

male, 14 kg and female, 7 kg [37-39]. Additionally, the ranging behaviour of primate

species that prefer to feed on patchy and clumped food sources, e.g., fruits and flowers,

such as northern pig-tailed and long-tailed macaques [16,25,26], is more influenced by food

distribution and abundance than that of primate species that prefer to feed on ubiquitous

food sources (i.e., leaves), such as proboscis monkeys [40]. Therefore, for macaques, the

location of foraging patches may have a stronger effect on sleeping site selection.

In the present study, to gain a more complete picture of the riverine utilization

patterns in the two species of sympatric macaques, we sought to 1) evaluate temporal

variation in their riverine usage and asses the factors affecting riverine usage, including

physical environment, i.e., river width and water level, and 2) describe their diets in

river-edge areas, with a comparison of those availability. In addition, the effects of

feeding niches and social structure on ranging behaviour are discussed by qualitatively

comparing the characteristics of riverine utilization of the two species of macaques with

those of proboscis monkeys reported in previous studies [17,21,23].

Methods

Study area and subjects

We performed the observations over two years from 2012 to 2014 in riverine forests along

the Menanggul River (average river width 0–4000 m from the river mouth: 19.9 m), a

tributary of the Kinabatangan River, Sabah, Borneo, Malaysia (118°30'E, 5°30'N). The

south side of the Menanggul River is covered extensively in natural forest, whereas the

north side has been deforested for oil palm plantations, except for a protected zone along

the river [41]. The mean minimum and maximum daily temperatures were approximately

24°C and 30°C, respectively, and the mean annual precipitation at the site was 2,474 mm

[6]. The riverine forest was inhabited by long-tailed and pig-tailed macaques, as well as

proboscis monkeys, silver langurs (*Trachypithecus cristatus*), Hose's langurs (*Presbytis*

hosei), maroon langurs (*Presbytis rubicunda*), Bornean gibbons (*Hylobates muelleri*) and

orangutans (*Pongo pygmaeus*) [32]. Long-tailed and pig-tailed macaques under observation

were well habituated to observers in boats, as this area is one of the main tourist attractions

in the region, with many boats and tourists visiting the Menanggul River since more than

10 years ago.

Data collection

Boat-based surveys

Surveys by boat in the late afternoon are considered the most effective method for

studying primates, including the two sympatric macaques in this region, because they

often sleep in riverside trees [32]. We therefore collected data on the distribution pattern of

the sympatric macaques in the late afternoon (16:00–19:00) for 434 days from June 2012

to July 2014 via boat-based surveys. We conducted the surveys along the river at a speed

of approximately 4–6 km/h. When we detected a group or individual macaque, we

switched off the boat engine to avoid disturbing them and paddled closer to record their

species and numbers. We divided the river into 50-metre sections from the river mouth to

4,000 m inland, recording the river sections where sightings of macaques were made.

When group members were distributed over several sections, the section containing the

largest number of individuals in the visual inspection was defined as the detected section.

Boat-based behavioural observations

We collected behavioural data from the adults and subadults in both macaque species

during the boat-based surveys. During the observation periods, we recorded the activity

of all visible primates at the time of detection by scan sampling [42] over 60 days from June

2012 to May 2014. We divided the behaviours into seven categories: feeding, grooming,

moving, resting, playing, fighting and other. Food items consisted of leaves, fruits,

flowers and other items, and food plants were taxonomically identified *in situ*.

Vegetation survey

We established a total of 16 transects (200–500 m × 3 m) on both sides of the river at 500

m intervals from the river mouth to 4,000 m. The total length of the 16 transects was

7,150 m, and the total surveyed area was 2.15 ha. We taxonomically identified trees ≥10

cm in diameter at breast height (DBH) and vines ≥5 cm in diameter that were located

within the transects [6]. Because these data did not include herbaceous climbers, we added

data on *Cayratia trifolia* (Vitaceae), an important food source for macaque species (see

results) in case the climber was entangled in the surveyed trees/vines.

Monthly availability survey for C. trifolia *fruits*

We carried out a fruit quantity survey of *C. trifolia*, which has fleshy, juicy, dark purple

and nearly spherical fruits ca. 1 cm in diameter [43]. It has been reported that sympatric

proboscis monkey at this study site consume *C. trifolia* [40]. Early in each monthly survey

from July 2012 to June 2014, we travelled by boat up to 4000 m from the river mouth and

counted all the visible mature and young fruits of *C. trifolia* on both riverbanks. Two

observers independently counted the number of fruits in a section, and the average was

defined as the availability of *C. trifolia* in the section. We judged the degree of fruit

maturation by its colour, with mature fruit being purple to black and young fruit being

green.

Rainfall, water level and river width

We measured daily rainfall every morning at base camp approximately 1.5 km from the

mouth of Menanggul River, using a tipping bucket rain gauge. We also recorded water

level and river width to evaluate the effects of river level on behaviours of the study

macaques. We installed a water level gauge at the mouth of Menanggul River, and

measured the water level at the end of the boat-based survey (17:00–19:00). We

measured the river width at the start and end points of each 50 m section with a laser

rangefinder, and the average value of the start and end points was used as a

representative value of the river width of the section.

Data analysis

To extract time-series characteristics of the increase or decrease in the number of *C.*

trifolia fruit, we conducted seasonal decomposition of time series by loess (STL) [44]

which is a filtering procedure for decomposing a seasonal times series into three

components: trend, seasonal, and remainder or residual.

We evaluated the effects of the availability of *C. trifolia* fruits, rainfall and river

width on the sighting frequency of the two macaque species counted during the boat-

based surveys, using a hierarchical Bayesian continuous-time structural equation model

(CtSEM). Models were fitted using the R package ctsem ver. 2.5.0 [45] with four chains

and 4,000 iterations. The CtSEM modelling addresses unequally spaced time intervals

in longitudinal data assessment. Since the number of survey days varied between

months and the survey days were not evenly spaced, the monthly mean values were not

strictly equally spaced time interval data. Through a hierarchical Bayesian framework,

CtSEM allows for the estimation of continuous time processes of a sample while

accounting for potential subject-level deviations by using population model estimates to

inform subject-level model priors [46]. The possible temporal autocorrelations among our

data are the total amount of *C. trifolia* fruits, and the monthly mean number of counted

macaques on each day. In addition, since there is a possibility that spatial

autocorrelation occurs between adjacent 50 m sections, the 500 m section was adopted

for examination of factors for sightings of macaques on riverbanks. As a result of the

evaluation of the spatial autocorrelation by Moran's *I* index, which is the most

commonly used coefficient in univariate autocorrelation analyses, such a trend was not

detected at each 500 m section for the monthly sighting frequency of the macaques, i.e.,

mean monthly number of counted pig-tailed (Moran *I* statistic index = −0.324 – −0.115,

p > 0.3) and long-tailed macaques (Moran *I* statistic standard deviate = −0.299 –

−0.085, *p* > 0.2), and availability of *C. trifolia* fruits (Moran *I* statistic standard deviate

= −0.224 – −0.101, *p* > 0.2). This indicates that the 500 m section is a unit that can be

analysed without considering spatial autocorrelation and is a suitable analytical unit for

subsequent analysis. Predictor and independent variables were z-standardized to build a

common metric. We performed the calculations using R ver. 3.6.1 [47]. In the

representation of the result of the model, SD refers to posterior standard deviation and

PCI refers to posterior credibility intervals. The PCI indicates the probability that the

parameter falls between the lower (2.5%) and upper (97.5%) limits.

Results

Boat-based survey: sighting frequency of primates and consumed food items

During the study period, there were a total of 3,180 detection events for six species of

primates, and 39,907 individuals were observed during boat-based surveys (Table 1).

Long-tailed and pig-tailed macaques accounted for 37.0% and 25.7% of the total

number of observed individuals, respectively (Table 1). We collected a total of 66 and

277 feeding records for pig-tailed and long-tailed macaques, respectively. Fruits and

flowers/buds of *C. trifolia* and *Dillenia excelsa* were by far the most important foods at

the riverbanks, which constituted 22.7%, and 50.9% (*C. trifolia*), and 42.4% and 18.8%

(*D. excelsa*) of the total feeding records in pig-tailed and long-tailed macaques,

respectively (Table 2).

Vegetation characteristics and food availability

We marked 1,645 trees and 497 vines (180 species, 125 genera, 52 families) along our

16 trails (for details, see [6]). *Cayratia trifolia* was entangled in only four of 1,645 marked

trees (0.24%), and was located in well-lit forest gaps caused by fallen trees. Conversely,

C. trifolia was clearly more abundant along the riverbanks, and was found in 12.5%–

72.5% of all 50-m river sections in each monthly survey. Contrary to the distribution

pattern of *C. trifolia*, *D. excelsa* was more abundant inside the forest: of 98 *D. excelsa*

plants in the vegetation transects, 84 (86%) were found in the inland forest (>50 m from

the riverbanks). The tendency was the same for *Mallotus muticus* (120 of 149 in the

inland forest), which was the third most common plant in feeding records of pig-tailed

and long-tailed macaques; *Albizia corniculate* (22 of 28 in the inland forest) and *Ficus*

spp. (20 of 23 in the inland forest), which were the fourth and fifth most common in the

feeding records of pig-tailed macaques; and *Antidesma thwaitesianum* (27 of 28 in the

inland forest) and *Xylosma sumatrana* (53 of 67 in the inland forest), which were the

fourth and fifth most common in the feeding records of long-tailed macaques.

The mean monthly number of counted *C. trifolia* fruits on the riverbanks was

23,283.7 (SD ±19,707.3; range 1,844–78,505). The mean monthly numbers of young and

mature fruits were 22,795.0 (±19,260.0; 1,806–76,002) and 488.6 (±572.4; 38–2,503),

respectively. STL based on the number of counted fruits in each monthly survey clarified

that the *C. trifolia* fruit availability seasonally fluctuated (Fig. 1). Fruit availability

declined in the middle of 2013 and then increased again, and tended to increase and

decrease every 4−6 months.

Factors affecting temporal variation in sighting frequency of macaques

The mean and 95% PCI of T_0 mean parameters (Table 3), representing the relationship

between the subject's initial states with their later states throughout the latent process,

were 0.408 (−0.089, 0.911), 0.313 (−0.199, 0.783) and 0.270 (−0.230, 0.805) for the

sighting frequency of pig-tailed and long-tailed macaques and *C. trifolia* fruits,

respectively. This indicated that there was no tendency for each parameter to increase or

decrease substantially over time, because for all parameters zero falls within the PCIs.

The manifest mean parameters (Table 3) represent the average level of the processes,

reflecting the intercepts of the sighting frequency of pig-tailed and long-tailed macaques

and availability of *C. trifolia* fruits.

The regression coefficients of the monthly mean sighting frequency of

macaques and monthly availability of *C. trifolia* fruits within each section denoted the

temporal autoregressive effects (Fig. 2); *C. trifolia* fruits had a temporal autocorrelation

that lasted approximately three months, but sighting frequency of macaques had no such

autocorrelation.

To assess the effects of the monthly availability of *C. trifolia* fruits on the

monthly mean sighting frequency of pig-tailed and long-tailed macaques, we evaluated

the drift parameters representing the cross effects (Table 3). The positive values showed

that the *C. trifolia* fruits had a positive effect on the sighting frequency of pig-tailed

(mean = 1.560, SD = 0.653, PCI = [0.131, 2.710]) and long-tailed (mean = 0.850, SD =

0.148, PCI = [0.581, 1.168]) macaques throughout the study period, indicating that

more macaques were sighted in areas where *C. trifolia* fruits were abundant.

Conversely, the sighting frequency of macaques did not affect the availability of *C.*

trifolia fruits. Additionally, the positive values indicated that the sighting frequency of

both macaques had positive effects on each other (pig-tailed to long-tail macaques,

mean = 0.984, SD = 0.367, PCI = [0.356, 1.802]; long-tailed to pig-tailed macaques,

mean = 0.753, SD = 0.281, PCI = [0.272, 1.373]).

Of the time-dependent/independent variables, we detected neither significantly

positive nor negative effects of the monthly rainfall or river width on the mean monthly

sighting frequency of both macaque species and the availability of *C. trifolia* fruits,

while the water level had a negative effect on those factors (Table 3).

Discussion

We found that ecological factors influenced the riverine habitat utilization of the two

sympatric macaque species in this study. One of the important factors was the availability

of *C. trifolia,* which was also the most consumed plant species by the macaques; their

temporal variation in sighting frequency at the riverbanks was positively related to the

abundance of *C. trifolia* fruits. Conversely, a previous study of proboscis monkeys at this

study site reported that food availability is not a fundamental factor for explaining their

riverine habitat utilization patterns in the late afternoon [17]. Differing dietary preference in

relation to the digestive physiology between the two macaque species and proboscis

monkeys may have created this inconsistency. Hindgut-fermenting primates (e.g.,

macaques) generally show a stronger preference for fruits than foregut-fermenting

primates (e.g., proboscis monkeys), allowing the latter to exploit a diet of leaves in greater

quantities [48,49]. Indeed, *C. trifolia* fruits are not the most preferred food by proboscis

monkeys at this study site [40].

Predation pressure may also be a factor that affects riverine habitat utilization

patterns for the two macaque species in this study, which was also shown for proboscis

monkeys [17]. The landscape of fear is an important driver of prey habitat utilization [50].

Although it is generally difficult to evaluate predation pressure on primates because of

how rare it is to directly observe cases of attempted or successful predation [34,51],

behavioural responses to predator presence were reported to have more far-reaching

consequences for prey ecology than the actual killing of individuals [12]. Indeed, despite

the abundance of food resources when studying these macaques, their use of riverbanks

was mainly limited to the late afternoon [17,32]; one reason for this could be related to their

nocturnal anti-predator strategy.

According to previous reports [20,25,30,52-54], clouded leopards (*Neofelis diardi*),

crocodiles (*Crocodylus porosus* and *Tomistoma schlegeli*), and pythons (*Python* spp.) may

be significant potential predators of macaques of any age or sex at this study site. It was

previously reported that proboscis monkeys were attacked by clouded leopards when they

were in trees [19]. Therefore, the studied macaques should also be exposed to a threat of

predation by clouded leopards, which generally show a strongly nocturnal activity pattern

[22], when the macaques sleep in trees during the night time; during this time, there is no

predation threat from crocodiles. As pythons also tend to move and search for prey during

the night, even in trees [55], threats of predation on macaques by pythons may be similar to

those by clouded leopards and may be predictable.

Riverine habitat utilization in the late afternoon and during sleeping periods at

night provides more effective protection against attacks from terrestrial predators such as

clouded leopards, because they can only approach the macaques from the landward side.

Indeed, several studies reported the use of riverine refugia by long-tailed and northern

pig-tailed macaques [16-18], possibly to reduce predation risk from such terrestrial predators.

Alternatively, the openness of the river banks may pose another problem: vulnerability to

predation by raptors. However, raptors that prey on adult diurnal primates are considered

to be absent from Southeast Asia [56], probably because there are few large raptor species

[57]. According to literature reviews [58,59], in the case of immature primates (e.g., infants and

juveniles), potential predators in the study area may include raptors such as black eagles

(*Ictinaetus malayensis*), crested serpent-eagles (*Spilornis cheela*), and bat hawks

(*Macheiramphus alcinus*) [19]. However, predation upon any primates by these animals was

not seen at the study site; thus, their predation pressure on macaques may be less

prominent. Therefore, late afternoon and night-time use of the riverbanks by macaques

may be a response to fear of nocturnal predators. Further studies that include longer-term

direct observations and nocturnal observations would provide direct data on predation

(e.g., capture rate, loss rate of group members, and contextual data such as age, sex, and

social status of prey) and degree of fear of macaques (e.g., vigilance behaviour inland and

on river banks). Such information could provide explicit insight into the nature of the

landscape of fear.

Contrary to the anti-predator strategy observed in proboscis monkeys, in which

they select sleeping sites in areas with narrow river widths [19,20], river width was not a

significant factor that predicted the sighting frequency of macaque species at the

riverbanks in this study. Although the two species of macaques were rarely observed

swimming in the river during this study, river crossing at narrower river sections have

been more commonly observed in proboscis monkeys [21,60]; this may be why river width

was not detected as a significant factor for the macaques. Both macaque species live in

larger groups with more males than proboscis monkeys [35,36], and larger groups are

generally more vigilant and are capable of detecting predators from longer distances,

which potentially reduces predation risk [33,61]. Therefore, the benefit of sleeping on the

riverbanks for the macaques may simply be the vantage point for detecting approaching

terrestrial predators like clouded leopards and pythons, rather than ease of crossing the

river. Additionally, to gain a better understanding of our riverine anti-predator hypothesis,

further studies should evaluate the differences and similarities of predation vulnerability

levels between the two macaque species in terms of their differences in

arboreality/terrestriality levels and riverine utilization frequency on the basis of accurate

population density estimates in the habitat. Furthermore, many studies emphasized the

importance of food resource proximity for sleeping site choice by various primate species

e.g., [18,30,62,63-66]. We do not deny the possibility that selecting sleeping sites on riverbanks

may also have a secondary effect of minimizing the macaques' foraging and traveling

costs by sleeping near their feeding areas (i.e., areas abundant in *C. trifolia*).

Echoing our observation of the negative effect of river level on macaque sighting

frequency at the riverbanks, Matsuda, et al. [23] noted such an effect in proboscis monkeys

at this study site; they suggested that this occurred because of reduced predation threats,

as terrestrial predators such as clouded leopards are prevented from foraging by deep

water covering the forest floor. For the macaques, the negative effect of river level could

be caused by decreased attractiveness of the dominant food resources on the riverbanks.

Because *C. trifolia* was mostly distributed along lower parts of the riverbanks at this study

site, these plants were under water or near the surface of the river when the water level

was high. The macaque species in this study may hesitate to forage for *C. trifolia* fruits

under such circumstances because of the risk of aquatic predator attacks, despite the

reduced terrestrial predator threat due to high-level river water. Indeed, while feeding on

C. trifolia fruits on low branches (1–3 m above the river), an adult male long-tailed

macaque was preyed upon by an estuarine crocodile (*Crocodylus porosus*) [20] at our study

site. As a result, the increased risks associated with *C. trifolia* foraging because of aquatic

predators may have diminished the value of *C. trifolia* as a food resource, and this led to

a low sighting frequency of the macaques. Additionally, high water levels may simply

make access to the riverbanks more difficult for both macaque species, although this is

unlikely: both macaque species were affected by water level, even though long-tailed

macaques are more arboreal than pig-tailed macaques [67].

This study showed that riverine utilization by pig-tailed and long-tailed

macaques was greatly influenced by temporal variation in food resource abundance and

predation pressure. In addition, qualitative comparisons of sympatric proboscis monkeys

suggested that the drivers of riverine utilization depend on the feeding niches of the

species and variations in how they cope with predation pressure due to differences in

behavioural patterns and social structure. The sighting frequency of both macaques in the

riverine habitat had a positive effect, and their dietary patterns in the riverine habitat were

similar; this indicated that their feeding niche separation is ambiguous, especially on the

riverbanks, although microhabitat segregation has been reported in these two closely

related, coexisting macaque species in East Kalimantan [26,31].

It should be noted, however, that we cannot deny the possibility of the

microhabitat segregation of these two macaque species in the inland habitat where they

were not observed for this study. Co-occurrence of these two macaque species on the

riverbanks at our study site may be due to the distinctive distribution of the food resource

C. trifolia. In fact, the sighting frequency of the macaques had no effect on the subsequent

abundance of *C. trifolia* fruits. This indicates that the fruits were super-abundant;

therefore, the effect of foraging by macaques on fruit abundance was nearly negligible.

The presence of super-abundant food would mitigate feeding competition and allow the

two macaque species to co-occur on the riverbanks. Tracking macaque groups into the

inland forest during other times of the day would reveal the importance of riverine

habitats in the diet of these species and provide further insights into the mechanisms of

coexistence. In addition to feeding competition, future studies should evaluate the effects

of competition for sleeping trees to elucidate coexistence mechanisms in sympatric

macaques, as we observed the two macaque species both sharing a sleeping tree and

competing for a sleeping tree.

Acknowledgments

We thank the Sabah Biodiversity Centre and the Sabah Wildlife Department for granting

permission to carry out this research and our research assistants, especially Shah, Asnih,

Jasrudy, and the Sabah Forestry Department for support in the field. We appreciate the

support of Dr. K. Fukaya in terms of statistical analyses. We thank Clio Reid, PhD, from

Edanz Group (https://en-author-services.edanzgroup.com/ac) for editing a draft of this

manuscript. Finally, we thank Dr. Julie Teichroeb (handling editor) and two anonymous

reviewers for their constructive comments which improved our manuscript. The animals

involved in this study were not handled for this study. This study was performed

following the Research Guidelines for the Study of Wild Primates of the Primate

Research Institute, Kyoto University and in compliance with applicable Malaysian laws.

Funding

This study was partially funded by the HOPE and Human Evolution Project of KUPRI,

a MEXT Grant-in-Aid for JSPS Fellows to YO (#11J04699), JSPS KAKENHI

(#26711027, # 19KK0191 and # 19H03308 to IM; # 22687002 to GH), publication

support program of Osaka University and Japan Science and Technology Agency Core

Research for Evolutional Science and Technology 17941861 (#JPMJCR17A4). This

project has also benefitted funding from the JSPS Strategic Young Overseas Visits

Program for Accelerating Brain Circulation (S2508, PI: Hirohisa Hirai).

Author contributions

455 YO, GH and IM conceptualized the initial idea; YO and IM performed the data

456 collections; AT, HB, AW and JT arranged the data collection in the wild; YO performed

457 and interpreted the statistical analyses; YO and IM drafted the manuscript; all authors

458 contributed to the final version of the manuscript.

Competing interests

461 The authors declare that they have no competing interests.

Data availability statement

464 Data in support of the findings of this study are available from the corresponding

465 authors by reasonable request.

References

1 Helfield, J. M. & Naiman, R. J. Keystone Interactions: Salmon and Bear in

Riparian Forests of Alaska. *Ecosystems* **9**, 167-180, doi:10.1007/s10021-004-0063-5

(2006).

2 Smith, R. A. & Kennedy, M. L. Food habits of the racoon (*Procyon lotor*). *Journal of*

the Tennessee Academy of Science **62**, 79-82 (1987).

3   Rees, E. E. *et al.* Assessing a landscape barrier using genetic simulation modelling:

implications for raccoon rabies management. *Prev Vet Med* **86**, 107-123,

doi:10.1016/j.prevetmed.2008.03.007 (2008).

4   Kempf, E. Patterns of water use in primates. *Folia Primatol (Basel)* **80**, 275-294,

doi:10.1159/000252586 (2009).

5   Nowak, K., Barnett, A. A. & Matsuda, I. *Primates in Flooded Habitats: Ecology and*

Conservation. (Cambridge University Press, 2019).

6   Matsuda, I. *et al.* in *Primates in Flooded Habitats: Ecology and Conservation*

(eds Katarzyna Nowak, Adrian A. Barnett, & Ikki Matsuda) 15-28 (Cambridge

University Press, 2019).

7   van Schaik, C. P. & Mirmanto, E. Spatial variation in the structure and litterfall of

a Sumatran rain forest. *Biotropica* **17**, 196–205, doi:10.2307/2388217 (1985).

8   Coley, P. D. Interspecific variation in plant anti-herbivore properties: the role of

habitat quality and rate of disturbance. *New Phytologist* **106**, 251-263,

doi:10.1111/j.1469-8137.1987.tb04693.x (1987).

9   Chapman, C. A., Chapman, L. J., Naughton-Treves, L., Lawes, M. J. & McDowell, L.

R. Predicting folivorous primate abundance: validation of a nutritional model. *Am J*

Primatol **62**, 55–69, doi:10.1002/ajp.20006 (2004).

10   Matsuda, I., Tuuga, A., Bernard, H., Sugau, J. & Hanya, G. Leaf selection by two

Bornean colobine monkeys in relation to plant chemistry and abundance. *Scientific*

Reports **3**, 1873, doi:10.1038/srep01873 (2013).

11   Hanya, G., Kiyono, M., Takafumi, H., Tsujino, R. & Agetsuma, N. Mature leaf

selection of Japanese macaques: effects of availability and chemical content.

Journal of Zoology **273**, 140-147, doi:10.1111/j.1469-7998.2007.00308.x (2007).

12   Laundré, J. W., Hernández, L. & Altendorf, K. B. Wolves, elk, and bison:

reestablishing the "landscape of fear" in Yellowstone National Park, U.S.A.

Canadian Journal of Zoology **79**, 1401-1409, doi:10.1139/z01-094 (2001).

13   Altendorf, K. B., Laundré, J. W., López González, C. A. & Brown, J. S. Assessing

Effects of Predation Risk on Foraging Behavior of Mule Deer. *Journal of*

Mammalogy **82**, 430-439, doi:10.1644/1545-1542(2001)082<0430:aeopro>2.0.co;2

(2001).

14   Laundré, J. W., Hernandez, L. & Ripple, W. J. The landscape of fear: ecological

implications of being afraid. *The Open Ecology Journal* **3**, 1-7,

doi:10.2174/1874213001003030001 (2010).

15   Gautier-Hion, A. in *Comparative Ecology and Behaviour of Primates: Proceedings of*

a Conference held at the Zoological Society   (eds R.P. Michael & J.H. Crook)

147–170 (Academic Press, 1973).

16   van Schaik, C. P., van Amerongen, A. & van Noordwijk, M. A. in *Evolution and*

ecology of macaque societies   (eds J.A.   Fa & D.G. Lindburg)   160-181

(Cambridge University Press, 1996).

17   Matsuda, I., Tuuga, A. & Bernard, H. Riverine refuging by proboscis monkeys

(*Nasalis larvatus*) and sympatric primates: Implications for adaptive benefits of the

riverine habitat. *Mammalian Biology* **76**, 165-171,

doi:10.1016/j.mambio.2010.03.005 (2011).

18   Albert, A., Savini, T. & Huynen, M. C. Sleeping site selection and presleep behavior

in wild pigtailed macaques. *Am J Primatol* **73**, 1222-1230, doi:10.1002/ajp.20993

(2011).

19   Matsuda, I., Tuuga, A. & Higashi, S. Clouded leopard (*Neofelis diardi*) predation on

521   proboscis monkeys (*Nasalis larvatus*) in Sabah, Malaysia. *Primates* **49**, 227-231,

522   doi:10.1007/s10329-008-0085-2 (2008).

523 20 Otani, Y., Tuuga, A., Bernard, H. & Matsuda, I. Opportunistic predation and

524   predation-related events on long-tailed macaque and proboscis monkey in

525   Kinabatangan, Sabah, Malaysia. *Journal of Tropical Biology and Conservation* **9**,

526   214-218 (2012).

527 21 Matsuda, I., Tuuga, A., Akiyama, Y. & Higashi, S. Selection of river crossing location

528   and sleeping site by proboscis monkeys (*Nasalis larvatus*) in Sabah, Malaysia. *Am J*

529   *Primatol* **70**, 1097-1101, doi:10.1002/ajp.20604 (2008).

530 22 Ross, J., Hearn, A. J., Johnson, P. J. & Macdonald, D. W. Activity patterns and

531   temporal avoidance by prey in response to Sunda clouded leopard predation risk.

532   *Journal of Zoology* **290**, 96-106, doi:10.1111/jzo.12018 (2013).

533 23 Matsuda, I., Tuuga, A. & Higashi, S. Effects of water level on sleeping-site selection

534   and inter-group association in proboscis monkeys: why do they sleep alone inland

535   on flooded days? *Ecological Research* **25**, 475-482, doi:10.1007/s11284-009-0677-3

536   (2010).

537 24 Mittermeier, R. A., Rylands, A. B. & Wilson, D. E. *Handbook of the mammals of the*

538   *world. Vol. 3. Primates*. (Lynx Edicions, 2013).

539 25 Kurland, J. A. A natural history of kra macaques (*Macaca fascicularis* Raffles, 1821)

540   at the Kutai Reserve, Kalimantan Timur, Indonesia. *Primates* **14**, 245-262,

541   doi:10.1007/bf01730823 (1973).

542 26 Yeager, C. P. Feeding ecology of the long-tailed macaque (*Macaca fascicularis*) in

543   Kalimantan Tengah, Indonesia. *International Journal of Primatology* **17**, 51-62,

544   doi:10.1007/bf02696158 (1996).

27   Choudhury, A. Ecology and behaviour of the pigtailed macaque *Macaca nemestrina*

leonina in some forests of Assam in North-East India. *The journal of the Bombay*

Natural History Society **105**, 279-291 (2008).

28   Albert, A. *Feeding and ranging behavior of northern pigtailed macaques (Macaca*

leonina): impact on their seed dispersal effectiveness and ecological contribution in

a tropical rainforest at Khao Yai National Park, Thailand PhD. thesis thesis,

Université de Liège, (2012).

29   Caldecott, J. O. *An ecological and behavioural study of the pig-tailed macaque.* Vol.

21 262 (Karger, 1985).

30   Gazagne, E. *et al.* When Northern Pigtailed Macaques (*Macaca leonina*) Cannot

Select for Ideal Sleeping Sites in a Degraded Habitat. *International Journal of*

Primatology **41**, 614-633, doi:10.1007/s10764-020-00173-4 (2020).

31   Rodman, P. S. Structural differentiation of microhabitats of sympatric *Macaca*

fascicularis and *M. nemestrina* in East Kalimantan, Indonesia. *International*

Journal of Primatology **12**, 357-375, doi:10.1007/bf02547617 (1991).

32   Matsuda, I., Otani, Y., Bernard, H., Wong, A. & Tuuga, A. Primate Survey in a

Bornean Flooded Forest: Evaluation of Best Approach and Best Timing. *Mammal*

Study **41**, 101-106, doi:10.3106/041.041.0201 (2016).

33   Alexander, R. D. The evolution of social behavior. *Annual Review of Ecology and*

Systematics **5**, 325-383, doi:10.1146/annurev.es.05.110174.001545 (1974).

34   Cheney, D. L. & Wrangham, R. W. in *Primate societies*   (eds B.B Smuts *et al.*)

227–239 (University of Chicago Press, 1987).

35   Hill, R. A. & Lee, P. C. Predation risk as an influence on group size in cercopithecoid

primates: implications for social structure. *Journal of Zoology* **245**, 447-456,

doi:10.1111/j.1469-7998.1998.tb00119.x (1998).

36   Oi, T. Population organization of wild pig-tailed macaques (*Macaca nemestrina*

nemestrina) in West Sumatra. *Primates* **31**, 15-31, doi:10.1007/bf02381027 (1990).

37   Sackett, G. P., Ruppenthal, G. C. & Davis, A. E. Survival, growth, health, and

reproduction following nursery rearing compared with mother rearing in pigtailed

monkeys (*Macaca nemestrina*). *Am J Primatol* **56**, 165-183, doi:10.1002/ajp.1072

(2002).

38   Gouzoules, H. & Gouzoules, S. Body size effects on the acoustic structure of pigtail

macaque (*Macaca nemestrina*) screams. *Ethology* **85**, 324-334, doi:10.1111/j.1439-

0310.1990.tb00411.x (2010).

39   Matsuda, I. *et al.* The nose is mightier than the tooth: larger male proboscis

monkeys have smaller canines. *bioRxiv*, doi:10.1101/848515 (2019).

40   Matsuda, I., Tuuga, A. & Higashi, S. The feeding ecology and activity budget of

proboscis monkeys. *American journal of primatology* **71**, 478-492,

doi:10.1002/ajp.20677 (2009).

41   Matsuda, I., Kubo, T., Tuuga, A. & Higashi, S. A Bayesian analysis of the temporal

change of local density of proboscis monkeys: implications for environmental effects

on a multilevel society. *Am J Phys Anthropol* **142**, 235-245, doi:10.1002/ajpa.21218

(2010).

42   Altmann, J. Observational study of behavior: sampling methods. *Behaviour* **49**,

227–267, doi:10.1163/156853974x00534. (1974).

43   Kumar, D., Kumar, S., Gupta, J., Arya, R. & Gupta, A. A review on chemical and

biological properties of *Cayratia trifolia* Linn. (Vitaceae). *Pharmacogn Rev* **5**, 184-

188, doi:10.4103/0973-7847.91117 (2011).

44   Cleveland, R. B., Cleveland, W. S., McRae, J. E. & Terpenning, I. STL: a seasonal-

trend decomposition procedure based on loess. *Journal of official statistics* **6**, 3-73

(1990).

45   Driver, C. C., Oud, J. H. L. & Voelkle, M. C. Continuous time structural equation

modeling with R Package ctsem. *Journal of Statistical Software* **77**,

doi:10.18637/jss.v077.i05 (2017).

46   Driver, C. C. & Voelkle, M. C. Hierarchical Bayesian continuous time dynamic

modeling. *Psychol Methods* **23**, 774-799, doi:10.1037/met0000168 (2018).

47   R: a language and environment for statistical computing (Foundation for Statistical

Computing, Vienna, 2019).

48   Tsuji, Y., Hanya, G. & Grueter, C. C. Feeding strategies of primates in temperate

and alpine forests: comparison of Asian macaques and colobines. *Primates*,

doi:10.1007/s10329-013-0359-1 (2013).

49   Campbell, C. J., Fuentes, A., MacKinnon, K. C., Bearder, S. K. & Stumpf, R. M.

Primates in perspective.   852 (Oxford University Press, 2011).

50   Smith, J. A., Donadio, E., Pauli, J. N., Sheriff, M. J. & Middleton, A. D. Integrating

temporal refugia into landscapes of fear: prey exploit predator downtimes to forage

in risky places. *Oecologia* **189**, 883-890, doi:10.1007/s00442-019-04381-5 (2019).

51   Miller, L. E. & Treves, A. in *Primates in perspective*   (eds C.J. Campbell *et al.*)

525–543 (Oxford University Press, 2007).

52   Galdikas, B. M. F. & Yeager, C. P. Brief report: Crocodile predation on a crab-eating

macaque in Borneo. *Am J Primatol* **6**, 49-51, doi:10.1002/ajp.1350060106 (1984).

53   van Schaik, C. P., van Noordwijk, M. A., Warsono, B. & Sutriono, E. Party size and

early detection of predators in sumatran forest primates. *Primates* **24**, 211-221,

doi:10.1007/bf02381083 (1983).

54   Khamcha, D. & Sukumal, N. Burmese python (*Python molurus*) predation on a pig-

tailed macaque (*Macaca nemestrina*) in Khao Yai National Park. *Hamadryad* **34**,

176-178, doi:10.13140/2.1.3857.7449 (2009).

55   van Schaik, C. & Mitrasetia, T. Changes in the behaviour of wild long-tailed

macaques (*Macaca fascicularis*) after encounters with a model python. *Folia*

Primatol (Basel) **55**, 104-108, doi:10.1159/000156506 (1990).

56   Terborgh, J. Mixed flocks and polyspecific associations: Costs and benefits of mixed

groups to birds and monkeys. *Am J Primatol* **21**, 87-100,

doi:10.1002/ajp.1350210203 (1990).

57   Hart, D. in *Primate Anti-Predator Strategies, Developments in Primatology:*

Progress and Prospects   (eds S.L. Gursky & K.A.I. Nekaris)   27–59 (Springer US,

2007).

58   Fam, S. D. & Nijman, V. *Spizaetus* hawk-eagles as predators of arboreal colobines.

Primates **52**, 105-110, doi:10.1007/s10329-011-0240-z (2011).

59   Prawiradilaga, D. M. Ecology and conservation of endangered Javan Hawk-eagle

Spizaetus bartelsi. Ornithological Science **5**, 177-186, doi:10.2326/1347-

0558(2006)5[177:eacoej]2.0.co;2 (2006).

60   Yeager, C. P. Possible antipredator behavior associated with river crossings by

proboscis monkeys (*Nasalis larvatus*). *American Journal of Primatology* **24**, 61-66,

doi:10.1002/ajp.1350240107 (1991).

61   van Schaik, C. P. & van Hooff, J. A. R. A. M. On the ultimate causes of primate

social systems. *Behaviour* **85**, 91-117, doi:10.1163/156853983x00057 (1983).

62   Anderson, J. R. Sleep, sleeping sites, and sleep-related activities: Awakening to

their significance. *American Journal of Primatology* **46**, 63–75,

doi:10.1002/(sici)1098-2345(1998)46:1<63::aid-ajp5>3.0.co;2-t (1998).

63 Chapman, C. A., Chapman, L. J. & McLaughlin, R. L. Multiple central place

foraging by spider monkeys: travel consequences of using many sleeping sites.

Oecologia **79**, 506-511, doi:10.1007/BF00378668 (1989).

64 Teichroeb, J. A., Holmes, T. D. & Sicotte, P. Use of sleeping trees by ursine colobus

monkeys (*Colobus vellerosus*) demonstrates the importance of nearby food.

Primates **53**, 287-296, doi:10.1007/s10329-012-0299-1 (2012).

65 Vessey, S. H. Night observations of free-ranging Rhesus monkeys. *Am J Phys*

Anthropol **38**, 613-619, doi:10.1002/ajpa.1330380276 (1973).

66 Dasilva, G. L. Postural changes and behavioural thermoregulation in *Colobus*

polykomos: the effect of climate and diet. *African Journal of Ecology* **31**, 226-241,

doi:10.1111/j.1365-2028.1993.tb00536.x (1993).

67 Rodman, P. S. Skeletal differentiation of *Macaca fascicularis* and *Macaca*

nemestrina in relation to arboreal and terrestrial quadrupedalism. *American*

Journal of Physical Anthropology **51**, 51-62, doi:10.1002/ajpa.1330510107 (1979).

Table and Figure legends

Table 1. Summary of detection events and the number of individuals sightings during

boat-based surveys (n = 434 days).

Table 2. Food items and parts consumed by southern pig-tailed and long-tailed

macaques with their observed frequency during boat-based surveys.

Table 3. Means, standard deviations (SD) and posterior credibility intervals (PCI) of the

CtSEM model. Asterisks (*) indicate that zero does not falls within the PCIs, i.e.,

positive or negative effects are indicated.

Figure 1. Decomposition plot of abundance of *Cayratia trifolia* fruits based on Seasonal

decomposition of time series by loess (STL). Numbers of *C. trifolia* fruits are

represented on the y-axis. Trend and seasonality indicate a relatively steady

increase or decrease over time, and a pattern that repeats, respectively.

Figure 2. Auto-regressive effects plotted for time intervals of $0 < \Delta t < 10$ months.

Parameters represent within-section persistence of the number of *Cayratia trifolia*

fruits, and the sighting frequency of pig-tailed and long-tailed macaques over time.

Solid lines represent auto-regressive effects for the number of *C. trifolia* fruits

over time, and dashed lines represents auto-regressive effects for the sighting

frequency of pig-tailed (a) and long-tailed macaques (b).

Table 1. Summary of detection events and the number of individual sightings during boat-based surveys (n = 434 days).

	Pig-tailed macaque	Long-tailed macaque	Proboscis monkey	Silver langur	Maroon langur	Orangutan
Total number of discovery event (times)	442	1357	1278	44	1	67
Total number of individual sighting (head)	10257	14785	14393	344	6	122
Averaged individual sighting (head/day ±SD)	23.63 ±24.07	34.07 ±23.05	33.16 ±23.86	0.79 ±2.67	0.01 ±0.29	0.28 ±0.91

Table 2. Food items and parts consumed by southern pig-tailed and long-tailed macaques with their observed frequency during boat-based surveys.

Species	Type	Pig-tailed macaque		Long-tailed macaque	
		Number of observation time	%	Number of observation time	%
Albizia corniculata	fruit	1	1.52		
	unkown	1	1.52		
Antidesma thwaitesianum	flower			10	3.61
	fruit			2	0.72
	leaf			1	0.36
Baccaurea stipulata	fruit	1	1.52		
Cayratia trifolia	fruit	15	22.73	141	50.90
	leaf			9	3.25
Dillenia excelsa	flower	24	36.36	45	16.25
	fruit	4	6.06	7	2.53
	leaf			2	0.72
	unkown			1	0.36
Eichhornia crassipes	stem			2	0.72
Ficus sp.	fruit	2	3.03	4	1.44
Gnetum gnemonoides	fruit			1	0.36
Mallotus muticus	flower	1	1.52	3	1.08
	fruit	3	4.55	8	2.89
	leaf	1	1.52	7	2.53
	unkown			1	0.36
Nauclea subdita	flower			1	0.36
	leaf	1	1.52		
Pternandra galeata	fruit	1	1.52	2	0.72
	leaf			2	0.72
Spatholobus cf. macropterus	leaf			1	0.36
Vitex pinnata	fruit			1	0.36
Xylosma sumatrana	flower			5	1.81
Ziziphus bornensis	fruit	1	1.52		
unkown	flower	6	9.09	16	5.78
	fruit	2	3.03	4	1.44
	leaf	2	3.03	1	0.36
Total		66		277	

Table 3. Means, standard deviations (SD) and posterior credibility intervals (PCI) of the CtSEM model.

Parameter	Mean number of pig-tailed			Mean number of long-tailed			Amount of *Cayratia trifolia* fruits		
	mean	SD	PCI [2.5%, 97.5%]	mean	SD	PCI [2.5%, 97.5%]	mean	SD	PCI [2.5%, 97.5%]
T_0 mean	0.408	0.253	-0.089, 0.911	0.313	0.251	-0.199, 0.783	0.270	0.262	-0.230, 0.805
Manifest means	-0.021	0.129	-0.277, 0.233	-0.030	0.078	-0.192, 0.188	-0.011	0.100	-0.211, 0.184
Drift Parameters									
Mean number of pig-tailed		–		0.984	0.367	0.356, 1.802 *	0.557	0.446	-0.264, 1.486
Mean number of long-tailed	0.753	0.281	0.272, 1.373 *		–		0.101	0.215	-0.309, 0.527
Amount of *Cayratia trifolia* fruits	1.560	0.653	0.131, 2.710 *	0.850	0.148	0.581, 1.167 *		–	
Effect of time dependent parameters									
Rainfall	0.008	0.054	-0.098, 0.115	-0.022	0.049	-0.119, 0.075	0.020	0.053	-0.084, 0.124
Water level	-0.179	0.052	-0.282, -0.077 *	-0.110	0.049	-0.205, -0.011 *	-0.174	0.053	-0.278, -0.071 *
Effect of time independent parameters									
River width	-0.466	0.257	-0.971, 0.045	-0.155	0.259	-0.638, 0.365	-0.225	0.258	-0.779, 0.245

Asterisks (*) indicate that zero does not falls within the PCIs, i.e., positive or negative effects are indicated.

Fig. 1

Fig. 2a

Fig. 2b