Type: Original Article

Herkogamy but not reciprocity is a better predictor of legitimate pollen transfer and

fruit set in *Jasminum malabaricum,* a self-compatible species with stigma height

dimorphism.

Shatarupa Ganguly*, Deepak Barua

Department of Biology, Indian Institute of Science Education and Research, Pune, India

*Author for correspondence

Email: shatarupa.ganguly23@gmail.com

Running title: Herkogamy but not reciprocity predicts legitimate pollen load and fruit set

Abstract

Background and aims: Reciprocity, the match between anther and stigma positions of

complementary morphs, and herkogamy, the separation between anthers and stigmas within a

flower, are morphological traits that define style length polymorphisms. While these traits are

thought to be critical in determining legitimate pollen transfer, the consequences of individual

level variation in reciprocity and herkogamy have rarely been examined. Additionally,

changes in sex organ positions within a flower will affect stigma-anther separation, but will

likely also alter the match of these organs with counterparts in the complementary morph.

Thus, variation in herkogamy is likely related to reciprocity, but the relationship between

these two fundamental traits remains unexplored.

Methods: We quantified reciprocity and herkogamy in individuals from naturally occurring

populations of *Jasminum malabaricum*, a species with stigma-height dimorphism. To

understand how variation in these two fundamental traits are related to legitimate pollination

we examined the relationship between reciprocity and herkogamy of individuals and

stigmatic pollen load, legitimate pollen fraction, and reproductive success. Additionally, we

examined the relationship between herkogamy and reciprocity to understand potential

consequences for avoiding self-pollen deposition and encouraging legitimate pollen

deposition.

Key Results: Contrary to expectations, reciprocity was not related to total pollen load,

legitimate pollen fraction or reproductive success. Herkogamy was negatively related to total

stigma load, but positively related to legitimate pollen fraction and fruit set in the long-styled

morph. Interestingly, we observed a negative relationship between herkogamy and reciprocity

in the long-styled morph.

Conclusions: Herkogamy was more important than reciprocity in this species, and likely

plays a significant role in stabilizing such polymorphic intermediates, allowing reciprocal

arrangements between complementary sex-organs to arise later in the evolutionary pathways

towards heterostyly. The negative relationship between herkogamy and reciprocity suggests

potential trade-offs between avoidance of self-pollen deposition and promotion of legitimate

pollen deposition.

Keywords: *Jasminum malabaricum* Wight*.,* herkogamy, reciprocity**, s**tigma-height

dimorphism; self-pollination, legitimate pollen deposition; fruit set; intra-population variation

Introduction

Reciprocity and herkogamy are the two defining morphological features in flowers of species

with style length polymorphisms. Reciprocity is the physical match between complementary

sex organ heights (Armbruster *et al.* 2017), while herkogamy is the separation of the male

and female sex organs within a flower (Webb and Lloyd 1986; Opedal 2018). However,

while reciprocity and herkogamy are widely assumed to play a critical role in the promotion

of legitimate pollen transfer that is required to maintain style length polymorphisms, few

studies have examined how variation in these traits is related to inter-morph pollen transfer

and reproductive success in plants with style length polymorphisms (Cesaro *et al.* 2004;

Thompson *et al.* 2012; Keller *et al.* 2014; Liu *et al.* 2016; Jacquemyn *et al.* 2018; Brys and

Jacquemyn 2020).

The reciprocal arrangement of sex organs in species with style length polymorphism enables

pollinators to pick up pollen from anthers on specific locations on their body from where it

can be accurately deposited on stigmas of the complementary morphs (Lloyd and Webb

1992). Empirical studies in heterostylous species have focused on population level estimates

of reciprocity, and these have shown that higher reciprocity is related to greater legitimate

pollen transfer (Lau and Bosque 2003; Baena-Díaz *et al.* 2012), and reproductive success

(Valois-Cuesta *et al.* 2011; Brys and Jacquemyn 2014; Keller *et al.* 2014; Zhou *et al.* 2015;

Ferrero *et al.* 2017). While limited, studies in species with stigma-height dimorphism have

also shown that population level estimates of reciprocity are positively related to reproductive

success (Thompson *et al.* 2003, 2012; Stehlik *et al.* 2006). There are no studies that have

examined individual level reciprocity in species with stigma-height dimorphism, and only

two in species with heterostyly (Jacquemyn *et al.* 2018; Brys and Jacquemyn 2020) that show

that individuals with higher reciprocity have higher proportions of legitimate pollen.

The presence of self and heteromorphic incompatibility in most species with style length

polymorphism makes it difficult to tease apart the contribution of reciprocity alone to

legitimate pollination and maintenance of the polymorphism. Additionally, even in the

presence of physiological incompatibility, reduced male and female fitness due to pollen

discounting and stigma clogging, and associated reductions in reproductive success, have

been observed (Shore and Barrett 1984; Barrett and Glover 1985; Martén-Rodríguez *et al.*

2013; Keller *et al.* 2014). In self-compatible species higher legitimate pollen fractions do not

guarantee greater legitimate offspring, and substantial numbers of illegitimate offspring can

result despite a higher legitimate fraction of pollen (Zhou *et al.* 2015). In such cases, the role

of herkogamy in ensuring more legitimate offspring becomes very important.

The function of herkogamy is to increase the fraction of legitimate pollen deposition on the

stigma by reducing interference between male and female organs, and reducing self and intra-

morph pollen deposition (Webb and Lloyd 1986; Barrett 2002). Herkogamy has been studied

extensively in monomorphic species that do not exhibit style length polymorphisms, and

results from these have shown that greater stigma-anther separation between sex organs

within flowers results in higher outcrossing rates (Ennos 1981; Brunet and Eckert 1998;

Takebayashi *et al.* 2006; Larrinaga *et al.* 2009; de Vos *et al.* 2012, 2018; Li *et al.* 2013).

However, empirical studies in species with style length polymorphisms are few, and these

show that herkogamy is related to decreased self-pollination (Nishihiro and Washitani 1998),

as well as increased legitimate pollen transfer and reproductive success in heterostylous

species (Liu et al. 2016). However, as with reciprocity, these studies focused on population

level estimates of herkogamy while ignoring intra-population variation in herkogamy (but see

Hernandez and Ornelas 2007, Keller et al. 2014). Population level estimates of herkogamy

were also observed to be associated with decreased self-pollination in species with stigma-height dimorphism (Cesaro *et al.* 2004). Again, as with reciprocity, studies on consequences of intra-population variation in herkogamy are lacking in species with stigma-height dimorphism.

It is widely acknowledged that there can be large variation in sex organ positions among individuals in populations with style length polymorphism, and this variation can have important consequences for reciprocity, herkogamy, pollen flow, and reproductive success for these individuals (Eckert and Barrett 1994; Nishihiro *et al.* 2000; Sanchez *et al.* 2008; Sampson and Krebs 2012; Costa *et al.* 2017). Population level estimates of reciprocity and herkogamy can hide biologically important variation among individuals, and understanding intra-population variation in these traits is important as selection ultimately acts at the level of the individuals. Additionally, changes in organ positions in an individual flower have obvious implications for stigma-anther separation but are likely to also alter the match of these organs with counterparts in the complementary morph and affect reciprocity for that individual. However, the relationship between these two fundamental traits remain unexplored and the consequences for the two distinct functions in avoiding self-pollen deposition and encouraging legitimate pollen deposition remain unknown. The lack of an individual-level reciprocity measure (Jacquemyn *et al.* 2018; Brys and Jacquemyn 2020) and the focus on population-level estimates for both reciprocity and herkogamy have precluded such studies.

In this study, we examined the variation in reciprocity and herkogamy among individuals of four naturally occurring isoplethic populations of *J. malabaricum* (Ganguly and Barua 2020) with stigma-height dimorphism. We estimated reciprocity for an individual as the average mismatch between the stigma of that individual and the anthers of all other individuals of the

complementary morph. We investigated the consequences of intra-population variation in

reciprocity and herkogamy for legitimate pollen transfer and reproductive success by

examining the relationship of individual reciprocity and herkogamy with stigmatic pollen

load and fruit set. Morph specific differences in stigmatic pollen load and reproductive

success were also quantified to aid in the interpretation of the results as they can influence the

above-mentioned relationships. We expected reciprocity to play an important role in

legitimate pollen transfer and fruit set. However, given that our study species exhibits stigma-

height dimorphism with low reciprocity and self-compatibility, we also expected herkogamy

to significantly influence legitimate pollen transfer. Additionally, we examined the

relationship between reciprocity and herkogamy across individuals to understand potential

consequences for avoiding self-pollen deposition and encouraging legitimate pollen

deposition.

Methods

<u>Study sites and species</u>: *Jasminum malabaricum* Wight. is a woody liana endemic to the

Western Ghats of India, and Sri Lanka (Singh & Karthikeyan 2000) that exhibits stigma-

height dimorphism (Ganguly 2019). This study was conducted between 2015 and 2017 on

four populations of *J. malabaricum*: namely, Trimbak (19.94° N, 73.54° E), Bhimashankar

(19.07° N, 73.55° E), Mulshi (18.50° N, 73.51° E) and Kaas (17.72° N, 73.81° E).

Pollen transfer and reproductive success can be influenced by morph specific differences in

various factors affecting mating opportunities. In all four study populations, the short-styled

morph had higher herkogamy, while reciprocity was higher between the long-styled stigmas

and the short-styled anthers (Ganguly and Barua 2020). With the exception of these two

traits, morphs did not differ in spatiotemporal mate availability, pollinator visitation, and did

not exhibit physiological incompatibility (Ganguly and Barua 2020). Since the mating

opportunities were similar for both morphs it is unlikely to be a confounding factor in our

investigations. The species exhibits pollen size dimorphism (Ganguly and Barua 2020) which

is rare in species with stigma-height dimorphism.

<u>Quantification of herkogamy and reciprocity</u>: As reported in a previous study (Ganguly and

Barua 2020), one flower from 30 individuals of each morph was collected from

Bhimashankar, Mulshi and Kaas, between March and April of 2015 and 2017, and preserved

in formalin acetic alcohol (FAA; 2.5%:2.5%:95%). In Trimbak, one flower was collected

from 20 long-, and 19 short-styled individuals because of the small population size and

limited availability of individuals. Herkogamy and reciprocity were also quantified for an

additional 70 individuals for a total of 100 individuals of each morph in the Bhimashankar

population. Anther and stigma heights were quantified using ImageJ (ver. 1.52a) from

scanned images of dissected flowers. Since anthers were attached to the corolla tube, the

distance from the base of the corolla tube to the mid-point of the anthers was measured as

anther height. The distance from the base of the ovary to the mid-point of the stigma was

measured as stigma height. Herkogamy was quantified for one flower per individual as the

difference in anther and stigma height of a flower (Opedal 2018). As we examined the

relationship of reciprocity with measures of female fitness like pollen load and fruit set,

reciprocity was calculated for one flower per individual as the mean of mismatches between

stigma height for that individual and anther height of every other individual of the

complementary morph. These pair-wise mismatches were averaged to give an estimate of

reciprocity for that individual (Ganguly and Barua 2020). We calculate mismatch, which is

the inverse of reciprocity and hence higher values denote lower reciprocity. The relationship

between herkogamy and reciprocity was examined using Pearson's correlation coefficient.

We use the term "mismatch" to refer to the spatial correspondence between anther and stigma

heights of reciprocal morphs to describe our results. However, we use the term "reciprocity"

to discuss the results to increase ease of interpretation.

<u>Quantification of stigma pollen load:</u> We quantified stigma pollen load for the same

individuals from the Bhimashankar (March 2016) and Kaas (April 2017) populations for

which floral morphometry was quantified. Flowers of *J. malabaricum* open between 7 p.m. to

9 p.m. and last for approximately 33 h (Ganguly and Barua 2020). Post 3 p.m. on the day

after anthesis, the flowers wilt and no pollinator activity is seen on these flowers (Ganguly

and Barua 2020). Hence, to allow for maximal pollen deposition, flowers were collected

between 2 and 3 p.m. the day after anthesis and preserved in formalin acetic alcohol (FAA;

2.5%:2.5%:95%). The quantification was done for 30 long- and 29 short-styled individuals

for Bhimashankar, and 32 long-, and 30 short-styled individuals for Kaas. Stigmas were

washed in 70% ethanol and softened in 4N NaOH for 12 h (Kearns and Inouye 1993).

Subsequently, they were washed in distilled water and passed through serial concentrations of

ethanol from 10% to 100% in steps of 15% to wash the oil from the pollen surface. The

stigmas were then resuspended in a series of ethanol dilutions of glycerol with increasing

concentration of glycerol from 10% to 90% in steps of 20%, squashed with a coverslip, and

washed and mounted in 100% glycerol.

To quantify the total number of pollen deposited on a stigma, images of stigma squashes for

two flowers per individual were taken at 40x magnification. An exhaustive pollen count was

performed for all the stigmas. Legitimate pollen fraction was estimated for one stigma for

each individual. Ten images were taken at 400x magnification for each stigma, and this

accounted for greater than 15% of the total pollen on that stigma. Both the pollen count and

measurement of pollen size were performed using ImageJ. As the equatorial and polar

diameters were strongly correlated, only the equatorial diameter of the pollen was used as a

measure of pollen size. The pollen were discriminated based on differences in diameter

between the morphs using the estimates of pollen size obtained previously. The short-styled

morph produced larger pollen but pollen production was not significantly different between

morphs (Ganguly and Barua 2020). The diameter of the long-styled pollen ranged between 34

µm to 55 µm and the short-styled pollen between 47 µm to 73 µm (Ganguly and Barua

2020). Approximately 35% of the pollen were in the range from 47 µm to 55 µm, which was

common to both morphs. The ratio of the frequency of pollen grains extracted from anthers of

the long- and short-styled morph in each size class of 1 µm was used to estimate the

probability of occurrence of pollen from each morph in this overlapping size range. The

pollen from the stigma which belonged to these size classes were distributed among the

morphs based on this probability of occurrence. The legitimate pollen fraction was calculated

as the percentage of the number of legitimate pollen identified in the total number of pollen

sampled. Differences between morphs in natural total stigma pollen load and the legitimate

pollen fraction were examined using a non-parametric Mann-Whitney U test as the values

were not normally distributed.

<u>Autogamous pollen deposition:</u> To investigate the function of herkogamy in decreasing self-

pollination, three mature buds were tagged and emasculated for five individuals of each

morph to quantify autogamous self-pollination in March 2017 in individuals from the

Bhimashankar population. Another three buds on the same individuals were tagged as

controls to account for individual differences in pollen deposition. Flowers were collected

between 2 p.m. and 3 p.m. on the day after anthesis. The total pollen load and the legitimate

fraction of pollen were quantified as described above. Wilcoxon's signed-rank test was used

to examine differences in total pollen load and the legitimate fraction between the control and

the emasculated flowers from the same individual. This non-parametric test was chosen as the

variables contained extreme values and so that control and treatment comparisons within the

same individual could be made.

<u>Quantification of fruit set</u>: Given the time-, and labour-intensive nature of this work, fruit set

was quantified in March 2015 and 2016 in only one of the four study populations

(Bhimashankar). This was done for the same individuals for which we had quantified

herkogamy and reciprocity and estimated stigma pollen load. This included 30 long- and 29

short-styled individuals in 2015, and 29 individuals of each morph in 2016. For both morphs,

20 mature buds were tagged per individual in the peak flowering season and followed till the

fruits matured. The percentage of total mature fruits formed from the 20 buds was taken as

the final fruit set. Differences in natural fruit set between the two morphs were examined

using a non-parametric Mann-Whitney U test as the values obtained were not normally

distributed. The relationships of herkogamy and reciprocity with the natural total pollen load,

the legitimate fraction of stigmatic pollen load and the fruit set were examined using

Spearman's rank correlation coefficient as most of the examined variables were counts and

percentages, and had extreme values and skewed distributions.

Results

There was substantial intra-population variation in estimates of mismatch and herkogamy

between individuals in the populations (Fig. 1, Table 1). The long-styled morph exhibited a

much larger range and higher coefficient of variation in mismatch as compared to the short-

styled morph in all four study populations (Table 1). Similarly, the long-styled morph

exhibited a larger range and higher coefficient of variation in herkogamy as compared to the

short-styled morph (Table 1).

There was high variation between individuals in natural total stigma pollen load, with counts

ranging from 13 to 5015 in the long-styled, and 38 to 1394 in the short-styled morph in the

Bhimashankar population, and from 6 to 1542 in the long-styled and 1 to 1350 in the short-

styled morph (Fig. 2 a, c) in the Kaas population. Although long-styled morphs had higher

mean stigma pollen load, no significant differences were detected between the two morphs in

either of the two populations examined (Fig. 2 a, c; Mann-Whitney U test: Bhimashankar $p =$

0.41 and Kaas $p = 0.11$; $n = 30$ and 29 individuals for Bhimashankar, and 32 and 30

individuals for Kaas for the long- and short-styled morphs). The legitimate pollen fraction

was significantly higher in the short- than in the long-styled morph for both populations (Fig

2 b, d; Mann-Whitney U test: Bhimashankar $p < 0.001$ and Kaas $p < 0.001$; n for long-styled

and short-styled morph was 30 and 29 individuals for Bhimashankar, and 32 and 30

individuals for Kaas).

In both morphs, emasculation resulted in a decrease in the total pollen deposition (Fig. 3 a;

Wilcoxon signed-rank test: long-styled $p = 0.04$ and short-styled $p = 0.04$; 3 flowers per

treatment for 5 individuals of each morph). In the short-styled morphs emasculation did not

affect the legitimate pollen fraction (Fig. 3 b; Wilcoxon signed-rank test: $p = 0.34$). In

contrast, in the long-styled morphs emasculation resulted in an increase in the legitimate

pollen fraction (Wilcoxon signed-rank test: $p = 0.04$). Fruit set was not significantly different

between the long-, and short-styled morphs, and this was consistent across the two years in

which we examined this in individuals from the Bhimashankar population (Fig. 4; Mann-

Whitney U test: year 2015 $p = 0.52$, $n = 30$ long- and 29 short-styled individuals; year 2016 p

$= 0.80$, $n = 29$ individuals of each morph).

Mismatch was not related to total pollen deposition or the legitimate pollen fraction. Counter-

intuitively, mismatch was positively related to fruit set in the short-styled morph in

Bhimashankar (Table 2 a). Herkogamy, on the other hand, was positively related to

disassortative pollen load and fruit set in the long-styled morph, but not in the short-styled

morph in Bhimashankar (Table 2 a). Herkogamy was negatively related to total pollen load in

the long-styled morph, but not related to pollen deposition in the short-styled morph in Kaas

(Table 2 b).

Herkogamy and mismatch showed a significant positive correlation in the long-styled morph

in all study populations (Fig. 5 a-d). However, such a positive relationship between

herkogamy and mismatch was seen in only one of the four study populations in the short-

styled morph (Fig. 5 e-h). Similar results were observed when examining this relationship in

a larger number of individuals from the Bhimashankar population with a significant positive

relationship between herkogamy and mismatch in the long-, but not the short-styled morph

(Supplementary Information Fig. S1 a, b).

Discussion

Surprisingly, individual level variation in reciprocity was not related to legitimate pollen

fraction or fruit set. In contrast, individuals of the long-styled morph with higher herkogamy

had a higher fraction of legitimate pollen and greater fruit set. Thus, herkogamy may be

relatively more important than reciprocity in increasing legitimate pollen deposition and

reproductive success in *Jasminum malabaricum*, a species with stigma height dimorphism

that lacks self-, and heteromorphic incompatibility. Additionally, we report here for the first

time a negative relationship between reciprocity and herkogamy, the two defining traits of

style length polymorphism. This suggests a potential trade-off between increasing legitimate

pollen deposition and avoidance of self-pollen deposition.

Intra-population variation in anther and stigma heights is expected to reduce reciprocity and

hence legitimate pollen transfer and reproductive success in species with style length

polymorphism (Sanchez *et al.* 2008; Armbruster *et al.* 2017). We observed large variation in

anther and stigma positions, and in the resultant reciprocity in individuals of all populations

of our study species. However, this observed variation in reciprocity was not associated with

any negative consequences and was unrelated to the legitimate fraction of pollen load or fruit

set. Previous studies have shown that higher herkogamy but not reciprocity was related to

higher legitimate pollen transfer and reproductive success (Keller *et al.* 2014). Additionally,

legitimate pollen transfer in some species with style length polymorphism is likely to be

primarily governed by effective long-tongued pollinators rather than sex organ positions

(Simón-Porcar *et al.* 2014).

The higher fraction of legitimate pollen received by the short-styled morph is likely due to

reduced self-pollen deposition as a consequence of higher herkogamy in this morph (Liu *et*

al. 2016). High legitimate pollen transfer has also been associated with efficient long-tongued

pollinators (Haddadchi 2013; Simón-Porcar *et al.* 2014) that are capable of reaching the

short-styled stigma. However, the higher legitimate pollen fractions observed in the short-

styled morph did not translate into higher fruit set. As both morphs lack physiological

incompatibility, similar total pollen loads resulted in equal fruit set between morphs. This is

congruent with an observation from a previous study, where despite much higher legitimate

pollen deposition, a considerable number of illegitimate offspring are formed due to the

absence of self- and heteromorphic incompatibility (Zhou *et al.* 2015).

In the long-styled morph, emasculation resulted in a decrease in total pollen load and an

increase in the legitimate pollen fraction. Thus, reduced self pollen deposition with increasing

herkogamy is likely to result in the negative relationship between herkogamy and total pollen

load, and the positive relationship between herkogamy and the legitimate pollen fraction

that was observed in the long-styled morph. The higher fruit set associated with higher

herkogamy in the same morph could be attributed to lower mechanical interference between

the male and female sex organs. The lack of any change in the legitimate pollen fraction with

emasculation in the short-styled morph implies that the higher herkogamy observed in this

morph was sufficient to avoid self-pollen deposition. This was also reported in *Persicaria*

japonica where emasculation had no effect on the deposition of incompatible pollen

(Nishihiro and Washitani 1998). This likely explains the absence of a relationship between

herkogamy and reproductive success in the short-styled morph, where due to the higher

herkogamy observed in this morph minor changes in herkogamy did not influence self-pollen

deposition or fruit set. Similar results were reported in *Primula* spp., where the legitimate

pollen fraction increased with herkogamy in the long-, but not the short-styled morph, and the

"threshold" effect as mentioned above was offered as a possible explanation (Keller *et al.*

2014).

Reciprocity was negatively related to herkogamy in individuals of the long-styled morph.

This relationship suggests a potential trade-off between maximizing legitimate pollen transfer

and avoiding self-pollination and will be useful in providing insights into the response of

these species under different ecological scenarios. For example, in situations of pollen

limitation due to the scarcity of mate and pollinators, and an absence of intra-morph

incompatibility or inbreeding depression (Baker *et al.* 2000), reproductive assurance is

important. If the relationship between reciprocity and herkogamy is negative, the individuals

with high reciprocity and low herkogamy will be selected for as they would have the highest

reproductive assurance in the population. A positive relationship can be maladaptive in such a

situation. Hence, such relationships can help us understand the change in the distribution of

anther and stigma heights in a population under different ecological scenarios. There was no

relationship between reciprocity and herkogamy in individuals of the short-styled morph with

the exception of the Mulshi population where, similar to the long-styled individuals, a

negative relationship was observed. The difference in the relationship between reciprocity

and herkogamy between morphs and among populations indicates the effect of morph

specific selection pressures and differences in local environmental factors among populations

(Santos-Gally *et al.* 2013). The short-styled morph does not show a relationship even at a

higher sample size implying that the lack of a relationship is unlikely due to insufficient

sample sizes.

Several studies have highlighted the functional significance of sex organ positions in species

with style length polymorphism (Faivre and McDade 2001; Sosenski *et al.* 2010; V. Ferrero

et al. 2011; Victoria Ferrero *et al.* 2011; Keller *et al.* 2012; Yuan *et al.* 2017), but the

emphasis of these has primarily been on reciprocity while herkogamy has been relatively

understudied (but see Nishihiro and Washitani 1998, Keller et al. 2014, Liu et al. 2016,

Faife-Cabrera et al. 2016, Barranco et al. 2019). Moreover, although it has been

acknowledged that intra-population variation in sex organ positions affects legitimate pollen

transfer and reproductive success (Eckert and Barrett 1994; Lau and Bosque 2003; Sanchez *et*

al. 2008; Armbruster *et al.* 2017), studies examining this are rare. This study reveals that

herkogamy can be more important than reciprocity in promoting legitimate pollen transfer

and fruit set. The result highlights the functional role of herkogamy and has especially

important implications for the maintenance of imperfectly reciprocal intermediates in the

pathway towards heterostyly. The results also suggest that intra-population variation in

herkogamy and reciprocity might not affect legitimate pollen transfer and reproductive

success across individuals and is perhaps the reason why species with style length

polymorphisms display high variation in sex organ positions. Examination of the intra-

population variation in reciprocity and herkogamy also allowed us to investigate the

relationship between the two which revealed a trade-off between self-pollination and

legitimate pollination. Further studies on the causes and consequences of intra-population

variation in reciprocity and herkogamy can substantially increase our understanding of the

coevolution of the dimensions of floral parts with pollinators.

Author contributions

SG and DB conceived the idea. SG conducted field studies to collect the data. SG analysed

the data. SG and DB wrote the manuscript.

Acknowledgements:

We thank the Maharashtra State Biodiversity Board and the Maharashtra Forest Department

for permits to work on *J. malabaricum*; Kalu Kurade, Ganpat Lokhare and Thakku Kurade

for assistance in the field; Neha Wadmare for help in quantifying floral morphology; Rashmi

Rai and Zakhiya P.C. for help in pollen size quantification; and Yashraj Chavhan, Souparna

Chakrabarty and Nitish for occasional assistance in fieldwork. This research was supported

by intra-mural funding from the Indian Institute of Science Education and Research Pune
(IISER Pune, Maharashtra, India).

References

Armbruster WS, Bolstad GH, Hansen TF, Keller B, Conti E, Pélabon C. **2017**. The measure and mismeasure of reciprocity in heterostylous flowers. *New Phytologist* **215**: 906–917.

Baena-Díaz F, Fornoni J, Sosenski P, *et al.* **2012**. Changes in reciprocal herkogamy during the tristyly-distyly transition in *Oxalis alpina* increase efficiency in pollen transfer: Functional changes in herkogamy. *Journal of Evolutionary Biology* **25**: 574–583.

Baker AM, Thompson JD, Barrett SC. **2000**. Evolution and maintenance of stigma-height dimorphism in Narcissus. II. Fitness comparisons between style morphs. *Heredity* **84**: 514–524.

Barranco D, Arroyo J, Santos-Gally R. **2019**. Avoiding sexual interference: herkogamy and dichogamy in style dimorphic flowers of *Narcissus broussonetii* (Amaryllidaceae). *AoB PLANTS* **11**.

Barrett SCH. **2002**. Sexual interference of the floral kind. *Heredity* **88**: 154–159.

Barrett SCH, Glover DE. **1985**. On the Darwinian Hypothesis of the Adaptive Significance of Tristyly. *Evolution* **39**: 766–774.

Brunet J, Eckert CG. **1998**. Effects of Floral Morphology and Display on Outcrossing in Blue Columbine, *Aquilegia caerulea* (Ranunculaceae). *Functional Ecology* **12**: 596–606.

Brys R, Jacquemyn H. **2014**. Disruption of the distylous syndrome in *Primula veris*. *Annals of Botany* **115**: 27–39.

Brys R, Jacquemyn H. **2020**. The impact of individual inaccuracy of reciprocal herkogamy on legitimate pollen deposition and seed set in a distylous self-incompatible herb. *Journal of Ecology* **108**: 81–93.

Cesaro AC, Barrett SCH, Maurice S, Vaissiere BE, Thompson JD. **2004**. An experimental evaluation of self-interference in *Narcissus assoanus*: functional and evolutionary implications. *Journal of evolutionary biology* **17**: 1367–1376.

Costa J, Castro S, Loureiro J, Barrett SCH. **2017**. Experimental insights on Darwin's cross-promotion hypothesis in tristylous purple loosestrife (*Lythrum salicaria*). *American Journal of Botany* **104**: 616–626.

Eckert CG, Barrett SC. **1994**. Tristyly, self-compatibility and floral variation in *Decodon verticillatus* (Lythraceae). *Biological Journal of the Linnean Society* **53**: 1–30.

Ennos RA. **1981**. Quantitative studies of the mating system in two sympatric species of *Ipomoea* (Convolvulaceae). *Genetica* **57**: 93–98.

Faife-Cabrera M, Ferrero V, Navarro L. **2016**. Relationship between herkogamy, incompatibility and reciprocity with pollen–ovule ratios in *Melochia* (Malvaceae). *Plant Biosystems - An International Journal Dealing with all Aspects of Plant Biology* **152**: 80–89.

Faivre AE, McDade LA. **2001**. Population-level variation in the expression of heterostyly in three species of Rubiaceae: does reciprocal placement of anthers and stigmas characterize heterostyly? *American Journal of Botany* **88**: 841–853.

Ferrero V, Barrett SCH, Rojas D, Arroyo J, Navarro L. **2017**. Associations between sex-

organ deployment and morph bias in related heterostylous taxa with different stylar

polymorphisms. *American Journal of Botany* **104**: 50–61.

Ferrero Victoria, Castro S, Sánchez JM, Navarro L. **2011**. Stigma–anther reciprocity,

pollinators, and pollen transfer efficiency in populations of heterostylous species of *Lithodora*

and *Glandora* (Boraginaceae). *Plant Systematics and Evolution* **291**: 267–276.

Ferrero V., Chapela I, Arroyo J, Navarro L. **2011**. Reciprocal style polymorphisms are not

easily categorised: the case of heterostyly in *Lithodora* and *Glandora* (Boraginaceae). *Plant*

Biology **13**: 7–18.

Ganguly S. **2019**. Variation in sex organ dimensions across individuals, morphs and species:

consequences for reciprocity, herkogamy, and reproductive success in species with style

length polymorphism. PhD. Thesis, Indian Institute of Science Education and Research Pune,

India.

Ganguly S, Barua D. **2020**. High herkogamy but low reciprocity characterises isoplethic

populations of *Jasminum malabaricum*, a species with stigma-height dimorphism. *Plant*

Biology. doi: 10.1111/plb.13127. I*n press*.

Haddadchi A. **2013**. Stylar polymorphism, reciprocity and incompatibility systems in

Nymphoides montana (Menyanthaceae) endemic to southeastern Australia. *Plant systematics*

and evolution **299**: 389–401.

Hernandez A, Ornelas JF. **2007**. Disassortative pollen transfer in distylous *Palicourea*

padifolia (Rubiaceae), a hummingbird-pollinated shrub. *Ecoscience* **14**: 8–16.

Jacquemyn H, Gielen M, Brys R. **2018**. Is sexual organ reciprocity related to legitimate pollen deposition in distylous *Pulmonaria* (Boraginaceae)? *Oikos* **127**: 1216–1224.

Keller B, Thomson JD, Conti E. **2014**. Heterostyly promotes disassortative pollination and reduces sexual interference in Darwin's primroses: evidence from experimental studies. *Functional Ecology* **28**: 1413–1425.

Keller B, de Vos JM, Conti E. **2012**. Decrease of sexual organ reciprocity between heterostylous primrose species, with possible functional and evolutionary implications. *Annals of botany* **110**: 1233–1244.

Larrinaga AR, Guitián P, Garrido JL, Guitián J. **2009**. Floral morphology and reproductive success in herkogamous *Narcissus cyclamineus* (Amaryllidaceae). *Plant Systematics and Evolution* **278**: 149–157.

Lau P, Bosque C. **2003**. Pollen flow in the distylous *Palicourea fendleri* (Rubiaceae): an experimental test of the disassortative pollen flow hypothesis. *Oecologia* **135**: 593–600.

Li X-X, Zou Y, Xiao C-L, Gituru RW, Guo Y-H, Yang C-F. **2013**. The differential contributions of herkogamy and dichogamy as mechanisms of avoiding self-interference in four self-incompatible *Epimedium* species. *Journal of evolutionary biology* **26**: 1949–1958.

Liu S-J, Wu L-Y, Huang S-Q. **2016**. Shortened anther–stigma distance reduces compatible pollination in two distylous *Primula* species. *Journal of Plant Ecology* **9**: 224–232.

Lloyd DG, Webb CJ. **1992**. The evolution of heterostyly In: Barrett SCH, ed. *Evolution and function of heterostyly*. Springer, 151–178.

Martén-Rodríguez S, Muñoz-Gamboa P, Delgado-Dávila R, Quesada M. **2013**. Asymmetric pollen transfer and reproductive success of the hawkmoth-pollinated distylous

tree *Palicourea tetragona* (Rubiaceae) at La Selva, Costa Rica. *Journal of Tropical Ecology* **29**: 501–510.

Nishihiro J, Washitani I. **1998**. Patterns and consequences of self-pollen deposition on stigmas in heterostylous *Persicaria japonica* (Polygonaceae). *American Journal of Botany* **85**: 352–359.

Nishihiro J, Washitani I, Thomson JD, Thomson BA. **2000**. Patterns and consequences of stigma height variation in a natural population of a distylous plant, *Primula sieboldii*. *Functional Ecology* **14**: 502–512.

Opedal ØH. **2018**. Herkogamy, a Principal Functional Trait of Plant Reproductive Biology. *International Journal of Plant Sciences*.

Sampson DA, Krebs RA. **2012**. Quantitative evaluation of reciprocal herkogamy in the distylous species, *Hedyotis caerulea* (Rubiaceae). *Plant systematics and evolution* **298**: 1361–1370.

Sanchez JM, Ferrero V, Navarro L. **2008**. A new approach to the quantification of degree of reciprocity in distylous (sensu lato) plant populations. *Annals of Botany* **102**: 463–472.

Santos-Gally R, Perez-Barrales R, Simon VI, Arroyo J. **2013**. The role of short-tongued insects in floral variation across the range of a style-dimorphic plant. *Annals of Botany* **111**: 317–328.

Shore JS, Barrett SCH. **1984**. The effect of pollination intensity and incompatible pollen on seed set in *Turnera ulmifolia* (Turneraceae). *Canadian Journal of Botany* **62**: 1298–1303.

Simón-Porcar VI, Santos-Gally R, Arroyo J. **2014**. Long-tongued insects promote

disassortative pollen transfer in style-dimorphic *Narcissus papyraceus* (Amaryllidaceae).

Journal of Ecology **102**: 116–125.

Sosenski P, Fornoni J, Molina-Freaner FE, Weller SG, Domínguez CA. **2010**. Changes in

sexual organ reciprocity and phenotypic floral integration during the tristyly–distyly

transition in *Oxalis alpina*. *New Phytologist* **185**: 829–840.

Stehlik I, Caspersen JP, Barrett SCH. **2006**. Spatial ecology of mating success in a

sexually polymorphic plant. *Proceedings of the Royal Society of London B: Biological*

Sciences **273**: 387–394.

Takebayashi N, Wolf DE, Delph LF. **2006**. Effect of variation in herkogamy on outcrossing

within a population of *Gilia achilleifolia*. *Heredity* **96**: 159–165.

Thompson JD, Barrett SC, Baker AM. **2003**. Frequency–dependent variation in

reproductive success in *Narcissus: implications* for the maintenance of stigma–height

dimorphism. *Proceedings of the Royal Society of London B: Biological Sciences* **270**: 949–

953.

Thompson JD, Cesaro AC, Arroyo J. **2012**. Morph Ratio Variation and Sex Organ

Reciprocity in Style-Dimorphic *Narcissus assoanus*. *International Journal of Plant Sciences*

173: 885–893.

Valois-Cuesta H, Soriano PJ, Ornelas JF. **2011**. Dimorphisms and self-incompatibility in

the distylous species *Palicourea demissa* (Rubiaceae): possible implications for its

reproductive output. *Journal of plant research* **124**: 137–146.

de Vos JM, Keller B, Isham ST, Kelso S, Conti E. **2012**. Reproductive implications of

herkogamy in homostylous primroses: variation during anthesis and reproductive assurance

in alpine environments. *Functional Ecology* **26**: 854–865.

de Vos JM, Keller B, Zhang L-R, Nowak MD, Conti E. **2018**. Mixed Mating in

Homostylous Species: Genetic and Experimental Evidence from an Alpine Plant with

Variable Herkogamy, Primula halleri. *International Journal of Plant Sciences* **179**: 87–99.

Webb CJ, Lloyd DG. **1986**. The avoidance of interference between the presentation of

pollen and stigmas in angiosperms II. Herkogamy. *New Zealand Journal of Botany* **24**: 163–

178.

Yuan S, Chen S, Deng X, Duan T, Luo Z, Zhang D. **2017**. Pollen–ovule ratios are strongly

correlated with floral reciprocity, in addition to sexual system, in *Mussaenda* (Rubiaceae).

Nordic Journal of Botany **35**: 395–403.

Zhou W, Barrett SC, Wang H, Li D-Z. **2015**. Reciprocal herkogamy promotes

disassortative mating in a distylous species with intramorph compatibility. *New Phytologist*

206: 1503–1512.

Tables

Table 1: Range (Min-Max) and coefficient of variation (CV) of individual mismatch and

herkogamy observed in the four study populations a) Trimbak, b) Bhimashankar, c) Mulshi

and d) Kaas. L and S stand for long- and short-styled morph respectively. The number of

individuals sampled for the long- and the short-styled morph for the populations are 20 and

19 for Trimbak; 30 and 30 for Bhimashankar, Mulshi, and Kaas.

		Mismatch		**Herkogamy**	
Population	**Morph**	**Min-Max (mm)**	**CV %**	**Min-Max (mm)**	**CV %**
a) Trimbak	L	1.25, 9.27	38.70	2.47, 9.49	23.90
	S	4.39, 6.47	10.80	5.91, 11.30	17.70
b) Bhimashankar	L	1.74, 7.80	36.70	2.70, 11.40	25.20
	S	4.22, 7.02	12.30	2.61, 12.50	22.90
c) Mulshi	L	1.27, 8.80	41.10	3.53, 11.40	22.20
	S	4.24, 6.92	12.00	5.56, 11.50	16.80
d) Kaas	L	2.10, 9.34	38.50	2.00, 11.30	29.60
	S	4.93, 7.42	10.50	6.51, 13.10	15.80

Table 2: Relationship of herkogamy and reciprocity to total stigma pollen load, legitimate

pollen fraction and fruit set in the Long-styled (L) and short-styled (S) morph in a)

Bhimashankar, and b) Kaas. Fruit set was estimated only for the Bhimashankar population. *P*

< 0.05 and *P* < 0.1 are denoted by * and Ψ, respectively, next to the Spearman's rank-order

correlation coefficients presented. For the Bhimashankar population *n* = 30 long-styled and

29 short-styled individuals for total pollen load and legitimate pollen fraction, and *n* = 29

individuals for each morph for fruit set. For the Kaas population *n* = 32 long-styled and 30

short-styled individuals.

Population	Trait	Morph	Total pollen load	Legitimate fraction	Fruit set
a) Bhimashankar	Mismatch	L	0.14	0.28	0.18
		S	0.03	-0.23	0.33Ψ
	Herkogamy	L	0.06	0.45*	0.39*
		S	-0.05	-0.20	0.06
b) Kaas	Mismatch	L	-0.19	0.04	
		S	0.19	-0.18	
	Herkogamy	L	-0.33Ψ	0.13	
		S	-0.27	-0.06	

Figures

Figure 1: (a) Mismatch and (b) herkogamy in the long-styled (dark grey boxes) and short-

styled (open boxes) morphs of the four study populations. The number of individuals sampled

for the long- and the short-styled morph for the populations are 20 and 19 for Trimbak; 30

and 30 for Bhimashankar, Mulshi, and Kaas. Outliers are denoted by +. The bold line in the

boxplots represents the median, the ends of the box represent the first and the third quartiles

and the whiskers represent 1.5 times the interquartile range.

Figure 2: The total stigma pollen load (a, c) and the legitimate pollen fraction (b, d) in the Bhimashankar and Kaas populations. The number of individuals sampled for the long- and short-styled morph is 30 and 29 for Bhimashankar, and 32 and 30 for Kaas. One extreme outlier value of greater than 5000 for total pollen load in a long-styled individual from the Bhimashankar population has been removed from the plot for better visualization of the differences between the morphs. The bold line in the boxplots represents the median, the ends of the box represent the first and the third quartiles and the whiskers represent 1.5 times the interquartile range.

Figure 3: (a) The total pollen load, and (b) the legitimate pollen fraction in control (non-

emasculated, grey boxes) and emasculated flowers (open boxes) to determine autogamous

pollen deposition. One point from the legitimate pollen fraction for the emasculated group of

long-styled morph (80%) has been removed from this plot for better visualization of the

differences between the control and emasculation treatments. Data are from three flowers

from five individuals (represented by the different symbols). The bold line in the boxplots

represents the median, the ends of the box represent the first and the third quartiles and the

whiskers represent 1.5 times the interquartile range.

Figure 4: Natural fruit set in long- (grey boxes) and short-styled (open boxes) individuals

from the Bhimashankar population measured as the percentage fruit formed per flower in

2015 and 2016. The number of individuals sampled for the long- and the short-styled morphs

are 30 and 29 respectively in 2015, and 29 individuals of each morph in 2016. The plus sign

represents outliers. The bold line in the boxplots represents the median, the ends of the box

represent the first and the third quartiles and the whiskers represent 1.5 times the interquartile

range.

Figure 5: Relationship between herkogamy and mismatch in the two morphs (long-styled

morph: a,b,c,d; short-styled morph: e,f,g,h) in the four study populations. Pearson's

correlation coefficient (*r*), and the corresponding *P*-value are shown where $P < 0.05$, and 'ns'

denotes not significant. The number of individuals sampled for the long- and the short-styled

morph for the populations are 20 and 19 for Trimbak; 30 and 30 for Bhimashankar, Mulshi,

and Kaas.