Prediction of complex phenotypes using the *Drosophila* metabolome

Palle Duun Rohde[1,*], Torsten Nygaard Kristensen[1], Pernille Sarup[2,3],

Joaquin Muñoz[1] & Anders Malmendal[4,*]

[1] *Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark.*

[2] *Department of Molecular Biology and Genetics, Aarhus University, Tjele, Denmark*

[3] *Nordic Seed A/S, 8300 Odder, Denmark*

[4] *Department of Science and Environment, Roskilde University, Roskilde, Denmark*

[*]*Corresponding authors: palledr@bio.aau.dk and amalm@ruc.dk*

Abstract

Understanding the genotype – phenotype map and how variation at different levels of biological organization are associated are central topics in modern biology. Fast developments in sequencing technologies and other molecular omic tools enable researchers to obtain detailed information on variation at DNA level and on intermediate endophenotypes, such as RNA, proteins, metabolites. This facilitates our understanding of the link between genotypes and molecular and functional organismal phenotypes. Here, we use the *Drosophila* Genetic Reference Panel and nuclear magnetic resonance (NMR) metabolomics to investigate the ability of the metabolome to predict organismal phenotypes. We do this by performing NMR metabolomics on four replicate pools of male flies from each of 170 isogenic lines. We show first that metabolite profiles are variable among the investigated lines and that this variation is highly heritable. Second, we identify genes associated with metabolome variation. Third, we show that for four of five traits related to behaviour and stress resistance the metabolome predicts phenotypes more accurately than genomic data, and that the metabolites driving the increased accuracy was trait specific. Our comprehensive characterization of population-scale diversity of metabolomes and its genetic basis illustrates that metabolites have large potential as predictors of organismal phenotypes. This finding has strong applied importance e.g. in human medicine and animal and plant breeding.

Keywords

Drosophila melanogaster; locomotor activity; stress resistance traits; NMR metabolomics; *Drosophila* Genetic Reference Panel

1. Introduction

Understanding how information encoded in the DNA is transcribed to RNA, translated to proteins and other downstream endophenotypes such as metabolites, and how this information dictates the organismal functional phenotype is in the core of several biological disciplines. While the revolutionary work that led to the discovery of this central flow of genetic information within biological systems was published more than 70 years ago[1] we are still making progress in understanding the genotype – phenotype map. This is aided by new technologies within molecular and systems biology, allowing researchers to obtain full genome sequences from individuals of any species, detailed information about expression levels of all genes, abundancies of proteins and metabolites etc. These omics tools have provided unforeseen knowledge about the genetic and environmental background of complex traits, which has revolutionized the field of genetics with strong impacts on multiple research disciplines including medicine, animal and plant breeding and evolutionary biology.[2–5]

One common goal of studies on genotype – phenotype associations is to understand to what extent a complex organismal phenotype such as behavioural traits, traits linked to reproduction, diseases or the ability to cope with stressful environmental conditions, can be predicted from DNA sequence information or from endophenotypes. If information obtained from levels upstream the functional organismal phenotype can accurately predict the phenotype this has wide ranging applications across life sciences, and this has proven useful in several cases.[6–10] However, studies have also demonstrated that predicting complex phenotypes based on genetic information can often be difficult and the predictive power in such studies is typically low.[11–13] Reasons for the difficulties to accurately predict the phenotype are many and include (i) that quantitative traits result from a complex interplay between a large number of genes, each with a small contribution to the phenotype, combined with environmental factors, (ii) despite progress the underlying genetic architectures for most traits of medical interest or for traits of relevance in agriculture or evolutionary biology are still not well-understood, (iii) the effect of so called candidate genes is often depending on the genetic background (epistasis) and they explain only a small proportion of the heritability, and (iv) genes and environments interacts in their effect on the phenotype. These factors give rise to substantial challenges in constructing and implementing genetic risk prediction models across biological disciplines.

Challenges with using DNA sequence variation to predict variation at the organismal phenotypic level have sparked an interest in using endophenotypes, such as transcript, protein or metabolite levels as predictors of complex functional phenotypes.[14–17] Endophenotypes influence and regulate the functional phenotype and in contrast to the genotype, which is fixed in an individual's lifetime, they are governed by interactions between the genome of an individual, and internal and external influences that range from the cellular level and to the wealth of external biotic and abiotic factors an individual is exposed to in its environment. Thus, endophenotypes have been proposed to constitute proximal links between variation at the genome level and the organismal phenotype and they may provide more accurate predictors of the functional phenotype compared to the genotype.[15,18]

We hypothesize, that the predictive value of endophenotypes for organismal functional phenotypes is linked to the proximity of the endophenotype to the organismal phenotype. Accepting this premise information on the

metabolome level should provide more accurate prediction than e.g. information at the gene-expression or protein levels. This is an emerging research field and we do not have many results on this yet. However, there are studies that support the hypothesis that data from gene expression or proteomics studies combined with genotype information improves prediction of several traits in e.g. *Drosophila melanogaster* and maize.[19–22] The metabolome is closer to the functional phenotype than transcriptomics and proteomics and recent study based on investigating 453 metabolites in 40 DGRP lines suggest that the metabolome might constitute a reliable predictor of organismal phenotypes and that the metabolome provide novel insights into the underpinnings of complex traits and its genetic basis.[18]

Here we elaborate on the findings by Zhou et al. (2019)[18] by using nuclear magnetic resonance (NMR) metabolomics to obtain information about fly metabolomes in pooled samples of whole male flies from 170 lines from the *Drosophila* Genetic Reference Panel (DGRP); a system of fully inbred sequenced lines of *D. melanogaster*.[23,24] NMR metabolomics constitute a highly reproducible technique that in contrast to mass spectrometry allows for metabolic profiling of the total complement of metabolites in a sample.[25] With this set-up, we test 1) to what extent the metabolome varies across investigated lines and whether this variation is heritable; 2) can we detect DNA sequence variants that are associated with variation in NMR feature intensity (mQTLs)?; and 3) can we use metabolomic data to increase prediction accuracy of five complex behavioural and stress tolerance phenotypes compared to when prediction is based on DNA sequence variation?

2. Materials and Methods

2.1 *Drosophila* lines, husbandry and collection

We used 170 inbred lines of the DGRP.[23,24] The DGRP lines were established by 20 consecutive generations of full sibling inbreeding from isofemales collected at the farmer's market in Raleigh, NC. Complete genome sequence of the DGRP lines have been obtained using Illumina platform, and is publicly available.[23,24]

The DGRP lines were maintained on standard *Drosophila* medium (see Kristensen *et al.* (2016)[26] for recipe details) in a 23 °C climate chamber at 50% relative humidity with a 12h:12-h light:dark cycle. For this study only male flies with an average age of 24 hours was used. Flies were collected from 170 DGRP lines in four replicates containing 20 flies per replicate.

2.2 *Drosophila* sample preparation

We prepared four replicates of 20 pooled male flies for NMR spectroscopy. Flies were snap frozen at three days of age and kept at –80°C. Samples were mechanically homogenized with a Kinematica, Pt 1200 (Buch & Holm A/S, Herlev, Denmark) in 1 mL of ice-cold acetonitrile (50%) for 45 s. Hereafter samples were centrifuged (10,000 g) for 10 min at 4°C and the supernatant (900 μL) was transferred to new tubes, snap frozen and stored at –80°C. The supernatant was lyophilized and stored at -80°C. Immediately before NMR measurements, samples were rehydrated in 200 mL of 50 mM phosphate buffer (pH 7.4) in D_2O, and 180 μL was transferred to a 3 mm NMR tube. The buffer contained 50 mg/L of the chemical shift reference 2,2-Dimethyl-2-silapentane-5-sulfonate-d_6, sodium salt (DSS), and 50 mg/L of sodium azide to prevent bacterial growth.

2.3 NMR experiments and spectral processing

NMR measurements were performed at 25°C on a Bruker Avance III HD 800 spectrometer (Bruker Biospin, Rheinstetten, Germany), operating at a [1]H frequency of 799.87 MHz, and equipped with a 3 mm TCI cold probe. [1]H NMR spectra were acquired using a standard NOESYPR1D experiment with a 100 ms delay. A total of 128 transients of 32 K data points spanning a spectral width of 20 ppm were collected.

The spectra were processed using Topspin (Bruker Biospin, Rheinstetten, Germany). An exponential line-broadening of 0.3 Hz was applied to the free-induction decay prior to Fourier transformation. All spectra were referenced to the DSS signal at 0 ppm, manually phased and baseline corrected. The spectra were aligned using *i*coshift.[27] The region around the residual water signal (4.87-4.70 ppm) was removed in order for the water signal not to interfere with the analysis. The high- and low-field ends of the spectrum, where no signals except the reference signal from DSS appear, were also removed (i.e., leaving data between 9.7 and 0.7 ppm). The spectra were normalized by probabilistic quotient area normalization.[28]

2.4 Quantitative genetics of NMR intensies

Each NMR feature (14,440 in total) was treated as a quantitative trait. For each NMR feature, we fitted a linear mixed model to partition the total phenotypic variation (i.e. one NMR feature) into genetic and environmental variation. Using the R package qgg[29] we fitted the model,

$$y = Xb + Zg + e \qquad (1)$$

where y was a vector of NMR intensities for a particular NMR feature, X and Z are design matrices linking fixed and random effects to the phenotype, b is a vector of the fixed effects (*Wolbachia* infection status, and major polymorphic inversions; *In2Lt, In2RNS, In2RY1, In2RY2, In3LP, In3LM, In3LY, In3RP, In3RK, In3RMo, In3RC*. Information available at http://dgrp2.gnets.ncsu.edu), g is a vector of the random genetic effects defined as $g \sim N(\mathbf{0}, G\sigma_g^2)$, and e is a vector of residual effects defined as $e \sim N(\mathbf{0}, I\sigma_e^2)$. The variance structure among the random effects are modelled as independent for the residual effects (by the identity matrix I), and for the genetic effects by the additive genomic relationship matrix G, which was constructed using all SNPs (minor allele frequency ≥ 0.05) as $G = WW'/m$, where m is the number of SNPs (i.e. 1,725,755), and W is a centered and scaled genotype matrix, where each column vector is $w_i = \frac{a_i - 2p_i}{\sqrt{2p_i(1-2p_i)}}$, p_i is the allele frequency of the ith SNP, and a_i is the ith column vector of the allele count matrix, A, which contains the genotypes coded as 0 or 2 counting the number of the minor allele (genotypes are available at http://dgrp2.gnets.ncsu.edu).

For each NMR feature we estimated the proportion of phenotypic variation explained by SNP variation as $\hat{h}_{\text{SNP}}^2 = \frac{\hat{\sigma}_g^2}{\hat{\sigma}_g^2 + \hat{\sigma}_e^2}$, where $\hat{\sigma}_g^2$ and $\hat{\sigma}_e^2$ are the estimated variance components from equation 1. The significance of each $\hat{h}_{\text{SNP}}^2$ was determined as $\hat{h}_{\text{SNP}}^2 - SE(\hat{h}_{\text{SNP}}^2) \times Z > 0$, where Z is the quantile function of the normal distribution at probability $P = 0.05/14,440$. Thus, the resulting set of heritability estimates are those estimates that differ significantly from zero when accounting for a total 14,440 statistical tests.

2.5 Mapping of metabolome QTL

mQTLs for mean NMR intensity was identified using linear regression,[30] using the function for single marker association analysis implemented in the qgg package.[29] The estimated genetic effects ($\hat{g}$, from equation 1) was used as line means, since these values represents the within DGRP line mean intensity of a single NMR feature adjusted for *Wolbachia*, chromosomal inversions, and polygenicity, which then was regressed on marker genotypes.

Significant mQTLs were identified as those SNP-NMR associations with a Bonferroni adjusted $P < 0.05/1{,}725{,}755 = 2.9 \times 10^{-8}$; i.e. genome-wide significant. Furthermore, we restricted the search for mQTLs to the set of NMR features with a significant heritability estimate.

2.6 Phenotypic predictions

Using the linear mixed model (BLUP; best linear unbiased prediction) framework we performed several phenotypic prediction models using either genomic (GBLUP) or metabolomic (MBLUP) information to investigate if the metabolome provides additional information that will increase the accuracy of prediction compared to genomic prediction. The DGRP has been characterised for a wide range of molecular, environmental stress resistance, morphological and behavioural phenotypes.[31,32] When the number of individual genotypes is low, *i.e.*, the number of DGRP lines, a large number of within-line replicates are required for accurate predictions.[33] Therefore, we restricted the set of quantitative traits to those where we had access to all individual observations, and where the average number of observations within-line >25 (Table 1).

The five test traits were initially adjusted for experimental factors (see references in Table 1), *Wolbachia* infection status, and major polymorphic inversions (*In2Lt*, *In2RNS*, *In3RP*, *In3RK*, *In3RMo*). The adjusted phenotypic values were obtained as: $\widetilde{y}_l = \hat{L}_l + \hat{e}_l$, where $\hat{L}_l$ is the estimated line effect for DGRP line l (*i.e.*, the BLUP value), and $\hat{e}_l$ is a vector containing the residuals for line l. Thus, y_l and $\widetilde{y}_l$ has the same dimension. The estimated line effects ($\hat{L}$) were assumed $L \sim N(0, I\sigma_L^2)$. The metabolome contains aggregated information both on the individual genotypes and environmental exposures. Thus, to avoid double counting the genomic variation as represented by SNP information by adding genomic information in both the two steps in the GBLUP and MBLUP analysis, we assumed the DGRP lines to be independent, modelled by the identity matrix I, instead of G in the first step.

For each quantitative trait (Table 1) we fitted two models, each for 100 randomly selected training sets (t, the training sets were the same for all prediction models) containing 90% of the DGRP lines:

$$\widetilde{y}_t = Z_t g_t + e_t, \tag{1}$$

$$\widetilde{y}_t = Z_t m_t + e_t, \tag{2}$$

where $\widetilde{y}_t$ is the adjusted phenotypic values for the DGRP lines in training set t, e_t is a vector of random residuals, Z_t is a design matrix linking the genomic (g_t) and metabolomic (m_t) effects to the phenotypes. The random genomic effects are $g_t \sim N(0, G_{[t,t]}\sigma_g^2)$, and the metabolomic effects are $m_t \sim N(0, M_{[t,t]}\sigma_m^2)$, where G is the additive genomic relationship matrix as specified previously, and M is the metabolomic relationship matrix. The metabolomic relationship matrix was computed as $M = QQ'/m_{\text{NMR}}$, where Q is a $n \times m_{\text{NMR}}$ matrix of

adjusted, centred and scaled NMR intensities ($m_{\text{NMR}} = 14{,}440$). Each column vector of $\boldsymbol{Q}$ contains the BLUP values from a mixed model where the phenotype was the corresponding NMR intensity, which was adjusted for *Wolbachia* infection status and major polymorphic inversions (*In2Lt*, *In2RNS*, *In3RP*, *In3RK*, *In3RMo*), using the identity matrix as covariance structure. This was done to obtain a data structure similar to the genomic data; namely one value of each DGRP line/NMR intensity.

The predicted phenotypes in validation set v ($\widehat{\boldsymbol{y}}_v$) was obtained using Equation 2 and 3 for GBLUP and MBLUP, respectively.

$$\widehat{\boldsymbol{y}}_v = (\boldsymbol{G}_{[v,t]}\hat{\sigma}_g^2)\big[\boldsymbol{G}_{[t,t]}\hat{\sigma}_g^2 + \boldsymbol{I}_{[t,t]}\hat{\sigma}_e^2\big]^{-1}(\widetilde{\boldsymbol{y}}_t - \mu_t), \qquad (2)$$

$$\widehat{\boldsymbol{y}}_v = (\boldsymbol{M}_{[v,t]}\hat{\sigma}_m^2)\big[\boldsymbol{M}_{[t,t]}\hat{\sigma}_m^2 + \boldsymbol{I}_{[t,t]}\hat{\sigma}_e^2\big]^{-1}(\widetilde{\boldsymbol{y}}_t - \mu_t). \qquad (3)$$

The prediction accuracy (PA) was quantified as the mean Pearson's correlation between observed and predicted phenotypes across training sets, $PA = \frac{1}{100}\sum_{t=1}^{100} Cor(\widetilde{\boldsymbol{y}}_v, \widehat{\boldsymbol{y}}_v)$. The accuracies of GBLUP and MBLUP was compared using a paired t-test.

2.7 NMR cluster-guided phenotypic predictions

From genomic prediction models we know that allowing marker effects to come from different distributions, e.g. grouping genetic variants into functional pathways, can increase the prediction accuracy markedly.[33–38] Therefore, we investigated if similar benefits could be achieved by partitioning the metabolome.

Using the $\boldsymbol{Q}$-matrix (i.e. the $n \times m_{\text{NMR}}$ matrix of adjusted, centred and scaled NMR intensities) we computed all pairwise Pearson's correlation coefficients and performed hierarchical clustering on the dissimilarity on the correlation coefficients using an unweighted pair group method with arithmetic mean agglomeration (Fig. 1). Using a range of total number of clusters $K_{cl} = \{25, 50, 75, 100, 125, 200\}$ we performed metabolomic feature best linear unbiased prediction (MFBLUP) which is an extension to the MBLUP model (Eq. 2) containing an additional metabolomic effect (Eq. 4, Fig. 1). For each K_{cl} level we estimated model parameters for each cluster (for $K_{cl} = 25$ we ran 25 models, and for $K_{cl} = 50$ we ran 50 models etc.) as follows:

$$\widetilde{\boldsymbol{y}}_t = \boldsymbol{Z}_t \boldsymbol{m}_t^{K_{cl\_k}} + \boldsymbol{Z}_t \boldsymbol{m}_t^r + \boldsymbol{e}_t, \qquad (4)$$

where the superscript K_{cl_k} indicate the total number of clusters (K_{cl}) and the cluster number (cl). The first metabolomic effect was defined as $\boldsymbol{m}_t^{K_{cl\_k}} \sim N(0, \boldsymbol{M}_{[t,t]}^{K_{cl\_k}} \sigma_{m^{K_{cl\_k}}}^2)$, where $\boldsymbol{M}_{[t,t]}^{K_{cl\_k}}$ corresponds to the metabolomic relationship of the DGRP lines within the training set (t) for the NMR features within cluster number cl among the K_{cl} clusters. The second metabolomic effect ($\boldsymbol{m}_t^r \sim N(\boldsymbol{0}, \boldsymbol{M}_{[t,t]}^r \sigma_{m^r}^2)$) is the random effects using all NMR features except those within the K_{cl_k} cluster.

Similar to the MBLUP model, the predicted phenotypes in the validation set v ($\widehat{\boldsymbol{y}}_v$) was obtained as

$$\widehat{\boldsymbol{y}}_v = (\boldsymbol{M}_{[v,t]}^{K_{cl\_k}}\hat{\sigma}_{m^{K_{cl_k}}}^2 + \boldsymbol{M}_{[v,t]}^r \hat{\sigma}_{m^r}^2)\big[\boldsymbol{M}_{[t,t]}^{K_{cl\_k}}\hat{\sigma}_{m^{K_{cl_k}}}^2 + \boldsymbol{M}_{[t,t]}^r\hat{\sigma}_{m^r}^2 + \boldsymbol{I}_{[t,t]}\hat{\sigma}_e^2\big]^{-1}(\widetilde{\boldsymbol{y}}_t - \mu_t). \qquad (5)$$

The prediction accuracy for each K_{cl_k} cluster was obtained as $PA = \frac{1}{100}\sum_{t=1}^{100} Cor(\widetilde{\boldsymbol{y}}_v, \widehat{\boldsymbol{y}}_v)$, and was compared (using paired t-test corrected for multiple testing by a false discovery rate (FDR) of <0.05) within and across K_{cl} clusters to identify the NMR features resulting in the largest prediction accuracy. We only considered

clusters to be signigicant if the FDR was below 0.05 , and if the proportion of variance captured by the cluster was larger than 1%, which is computed as $\left(\hat{\sigma}^2_{m^{K_{cl_k}}}/(\hat{\sigma}^2_{m^{K_{cl_k}}} + \hat{\sigma}^2_{m^r})\right) > 1\%$.

Finaly, to investigate if we could increase the predive performance further, we took all the clusters that increased the trait-specific predictive performance (including clusters where the variance captured was below 1%), ranked them by their predictive performance, and ran a new series of prediction models, adding the NMR features sequential to the model based on the cluster's predictive performance (high to low).

3 Results

3.1 The metabolome of *D. melanogaster*

Using [1]H NMR we quantified the metabolome of males from 170 DGRP lines in four biological replicates (Supplementary Data file S1). For each of the 14,440 NMR features we estimated the proportion of variation in NMR intensity explained by common genetic variants (h^2, Fig. 2A). In total, 39% of all NMR features had a significant heritability estimate (Fig. 2B), of which the average heritability was 0.26 (0.13 across all features).

For those NMR features where we observed a significant heritability estimate, we next sought to identify genetic variants associated with NMR feature intensity, namely mQTLs. We found a total of 5,782 genome-wide significant mQTLs (Supplementary Table 1) covering 1249 NMR features and 1468 unique genetic variants. The significant mQTLs were found within 963 genes, with the gene *Coronin* having 488 genome-wide associations (Supplementary Table S1), which is extreme compared to the average of 6.0 significant associations per gene.

3.2 Phenotypic predictions

To test the predictive performance of the metabolome we obtained data from five previously published complex phenotypes (Table 1); two behavioural traits and three stress resistance traits. We constructed relationship matrices based on genomic and on metabolomic information and performed genomic/metabolomic best linear unbiased prediction (GBLUP/MBLUP). For each trait we used 90% of the data to estimate the parameters using either the genomic or metabolomic relationship matrices and used the estimated parameters to predict the remaining 10% of the data. This was repeated on 100 random data divisions.

The mean prediction accuracy for the two behavioural traits, locomotor activity without and with treatment of Ritalin, was below 0.1 when based on genomic information (Fig. 3A and Fig. 3C). Using metabolomic information the predictive performance was increased to above 0.4 (Fig. 3A, Fig. 3C, and Suplementary Table S2). By using the *Drosophila* metabolome, we could also increase the predictive accuracy of the two environmental stress resistance traits, chill come recovery and starvation resistance (Fig. 3E and Fig. 3I, Supplementary Table S2). However, for the trait, startle response (a behavioural response to a physical disturbance), prediction using genomic information was superior over the metabolome (Fig. 3G).

We computed Pearson's correlation coefficient among all NMR features and performed hierachical clusting (Supplementary Fig. S1). We then ran the two component NMR cluster-guided prediction model (MFBLUP) where the first component was based on NMR features within one cluster, and the second component was based on the remaining NMR features (Fig. 1). We tested all clusters from the hierachical clustering using different

number of total clusters; $K_{cl} = \{25, 50, 75, 100, 125, 200\}$ (Fig. 1). By the extension of the MBLUP model we could increase the predictive performance of all five quantitative traits by 17%-185% (Fig. 3, Supplementary Table S2). Interestingly, the largest improvement in prediction accuracy was obtained at different cluster levels for the five traits (Fig. 3 and Supplementary Figure S2). For locomotor activity the largest improvement in prediction accuracy was obtained at cluster level $K_{cl} = 100$ (Fig. 3B), with cluster 1 as the only cluster that had significantly increased predictive accuracy (Supplementary Table S3 and Supplementary Fig. 3). For locomotor activity (Ritalin treatment), startle response and starvation resistance, cluster level $K_{cl} = 200$ contained the clusters that gave the highest predictive performance (Fig. 3 and Supplementary Fig. S2). For activity (with Ritalin) cluster 121 and 74 (Supplementary Table S3) increased the predictive performance. Combining the two clusters increased the prediction accuracy insignificantly by 1% (Fig. 3C, Supplementary Figure S5). Eight clusters increased the predictive performance (that also captured >1% feature variance) for startle response (1, 2, 5, 14, 99, 112, 145 and 187, Supplementary Table S3 and Fig. S3), and by combining cluster 1, 2, 5, 112 and 145 further increased the predition accuracy (Fig 3G, Supplementary Table S2, Supplementary Figure S7). Startle response was the only trait where metabolomic prediction performed worse than genomic prediction (Figure 3G); however, by combining the five clusters with the highest prediction accuracy the metabolomic prediction performed better than genomic prediction (Supplementary Table S2). Five clusters (21, 65, 83, 95 and 140) increased the accuracy of prediction (and captured >1% feature variance) for starvation resistance (Supplementary Fig. 3), and the joint effect of cluster 21, 65, 83 and 95 further insignificantly increased the accuracy (Fig. 3I, Supplementary Table S3, Supplementary Figure S8). Finally, for chill coma recovery the maximum prediction accuracy was obtained at cluster level $K_{cl} = 50$ (Fig. 3F), where cluster 5, 27 and 35 was significant explaining >1% of the NMR feature variance (Supplementary Table S3 and Supplementary Figure S3). Combining the two clusters with the highest accuracy led to insignificant increased in accuracy (Fig. 3E, Supplementary Table S2 and Fig. S8).

The contributions to the predictive clusters from the metabolite NMR-spectra were mapped on to the NMR spectrum (Fig. 4). Interestingly, none of the clusters contained signals from the highest concentration metabolites. Rather they contain signals from metabolites at lower abundance or very broad signals, suggesting contributions from small metabolites bound to larger molecules or larger molecules themselves. Out of the 18 (Table 2) clusters, 7 (1, 2, 5, 21, 32, 45, 65; Figure 4A) contained signals in a region where aromatic compunds with quartenary nitrogens, such as NAD and nicotinamide ribotide appears. These clusters showed high prediction accuracy for locomotor activity , chill coma recovery and startle response. Four clusters (32, 74, 83, 95; Figure 4A) contained signals in a region where other heterocycles, such as adenosine, appears. Furthermore, 6 clusters (112, 121, 129, 139, 145, 154; Figure 4B) contained signals in a region where aromatic groups from aminoacids like histidine and tyrosine appear. Most of these are important for locomotor activity with Ritalin and starvation resistance. There is also one cluster (171 ; Figure 4C) that contained signals in a region where signals from sugars appear, and another (187 ; Figure 4D) that contained signals in a region where amino acid CH_2 groups appear. Out of these clusters 1 and 187 clearly contained signals that would usually be identified as baseline, while clusters 2, 5, 32, 112, 129, 139, 145 contained mostly well resolved, though often low intensity, signals. The remaining signals are somewhere in between. Only cluster 32 could be matched to a known

metabolite and its signals are assigned as coming from the nicotinamide group of NADP. The other metabolites could not be found in currently available databases with NMR characteristics of metabolites.[39-41]

4. Discussion

Prediction of phenotypic trait values using genetic markers has been a central element in plant and livestock breeding for decades,[42-44] and more recently this strategy has emerged within human genetics attempting to accurately predict e.g. disease risk from DNA information.[11,45-47] However, the predictive value from genotyped genetic variants is often low,[11,48] and this is problematic when aiming to predict complex phenotypes such as many diseases, behaviors or production traits. Therefore, there is a potential to further optimize the applicability of these methods. Here, we obtained full metabolome profiles of males from 170 DGRP lines in four biological replicates to investigate if an endophenotype, in our case the metabolome, has improved predictive power compared with a situation with only genome information available.

We found that the *Drosophila* metabolome was highly variable with more than 39% of the NMR features having a significant heritability estimate (Fig. 2), displaying same level of genetically determined variability as the *Drosophila* transcriptome.[30] It has previously been shown that metabolome variation appears to have a genetic signature,[18] but also that the metabolome is highly variable among sexes[18,49] and change with age.[50-52]. Our findings confirm that variation in metabolite abundance is genetically controlled. The metabolome can therefore be influenced by evolutionary forces like any other phenotypic trait and this variation can be utilized e.g. in livestock and plant breeding where specific metabolites may be of interests.[53-55]

Metabolite-wide association studies, which is the mapping of metabolite QTLs, seeks the same as genome-wide and transcriptome-wide association studies, namely to identify genetic variants associated with variation in a functional character or an endophenotype.[56,57] Here we mapped the individual data points in the metabolome spectra with SNP genotypes and identified abundant metabolite feature – SNP associations (Supplementary Table S1). We identified more than 900 genes associated with metabolome variation, where the average number of associations per gene was 6. Four genes had more than 100 feature – SNP associations. The *Drosophila* gene *Coronin* was the gene containing most significant associations with metabolome variation (488 metabolomic features). *Coronin* is involved in muscle morphogenesis[58]. The human orthologue of *Coronin* is *coronin 1C* and this gene has previously been associated with lipoprotein and cholesterol levels[59,60]. The second most associated gene was *sidestep* (133 NMR features), which controls the migration of motor axons in the developing fly[61]. Across model organisms and humans *sidestep* has no apparent orthologous genes[62]. Another gene with many (123) associated NMR features was *CG43373*, which has the human ortholog adenylate cyclase 5 (*ADCY5*), that in multiple studies have been associated with type II diabetes mellitus,[63-65] body mass index,[66] blood glucose[67] and cholesterol levels[68,69]. The last gene with more than 100 associated NMR features is *CCHa2r* that encodes a neuropeptide[70]. In humans, *CCHa2r* is the bombesin receptor subtype 3, which regulates metabolic rate[71] and glucose metabolism.[72] These results clearly indicate that the human orthologues of the top associated genes are involved in metabolic processes supporting that the metabolomic feature-SNP associations are biological relevant and not being statistical artefacts. The genes shortly presented here and

other genes strongly associated with metabolome variation (Supplementary Table S1) are candidates for further studies such as knock in and knock out technologies to verify their importance at the functional phenotypic level.

Accurate phenotypic prediction of any trait requires large sample sizes to reliable estimate model parameters, and some measure that can describe the covariance structure among individuals, such as genetic variants or other molecular variation. Although the DGRP system appears to lack power for mapping and prediction studies because of the limited number of inbred lines, it gains statistical power because it is possible to obtain repeated measures on a large number of individuals with the same genetic background resulting in very accurate within-DGRP-line measures.[32]

The main aim of the current study was to investigate the predictive power of the *Drosophila* metabolome compared with prediction models using genomic polymorphism data. We showed that for four out of five quantitative traits using the *Drosophila* metabolome for phenotypic predictions significantly improved the accuracy of prediction compared to using the *Drosophila* genome (Fig. 3). The extent to which prediction was increased varied across traits, but the trend was clear across all five traits; partitioning the metabolome by highly correlated NMR featues increased the predictive performance (Fig. 3). Despite lack of significance for startle response (Fig. 3G), the increase was borderline significant (*P* value = 0.09).

Recently, Zhou et al. (2019)[18] measured metabolite variation in 453 metabolites using 40 DGRP lines, and concluded that if the sample size was larger the accuracy of metabolome predictions could be improved. This is exactly what we have shown here; that the metabolome can increase the accuracy of phenotypic prediction. The increased power relative to the previous study may originate not only from the larger numbers of lines, but also from the higher reproducibility of NMR metabolomics. Moreover, we have demonstrated that our novel approach, NMR cluster-guided phenotypic prediction, identifies metabolite modules of NMR features which are trait specific (Table 2 and Fig. 3), which significantly increased the prediction accuracy compared to when using the entire metabolome (Fig. 3).

Our results clearly demonstrate the added value of performing predictions of functional phenotypes using NMR metabolomics compared with SNP genotypes. These results, together with others, truly open the doors for applying metabolomics in different disciplines, for example in the human health sector or in animal breeding. Metabolites can be easily quantified in biofluids from livestock, human blood donors or patients that have blood samples taken on a regular basis. This entails a clear advantage over other methods in terms of translatability.[73] Recently, studies have shown that the metabolomic signatures of blood from cattle[74] and pigs[75] can be used to accurately separate animals by their feed efficiency which is one of the most economically important traits in livestock production. Studies of the human metabolome and its relevance to human health has also clearly increased the last five years,[76,77] and is very likely to be one of the cornerstones in an implementation of personalised medicine. With the emergence of large human cohort projects, like the UK Biobank[78], FinnGenn[79] and BioBank Japan[80] etc., the data samples are reaching a sample level that could provide ground-breaking research if full metabolomic profiles were obtained.

The fact that many of the NMR signals with high predictivity come from unidentified and/or larger metabolites of lower concentration shows that despite the inherently low sensitivity of NMR, the high

reproducibility of the method allows for a high accumulated sensitivity to be obtained for the combined data set. It also shows the need for a deeper investigation of the *Drosophila* metabolome by NMR. From a metabolic signalling perspective it also makes sense that it is the less abundant metabolites that are important in these processes.

In conclusion we have convincingly shown that metabolomic approaches have large potential for predicting functional phenotypes. Obviously, the generality and repeatability of these findings should be verified in different genetic backgrounds, in non-model species and in samples that are easy to generate from livestock, humans and plants. However, our main findings namely that metabolite profiles are highly heritable, that specific genes are associated with metabolome variation and that the metabolome predicts phenotypes more accurately than genomic data are robust and we do not see reasons why these results should be limited to the investigated population of *D. melanogaster*.

5. Data availability

The DGRP genotypes, chromosomal inversions, *Wolbachia* infection status, and the phenotypic values for startle response, starvation resistance and chill coma recovery can be obtained from http://dgrp2.gnets.ncsu.edu/. The adjusted metabolome data can be found in the supplementary material and the locomotor activity measurements in the origical publication.[81]

6. Acknowledgements

The DGRP lines were obtained from the Bloomington Drosophila Stock Center (NIH P40OD018537, http://flystocks.bio.indiana.edu). We thank Helle Blendstrup, Susan Marie Hansen and Michael Ørsted from Aalborg University for assistance in fly maintenance and sample collection. All of the computing for this project was performed on the GenomeDK cluster. We would like to thank GenomeDK and Aarhus University for providing computational resources and support that enabled us to perform the analyses presented in the paper. The study was supported by the Danish Natural Science Research Council through a grant to TNK (DFF-8021-00014B), and by a grant from the Lundbeck Foundation (R287-2018-735) to PDR.

7. References

1.   Crick, F. 1970, Central dogma of molecular biology. *Nature*, **227**, 561–3.

2.   Pinu, F. R., Beale, D. J., Paten, A. M., et al. 2019, Systems biology and multi-omics integration: Viewpoints from the metabolomics research community. *Metabolites*, **9**, 1–31.

3.   Hasin, Y., Seldin, M., and Lusis, A. 2017, Multi-omics approaches to disease. *Genome Biol.*, **18**, 1–15.

4.   Elmer, K. R. 2016, Genomic tools for new insights to variation, adaptation, and evolution in the salmonid fishes: a perspective for charr. *Hydrobiologia*, **783**, 191–208.

5.   Dekkers, J. C. M. 2012, Application of genomics tools to animal breeding. *Curr. Genomics*, **13**, 207–12.

6.   Buckler, E. S., Holland, J. B., Bradbury, P. J., et al. 2009, The genetic architecture of maize flowering time. *Science*, **325**, 714–8.

7.   Hayes, B., and Goddard, M. 2010, Genome-wide association and genomic selection in animal breeding. *Genome*, **53**, 876–83.

8.   Desta, Z. A., and Ortiz, R. 2014, Genomic selection: Genome-wide prediction in plant improvement. *Trends Plant Sci.*, **19**, 592–601.

9.   Hickey, J. M., Chiurugwi, T., Mackay, I., and Powell, W. 2017, Genomic prediction unifies animal and plant breeding programs to form platforms for biological discovery. *Nat. Genet.*, **49**, 1297–303.

10.   Grinberg, N. F., Orhobor, O. I., and King, R. D. 2019, An evaluation of machine-learning for predicting phenotype: studies in yeast, rice, and wheat. *Mach. Learn.*, **109**, 251–77.

11.   Schrodi, S. J., Mukherjee, S., Shan, Y., et al. 2014, Genetic-based prediction of disease traits: Prediction is very difficult, especially about the future. *Front. Genet.*, **5**, 1–18.

12.   Sun, S., Yang, F., Tan, G., et al. 2016, An extended set of yeast-based functional assays accurately identifies human disease mutations. *Genome Res.*, **26**, 670–80.

13.   Märtens, K., Hallin, J., Warringer, J., Liti, G., and Parts, L. 2016, Predicting quantitative traits from genome and phenome with near perfect accuracy. *Nat. Commun.*, **7**, 1–8.

14.   Scoriels, L., Salek, R. M., Goodby, E., et al. 2015, Behavioural and molecular endophenotypes in psychotic disorders reveal heritable abnormalities in glutamatergic neurotransmission. *Transl. Psychiatry*, **5**, e540.

15.   te Pas, M. F. W., Madsen, O., Calus, M. P. L., and Smits, M. A. 2017, The importance of endophenotypes to evaluate the relationship between genotype and external phenotype. *Int. J. Mol. Sci.*, **18**, 1–22.

16.   Hayes, B. J., Panozzo, J., Walker, C. K., et al. 2017, Accelerating wheat breeding for end-use quality with multi-trait genomic predictions incorporating near infrared and nuclear magnetic resonance-derived phenotypes. *Theor. Appl. Genet.*, **130**, 2505–19.

17.   Van Der Ende, M. Y., Said, M. A., Van Veldhuisen, D. J., Verweij, N., and Van Der Harst, P. 2018, Genome-wide studies of heart failure and endophenotypes: Lessons learned and future directions. *Cardiovasc. Res.*, **114**, 1209–25.

18.   Zhou, S., Morgante, F., Geisz, M., Ma, J., Anholt, R., and Mackay, T. 2019, Systems genetics of the Drosophila metabolome. *Genome Res.*, gr.243030.118.

19.   Wang, P. I., and Marcotte, E. M. 2010, It's the machine that matters: Predicting gene function and phenotype from protein networks. *J. Proteomics*, **73**, 2277–89.

20.   Harel, T., Peshes-Yaloz, N., Bacharach, E., and Gat-Viks, I. 2019, Predicting phenotypic diversity from molecular and genetic data. *Genetics*, **213**, 297–311.

21.   Azodi, C. B., Pardo, J., VanBuren, R., de Los Campos, G., and Shiu, S. H. 2020, Transcriptome-based prediction of complex traits in maize. *Plant Cell*, **32**, 139–51.

22.   Li, Z., Gao, N., Martini, J. W. R., and Simianer, H. 2019, Integrating gene expression data into genomic prediction. *Front. Genet.*, **10**, 1–11.

23.   Huang, W., Massouras, A., Inoue, Y., et al. 2014, Natural variation in genome architecture among 205 *Drosophila melanogaster* Genetic Reference Panel lines. *Genome Res.*, **24**, 1193–208.

24.   Mackay, T. F. C., Richards, S., Stone, E. A., et al. 2012, The *Drosophila melanogaster* Genetic Reference Panel. *Nature*, **482**, 173–8.

25.   Emwas, A.-H. M. 2015, The Strengths and Weaknesses of NMR Spectroscopy and Mass Spectrometry with Particular Focus on Metabolomics Research In: Bjerrum, J. T., (ed.), *Metabonomics: Methods and Protocols*.

Springer New York, New York, NY, pp. 161–93.

26. Kristensen, T. N., Henningsen, A. K., Aastrup, C., et al. 2016, Fitness components of *Drosophila melanogaster* developed on a standard laboratory diet or a typical natural food source. *Insect Sci.*, **23**, 771–9.

27. Savorani, F., Tomasi, G., and Engelsen, S. B. 2010, icoshift: A versatile tool for the rapid alignment of 1D NMR spectra. *J. Magn. Reson.*, **202**, 190–202.

28. Dieterle, F., Ross, A., Schlotterbeck, G., and Senn, H. 2006, Probabilistic quotient normalization as robust method to account for dilution of complex biological mixtures. Application in 1H NMR metabonomics. *Anal. Chem.*, **78**, 4281–90.

29. Rohde, P. D., Fourie Sørensen, I., and Sørensen, P. 2020, qgg: an R package for large-scale quantitative genetic analyses. *Bioinformatics*, **36**, 2614–5.

30. Huang, W., Carbone, M. A., Magwire, M. M., et al. 2015, Genetic basis of transcriptome diversity in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci.*, **112**, 6010–6019.

31. Anholt, R. R. H., and Mackay, T. F. C. 2018, The road less traveled: from genotype to phenotype in flies and humans. *Mamm. Genome*, **29**, 5–23.

32. Mackay, T. F. C., and Huang, W. 2018, Charting the genotype–phenotype map: lessons from the *Drosophila melanogaster* Genetic Reference Panel. *Wiley Interdiscip. Rev. Dev. Biol.*, **7**, 1–18.

33. Edwards, S. M., Sørensen, I. F., Sarup, P., Mackay, T. F. C., and Sørensen, P. 2016, Genomic prediction for quantitative traits is improved by mapping variants to gene ontology categories in *Drosophila melanogaster*. *Genetics*, **203**, 1871–83.

34. Speed, D., and Balding, D. J. 2014, MultiBLUP: Improved SNP-based prediction for complex traits. *Genome Res.*, **24**, 1550–7.

35. Rohde, P. D., Gaertner, B., Ward, K., Sørensen, P., and Mackay, T. F. C. 2017, Genomic analysis of genotype-by-social environment interaction for *Drosophila melanogaster*. *Genetics*, **206**, 1969–84.

36. Rohde, P. D., Østergaard, S., Kristensen, T. N., et al. 2018, Functional validation of candidate genes detected by genomic feature models. *Genes, Genomes Genet.*, **8**, 1659–68.

37. Sørensen, I. F., Edwards, S. M., Rohde, P. D., and Sørensen, P. 2017, Multiple trait covariance association test identifies gene ontology categories associated with chill coma recovery time in *Drosophila melanogaster*. *Sci. Rep.*, **7**, 2413.

38. Fang, L., Sahana, G., Ma, P., et al. 2017, Exploring the genetic architecture and improving genomic prediction accuracy for mastitis and milk production traits in dairy cattle by mapping variants to hepatic transcriptomic regions responsive to intra-mammary infection. *Genet. Sel. Evol.*, **49**.

39. Ulrich, E. L., Akutsu, H., Doreleijers, J. F., et al. 2008, BioMagResBank. *Nucleic Acids Res.*, **36**, 402–8.

40. Cui, Q., Lewis, I. A., Hegeman, A. D., et al. 2008, Metabolite identification via the Madison Metabolomics Consortium Database [3]. *Nat. Biotechnol.*, **26**, 162–4.

41. Wishart, D. S., Feunang, Y. D., Marcu, A., et al. 2018, HMDB 4.0: The human metabolome database for 2018. *Nucleic Acids Res.*, **46**, D608–17.

42. Meuwissen, T. H. E., Hayes, B. J., and Goddard, M. E. 2001, Prediction of total genetic value using genome-wide dense marker maps. *Genetics*, **157**, 1819–29.

43. Van Arendonk, J. A. M., Tier, B., and Kinghorn, B. P. 1994, Use of multiple genetic markers in prediction of breeding values. *Genetics*, **137**, 319–29.

44. Goddard, M. E., Wray, N. R., Verbyla, K., and Visscher, P. M. 2009, Estimating effects and making predictions from genome-wide marker data. *Stat. Sci.*, **24**, 517–29.

45. Wray, N. R., Goddard, M. E., and Visscher, P. M. 2008, Prediction of individual genetic risk of complex disease. *Curr. Opin. Genet. Dev.*, **18**, 257–63.

46. Wray, N. R., Kemper, K. E., Hayes, B. J., Goddard, M. E., and Visscher, P. M. 2019, Complex trait prediction from genome data: Contrasting EBV in livestock to PRS in humans. *Genetics*, **211**, 1131–41.

47. Hall, W., Morley, K., and Lucke, J. 2004, The prediction of disease risk in genomic medicine: Scientific prospects and implications for public policy and ethics. *EMBO Rep.*, **5**.

48. Patron, J., Serra-Cayuela, A., Han, B., Li, C., and Wishart, D. S. 2019, Assessing the performance of genome-wide association studies for predicting disease risk. *PLoS One*, **14**, 1–24.

49. Li, Z., Zhang, Y., Hu, T., et al. 2018, Differential metabolomics analysis allows characterization of diversity of metabolite networks between males and females. *PLoS One*, **13**, 1–10.

50. Hoffman, J. M., Soltow, Q. A., Li, S., Sidik, A., Jones, D. P., and Promislow, D. E. L. 2014, Effects of age, sex, and genotype on high-sensitivity metabolomic profiles in the fruit fly, *Drosophila melanogaster*. *Aging Cell*, **13**, 596–604.

51. Yoshida, R., Tamura, T., Takaoka, C., et al. 2010, Metabolomics-based systematic prediction of yeast lifespan and its application for semi-rational screening of ageing-related mutants. *Aging Cell*, **9**, 616–25.

52. Lawton, K. A., Berger, A., Mitchell, M., et al. 2008, Analysis of the adult human plasma metabolome. *Pharmacogenomics*, **9**, 383–97.

53. Gamboa-Becerra, R., Hernández-Hernández, M. C., González-Ríos, Ó., et al. 2019, Metabolomic markers for the early selection of coffea canephora plants with desirable cup quality traits. *Metabolites*, **9**, 214.

54. Browne, R. A., and Brindle, K. M. 2007, 1H NMR-based metabolite profiling as a potential selection tool for breeding passive resistance against Fusarium head blight (FHB) in wheat. *Mol. Plant Pathol.*, **8**, 401–10.

55. Goldansaz, S. A., Guo, A. C., Sajed, T., Steele, M. A., Plastow, G. S., and Wishart, D. S. 2017, Livestock metabolomics and the livestock metabolome: A systematic review. *PLoS One*, **12**, 1–26.

56. Holmes, E., Loo, R. L., Stamler, J., et al. 2008, Human metabolic phenotype diversity and its association with diet and blood pressure. *Nature*, **453**, 396–400.

57. Bictash, M., Ebbels, T. M., Chan, Q., et al. 2010, Opening up the "black box": Metabolic phenotyping and metabolome-wide association studies in epidemiology. *J. Clin. Epidemiol.*, **63**, 970–9.

58. Schnorrer, F., Schönbauer, C., Langer, C. C. H., et al. 2010, Systematic genetic analysis of muscle morphogenesis and function in *Drosophila*. *Nature*, **464**, 287–91.

59. Siewert, K. M., and Voight, B. F. 2018, Bivariate genome-wide association scan identifies 6 novel loci associated with lipid levels and coronary artery disease. *Circ. Genomic Precis. Med.*, **11**, e002239.

60. Wakil, S. M., Ram, R., Muiya, N. P., et al. 2016, A common variant association study reveals novel susceptibility loci for low HDL-cholesterol levels in ethnic Arabs. *Clin. Genet.*, **90**, 518–25.

61. Siebert, M., Banovic, D., Goellner, B., and Aberle, H. 2009, *Drosophila* motor axons recognize and follow a Sidestep-labeled substrate pathway to reach their target fields. *Genes Dev.*, **23**, 1052–62.

62. Hu, Y., Flockhart, I., Vinayagam, A., et al. 2011, An integrative approach to ortholog prediction for disease-focused and other functional studies. *BMC Bioinformatics*, **12**, 357.

63. Qi, Q., Stilp, A. M., Sofer, T., et al. 2017, Genetics of type 2 diabetes in U.S. Hispanic/Latino individuals:

Results from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Diabetes*, **66**, 1419–25.

64. Mahajan, A., Go, M. J., Zhang, W., et al. 2014, Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. *Nat. Genet.*, **46**, 234–44.

65. Bonàs-Guarch, S., Guindo-Martínez, M., Miguel-Escalada, I., et al. 2018, Re-analysis of public genetic data reveals a rare X-chromosomal variant associated with type 2 diabetes. *Nat. Commun.*, **9**, 1–14.

66. Locke, A. E., Kahali, B., Berndt, S. I., et al. 2015, Genetic studies of body mass index yield new insights for obesity biology. *Nature*, **518**, 197–206.

67. Manning, A. K., Hivert, M. F., Scott, R. A., et al. 2012, A genome-wide approach accounting for body mass index identifies genetic variants influencing fasting glycemic traits and insulin resistance. *Nat. Genet.*, **44**, 659–69.

68. Liu, D. J., Peloso, G. M., Yu, H., et al. 2017, Exome-wide association study of plasma lipids in >300,000 individuals. *Nat. Genet.*, **49**, 1758–66.

69. Hoffmann, T. J., Theusch, E., Haldar, T., et al. 2018, A large electronic-health-record-based genome-wide study of serum lipids. *Nat. Genet.*, **50**, 401–13.

70. Hansen, K. K., Hauser, F., Williamson, M., Weber, S. B., and Grimmelikhuijzen, C. J. P. 2011, The *Drosophila* genes CG14593 and CG30106 code for G-protein-coupled receptors specifically activated by the neuropeptides CCHamide-1 and CCHamide-2. *Biochem. Biophys. Res. Commun.*, **404**, 184–9.

71. Xiao, C., Piñol, R. A., Carlin, J. L., et al. 2017, Bombesin-like receptor 3 (Brs3) expression in glutamatergic, but not GABAergic, neurons is required for regulation of energy metabolism. *Mol. Metab.*, **6**, 1540–50.

72. Feng, Y., Guan, X. M., Li, J., et al. 2011, Bombesin receptor subtype-3 (BRS-3) regulates glucose-stimulated insulin secretion in pancreatic islets across multiple species. *Endocrinology*, **152**, 4106–15.

73. Fontanesi, L. 2016, Metabolomics and livestock genomics: Insights into a phenotyping frontier and its applications in animal breeding. *Anim. Front.*, **6**, 73–9.

74. Novais, F. J., Pires, P. R. L., Alexandre, P. A., et al. 2019, Identification of a metabolomic signature associated with feed efficiency in beef cattle. *BMC Genomics*, **20**, 1–10.

75. Carmelo, V. A. O., Banerjee, P., da Silva Diniz, W. J., and Kadarmideen, H. N. 2020, Metabolomic networks and pathways associated with feed efficiency and related-traits in Duroc and Landrace pigs. *Sci. Rep.*, **10**, 1–14.

76. Rangel-Huerta, O. D., Pastor-Villaescusa, B., and Gil, A. 2019, *Are we close to defining a metabolomic signature of human obesity? A systematic review of metabolomics studies*. *Metabolomics*. Springer US.

77. Zhang, X. W., Li, Q. H., Xu, Z. Di, and Dou, J. J. 2020, Mass spectrometry-based metabolomics in health and medical science: A systematic review. *RSC Adv.*, **10**, 3092–104.

78. Bycroft, C., Elliott, L. T., Young, A., et al. 2018, The UK Biobank resource with deep phenotyping and genomic data. *Nature*, **562**, 203–9.

79. Mars, N., Koskela, J. T., Ripatti, P., et al. 2020, Polygenic and clinical risk scores and their impact on age at onset and prediction of cardiometabolic diseases and common cancers. *Nat. Med.*, **26**, 549–57.

80. Nagai, A., Hirata, M., Kamatani, Y., et al. 2017, Overview of the BioBank Japan Project: Study design and profile. *J. Epidemiol.*, **27**, S2–8.

81. Rohde, P. D., Jensen, I. R., Sarup, P. M., et al. 2019, Genetic signatures of drug response variability in *Drosophila melanogaster*. *Genetics*, **213**, 633–50.

Tables

Table 1 Quantitative traits used in phenotypic predictions. Locomotor activity was assessed as distance moved per unit time (the trait was assessed with and without exposure to Ritalin (methylphenidate)). Chill come recovery is time to recover from a chill-induced coma. Startle response is a measured as the time taken to reach a certain distance following disturbance. Starvation resistance is assessed as time to death when deprived of nutrients.

Trait	Number of DGRP lines	Number of observations*	H^{2**}	Lines for predictions***
Locomotor activity [81]	172	30	0.43	169
Locomotor activity w/ Ritalin [81]	172	30	0.45	169
Chill coma recovery [24]	159	101	0.42	132
Startle response [24]	166	40	0.50	137
Starvation resistance [24]	197	49	0.52	166

* Average number of individual observations per DGRP line.

** Estimated broad-sense heritability

*** The number of DGRP lines that are in common between phenotype and the metabolomic data.

Table 2 Predictive NMR feature clusters.

Trait	Clusters	Cluster level
Locomotor activity	1	K100
Locomotor activity w/ Ritalin	74, 121	K200
Chill coma recovery	5, 27, 35	K50
Startle response	1, 2, 5, 14, 99, 112, 145, 187	K200
Starvation resistance	21, 65, 83, 95, 140	K200

NMR feature IDs for each cluster can be found in Supplementary Data File S2.

Figures

1: Compute all pairwise NMR feature correlations

2: Hierachical clustering of NMR features

3: Define K clusters

$$y = m^{K_1} + m^r + e$$

4: For each K cluster run MFBLUP with first component based on NMR features within a cluster, redo for all K clusters

5: Repeat step 2 and 3 for the seven levels of K clusters

- 25 clusters
- 50 clusters
- 75 clusters
- 100 clusters
- 125 clusters
- 150 clusters
- 200 clusters

Figure 1. Conceptual illustration of the NMR cluster-guided phenotypic predictions. All pairwise correlations among the NMR featues were computed, which was used in a hierachical clustering of NMR features. The dendrogram was then sequential cut into K clusters (25, 50, 75, 100, 125, 150 and 200 clusters), and each individual cluster was then used in the MFBLUP model. NMR features within one cluster was used to construct a metabolomic relationship matrix that was used as covariance matrix in the MFBLUP model. The MFBLUP model was fitted for all clusters within the seven levels of K clusters.

Figure 2. Genetic variation for the *D. melanogaster* metabolome. Panel **A** shows in solid blue line the average NMR intensity across all DGRP lines (intensity axis not shown) as function of chemical shift. For each NMR feature we estimated h^2; the points in grey represent non-significant estimates of h^2, and points in green are significant estimates of h^2. Panel **B** is a histogram of the significant heritability estimates.

Figure 3. Prediction accuracies using genomic and metabolomic data. Panel **A**, **C**, **E**, **G** and **I** shows the maximum mean prediction (+ standard error) accuracies for the different models and traits. GBLUP and MBLUP are based on single component prediction models, whereas the two MFBLUP models are based on two component. The maxium prediction accuracy obtained across all levels of maximum clusters ($K_{cl} = \{25, 50, 75, 100, 125, 200\}$) is shown in the MFBLUP bar. The prediction accuracy when combining all the significant clusters is shown in the MFBLUP-2 bar. Significant improved predictive performance is indicated by asterisks above the bars, see Supplementary Table S3 for all comparisons. Panel **B**, **D**, **F**, **H**, and **J** shows for each level of total clusters the mean prediction accuracy at each individual sub-cluster. The maxium prediction accuracy within each cluster level is indicated with colored squares and cluster number. Squares in grey illustrates multiple clusters at the same point (numbers show which). The individual cluster that across all cluster levels result in the highest prediction accuracy is indicated with a circle; which is the point that is used in panels **A**, **C**, **E**, **G** and **I** (MFBLUP).

Figure 4. Contributions to predictive clusters from the metabolite NMR-spectra. Clusters are indicated with colored dots. Panel **A** shows clusters: 1, 2, 5, 21, 32, 45, 65, 74, 83 and 95; panel **B** clusters: 112, 121, 129, 139, 145 and 154; panel **C** cluster: 171; and panel **D** cluster: 187. Selected major metabolites in these regions are identified. The location of the nicotinamide ribotide signals resonating at the highest ppm values are also indicated in panel **A**.