scGRNom: a computational pipeline of integrative multi-omics analyses for predicting cell-type disease genes and regulatory networks

Ting Jin[1,2,*], Peter Rehani[2,3*], Mufang Ying[4,*], Jiawei Huang[4], Shuang Liu[2], Panagiotis Roussos[5,6], Daifeng Wang[1,2,7,#]

[1]Department of Biostatistics and Medical Informatics, University of Wisconsin – Madison, Madison, WI, 53706, USA
[2]Waisman Center, University of Wisconsin – Madison, Madison, WI, 53705, USA
[3]Department of Integrative Biology, University of Wisconsin - Madison, Madison, WI, 53706, USA
[4]Department of Statistics, University of Wisconsin - Madison, Madison, WI, 53706, USA
[5]Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA
[6]Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, USA
[7]Department of Computer Sciences, University of Wisconsin – Madison, Madison, WI, 53706, USA

[*]equal contribution
[#]corresponding author
Short title: cell-type disease genes and networks from multi-omics
Keywords: Single-cell genomics; single-cell multi-omics integration; cell-type gene regulatory network; cell-type disease risk genes; cross-disease functional genomics; schizophrenia; Alzheimer's disease

Abstract

Understanding cell-type-specific gene regulatory mechanisms from genetic variants to diseases remains challenging. To address this, we developed an open-source computational pipeline, scGRNom, to predict the cell-type disease genes and regulatory networks from multi-omics data, including cell-type chromatin interactions, epigenomics, and single-cell transcriptomics. With applications to Schizophrenia and Alzheimer's Disease, our predicted cell-type regulatory networks link transcription factors and enhancers to disease genes for excitatory and inhibitory neurons, microglia, and oligodendrocytes. The enrichments of cell-type disease genes reveal cross-disease and disease-specific functions and pathways. Finally, machine learning analysis found that cell-type disease genes shared by diseases have improved clinical phenotype predictions.

Background

Recent Genome Wide Association Studies (GWAS) studies have identified a variety of genetic risk variants associated with multiple brain diseases. For example, a recent study found 109 pleiotropic loci significantly associated with at least two brain disorders [1]. Many cross-disease common genetic risk factors have revealed many shared functional consequences in clinical presentations [2]. Recent studies have also revealed shared symptoms at both psychiatric and physical levels between neurodegenerative and neuropsychiatric diseases [3]. For instance, 97% of Alzheimer's disease patients develop neuropsychiatric symptoms throughout the disease [4]. Besides, additional insights into each disease's progression and causes have further demonstrated the highly interlinked nature of both disease types [5]. However, our understanding of the molecular mechanisms of genetic variants between diseases remains elusive, particularly at the cell-type levels.

Alzheimer's disease (AD) and Schizophrenia (SCZ) are representative neurodegenerative and neuropsychiatric diseases, respectively. Both are significantly associated with genetic variants and have complex underlying cellular and molecular mechanisms from genotype to phenotype [6,7]. Notably, AD is physiologically characterized by accumulations of amyloid beta plaques and neurofibrillary tau protein tangles in the brain. Amyloid beta plaques primarily originate from the apolipoprotein E encoding gene APOE and its multiple variants. The APOE gene is a single step in the broader amyloidogenic processing pathway (APP), and additional genes involved in the process contribute to the regulation of amyloid beta production. Much work has identified major genes of interest involved in the APP. However, a distinct need still exists to further explore these disease loci to understand better the interplay between their regulatory elements and eventual amyloid beta creation and accumulation. Similarly, neurofibrillary tau tangles are associated with many genetic loci and require a study of the highly complex molecular mechanisms required to achieve disease pathology. Further, the downstream effects from both amyloid beta and neurofibrillary tangles within and between various cell types add additional complexity toward linking specific regulatory events and elements with clinical pathology [8,9].

Also, SCZ is a neuropsychiatric disorder characterized by disruptions in dopamine, glutamate, and GABA-based receptor signaling pathways. Pathologically, the direct connection is less clear between known molecular abnormalities and observed physical changes through neurological imaging studies. Thus, attempting to understand interactions between activation of known risk genes and higher-level pathway disruptions may help elucidate causes for structural shifts in SCZ patients. At a psychiatric level, the alterations to various cortical structures create multiple forms of symptoms, including positive (e.g., Hallucinogenic episodes) and negative (e.g., Anti-social tendencies). Finally, GWAS for AD cohorts has revealed multiple conserved genetic loci that could encode shared risk factors between the two diseases [10]. Clinical presentations, specifically psychiatric effects, create a crucial point of intersection to be explored where general psychosis was found in up to 60% of AD patients, including hallucination events as well as other effects mirroring those of the positive symptoms found in SCZ patients [11]. Thus, studying shared risk variants and genes between both diseases may help elucidate functional genomics of interest in both diseases. And it can further uncover cross-disease and disease-specific mechanisms between neurodegenerative and neuropsychiatric diseases.

81
Recently, advances in single-cell sequencing technologies have generated a great deal of
excitement and interest in studying functional genomics at cellular resolution. For example,
scRNA-seq and scATAC-seq techniques have measured the transcriptomics and epigenomics of
individual cells in the human brain [12,13]. Further computational analyses have clustered cells
into many cell types [12]. The cells in the same type share similar transcriptional activities such
as gene expression and genomic functions. Differential gene expression across cell types is a
complex, multi-gene dynamic process that tightly regulates and controls functions and is
governed by gene regulatory factors such as transcriptional factors (TFs) and non-coding
regulatory elements. These factors cooperate as a gene regulatory network (GRN) to facilitate the
correct cellular and molecular functions on the genome-scale. Disrupted cooperation can give
rise to abnormal gene expression, such as those present in diseases. Therefore, GRN has been
used as robust system to infer genomic functions and molecular mechanisms, especially for
human diseases [14].
95
Recent analyses have also revealed that brain disease risk variants are located in non-coding
regulatory elements (e.g., enhancers). The risk genes likely have cell-type-specific effects for
both neuronal and non-neuronal cell types [15,16]. Besides, recent single-cell studies suggest
changes to cell-type-specific gene expression in brain diseases [8,9]. However, our
understanding of the gene regulatory networks driving cell-type and disease-specific gene
expression, especially across diseases, remains elusive. To address this issue, ones developed
several computational methods to predict cell-type GRNs [17], such as PIDC [18], GENIE3 [19],
and GRNBoost2 [20]. However, these methods typically only use single omics (e.g.,
transcriptomics) and predict networks based on statistical associations (e.g., co-expression),
providing insufficient mechanistic insights into gene regulation at the cellular resolution. For
instance, how the disease variants affect the transcription factor binding sites (TFBSs) on the
distal regulatory elements (e.g., enhancers) that control disease genes is still unclear, especially
at the cell-type level. Thus, it is essential to integrate emerging multi-omics data to understand
cell-type gene regulation, especially involving non-coding regulatory elements. Recent studies
have shown that integrating multi-omics data can reduce the impact of noise from a single omics
data and achieve better prediction accuracy [21].
112
To explore these ideas, we developed a computational pipeline, scGRNom, by integrating multi-
omics data to predict cell-type gene regulatory networks (GRNs) linking TFs, regulatory
elements (e.g., enhancers and promoters), and target genes. In particular, we applied scGRNom
to the multi-omics data at the cellular resolution, such as chromatin interactions, epigenomics,
and single-cell transcriptomics of primary cell types in the human brain, including different
excitatory and inhibitory neuronal types, microglia, and oligodendrocyte. Our predictions have
high overlapping with state-of-the-art methods for revealing TFs and target genes [17], but they
provide additional information on cell-type gene regulation, such as linking the regulatory
elements to the genes. We also found that the enhancers in our cell-type GRNs are enriched with
GWAS SNPs in human brain diseases, including psychiatric disorders and AD. Thus, we further
linked the GWAS SNPs that interrupt TFBSs to cell-type disease genes based on the cell-type
GRNs for SCZ and AD and found cross-disease and disease-specific genomic functions at the
cell-type level. Finally, we found that the cell-type disease genes shared by AD and SCZ have
improved predicting clinical phenotypes in AD, like disease staging and cognitive impairment.

Methods and Materials

Figure 1. The computational pipeline, scGRNom, for predicting the gene regulatory network via multi-omics data. The pipeline inputs the chromatin interactions (e.g., from Hi-C) of regulatory elements (e.g., enhancer-promoter), identifies the transcription factor binding sites (TFBSs) on interacting regulatory elements, predicts TF-target gene expression relationships (e.g., high coefficients from Elastic net regression), and finally outputs a gene regulatory network linking TFs (cyan), regulatory elements (purple) to target genes (green).

Predicting gene regulatory networks from multi-omics data

scGRNom is a computational pipeline in R (https://github.com/daifengwanglab/scGRNom) to (I) integrate multi-omics datasets for predicting gene regulatory networks linking transcription factors, non-coding regulatory elements, and target genes, and (II) identify cell-type disease genes and regulatory elements. First, for predicting gene regulatory networks from multi-omics, scGRNom has three steps (**Figure 1**), each of which is available as an R function:

Step1: Finding chromatin interactions. The function, *scGRNom_interaction,* inputs the chromatin interaction data (e.g., Hi-C) and predicts all possible interactions between enhancers and promoters in the data or the user-provided list - for example, those from Topologically Associating Domains (TADs) in Hi-C data. In addition, the function uses an R package, *GenomicInteractions* [22], to annotate interacting regions and link them to genes;

Step 2: Inferring the transcription factor binding sites on interacting regions. The function, *scGRNom_getTF,* infers the transcription factor binding sites (TFBS) based on consensus binding site sequences in the enhancers and promoters that potentially interact from the previous step *scGRNom_interaction*. It outputs a reference gene regulatory network linking these TF, enhancers, and/or promoters of genes. In particular, this function uses *TFBSTools* [23] to obtain the position weight matrices of the TFBS motifs from the JASPAR database [24] and predicts the TFBS locations on the enhancers and promoters via mapping TF motifs. The function further links TFs with binding sites on all possible interacting enhancers and promoters and outputs the

149   reference regulatory network. Furthermore, this function can run on a parallel computing version
150   via an R package, *motifmatchr* [25] for computational speed-up;
151
<u>Step 3: Predicting the gene regulatory network</u>. The function, *scGRNom_getNt* predicts the final
gene regulatory network based on the TF-target gene expression relationships in the reference
network. The reference gene regulatory network from the previous step provides all possible
regulatory relationships (wires) between TF, enhancers, and target genes. However, the
chromatin interacting regions are broad, so that many TFs likely have binding sites on them.
Also, changes in gene expression may trigger different regulatory wires in the reference network.
To refine our maps and determine the activity status of regulatory wires, we apply elastic net
regression, a machine learning method that has successfully modeled TF-target gene expression
relationships in the gene regulatory networks by our previous work [9]. Further, suppose the
chromatin accessibility information is available (e.g., from scATAC-seq data for a cell type). In
that case, the function can also filter the enhancers based on their chromatin accessibilities and
then output the network links only having the enhancers with high accessibility (e.g., overlapped
with scATAC-seq peaks). The parameter "open_chrom" inputs a list of user-defined chromatin
accessible regions.
166
Mathematically, given a gene expression dataset and a reference network (e.g., from
scGRNom_getTF), the function uses the TF expression to predict each target gene expression and
finds the TF with high regression coefficients. Given a target gene, let $y \in \mathcal{R}^n$ be a gene
expression vector modeling its expression values across *n* samples (e.g., *n* cells from single cell
data), and $X \in \mathcal{R}^{n \times m}$ be the gene expression matrix of *m* TFs across *n* samples. Those *m* TFs
should link to the target gene from the reference network, implying possible regulatory
relationships to the gene. The elastic net regression model then aims to find the optimal
coefficients for *m* TFs $c^* \in \mathcal{R}^m$ to solve the following optimization problem:

$$c^* = argmin_c(\|y - Xc\|^2 + \alpha\|c\|^2 + \beta\|c\|_1),$$

where α and β are parameters to adjust the contributions from L_2 and L_1 regularizations of $c \in$
$\mathcal{R}^m$. The samples are randomly divided into the training and testing sets by the parameter,
train_ratio (e.g., if train_ratio=0.7, then 70% training and 30% testing data). The optimal TF
coefficients, c^* are estimated by the training data. Also, for the model evaluation, the mean
square error (MSE) of the regression is calculated and reported by $\|y_{test} - X_{test}c^*\|^2$ using the
testing data. Further, the top TFs with high coefficients can be either selected by absolute
coefficient values (the parameter, cutoff_absolute) or a percentage from all *m* TFs (the
parameter, cutoff_percentage). Finally, the function outputs a final gene regulatory network
linking the top TFs as well as their linked enhancers (from the reference network, if any) to all
possible target genes.

Identifying cell-type disease genes and regulatory elements

In addition to predicting gene regulatory networks, the pipeline also provides another function,
scGRNom_disGenes, for identifying cell-type disease genes and regulatory elements (e.g.,
enhancers, promoters). This function's input includes a cell-type gene regulatory network and a
list of GWAS SNPs associated with a disease (**Figure 2**). The function uses an R package,
GenomicRanges [26], to overlap these disease SNPs with the enhancers and promoters of the
input cell-type gene regulatory network, and then find the ones that interrupt the binding sites of

all possible TFs (TFBSs) on the enhancers and promoters by *motifbreakR* [27]. It finally maps
the overlapped enhancers or promoters and TFs with interrupted TFBSs onto the input network
to find the linked genes and enhancers/promoters as the output cell-type disease genes and
regulatory elements.

Figure 2. Identification of cell-type disease genes and regulatory elements. The function of our pipeline, *scGRNom_disGenes,* predicts cell-type disease genes and regulatory elements. First, it inputs a cell-type GRN (top right) and disease-associated SNPs from GWAS (top left). Second, the function identifies the disease SNPs that interrupt the binding sites of the TFs on the enhancers or promoters that link to the target genes in the input GRN (Middle). Finally, the function outputs a list of the target genes and regulatory elements (enhancers or promoters) linked by disease SNPs (Bottom). Red star: SNP. Cyan triangle: TF. Purple ellipse: Enhancer. Green square: cell-type disease gene.

Application to multi-omics data and data processing for predicting cell-type gene regulatory networks in the human brain

We applied our computational pipeline to the multi-omics datasets at the cellular resolution in
the human brain. We predicted cell-type gene regulatory networks for major cell types:
excitatory and inhibitory neurons, microglia, and oligodendrocyte for the human brain [12]. The
excitatory neuronal types include Ex1, Ex2, Ex3e, Ex4, Ex5b, Ex6a, Ex6b, Ex8, and Ex9. The
inhibitory neuronal types include In1a, In1b, In1c, In3, In4a, In4b, In6a, In6b, In7, and In8. We
first input recently published cell-type chromatin interactome data in the human brain [15] to
scGRNom_interaction to reveal all possible interactions from enhancers to gene promoters in the
neuronal, microglia, and oligodendrocyte types. The genome annotation was from
TxDb.Hsapiens.UCSC.hg19.knownGene [28]. We then predicted a reference regulatory network
for each of these cell types using *scGRNom_getTF*. Finally, given a cell type, we input the single
cell gene expression data for the type and the reference network from *scGRNom_getTF* to
scGRNom_getNt for predicting the cell-type gene regulatory network (GRN).
211

Specifically, to make our predicted networks comparable across cell types, we made the
following data preparation and processing steps. First, although different studies have generated
increasing numbers of single cell data (e.g., for microglia [8]), we used the data from one study
(GSE97942) [12] that includes the gene expression data (UMI) of individual cells of primary cell
types, all from one postmortem tissue of human frontal cortex, aiming not to introduce additional
batches from different studies, which helps our further comparative analyses across cell types. In
addition, we filtered the genes that express in less than 100 cells. We then normalized gene
expression by *Seurat 4.0* [29] for further removing noises and batches across cell types. We also
applied the method *MAGIC* [30] to impute the single cell gene expression of all cells to address
potential dropout issues. Then for each cell type, we removed the lowly expressed genes with
log10(sum of imputed gene expression levels of the cells of the cell type+1) < 1. The numbers of
genes and cells for each cell type used for prediction are available in Supplemental File 1.
224
For predicting a cell type GRN using the Elastic net regression model, we randomly split the
cells into 70% training and 30% testing sets. We then selected the best Elastic net model that
minimized the mean square error (MSE). We then filtered the target genes based on the goodness
of fit by MSEs. In particular, we removed the target genes predicted by Elastic net regression
with MSE > 0.1 and also TF-target gene with absolute Elastic net coefficient < 0.01. The output
cell-type GRN consists of the network edges that link TFs, enhancers (if any), and target genes
(TGs), as well as the Elastic net coefficient of TF-TGs for each edge. Finally, we provided two
versions of each cell-type GRN (Supplemental File 2):
(I) the edges that include the enhancers that overlap the cell-type open chromatin regions
234   predicted by recent scATAC-seq data (broad excitatory and inhibitory neurons, microglia
235   and oligodendrocyte) [13];
236   (II) the edges that only include the top 10% TFs with absolute coefficients for each target
237   gene without considering cell-type open chromatin regions (for scATAC-seq data might
238   be likely noisy and no open chromatin regions available for neuronal subtypes).

Comparison with state-of-the-art methods

We compared our scGRNom predictions with existing state-of-the-art methods. In particular, we
input the single-cell gene expression data for each cell type to a recently published benchmark
framework, BEELINE [17], and predicted the cell-type regulatory networks using three of the
most consistent and highly accurate methods, PIDC, GENIE3, and GRNBoost2. These methods
only input gene expression data to predict all possible TF-target gene (TG) regulatory links based
on their expression relationships, without considering the regulatory elements or open chromatin
regions at the cell-type level. Thus, after applying to our processed single cell gene expression
data, they generated more network edges than us because scGRNom only keeps the TF-TG links
in which TFs have binding sites on the regulatory elements (e.g., enhancers and promoters).
Therefore, to make these networks comparable with ours, we extended our networks by selecting
up to the top 30% TFs for each TG and then checked if our TF-TG links predicted by the state-
of-the-art method (also up to top 30% TFs for each TG selected for each method). In particular,
given a cell-type GRN by scGRNom, we selected top K TFs per target gene (TG) to see if the
TF-TG pairs were also predicted by PIDC, GENIE3, or GRNBoost2 (again picking top K TFs
per TG). We varied K values from 0% to 30% and then calculated the percentages of the
scGRNom's TF-TG pairs that can be predicted by one of those methods (Figure S1).

GWAS SNPs and heritability enrichment analyses for the enhancers in the cell-type gene regulatory networks in the human brain

Genome Wide Association Studies (GWAS) have identified a variety of genetic risk variants, including single nucleotide polymorphisms (SNPs) that are significantly associated with diseases and phenotypes (i.e., the traits). For example, recent GWAS studies have identified many SNPs associated with Alzheimer's Disease (AD, 2357 credible SNPs) [6] and Schizophrenia (SCZ, 6105 credible SNPs) [7]. In addition to the credible SNPs, we also included additional SNPs with p<5e-5 from the AD and SCZ GWAS summary statistics for linking potential additional cell-type disease genes [31]. We applied the *partitioned linkage disequilibrium score regression (LDSC)* [32] to evaluate the heritability explained by the enhancers of our cell-type gene regulatory networks for GWAS SNPs. In particular, our heritability enrichment analyses used the GWAS summary statistics for the diseases or traits: Schizophrenia (SCZ) [7], Alzheimer's disease (AD) [6], Autism spectrum disorder (ASD) [33], Bipolar disorder (BPD) [34], Amyotrophic lateral sclerosis (ALS) [35], Major depressive disorder (MDD) [36], Intelligence [37], Multiple sclerosis (MS) [38], Parkinson's disease (PD) [39], Attention deficit hyperactivity disorder (ADHD) [40], Education [41], Type 2 diabetes (T2D) [42], Inflammatory bowel disease (IBD) [43], Coronary artery disease (CAD) [44]. Also, we provided the numbers of GWAS SNPs for AD and SCZ that interrupt the binding sites of at least one of all possible TFBSs and the binding sites of the regulatory TFs in each cell-type GRN (Table S1).

Identification and enrichment analyses of cell-type disease genes

Given a cell-type and a disease type (AD or SCZ), we input the cell-type GRNs (both versions) and the GWAS SNPs for the disease to the *scGRNom_disGenes* function for identifying the cell-type disease genes. We identified the cell type disease genes (AD and SCZ) for all the cell types as above, including excitatory and inhibitory neuronal subtypes, microglia, and oligodendrocyte (Supplemental File 3). We also merged the disease genes for excitatory/inhibitory neuronal subtypes as the broad excitatory/inhibitory neuronal disease genes. We used the web app, *Metascape* [45] to find the enrichments of cell-type disease genes such as KEGG pathways, GO terms, protein-protein interactions, and diseases (via DisGeNET). Enrichment p-values shown in this paper were adjusted using the Benjamin-Hochberg (B-H) correction. Also, we looked at the expression levels of our cell-type disease genes in the disease samples. In particular, we compared the published population gene expression data in AD for single cell expression [8].

Machine learning prediction of clinical phenotypes from cell-type disease genes

Finally, we used the machine learning approach to predict clinical phenotypes from our cell-type disease genes using the population data of the ROSMAP project, an independent AD cohort [46]. Given a clinical phenotype, we assume that $X_i \in R^d$ represents the expression data of d disease genes for the i^{th} individual in the cohort and $Y_i \in \{0,1\}$ represents the binarized class of the i^{th} individual's phenotype with $i \in 1, \dots, n$ individuals for training. We then found the optimal logistic regression model with the parameters $\{\beta_0^*, \beta_1^* \in R^d\}$ to classify the phenotype from the disease gene expression data via minimizing the following loss function:

296
$$\{\beta_0^*, \beta_1^*\} = argmax_{\beta_0,\beta_1} \sum_{i=1}^{n} -Y_i \log(\beta_0 + \beta_1^T X_i) - (1 - Y_i) \log(1 - \beta_0 - \beta_1^T X_i)$$

, where $(.)^T$ is the transpose operation. Also, we performed Cross-Validation (K=5) for the
individual samples with 80% training and 20% testing sets. We also balanced the class sample
size in each training set by the weighting method [47] so that the baseline of the classification
accuracy is 50% for two classes. We used the individuals from the training sets to train the
classification model. We then used the individuals from the testing sets to evaluate the
classification performance, i.e., accuracy. In particular, we predicted four clinical phenotypes in
ROSMAP: Braak Stages that measure the severity of neurofibrillary tangle (NFT) pathology
(Braak early stages (0,1,2,3) vs. late stages (4, 5, 6)), CERAD scores that measure neuritic
plaques (no AD vs. AD), the diagnosis of cognitive status (DCFDX, no or mild cognitive
impairments (1,2,3) vs. Alzheimer's dementia (4, 5)) and the cognitive status at the time of death
(COGDX, no or mild cognitive impairments (1,2,3) vs. Alzheimer's dementia (4, 5)).

Results

Predicting cell-type gene regulatory networks in the human brain

We applied the scGRNom pipeline to the multi-omics data for the human brain, including cell-
type chromatin interactions [15], transcription factor binding sites [23], single-cell
transcriptomics [9] and recent cell-type open chromatin regions [13] (Methods and Materials). In
particular, we predicted the cell-type gene regulatory networks (GRNs) for both glial and
neuronal types, including microglia, oligodendrocyte, excitatory neuronal subtypes (Ex1, Ex2,
Ex3e, Ex4, Ex5b, Ex6a, Ex6b, Ex8 and Ex9, and inhibitory neuronal subtypes (In1a, In1b, In1c,
In3, In4a, In4b, In6a, In6b, In7, and In8). Each cell-type GRN links TFs to enhancers to target
genes (TGs), and has two versions that (I) only include the network edges with open cell-type
enhancers predicted by recent scATAC-seq data [13] and (II) only include the edges with top
10% TFs with highest absolute coefficients for each target gene without using cell-type open
chromatin regions (due to scATAC-seq data might be likely noisy and not specific to neuronal
subtypes). For instance, we found that the microglia GRN with open chromatin regions consists
of 47353 edges linking 180 TFs, 1893 microglia open enhancers, 1236 TGs. All cell-type GRNs
are available in Supplemental File 2. The network statistics such as numbers of cells, edges, TFs,
enhancers and TGs for all cell types are available in Supplemental File 1.
325
Our GRNs reveal many known cell-type-specific regulations. **Figure 3A** visualizes the
subnetworks among TFs for select cell types (i.e., TGs are also TFs). For example, two known
TFs, MEF2A and RFX3, that control microglia phenotypes play hub roles in the microglia
network [48,49]. The nuclear factors NFIA, NFIX, and FOXP1 controlling neural differentiation
and gliogenesis are also hub genes in the oligodendrocyte network [50–52]. MEF2C regulating
inhibitory and excitatory synapses is a central node in both excitatory and inhibitory networks
(e.g., Ex1 and In6b) [53]. In addition to cell-type TFs, we also observed the cell-type-specific
expression relationships between TFs and TGs (high correlation). For instance, in **Figure 3B**,
E2F3-LRRK2, STAT2-FBXO32, IRF2-DYNC1, and ATF4-EPB41L1 show cell-type-specific
expression relationships across the cells of microglia, oligodendrocyte, Ex1, and In6b types.
336

In addition, we compared our predicted cell-type gene regulatory networks with existing state-of-
the-art methods for predicting cell-type gene regulatory networks, particularly those that are
consistent and highly accurate PIDC, GENIE3, and GRNBoost2 benchmarked by BEELINE [17]
(Materials and Methods). The percentages of the overlapped TF-TG links of the cell-type
network between scGRNom (both versions) and the state-of-the-art methods are over 50%
(Methods, **Figure S1**). This suggests a high consistency between scGRNom and these methods.
However, these methods predicted TF-TG links without providing information on regulatory
elements like enhancers. Thus, we looked further at the enhancers in our cell-type GRNs and
found that they have significantly high heritability enrichments for GWAS SNPs of multiple
brain diseases and traits ($p<0.05$, Materials and Methods). For example, the enhancers in our
excitatory-neuron and inhibitory-neuron GRNs have high enrichments for AD, SCZ, Major
depressive disorder, intelligence, education (**Figure 4A, Figure S2**). Besides, these brain-cell-
type enhancers do not have significant enrichment of GWAS SNPs for non-brain diseases.
Therefore, the heritability enrichment analysis suggests that the enhancers in our cell-type GRNs
have potential pleiotropic roles associated with multiple brain diseases or traits. Finally, we also
compared our cell-type networks with public GRN databases such as TRRUST [54], Dorothea
[55], and RegNetwork [56]. Those GRNs were primarily inferred by integrating different studies
from the literature (e.g., via physical interactions, co-expressed genes at the bulk tissue level) and
thus may not be specific for the neuronal and glial cell types in the human brain. However, we
still found many overlapped network edges (TF-TG) with our cell-type gene regulatory
networks. For example, we found that 319 TRRUST edges, 5935 Dorothea edges, 106
RegNetwork edges overlap with at least one of our cell-type GRNs, implying their potential
associations with the human brain's cell types.
360

Figure 3. Cell-type gene regulatory networks (GRNs) in the human brain. (**A**) The subnetworks of select cell-type GRNs among TFs (i.e., TG is also TF): Ex1 (Top left), Microglia (Top right), In6b (Bottom left), Oligodendrocyte (Bottom right). (**B**) The expression levels of the cells for select TFs and TGs across the four cell types in (**A**). TF: x-axis. TG: y-axis. The select TF-TG pairs from top to bottom are E2F3-LRRK2, STAT2-FBXO32, IRF2-DYNC1, and ATF4-EPB41L1. The red darkness corresponds to the expression level.

Identifying cell-type disease genes in AD and SCZ for neuronal and glial
types
We used these cell-type GRNs to link GWAS SNPs with disease risk genes for each cell type,
advancing knowledge on cross-disease and disease-specific interplays among genetic,
transcriptional, and epigenetic risks at cellular resolution. In particular, we chose SCZ and AD,
two majorly represented neuropsychiatric and neurodegenerative diseases with potential
convergent underlying mechanisms [4], and we linked a number of cell-type disease genes
(Methods, Supplemental File 3) and performed their enrichments (Supplemental File 4). We
found that many disease genes present in one or a few cell types only, suggesting potential cell-
type-specific contributions to AD and SCZ (**Figure S3**). As shown in **Figure 4B**, the cell-type
AD genes are also significantly enriched with known disease genes for AD and other dementia
diseases such as brain atrophy and cerebral amyloid angiopathy (p<0.01). For example, the
microglia AD genes, including APP, CLU, BACE2, and BIN1, are specifically enriched for other
diseases such as neurofibrillary degeneration, Parkinson's dementia, and aging memory
impairment. Also, the cell-type SCZ genes are enriched for various disorders such as
neurodevelopmental, mental and mood disorders, and depression ($p < 0.01$, **Figure 4C**).
Furthermore, many cell-type disease genes have corresponding expression activities in the
disease samples. For example, 72 excitatory AD genes (86%) and 65 inhibitory AD genes (80%)
(plus 22 oligodendrocyte AD genes and three microglia AD genes) are significantly differentially
expressed in the corresponding cell types in AD individuals, respectively ($p < 0.05$) [8].
382
The functional enrichment analyses for our cell-type disease genes also uncover known genomic
functions and pathways at the cell-type level. For instance, the microglia AD genes are enriched
with amyloid beta formation and clearance [57], MAPK signaling [58] and neuron death [59]
(**Figure 4D**), and the oligodendrocyte AD genes are enriched with Tau protein binding [60]. This
is vital to understanding a multitude of diseases that commonly demonstrate atrophy of cortical
tissue as a hallmark feature. In SCZ genes, we also found that multiple key hallmark pathways
were enriched, such as dopaminergic synapse [61], trans-synaptic signaling [62], and synapse
organization [63] for excitatory SCZ genes (**Figure 4E**). For inhibitory SCZ genes, we observed
that MAPK family signaling [64], regulation of NMDA receptor activity [65], dopaminergic
synapses [61] and neurotransmission [65] are enriched.
393
394
395

Figure 4. Enrichments of cell-type enhancers and disease genes. (**A**) Partitioned heritability enrichment of GWAS SNPs associated with various diseases and traits (bar) on the enhancers of In1a GRN. Bar height is -log10(p value) of the enrichment. The diseases and traits are Schizophrenia (SCZ), Alzheimer's disease (AD), Autism spectrum disorder (ASD), Bipolar disorder (BPD), Amyotrophic lateral sclerosis (ALS), Major depressive disorder (MDD), Intelligence, Multiple sclerosis (MS), Parkinson's disease (PD), Attention deficit hyperactivity disorder (ADHD), Education, Type 2 diabetes (T2D), Inflammatory bowel disease (IBD), Coronary artery disease (CAD). The red line represents p value = 0.05. (**B**) The disease enrichments of cell-type AD genes. Rows are diseases/traits. Columns are cell types. Darkness is proportional to -log10(p) of the enriched term. (**C**) The disease enrichments of cell-type SCZ genes. Rows are diseases/traits. Columns are cell types. Darkness is proportional to -log10(p) of the enriched term. (**D**) The enrichments of functions and pathways (e.g., GO, KEGG, REACOME) in the microglia AD genes. Bar darkness and length are proportional to -log10(p) of the enriched term. (**E**) The enrichments of functions and pathways (e.g., GO, KEGG, REACOME) in the excitatory neuronal SCZ genes. Bar darkness and length are proportional to -log10(p) of the enriched term.

Comparative analyses reveal the interplays between genomic functions, pathways, cell types, and diseases

In addition to cell-type-specific pathways in these diseases, we also identified those involving multiple cell types in each disease, implying that potential cell-type interactions drive the disease pathology. For example, the enrichment of SCZ primarily includes changes in synapse structures and cell shaping and differentiation (**Figure 5A**). Clinically, this is consistent with the consensus that SCZ is strictly neuropsychiatric as opposed to degenerative. In particular, cell morphogenesis and regulation of neuron differentiation are enriched in all four major cell type SCZ genes ($p < 0.01$). Early life neurodevelopmental genetic markers may suggest causal links with alterations in hippocampal cell differentiation points on the front of cell morphogenesis, leading to cascades of downstream effects [66]. This has primarily been studied and modeled within the scope of iPSC-based analyses, which make correlations and connections to the clinical presentation more difficult due to the additional abstraction from the standard pathology-based analysis. Also, the BDNF signaling pathway that potentially relates to intercellular communications is enriched as well in multiple cell types ($p < 0.01$) [67]. Finally, we also observed that protein-protein interactions (PPIs) are enriched among the disease genes of the cell types at a higher level. As shown in **Figure 5B**, the SCZ genes for dopaminergic synapse, NMDA receptors, glutamate binding, and activation are shared by multiple cell types and have strong PPIs, implying protein-level cross-type coordination [68]. In AD, multiple pathways were significantly enriched across various cell types (**Figure 5C**). For instance, the catabolic process for the AD key player, amyloid precursor protein (APP) is enriched with both glial and neuronal types ($p < 0.01$) [69].

Furthermore, we found that cross-disease conserved functions/pathways are involved in one or multiple cell types, revealing potential novel functional interplays across cell types and diseases (**Figure 5D**). For example, the cell morphogenesis involved in differentiation is enriched in both AD and SCZ neuronal genes. Another example is that the vesicle-mediated transport is enriched for both AD and SCZ microglia genes. In total, we found 11 microglia genes shared by AD and SCZ. In particular, the VEGF signaling pathway is enriched in the SCZ microglia genes. In the general theme of AD pathology, increased VEGF expression has been linked to worse cognitive outcomes in post-mortem analysis [70]. Similarly, multiple meta analyses have revealed

differential expression levels between healthy controls and SCZ patients [71]. However, little has
been done to link potential gene function to cell-type level interactions and pathways. Here, these
SCZ-AD shared microglia genes may help explain shared higher-level dysfunction between both
diseases as evidenced by higher expression. Also, we found that some functions involve different
cell types across diseases. The dendrite development has been found in both SCZ and AD
pathology [72,73]. We found that it is mainly enriched with microglia in AD but neuronal types
in SCZ.
434
More interestingly, when exploring the interactions between cell types that change between
diseases, the disease-specific pathologies enter to explain the cause of discrepancies. In
particular, for AD, it is shown that phagocytic microglia are activated during the early stages of
synaptic decline, leading to eventual neuroinflammation and programmed cell death [74]. For
SCZ, the oligodendrocyte enrichment reveals similar intercellular mechanisms between
excitatory and inhibitory neurons, specifically those regulating neuron differentiation (**Figure
5A**), providing potential direction for future exploration and validation of the communication
role of oligodendrocytes [75].

Figure 5. Cross-cell-type conserved and cell-type-specific functions, pathways and protein-protein interactions in Schizophrenia and Alzheimer's Disease. Darkness in heatmaps is proportional to -log10(p) of the enrichment. (**A**) The enrichments of select conserved and specific functions/pathways (e.g., GO, KEGG, REACTOME) across cell-type disease genes in Schizophrenia. (**B**) The enrichments of protein-protein interactions among major cell-type disease genes in Schizophrenia: Excitatory neuron (broad, red), Inhibitory neuron (broad, blue), Microglia (green), Oligodendrocyte (Purple). (**C**) The enrichments of select conserved and specific functions/pathways across cell-type disease genes in Alzheimer's Disease. (**D**) The enrichments of select conserved and specific functions/pathways across cell-type disease genes between Schizophrenia and Alzheimer's Disease.

Prediction of clinical phenotypes from cell-type disease genes

Finally, we want to investigate the clinical applications of our cell-type disease genes. To this end, we looked at the population-level gene expression data for Alzheimer's disease in the ROSMAP cohort [46]. In particular, we first found that many cell-type AD genes have significantly associated expression levels with clinical phenotypes across individuals in ROSMAP. For example, out of 195 cell-type AD genes, we found that 72 genes are significantly associated with the Braak Stages that measure the severity of neurofibrillary tangle (NFT) pathology (ANOVA $p < 0.05$), 78 genes with the CERAD scores that measure neuritic plaques (ANOVA $p < 0.05$), 89 genes with the diagnosis of cognitive status (DCFDX) (ANOVA $p < 0.05$), 73 genes with the cognitive status at the time of death (COGDX) (ANOVA $p < 0.05$) and 92 genes with the Mini-Mental State Examination (MMSE) scores (Pearson correlation $p < 0.05$). In total, 135 cell-type AD genes were significantly associated with at least one clinical phenotype in ROSMAP (Supplemental File 5).

Figure 6. Prediction accuracy of AD clinical phenotypes from disease genes. The AD population data for prediction was from the ROSMAP cohort [46]. AD clinical phenotypes include Braak – stages that measure the severity of neurofibrillary tangle (NFT) pathology, Cerad – scores that measure neuritic plaques, Cogdx – cognitive status at the time of death, and Dcfdx - the diagnosis of cognitive status. The bar height represents the average accuracy of cross-validation (K=5) from the prediction using logistic regression (Methods). Red: scGRNom's cell-type disease genes shared by AD and SCZ (SCZ-AD genes). Green: AD genes from GWAS [6]. Blue: randomly select genes (same number as SCZ-AD genes).

In addition to statistically significant associations between our cell-type disease genes and clinical phenotypes, we also applied the machine learning approach to predict clinical phenotypes from these cell-type disease genes using ROSMAP data (Methods). Specifically, for each clinical phenotype, we used the logistic regression model to classify individual states of the phenotype (as classes) from their expression data of 53 AD-SCZ shared cell-type genes. We

performed the cross-validation (K=5) for the individual samples with 80% training and 20%
testing sets. As shown in Fig. 6, our average accuracy values for classifying all four major
clinical phenotypes: Braak Stages, CERAD scores, DCFDX status, COGDX status are much
higher than the baselines (50%), the random select genes, and AD genes from the latest GWAS
study [6]. This suggests that using cell-type disease genes shared by AD and SCZ has improved
predicting those clinical phenotypes, especially for cognitive related ones in AD.

Discussion

This paper focuses on the scGRNom's application to single cell data for Alzheimer's and
schizophrenia. However, scGRNom is general-purpose for understanding functional genomics
and gene regulation in other diseases. Besides the cell types, the pipeline also predicts the gene
regulatory networks for cell clusters (unknown cell types) and bulk tissue types. Furthermore,
recent eQTL studies have identified various SNPs associating with gene expression in multiple
brain tissue types using the population data such as GTEx [76] and PsychENCODE [9].
Although those brain eQTLs suggest the SNP-gene association at the bulk tissue level, we still
found that several eQTLs in PsychENCODE match our linked GWAS SNPs and cell-type
disease genes, e.g., SNPs chr2:127846321 for and chr20:43598154 for STK4, two microglia
disease genes for AD and SCZ, respectively. This suggests potential cell-type effects of these
human brain eQTLs. Thus, increasing single cell data at the population level allows us to predict
the cell-type eQTLs [77], which will likely help understand cell-type gene regulation and refine
linking disease genes at the cell-type level.
482
For linking disease genes, we primarily used the interrupted TFBSs by GWAS SNPs. Future
studies utilizing scGRNom would be able to take advantage of the ever-growing number of
GWAS for a wide variety of diseases as well as single cell data. However, additional information
can also help link GWAS SNPs to disease genes. For example, existing tools such as FUMA [78]
have linked GWAS loci to genes by integrating information from multiple resources, providing
more functional linkages from genotype to genes to phenotype in human diseases. Thus,
incorporating multiple GWAS data (e.g., various brain regions) exposes key areas of observed
phenotypes. Previous studies have demonstrated the caution that must be exercised when
attempting to correlate GWAS data with clinical phenotypes, and methods such as our analysis
mitigate these effects [16]. A similar methodology as outlined could be used where common loci
within each set of summary statistics are incorporated and established before integration into the
cell type GRNs, thus linking neuronal spatial information with known mutation sites in patients
along with potentially cell-type-specific functionality. Expanding past the cell types examined
here into additional multicellular analyses is also possible given expanded interactome data.
Notably, this would allow for further investigation of complex neuropsychological diseases and
cases where the line between different clinical classifications becomes blurred and leads to
additional complications about clinically relevant genetic therapies. One such example includes
Autism Spectrum Disorder (ASD), where clinical presentations can vary in multiple axes of
severity, creating a broad spectrum of phenotypes. In such cases, potentially linking specific
symptoms or aspects of a particular subset of ASD to particular brain regions and cell types
allows for a better-informed picture of functional consequences associated with genetic mutation
sites. Such connections could aid in determining genetic risk factors associated with variations in
edge case patients; they also create the opportunity to take advantage of Induced Pluripotent

Stem Cell (iPSC) technology using genetic engineering technologies to create point mutations
matching computationally identified genes. Moreover, because the single cell multi-omics data
we used for predicting cell-type GRNs are not specific for particular diseases (AD or SCZ), we
used the SNPs disrupting all possible TFs on the enhancers and promoters from our cell type
GRNs to link at large cell-type disease genes. However, the scGRNom pipeline is general-
purpose and able to work for incoming disease-specific single cell multi-omics, and link to cell-
type disease genes via interrupted regulatory TFs in the diseases.
513
Machine learning has also been widely used to analyze multi-omics, such as multiview learning
and deep learning [9,79]. Multiview learning has great potential for understanding functional
multi-omics and revealing nonlinear interactions across omics. Therefore, integrating such
emerging machine learning approaches will enable identifying different cross-omic patterns,
especially for increasing single-cell multi-omics data and providing more comprehensive
mechanistic insights in cell-type gene regulation and linking to disease genes. For example, this
means adding more omics such as methylation data that reflect epigenetic changes that may
occur due to wide variations of inherited and environmental factors [80]. At a deeper functional
level, variations in methylation have been attributed to alterations in splicing activity, ultimately
impacting the regulation and expression of key genes [81]. Additionally, integrating proteomic
data at a single-cell level enhances the broader picture formed through additional data sources
even further [82]. Lastly, expanding past simple methylation and proteomics allows for the
ability to include all forms of data incorporated through the single-cell cytometry [83].

Conclusions

We developed a computational pipeline, scGRNom, to integrate multi-omics data and predict
gene regulatory networks (GRNs), which link TFs, non-coding regulatory elements (e.g.,
enhancers), and target genes. With applications to the data from single-cell multi-omics of the
human brain, we predicted cell-type GRNs for both neuronal (e.g., excitatory, inhibitory) and
glial cell types (e.g., microglia, oligodendrocyte). Further, scGRNom can input cell-type GRNs
and disease risk variants to link disease genes at the cell-type level, such as brain diseases like
Alzheimer's and schizophrenia. These disease genes revealed conserved and specific genomic
functions across neuropsychiatric and neurodegenerative diseases, providing cross-disease
regulatory mechanistic insights at the cellular resolution. Although this paper focuses on
Alzheimer's and schizophrenia, scGRNom is general-purpose for understanding functional
genomics and gene regulation in other diseases, at either bulk tissue or cell-type levels. Finally,
scGRNom is open-source available at https://github.com/daifengwanglab/scGRNom.
540

Supplementary information

542   Supplemental materials – supplemental figures and table
543   Supplemental File 1 – statistics of cell-type gene regulatory networks
544   Supplemental File 2 – cell-type gene regulatory networks (TF, enhancer, target gene)
545   Supplemental File 3 – cell-type disease genes for AD and SCZ
546   Supplemental File 4 – enrichments of cell-type disease genes for AD and SCZ
547

548   Supplemental File 5 – cell-type disease genes associated with clinical phenotypes in AD

Data availability

Our pipeline for predicting gene regulatory networks via multi-omics is open-source available, as well as a tutorial at https://github.com/daifengwanglab/scGRNom. All data supporting this study are included in this paper, the supplemental files, and/or available from the corresponding author on request.

Author contributions

D.W. conceived and designed the study. T.J. and M.Y. implemented the software. T.J., P.R., M.Y., J.H., S.L., and D.W. analyzed the data. T.J., P.R., M.Y. , Pa.R., and D.W wrote the manuscript. All authors read and approved the final manuscript.

Competing interests

None declared.

Funding

This work was supported by the grants of National Institutes of Health, R01AG067025, R21CA237955 and U01MH116492 to D.W., U54HD090256 to Waisman Center, and the start-up funding for D.W. from the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin–Madison.

References

1. Cross-Disorder Group of the Psychiatric Genomics Consortium. Electronic address: plee0@mgh.harvard.edu, Cross-Disorder Group of the Psychiatric Genomics Consortium. Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. Cell. 2019;179:1469-1482.e11.

2. Brainstorm Consortium, Anttila V, Bulik-Sullivan B, Finucane HK, Walters RK, Bras J, et al. Analysis of shared heritability in common disorders of the brain. Science. 2018;360.

3. Ciccocioppo F, Bologna G, Ercolino E, Pierdomenico L, Simeone P, Lanuti P, et al. Neurodegenerative diseases as proteinopathies-driven immune disorders. Neural Regen Res. 2020;15:850–6.

4. Steinberg M, Shao H, Zandi P, Lyketsos CG, Welsh-Bohmer KA, Norton MC, et al. Point and 5-year period prevalence of neuropsychiatric symptoms in dementia: the Cache County Study. Int J Geriatr Psychiatry. 2008;23:170–7.

5. Cummings J, Ritter A, Rothenberg K. Advances in Management of Neuropsychiatric
Syndromes in Neurodegenerative Diseases. Curr Psychiatry Rep. 2019;21:79.

6. Jansen IE, Savage JE, Watanabe K, Bryois J, Williams DM, Steinberg S, et al. Genome-wide
meta-analysis identifies new loci and functional pathways influencing Alzheimer's disease risk.
Nat Genet. 2019;51:404–13.

7. Pardiñas AF, Holmans P, Pocklington AJ, Escott-Price V, Ripke S, Carrera N, et al. Common
schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong
background selection. Nat Genet. 2018;50:381–9.

8. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, et al. Single-cell
transcriptomic analysis of Alzheimer's disease. Nature. 2019;570:332–7.

9. Wang D, Liu S, Warrell J, Won H, Shi X, Navarro FCP, et al. Comprehensive functional genomic
resource and integrative model for the human brain. Science. 2018;362.

10. DeMichele-Sweet M a. A, Weamer EA, Klei L, Vrana DT, Hollingshead DJ, Seltman HJ, et al.
Genetic risk for schizophrenia and psychosis in Alzheimer disease. Mol Psychiatry. 2018;23:963–
72.

11. Murray PS, Kumar S, Demichele-Sweet MAA, Sweet RA. Psychosis in Alzheimer's disease.
Biol Psychiatry. 2014;75:542–52.

12. Lake BB, Chen S, Sos BC, Fan J, Kaeser GE, Yung YC, et al. Integrative single-cell analysis of
transcriptional and epigenetic states in the human adult brain. Nat Biotechnol. 2018;36:70–80.

13. Corces MR, Shcherbina A, Kundu S, Gloudemans MJ, Frésard L, Granja JM, et al. Single-cell
epigenomic analyses implicate candidate causal variants at inherited risk loci for Alzheimer's
and Parkinson's diseases. Nat Genet. 2020;52:1158–68.

14. Marbach D, Lamparter D, Quon G, Kellis M, Kutalik Z, Bergmann S. Tissue-specific regulatory
circuits reveal variable modular perturbations across complex diseases. Nat Methods.
2016;13:366–70.

15. Nott A, Holtman IR, Coufal NG, Schlachetzki JCM, Yu M, Hu R, et al. Brain cell type-specific
enhancer-promoter interactome maps and disease-risk association. Science. 2019;366:1134–9.

16. Sey NYA, Hu B, Mah W, Fauni H, McAfee JC, Rajarajan P, et al. A computational tool (H-
MAGMA) for improved prediction of brain-disorder risk genes by incorporating brain chromatin
interaction profiles. Nat Neurosci. 2020;

17. Pratapa A, Jalihal AP, Law JN, Bharadwaj A, Murali TM. Benchmarking algorithms for gene
regulatory network inference from single-cell transcriptomic data. Nat Methods. 2020;17:147–
54.

18. Chan TE, Stumpf MPH, Babtie AC. Gene Regulatory Network Inference from Single-Cell Data
Using Multivariate Information Measures. Cell Syst. 2017;5:251-267.e3.

19. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks from
expression data using tree-based methods. PloS One. 2010;5.

20. Moerman T, Aibar Santos S, Bravo González-Blas C, Simm J, Moreau Y, Aerts J, et al.
GRNBoost2 and Arboreto: efficient and scalable inference of gene regulatory networks.
Bioinforma Oxf Engl. 2019;35:2159–61.

21. Li Y, Wu F-X, Ngom A. A review on machine learning principles for multi-view biological data
integration. Brief Bioinform. 2018;19:325–40.

22. Harmston, N., Ing-Simmons, E., Perry, M., Baresic, A., Lenhard, B. GenomicInteractions: R
package for handling genomic interaction data [Internet]. 2020. Available from:
https://github.com/ComputationalRegulatoryGenomicsICL/GenomicInteractions/

23. Tan G, Lenhard B. TFBSTools: an R/bioconductor package for transcription factor binding
site analysis. Bioinforma Oxf Engl. 2016;32:1555–6.

24. Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, et al.
JASPAR 2020: update of the open-access database of transcription factor binding profiles.
Nucleic Acids Res. 2020;48:D87–92.

25. Schep, Alicia. motifmatchr: Fast Motif Matching in R [Internet]. 2019. Available from:
https://www.bioconductor.org/packages/release/bioc/html/motifmatchr.html

26. Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for
computing and annotating genomic ranges. PLoS Comput Biol. 2013;9:e1003118.

27. Coetzee SG, Coetzee GA, Hazelett DJ. motifbreakR: an R/Bioconductor package for
predicting variant effects at transcription factor binding sites. Bioinforma Oxf Engl.
2015;31:3847–9.

28. Carlson, Marc. TxDb.Hsapiens.UCSC.hg19.knownGene: Annotation package for TxDb
object(s) [Internet]. Bioconductor; 2015. Available from:
https://bioconductor.org/packages/release/data/annotation/html/TxDb.Hsapiens.UCSC.hg19.k
nownGene.html

29. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al. Comprehensive
Integration of Single-Cell Data. Cell. 2019;177:1888-1902.e21.

30. van Dijk D, Sharma R, Nainys J, Yim K, Kathail P, Carr AJ, et al. Recovering Gene Interactions
from Single-Cell Data Using Data Diffusion. Cell. 2018;174:716-729.e27.

31. Panagiotou OA, Ioannidis JPA, for the Genome-Wide Significance Project. What should the
genome-wide significance threshold be? Empirical replication of borderline genetic
associations. Int J Epidemiol. 2012;41:273–86.

32. Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh P-R, et al. Partitioning
heritability by functional annotation using genome-wide association summary statistics. Nat
Genet. 2015;47:1228–35.

33. Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with
shared effects on five major psychiatric disorders: a genome-wide analysis. Lancet Lond Engl.
2013;381:1371–9.

34. Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide
association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. Nat
Genet. 2011;43:977–83.

35. van Rheenen W, Shatunov A, Dekker AM, McLaughlin RL, Diekstra FP, Pulit SL, et al.
Genome-wide association analyses identify new risk variants and the genetic architecture of
amyotrophic lateral sclerosis. Nat Genet. 2016;48:1043–8.

36. Howard DM, Adams MJ, Clarke T-K, Hafferty JD, Gibson J, Shirali M, et al. Genome-wide
meta-analysis of depression identifies 102 independent variants and highlights the importance
of the prefrontal brain regions. Nat Neurosci. 2019;22:343–52.

37. Savage JE, Jansen PR, Stringer S, Watanabe K, Bryois J, de Leeuw CA, et al. Genome-wide
association meta-analysis in 269,867 individuals identifies new genetic and functional links to
intelligence. Nat Genet. 2018;50:912–9.

38. International Multiple Sclerosis Genetics Consortium, Wellcome Trust Case Control
Consortium 2, Sawcer S, Hellenthal G, Pirinen M, Spencer CCA, et al. Genetic risk and a primary
role for cell-mediated immune mechanisms in multiple sclerosis. Nature. 2011;476:214–9.

39. Blauwendraat C, Heilbron K, Vallerga CL, Bandres-Ciga S, von Coelln R, Pihlstrøm L, et al.
Parkinson's disease age at onset genome-wide association study: Defining heritability, genetic
loci, and α-synuclein mechanisms. Mov Disord Off J Mov Disord Soc. 2019;34:866–75.

40. Demontis D, Walters RK, Martin J, Mattheisen M, Als TD, Agerbo E, et al. Discovery of the
first genome-wide significant risk loci for attention deficit/hyperactivity disorder. Nat Genet.
2019;51:63–75.

41. Okbay A, Beauchamp JP, Fontana MA, Lee JJ, Pers TH, Rietveld CA, et al. Genome-wide
association study identifies 74 loci associated with educational attainment. Nature.
2016;533:539–42.

42. Morris AP, Voight BF, Teslovich TM, Ferreira T, Segrè AV, Steinthorsdottir V, et al. Large-
scale association analysis provides insights into the genetic architecture and pathophysiology of
type 2 diabetes. Nat Genet. 2012;44:981–90.

43. Jostins L, Ripke S, Weersma RK, Duerr RH, McGovern DP, Hui KY, et al. Host-microbe
interactions have shaped the genetic architecture of inflammatory bowel disease. Nature.
2012;491:119–24.

44. Schunkert H, König IR, Kathiresan S, Reilly MP, Assimes TL, Holm H, et al. Large-scale
association analysis identifies 13 new susceptibility loci for coronary artery disease. Nat Genet.
2011;43:333–8.

45. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al. Metascape
provides a biologist-oriented resource for the analysis of systems-level datasets. Nat Commun.
2019;10:1523.

46. De Jager PL, Ma Y, McCabe C, Xu J, Vardarajan BN, Felsky D, et al. A multi-omic atlas of the
human frontal cortex for aging and Alzheimer's disease research. Sci Data. 2018;5:180142.

47. King G, Zeng L. Logistic Regression in Rare Events Data. Polit Anal. 2001;9:137–63.

48. Holtman IR, Skola D, Glass CK. Transcriptional control of microglia phenotypes in health and
disease. J Clin Invest. 2017;127:3220–9.

49. Zeisel A, Munoz-Manchado AB, Codeluppi S, Lonnerberg P, La Manno G, Jureus A, et al. Cell
types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. Science.
2015;347:1138–42.

50. Deneen B, Ho R, Lukaszewicz A, Hochstim CJ, Gronostajski RM, Anderson DJ. The
Transcription Factor NFIA Controls the Onset of Gliogenesis in the Developing Spinal Cord.
Neuron. 2006;52:953–68.

51. Zhou B, Osinski JM, Mateo JL, Martynoga B, Sim FJ, Campbell CE, et al. Loss of NFIX
Transcription Factor Biases Postnatal Neural Stem/Progenitor Cells Toward Oligodendrogenesis.
Stem Cells Dev. 2015;24:2114–26.

52. Pearson CA, Moore DM, Tucker HO, Dekker JD, Hu H, Miquelajáuregui A, et al. Foxp1
Regulates Neural Stem Cell Self-Renewal and Bias Toward Deep Layer Cortical Fates. Cell Rep.
2020;30:1964-1981.e3.

53. Harrington AJ, Raissi A, Rajkovich K, Berto S, Kumar J, Molinaro G, et al. MEF2C regulates
cortical inhibitory and excitatory synapses and behaviors relevant to neurodevelopmental
disorders. eLife. 2016;5.

54. Han H, Cho J-W, Lee S, Yun A, Kim H, Bae D, et al. TRRUST v2: an expanded reference
database of human and mouse transcriptional regulatory interactions. Nucleic Acids Res.
2018;46:D380–6.

55. Garcia-Alonso L, Holland CH, Ibrahim MM, Turei D, Saez-Rodriguez J. Benchmark and
integration of resources for the estimation of human transcription factor activities. Genome
Res. 2019;29:1363–75.

56. Liu Z-P, Wu C, Miao H, Wu H. RegNetwork: an integrated database of transcriptional and
post-transcriptional regulatory networks in human and mouse. Database. 2015;2015:bav095.

57. Baik SH, Kang S, Son SM, Mook-Jung I. Microglia contributes to plaque growth by cell death
due to uptake of amyloid β in the brain of Alzheimer's disease mouse model. Glia.
2016;64:2274–90.

58. Kheiri G, Dolatshahi M, Rahmani F, Rezaei N. Role of p38/MAPKs in Alzheimer's disease:
implications for amyloid beta toxicity targeted therapy. Rev Neurosci. 2018;30:9–30.

59. Marín-Teva JL, Cuadros MA, Martín-Oliva D, Navascués J. Microglia and neuronal cell death.
Neuron Glia Biol. 2011;7:25–40.

60. Barbier P, Zejneli O, Martinho M, Lasorsa A, Belle V, Smet-Nocca C, et al. Role of Tau as a
Microtubule-Associated Protein: Structural and Functional Aspects. Front Aging Neurosci.
2019;11:204.

61. McCutcheon RA, Krystal JH, Howes OD. Dopamine and glutamate in schizophrenia: biology,
symptoms and treatment. World Psychiatry Off J World Psychiatr Assoc WPA. 2020;19:15–33.

62. Berdenis van Berlekom A, Muflihah CH, Snijders GJLJ, MacGillavry HD, Middeldorp J, Hol
EM, et al. Synapse Pathology in Schizophrenia: A Meta-analysis of Postsynaptic Elements in
Postmortem Brain Studies. Schizophr Bull. 2020;46:374–86.

63. Osimo EF, Beck K, Reis Marques T, Howes OD. Synaptic loss in schizophrenia: a meta-
analysis and systematic review of synaptic protein and mRNA measures. Mol Psychiatry.
2019;24:549–61.

64. McGuire JL, Depasquale EA, Funk AJ, O'Donnovan SM, Hasselfeld K, Marwaha S, et al.
Abnormalities of signal transduction networks in chronic schizophrenia. NPJ Schizophr.
2017;3:30.

65. Kehrer C, Maziashvili N, Dugladze T, Gloveli T. Altered Excitatory-Inhibitory Balance in the
NMDA-Hypofunction Model of Schizophrenia. Front Mol Neurosci. 2008;1:6.

66. Ahmad R, Sportelli V, Ziller M, Spengler D, Hoffmann A. Tracing Early Neurodevelopment in
Schizophrenia with Induced Pluripotent Stem Cells. Cells. 2018;7.

67. Sasi M, Vignoli B, Canossa M, Blum R. Neurobiology of local and intercellular BDNF signaling.
Pflüg Arch - Eur J Physiol. 2017;469:593–610.

68. Fabiani C, Antollini SS. Alzheimer's Disease as a Membrane Disorder: Spatial Cross-Talk
Among Beta-Amyloid Peptides, Nicotinic Acetylcholine Receptors and Lipid Rafts. Front Cell
Neurosci. 2019;13:309.

69. O'Brien RJ, Wong PC. Amyloid precursor protein processing and Alzheimer's disease. Annu
Rev Neurosci. 2011;34:185–204.

70. Mahoney ER, Dumitrescu L, Moore AM, Cambronero FE, De Jager PL, Koran MEI, et al. Brain
expression of the vascular endothelial growth factor gene family in cognitive aging and
alzheimer's disease. Mol Psychiatry. 2019;

71. Misiak B, Stramecki F, Stańczykiewicz B, Frydecka D, Lubeiro A. Vascular endothelial growth
factor in patients with schizophrenia: A systematic review and meta-analysis. Prog
Neuropsychopharmacol Biol Psychiatry. 2018;86:24–9.

72. Glausier JR, Lewis DA. Dendritic spine pathology in schizophrenia. Neuroscience.
2013;251:90–107.

73. Cochran JN, Hall AM, Roberson ED. The dendritic hypothesis for Alzheimer's disease
pathophysiology. Brain Res Bull. 2014;103:18–28.

74. Fakhoury M. Microglia and Astrocytes in Alzheimer's Disease: Implications for Therapy. Curr
Neuropharmacol. 2018;16:508–18.

75. Raabe FJ, Slapakova L, Rossner MJ, Cantuti-Castelvetri L, Simons M, Falkai PG, et al.
Oligodendrocytes as A New Therapeutic Target in Schizophrenia: From Histopathological
Findings to Neuron-Oligodendrocyte Interaction. Cells. 2019;8.

76. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. Nat Genet. 2013;45:580–
5.

77. Kim-Hellmuth S, Aguet F, Oliva M, Muñoz-Aguirre M, Kasela S, Wucher V, et al. Cell type–
specific genetic regulation of gene expression across human tissues. Science.
2020;369:eaaz8528.

78. Watanabe K, Taskesen E, van Bochoven A, Posthuma D. Functional mapping and annotation
of genetic associations with FUMA. Nat Commun. 2017;8:1826.

79. Nguyen ND, Wang D. Multiview learning for understanding functional multiomics. PLoS
Comput Biol. United States; 2020;16:e1007677.

80. Lacal I, Ventura R. Epigenetic Inheritance: Concepts, Mechanisms and Perspectives. Front
Mol Neurosci. 2018;11:292.

81. Linker SM, Urban L, Clark SJ, Chhatriwala M, Amatya S, McCarthy DJ, et al. Combined single-
cell profiling of expression and DNA methylation reveals splicing regulation and heterogeneity.
Genome Biol. 2019;20:30.

82. Marx V. A dream of single-cell proteomics. Nat Methods. 2019;16:809–12.

83. Brummelman J, Haftmann C, Núñez NG, Alvisi G, Mazza EMC, Becher B, et al. Development,
application and computational analysis of high-dimensional fluorescent antibody panels for
single-cell flow cytometry. Nat Protoc. 2019;14:1946–69.