1

A Platform Incorporating Trimeric Antigens into Self-Assembling Nanoparticles Reveals SARS-CoV-2-Spike Nanoparticles to Elicit Substantially Higher Neutralizing Responses than Spike Alone

Baoshan Zhang[1*], Cara W. Chao[1*], Yaroslav Tsybovsky[2], Olubukola M. Abiona[1], Geoffrey B. Hutchinson[1], Juan I. Moliva[1], Adam S. Olia[1], Amarendra Pegu[1], Emily Phung[1,3], Guillaume Stewart-Jones[1], Raffaello Verardi[1], Lingshu Wang[1], Shuishu Wang[1], Anne Werner[1], Eun Sung Yang[1], Christina Yap[1], Tongqing Zhou[1], John R. Mascola[1], Nancy J. Sullivan[1], Barney S. Graham[1], Kizzmekia S. Corbett[1] and Peter D. Kwong[1¶]

[1] Vaccine Research Center, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland, USA.

[2] Electron Microscopy Laboratory, Cancer Research Technology Program, Leidos Biomedical Research Inc., Frederick National Laboratory for Cancer Research, Frederick, Maryland, USA.

[3] Institute for Biomedical Sciences, George Washington University, Washington, DC, USA.

[*] These authors contributed equally to this work.

[¶] Corresponding author (PDK) Email: pdkwong@nih.gov

Abstract

Antigens displayed on self-assembling nanoparticles can stimulate strong immune responses and have been playing an increasingly prominent role in structure-based vaccines. However, the development of such immunogens is often complicated by inefficiencies in their production. To alleviate this issue, we developed a plug-and-play platform using the spontaneous isopeptide-bond formation of the SpyTag:SpyCatcher system to display trimeric antigens on self-assembling nanoparticles, including the 60-subunit *Aquifex aeolicus* lumazine synthase (LuS) and the 24-subunit *Helicobacter pylori* ferritin. LuS and ferritin coupled to SpyTag expressed well in a mammalian expression system when an *N*-linked glycan was added to the nanoparticle surface. The respiratory syncytial virus fusion (F) glycoprotein trimer – stabilized in the prefusion conformation and fused with SpyCatcher – could be efficiently conjugated to LuS-SpyTag or ferritin-SpyTag, enabling multivalent display of F trimers with prefusion antigenicity. Similarly, F-glycoprotein trimers from human parainfluenza virus-type 3 and spike-glycoprotein trimers from SARS-CoV-2 could be displayed on LuS nanoparticles with decent yield and antigenicity. Notably, murine vaccination with the SARS-CoV-2 spike-LuS nanoparticles elicited ~25-fold higher neutralizing responses, weight-per-weight relative to spike alone. The versatile platform described here thus allows for multivalent plug-and-play presentation on self-assembling nanoparticles of trimeric viral antigens, with SARS-CoV-2 spike-LuS nanoparticles inducing particularly potent neutralizing responses.

Introduction

Self-assembling nanoparticles are playing an increasingly prevalent role in vaccine development as vaccine vehicles and immunomodulators. The appeal of nanoparticle immunogens lies in their inherent multivalent display of antigens, which is known to elicit robust B cell responses (reviewed in [1]). There have been numerous efforts to genetically fuse viral immunogens to nanoparticles within prokaryotic and eukaryotic systems, utilizing direct genetic fusion of antigenic molecules with self-assembling nanoparticle monomers[2-4], chemical conjugation[5,6], and a spontaneous intramolecular isopeptide bond formation with the SpyTag:SpyCatcher system[7,8], and some of these nanoparticles are now entering clinical trials[2,9,10].

Another important factor to consider in viral immunogen design is glycosylation. Viral pathogens are often heavily glycosylated, often as a means to evade the human immune system. Moreover, many viral antigens require glycosylation to be stably expressed and correctly folded. Although several studies have described plug-and-play nanoparticle systems[11-16], many use prokaryotic expression systems, which are not suitable to produce correctly glycosylated antigens. Furthermore, *N*-glycans can be manipulated in immunogen design to selectively occlude unwanted epitopes as well as to improve the solubility and stability of immunogens[17-19]. Another factor to consider is that metastable type 1 fusion machines are prevalent vaccine targets[20].

Here we developed a modular self-assembling nanoparticle platform that allows for the plug-and-play display of trimeric viral glycoproteins on nanoparticle surfaces, utilizing the SpyTag:SpyCatcher system. We assessed this system with three prefusion (preF)-stabilized viral trimeric glycoproteins: respiratory syncytial virus fusion (RSV F) glycoprotein[21], human parainfluenza virus type 3 fusion glycoprotein (PIV3 F)[22], and SARS-CoV-2 spike glycoprotein[23,24]. SpyTag-coupled nanoparticles could be expressed with sufficient yield of soluble proteins from a mammalian expression system after the addition of nanoparticle surface glycans.

The nanoparticle-formatted trimers exhibited improved antigenicity versus soluble trimers for apical epitopes, and we explicitly tested the immunogenicity for the nanoparticle-formatted trimeric antigen from SARS-CoV-2 in mice. Overall, protein antigens and nanoparticle scaffolds could be produced independently before conjugation, thereby expediting the otherwise generally cumbersome process of making and troubleshooting immunogens genetically fused to self-assembling nanoparticle subunits. Such a modular nanoparticle assembly platform may thus be a useful tool for plug-and-play screening of trimeric viral immunogens in a multivalent highly immunogenic context, and we provide proof-of-principle for increased immunogenicity of a nanoparticle-displayed SARS-CoV-2 spike.

Results

Expression of LuS- and ferritin-nanoparticle scaffolds with SpyTag requires the addition of an *N*-linked glycan

To construct a reliable plug-and-play platform for nanoparticle presentation of antigens, we chose *Aquifex aeolicus* lumazine synthase (LuS)[25] and *Helicobacter pylori* ferritin[26] as nanoparticle scaffolds with SpyTag:SpyCatcher conjugation system[15] to display antigens on nanoparticle surface. The SpyTag:SpyCatcher system is highly specific and stable with an isopeptide bond and has been used for conjugation of antigens on nanoparticle surfaces[7,27] (Fig. 1a). LuS and ferritin have served as scaffolds for nanoparticle immunogens in several clinical studies: for LuS see https://www.clinicaltrials.gov/ct2/show/NCT03699241[28]; for ferritin, see https://www.clinicaltrials.gov/ct2/show/NCT03547245[10,29,30]. The N terminus of both ferritin and LuS are exposed to the nanoparticle surface and are thus accessible for SpyTag or SpyCatcher attachment (Fig. 1b). The C terminus of LuS is also accessible on the nanoparticle surface and can be used to display purification tags. We designed mammalian expression constructs expressing

90   fusion proteins of SpyTag or SpyCatcher with LuS or ferritin. The constructs included both His-

91   and Strep-tags for purification purposes, along with a signal peptide for secretion of the expressed

92   proteins into supernatant medium (Fig. 1b).

93   Initial constructs yielded low levels of soluble proteins for the nanoparticle-SpyTag or

SpyCatcher fusion proteins. To improve protein solubility and expression, we added glycans to the

surface of the nanoparticles, designing a panel of LuS and ferritin constructs with SpyTag and

SpyCatcher (Table 1 and Supplementary Table S1). For LuS constructs, we added a glycosylation

site at position 71 (PDB 1HQK numbering). For ferritin constructs, two potential glycosylation

sites (96 and 148) were tested. The addition of N-linked glycosylation sites facilitated expression

of soluble nanoparticles in the cell culture supernatant. Three of the constructs produced

appreciable yields of well-assembled nanoparticles, LuS with N71 and SpyTag at N-terminus

(hereafter referred to as LuS-N71-SpyTag), ferritin with N96 and SpyTag, and ferritin S148

(glycan at N146) and SpyTag (Table 1). Of the two ferritin constructs, the ferritin with N96 and

SpyTag had a higher yield and was chosen for further studies (hereafter referred to as ferritin-N96-

SpyTag). Size exclusion chromatography (SEC) and electron microscopy (EM) analyses indicated

that LuS-N71-SpyTag formed a homogeneous nanoparticle population in solution (Fig. 1c,d). The

ferritin-N96-SpyTag sample comprised mainly intact nanoparticles with some minor unassembled

species (Fig. 1c,d). Negative-stain electron microscopy (EM) images indicated both nanoparticles

to be well-assembled with expected sizes[25,26] (Fig. 1d). Two-dimensional class average revealed

more detailed structural features of the nanoparticles, which were consistent with previously

published structures of the two nanoparticles. These data indicated the ferritin and LuS

nanoparticles were compatible with the SpyTag and glycosylation site addition. These alterations

were well tolerated, allowing for robust nanoparticle assembly. To verify the glycosylation of

LuS- and ferritin-SpyTag nanoparticles, we performed PNGase F digestion and checked for glycan

cleavage through SDS-PAGE (Fig. 1e). Both nanoparticles showed a band shift in the presence of

PNGase F, indicating the presence of *N*-liked glycan on the nanoparticles and its removal by the

amidase digestion. While the glycan cleavage in LuS-N71-SpyTag is distinct, it is less apparent in

ferritin-N96-SpyTag, likely due to incomplete glycosylation of ferritin-N96-SpyTag and multiple

bands of ferritin on SDS-PAGE. Ferritin has been observed to exhibit a single band on SDS-PAGE

in some studies[2] but multiple bands in others[16,31], presumably due to protease cleavage at the C

terminus or incomplete glycosylation. However, these different sized ferritin molecules assembled

correctly as nanoparticles with expected dimensions as indicated by SEC and EM (Fig. 1c,e).

Conjugation of RSV F-SpyCatcher to LuS-N71-SpyTag displays prefusion RSV F 'DS2'-trimers homogeneously on the surface of the LuS-N71-Spylinked-RSV F nanoparticle

With no effective licensed vaccine against respiratory syncytial virus (RSV), the causative

agent for a prevalent childhood disease that results in approximately 60,000 hospitalizations and

10,000 annual deaths in the US, developing an immunogen capable of eliciting protection is of

paramount importance[32]. Previous efforts in developing a vaccine capable of eliciting protective

antibodies have resulted in the identification of RSV trimers stabilized in its prefusion

conformation, RSV F DS-Cav1 (DS-Cav1), and RSV F DS2 (DS2)[21,33]. DS2 was shown to elicit

higher RSV neutralization responses than DS-Cav1[33]. With this as our motivation and as a test

case for our nanoparticle SpyTag:SpyCatcher system, we investigated the feasibility of displaying

DS2 in the context of nanoparticle immunogens.

We prepared DS2 coupled to SpyCatcher (hereafter referred to as RSV F-SpyCatcher) by

genetic engineering to append SpyCatcher to the C-terminus of RSV F after a 3 residue (GSG)

linker (Supplementary Table S1). After expression and purification, we conjugated the purified

RSV F-SpyCatcher to the purified 60-mer LuS-N71-SpyTag nanoparticle (Fig. 2a). SEC profiles

of the two components revealed that LuS-N71-SpyTag eluted around 13 mL and RSV F-

SpyCatcher eluted near 15 mL on Superdex 200 Increase 10/300 column (GE Health Sciences)

(Fig. 2b). The conjugated LuS-N71-SpyLinked-RSV F-SpyCatcher nanoparticle (LuS-N71-

SpyLinked-RSV F) eluted in a new peak at ~10 mL by SEC (Fig. 2b). SDS-PAGE showed the

appearance of species of larger molecular weight of ~90 kDa in the conjugation mixture, followed

by bands of residual LuS-N71-SpyTag monomer and RSV F-SpyCatcher components at 20 kDa

and 60 kDa, respectively (Fig. 2c), confirming the success of the conjugation reaction. To estimate

the conjugation efficiency, we measured the intensity of each band on the SDS-PAGE gel image

of the conjugated nanoparticle product (Fig. 2c), as a surrogate of mass for each component.

Taking into consideration the molecular weight of each component, we calculated the molar ratio

of each component to total protein in the sample. We estimated 67% of all the LuS nanoparticle

subunit was conjugated to RSV F trimer. To verify particle integrity after conjugation, we

performed negative stain EM following SEC purification. LuS-N71 -SpyTag conjugated with RSV

F-SpyCatcher efficiently produced uniform particles with a core diameter of 18.2 ± 1.0 nm

decorated with trimer spike of 11.4 ± 0.8 nm in length (Fig. 2d). We then confirmed the prefusion

state of the LuS-N71-SpyLinked RSV F nanoparticle through surface plasmon resonance using

RSV prefusion F specific antibodies D25 (site Ø) and MPE8 (site III) (Fig. 2e)[21]. Notably, RSV F

on nanoparticles showed an enhanced on-rate to the apex-targeting D25 antibody and reduced on-

rate to the equatorial targeting MPE8 versus trimeric RSV F, a crucial antigenic characteristic

signifying appropriate nanoparticle display[3].

Conjugation of RSV prefusion F-SpyCatcher to ferritin-SpyTag produces uniform ferritin-

RSV F nanoparticles

Having produced successfully the LuS-N71-SpyLinked-RSV F nanoparticle, we next set

out to conjugate the 24-mer ferritin-N96-SpyTag with RSV F-SpyCatcher in the same manner

(Fig. 3a). SEC of ferritin-N96-SpyTag nanoparticle showed a peak at around 16-17 mL, slightly

slower than RSV F-SpyCatcher (Fig. 3b). Negative stain EM revealed that ferritin-N96-SpyTag

formed nanoparticle of the expected size (Fig. 1c). The conjugation mixture of ferritin-N96-

SpyTag with RSV F-SpyCatcher exhibited a peak at ~10 mL (void volume of the SEC column),

suggesting successful formation of the conjugation product (referred to as ferritin-RSV F) (Fig.

3b). SDS-PAGE demonstrated the appearance of a new band at ~90 kDa, the expected size of

ferritin-N96-SpyLinked-RSV F nanoparticle, with residual ferritin-N96-SpyTag at around 20 kDa

(Fig. 3c). Using the same method as for LuS-N71-SpyLinked-RSV F above, we estimated 85% of

all the ferritin nanoparticle subunit was conjugated to RSV F trimer. To confirm the formation of

ferritin-N96-SpyLinked-RSV F nanoparticle, we performed negative stain EM, which showed

well-formed nanoparticles with the expected size and shape, displaying trimer spikes around the

ferritin nanoparticle (Fig. 3d).

To verify the conserved prefusion state of the conjugated RSV F trimer, we measured the

binding of ferritin-N96-SpyLinked-RSV F to D25 and MPE8 IgGs through SPR (Fig. 3e).

Importantly, we observed the on-rate to increase for D25, which recognizes an epitope at the

trimer apex, but the on-rate to decrease for MPE8, which recognizes an equatorial epitope on the

trimer, similar to the observation for LuS-N71-SpyLinked-RSV F.

Displaying PIV3 F glycoprotein trimer on LuS nanoparticle via SpyTag:SpyCatcher

conjugation improves antibody binding to the trimer apex

To demonstrate the plug-and-play versatility of the SpyTag:SpyCatcher nanoparticle

system, we produced PIV3 F glycoprotein trimer[34] as a fusion protein with SpyCatcher at the C

terminus and conjugated with the LuS-N71-SpyTag nanoparticle (Fig. 4, a-c, Supplementary Table

S1), similar to that described above for the conjugation of RSV F-SpyCatcher. PIV3 is a prevalent

human parainfluenza virus that causes respiratory illnesses, especially in infants and young

children[35,36]. The conjugation mixture of PIV3 F-SpyCatcher and LuS-N71-SpyTag was loaded

onto the SEC column to purify the conjugated nanoparticle product LuS-N71-SpyLinked-PIV3 F

from unconjugated nanoparticles and the PIV3 F SpyCatcher trimer (Fig. 4b). SDS-PAGE analysis

revealed that the conjugated product had the expected molecular weight and there was not

unconjugated PIV3 F-SpyCatcher in the conjugation mixture (Fig. 4c). Using the same method as

for LuS-N71-SpyLinked-RSV F in the previous section, we estimated 68% of all the LuS

nanoparticle subunit was conjugated to PIV3 F trimer. The PIV3 F conjugated nanoparticle was

further verified through negative stain EM, which showed well defined trimer spikes decorating

the LuS nanoparticle with the expected size (Fig. 4d).

Having produced nanoparticles of LuS-N71-SpyTag conjugated with PIV3 F-SpyCatcher,

we next evaluated binding of LuS-N71-SpyLinked-PIV3 F with antibodies PIA174 and PIA75[34],

using SPR (Fig. 4e). The head-targeting antibody PIA174 showed an improved binding to LuS-

N71-SpyLinked-PIV3 F relative to its binding to PIV3 F-SpyCatcher. The stem-targeting antibody

PIA75, however, showed decreased binding to LuS-N71-SpyLinked-PIV3 F compared with PIV3

F-SpyCatcher.

Conjugation of SARS-CoV-2 spike trimer to LuS nanoparticle via SpyTag:SpyCatcher

displays the spike trimers homogeneously on the nanoparticle surface

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) caused the COVID-19

pandemic that is ongoing worldwide[37]. An effective vaccine against SARS-CoV-2 and related

coronaviruses is urgently needed. The SARS-CoV-2 spike glycoprotein trimer mediates virus-cell

membrane fusion and thus a target for vaccine development[23,24]. To test the versatility of our plug-

and-play SpyTag:SpyCatcher nanoparticle system, we expressed and purified SARS-CoV-2 spike

fused with a C-terminal SpyCatcher and conjugated to the LuS-N71-SpyTag nanoparticle (Fig. 5,

a-c, Supplementary Table S1). For this construct, we used the prefusion stabilized version of spike

developed by McLellan and colleagues[24], which included GSAS and PP mutations and the T4

phage fibritin trimerization domain along with a single-chain Fc tag as described by Zhou and

colleagues[38].

The conjugation mixture was loaded onto an SEC column to purify the conjugated

nanoparticle product LuS-N71-SpyLinked-CoV spike from unconjugated LuS-N71-SpyTag and

SARS-CoV-2 spike-SpyCatcher (Fig. 5b). SDS-PAGE analysis revealed the conjugated product to

have the expected molecular weight, and unconjugated spike-SpyCatcher was not observed after

conjugation (Fig. 5c). Using the same method as for LuS-N71-SpyLinked-RSV F, we estimated

91% of all the LuS nanoparticle subunit was conjugated to the spike trimer. Negative stain EM

showed LuS-N71-SpyLinked-CoV-2 spike nanoparticle to exhibit the expected size with spike

trimers displaying on the LuS nanoparticle surface (Fig. 5d). SPR measurements showed LuS-

N71-SpyLinked-SARS-CoV-2 Spike to bind to CR3022[39,40], an antibody targeting the receptor-

binding domain (RBD), indicating successful nanoparticle presentation of the spike trimer using

the LuS-SpyTag:SpyCatcher system.

LuS-N71-SpyLinked-nanoparticle display increases potential of SARS-CoV-2 spike to elicit

neutralizing antibodies

To assess immunogenicity, we injected mice with the LuS-N71-SpyLinked-CoV-2 spike

nanoparticle or spike trimers (stabilized by 2P mutation)[24,41], or mock (LuS-N71-SpyTag)

nanoparticles at weeks 0 and 3 (Fig. 6a). Serum samples were collected two weeks after each

immunization. After the first immunization, at the lowest immunogen dose of 0.08 µg, spike

nanoparticle-immune sera exhibited an anti-SARS-CoV-2 spike ELISA geometric mean titer of

5,116, whereas only 1 out of 10 trimeric spike-immunized sera exhibited a measurable titer (Fig.

6b); after a second immunization, titers for the spike nanoparticle-immune sera increased

substantially, by approximately 25-fold. Immunizations with higher doses of spike nanoparticle

(0.4 and 2.0 µg) increased titers more incrementally, both at week 2 and at week 5. By contrast,

increases in dose of the spike trimer raised ELISA titers more substantially, with two of the mice

in the 2.0 µg spike-trimer immune sera reaching the assay upper limit of detection with a titer of

1,638,400 (Fig. 6b).

Importantly, pseudovirus neutralization assays revealed the LuS-N71-SpyLinked-CoV-2

spike nanoparticle to elicit potent neutralization responses with geometric mean ID_{50} titers of 413,

1820, and 1501 for immunization doses of 0.08, 0.4, and 2 µg, respectively (Fig. 6c). In

comparison, two doses of trimeric spike elicited neutralization titers at the 0.4 and 2 µg doses with

a geometric mean ID_{50} of 49 and 315, respectively, with no measurable neutralization at the 0.08

µg dose. In essence, 0.08 µg of spike nanoparticle elicited a neutralization response that was

higher, though statistically indistinguishable from 2 µg of trimeric spike. This indicated ~25-fold

higher immunogenicity on a weight-by-weight basis for the spike nanoparticle versus spike alone,

suggesting a substantial "dose-sparing" effect. Overall, presentation of the SARS-CoV-2 spike on

the LuS nanoparticle surface significantly improved its immunogenicity and required a lower

immunogen dose to elicit potent neutralization responses compared with the trimeric form.

Discussion

Nanoparticle-based immunogens can induce potent neutralizing antibodies[2,3,42] and thus

may be promising vaccine candidates. To develop nanoparticle vaccine immunogens, rapid and

efficient methods would help produce nanoparticle scaffolds that can be mixed and matched with

different immunogens. Previous efforts utilizing the spontaneous isopeptide bond formation with

the SpyTag:SpyCatcher system for nanoparticle surface display of immunogens[11-16] have proven

the versatility of this system for antigen display. However, none of these previously published

reports utilized mammalian expression allowing for post-translational modifications, such as *N*-

linked glycosylation. Here, we describe two nanoparticle platforms, lumazine synthase and ferritin,

for the display of trimeric viral protein immunogens using the SpyTag:SpyCatcher system. By

adding *N*-linked glycosylation sites to nanoparticle monomers, we were able to produce SpyTag-

coupled nanoparticles using mammalian cell culture.

266   LuS and ferritin nanoparticle platforms vary in the number of molecules displayed on the

surface. LuS-N71-SpyTag contains 60 SpyTags whereas ferritin-N96-SpyTag has 24 displayed on

surface, available for SpyCatcher-carrying molecules to couple to. Both platforms showed efficient

conjugation of trimeric immunogens and formed nanoparticle rapidly under physiological

conditions for RSV F, PIV3 F and SARS-CoV-2 spike trimers. One advantage of the LuS

nanoparticle is the high efficiency of its particle assembly. The glycosylated LuS-N71-SpyTag

assembled into a homogenous particle that exhibited a single peak in size exclusion

chromatography.

274   To demonstrate the versatility of our SpyTag-displaying nanoparticles in immunogen

development, we conjugated them to three viral antigens of vaccine interest, the DS2-preF

stabilized RSV F[33], a DS2-stabilized version of PIV3 F[34], and the 2P-stabilized version of SARS-

CoV-2 spike[24]. In each of these, we appended the SpyCatcher after the 'foldon' heterologous

trimeric stabilization motif[43]. Conjugation of SpyTag-nanoparticles with SpyCatcher-coupled RSV

F, PIV3 F and SARS-CoV-2 spike trimers resulted in proper particle assembly. In all three cases,

we observed high conjugation efficiency.

Our antigenicity analyses indicate that presentation of trimeric antigens from viral pathogens on self-assembling nanoparticles needs to take into consideration the accessibility of the antigenic epitopes. When a trimer protein is conjugated to a nanoparticle, such as LuS or ferritin in this study, the trimer molecules are densely displayed on the nanoparticle surface. As a result of the dense display, the epitopes near the nanoparticle surface, such as those at the stem region of the trimers in this study, are less accessible to antibodies than the epitopes on the apex of trimer molecules. Consequently, we observed an increased level of antibody binding to epitopes on the apex and a decreased level of antibody binding to epitopes on the equatorial or stem region of RSV F and PIV3 F trimer molecules (Figs. 2e, 3e and 4e).

The increased antigenicity of the apical epitopes on the trimer conjugated to nanoparticles is expected to yield increased immunogenicity – especially at lower dose, and we provide proof-of-principle for this with murine immunization studies with LuS-N71-SpyLinked-CoV-2 spike as compared to soluble spike. As observed in prior studies[2,3], nanoparticle immunogens elicited stronger immune responses than the corresponding trimers at low immunogen doses: at the 0.08 µg dose after two immunizations, spike nanoparticle elicited neutralization response with ID_{50} of 413, whereas trimeric spike elicited an equivalent neutralization titer only at the 25-fold higher dose of 2 µg. At 0.4 µg, spike nanoparticle elicited ~37-fold higher ID_{50} than trimeric spike. However, at a high dose of 2 µg, spike nanoparticle-elicited neutralization response appeared to plateau – at a level ~5-fold higher in neutralization titer than the trimeric immunogen. Similar increases in immunogenicity and with dose-sparing have been recently reported for nanoparticles incorporating the receptor-binding domain (RBD) of the spike[44]. Overall, multivalent presentation of trimeric antigens on nanoparticle can significantly improve their immunogenicity, allowing for elicitation of potent immune responses at a relatively low immunogen dose. Our

SpyTag:SpyCatcher system provides a versatile platform for preparation of such nanoparticle

immunogens from trimeric antigens.

It will be interesting to see if the plug-and-display technology described here will allow

for the incorporation of different molecules on multiple nanoparticles. Such molecules could

include not only trimeric viral immunogens, but immunostimulatory components, or molecules

targeting antigen presenting cells. Thus, the LuS- and ferritin-SpyTag displaying nanoparticles

described here may be amendable to mix-and-match display of immunogens and of

immunostimulatory or targeting components.

Materials and Methods

Protein production and purification

The amino acid sequences of protein expression constructs are listed in Supplementary Table S1.

For protein expression, 3 ml of Turbo293 transfection reagent (Speed BioSystems) was mixed

with 50 ml Opti-MEM medium (Life Technology) and incubated at room temperature (RT) for 5

min. 1 mg plasmid DNAs was mixed with 50 ml of Opti-MEM medium in a separate tube, and the

mixture added to the Turbo293 Opti-MEM mixture. The transfection mixture was incubated for 15

min at RT then added to 800 ml of Expi293 cells (Life Technology) at 2.5 million cells/ml. The

transfected cells were incubated overnight in a shaker incubator at 9% CO_2, 37 °C, and 120 rpm.

On the second day, about 100 ml of Expi293 expression medium was added. On day 5 post

transfection, supernatants were harvested, filtered. Proteins were purified from the supernatant

using Ni-NTA and strep chromatography. SARS-CoV-2 spike-SpyCatcher was expressed as a

fusion protein with a single-chain Fc purification tag and purified using Protein A

chromatography. SARS-CoV-2 spike-SpyCatcher protein was cleaved off from Protein A column

by HRV3C protease. All proteins were further purified by size exclusion chromatography on

Superdex 200 Increase 10/300 GL in PBS.

LuS- and ferritin- SpyTag conjugations

A 1:1 molar ratio of LuS- or ferritin-SpyTag and immunogen-SpyCatcher components were

combined and incubated at ambient temperature for approximately 3 hours, followed by size

exclusion column on Superdex200 Increase 10/300 GL in PBS to separate conjugated products

from residual components. The conjugated nanoparticle product was then run through SDS-PAGE

to verify conjugation and analyzed by negative-stain EM.

Negative-stain electron microscopy (EM)

Samples were diluted to 0.02-0.05 mg/ml with a buffer containing 10 mM HEPES, pH 7, and 150

mM NaCl. A 4.7-µl drop of the diluted sample was applied to a glow-discharged carbon-coated

copper grid for approximately 15 s. The drop was removed using blotting paper, and the grid was

washed three times with 4.7-µl drops of the same buffer. Adsorbed proteins were negatively

stained by applying consecutively three 4.7-µl drops of 0.75% uranyl formate and removing each

drop with filer paper. Micrographs were collected using SerialEM[45] on an FEI Tecnai T20 electron

microscope operated at 200 kV and equipped with an Eagle CCD camera or using EPU on a

ThermoFisher Talos F200C electron microscope operated at 200 kV and equipped with a Ceta

CCD camera. The pixel size was 0.44 and 0.25 nm for Tecnai T20 and Talos F200C, respectively.

Particles were picked automatically using in-house written software (Y.T., unpublished).

Reference-free 2D classification was performed with Relion 1.4[46] and SPIDER[47]. The dimensions

of VLP cores and spikes were measured with e2display.py program from EMAN2.1 software

package[48] using a representative micrograph (LuS-N71-SpyLinked–CoV-2 S) or 2D class average

images (all other VLPs). For CoV-2 nanoparticles we observed increased structural content at low

pH, whereas the other micrographs were collected at physiological pH.

Surface plasmon resonance (SPR)

To prepare the chips (GE Healthcare Life Sciences CM5 chips), antibody IgG were immobilized

onto the chip by amine coupling to ~100-1000 response units (RU) depending the level of binding

of each trimer and nanoparticle pair to antibodies. To measure binding of SpyCatcher proteins and

and nanoparticles, a dilution series of SpyCatcher-linked proteins and nanoparticles were flowed

through the IgG sensor chip for 200 s followed by 800 s of dissociation at a flow rate of 30

µL/min. The starting trimer concentration for each sample was 200 nM. Sensor chip surfaces were

regenerated after each injection following manufacture instructions with Glycine 2.5 (GE

Healthcare Life Sciences 10 mM glycine -HCl at pH 2.5) at a flow rate of 40 µL/min for 180 s.

Mouse immunization

Mouse experiments were carried out in compliance with National Institutes of Health regulations

and approval from the Animal Care and Use Committee of the Vaccine Research Center. Six week

old female BALB/cJ mice (Jackson Laboratories) were inoculated intramuscularly with Sigma

Adjuvant System, at weeks 0 and 3, as detailed previously[49]. Serum was collected 2 weeks post-

prime and post-boost for measurements of antibody responses as detailed hereafter.

Enzyme-linked immunosorbent assay (ELISA)

ELISA experiments were carried out as previously described[41]. Briefly, Nunc Maxisorp ELISA

plates (ThermoFisher) were coated with 100 ng/well of stabilized soluble SARS-CoV-2 spike

protein[24] (with His-tag cleaved to remove potential cross-reactivity) in 1X PBS at 4 °C for 16 hr.

To eliminate fold-on-specific binding, 50 µg/mL of fold-on protein was added to serial dilutions of

heat-inactivated sera for 1 hr at room temperature (RT). After blocking in PBS-Tween (PBST)

supplemented with 5% nonfat milk, plates were incubated with sera for 1 hr at RT. After blocking

in PBS-Tween (PBST) supplemented with 5% nonfat milk, plates were incubated with serial

dilutions of heat-inactivated sera for 1 hr at RT. Secondary antibody, goat anti-mouse IgG

conjugated to horseradish peroxidase (ThermoFisher), was then added, followed by excitation with

3,5,3′5′-tetramethylbenzidine substrate (KPL). Each step in this procedure was followed by

standard washes in PBST. Endpoint titers were calculated as the dilution factor that resulted in an

optical density exceeding 4X background (secondary antibody alone).

Lentivirus-based pseudovirus neutralization assay

The pseudovirus neutralization assay was performed as described previously[41,50]. To produce

SARS-CoV-2 pseudovirus, a codon-optimized CMV/R-SARS-CoV-2 spike (Wuhan-1, Genbank

#: MN908947.3) plasmid, was constructed and co-transfected with plasmids encoding luciferase

reporter, human transmembrane protease serine 2 (TMPRSS2)[51], and lentivirus backbone into

HEK293T/17 cells (ATCC #CRL-11268), as previously described[52]. Heat-inactivated serum was

mixed with the pseudovirus, incubated at 37 °C, and then added to ACE-2-expressing 293T cells.

Cells were lysed after 72 hr, and luciferase activity was measured. Percent neutralization was

calculated with uninfected cells as 100% neutralization and cells infected with only pseudovirus as

0% neutralization. ID_{50} titers were determined using a log (agonist) vs. normalized response

(variable slope) nonlinear function in Prism v8 (GraphPad).

Data availability

All relevant data are within the paper and its Supporting Information files.

References

1   Lopez-Sagaseta, J., Malito, E., Rappuoli, R. & Bottomley, M. J. Self-assembling protein
nanoparticles in the design of vaccines. *Comput Struct Biotechnol J* **14**, 58-68,
doi:10.1016/j.csbj.2015.11.001 (2016).
2   Kanekiyo, M. *et al.* Self-assembling influenza nanoparticle vaccines elicit broadly neutralizing
H1N1 antibodies. *Nature* **499**, 102-106, doi:10.1038/nature12202 (2013).
3   Marcandalli, J. *et al.* Induction of Potent Neutralizing Antibody Responses by a Designed
Protein Nanoparticle Vaccine for Respiratory Syncytial Virus. *Cell* **176**, 1420-1431 e1417,
doi:10.1016/j.cell.2019.01.046 (2019).
4   Swanson, K. A. *et al.* A respiratory syncytial virus (RSV) F protein nanoparticle vaccine
focuses antibody responses to a conserved neutralization domain. *Sci Immunol* **5**,
doi:10.1126/sciimmunol.aba6466 (2020).
5   Cavelti-Weder, C. *et al.* Development of an Interleukin-1beta Vaccine in Patients with Type 2
Diabetes. *Mol Ther* **24**, 1003-1012, doi:10.1038/mt.2015.227 (2016).
6   Cornuz, J. *et al.* A vaccine against nicotine for smoking cessation: a randomized controlled
trial. *PLoS One* **3**, e2547, doi:10.1371/journal.pone.0002547 (2008).
7   Zakeri, B. *et al.* Peptide tag forming a rapid covalent bond to a protein, through engineering a
bacterial adhesin. *Proc Natl Acad Sci U S A* **109**, E690-697, doi:10.1073/pnas.1115485109
(2012).
8   Marini, A. *et al.* A Universal Plug-and-Display Vaccine Carrier Based on HBsAg VLP to
Maximize Effective Antibody Response. *Front Immunol* **10**, 2931,
doi:10.3389/fimmu.2019.02931 (2019).
9   He, L. *et al.* Presenting native-like trimeric HIV-1 antigens with self-assembling nanoparticles.
Nat Commun **7**, 12041, doi:10.1038/ncomms12041 (2016).
10 Yassine, H. M. *et al.* Hemagglutinin-stem nanoparticles generate heterosubtypic influenza
protection. *Nat Med* **21**, 1065-1070, doi:10.1038/nm.3927 (2015).
11 Escolano, A. *et al.* Immunization expands B cells specific to HIV-1 V3 glycan in mice and
macaques. *Nature* **570**, 468-473, doi:10.1038/s41586-019-1250-z (2019).
12 Govasli, M. L., Diaz, Y. & Puntervoll, P. Virus-like particle-display of the enterotoxigenic
Escherichia coli heat-stable toxoid STh-A14T elicits neutralizing antibodies in mice. *Vaccine*
37, 6405-6414, doi:10.1016/j.vaccine.2019.09.004 (2019).
13 Janitzek, C. M. *et al.* Bacterial superglue generates a full-length circumsporozoite protein
virus-like particle vaccine capable of inducing high and durable antibody responses. *Malar J*
15, 545, doi:10.1186/s12936-016-1574-1 (2016).
14 Thrane, S. *et al.* Bacterial superglue enables easy development of efficient virus-like particle
based vaccines. *J Nanobiotechnology* **14**, 30, doi:10.1186/s12951-016-0181-1 (2016).
15 Brune, K. D. *et al.* Plug-and-Display: decoration of Virus-Like Particles via isopeptide bonds
for modular immunization. *Sci Rep* **6**, 19234, doi:10.1038/srep19234 (2016).
16 Wang, W. *et al.* Ferritin nanoparticle-based SpyTag/SpyCatcher-enabled click vaccine for
tumor immunotherapy. *Nanomedicine* **16**, 69-78, doi:10.1016/j.nano.2018.11.009 (2019).
17 Duan, H. *et al.* Glycan Masking Focuses Immune Responses to the HIV-1 CD4-Binding Site
and Enhances Elicitation of VRC01-Class Precursor Antibodies. *Immunity* **49**, 301-311 e305,
doi:10.1016/j.immuni.2018.07.005 (2018).
18 Ingale, J. *et al.* Hyperglycosylated stable core immunogens designed to present the CD4
binding site are preferentially recognized by broadly neutralizing antibodies. *J Virol* **88**,
14002-14016, doi:10.1128/JVI.02614-14 (2014).

19 Pantophlet, R., Wilson, I. A. & Burton, D. R. Hyperglycosylated mutants of human
immunodeficiency virus (HIV) type 1 monomeric gp120 as novel antigens for HIV vaccine
design. *J Virol* **77**, 5889-5901 (2003).
20 Rey, F. A. & Lok, S. M. Common Features of Enveloped Viruses and Implications for
Immunogen Design for Next-Generation Vaccines. *Cell* **172**, 1319-1334,
doi:10.1016/j.cell.2018.02.054 (2018).
21 McLellan, J. S. *et al.* Structure-based design of a fusion glycoprotein vaccine for respiratory
syncytial virus. *Science* **342**, 592-598, doi:10.1126/science.1243283 (2013).
22 Stewart-Jones, G. B. *et al.* Trimeric HIV-1-Env Structures Define Glycan Shields from Clades
A, B, and G. *Cell* **165**, 813-826, doi:10.1016/j.cell.2016.04.010 (2016).
23 Walls, A. C. *et al.* Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
Glycoprotein. *Cell* **181**, 281-292 e286, doi:10.1016/j.cell.2020.02.058 (2020).
24 Wrapp, D. *et al.* Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
Science **367**, 1260-1263, doi:10.1126/science.abb2507 (2020).
25 Zhang, X., Meining, W., Fischer, M., Bacher, A. & Ladenstein, R. X-ray structure analysis
and crystallographic refinement of lumazine synthase from the hyperthermophile Aquifex
aeolicus at 1.6 A resolution: determinants of thermostability revealed from structural
comparisons. *J Mol Biol* **306**, 1099-1114, doi:10.1006/jmbi.2000.4435 (2001).
26 Cho, K. J. *et al.* The crystal structure of ferritin from Helicobacter pylori reveals unusual
conformational changes for iron uptake. *J Mol Biol* **390**, 83-98, doi:10.1016/j.jmb.2009.04.078
(2009).
27 Reddington, S. C. & Howarth, M. Secrets of a covalent interaction for biomaterials and
biotechnology: SpyTag and SpyCatcher. *Curr Opin Chem Biol* **29**, 94-99,
doi:10.1016/j.cbpa.2015.10.002 (2015).
28 Jardine, J. G. *et al.* HIV-1 broadly neutralizing antibody precursor B cells revealed by
germline-targeting immunogen. *Science* **351**, 1458-1463, doi:10.1126/science.aad9195 (2016).
29 Corbett, K. S. *et al.* Design of Nanoparticulate Group 2 Influenza Virus Hemagglutinin Stem
Antigens That Activate Unmutated Ancestor B Cell Receptors of Broadly Neutralizing
Antibody Lineages. *mBio* **10**, doi:10.1128/mBio.02810-18 (2019).
30 Darricarrere, N. *et al.* Development of a Pan-H1 Influenza Vaccine. *J Virol* **92**,
doi:10.1128/JVI.01349-18 (2018).
31 Morris, C. D. *et al.* Differential Antibody Responses to Conserved HIV-1 Neutralizing
Epitopes in the Context of Multivalent Scaffolds and Native-Like gp140 Trimers. *mBio* **8**,
doi:10.1128/mBio.00036-17 (2017).
32 Graham, B. S. Immunological goals for respiratory syncytial virus vaccine development. *Curr*
Opin Immunol **59**, 57-64, doi:10.1016/j.coi.2019.03.005 (2019).
33 Joyce, M. G. *et al.* Iterative structure-based improvement of a fusion-glycoprotein vaccine
against RSV. *Nat Struct Mol Biol* **23**, 811-820, doi:10.1038/nsmb.3267 (2016).
34 Stewart-Jones, G. B. E. *et al.* Structure-based design of a quadrivalent fusion glycoprotein
vaccine for human parainfluenza virus types 1-4. *Proc Natl Acad Sci U S A* **115**, 12265-12270,
doi:10.1073/pnas.1811980115 (2018).
35 Jain, S. *et al.* Community-Acquired Pneumonia Requiring Hospitalization among U.S. Adults.
N Engl J Med **373**, 415-427, doi:10.1056/NEJMoa1500245 (2015).
36 Jain, S. *et al.* Community-acquired pneumonia requiring hospitalization among U.S. children.
N Engl J Med **372**, 835-845, doi:10.1056/NEJMoa1405870 (2015).
37 Cucinotta, D. & Vanelli, M. WHO Declares COVID-19 a Pandemic. *Acta Biomed* **91**, 157-
160, doi:10.23750/abm.v91i1.9397 (2020).

38 Zhou, T. *et al.* Structure-Based Design with Tag-Based Purification and In-Process
Biotinylation Enable Streamlined Development of SARS-CoV-2 Spike Molecular Probes.
bioRxiv, 2020.06.22.166033, doi:10.1101/2020.06.22.166033 (2020).
39 ter Meulen, J. *et al.* Human monoclonal antibody combination against SARS coronavirus:
synergy and coverage of escape mutants. *PLoS Med* **3**, e237,
doi:10.1371/journal.pmed.0030237 (2006).
40 Yuan, M. *et al.* A highly conserved cryptic epitope in the receptor-binding domains of SARS-
CoV-2 and SARS-CoV. *Science*, doi:10.1126/science.abb7269 (2020).
41 Corbett, K. S. *et al.* SARS-CoV-2 mRNA vaccine design enabled by prototype pathogen
preparedness. *Nature*, doi:10.1038/s41586-020-2622-0 (2020).
42 Rts, S. C. T. P. Efficacy and safety of RTS,S/AS01 malaria vaccine with or without a booster
dose in infants and children in Africa: final results of a phase 3, individually randomised,
controlled trial. *Lancet* **386**, 31-45, doi:10.1016/S0140-6736(15)60721-8 (2015).
43 Efimov, V. P. *et al.* Fibritin encoded by bacteriophage T4 gene wac has a parallel triple-
stranded alpha-helical coiled-coil structure. *J Mol Biol* **242**, 470-486,
doi:10.1006/jmbi.1994.1595 (1994).
44 Walls, A. C. *et al.* Elicitation of potent neutralizing antibody responses by designed protein
nanoparticle vaccines for SARS-CoV-2. *bioRxiv*, 2020.08.11.247395,
doi:10.1101/2020.08.11.247395 (2020).
45 Mastronarde, D. N. Automated electron microscope tomography using robust prediction of
specimen movements. *J Struct Biol* **152**, 36-51, doi:10.1016/j.jsb.2005.07.007 (2005).
46 Scheres, S. H. RELION: implementation of a Bayesian approach to cryo-EM structure
determination. *J Struct Biol* **180**, 519-530, doi:10.1016/j.jsb.2012.09.006 (2012).
47 Frank, J. *et al.* SPIDER and WEB: processing and visualization of images in 3D electron
microscopy and related fields. *J Struct Biol* **116**, 190-199, doi:10.1006/jsbi.1996.0030 (1996).
48 Tang, G. *et al.* EMAN2: an extensible image processing suite for electron microscopy. *J Struct*
Biol **157**, 38-46, doi:10.1016/j.jsb.2006.05.009 (2007).
49 Pallesen, J. *et al.* Immunogenicity and structures of a rationally designed prefusion MERS-
CoV spike antigen. *Proc Natl Acad Sci U S A* **114**, E7348-E7357,
doi:10.1073/pnas.1707304114 (2017).
50 Corbett, K. S. *et al.* Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in
Nonhuman Primates. *New England Journal of Medicine*, doi:10.1056/NEJMoa2024671
(2020).
51 Bottcher, E. *et al.* Proteolytic activation of influenza viruses by serine proteases TMPRSS2
and HAT from human airway epithelium. *J Virol* **80**, 9896-9898, doi:10.1128/JVI.01118-06
(2006).
52 Wang, L. *et al.* Evaluation of candidate vaccine approaches for MERS-CoV. *Nature*
Communications **6**, 7712, doi:10.1038/ncomms8712 (2015).
530

Acknowledgements

We thank T. Beaumont and H. Spits for antibody D25, A. Lanzavecchia for antibodies PIA174,

PIA75 and MPE8, J. Stuckey for assistance with figures, and members of the Vaccine Research

Center for discussions or comments on the manuscript. We thank members of the NIH NIAID

VRC Translational Research Program for technical assistance with mouse experiments. Support

for this work was provided by the Intramural Research Program of the Vaccine Research Center,

National Institute of Allergy and Infectious Diseases, National Institutes of Health. This project

has been funded in part with Federal funds from Frederick National Laboratory for Cancer

Research, NIH, under Contract No. HHSN261200800001E (Y. Tsybovsky). K.S. Corbett is the

recipient of a research fellowship that was partially funded by the Undergraduate Scholarship

Program, Office of Intramural Training and Education, Office of the Director, NIH. The content of

this publication does not necessarily reflect the views or policies of the Department of Health and

Human Services, nor does mention of trade names, commercial products, or organizations imply

endorsement by the U.S. Government.

Author contributions

B.Z. and C.W.C. designed research with B.Z. heading protein design and production; B.Z.,

C.W.C., A.S.O., and R.V. produced nanoparticle and trimer proteins; C.W.C. and B.Z. prepared

trimer-coupled nanoparticles and performed antigenic assessments; Y.T. performed negative-stain

EM; S.W. assisted with manuscript assembly; T.Z. provided design for SARS-CoV-2 spike

protein; G.S-J. provided the design for PIV3 F protein; A.P., L.W. and E.S.Y. provided

pseudovirus; G.B.H. carried out immunizations; O.M.A. and A.W. performed ELISA; J.I.M.

performed neutralization assay; B.S.G. and K.S.C. designed mouse experiment and oversaw

ELISA experiment; J.R.M. oversaw pseudovirus preparation; N.J.S., B.S.G., and K.S.C. oversaw

neutralization assay; E.P. and C.Y. performed data analyses for immunoassays; P.D.K. oversaw

the project with B.Z., C.W.C., S.W., and P.D.K. writing the paper, and all other authors providing

revisions and comments.

Competing interests

K.S.C. and B.S.G. are inventors on International Patent Application No. WO/2018/081318 entitled

"Prefusion Coronavirus Spike Proteins and Their Use." K.S.C., O.M.A., G.B.H., and B.S.G. are

inventors on US Patent Application No. 62/972,886 entitled "2019-nCoV Vaccine".

Figure legends

Fig. 1. LuS- and ferritin-nanoparticle scaffolds with N-linked glycan and SpyTag express well as assembled nanoparticles in mammalian cells

(**a**) Schematic diagram showing the separate SpyTag and SpyCatcher to combine through an isopeptide bond as a means to covalently link molecules attached to SpyTag and molecules attached to SpyCatcher. (**b**) Design of expression constructs to produce activated nanoparticles with SpyTag in mammalian cells for conjugating antigens on the nanoparticle surface. Upper panel shows the DNA construct. A SpyTag was placed at the N-terminus of the nanoparticle sequence after the cleavable signal peptide. His and Strep tags were placed at the C-terminus of the LuS nanoparticle. An N-linked glycosylation site was engineered in the nanoparticle sequence to facilitate protein expression (see Table 1 and Supplementary Table S1 for more details). Lower panels show the expected structures of the LuS-N71-SpyTag and ferritin-N96-SpyTag monomers and assembled nanoparticles. Both glycan and SpyTag are expected to be on the nanoparticle surface. (**c**) Size exclusion chromatograms confirmed the correct sizes of the nanoparticles. The samples were loaded on a Superdex 200 Increase 10/300 GL column in PBS. Initial run of ferritin-96N-SpyTag nanoparticle revealed a tail of small molecular weight species; the chromatogram shown here is the re-run main peak. (**d**) SDS-PAGE of LuS-N71-SpyTag and ferritin-N96-SpyTag in the presence or absence of PNGase F. The position of PNGase F is marked. The multiple bands for ferritin are likely due to proteolytic cleavage and incomplete glycosylation (see text). (**e**) Negative stain EM images (left panels) and 2D class averages (right panels) of LuS-N71-SpyTag and ferritin-N96-SpyTag show the correct assembly of the purified nanoparticles with expected sizes.

Fig. 2. Conjugation of RSV F-SpyCatcher to LuS-SpyTag displays prefusion RSV F trimer

homogenously on the surface of the LuS-N71-SpyLinked-RSV F nanoparticle

(**a**) Schematic diagram showing conjugation of SpyTag-coupled LuS to SpyCatcher-coupled RSV

prefusion F trimer to make LuS-N71-SpyLinked-RSV F nanoparticle. (**b**) SEC profiles of LuS-

N71-SpyTag, RSV F-SpyCatcher, and the conjugated product LuS-N71-SpyLinked-RSV F on a

Superdex 200 Increase 10/300 GL column in PBS. (**c**) SDS-PAGE of LuS-N71-SpyTag (lane 1),

RSV F-SpyCatcher (lane 2), and the conjugated LuS-N71-SpyLinked-RSV F nanoparticle product

(lane 3), in the presence of DTT. (**d**) Negative stain EM images of the LuS-N71-SpyLinked-RSV

F nanoparticle after SEC purification, showing (left panel) a representative micrograph and (right

panel) the 2D class averages. (**e**) Surface plasmon resonance of RSV F-SpyCatcher and LuS-N71-

SpyLinked-RSV F nanoparticle with prefusion-specific D25 IgG (site Ø) and MPE8 IgG (site III),

with IgG coupled to chip and nanoparticle in solution. A concentration series from 200 nM to 1.56

nM of RSV F either as trimer (left) or couped to nanoparticle (right) was measured; k_a values are

provided as these have been found to correlate with immunogenicity[3].

Fig. 3. Conjugation of RSV prefusion F-SpyCatcher to ferritin-SpyTag produces uniform

ferritin-RSV F nanoparticles

(**a**) Schematic diagram showing the conjugation process of ferritin-N96-SpyTag and RSV F–

SpyCatcher to make ferritin-N96-SpyLinked-RSV F nanoparticle. (**b**) SEC profiles of ferritin-

N96-SpyTag, RSV F-SpyCatcher, and the conjugation reaction mixture on a Superdex 200

Increase 10/300 GL column in PBS. (**c**) SDS-PAGE of ferritin-N96-SpyTag (lane 1), RSV F-

SpyCatcher (lane 2), and the conjugated ferritin-N96-SpyLinked-RSV prefusion F SpyCatcher

nanoparticle product (lane 3), in the presence of DTT. Ferritin exhibited multiple bands due to

proteolytic cleavage and incomplete glycosylation (see text). (**d**) Negative stain EM images of the

ferritin-RSV F nanoparticle after SEC purification, showing (left panel) a representative

micrograph and (right panel) the 2D class averages. (**e**) SPR of RSV F-SpyCatcher and Ferritin-

N96-SpyLinked-RSV F nanoparticle with prefusion-specific D25 IgG (site Ø) and MPE8 IgG (site

III) using immobilized IgG on sensor chip with nanoparticle and trimer in solution. A

concentration series from 200 nM to 1.56 nM of RSV F either as trimer (left) or coupled to

nanoparticle (right) was measured; k_a values are provided.

Fig. 4. Conjugation of PIV3 F-SpyCatcher to LuS-SpyTag displays prefusion PIV3 F trimer

homogenously on the surface of the LuS-N71-SpyLinked-PIV3 F nanoparticle

(**a**) Schematic of the conjugation between LuS-N71-SpyTag and PIV3 F-SpyCatchert to produce

LuS-N71-SpyLinked-PIV3 F nanoparticle (**b**) SEC profiles of PIV3 F-SpyCatcher, LuS-SpyTag,

and the conjugated product LuS-N71-SpyLinked-PIV3 F on a Superdex 200 Increase 10/300 GL

in PBS. (**c**) SDS-PAGE of LuS-N71 (lane 1), PIV3 F-SpyCatcher (lane 2), and LuS-N71-

SpyLinked-PIV3 F conjugation mixture (lanes 3) in the presence of DTT. (**d**) Negative stain EM

of LuS-N71-SpyLinked-PIV3 following SEC showing a representative micrograph (left panel) and

2D class averages (right panel). (**e**) SPR measurements of PIV3 F-SpyCatcher and LuS-N71-

SpyLinked-PIV3 F were performed using IgG coupled chips with nanoparticle and timer in

solution. A concentration series from 200 nM to 1.56 nM of PIV3 F either as trimer (left) or

coupled to nanoparticle (right) was measured; k_a values are provided.

Fig. 5. Conjugation of SARS-CoV-2 spike trimer to LuS-SpyTag displays SARS-CoV-2 spike

trimer on the surface of the LuS-N71-SpyLinked-CoV-2 spike nanoparticle

(**a**) Schematic diagram showing conjugation of SpyTag-coupled LuS to SpyCatcher-coupled

SARS-CoV-2 spike trimer to make LuS-N71-SpyLinked-CoV-2 spike nanoparticle. (**b**) SEC

profiles of LuS-N71-SpyTag, SARS-CoV-2 spike-SpyCatcher, and the conjugated product LuS-

N71-SpyLinked-CoV-2 spike on a Superdex 200 Increase 10/300 GL column in PBS. (**c**) SDS-

PAGE of LuS-N71-SpyTag (lane 1), SARS-CoV-2 spike-SpyCatcher (lane 2), and the conjugation

mixture of LuS-N71-SpyTag with SARS-CoV-2 spike-SpyCatcher (lane 3) in the presence of

DTT. The conjugation mixture (lane 3) shows the conjugated LuS-N71-SpyLinked-CoV-2 spike

nanoparticle with minor excess of LuS-N71-SpyTag. (**d**) Negative stain EM of the LuS-N71-

SpyLinked-CoV-2 spike nanoparticle after SEC purification showing representative micrographs

(left panel) and 2D class average (right panel). (**e**) SPR response curves for LuS-N71-SpyLinked-

CoV-2 spike nanoparticle binding with RBD-targeting antibody CR3022 IgG, with IgG coupled to

chip and nanoparticle in solution. Because SARS-CoV-2 spike-SpyCatcher showed non-specific

binding only the coupled nanoparticle is shown. A series of nanoparticle concentrations was

analyzed in which the concentration of spike coupled to the nanoparticle ranged from 200 nM to

1.56 nM. Observed k_a value provided.

Fig. 6. Immunogenicity of LuS-N71-SpyLinked-CoV-2 spike

(**a**) Schematic immunization procedures for SARS-CoV-2 spike immunogens. (**b**) Serum

assessment of anti-SARS-CoV-2 spike ELISA titers. Immunization groups are color-coded.

Vertical dotted lines separate immunogen dose groups and weeks post prime. Starting reciprocal

serum dilution (100) is indicated with a horizontal dashed line. ELISA titer from each animal is

shown as an individual dot. Triangle-shape dot provided for ELISA titers at assay maximum.

Geometric means indicated by black horizontal lines. Note that the three animals immunized with

0.08 µg LuS-N71-SpyTag, which showed high ELISA titers at week 5, were the same three

animals of this control group that showed detectable neutralization. (**c**) Neutralization titer from

each animal at week 5 is shown as an individual dot, and geometric means are indicated by black

horizontal lines with values provided for each group. Immunization groups are color-coded as in

panel **b**. Limit of detection (titer = 40) indicated with a horizontal dashed line. *P* values determined

by two-tailed Mann-Whitney tests. * indicates $P \leq 0.05$, ** indicates $P \leq 0.01$, *** indicates $P \leq$

0.001 and **** indicates $P \leq 0.0001$.

Table 1. LuS- and ferritin-nanoparticles with SpyTag require the addition of *N*-linked

glycans for expression.

Construct ID	SpyTag	SpyCatcher	Position of glycan	Expression level (mg/L)
Lumazine synthase				
LuS-SpyTag no glycan	x		None	<0.1
LuS-N71-SpyTag*	x		71	**3.0**
LuS-C-SpyCatcher no glycan		x	None	<0.1
LuS-C71-SpyCatcher		x	71	<0.1
LuS-N-SpyCatcher no glycan		x	None	<0.1
LuS-N71-SpyCatcher		x	71	<0.1
Ferritin				
Ferritin-SpyTag no glycan	x		None	<0.1
Ferritin-N96-SpyTag*	x		96	**2.5**
Ferritin-S148-SpyTag*	x		146	**1.0**
Ferritin-SpyCatcher no glycan		x	None	<0.1
Ferritin-N96-SpyCatcher		x	96	<0.1
Ferritin-S148-SpyCatcher		x	146	<0.1

665 * These constructs showed suitable expression levels.
666

a

Protein A-SpyTag + Protein B-SpyCatcher → CnaB2

b

Signal peptide | SpyTag | Glycosylation site (N71) | His/Strep tags

N ――――――――――――――――――――――――――――――――――――――― C

Nanoparticle-SpyTag mammalian cell expression construct

Glycan
SpyTag
N71

LuS-N71-SpyTag (Monomer)

LuS-N71-SpyTag nanoparticle (60-mer)

Glycan
SpyTag
N96

Ferritin-N96-SpyTag (Monomer)

Ferritin-N96-SpyTag nanoparticle (24-mer)

c

LuS–N71-SpyTag

A280 (mAU) vs Elution volume (mL)

Ferritin–N96-SpyTag

A280 (mAU) vs Elution volume (mL)

d

PNGase F   1 2 3 4
           + + - -

1. LuS-N71-SpyTag + PNGase F
2. Ferritin-N96-SpyTag + PNGase F
3. LuS-N71-SpyTag
4. Ferritin-N96-SpyTag

98 —
62 —
49 —
38 —   PNGase F
28 —   Glycosylated and deglycosylated nanoparticle monomers
17 —

e

LuS-N71-SpyTag

50 nm   10 nm

Ferritin-N96-SpyTag

50 nm   10 nm

a LuS-N71-SpyTag · RSV F-SpyCatcher · LuS-N71-SpyLinked-RSV F

160 Å   150 Å

b

A280 (mAU)

LuS-N71-SpyLinked-RSV F
LuS-N71-SpyTag
RSV F-SpyCatcher

Elution volume (mL)

c

1 2 3

LuS-N71-SpyLinked-RSV F
RSV F-SpyCatcher
LuS-N71-SpyTag

1. LuS-N71-SpyTag
2. RSV F-SpyCatcher
3. LuS-N71-SpyLinked-RSV F

d LuS-N71-SpyLinked–RSV F

50 nm   10 nm

Core diameter: 18.2 ± 1.0 nm
Spike length: 11.4 ± 0.8 nm

e

Response Units (RU)

RSV F-SpyCatcher
ka: 2.18E+04 1/Ms

LS-N71-SpyLinked-RSV F
ka: 1.33E+05 1/Ms

ka: 7.32E+04 1/Ms

ka: 5.17E+04 1/Ms

Time (s)

D25 (site Ø)

MPE8 (site III)

a Ferritin-N96-SpyTag   RSV F–SpyCatcher   Ferritin-N96-SpyLinked-RSV F

108 Å

c

98 —
62 —
49 —
38 —
28 —

17 —

14 —
6 —

Ferritin-N96-SpyLinked-RSV F
RSV F-SpyCatcher

Ferritin-N96-SpyTag

1. Ferritin-N96-SpyTag
2. RSV F-SpyCatcher
3. Ferritin-N96-SpyLinked-RSV F

b

Ferritin-N96-SpyLinked-RSV F
RSV F–SpyCatcher
Ferritin-N96-SpyTag

A280 (mAU)

Elution volume (mL)

d Ferritin-N96-SpyLinked-RSV F

50 nm   10 nm

Core diameter: 15.5 ± 1.2 nm
Spike length: 12.9 ± 0.3 nm

e

RSV F-SpyCatcher
ka: 2.18E+04 1/Ms

Feritin-N96-SpyLinked-RSV F
ka: 1.69E+05 1/Ms

ka: 7.32E+04 1/Ms

ka: 3.82E+04 1/Ms

Response Units (RU)

Time (s)

D25 (site Ø)

MPE8 (site III)

a LuS-N71-SpyTag   PIV3 F-SpyCatcher   LuS-N71-SpyLinked-PIV3 F

+ → 145 Å

b

A280 (mAU)

LuS-N71-SpyTag

PIV3 F-SpyCatcher

LuS-N71-SpyLinked-PIV3 F

Elution volume (mL)

c

1   2   3

98
62   LuS-N71-SpyLinked-PIV3 F
49   PIV3 F-SpyCatcher
38
28   LuS-N71-SpyTag
17
14

1. LuS-N71-SpyTag
2. PIV3 F-SpyCatcher
3. LuS-N71-SpyLinked-PIV3 F

d LuS-N71-SpyLinked–PIV3 F

50 nm   Core diameter: 18.6 ± 1.1 nm   10 nm
Spike length: 12.6 ± 1.3 nm

e

PIV3 F-SpyCatcher
ka: 1.34E+05 1/Ms

LS-N71-SpyLinked-PIV3 F
ka: 1.16E+05 1/Ms

Response Units (RU)

ka: 1.15E+05 1/Ms

ka: 2.38E+03 1/Ms

Time (s)   Time (s)

PIA174 (head)

PIA75 (stem)

a LuS-N71-SpyTag

SARS-CoV-2 spike-SpyCatcher

LuS-N71-SpyLinked-CoV-2 spike

160 Å

200 Å

b

LuS-N71-SpyLinked-CoV-2 spike

LuS-N71-SpyTag

SARS-CoV-2 spike-SpyCatcher

A280 (mAU)

1500

1000

0   10   20

Elution volume (mL)

c

1   2   3   LuS-N71-SpyLinked-CoV-2 spike

SARS-CoV-2 spike-SpyCatcher

28   LuS-N71-SpyTag

1. LuS-N71-SpyTag
2. SARS-CoV-2 spike-SpyCatcher
3. LuS-N71-SpyLinked-CoV-2 spike

d LuS-N71-SpyLinked–CoV-2 spike

50 nm   Spike length: 22.5 ± 3.4 nm   20 nm

e SPR of LuS-N71-SpyLinked-CoV-2 spike

ka: 7.24E+05 1/Ms

CR3022 (RBD)

Response Units (RU)

0   200 400 600 800 1000

Time (s)