Disentangling interactions between mercury, immunity, and infection in a Neotropical bat

community

Daniel J. Becker[1,2*], Kelly A. Speer[3-5], Jennifer M. Korstian[6], Dmitriy V. Volokhov[7], Hannah F.

Droke[8], Alexis M. Brown[9], Catherene L. Baijnauth[10], Ticha Padgett-Stewart[11], Hugh G.

Broders[12], Raina K. Plowright[11], Thomas R. Rainwater[13-15], M. Brock Fenton[16], Nancy B.

Simmons[17], Matthew M. Chumchal[18]

[1]Department of Biology, Indiana University, Bloomington, IN, USA

[2]Center for the Ecology of Infectious Disease, University of Georgia, Athens, GA, USA

[3]Richard Gilder Graduate School, American Museum of Natural History, New York, NY, USA

[4]Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian

Institution, Washington, D.C., USA

[5]Center for Conservation Genomics, Smithsonian Conservation Biology Institute, National

Zoological Park, Washington, D.C., USA

[6]Department of Biological Sciences, Texas Tech University, Lubbock, TX, USA

[7]Center for Biologics Evaluation and Research, Food and Drug Administration, Silver Spring,

MD, USA

[8]Department of Global and Planetary Health, University of South Florida, Tampa, FL, USA

[9]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA

[10]Sackler Institute of Comparative Genomics, American Museum of Natural History, New York,

NY, USA

[11]Department of Microbiology and Immunology, Montana State University, Bozeman, MT, USA

[12]Department of Biology, University of Waterloo, Waterloo, ON, Canada

[13]Department of Forestry and Environmental Conservation, Clemson University, Clemson, SC,

USA

[14]Baruch Institute of Coastal Ecology and Forest Science, Clemson University, Georgetown, SC,

USA

[15]Tom Yawkey Wildlife Center, Georgetown, SC, USA

[16]Department of Biology, Western University, London, Ontario, Canada

31 [17]Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural

32 History, New York, NY, USA

33 [18]Department of Biology, Texas Christian University, Fort Worth, TX, USA

34 [*]Corresponding author, danbeck@iu.edu

35 **Keywords:** ecoimmunology; disease ecology; ecotoxicology; heavy metals; bacterial pathogens;

36 Chiroptera, Latin America

37 **Running head:** Mercury, immunity, and infection in bats

Abstract

1. Contaminants such as mercury are pervasive and can have immunosuppressive effects on wildlife. Impaired immunity could be important for forecasting pathogen spillover risks, as many land-use changes that generate mercury contamination can also bring wildlife reservoir hosts into close contact with humans and domestic animals. However, the interactions between contaminants, immunity, and infection are difficult to study in natural systems, and empirical tests of possible directional relationships remain rare.

2. We capitalized on extreme mercury variation in a diverse Neotropical bat community in Belize to test association between contaminants, immunity, and infection. By comparing a previous dataset of bats sampled in 2014 with new data from 2017, we first confirmed bat species with more aquatic prey in their diet had higher fur mercury. We then tested covariation between mercury and cellular immunity and determined if such relationships mediated associations between mercury and common bacterial pathogens (hemoplasmas and *Bartonella*). Lastly, as bat ecology can dictate exposure to mercury and pathogens, we also assessed species-specific patterns in these mercury–infection relationships.

3. Across the bat community, individuals with higher mercury had fewer neutrophils but not lymphocytes, suggesting stronger associations with innate immunity. However, the odds of infection for both pathogens were generally lowest in bats with high mercury, and relationships between mercury and immunity did not mediate infection patterns. When considering species-specific relationships between mercury and infection, mercury was associated with especially low odds for *Pteronotus mesoamericanus* and *Dermanura phaeotis* for hemoplasmas. Yet for *Bartonella*, mercury was associated with particularly low odds in the genus *Pteronotus* but especially high odds within the Stenodermatinae.

4. *Synthesis and application*: Lower infection risk in bats with high mercury despite weaker innate defense suggests contaminant-driven loss of pathogen habitat (i.e., anemia) or vector mortality as possible causal mechanisms. Strong positive associations between mercury and *Bartonella* in the Stenodermatinae, however, alongside negative associations between mercury and neutrophils in this clade, suggest these bats could be especially vulnerable to immunosuppression. Greater attention to these potential causal pathways could help disentangle the complex relationships between environmental contaminants, immunity, and infection in natural systems and forecast disease risks.

Introduction

Wildlife are commonly exposed to a range of contaminants that are ubiquitous in the environment, including heavy metals, organic compounds, and pesticides (Smith et al., 2007). Contaminants can be novel stressors that have direct impacts such as mortality (Davidson, 2004) and more subtle consequences such as immunosuppression (Grasman, 2002). For example, even relatively low concentrations of mercury (Hg), a neurotoxic heavy metal, can impair wildlife immunity (Scheuhammer, Meyer, Sandheinrich, & Murray, 2007). When immunosuppression manifests in increased susceptibility to pathogens, environmental gradients of contaminants could increase infection prevalence or intensity in wild hosts (Becker, Albery, et al., 2020). These elevated infection risks could be especially relevant in the context of environmental changes such as gold mining and agricultural land clearing, both of which allow contaminants like Hg to bioaccumulate in aquatic and terrestrial food webs (Almeida, Lacerda, Bastos, & Herrmann, 2005; Palheta & Taylor, 1995). As land conversions such as these also facilitate novel interactions between wildlife, domestic animals, and humans, impaired immunity could further increase pathogen spillover risks (Borremans, Faust, Manlove, Sokolow, & Lloyd-Smith, 2019).

The interactions between contaminants, immunity, and infection are notoriously difficult to study in natural systems, and empirical tests of possible directional relationships remain rare (Ross, 2002). Without captive or field experiments, natural systems must demonstrate sufficient variation in contaminant exposure and infection status to permit inference, and these criteria may be hard to meet in practice. For example, recent work on urbanized bobcats demonstrated strong effects of anticoagulant exposure on immunity in ways that should increase susceptibility, but pathogens of clinical relevance were generally rare, making it difficult to draw epidemiological inferences (Serieys et al., 2018). Additionally, effects may be difficult to detect if contaminant concentrations have low variance or are below toxicity thresholds (Fisk et al., 2005). Other examples suggest contaminants may instead decrease infection risks (Prüter et al., 2018), but the degree to which such patterns might be mediated by contaminant effects on immunity are unclear. Field-based assessments of how variable contaminant concentrations are associated with both immunity and common pathogens are necessary to disentangle directional relationships.

Here, we capitalized on high variation in Hg and infection across species in a diverse Neotropical bat community in Belize to test associations between contaminants, immunity, and infection. Although Hg concentrations are typically highest in aquatic animals owing to

bioaccumulation (Chumchal et al., 2011), such contaminants can transfer into terrestrial

ecosystems through diet (Cristol et al., 2008). Neotropical bats are ecologically diverse (Gunnell

& Simmons, 2012; Rojas, Vale, Ferrero, & Navarro, 2011), and their diet variation enables

strong heterogeneity in Hg exposure. Specifically, the degree to which species feed on aquatic

prey (or prey with at least some life stages linked to aquatic ecosystems) determines bat fur

concentrations of Hg, such that insectivores and species feeding on amphibians and fish have

greater diet exposure than frugivores and sanguivores (Becker, Chumchal, et al., 2018). Such

variation should produce strong associations with immunity, as even sublethal Hg concentrations

correlate with immunity (Becker et al., 2017). Because bats can be vulnerable to extracellular

pathogens and can also harbor viral and bacterial zoonoses (Brook & Dobson, 2015), immune

differences driven by dietary variation in Hg could have health implications for disease risks to

and from bats. This sampled region in Belize is also undergoing intensive land clearance for

agriculture similar to much of Latin America (Patterson, 2016), and thus analyses of Hg,

immunity, and infection can also help assess how land use affects wildlife and human health.

Here we built upon our prior studies of Hg and of infectious disease in Neotropical bats

(e.g., Becker, Chumchal, et al., 2018; Becker, Speer, et al., 2020) to address three study aims.

First, we assessed whether bat dietary connectivity to aquatic ecosystems is the primary driver of

Hg bioaccumulation despite spatiotemporal variation. We combined historic and new data on Hg

concentrations in bat fur, an indication of long-term metal exposure (Flache et al., 2015), and

compared contaminant load over a three-year period and two sites. Second, we tested whether

elevated bat Hg is correlated with impaired cellular immunity, using blood samples. Third, we

examined how infection with two common bacterial pathogens is associated with Hg and tested

if patterns were mediated by immunological relationships. We predicted long-term Hg exposure

would be highest in bats more reliant on aquatic food webs and that individuals of these species

would have lower measures of cellular immunity, which could manifest in higher infection risks.

Methods

Bat sampling

During April 28 to May 4 2014 and April 24 to May 6 2017, we sampled 249 bats from 29

species captured in two areas of the Orange Walk District of Belize: Lamanai Archeological

Reserve (LAR) and Ka'Kabish (KK). The LAR is bordered by the New River Lagoon, forest,

and agriculture, while KK is a remnant forest patch surrounded by agriculture located 10 km

away. At least 44 of the 70 bat species in Belize have been recorded in this region (Herrera,

Duncan, Clare, Fenton, & Simmons, 2018; Reid, 1997). Bats were captured with mist nets from

19:00 until 22:00, and harp traps were also set from 18:00 to 05:00.

Bats were placed in individual cloth bags until processing and were identified to species

based on morphology (Reid, 1997). For Hg analysis, we trimmed <10 mg of fur from the dorsal

or ventral region. Scissors were cleaned with ethanol between processed bats, and samples were

stored in individual cryovials or Ziploc bags. From a subset of bats sampled in 2017, we

collected 3–30 μL of blood by lancing the propatagial vein with sterile needles, followed by

collection using heparinized capillary tubes. Thin blood smears were prepared on glass slides and

stained with Wright–Geimsa (Astral Diagnostics Quick III) to characterize cellular immunity.

Remaining blood was stored on Whatman FTA cards (room temperature) or RNAlater (room

temperature for four weeks and then –20°C) to preserve bacterial DNA. All bats were released

after processing. Sampling followed guidelines for safe and humane handling of bats issued by

the American Society of Mammalogists (Sikes, Care, & Mammalogists, 2016) and was approved

by the University of Georgia Animal Care and Use Committee (A2014 04-016-Y3-A5).

Sampling was authorized by the Belize Forest Department under permits CD/60/3/14(27),

WL/2/1/17(16), and WL/2/1/17(19). Sample size for Hg varied by year (2014=98, 2017=148)

and by site (LAR=163, KK=83) and ranged from 1–58 individuals per species ($\bar{x}$=30 ± 29 SE).

Fur Hg analysis

Bat fur was analyzed for total Hg (THg) at the Texas Christian University Aquatic Ecology

Laboratory. THg data from 2014 were published previously (Becker, Chumchal, et al., 2018).

Fur samples were rinsed in a 2:1 chloroform:methanol solution and dried overnight in a fume

hood and reported on a fresh weight basis. We quantified THg with a direct Hg analyzer (DMA-

80, Milestone, CT, USA) and analyzed National Research Council Canada reference material

DORM 4 (certified value = 0.412 ± 0.036 mg/kg) every 10 samples for quality control; mean

recovery was 94.32 ± 0.96% for 2017 data. Limited fur resulted in some samples falling below

detection limit (DL), which was higher in 2014 (~0.48 ng). THg values below DL were

estimated as 50% DL, and we used the 2014 DL to standardize concentrations (Rainwater et al.,

2005). Fur THg was expressed in mg/kg and $\log_{10}$-transformed prior to statistical analyses. THg

is a proxy for methylmercury, which comprises 71-95% of Hg in bat fur (Yates et al., 2013).

Statistical analysis of fur THg

To first link bats to aquatic food webs (i.e., the primary source of dietary Hg exposure; Becker,

Chumchal, et al., 2018), we used the EltonTraits database to classify bat species according to the

proportion of diet consisting of potentially aquatic prey: invertebrates, ectothermic tetrapods, and

fish (Wilman et al., 2014). We used phylogenetic generalized linear mixed models (GLMMs) to

test how THg varied with bat diet, site (LAR and KK), and year (2014 and 2017). We fit

candidate models that considered all fixed effects and their two- and three-way interactions. We

fit the phylogenetic GLMM using the *brms* package in R, default priors, and Gaussian errors

(Bürkner, 2017). We included random effects for species and phylogeny, the latter of which used

a phylogenetic covariance matrix derived from the Open Tree of Life through the *rotl* and *ape*

packages (Michonneau, Brown, & Winter, 2016; Paradis, Claude, & Strimmer, 2004). We ran

four chains for 20,000 iterations with a burn-in period of 10,000, thinned every 10 steps, for a

total 4,000 samples. We compared GLMMs using leave-one-out cross-validation (LOOIC) and

assessed fit with a Bayesian R^2, including the total modeled variance and that attributed to only

the fixed effects (Gelman, Goodrich, Gabry, & Vehtari, 2019; Vehtari, Gelman, & Gabry, 2017).

We then estimated fixed effects per predictor (means and 95% highest density intervals [HDI])

and visualized fitted values with 100 random draws from the GLMM posterior distribution.

Quantifying immunity and bacterial infection

Using blood smears, we first estimated total white blood cell (WBC) counts as the mean number

of leukocytes from 10 fields of view (400X) with a light microscope (Schneeberger, Czirják, &

Voigt, 2013). We next used differential WBC counts (1000X) to quantify the relative abundance

of neutrophils and lymphocytes from 100 leukocytes. Neutrophils are components of the innate

immune system, whereas lymphocytes are involved in adaptive responses like immunoglobulin

production (Lanier, 2013). We derived absolute leukocyte counts by multiplying the total and

differential WBC data. Elevated absolute WBC counts can indicate a more robust cellular

defense or an inflammatory response to acute infection (French, Fokidis, & Moore, 2008).

To detect bacterial pathogens, we extracted genomic DNA from blood using QIAamp

DNA Investigator Kits and DNeasy Blood and Tissue Kits (Qiagen). Before extracting

RNAlater-preserved blood, samples were vortexed with 1 mL 1X phosphate-buffered saline and

centrifuged, from which 200 μL from the bottom of the tube was used for extraction. We used

PCR to test for hemoplasmas (targeting the 16S rRNA gene) and *Bartonella* spp. (targeting the

gltA gene) using previous diagnostic protocols (Bai, Gilbert, Fox, Osikowicz, & Kosoy, 2016;

Volokhov et al., 2017). Both pathogens are commonly in Neotropical bats (Ikeda et al., 2017),

and infection status can strongly correlate with bat immunity (Becker, Czirják, et al., 2018).

Hemoplasma data and sequences have been published previously (Becker, Speer, et al., 2020).

Analyses of THg, immunity, and infection

We used phylogenetic GLMMs to test the overall relationship between THg and both absolute

neutrophil and lymphocyte counts. Each model included THg as the fixed effect, with species

and phylogeny as random effects. We used a Gaussian distribution for $\log_{10}$-transformed WBC

counts. We next fit phylogenetic GLMMs with binomial errors to test associations between THg

and infection with hemoplasmas and *Bartonella*. We then used causal mediation analyses to test

support for directional relationships between THg, immunity, and infection (Imai, Keele, &

Tingley, 2010). This estimates how much of a directional relationship (i.e., outcome model) is

mediated indirectly by a third variable (i.e., mediator model). Mediator models were each of the

GLMMs predicting immunity by THg. For outcome models, we fit two GLMMs per pathogen;

each modeled infection as a function of THg and each WBC count. We used the *brms* package to

estimate the proportion of the relationship between THg and infection mediated by immunity.

Because bat ecology likely dictates exposure to contaminants and pathogens, we assessed

species-specific patterns in Hg–infection relationships. We fit logistic regressions for each bat

species and pathogen when sample sizes were sufficient (e.g., *n*>2). As small samples can bias

odds ratio estimates, we used the *logistf* package to implement Firth's bias reduction (Heinze &

Schemper, 2002). For species with no variance in infection, we assigned log odds of zero. We

estimated phylogenetic signal (Pagel's λ) in the log odds using the *caper* package and used

phylogenetic generalized least squares to test if log odds covaried with sample size (Orme,

2013). We lastly used phylogenetic factorization to identify clades with different log odds at

various taxonomic depths. We used the *taxize* and *phylofactor* packages to obtain taxonomic

information for each bat species and to partition the log odds as a Gaussian response in a GLM

(Chamberlain & Szöcs, 2013; Washburne et al., 2019). We weighted by sample size per species

and used Holm's sequentially rejective test to determine the number of significant bat clades.

Results

Neotropical bat fur THg

Expanding our initial studies of fur Hg across this Neotropical bat community in 2014, we found

that THg varied up to five orders of magnitude across species, sites, and years (Fig. 1). Our top

model included interactions between diet and year and between year and site (Table 1). The

proportion of potentially aquatic prey in diet positively predicted THg but more so in 2014

(β=0.019, 95% HDI: 0.011–0.026) than in 2017 (β=0.016, 95% HDI: 0.008–0.023; Fig. 2A). We

also identified strong spatiotemporal variation, such that fur THg increased between 2014 and

2017 for KK ($\tilde{x}$=0.46 to 0.66) but slightly decreased for LAR ($\tilde{x}$=0.52 to 0.49).

Associations with immunity and infection

Across bats sampled in 2017 with Hg and WBC data (n=52), higher THg was associated with

fewer neutrophils (β=–0.48, 95% HDI:–0.89 to –0.12) but not lymphocytes (β=–0.36, 95% HDI:

–0.83 to 0.03; Fig. 2B). However, higher THg was also associated with lower odds of infection

for hemoplasmas (OR=0.19, 95% HDI: 0.04–0.79, n=134) and *Bartonella* (OR=0.18, 95% HDI:

0.03–0.84, n=117; Fig. 3A). Accordingly, the relationship between THg and leukocytes did not

mediate relationships between THg and infection.

When we analyzed relationships between THg and infection per species and pathogen,

we only found significant protective effects for *Pteronotus mesoamericanus* and hemoplasmas

after adjusting for multiple comparisons (ln(OR)=–10.20, p=0.001; Table 2). We also detected a

strong negative association between THg and *Bartonella* in this species (ln(OR)=–3.34), but no

log odds were significantly different from zero after adjustment. Most species instead had null

relationships between THg and infection (i.e., 36% for hemoplasmas and 43% for *Bartonella*) or

relatively weaker negative effects (e.g., *Dermanura phaeotis*, ln(OR)=–7.58 for hemoplasmas; *P.*

fulvus, ln(OR)=–3.35 for *Bartonella*). These negative THg–infection associations were identified

for 29% of species for both pathogens. We also estimated similar proportions of positive THg

associations for hemoplasmas (36%) and *Bartonella* (29%). *Desmodus rotundus* had the

strongest positive THg effect size for hemoplasma infection (ln(OR)=2.40), whereas *Dermanura*

watsoni had the largest positive THg effect size for *Bartonella* infection (ln(OR)=8.9).

Comparative analyses of the log odds ratios across bat species revealed no phylogenetic

signal for the relationship between THg and hemoplasmas (λ=0) but strong phylogenetic signal

for the relationship between THg and *Bartonella* (λ=0.84). Log odds ratios were not associated

with sample size for hemoplasmas ($F_{1,12}$=1.46, *p*=0.25) or *Bartonella* ($F_{1,12}$=0.25, *p*=0.63).

Phylogenetic factorization further identified species-specific or taxonomic patterns in the

magnitude and direction of effect size. For hemoplasmas, the odds of infection were significantly

lower for *Pteronotus mesoamericanus* and *Dermanura phaeotis* when compared to the remaining

sampled bat phylogeny. For *Bartonella*, however, the odds of infection were significantly lower

for the genus *Pteronotus* (mean ln(OR)=–3.34) and significantly higher for the subfamily

Stenodermatinae (*Dermanura* spp. and *Sturnira parvidens*, mean ln(OR)=5.79; Fig. 3B). Post-

hoc GLMs showed that bats in the Stenodermatinae had especially negative associations between

THg and neutrophils (β=–2.14, *p*=0.001) but not lymphocytes (β=–0.43, *p*=0.63), but these

analyses were limited by small sample size (*n*=7).

Discussion

Interactions between contaminants, immunity, and infection are difficult to disentangle in natural

systems, but quantifying their proposed causal relationships can inform how land-use change

affects wildlife health and human disease risks. By capitalizing on a diverse Neotropical bat

system with high variation in Hg bioaccumulation and bacterial pathogens, we found higher THg

was associated with fewer neutrophils but also lower odds of infection. Our species-specific and

taxonomic analyses showed THg had protective effects for hemoplasmas and *Bartonella* in the

genus *Pteronotus*, whereas THg was associated with fewer neutrophils and elevated infection in

the subfamily Stenodermatinae. These contrasting relationships suggest contaminant-driven loss

of pathogen habitat (i.e., anemia) or vector mortality versus immunosuppression as possible

causal mechanisms, respectively, and identify clades of Neotropical bats that may be especially

resilient or vulnerable to infection risks stemming from long-term Hg exposure.

Expanding our prior analyses of this Neotropical bat community and global patterns of

THg in bat fur (Becker, Chumchal, et al., 2018) with larger within-species sample sizes, we first

demonstrated that Hg exposure increases as a function of potentially aquatic prey (or prey with at

least some life stages linked to aquatic ecosystems) in diet despite spatial and temporal variation

in THg. Consistent positive associations with diets linked to aquatic ecosystems across sites and

years provides additional support for trophic transfer of Hg through foraging (Cristol et al., 2008;

Speir et al., 2014). This diet-mediated connectivity to aquatic ecosystems likely underlies other

cases of guild-specific Hg bioaccumulation in bat communities (Carrasco-Rueda, Loiselle, &

Frederick, 2020; Korstian, Chumchal, Bennett, & Hale, 2018). In many of these regions, bat diet

exposure to Hg is driven by land-use changes such as gold mining and agriculture (Carrasco-

Rueda et al., 2020; Costantini, Czirják, Bustamante, Bumrungsri, & Voigt, 2019), whereas

atmospheric deposition is often the primary source of Hg in regions located further from

anthropogenic point sources (Chételat et al., 2018; Korstian et al., 2018). The latter is a likely

source of Hg in the Belize system (Becker, Chumchal, et al., 2018), although intensified

agriculture and slash-and-burn practices in this region could provide other Hg inputs and may

explain why bat THg increased between 2014 and 2017 in KK but not LAR (Patterson, 2016).

Across bat species, fur THg was negatively correlated with neutrophil counts, which

mirrors captive results and suggests impaired innate immunity (Lalancette, Morin, Measures, &

Fournier, 2003). Previously, vampire bats sampled in Belize with high fur THg had weaker

innate defense (i.e., bacterial killing ability; Becker et al., 2017). Wrinkle-lipped free-tailed bats

with higher Hg exposure also had weaker innate immunity (i.e., bacterial killing ability,

lysozyme and haptoglobin concentrations; Costantini et al., 2019). Most individual bats for

which we had both Hg and immunity data showed THg below toxicity and subclinical thresholds

of 5–10 mg/kg (Nam et al., 2012), which suggests innate immunity could be weakened at

sublethal contaminant concentrations (Lewis, Cristol, Swaddle, Varian-Ramos, & Zwollo, 2013).

Alternatively, sublethal effects of THg could combine with other stressors (e.g., reproduction,

food and roost availability) to impair immunity. Additional immune measures across broader

land-use gradients would help characterize functional differences in defense in relation to THg

concentrations in anthropogenic habitats (Becker, Albery, et al., 2020; Costantini et al., 2019).

Although neutrophils were lowest in bats with high Hg exposure, the odds of infection

with hemoplasmas and *Bartonella* decreased with fur THg. As in other mammals, bats

challenged with bacteria produce neutrophils as part of the innate immune response (Weise,

Czirják, Lindecke, Bumrungsri, & Voigt, 2017). Elevated neutrophil counts could be protective,

as implied through negative associations between innate immunity and these two pathogens in

vampire bats (Becker, Czirják, et al., 2018). Accordingly, relationships between THg and

immunity did not mediate the relationships between THg and infection, which likely indicates

immunosuppression is not a causal mechanism. One alternative may involve contaminant-

mediated pathogen mortality; for example, lead exposure likely caused helminth mortality in

avian hosts (Prüter et al., 2018). As intracellular pathogens, hemoplasmas and *Bartonella* both

rely on red blood cells. Hg can lower erythrocyte counts (Shaw, Dash, & Panigrahi, 1991), which

could reduce resources available to both bacteria. We did not measure anemia or Hg in blood;

however, fur THg strongly correlates with blood THg, despite the former being orders of

magnitude higher than the latter (Wada, Yates, Evers, Taylor, & Hopkins, 2010). As *Bartonella*

and possibly hemoplasmas likely depend in part on vector-borne transmission (Becker, Bergner,

et al., 2018; Willi et al., 2007), ectoparasite mortality from contaminants in hosts or the

environment could also explain observed infection patterns (e.g., as found for some avian

ectoparasites; Eeva & Klemola, 2013). Lastly, we did not reliably age bats, but older animals

typically have higher fur THg (Korstian et al., 2018). If older bats also have stronger adaptive

immunity (e.g., antibody titers in *Saccopteryx bilineata*; Schneeberger, Courtiol, Czirják, &

Voigt, 2014), age could explain negative associations between THg and infection.

Our species-specific analyses suggested that possibly protective effects of THg on

infections primarily occurred in the genus *Pteronotus*, with strong negative effects for *P.*

mesoamericanus for both pathogens. Both sampled species in this genus had high THg, being

insectivores that often eat dipterans and coleopteran (Salinas-Ramos, Herrera Montalvo, León-

Regagnon, Arrizabalaga-Escudero, & Clare, 2015). Given this strong clade-based signal,

Pteronotus bats may be particularly resilient to disease impacts of Hg exposure. Such effects

could be in part mediated by high host-specificity of bat flies for this genus (Ter Hofstede,

Fenton, & Whitaker, 2004), particularly if high THg increases vector mortality or if aspects of

bat fly biology impact vector competence (Weiss & Aksoy, 2011). In contrast, we also identified

several species with positive relationships between THg and infection. For hemoplasmas, higher

THg was associated with hemoplasma risk in vampire bats. As this species frequently feeds on

humans and livestock (Streicker & Allgeier, 2016), focused pathogen surveillance would be

particularly important in habitats that are contaminated by agriculture or mining. Additionally,

the Stenodermatinae displayed strong positive associations between THg and *Bartonella*. These

frugivorous bats also showed negative correlations between THg and neutrophils, suggesting Hg-

mediated immunosuppression. Such species may be especially vulnerable to infection from Hg

exposure and could play key roles in maintaining *Bartonella* infection cycles between bats and

ectoparasites in environments with high Hg contamination (Judson, Frank, & Hadly, 2015).

From another perspective, because the Stenodermatinae and *Pteronotus* had lower and higher fur

THg, respectively, low Hg exposure could increase susceptibility while higher concentrations

instead cause anemia and facilitate protective effects against erythrocyte-dependent pathogens.

Beyond bats and their pathogens, our study more broadly emphasizes the need to assess

potential causal relationships between contaminants and infectious diseases in natural systems. In

particular, we provide a novel perspective on integrating approaches from ecotoxicology with

those of disease ecology to disentangle the complex relationships between contaminants,

immunity, and infection. Future work that carefully integrates data on contaminant exposure,

multiple immune measures, and pathogen diversity (e.g., using metagenomics approaches;

Bergner et al., 2019) across site gradients of anthropogenic intensity could help identify

particular habitats, host species, and infections for which disease risks are highest.

Acknowledgements

For assistance with field logistics and permits, we thank Mark Howells, Neil Duncan, and staff

of the Lamanai Field Research Center. We also thank the many colleagues who helped net bats

during 2014 and 2017 bat research in Belize as well as Susan Perkins for laboratory reagents.

Data accessibility

Data from 2017 will be deposited in Dryad Digital Repository upon acceptance. Data from 2014

have been published previously (Becker, Chumchal, et al., 2018).

Author contributions

DJB, KAS, AMB, HGB, and NBS collected samples; JMK, DJB, KAS, DVV, AMB, CLB, HD,

and TPS analyzed samples; DJB analyzed the data; and HGB, RKP, TRR, MBF, NBS, and

MMC provided funding and logistical support. DJB wrote the paper with input from authors.

Funding

DJB was funded by the ARCS Foundation and Explorer's Club. Field work by DJB and KAS was supported by grants from the American Museum of Natural History Theodore Roosevelt Memorial Fund. Lab work by KAS and CLB was funded by the Richard Gilder Graduate School Student Research Fellowship. RKP was supported by NSF DEB-1716698, the Defense Advanced Research Projects Agency PREEMPT program Cooperative Agreement D18AC00031, U.S. National Institutes of General Medical Sciences IDeA Program (P20GM103474 and P30GM110732), and the USDA National Institute of Food and Agriculture (Hatch project 1015891). NBS was supported by the American Museum of Natural History Taxonomic Mammalogy Fund. JMK and MMC were supported by a Texas Christian University Research and Creative Activities Fund Award, and HGB was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant. TRR was supported by the Yawkey Foundation and Clemson University. The views, opinions, and/or findings expressed are those of the authors and should not be interpreted as representing the official views or policies of the Department of Defense or the U.S. Government.

391 **Tables**

392 Table 1. Competing phylogenetic GLMMs predicting log fur THg concentrations across the

393 Belize bat community (n=249). Models are ranked by ΔLOOIC with LOOIC SE, Akaike weights

394 (w_i) and Bayesian R^2 estimates. All models include random effects for species and phylogeny.

Fixed effects	LOOIC	SE	ΔLOOIC	w_i	R^2_m	R^2_c
~ diet + year + site + year:site + diet:year	76.21	38.83	0.00	0.39	0.80	0.92
~ diet + year + site + year:site + diet:year + diet:site	77.22	38.12	1.01	0.24	0.79	0.92
~ diet * year * site	77.64	37.03	1.43	0.19	0.79	0.92
~ diet * year + site	78.78	39.63	2.57	0.11	0.80	0.92
~ diet + year + site + diet:site + diet:year	79.57	38.96	3.36	0.07	0.79	0.92
~ diet + year + site + diet:site + year:site	86.87	33.20	10.66	>0.01	0.77	0.91
~ diet + year * site	89.33	33.93	13.13	>0.01	0.78	0.91
~ diet *site + year	89.97	33.94	13.77	>0.01	0.77	0.91
~ diet + year + site	94.43	34.22	18.23	>0.01	0.78	0.91

397 Table 2. Results of species-specific logistic regressions (using Firth's bias reduction method)

398 between bat fur THg concentrations and infection status per each bacterial pathogen.

	Hemoplasmas				*Bartonella*			
Bat species	**ln(OR)**	***p***	***σ^2***	***n***	**ln(OR)**	***p***	***σ^2***	***n***
Eptesicus furinalis	-1.545	0.4569	0.214	8	0	1	0	7
Bauerus dubiaquercus	0	1	0	3	0	1	0	3
Dermanura watsoni	-0.128	0.984	0.267	6	8.904	0.3177	0.167	6
Dermanura phaeotis	-7.578	0.2562	0.278	9	4.27	0.5789	0.214	8
Sturnira parvidens	1.099	0.7923	0.257	15	4.198	0.4613	0.194	9
Carollia sowelli	0.213	0.9427	0.178	10	-1.987	0.5159	0.233	10
Trachops cirrhosus	0	1	0	3	0	1	0	3
Chrotopterus auritus	0	1	0	3	0	1	0	3
Glossophaga soricina	0.665	0.8421	0.218	11	-2.373	0.4911	0.194	9
Desmodus rotundus	2.395	0.3328	0.208	22	0.178	0.933	0.208	22
Pteronotus mesoamericanus	-10.201	0.0001	0.227	22	-3.339	0.0159	0.25	16
Pteronotus fulvus	1.653	0.65	0.25	4	-3.346	0.4073	0.25	4
Rhynchonycteris naso	0	1	0	4	0	1	0	3
Saccopteryx bilineata	0	1	0	8	0	1	0	8

Figures and legends

Figure 1. THg concentrations in bat fur across Neotropical bat species sampled in 2014 and 2017

across two sites in Belize. Boxplots are colored by the proportion of potentially aquatic prey in

bat diets from the EltonTraits database. THg concentrations are displayed on a $\log_{10}$ scale.

408   Figure 2. Dietary drivers of THg concentrations in Neotropical bat fur (A) and cellular immunity

409   correlates of THg (B). Points indicate individual bats and are colored by sampling year in Belize.

410   Lines display 100 random draws from the posterior distribution of the main phylogenetic

411   GLMMs. THg concentrations and absolute leukocyte counts are displayed on a $\log_{10}$ scale.

References

Almeida, M. D., Lacerda, L. D., Bastos, W. R., & Herrmann, J. C. (2005). Mercury loss from soils following conversion from forest to pasture in Rondônia, Western Amazon, Brazil. *Environmental Pollution*, *137*(2), 179–186.

Bai, Y., Gilbert, A., Fox, K., Osikowicz, L., & Kosoy, M. (2016). Bartonella rochalimae and b. vinsonii subsp. berkhoffii in wild carnivores from colorado, usa. *Journal of Wildlife Diseases*, *52*(4), 844–849. doi: 10.7589/2016-01-015

Becker, D. J., Albery, G. F., Kessler, M. K., Lunn, T. J., Falvo, C. A., Czirják, G. Á., … Plowright, R. K. (2020). Macroimmunology: The drivers and consequences of spatial patterns in wildlife immune defence. *Journal of Animal Ecology*, *89*(4), 972–995. doi: 10.1111/1365-2656.13166

Becker, D. J., Bergner, L. M., Bentz, A. B., Orton, R. J., Altizer, S., & Streicker, D. G. (2018). Genetic diversity, infection prevalence, and possible transmission routes of Bartonella spp. in vampire bats. *PLOS Neglected Tropical Diseases*, *12*(9), e0006786. doi: 10.1371/journal.pntd.0006786

Becker, D. J., Chumchal, M. M., Bentz, A. B., Platt, S. G., Czirják, G. Á., Rainwater, T. R., … Streicker, D. G. (2017). Predictors and immunological correlates of sublethal mercury exposure in vampire bats. *Royal Society Open Science*, *4*(4), 170073. doi: 10.1098/rsos.170073

Becker, D. J., Chumchal, M. M., Broders, H. G., Korstian, J. M., Clare, E. L., Rainwater, T. R., … Fenton, M. B. (2018). Mercury bioaccumulation in bats reflects dietary connectivity to aquatic food webs. *Environmental Pollution*, *233*, 1076–1085. doi: 10.1016/j.envpol.2017.10.010

Becker, D. J., Czirják, G. Á., Volokhov, D. V., Bentz, A. B., Carrera, J. E., Camus, M. S., …

449   Streicker, D. G. (2018). Livestock abundance predicts vampire bat demography, immune

450     profiles and bacterial infection risk. *Phil. Trans. R. Soc. B*, *373*(1745), 20170089. doi:

451     10.1098/rstb.2017.0089

452 Becker, D. J., Speer, K. A., Brown, A. M., Fenton, M. B., Washburne, A. D., Altizer, S., …

453     Volokhov, D. V. (2020). Ecological and evolutionary drivers of hemoplasma infection

454     and bacterial genotype sharing in a Neotropical bat community. *Molecular Ecology*,

455     *n/a*(n/a). doi: 10.1111/mec.15422

456 Bergner, L. M., Orton, R. J., Filipe, A. da S., Shaw, A. E., Becker, D. J., Tello, C., … Streicker,

457     D. G. (2019). Using noninvasive metagenomics to characterize viral communities from

458     wildlife. *Molecular Ecology Resources*, *19*(1), 128–143. doi: 10.1111/1755-0998.12946

459 Borremans, B., Faust, C., Manlove, K. R., Sokolow, S. H., & Lloyd-Smith, J. O. (2019). Cross-

460     species pathogen spillover across ecosystem boundaries: mechanisms and theory.

461     *Philosophical Transactions of the Royal Society B*, *374*(1782), 20180344.

462 Brook, C. E., & Dobson, A. P. (2015). Bats as 'special'reservoirs for emerging zoonotic

463     pathogens. *Trends in Microbiology*, *23*(3), 172–180.

464 Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal*

465     *of Statistical Software*, *80*(1), 1–28.

466 Carrasco-Rueda, F., Loiselle, B. A., & Frederick, P. C. (2020). Mercury bioaccumulation in

467     tropical bats from a region of active artisanal and small-scale gold mining. *Ecotoxicology*,

468     1–11.

469 Chamberlain, S. A., & Szöcs, E. (2013). taxize: taxonomic search and retrieval in R.

470     *F1000Research*, *2*.

471 Chételat, J., Hickey, M. B. C., Poulain, A. J., Dastoor, A., Ryjkov, A., McAlpine, D., …

Zanuttig, M. (2018). Spatial variation of mercury bioaccumulation in bats of Canada

linked to atmospheric mercury deposition. *Science of The Total Environment*, *626*, 668–

677. doi: 10.1016/j.scitotenv.2018.01.044

Chumchal, M. M., Rainwater, T. R., Osborn, S. C., Roberts, A. P., Abel, M. T., Cobb, G. P., …

Bailey, F. C. (2011). Mercury speciation and biomagnification in the food web of Caddo

Lake, Texas and Louisiana, USA, a subtropical freshwater ecosystem. *Environmental*

Toxicology and Chemistry, *30*(5), 1153–1162.

Costantini, D., Czirják, G. Á., Bustamante, P., Bumrungsri, S., & Voigt, C. C. (2019). Impacts of

land use on an insectivorous tropical bat: The importance of mercury, physio-

immunology and trophic position. *Science of The Total Environment*, *671*, 1077–1085.

doi: 10.1016/j.scitotenv.2019.03.398

Cristol, D. A., Brasso, R. L., Condon, A. M., Fovargue, R. E., Friedman, S. L., Hallinger, K. K.,

… White, A. E. (2008). The Movement of Aquatic Mercury Through Terrestrial Food

Webs. *Science*, *320*(5874), 335–335. doi: 10.1126/science.1154082

Davidson, C. (2004). Declining downwind: amphibian population declines in California and

historical pesticide use. *Ecological Applications*, *14*(6), 1892–1902.

Eeva, T., & Klemola, T. (2013). Variation in prevalence and intensity of two avian ectoparasites

in a polluted area. *Parasitology*, *140*(11), 1384–1393.

Fisk, A. T., de Wit, C. A., Wayland, M., Kuzyk, Z. Z., Burgess, N., Letcher, R., … Muir, D. C.

G. (2005). An assessment of the toxicological significance of anthropogenic

contaminants in Canadian arctic wildlife. *Science of The Total Environment*, *351–352*,

57–93. doi: 10.1016/j.scitotenv.2005.01.051

Flache, L., Becker, N. I., Kierdorf, U., Czarnecki, S., Düring, R.-A., & Encarnação, J. A. (2015).

Hair samples as monitoring units for assessing metal exposure of bats: a new tool for risk

assessment. *Mammalian Biology-Zeitschrift Für Säugetierkunde*, *80*(3), 178–181.

French, S. S., Fokidis, H. B., & Moore, M. C. (2008). Variation in stress and innate immunity in

the tree lizard (Urosaurus ornatus) across an urban–rural gradient. *Journal of*

Comparative Physiology B, *178*(8), 997–1005. doi: 10.1007/s00360-008-0290-8

Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian Regression

Models. *The American Statistician*, *73*(3), 307–309. doi:

10.1080/00031305.2018.1549100

Grasman, K. A. (2002). Assessing immunological function in toxicological studies of avian

wildlife. *Integrative and Comparative Biology*, *42*(1), 34–42.

Gunnell, G. F., & Simmons, N. B. (2012). *Evolutionary history of bats: fossils, molecules and*

morphology. Cambridge University Press.

Heinze, G., & Schemper, M. (2002). A solution to the problem of separation in logistic

regression. *Statistics in Medicine*, *21*(16), 2409–2419.

Herrera, J. P., Duncan, N., Clare, E., Fenton, M. B., & Simmons, N. (2018). Disassembly of

Fragmented Bat Communities in Orange Walk District, Belize. *Acta Chiropterologica*,

20(1), 147–159. doi: 10.3161/15081109ACC2018.20.1.011

Ikeda, P., Seki, M. C., Carrasco, A. O. T., Rudiak, L. V., Miranda, J. M. D., Gonçalves, S. M.

M., … others. (2017). Evidence and molecular characterization of Bartonella spp. and

hemoplasmas in neotropical bats in Brazil. *Epidemiology & Infection*, 1–15.

Imai, K., Keele, L., & Tingley, D. (2010). A general approach to causal mediation analysis.

Psychological Methods, *15*(4), 309–334. doi: 10.1037/a0020761

Judson, S. D., Frank, H. K., & Hadly, E. A. (2015). Bartonellae are Prevalent and Diverse in

Costa Rican Bats and Bat Flies. *Zoonoses and Public Health*, *62*(8), 609–617. doi:

10.1111/zph.12188

Korstian, J. M., Chumchal, M. M., Bennett, V. J., & Hale, A. M. (2018). Mercury contamination

in bats from the central United States. *Environmental Toxicology and Chemistry*, *37*(1),

160–165.

Lalancette, A., Morin, Y., Measures, L., & Fournier, M. (2003). Contrasting changes of

sensitivity by lymphocytes and neutrophils to mercury in developing grey seals.

Developmental & Comparative Immunology, *27*(8), 735–747.

Lanier, L. L. (2013). Shades of grey—the blurring view of innate and adaptive immunity. *Nature*

Reviews Immunology, *13*(2), 73.

Lewis, C. A., Cristol, D. A., Swaddle, J. P., Varian-Ramos, C. W., & Zwollo, P. (2013).

Decreased immune response in zebra finches exposed to sublethal doses of mercury.

Archives of Environmental Contamination and Toxicology, *64*(2), 327–336.

Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl : an R package to interact with the

Open Tree of Life data. *Methods in Ecology and Evolution*. doi: 10.1111/2041-

210X.12593

Nam, D.-H., Yates, D., Ardapple, P., Evers, D. C., Schmerfeld, J., & Basu, N. (2012). Elevated

mercury exposure and neurochemical alterations in little brown bats (Myotis lucifugus)

from a site with historical mercury contamination. *Ecotoxicology*, *21*(4), 1094–1101. doi:

10.1007/s10646-012-0864-9

Orme, D. (2013). The caper package: comparative analysis of phylogenetics and evolution in R.

R Package Version, *5*(2). Retrieved from

ftp://ctan.mirrorcatalogs.com/cran/web/packages/caper/vignettes/caper.pdf

Palheta, D., & Taylor, A. (1995). Mercury in environmental and biological samples from a gold

mining area in the Amazon region of Brazil. *Science of the Total Environment*, *168*(1),

63–69.

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: analyses of phylogenetics and evolution in

R language. *Bioinformatics*, *20*(2), 289–290.

Patterson, C. (2016). *Deforestation, agricultural intensification, and farm resilience in Northern*

Belize: 1980-2010 (Thesis, University of Otago). Retrieved from

https://ourarchive.otago.ac.nz/handle/10523/6858

Prüter, H., Franz, M., Auls, S., Czirják, G. Á., Greben, O., Greenwood, A. D., … Krone, O.

(2018). Chronic lead intoxication decreases intestinal helminth species richness and

infection intensity in mallards (Anas platyrhynchos). *Science of the Total Environment*,

644, 151–160.

Rainwater, T. R., Reynolds, K. D., Cañas, J. E., Cobb, G. P., Andersonv, T. A., McMurry, S. T.,

& Smith, P. N. (2005). Organochlorine pesticides and mercury in cottonmouths

(Agkistrodon piscivorus) from northeastern Texas, USA. *Environmental Toxicology and*

Chemistry, *24*(3), 665–673. doi: 10.1897/04-223R.1

Reid, F. (1997). *A Field Guide to the Mammals of Central America and Southeast Mexico*. OUP

USA.

Rojas, D., Vale, Á., Ferrero, V., & Navarro, L. (2011). When did plants become important to

leaf-nosed bats? Diversification of feeding habits in the family Phyllostomidae.

Molecular Ecology, *20*(10), 2217–2228. doi: 10.1111/j.1365-294X.2011.05082.x

Ross, P. S. (2002). The role of immunotoxic environmental contaminants in facilitating the

emergence of infectious diseases in marine mammals. *Human and Ecological Risk*

Assessment: An International Journal, *8*(2), 277–292.

Salinas-Ramos, V. B., Herrera Montalvo, L. G., León-Regagnon, V., Arrizabalaga-Escudero, A.,

& Clare, E. L. (2015). Dietary overlap and seasonality in three species of mormoopid bats

from a tropical dry forest. *Molecular Ecology*, *24*(20), 5296–5307.

Scheuhammer, A. M., Meyer, M. W., Sandheinrich, M. B., & Murray, M. W. (2007). Effects of

environmental methylmercury on the health of wild birds, mammals, and fish. *AMBIO: A*

Journal of the Human Environment, *36*(1), 12–19.

Schneeberger, K., Courtiol, A., Czirják, G. Á., & Voigt, C. C. (2014). Immune Profile Predicts

Survival and Reflects Senescence in a Small, Long-Lived Mammal, the Greater Sac-

Winged Bat (Saccopteryx bilineata). *PLoS ONE*, *9*(9), e108268. doi:

10.1371/journal.pone.0108268

Schneeberger, K., Czirják, G. Á., & Voigt, C. C. (2013). Measures of the Constitutive Immune

System Are Linked to Diet and Roosting Habits of Neotropical Bats. *PLoS ONE*, *8*(1),

e54023. doi: 10.1371/journal.pone.0054023

Serieys, L. E. K., Lea, A. J., Epeldegui, M., Armenta, T. C., Moriarty, J., VandeWoude, S., …

Uittenbogaart, C. H. (2018). Urbanization and anticoagulant poisons promote immune

dysfunction in bobcats. *Proc. R. Soc. B*, *285*(1871), 20172533. doi:

10.1098/rspb.2017.2533

Shaw, B. P., Dash, S., & Panigrahi, A. K. (1991). Effect of methyl mercuric chloride treatment

on haematological characteristics and erythrocyte morphology of Swiss mice.

Environmental Pollution, *73*(1), 43–52.

Sikes, R. S., Care, A., & Mammalogists, U. C. of the A. S. of. (2016). 2016 Guidelines of the

American Society of Mammalogists for the use of wild mammals in research and

education. *Journal of Mammalogy*, *97*(3), 663–688.

Smith, P. N., Cobb, G. P., Godard-Codding, C., Hoff, D., McMurry, S. T., Rainwater, T. R., &

Reynolds, K. D. (2007). Contaminant exposure in terrestrial vertebrates. *Environmental*

Pollution, *150*(1), 41–64. doi: 10.1016/j.envpol.2007.06.009

Speir, S. L., Chumchal, M. M., Drenner, R. W., Cocke, W. G., Lewis, M. E., & Whitt, H. J.

(2014). Methyl mercury and stable isotopes of nitrogen reveal that a terrestrial spider has

a diet of emergent aquatic insects. *Environmental Toxicology and Chemistry*, *33*(11),

2506–2509. doi: 10.1002/etc.2700

Streicker, D. G., & Allgeier, J. E. (2016). Foraging choices of vampire bats in diverse

landscapes: potential implications for land-use change and disease transmission. *Journal*

of Applied Ecology, *53*(4), 1280–1288. doi: 10.1111/1365-2664.12690

Ter Hofstede, H. M., Fenton, M. B., & Whitaker, J. O. (2004). Host and host-site specificity of

bat flies (Diptera: Streblidae and Nycteribiidae) on Neotropical bats (Chiroptera).

Canadian Journal of Zoology, *82*(4), 616–626.

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-

one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

Volokhov, D. V., Becker, D. J., Bergner, L. M., Camus, M. S., Orton, R. J., Chizhikov, V. E., …

Streicker, D. G. (2017). Novel hemotropic mycoplasmas are widespread and genetically

diverse in vampire bats. *Epidemiology & Infection*, *145*(15), 3154–3167.

Wada, H., Yates, D. E., Evers, D. C., Taylor, R. J., & Hopkins, W. A. (2010). Tissue mercury

concentrations and adrenocortical responses of female big brown bats (Eptesicus fuscus)

near a contaminated river. *Ecotoxicology*, *19*(7), 1277–1284. doi: 10.1007/s10646-010-

0513-0

Washburne, A. D., Silverman, J. D., Morton, J. T., Becker, D. J., Crowley, D., Mukherjee, S., …

Plowright, R. K. (2019). Phylofactorization: a graph partitioning algorithm to identify

phylogenetic scales of ecological data. *Ecological Monographs*, *0*(0), e01353. doi:

10.1002/ecm.1353

Weise, P., Czirják, G. A., Lindecke, O., Bumrungsri, S., & Voigt, C. C. (2017). Simulated

bacterial infection disrupts the circadian fluctuation of immune cells in wrinkle-lipped

bats (Chaerephon plicatus). *PeerJ*, *5*, e3570. doi: 10.7717/peerj.3570

Weiss, B., & Aksoy, S. (2011). Microbiome influences on insect host vector competence. *Trends*

in Parasitology, *27*(11), 514–522.

Willi, B., Boretti, F. S., Tasker, S., Meli, M. L., Wengi, N., Reusch, C. E., … Hofmann-

Lehmann, R. (2007). From Haemobartonella to hemoplasma: Molecular methods provide

new insights. *Veterinary Microbiology*, *125*(3–4), 197–209. doi:

10.1016/j.vetmic.2007.06.027

Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., & Jetz, W. (2014).

EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals.

Ecology, *95*(7), 2027–2027. doi: 10.1890/13-1917.1

Yates, D. E., Adams, E. M., Angelo, S. E., Evers, D. C., Schmerfeld, J., Moore, M. S., …

O'Driscoll, N. J. (2013). Mercury in bats from the northeastern United States.

Ecotoxicology, *23*(1), 45–55. doi: 10.1007/s10646-013-1150-1