Disentangling interactions among mercury, immunity, and infection in a Neotropical bat community

Daniel J. Becker[1*], Kelly A. Speer[2-4], Jennifer M. Korstian[5], Dmitriy V. Volokhov[6], Hannah F. Droke[7], Alexis M. Brown[8], Catherene L. Baijnauth[9], Ticha Padgett-Stewart[10], Hugh G. Broders[11], Raina K. Plowright[10], Thomas R. Rainwater[12-14], M. Brock Fenton[15], Nancy B. Simmons[16], Matthew M. Chumchal[17]

[1]Department of Biology, University of Oklahoma, Norman, OK, USA

[2]Richard Gilder Graduate School, American Museum of Natural History, New York, NY, USA

[3]Department of Invertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, D.C., USA

[4]Center for Conservation Genomics, Smithsonian Conservation Biology Institute, National Zoological Park, Washington, D.C., USA

[5]Department of Biological Sciences, Texas Tech University, Lubbock, TX, USA

[6]Center for Biologics Evaluation and Research, Food and Drug Administration, Silver Spring, MD, USA

[7]Department of Global and Planetary Health, University of South Florida, Tampa, FL, USA

[8]Department of Ecology and Evolution, Stony Brook University, Stony Brook, NY, USA

[9]Sackler Institute of Comparative Genomics, American Museum of Natural History, New York, NY, USA

[10]Department of Microbiology and Immunology, Montana State University, Bozeman, MT, USA

[11]Department of Biology, University of Waterloo, Waterloo, ON, Canada

[12]Department of Forestry and Environmental Conservation, Clemson University, Clemson, SC, USA

[13]Baruch Institute of Coastal Ecology and Forest Science, Clemson University, Georgetown, SC, USA

[14]Tom Yawkey Wildlife Center, Georgetown, SC, USA

[15]Department of Biology, Western University, London, Ontario, Canada

[16]Department of Mammalogy, Division of Vertebrate Zoology, American Museum of Natural History, New York, NY, USA

32 [17]Department of Biology, Texas Christian University, Fort Worth, TX, USA

33 [*]Corresponding author, danbeck@ou.edu

34 **Keywords:** ecoimmunology; disease ecology; ecotoxicology; heavy metals; bacterial pathogens;

35 Chiroptera, Latin America

36 **Running head:** Mercury, immunity, and infection in bats

Abstract (350/350 words)

1. Contaminants such as mercury are pervasive and can have immunosuppressive effects on wildlife. Impaired immunity could be important for forecasting pathogen spillover risks, as many land-use changes that generate mercury contamination also bring wildlife into close contact with humans and domestic animals. However, the interactions among contaminants, immunity, and infection are difficult to study in natural systems, and empirical tests of possible directional relationships remain rare.

2. We capitalized on extreme mercury variation in a diverse bat community in Belize to test association among contaminants, immunity, and infection. By comparing a previous dataset of bats sampled in 2014 with new data from 2017, representing a period of rapid agricultural land conversion, we first confirmed bat species more reliant on aquatic prey had higher fur mercury. Bats in the agricultural habitat also had higher mercury in recent years. We then tested covariation between mercury and cellular immunity and determined if such relationships mediated associations between mercury and common bacterial pathogens . As bat ecology can dictate exposure to mercury and pathogens, we also assessed species-specific patterns in mercury–infection relationships.

3. Across the bat community, individuals with higher mercury had fewer neutrophils but not lymphocytes, suggesting stronger associations with innate immunity. However, the odds of infection for hemoplasmas and *Bartonella* spp. were generally lowest in bats with high mercury, and relationships between mercury and immunity did not mediate infection patterns. Mercury also showed species- and clade-specific relationships with infection, being associated with especially low odds for hemoplasmas in *Pteronotus mesoamericanus* and *Dermanura phaeotis*. For *Bartonella* spp., mercury was associated with particularly low odds in the genus *Pteronotus* but high odds in the Stenodermatinae.

4. *Synthesis and application*: Lower general infection risk in bats with high mercury despite weaker innate defense suggests contaminant-driven loss of pathogen habitat (i.e., anemia) or vector mortality as possible causes. Greater attention to these potential pathways could help disentangle relationships among contaminants, immunity, and infection in anthropogenic habitats and help forecast disease risks. Our results also suggest contaminants may increase infection risk in some taxa but not others, emphasizing the importance of considering surveillance and management at different phylogenetic scales.

Introduction

Wildlife are commonly exposed to many contaminants that are ubiquitous in the environment, including heavy metals, organic compounds, and pesticides (Smith et al., 2007). Contaminants can be novel stressors that have direct impacts such as mortality (Davidson, 2004) and more subtle consequences such as immunosuppression (Grasman, 2002). For example, even relatively low concentrations of mercury (Hg), a neurotoxic heavy metal, can impair wildlife immunity (Scheuhammer, Meyer, Sandheinrich, & Murray, 2007). When immunosuppression manifests in increased susceptibility to pathogens, environmental gradients of contaminants could increase infection prevalence or intensity in wild hosts (Becker, Albery, et al., 2020). These elevated infection risks could be especially relevant in the context of environmental changes such as gold mining and agricultural land clearing, both of which are associated with bioaccumulation of contaminants like Hg in aquatic and terrestrial food webs (Farella, Davidson, Lucotte, & Daigle, 2007; Palheta & Taylor, 1995). As land conversions such as these also facilitate novel interactions among wildlife, domestic animals, and humans, impaired immunity could further increase pathogen spillover risks (Borremans, Faust, Manlove, Sokolow, & Lloyd-Smith, 2019).

The interactions among contaminants, immunity, and infection are notoriously difficult to study in natural systems, and empirical tests of possible directional relationships remain rare (Ross, 2002). Without captive or field experiments, natural systems must demonstrate sufficient variation in contaminant exposure and infection status to permit inference, and these criteria may be hard to meet in practice. For example, recent work on urbanized bobcats demonstrated strong effects of anticoagulant exposure on immunity in ways that should increase susceptibility, but pathogens of clinical relevance were generally rare, making it difficult to draw epidemiological inferences (Serieys et al., 2018). Additionally, effects may be difficult to detect if contaminant concentrations have low variance or are below toxicity thresholds (Fisk et al., 2005). Other examples suggest contaminants may instead decrease infection risks (Prüter et al., 2018), but the degree to which such patterns might be mediated by contaminant effects on immunity are unclear. Field-based assessments of how variable contaminant concentrations are associated with immunity and common pathogens are necessary to disentangle directional relationships.

Here, we capitalized on high variation in Hg and infection across species in a diverse Neotropical bat community in Belize to test associations among contaminants, immunity, and infection. Hg concentrations are typically highest in aquatic animals because methylmercury

(MeHg), the bioaccumulative form of Hg, is produced in aquatic ecosystems (Chumchal et al.,

2011). However, such contaminants can transfer into terrestrial ecosystems when terrestrial

consumers feed on aquatic prey contaminated with MeHg (Cristol et al., 2008). Neotropical bats

are ecologically diverse (Gunnell & Simmons, 2012; Rojas, Vale, Ferrero, & Navarro, 2011),

and their diet variation enables strong heterogeneity in Hg exposure. Specifically, our past work

showed that how often species feed on aquatic prey (or prey with some life stages linked to

aquatic ecosystems) determines bat fur Hg, such that insectivores and species feeding on

amphibians and fish have greater diet exposure than frugivores and sanguivores (Becker,

Chumchal, et al., 2018). Such variation should produce strong associations with immunity, as

even sublethal Hg concentrations correlate with immunity (Becker et al., 2017). Because bats can

be vulnerable to extracellular pathogens and can also harbor viral and bacterial zoonoses (Brook

& Dobson, 2015), immune differences driven by dietary variation in Hg could have implications

for disease risks to and from bats. Lastly, this region in Belize is also undergoing intensive land

clearance for agriculture similar to much of Latin America (Patterson, 2016), and thus analyses

of Hg, immunity, and infection could help assess how land use affects wildlife and human health.

Here we built upon our prior studies of Hg and of infectious disease in Neotropical bats

(e.g., Becker, Chumchal, et al., 2018; Becker, Speer, et al., 2020) to address three study aims.

First, we combined historic and new data on Hg concentrations in bat fur, an indication of long-

term metal exposure (Flache et al., 2015), and compared contaminant load over a three-year

period and two sites. This greater within-species sample size allowed us to more robustly assess

whether bat dietary connectivity to aquatic ecosystems predicts Hg bioaccumulation and if such

patterns persist despite spatiotemporal variation. Additionally, because agriculture can increase

environmental concentrations of Hg (Costantini, Czirják, Bustamante, Bumrungsri, & Voigt,

2019; Farella et al., 2007), we also tested whether more recently sampled bats within this rapidly

changing landscape had higher Hg exposure. Second, using blood samples, we tested whether

elevated bat Hg is correlated with impaired immune function. Expanding our species-specific

analyses in vampire bats (*Desmodus rotundus*; Becker et al., 2017) across the entire bat

community, we predicted that individuals with high Hg would have lower measures of cellular

immunity. Third, we assessed infection with two bacterial pathogens, hemoplasmas and

Bartonella spp. Both are common in Neotropical bats (Ikeda et al., 2017), including in these

Belize sites (Becker, Bergner, et al., 2018; Becker, Speer, et al., 2020), and we previously

showed infection can correlate with immunity in vampire bats specifically (Becker, Czirják, et

al., 2018). However, how Hg shapes infection risk, and if such patterns are mediated by

immunological relationships, is unknown. If Hg is associated with lower cellular immunity

across bat species, we would expect greater concentrations to manifest in higher infection risks.

Methods

Bat sampling

During April 28 to May 4 2014 and April 24 to May 6 2017, we sampled 249 bats from 29

species captured in two areas of the Orange Walk District of Belize: Lamanai Archeological

Reserve (LAR) and Ka'Kabish (KK). The LAR is bordered by the New River Lagoon, forest,

and agriculture, while KK is a remnant forest patch surrounded by agriculture located 10 km

away. At least 44 of the 70 bat species in Belize have been recorded in this region (Herrera,

Duncan, Clare, Fenton, & Simmons, 2018; Reid, 1997). Bats were captured with mist nets from

19:00 until 22:00, and harp traps were also set from 18:00 to 05:00.

Bats were placed in individual cloth bags until processing and were identified to species

based on morphology, including but not limited to body mass and forearm length (Reid, 1997).

For Hg analysis, we trimmed <10 mg of fur from the dorsal or ventral region. Scissors were

cleaned with ethanol between processed bats, and samples were stored in individual cryovials or

Ziploc bags. From a subset of bats sampled in 2017, we collected 3–30 μL of blood by lancing

the propatagial vein with sterile needles (23–30G; size and volume were dependent on bat mass),

followed by collection using heparinized capillary tubes. Thin blood smears were prepared on

glass slides and stained with Wright–Geimsa (Astral Diagnostics Quick III) to characterize

cellular immunity. Remaining blood was stored on Whatman FTA cards (room temperature) or

RNAlater (room temperature for four weeks and then –20°C) to preserve DNA. All bats were

released after processing. Sampling followed guidelines for safe and humane handling of bats

issued by the American Society of Mammalogists (Sikes, Care, & Mammalogists, 2016) and was

approved by the University of Georgia Animal Care and Use Committee (A2014 04-016-Y3-

A5). Sampling was authorized by the Belize Forest Department under permits CD/60/3/14(27),

WL/2/1/17(16), and WL/2/1/17(19). Sample size for Hg varied by year (2014=98, 2017=148)

and site (LAR=163, KK=83) and ranged from 1–58 individuals per species ($\overline{x}$=30 ± 29 SE).

Fur Hg analysis

Bat fur was analyzed for total Hg (THg) at the Texas Christian University Aquatic Ecology

Laboratory. THg data from 2014 were published previously (Becker, Chumchal, et al., 2018).

Fur samples were rinsed in a 2:1 chloroform:methanol solution and dried overnight in a fume

hood and reported on a fresh weight basis. We quantified THg with a direct Hg analyzer (DMA-

80, Milestone, CT, USA) and analyzed National Research Council Canada reference material

DORM 4 (certified value = 0.412 ± 0.036 mg/kg) every 10 samples for quality control; mean

recovery was 94.32 ± 0.96% for 2017 data. Limited fur resulted in some samples falling below

detection limit (DL), which was higher in 2014 (~0.48 ng). THg values below DL were

estimated as 50% DL, and we used the 2014 DL to standardize concentrations (Rainwater et al.,

2005). Fur THg was expressed in mg/kg and $\log_{10}$-transformed prior to statistical analyses. THg

is a proxy for MeHg, which comprises 71–95% of Hg in bat fur (Yates et al., 2013).

Statistical analysis of fur THg

To first link bats to aquatic food webs (i.e., the primary source of dietary Hg exposure; Becker,

Chumchal, et al., 2018), we used the EltonTraits database to classify bat species according to the

proportion of diet consisting of potentially aquatic prey: invertebrates, ectothermic tetrapods, and

fish (Wilman et al., 2014). We used phylogenetic generalized linear mixed models (GLMMs) to

test how THg varied with bat diet, site (LAR and KK), and year (2014 and 2017). We fit

candidate models that considered all fixed effects and their two- and three-way interactions. We

fit the phylogenetic GLMM using the *brms* package in R, default priors, and Gaussian errors

(Bürkner, 2017). We included random effects for species and phylogeny, the latter of which used

a phylogenetic covariance matrix derived from the Open Tree of Life through the *rotl* and *ape*

packages (Michonneau, Brown, & Winter, 2016; Paradis, Claude, & Strimmer, 2004). We ran

four chains for 20,000 iterations with a burn-in period of 10,000, thinned every 10 steps, for a

total 4,000 samples. We compared GLMMs using leave-one-out cross-validation (LOOIC) and

assessed fit with a Bayesian R^2, including the total modeled variance and that attributed to only

the fixed effects (Gelman, Goodrich, Gabry, & Vehtari, 2019; Vehtari, Gelman, & Gabry, 2017).

We then estimated fixed effects per predictor (means and 95% highest density intervals [HDI])

and visualized fitted values using 100 random draws from the GLMM posterior distribution.

Quantifying immunity and bacterial infection

Using blood smears, we first estimated total white blood cell (WBC) counts as the mean number of leukocytes from 10 fields of view (400X) with a light microscope (Schneeberger, Czirják, & Voigt, 2013). We next used differential WBC counts (1000X) to quantify the relative abundance of neutrophils and lymphocytes from 100 leukocytes. Neutrophils are components of the innate immune system, whereas lymphocytes are involved in adaptive responses like immunoglobulin production (Lanier, 2013). We derived absolute neutrophil and lymphocyte counts separately by multiplying total and differential WBCs. Elevated WBC counts can indicate a more robust cellular defense or an inflammatory response to acute infection.

To detect bacterial pathogens, we extracted genomic DNA from blood using QIAamp DNA Investigator Kits and DNeasy Blood and Tissue Kits (Qiagen). Before extracting RNAlater-preserved blood, samples were vortexed with 1 mL 1X phosphate-buffered saline and centrifuged for three minutes at 4000 RPM and 12 °C to prevent bacteria from floating in RNAlater; 200 μL from the bottom of the tube was then used for extraction. We used PCR to test for hemoplasmas (targeting the 16S rRNA gene) and *Bartonella* spp. (targeting the *gltA* gene) using previous diagnostic protocols (Bai, Gilbert, Fox, Osikowicz, & Kosoy, 2016; Volokhov et al., 2017). Hemoplasma data and sequences have been published previously (Becker, Speer, et al., 2020). Efforts to characterize *Bartonell*a spp. in this bat community are ongoing, but prior studies of Belize vampire bats indicate high *gltA* sequence similarity to sequences from vampire bats in Mexico, Neotropical bat flies, and other Neotropical bats (Becker, Bergner, et al., 2018).

Analyses of THg, immunity, and infection

We first used phylogenetic GLMMs to test the overall relationship between THg and both absolute neutrophil and lymphocyte counts. Each model included THg as the fixed effect, alongside sex and body condition (mass/forearm length) as covariates that could also affect leukocyte counts, with species and phylogeny as random effects. We used a Gaussian distribution for $\log_{10}$-transformed WBC counts. We next fit phylogenetic GLMMs with binomial errors to test associations between THg and infection with hemoplasmas and *Bartonella* spp. We then used causal mediation analyses (CMA) to test support for directional relationships among THg, immunity, and infection (Imai, Keele, & Tingley, 2010). Similar to structural equation modeling, CMA decomposes the hypothesized causal relationship between a predictor (i.e.,

THg) and a response (i.e., infection) into the direct effect and the indirect effect mediated

through a third variable (i.e., immunity; Fig. S1). CMA then estimates the proportion of the total

effect mediated through the indirect effect using a mediator model, which was each of the

GLMMs predicting leukocyte counts by THg, and an outcome model. Here, we fit two GLMMs

per pathogen that modeled infection as a function of THg and each WBC count. We then used

the *brms* and *sjstats* packages to estimate these direct and indirect effects and in turn the

proportion of the total relationship (THg and infection) mediated by the indirect relationship

between THg and immunity and immunity and infection .

Because bat ecology likely dictates exposure to contaminants and pathogens, we assessed

species-specific patterns in Hg–infection relationships. We fit logistic regressions for each bat

species and pathogen when sample sizes were greater than two individuals. As these small

samples can bias odds ratio estimates, we used the *logistf* package to implement Firth's bias

reduction (Heinze & Schemper, 2002). For species with no variance in infection, we assigned log

odds of zero. Across these species, we next estimated phylogenetic signal (Pagel's λ) in the log

odds using the *caper* package and used phylogenetic generalized least squares to test if log odds

covaried with sample size (Orme, 2013). We then used phylogenetic factorization to identify

clades with different log odds at various taxonomic depths. We used the *taxize* and *phylofactor*

packages to obtain taxonomic information for each species and partition log odds as a Gaussian

response in a GLM (Chamberlain & Szöcs, 2013; Washburne et al., 2019). We included sample

size as a weighting variable and used Holm's sequentially rejective test to determine the number

of significant clades.

Results

Neotropical bat fur THg

Expanding our initial studies of fur THg across this Neotropical bat community in 2014, we

found that THg varied up to five orders of magnitude across species, sites, and years (Fig. 1).

Our top GLMM included interactions between diet and year and between year and site (Table 1).

Across years, the proportion of potentially aquatic prey in diet positively predicted THg but more

so in 2014 (β=0.019, 95% HDI: 0.011–0.026) than in 2017 (β=0.016, 95% HDI: 0.008–0.023;

Fig. 2A). We also identified strong spatiotemporal variation, such that fur THg increased

between 2014 and 2017 for KK ($\tilde{x}$=0.46 to 0.66) but slightly decreased for LAR ($\tilde{x}$=0.52 to

0.49).

Associations with immunity and infection

Across bats sampled in 2017 with Hg and WBC data (*n*=52), our GLMMs showed that higher

THg was associated with fewer neutrophils (β=–0.53, 95% HDI:–0.94 to –0.17) but not

lymphocytes (β=–0.37, 95% HDI: –0.85 to 0.02; Fig. 2B) after adjusting for sex and condition;

these two covariates had weak associations with leukocytes (Table S1). However, higher THg

was also associated with lower odds of infection for hemoplasmas (OR=0.19, 95% HDI: 0.04–

0.79, *n*=134) and *Bartonella* spp. (OR=0.18, 95% HDI: 0.03–0.84, *n*=117; Fig. 3A).

Accordingly, CMAs found that the relationship between THg and either WBC did not mediate

any of the relationships between THg and infection for both pathogens (Table S2).

When we used logistic regression to analyze individual relationships between THg and

infection per bat species and pathogen, we only found significant protective effects for

Pteronotus mesoamericanus and hemoplasmas after adjusting for multiple comparisons

(ln(OR)=–10.20, *p*=0.001; Table 2). We also detected a strong negative association between THg

and *Bartonella* in this species (ln(OR)=–3.34), but no log odds were significantly different from

zero after adjustment. Most species instead had null relationships between THg and infection

(i.e., 36% for hemoplasmas and 43% for *Bartonella* spp.) or relatively weaker negative effects

(e.g., *Dermanura phaeotis*, ln(OR)=–7.58 for hemoplasmas; *P. fulvus*, ln(OR)=–3.35 for

Bartonella spp.). These negative THg–infection associations were identified for 29% of species

for both pathogens. We also estimated similar proportions of positive THg associations for

hemoplasmas (36%) and *Bartonella* spp. (29%). *Desmodus rotundus* had the strongest positive

THg effect size for hemoplasma infection (ln(OR)=2.40), whereas *Dermanura watsoni* had the

largest positive THg effect size for *Bartonella* spp. infection (ln(OR)=8.9).

Comparative analyses of the log odds ratios across bat species revealed no phylogenetic

signal for the relationship between THg and hemoplasmas (λ=0) but strong phylogenetic signal

for the relationship between THg and *Bartonella* spp. (λ=0.84). Log odds ratios were not

associated with sample size for hemoplasmas ($F_{1,12}$=1.46, *p*=0.25) or *Bartonella* spp. ($F_{1,12}$=0.25,

p=0.63). Phylogenetic factorization further identified species-specific or taxonomic patterns in

the magnitude and direction of effect size. For hemoplasmas, the odds of infection were

significantly lower for *Pteronotus mesoamericanus* and *Dermanura phaeotis* when compared to

the remaining sampled bat phylogeny. For *Bartonella* spp., however, the odds of infection were

significantly lower for the genus *Pteronotus* (mean ln(OR)=–3.34) and significantly higher for

the subfamily Stenodermatinae (*Dermanura* spp. and *Sturnira parvidens*, mean ln(OR)=5.79;

Fig. 3B). Post-hoc GLMs showed that bats in the Stenodermatinae had especially negative

associations between THg and neutrophils (β=–2.14, *p*=0.001) but not lymphocytes (β=–0.43,

p=0.63), although these analyses were limited by small sample size (*n*=7).

Discussion

Interactions among contaminants, immunity, and infection are difficult to disentangle in natural

systems, but quantifying their proposed causal relationships can inform how land-use change

affects wildlife health and human disease risks. By capitalizing on a diverse Neotropical bat

system with high variation in Hg bioaccumulation and bacterial pathogens, we found higher THg

was associated with fewer neutrophils but also lower odds of infection across the host

community. However, our species-specific and taxonomic analyses showed THg had protective

effects for hemoplasmas and *Bartonella* spp. in the genus *Pteronotus*, whereas THg was

associated with fewer neutrophils and elevated infection in the subfamily Stenodermatinae.

These contrasting relationships suggest contaminant-driven loss of pathogen habitat (i.e.,

anemia) or vector mortality versus immunosuppression as possible causal mechanisms,

respectively, and identify clades of bats that may be especially resilient or vulnerable to infection

risks from Hg exposure. Such findings more generally suggest contaminants may increase

infection risk in some taxa but not others, emphasizing the importance of considering

surveillance and management at different phylogenetic scales (Graham, Storch, & Machac,

2018).

Expanding our prior analyses of this Neotropical bat community and global patterns of

THg in bat fur (Becker, Chumchal, et al., 2018) with larger within-species sample sizes, we first

demonstrated that Hg exposure increases with potentially aquatic prey (or prey with some life

stages linked to aquatic ecosystems) in diet despite spatial and temporal variation in THg.

Positive associations with diets linked to aquatic ecosystems across sites and years provides

additional support for trophic transfer of Hg through foraging (Cristol et al., 2008; Ortega-

Rodriguez et al., 2019; Speir et al., 2014). This diet-mediated connectivity to aquatic ecosystems

likely underlies other cases of guild-specific Hg bioaccumulation in bat communities (Carrasco-

Rueda, Loiselle, & Frederick, 2020; Korstian, Chumchal, Bennett, & Hale, 2018). In many of

these regions, bat dietary exposure to Hg is driven by land-use changes such as gold mining and

agriculture (Carrasco-Rueda et al., 2020; Costantini et al., 2019), whereas atmospheric

deposition is often the primary source of Hg in regions located further from anthropogenic point

sources (Chételat et al., 2018; Korstian et al., 2018). The latter is a likely source of Hg in the

Belize system (Becker, Chumchal, et al., 2018). However, intensified agriculture and especially

slash-and-burn practices could provide other Hg inputs and may explain why bat THg increased

between 2014 and 2017 in KK, the more agricultural site, but not in the protected LAR (Farella

et al., 2007; Patterson, 2016).

Across bat species, fur THg was negatively correlated with neutrophil counts, which

mirrors captive results and suggests impaired innate immunity (Lalancette, Morin, Measures, &

Fournier, 2003). Previously, vampire bats sampled in Belize with high fur THg had weaker

innate defense (i.e., bacterial killing ability; Becker et al., 2017). Wrinkle-lipped free-tailed bats

with higher Hg exposure also had weaker innate immunity (i.e., bacterial killing ability,

lysozyme and haptoglobin concentrations; Costantini et al., 2019). Most individual bats for

which we had both Hg and immunity data showed THg below toxicity and subclinical thresholds

of 5–10 mg/kg (Nam et al., 2012), which suggests innate immunity could be weakened at

sublethal contaminant concentrations (Lewis, Cristol, Swaddle, Varian-Ramos, & Zwollo, 2013).

Alternatively, sublethal effects of THg could combine with other stressors (e.g., reproduction,

food and roost availability) to impair immunity. Additional immune measures across broader

land-use gradients would help characterize functional differences in defense in relation to THg

concentrations in anthropogenic habitats (Becker, Albery, et al., 2020; Costantini et al., 2019).

Although neutrophils were lowest in bats with high Hg exposure, the odds of infection

with hemoplasmas and *Bartonella* spp. decreased with fur THg across the bat community. As in

other mammals, bats challenged with bacteria produce neutrophils as part of the innate immune

response (Weise, Czirják, Lindecke, Bumrungsri, & Voigt, 2017). Elevated neutrophil counts

could be protective, as implied through negative associations between innate immunity and these

two pathogens in vampire bats (Becker, Czirják, et al., 2018). Accordingly, relationships

between THg and immunity did not mediate the relationships between THg and infection, which

likely indicates immunosuppression is not a causal mechanism. One alternative may involve

contaminant-mediated pathogen mortality; for example, lead exposure likely caused helminth

mortality in avian hosts (Prüter et al., 2018). As facultative intracellular pathogens, hemoplasmas

and *Bartonella* spp. both infect red blood cells. Hg can lower erythrocyte counts (Shaw, Dash, &

Panigrahi, 1991), which could reduce resources available to both bacteria. We did not measure

anemia or Hg in blood; however, fur THg strongly correlates with blood THg, despite the former

being orders of magnitude higher than the latter (Wada, Yates, Evers, Taylor, & Hopkins, 2010).

As *Bartonella* spp. and possibly hemoplasmas likely depend in part on vector-borne transmission

(Becker, Bergner, et al., 2018; Willi et al., 2007), ectoparasite mortality from contaminants in

hosts or the environment could also explain observed infection patterns (e.g., as found for some

avian ectoparasites; Eeva & Klemola, 2013). Lastly, we did not reliably age all bats, precluding

age from our analyses; however older animals typically have higher fur THg (Korstian et al.,

2018). If older bats also have stronger adaptive immunity (e.g., antibody titers in *Saccopteryx*

bilineata; Schneeberger, Courtiol, Czirják, & Voigt, 2014), age could explain negative

associations between THg and infection.

Our species-specific analyses suggested contaminants may have variable impacts on

infection risk depending on taxon. In particular, we identified possibly protective effects of THg

on infections primarily in the genus *Pteronotus*, with strong negative effects for *P.*

mesoamericanus for both pathogens. Both species in this genus had high THg, being insectivores

that often eat dipterans and coleopterans (Salinas-Ramos, Herrera Montalvo, León-Regagnon,

Arrizabalaga-Escudero, & Clare, 2015). Given this clade-based signal, *Pteronotus* bats may be

particularly resilient to disease impacts of Hg. Such effects could be in part mediated by high

host-specificity of bat flies for this genus (Ter Hofstede, Fenton, & Whitaker, 2004), particularly

if high THg increases vector mortality or if aspects of bat fly biology impact vector competence

(Weiss & Aksoy, 2011). In contrast, we also identified taxa with positive relationships between

THg and infection. For hemoplasmas, higher THg was associated with greater risk in vampire

bats. As this species frequently feeds on humans and livestock (Streicker & Allgeier, 2016),

pathogen surveillance would be particularly important in habitats contaminated by agriculture or

mining. Additionally, the Stenodermatinae displayed strong positive associations between THg

and *Bartonella* spp. These frugivores also showed negative correlations between THg and

neutrophils, suggesting Hg-mediated immunosuppression. Such species may thus be especially

vulnerable to infection from Hg exposure and could play key roles in maintaining *Bartonella*

spp. infection cycles between bats and ectoparasites in contaminated environments (Judson,

Frank, & Hadly, 2015). From another perspective, because the Stenodermatinae and *Pteronotus*

had lower and higher fur THg, respectively, low Hg exposure could increase susceptibility while

higher concentrations instead cause anemia and facilitate protective effects against erythrocytic

pathogens. More generally, these results highlight the importance of considering surveillance and

management of Hg exposure (e.g., through possible land use drivers) at different phylogenetic

scales, such as species, genus, or subfamily.

Beyond bats and their pathogens, our study more broadly emphasizes the need to assess

potential causal relationships between contaminants and infectious diseases in natural systems. In

particular, we provide a novel perspective on integrating approaches from ecotoxicology with

those of disease ecology to disentangle the complex relationships among contaminants,

immunity, and infection. Future work that carefully integrates data on contaminant exposure,

specific diet composition, multiple immune measures, and pathogen diversity (e.g., with

metagenomics; Bergner et al., 2019) across site gradients of anthropogenic intensity could help

identify habitats, host clades, and infections for which disease risks are highest.

Acknowledgements

For assistance with field logistics and permits, we thank Mark Howells, Neil Duncan, and staff

of the Lamanai Field Research Center. We also thank the many colleagues who helped net bats

during 2014 and 2017 bat research in Belize as well as Susan Perkins for laboratory reagents.

Lastly, we thank two anonymous reviewers for constructive feedback on this manuscript.

Data accessibility

Data from 2017 will be deposited in Dryad Digital Repository upon acceptance. Data from 2014

have been published previously (Becker, Chumchal, et al., 2018).

Author contributions

DJB, KAS, AMB, HGB, and NBS collected samples; JMK, DJB, KAS, DVV, AMB, CLB, HD,

and TPS analyzed samples; DJB analyzed the data; and HGB, RKP, TRR, MBF, NBS, and

MMC provided funding and logistical support. DJB wrote the manuscript with input from all

authors.

Funding

DJB was funded by the ARCS Foundation and Explorer's Club. Field work by DJB and KAS was supported by grants from the American Museum of Natural History Theodore Roosevelt Memorial Fund. Lab work by KAS and CLB was funded by the Richard Gilder Graduate School Student Research Fellowship. RKP was supported by NSF DEB-1716698, the Defense Advanced Research Projects Agency PREEMPT program Cooperative Agreement D18AC00031, U.S. National Institutes of General Medical Sciences IDeA Program (P20GM103474 and P30GM110732), and the USDA National Institute of Food and Agriculture (Hatch project 1015891). NBS was supported by the American Museum of Natural History Taxonomic Mammalogy Fund. JMK and MMC were supported by a Texas Christian University Research and Creative Activities Fund Award, and HGB was supported by a Natural Sciences and Engineering Research Council of Canada Discovery Grant. TRR was supported by the Yawkey Foundation and Clemson University. The views, opinions, and/or findings expressed are those of the authors and should not be interpreted as representing the official views or policies of the Department of Defense or the U.S. Government.

Tables

Table 1. Competing phylogenetic GLMMs predicting log fur THg concentrations across the Belize bat community (n=249). Models are ranked by ΔLOOIC with LOOIC SE, Akaike weights (w_i) and Bayesian R^2 estimates. All models include random effects for species and phylogeny.

Fixed effects	LOOIC	SE	ΔLOOIC	w_i	R^2_m	R^2_c
~ diet + year + site + year:site + diet:year	76.21	38.83	0.00	0.39	0.80	0.92
~ diet + year + site + year:site + diet:year + diet:site	77.22	38.12	1.01	0.24	0.79	0.92
~ diet * year * site	77.64	37.03	1.43	0.19	0.79	0.92
~ diet * year + site	78.78	39.63	2.57	0.11	0.80	0.92
~ diet + year + site + diet:site + diet:year	79.57	38.96	3.36	0.07	0.79	0.92
~ diet + year + site + diet:site + year:site	86.87	33.20	10.66	>0.01	0.77	0.91
~ diet + year * site	89.33	33.93	13.13	>0.01	0.78	0.91
~ diet *site + year	89.97	33.94	13.77	>0.01	0.77	0.91
~ diet + year + site	94.43	34.22	18.23	>0.01	0.78	0.91

430   Table 2. Results of species-specific logistic regressions (using Firth's bias reduction method)

431   between bat fur THg concentrations and infection status per each bacterial pathogen.

Bat species	Hemoplasmas				Bartonella			
	ln(OR)	p	σ^2	n	ln(OR)	p	σ^2	n
Eptesicus furinalis	-1.545	0.4569	0.214	8	0	1	0	7
Bauerus dubiaquercus	0	1	0	3	0	1	0	3
Dermanura watsoni	-0.128	0.984	0.267	6	8.904	0.3177	0.167	6
Dermanura phaeotis	-7.578	0.2562	0.278	9	4.27	0.5789	0.214	8
Sturnira parvidens	1.099	0.7923	0.257	15	4.198	0.4613	0.194	9
Carollia sowelli	0.213	0.9427	0.178	10	-1.987	0.5159	0.233	10
Trachops cirrhosus	0	1	0	3	0	1	0	3
Chrotopterus auritus	0	1	0	3	0	1	0	3
Glossophaga soricina	0.665	0.8421	0.218	11	-2.373	0.4911	0.194	9
Desmodus rotundus	2.395	0.3328	0.208	22	0.178	0.933	0.208	22
Pteronotus mesoamericanus	-10.201	0.0001	0.227	22	-3.339	0.0159	0.25	16
Pteronotus fulvus	1.653	0.65	0.25	4	-3.346	0.4073	0.25	4
Rhynchonycteris naso	0	1	0	4	0	1	0	3
Saccopteryx bilineata	0	1	0	8	0	1	0	8

Figures and legends

 Figure 1. THg concentrations in bat fur across Neotropical bat species sampled in 2014 and 2017

 across two sites in Belize. Boxplots are colored by the proportion of potentially aquatic prey in

 bat diets from the EltonTraits database. THg concentrations are displayed on a $\log_{10}$ scale.

440   Figure 2. Dietary drivers of THg concentrations in Neotropical bat fur (A) and cellular immunity

441   correlates of THg (B). Points indicate individual bats and are colored by sampling year in Belize.

442   Lines display 100 random draws from the posterior distribution of the main phylogenetic

443   GLMMs. THg concentrations and absolute leukocyte counts are displayed on a log$_{10}$ scale.

References

Bai, Y., Gilbert, A., Fox, K., Osikowicz, L., & Kosoy, M. (2016). Bartonella rochalimae and b.

Vinsonii subsp. Berkhoffii in wild carnivores from colorado, usa. *Journal of Wildlife*

Diseases, *52*(4), 844–849. doi: 10.7589/2016-01-015

Becker, D. J., Albery, G. F., Kessler, M. K., Lunn, T. J., Falvo, C. A., Czirják, G. Á., …

Plowright, R. K. (2020). Macroimmunology: The drivers and consequences of spatial

patterns in wildlife immune defence. *Journal of Animal Ecology*, *89*(4), 972–995. doi:

10.1111/1365-2656.13166

Becker, D. J., Bergner, L. M., Bentz, A. B., Orton, R. J., Altizer, S., & Streicker, D. G. (2018).

Genetic diversity, infection prevalence, and possible transmission routes of Bartonella

spp. In vampire bats. *PLOS Neglected Tropical Diseases*, *12*(9), e0006786. doi:

10.1371/journal.pntd.0006786

Becker, D. J., Chumchal, M. M., Bentz, A. B., Platt, S. G., Czirják, G. Á., Rainwater, T. R., …

Streicker, D. G. (2017). Predictors and immunological correlates of sublethal mercury

exposure in vampire bats. *Royal Society Open Science*, *4*(4), 170073. doi:

10.1098/rsos.170073

Becker, D. J., Chumchal, M. M., Broders, H. G., Korstian, J. M., Clare, E. L., Rainwater, T. R.,

… Fenton, M. B. (2018). Mercury bioaccumulation in bats reflects dietary connectivity to

aquatic food webs. *Environmental Pollution*, *233*, 1076–1085. doi:

10.1016/j.envpol.2017.10.010

Becker, D. J., Czirják, G. Á., Volokhov, D. V., Bentz, A. B., Carrera, J. E., Camus, M. S., …

Streicker, D. G. (2018). Livestock abundance predicts vampire bat demography, immune

profiles and bacterial infection risk. *Phil. Trans. R. Soc. B*, *373*(1745), 20170089. doi:

10.1098/rstb.2017.0089

Becker, D. J., Speer, K. A., Brown, A. M., Fenton, M. B., Washburne, A. D., Altizer, S., …

Volokhov, D. V. (2020). Ecological and evolutionary drivers of haemoplasma infection

and bacterial genotype sharing in a Neotropical bat community. *Molecular Ecology*,

29(8), 1534–1549. doi: 10.1111/mec.15422

Bergner, L. M., Orton, R. J., Filipe, A. da S., Shaw, A. E., Becker, D. J., Tello, C., … Streicker,

D. G. (2019). Using noninvasive metagenomics to characterize viral communities from

wildlife. *Molecular Ecology Resources*, *19*(1), 128–143. doi: 10.1111/1755-0998.12946

Borremans, B., Faust, C., Manlove, K. R., Sokolow, S. H., & Lloyd-Smith, J. O. (2019). Cross-

species pathogen spillover across ecosystem boundaries: Mechanisms and theory.

Philosophical Transactions of the Royal Society B, *374*(1782), 20180344.

Brook, C. E., & Dobson, A. P. (2015). Bats as 'special'reservoirs for emerging zoonotic

pathogens. *Trends in Microbiology*, *23*(3), 172–180.

Bürkner, P.-C. (2017). brms: An R package for Bayesian multilevel models using Stan. *Journal

of Statistical Software*, *80*(1), 1–28.

Carrasco-Rueda, F., Loiselle, B. A., & Frederick, P. C. (2020). Mercury bioaccumulation in

tropical bats from a region of active artisanal and small-scale gold mining. *Ecotoxicology*,

1–11.

Chamberlain, S. A., & Szöcs, E. (2013). taxize: Taxonomic search and retrieval in R.

F1000Research, *2*.

Chételat, J., Hickey, M. B. C., Poulain, A. J., Dastoor, A., Ryjkov, A., McAlpine, D., …

Zanuttig, M. (2018). Spatial variation of mercury bioaccumulation in bats of Canada

linked to atmospheric mercury deposition. *Science of The Total Environment*, *626*, 668–

677. doi: 10.1016/j.scitotenv.2018.01.044

Chumchal, M. M., Rainwater, T. R., Osborn, S. C., Roberts, A. P., Abel, M. T., Cobb, G. P., …

Bailey, F. C. (2011). Mercury speciation and biomagnification in the food web of Caddo

Lake, Texas and Louisiana, USA, a subtropical freshwater ecosystem. *Environmental*

Toxicology and Chemistry, *30*(5), 1153–1162.

Costantini, D., Czirják, G. Á., Bustamante, P., Bumrungsri, S., & Voigt, C. C. (2019). Impacts of

land use on an insectivorous tropical bat: The importance of mercury, physio-

immunology and trophic position. *Science of The Total Environment*, *671*, 1077–1085.

doi: 10.1016/j.scitotenv.2019.03.398

Cristol, D. A., Brasso, R. L., Condon, A. M., Fovargue, R. E., Friedman, S. L., Hallinger, K. K.,

… White, A. E. (2008). The Movement of Aquatic Mercury Through Terrestrial Food

Webs. *Science*, *320*(5874), 335–335. doi: 10.1126/science.1154082

Davidson, C. (2004). Declining downwind: Amphibian population declines in California and

historical pesticide use. *Ecological Applications*, *14*(6), 1892–1902.

Eeva, T., & Klemola, T. (2013). Variation in prevalence and intensity of two avian ectoparasites

in a polluted area. *Parasitology*, *140*(11), 1384–1393.

Farella, N., Davidson, R., Lucotte, M., & Daigle, S. (2007). Nutrient and mercury variations in

soils from family farms of the Tapajós region (Brazilian Amazon): Recommendations for

better farming. *Agriculture, Ecosystems & Environment*, *120*(2–4), 449–462.

Fisk, A. T., de Wit, C. A., Wayland, M., Kuzyk, Z. Z., Burgess, N., Letcher, R., … Muir, D. C.

G. (2005). An assessment of the toxicological significance of anthropogenic

contaminants in Canadian arctic wildlife. *Science of The Total Environment*, *351–352*,

57–93. doi: 10.1016/j.scitotenv.2005.01.051

Flache, L., Becker, N. I., Kierdorf, U., Czarnecki, S., Düring, R.-A., & Encarnação, J. A. (2015).

Hair samples as monitoring units for assessing metal exposure of bats: A new tool for

risk assessment. *Mammalian Biology-Zeitschrift Für Säugetierkunde*, *80*(3), 178–181.

Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian Regression

Models. *The American Statistician*, *73*(3), 307–309. doi:

10.1080/00031305.2018.1549100

Graham, C. H., Storch, D., & Machac, A. (2018). Phylogenetic scale in ecology and evolution.

Global Ecology and Biogeography, *27*(2), 175–187. doi: 10.1111/geb.12686

Grasman, K. A. (2002). Assessing immunological function in toxicological studies of avian

wildlife. *Integrative and Comparative Biology*, *42*(1), 34–42.

Gunnell, G. F., & Simmons, N. B. (2012). *Evolutionary history of bats: Fossils, molecules and*

morphology. Cambridge University Press.

Heinze, G., & Schemper, M. (2002). A solution to the problem of separation in logistic

regression. *Statistics in Medicine*, *21*(16), 2409–2419.

Herrera, J. P., Duncan, N., Clare, E., Fenton, M. B., & Simmons, N. (2018). Disassembly of

Fragmented Bat Communities in Orange Walk District, Belize. *Acta Chiropterologica*,

20(1), 147–159. doi: 10.3161/15081109ACC2018.20.1.011

Ikeda, P., Seki, M. C., Carrasco, A. O. T., Rudiak, L. V., Miranda, J. M. D., Gonçalves, S. M.

M., … others. (2017). Evidence and molecular characterization of Bartonella spp. And

hemoplasmas in neotropical bats in Brazil. *Epidemiology & Infection*, 1–15.

Imai, K., Keele, L., & Tingley, D. (2010). A general approach to causal mediation analysis.

Psychological Methods, *15*(4), 309–334. doi: 10.1037/a0020761

Judson, S. D., Frank, H. K., & Hadly, E. A. (2015). Bartonellae are Prevalent and Diverse in

Costa Rican Bats and Bat Flies. *Zoonoses and Public Health*, *62*(8), 609–617. doi:

550   10.1111/zph.12188

551 Korstian, J. M., Chumchal, M. M., Bennett, V. J., & Hale, A. M. (2018). Mercury contamination

552   in bats from the central United States. *Environmental Toxicology and Chemistry*, *37*(1),

553   160–165.

554 Lalancette, A., Morin, Y., Measures, L., & Fournier, M. (2003). Contrasting changes of

555   sensitivity by lymphocytes and neutrophils to mercury in developing grey seals.

556   *Developmental & Comparative Immunology*, *27*(8), 735–747.

557 Lanier, L. L. (2013). Shades of grey—The blurring view of innate and adaptive immunity.

558   *Nature Reviews Immunology*, *13*(2), 73.

559 Lewis, C. A., Cristol, D. A., Swaddle, J. P., Varian-Ramos, C. W., & Zwollo, P. (2013).

560   Decreased immune response in zebra finches exposed to sublethal doses of mercury.

561   *Archives of Environmental Contamination and Toxicology*, *64*(2), 327–336.

562 Michonneau, F., Brown, J. W., & Winter, D. J. (2016). rotl: An R package to interact with the

563   Open Tree of Life data. *Methods in Ecology and Evolution*. doi: 10.1111/2041-

564   210X.12593

565 Nam, D.-H., Yates, D., Ardapple, P., Evers, D. C., Schmerfeld, J., & Basu, N. (2012). Elevated

566   mercury exposure and neurochemical alterations in little brown bats (Myotis lucifugus)

567   from a site with historical mercury contamination. *Ecotoxicology*, *21*(4), 1094–1101. doi:

568   10.1007/s10646-012-0864-9

569 Orme, D. (2013). The caper package: Comparative analysis of phylogenetics and evolution in R.

570   *R Package Version*, *5*(2). Retrieved from

571   ftp://ctan.mirrorcatalogs.com/cran/web/packages/caper/vignettes/caper.pdf

572 Ortega-Rodriguez, C. L., Chumchal, M. M., Drenner, R. W., Kennedy, J. H., Nowlin, W. H.,

Barst, B. D., … Lauck, K. C. (2019). Relationship Between Methylmercury

Contamination and Proportion of Aquatic and Terrestrial Prey in Diets of Shoreline

Spiders. *Environmental Toxicology and Chemistry*, *38*(11), 2503–2508.

Palheta, D., & Taylor, A. (1995). Mercury in environmental and biological samples from a gold

mining area in the Amazon region of Brazil. *Science of the Total Environment*, *168*(1),

63–69.

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of phylogenetics and evolution in

R language. *Bioinformatics*, *20*(2), 289–290.

Patterson, C. (2016). *Deforestation, agricultural intensification, and farm resilience in Northern*

Belize: 1980-2010 (Thesis, University of Otago). University of Otago. Retrieved from

https://ourarchive.otago.ac.nz/handle/10523/6858

Prüter, H., Franz, M., Auls, S., Czirják, G. Á., Greben, O., Greenwood, A. D., … Krone, O.

(2018). Chronic lead intoxication decreases intestinal helminth species richness and

infection intensity in mallards (Anas platyrhynchos). *Science of the Total Environment*,

644, 151–160.

Rainwater, T. R., Reynolds, K. D., Cañas, J. E., Cobb, G. P., Andersonv, T. A., McMurry, S. T.,

& Smith, P. N. (2005). Organochlorine pesticides and mercury in cottonmouths

(Agkistrodon piscivorus) from northeastern Texas, USA. *Environmental Toxicology and*

Chemistry, *24*(3), 665–673. doi: 10.1897/04-223R.1

Reid, F. (1997). *A Field Guide to the Mammals of Central America and Southeast Mexico*. OUP

USA.

Rojas, D., Vale, Á., Ferrero, V., & Navarro, L. (2011). When did plants become important to

leaf-nosed bats? Diversification of feeding habits in the family Phyllostomidae.

Molecular Ecology, *20*(10), 2217–2228. doi: 10.1111/j.1365-294X.2011.05082.x

Ross, P. S. (2002). The role of immunotoxic environmental contaminants in facilitating the

emergence of infectious diseases in marine mammals. *Human and Ecological Risk*

Assessment: An International Journal, *8*(2), 277–292.

Salinas-Ramos, V. B., Herrera Montalvo, L. G., León-Regagnon, V., Arrizabalaga-Escudero, A.,

& Clare, E. L. (2015). Dietary overlap and seasonality in three species of mormoopid bats

from a tropical dry forest. *Molecular Ecology*, *24*(20), 5296–5307.

Scheuhammer, A. M., Meyer, M. W., Sandheinrich, M. B., & Murray, M. W. (2007). Effects of

environmental methylmercury on the health of wild birds, mammals, and fish. *AMBIO: A*

Journal of the Human Environment, *36*(1), 12–19.

Schneeberger, K., Courtiol, A., Czirják, G. Á., & Voigt, C. C. (2014). Immune Profile Predicts

Survival and Reflects Senescence in a Small, Long-Lived Mammal, the Greater Sac-

Winged Bat (Saccopteryx bilineata). *PLoS ONE*, *9*(9), e108268. doi:

10.1371/journal.pone.0108268

Schneeberger, K., Czirják, G. Á., & Voigt, C. C. (2013). Measures of the Constitutive Immune

System Are Linked to Diet and Roosting Habits of Neotropical Bats. *PLoS ONE*, *8*(1),

e54023. doi: 10.1371/journal.pone.0054023

Serieys, L. E. K., Lea, A. J., Epeldegui, M., Armenta, T. C., Moriarty, J., VandeWoude, S., …

Uittenbogaart, C. H. (2018). Urbanization and anticoagulant poisons promote immune

dysfunction in bobcats. *Proc. R. Soc. B*, *285*(1871), 20172533. doi:

10.1098/rspb.2017.2533

Shaw, B. P., Dash, S., & Panigrahi, A. K. (1991). Effect of methyl mercuric chloride treatment

on haematological characteristics and erythrocyte morphology of Swiss mice.

Environmental Pollution, *73*(1), 43–52.

Sikes, R. S., Care, A., & Mammalogists, U. C. of the A. S. of. (2016). 2016 Guidelines of the

American Society of Mammalogists for the use of wild mammals in research and

education. *Journal of Mammalogy*, *97*(3), 663–688.

Smith, P. N., Cobb, G. P., Godard-Codding, C., Hoff, D., McMurry, S. T., Rainwater, T. R., &

Reynolds, K. D. (2007). Contaminant exposure in terrestrial vertebrates. *Environmental*

Pollution, *150*(1), 41–64. doi: 10.1016/j.envpol.2007.06.009

Speir, S. L., Chumchal, M. M., Drenner, R. W., Cocke, W. G., Lewis, M. E., & Whitt, H. J.

(2014). Methyl mercury and stable isotopes of nitrogen reveal that a terrestrial spider has

a diet of emergent aquatic insects. *Environmental Toxicology and Chemistry*, *33*(11),

2506–2509. doi: 10.1002/etc.2700

Streicker, D. G., & Allgeier, J. E. (2016). Foraging choices of vampire bats in diverse

landscapes: Potential implications for land-use change and disease transmission. *Journal*

of Applied Ecology, *53*(4), 1280–1288. doi: 10.1111/1365-2664.12690

Ter Hofstede, H. M., Fenton, M. B., & Whitaker, J. O. (2004). Host and host-site specificity of

bat flies (Diptera: Streblidae and Nycteribiidae) on Neotropical bats (Chiroptera).

Canadian Journal of Zoology, *82*(4), 616–626.

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-

one-out cross-validation and WAIC. *Statistics and Computing*, *27*(5), 1413–1432.

Volokhov, D. V., Becker, D. J., Bergner, L. M., Camus, M. S., Orton, R. J., Chizhikov, V. E., …

Streicker, D. G. (2017). Novel hemotropic mycoplasmas are widespread and genetically

diverse in vampire bats. *Epidemiology & Infection*, *145*(15), 3154–3167.

Wada, H., Yates, D. E., Evers, D. C., Taylor, R. J., & Hopkins, W. A. (2010). Tissue mercury

concentrations and adrenocortical responses of female big brown bats (Eptesicus fuscus)

near a contaminated river. *Ecotoxicology*, *19*(7), 1277–1284. doi: 10.1007/s10646-010-

0513-0

Washburne, A. D., Silverman, J. D., Morton, J. T., Becker, D. J., Crowley, D., Mukherjee, S., …

Plowright, R. K. (2019). Phylofactorization: A graph partitioning algorithm to identify

phylogenetic scales of ecological data. *Ecological Monographs*, *0*(0), e01353. doi:

10.1002/ecm.1353

Weise, P., Czirják, G. A., Lindecke, O., Bumrungsri, S., & Voigt, C. C. (2017). Simulated

bacterial infection disrupts the circadian fluctuation of immune cells in wrinkle-lipped

bats (Chaerephon plicatus). *PeerJ*, *5*, e3570. doi: 10.7717/peerj.3570

Weiss, B., & Aksoy, S. (2011). Microbiome influences on insect host vector competence. *Trends*

in Parasitology, *27*(11), 514–522.

Willi, B., Boretti, F. S., Tasker, S., Meli, M. L., Wengi, N., Reusch, C. E., … Hofmann-

Lehmann, R. (2007). From Haemobartonella to hemoplasma: Molecular methods provide

new insights. *Veterinary Microbiology*, *125*(3–4), 197–209. doi:

10.1016/j.vetmic.2007.06.027

Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., & Jetz, W. (2014).

EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals.

Ecology, *95*(7), 2027–2027. doi: 10.1890/13-1917.1

Yates, D. E., Adams, E. M., Angelo, S. E., Evers, D. C., Schmerfeld, J., Moore, M. S., …

O'Driscoll, N. J. (2013). Mercury in bats from the northeastern United States.

Ecotoxicology, *23*(1), 45–55. doi: 10.1007/s10646-013-1150-1