Ultra-rapid metagenotyping of the human gut microbiome

Zhou Jason Shi[1,2], Boris Dimitrov[3], Chunyu Zhao[1], Stephen Nayfach[4,5,*] and Katherine S. Pollard[1,2,6,*]

[1]Chan Zuckerberg Biohub, Data Science, San Francisco, CA, [2]Gladstone Institutes, San Francisco, CA,
[3]Chan Zuckerberg Initiative, CA, [4]Department of Energy, Joint Genome Institute, Walnut Creek, CA,
[5]Lawrence Berkeley National Laboratory, Environmental Genomics and Systems Biology Division,
Berkeley, CA, [6]University of California San Francisco

*e-mail: snayfach@lbl.gov; katherine.pollard@gladstone.ucsf.edu

Abstract

Sequence variation is used to quantify population structure and identify genetic determinants of phenotypes that vary within species. In the human microbiome and other environments, single nucleotide polymorphisms (SNPs) are frequently detected by aligning metagenomic sequencing reads to catalogs of genes or genomes. But this requires high-performance computing and enough read coverage to distinguish SNPs from sequencing errors. We solved these problems by developing the GenoTyper for Prokaytotes (GT-Pro), a suite of novel methods to catalog SNPs from genomes and use exact k-mer matches to perform ultra-fast reference-based SNP calling from metagenomes. Compared to read alignment, GT-Pro is more accurate and two orders of magnitude faster. We discovered 104 million SNPs in 909 human gut species, characterized their global population structure, and tracked pathogenic strains. GT-Pro democratizes strain-level microbiome analysis by making it possible to genotype hundreds of metagenomes on a personal computer.

Software availability: GT-Pro is available at https://github.com/zjshi/gt-pro.

Introduction

Microbial species harbor extensive genetic variation, including single nucleotide polymorphisms (SNPs), structural variants (SVs), and mobile genetic elements. SNPs in particular are useful for population genetic analyses[1], such as tracking transmission of strains between environments or locations, reconstructing strain phylogenetic relationships, resolving mixtures of genotypes within a host, and depicting population diversity or structure along environmental gradients. Additionally, SNPs can result in changes in protein function. For example, a single SNP in the Dadh gene of the human commensal *Eggerthella lenta* can predict activity of levodopa, the primary medication used to treat Parkinson's disease[2]. Quantifying intra-species genomic variation in the human microbiome is a prerequisite to the potential application of microbiome genomics to precision medicine.

Several approaches exist for identifying SNPs in microbiomes. The gold standard[3] is to sequence individual isolate genomes and identify mismatches in whole-genome alignments. In contrast, metagenomes are a rich source of strain level diversity for uncultivated taxa. In a landmark study, Schloissnig et al[4]. discovered 10.3 million SNPs for 101 human gut species by aligning short reads from shotgun metagenomes to reference genomes. This approach is known as "metagenotyping" and has since been featured in several tools, including Constrains[5], MIDAS[6], metaSNV[7], DESMAN[8] and StrainPhlAn[9]. While algorithms for read alignment have improved, the approach is still computationally costly. Exact matching algorithms such as Kraken[10], CLARK[11] and bfMEM[12], have been developed as a more efficient solution to the read mapping problem, achieving speedups by orders of magnitude. However, these tools have thus far been used to quantify the abundance of microbiome taxa, rather than identify intra-species genetic variation. Genotyping by exact matches between reads and short sequences covering SNPs was

51 implemented in the method LAVA[13] for human whole-genome sequencing data. Our goal was to extend
52 this approach to metagenomes by addressing the challenges presented by complex mixtures of species and
53 strains within microbiome samples, while also making software that could run on a personal computer.
54
55 **Results**
56
57 *A novel framework for* in silico *genotyping of microbiome species*
58
We introduce the GenoTyper for PROkaryotes (GT-Pro), which is a novel computational pipeline that
utilizes an exact matching algorithm to perform ultra-rapid and accurate genotyping of known SNPs from
metagenomes. Our proof-of-principle initial implementation of this approach focuses on the human gut
microbiome. We created a reference database of 104 million common SNPs that we identified using
112,904 high-quality genomes from 909 human gut microbiome species. Then we used this catalog to
perform reference-based SNP calling for 25,133 publicly available metagenomes, providing insight into
strain variation across individuals and geographic regions. Our results demonstrate the feasibility of
performing large-scale metagenotyping without need for high-performance computing.
67
To overcome the low throughput, sensitivity and species coverage of current alignment-based
metagenotyping methods, we developed the GT-Pro framework (Fig. 1). Our key innovations are (i)
capturing the majority of common variation found in microbiome genomes with a compact database of
SNP covering k-mers (sck-mers), (ii) selecting highly species-specific sck-mers, overcoming high false
positives associated with k-mer exact-matching methods, and (iii) developing and optimizing algorithms
and data structures for exact matching of metagenomic sequencing reads to these sck-mers, enabling
SNPs to be detected rapidly and accurately in microbiome samples. Building a version of GT-Pro for a
given environment involves 1) discovering common SNPs in assembled genomes for each species, 2)
optionally identifying linkage disequilibrium (LD) blocks and "tag" SNPs that capture most variation
within each block, and 3) designing species-specific sck-mers. We focus on common SNPs) because this
allows us to create a virtual genotyping "array" that is a data structure small enough to fit in computer
memory while stll capturing the majority of prevalent genetic variation for each of many species.
80
81 *A database of common SNPs for bacterial species in the gut microbiome*
82
As a case study, we applied GT-Pro to the human gut microbiome due to the large number of microbial
genomes from this environment and its important role in human health. To construct a SNP catalog, we
used 112,904 high-quality genomes (>= 90% completeness and <= 5% contamination[14]) from 909 species
(minimum = 10 genomes, median = 35 genomes) that we downloaded from the Unified Gastrointestinal
Genomes (UHGG) resource[15] (Fig. S1, S2 and Table S1). These include both metagenome-assembled
genomes[16–18] (i.e. MAGs, 94.1%) as well as cultivated isolates (5.9%) and were derived from
geographically and phenotypically diverse human subjects. We performed whole-genome alignments for
each species, revealing 104,171,172 common, core-genome SNPs (minor allele frequency >= 1%, site
prevalence >= 90%), the vast majority of which (93.4%) were bi-allelic (Fig. 2a, S3a and S4). An
extremely low fraction of SNPs (<0.2%) either disrupted a stop codon or introduced a premature one,
which is one indicator of false positives (Fig. 2a). For context, this catalog is 10-fold larger than the one
established by Schloissnig et al. and 1.22-fold larger than the catalogue of all human SNPs[19] (Fig. S1).
Consistent with previous reports[4], SNP density, nucleotide diversity, and the rate of nonsynonymous
versus synonymous mutations (pN/pS) varied across species and phyla (Fig. 2b and Fig. S5-8), which
may reflect differences in selective pressures, population sizes, or transmission modes.
98
We hypothesized that the SNP database could be greatly compressed by clustering SNPs into linkage
disequilibrium (LD) blocks that co-vary across reference genomes (Fig. S9) and selecting a single "tag"

SNP per LD block. A similar strategy is commonly used when designing genotyping arrays in human
genetics. Using single-linkage clustering ($R^2 > 0.81$), the 104 million SNPs were clustered into 6.8
million LD blocks, representing a >15-fold reduction in database size and revealing a remarkable degree
of local genomic structure. Our choice of R^2 is motivated by thresholds used for high confidence SNP
imputation in other species and the fact that discovery of LD blocks stabilizes in this range for gut species
(Fig. S10). On average LD blocks spanned ~4.3Kbp and ~23.5 SNPs, though the number and size of LD
blocks varied considerably across bacterial species (Fig. 2c, S5c and S11a and b). As expected, linkage
between SNPs decayed with increasing genomic distance (Fig. 2d-f), though decay rates differed
substantially across species (Fig. 2d-e). Altogether, these differences in genetic diversity and structure
across species likely reflect variation in recombination rates and/or the number and relatedness of
sequenced genomes.
112
113 *Species-specific kmers enable accurate and efficient identification of SNPs*
114
Having constructed a large SNP catalog of the gut microbiome, we next used GT-Pro to identify k-mers
that could unique identify each SNP from shotgun metagenomes. We empirically determined that length
k=31 ensured high specificity while limiting compute and memory requirements. Of the ~13.3 billion
candidate 31-mers that overlapped a SNP (124 per SNP), we identified 5.7 billion that were unique. These
kmers overlapped 51% of the 104 million SNPs for 65% of LD blocks (mean 108 sck-mers per SNP, >1
sck-mer for 97% of species, Fig. S1 and S12). We refer to these as species-specific, SNP-covering kmers
(sck-mers). Species with few or no SNPs that can be genotyped with this strategy include those with a
very close relative and are most common within Actinobacteria (Fig. 2g and S3b). While only 50% of
SNPs were tagged by a sck-mer, they capture 83% of the within species variation compared to whole-
genome average nucleotide identity, and achieve a much higher level of resolution compared to individual
taxonomic markers (Fig. S13). Due to the large scale of the database, GT-Pro uses a highly efficient data
structure to store the sck-mers, requiring only 13 GB of RAM and permitting GT-Pro to run on most
modern personal computers (Fig. S14 and S15). We also created a low memory version of the GT-Pro
database (< 4 GB RAM) which just stores sck-mers for a single "tag" SNP per LD block (Methods) and
still captures the majority of within species variation (Fig. S13).
130
131 *Optimized k-mer exact matching accelerates metagenotyping 100-fold*
132
To search for exact matches between billions of k-mers among metagenome reads and billions of sck-
mers in the GT-Pro database, it is crucial to have a highly efficient search algorithm with low RAM and
I/O requirements. To this end, we developed an exact match algorithm that leverages data structures
optimized for this specific application (Fig. S16). Our approach is similar to a multi-index search with
three main steps operating on bit encoded k-mers (2 bits per base) (Fig. S16a). After generating all k-mers
in each metagenomic sequencing read, GT-Pro uses a l-bit Bloom filter on the first l<k bits of each k-mer
to quickly rule out the vast majority of read k-mers that have no chance to match database sck-mers
because they do not share an l-mer. For the k-mers that pass through the l-bit filter, the algorithm recruits
an m-bit (last m bits of encoded k-mer) index to serve as secondary filter that locates a bucket of pre-
sorted sck-mers in the database containing all possible exact matches to the full k-mer. Finally, the
algorithm invokes a sequential search for exact matches between the full k-mer and these only the sck-
mers in this bucket.
145
We next evaluated GT-Pro computational performance. First, we measured both speed and peak RAM
use while tuning the values of l and m, two parameters derived from the l-bit and m-bit filter that are
expected to have a large impact on performance due to their direct relationships with query speed and
peak RAM use. In general, both performance metrics increase with higher values of l and m (Fig. 3a).
Within the range of the tested parameters, we found best speed and peak RAM use with l = 30 and m = 35

in the laptop environment (26.5GB RAM) and the with $l = 32$ and $m = 36$ on a server (56.55 GB RAM).
In a boundary case ($l = 30$ and $m = 36$) on the laptop where the peak RAM use hit the hardware limit,
speed drops >87%. These results demonstrate that the values of l and m should be carefully chosen based
on the hardware for optimal performance, which is handled automatically by GT-Pro.
155
We then compared the computational performance of GT-Pro to traditional read alignment method as
baseline (Fig. 3b). We arbitrarily selected a total of 40 stool metagenomes from a Tanzanian cohort[20]
(Table S7) for the evaluation. For alignment, SNPs were called mapping reads to database from GT-Pro
and an independent one (metaSNV[7]). Although GT-pro had a larger peak RAM use than alignment
method (<10GB), was 100x faster on a server and 10x faster on a laptop where peak RAM use was
26.5GB (Fig. 3b).
162
163
164 *Accurate identification of SNPs from simulated metagenomes*
165
We next evaluated the accuracy of SNP calling with GT-Pro compared to alignment using simulated
metagenomes. Towards this goal, we generated Illumina sequencing reads *in silico* from 978 human gut
isolates[21] and identified the ground truth set of SNPs based on whole-genome alignment. We first
simulated reads from individual isolates with sequencing coverages ranging from 0.001x to 15x (Table
S2). In this simplified scenario, genotyping errors can result from sequencing errors, insufficient
coverage, or incorrect read or k-mer mapping. Across isolates and coverage levels, the false discovery
rate (FDR) of genotypes was on average lower for GT-Pro (median= 0.7%, IQR=1.1%) compared to read
alignment (median= 2.2%, IQR=4.7%) (Fig. 4a) while the median sensitivity of GT-Pro tended to be
consistently higher (4.1-17.6%) at all coverages (Fig. 4b). While read alignment methods typically use a
minimum coverage threshold to avoid false positives from sequencing error (e.g. >10x), that would have
further decreased the sensitivity in this experiment.
177
Next, we simulated metagenomes containing pairs of conspecific isolates to evaluate performance on
samples with strain mixtures, exploring a range of coverage ratios from 0.001x to 15x, where one strain is
always at 15x coverage and the other varying (Table S3 and S4). In terms of detecting heterozygous sites
(strains with different alleles), the false discovery rate (FDR) of GT-Pro (median= 0.9%, IQR=0.7%) was
slightly higher compared to alignment (median= 0.3%, IQR=0.6%) (Fig. 4c), however, median sensitivity
was higher (50.5-81.6%) for GT-Pro at all coverage ratios (Fig. 4d). A higher FDR for GT-Pro is likely
caused by sequencing errors that match the alternative allele by chance, which also could cause a slightly
lower sensitivity for GT-Pro at homozygous sites (Fig. S17).
186
To evaluate genotype calls imputed from tag SNPs, we found low FDR<5% comparing to true genotypes
for the vast majority of isolates (>95%) (Fig. 4e). SNPs belonging to an LD block were 5 times more
likely to be detected (non-zero read count) when their tag SNPs were also detected than when they were
not (Fig. S18). To show that GT-Pro is highly quantitative, we compared average coverage at SNPs in the
GT-Pro output to the known genome coverage using metagenomes we simulated from individual isolates
and pairs of conspecific isolates. Even at low sequencing coverage (<1X), GT-Pro was able to accurately
estimate the true coverage of each species (Fig. 4f) and the ratio between two strains (Fig. 4g). These
results suggest that GT-Pro allele calls and counts could be used to impute genotypes and estimate
relative abundances of species and strains accurately.
196
197 *Accurate metagenotyping and gene imputation from gut metagenomes*
198
To compare GT-Pro to existing approaches, we metagenotyped gut metagenomes[16,20,22,23] (Table S5-10)
with alignment and compared the number of genotyped SNPs plus estimates of allele frequencies and

genetic distances. We found that GT-Pro genotyped more species and SNPs per metagenome (Fig. S19a-
c), despite being limited to species with ≥10 genomes. This is likely due to GT-Pro having better
sensitivity for low coverage species and using a human gut focused database (comparing to metaSNV).
For species genotyped by both methods, within-sample heterozygosity (Fig. S18) and across-sample
allele presence and frequency (Fig. 5a-d) were highly correlated. For high coverage species, alignment
method detected some SNPs absent from the GT-Pro database, whereas GT-Pro detected more sites as
polymorphic in medium and low coverage species (Fig. 5a-d). Despite these differences in genotyped
sites, GT-Pro and alignment produce highly similar estimates of pairwise genetic distances (Jaccard
index) between samples, likely because rare variants missed by GT-Pro but with sufficient coverage to be
genotyped with alignment-based methods represent a small fraction of overall genetic diversity. For
comparison, we repeated this analysis using only SNPs in the 16S gene and observed much lower genetic
differences between samples (Fig. S21), emphasizing that GT-Pro provides strain resolution close to that
of alignment and greatly exceeding that of marker gene approaches. Altogether these results are
consistent with our simulations and underscore the high sensitivity of GT-Pro.
215
Next, we sought to determine if GT-Pro SNPs could be used to infer the presence of nearby genes or
operons, thereby serving as biomarkers for structural variants. As a case study, we used GT-Pro SNPs in
flanking genes to predict presence/absence of toxicity controlling genes in *Clostridium difficile (C.*
difficile). We used the GT-Pro SNPs from two 5' (CD2601 and CD2602), one 3' (trpS) gene and
intergenic region to train a Random Forest classifier to predict the presence of the genomic region
(CdtLoc) of three toxin genes (CD196_cdtA, CD630_cdtAB and cdtR) in a set of *C. difficile* isolate
genomes downloaded independently from NCBI (Fig. S22a and S23a). In another example, we
demonstrated that SNPs (cdd2, cdu1, and intergenic) flanking a pathogenicity locus (PaLoc) region could
predict its presence (Fig. S22b and S23b). Next we applied these models to GT-Pro metagenomes from
7,459 samples (Fig. 5e and f). Our predictions of CdtLoc and PaLoc region presence were highly
correlated with estimated presence based on read alignment to the *C. difficile* genome, especially in
metagenomes where this species was more abundant, and weaker predictions were made when not all of
the genes in CdtLoc or PaLoc were present (Fig. S24a and b). These results show that GT-Pro can detect
structural and strain variants when they are in high LD with flanking common SNPs.
230
231 *Depicting novel and global intra-species genetic structure with GT-Pro*
232
To evaluate the commonality of SNPs in GT-Pro database and how GT-Pro perform in metagenotying
unknown metagenomes. We next used GT-Pro's common SNPs to perform dimension reduction on the
genomes in the database as well as metagenomic samples from a North American IBD cohort[24] (n=220;
Table S11) that did not contribute genomes to the GT-Pro database. Looking for evidence of subspecies
genetic structure, we observed that for most species the metagenomes clustered with the genomes (Fig. 6a
and b), suggesting that GT-Pro's database represents the common diversity across diverse metagenomes.
For a few species, however, we observed clusters comprised only of metagenomes (Fig. 6c and d),
demonstrating that novel subspecies genetic structure can be discovered using GT-Pro common SNPs.
241
Having shown GT-Pro is faster and at least as accurate as alignment-based methods for genotyping
common SNPs from metagenomes, we leveraged GT-Pro metagenotypes to conduct the most
geographically diverse intra-species genetic variation meta-analysis to date, encompassing 51.8 million
SNPs for 881 species found in 7,459 gut samples from 31 locations across six continents (Table S13).
Consistent with prior studies[4,6,9], we observed much less allele sharing between hosts (median=0.03,
IQR=0.05) than within a host over time (median=0.38, IQR=0.4), and that intra-host allele sharing varies
greatly between species and hosts (Fig. S25). Inter-host allele sharing differed across countries and
continents (Fig. S26a and b), generally decreasing with geographic distance (Fig. S26a and b and S27a
and b) and varying across species (Fig. S28). Our results also show clear associations with degree of

industrialization as well as relatedness of hosts (e.g., hosts within villages in Fiji share more alleles than
unrelated hosts in North American cities) (Fig. 6e). To identify gut species with high levels of inter-
continental population differentiation, we calculated FST for 78 prevalent and well-detected (see
methods) species and observed large differences in the degree of differentiation across species (Fig. 6f).
Species with high FST show distinct clusters of hosts, some but not all of which correlate with geography
(Fig 6g), consistent with lifestyle and environment playing a role in which strains colonize a host. In
contrast, hosts do not cluster as clearly based on species relative abundance (Fig 6h), emphasizing that
metagenotypes may reveal microbiome-host associations missed in abundance analyses.
259
260
261
262 **Discussion**
263
Here, we greatly extended the gut microbiome genomic variation landscape by identifying more than 100
million common core-genome SNPs from 909 bacterial species. As our solution to the bioinformatics
challenge of metagenotyping, GT-Pro avoids computationally costly alignment and overcomes computing
barriers. It performs strain-level analyses of microbiomes with improved accuracy, especially for low
coverage species. Studies of microbiome genetic variation on a laptop or at the scale of human genome-
wide association studies will be computationally feasible with GT-Pro.
270
It should be noted that our method comes with several limitations. First, the GT-Pro database does not
capture all human gut microbial diversity. While we used 909 species, we could not use the majority of
the UHGG species due to limited availability of high-quality genomes. Second, GT-Pro is analogous to a
genotyping array and hence does not identify novel SNPs, which require other methods, such as
alignment-based SNP calling or single-cell genome sequencing. For some species, the common SNP pool
is expected to expand through additional genome sequencing. Third, a small number of species lacked
species-specific sck-mers due to the presence of highly related species in the genome collection. Separate
strategies such as using longer k-mers or less common SNPs could enable GT-Pro metagenotyping for
these species. Fourth, although we were very selective in the choice of genomes and SNPs used for
building GT-Pro, it is impossible to exclude all imperfections (e.g. incompleteness, contaminations and
species misclassification) in the genome assemblies that could contribute to false SNP calls. Finally, GT-
Pro does not directly genotype structural variants, which contribute significantly to intra-species genetic
diversity[25]. However, we did show that GT-Pro can be used to impute insertions and deletions in high LD
with common SNPs. Despite these caveats, we showed that the GT-Pro framework is general, accurate
and sensitive for identifying genetic variation in metagenomes.
286
We envision several directions for future work. First, this study applied the GT-Pro approach to human
gut prokaryotic species, and the framework could easily be expanded to other kingdoms and
environments. Another extension is to develop alignment-free metagenotyping for short indels and
structural variants. This study barely scratches the surface in terms of interpreting microbiome genetic
variation. Towards leveraging microbiomes in precision medicine, it will be critical to comprehensively
identify SNPs that are associated with disease and other traits (e.g. pathogenicity, antimicrobial
resistance, drug degradation). We anticipate that GT-Pro will also be useful for detecting contamination,
recombination, and horizontal gene transfer events, as well as tracking variants or strains over time, host
lifestyle and geography.
296
297

298   **Reference**
299
1.   Garud, N. R. & Pollard, K. S. Population Genetics in the Human Microbiome. *Trends Genet.* **36**,
53–67 (2020).
2.   Maini Rekdal, V., Bess, E. N., Bisanz, J. E., Turnbaugh, P. J. & Balskus, E. P. Discovery and
inhibition of an interspecies gut bacterial pathway for Levodopa metabolism. *Science* **364**, eaau6323
(2019).
3.   Treangen, T. J., Ondov, B. D., Koren, S. & Phillippy, A. M. The Harvest suite for rapid core-
genome alignment and visualization of thousands of intraspecific microbial genomes. *Genome Biol.* **15**,
524 (2014).
4.   Schloissnig, S. *et al.* Genomic variation landscape of the human gut microbiome. *Nature* **493**, 45–
50 (2013).
5.   Luo, C. *et al.* ConStrains identifies microbial strains in metagenomic datasets. *Nat. Biotechnol.*
33, 1045–1052 (2015).
6.   Nayfach, S., Rodriguez-Mueller, B., Garud, N. & Pollard, K. S. An integrated metagenomics
pipeline for strain profiling reveals novel patterns of bacterial transmission and biogeography. *Genome*
Res. **26**, 1612–1625 (2016).
7.   Costea, P. I. *et al.* metaSNV: A tool for metagenomic strain level analysis. *PLOS ONE* **12**,
e0182392 (2017).
8.   Quince, C. *et al.* DESMAN: a new tool for de novo extraction of strains from metagenomes.
Genome Biol. **18**, 181 (2017).
9.   Truong, D. T., Tett, A., Pasolli, E., Huttenhower, C. & Segata, N. Microbial strain-level
population structure and genetic diversity from metagenomes. *Genome Res.* **27**, 626–638 (2017).
10.   Wood, D. E. & Salzberg, S. L. Kraken: ultrafast metagenomic sequence classification using exact
alignments. *Genome Biol.* **15**, R46 (2014).
11.   Ounit, R., Wanamaker, S., Close, T. J. & Lonardi, S. CLARK: fast and accurate classification of
metagenomic and genomic sequences using discriminative k-mers. *BMC Genomics* **16**, 236 (2015).
12.   Liu, Y., Zhang, L. Y. & Li, J. Fast detection of maximal exact matches via fixed sampling of
query K-mers and Bloom filtering of index K-mers. *Bioinformatics* **35**, 4560–4567 (2019).
13.   Shajii, A., Yorukoglu, D., William Yu, Y. & Berger, B. Fast genotyping of known SNPs through
approximate k-mer matching. *Bioinforma. Oxf. Engl.* **32**, i538–i544 (2016).
14.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.*
25, 1043–1055 (2015).
15.   Almeida, A. *et al.* A unified sequence catalogue of over 280,000 genomes obtained from the
human gut microbiome. *bioRxiv* 762682 (2019) doi:10.1101/762682.
16.   Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000
Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**, 649-662.e20 (2019).
17.   Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New insights from
uncultivated genomes of the global human gut microbiome. *Nature* **568**, 505–510 (2019).
18.   Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**, 499–504
(2019).
19.   Auton, A. *et al.* A global reference for human genetic variation. *Nature* **526**, 68–74 (2015).
20.   Smits, S. A. *et al.* Seasonal cycling in the gut microbiome of the Hadza hunter-gatherers of
Tanzania. *Science* **357**, 802 (2017).
21.   Zou, Y. *et al.* 1,520 reference genomes from cultivated human gut bacteria enable functional
microbiome analyses. *Nat. Biotechnol.* **37**, 179–185 (2019).
22.   Turnbaugh, P. J. *et al.* The Human Microbiome Project. *Nature* **449**, 804–810 (2007).
23.   Bäckhed, F. *et al.* Dynamics and Stabilization of the Human Gut Microbiome during the First
Year of Life. *Cell Host Microbe* **17**, 852 (2015).

24.   Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in inflammatory bowel
disease. *Nat. Microbiol.* **4**, 293–305 (2019).
25.   Tierney, B. T. *et al.* The Landscape of Genetic Content in the Gut and Oral Human Microbiome.
Cell Host Microbe **26**, 283-295.e8 (2019).
352

353
354 **Figure legends**
355
Figure 1. In sillico metagenotyping framework
Our method starts with a whole genome sequence collection and identifies species with sufficient high-
quality genomes to call SNPs. For each species, a representative genome is chosen based on pairwise
Average Nucleotide Identity (ANI) plus assembly quality metrics. SNPs are called per species based upon
whole genome alignment of conspecific genomes to the representative genome. Common (MAF > 1%)
bi-allelic SNPs are selected for genotyping. Up to 4 X k candidate k-mers are extracted per SNP site,
covering both the reference and alternative allele on forward and reverse complementary strands (sck-
mers; k=31 in this study). These candidate sck-mers are iteratively filtered through species-specificity
filters of all unique k-mers present in the genomes of every other species, including species with
insufficient high-quality genomes for genotyping. Only SNPs with sck-mers for both the reference and
alternative allele are retained. SNPs are clustered based on pairwise linkage disequilibrium (LD). LD
blocks are detected with a threshold of mean r2 > 0.81, and we select a tag SNP with species-specific sck-
mers and the highest LD to other SNPs in the block. Optimized algorithms and compressed
representations of sck-mer data enable rapid metagenotyping. Further details in Methods and Figure 3.
370
Figure 2. Genetic landscape of 909 human gut species.
(a) Summary of common SNP characteristics across all species (from left to right): at most SNPs only
two alleles are observed, bi-allelic SNPs are mostly within protein-coding genes, these are largely
synonymous, and the non-synonymous ones rarely disrupt or introduce a stop codon. (b and c) Phyla
differ in their median SNP density and average LD block size with significant variation in density across
species within each phylum. (d) Rate of LD distance decay across gut bacterial species. (b-e) are colored
by bacterial phylum and share the same color scheme. (e) Examples of LD distance decay for individual
species. From top to bottom are three species (species id: 102446, 101694 and 102831) with increasingly
fast LD distance decay, suggesting higher recombination rates. Curves represent the fitted exponential
decay model. (f) Visualization of two distinct haplotype landscapes from (upper) species *Alistipes*
putredinis (species id: 101302) and (lower) *Bacteroides xylanisolvens* (species id: 101345). Base axis
represents and is ordered by genomic coordinate. Color indicates magnitude of LD between pairs of
SNPs. The examples have the same genomic span (10,000 bp). (g) Distribution across species of the
percentage of SNPs that can be genotyped by GT-Pro either directly ("without LD blocks") or by
imputation using genotyped tag SNPs ("with LD blocks"). For a typical species, ~75% of SNPs can be
genotyped directly and ~95% can be imputed.
387
Figure 3. Computational performance evaluation of GT-Pro.
(a) Computational performance of GT-Pro in laptop (left) and server (right) environments across values
for l (Bloom filter size parameter) and m (m-index size parameter). Color gradient: processing speed,
circle size: peak RAM use, black box: optimal l and m for each computing environment. (b) Comparison
of speed (upper) and peak RAM usage (lower) between GT-Pro and alignment-based metagenotyping
(metaSNV and MIDAS; see methods). We ran GT-Pro on both server (green) and laptop (yellow)
environments, while alignment-based methods were run only in the server (grey) environment due to not
being optimized for personal computers. Peak RAM usage exceeds RAM needed to store the database due
to intermediate calculations, such as applying filters.
397
Figure 4. Metagenotyping accuracy evaluation of GT-Pro using simulations.
Accuracy comparisons of GT-Pro and alignment-based metagenotyping across species based on reads
simulated from isolate genomes with sequencing error. (a) False discovery rate at a combination of
sequencing coverage ranged from 0.001x to 15x. Each observation is the result from a metagenome
containing reads from one isolate. False discoveries are genotype calls that do not match the genome from

which reads were simulated. (b) Sensitivity across coverage levels from the simulations in (a). Sensitivity
is the probability of detecting SNPs present in the isolate genome. (c) False discovery rate at
heterozygous sites in metagenomes containing reads from two isolates of each species. A combination of
sequencing coverage ratio between two isolates was simulated by fixing a more abundant isolate at 15x
coverage in all simulations, and varying the other isolate's coverage from 0.001x to 15x (coverage ratio =
0.001:15 to 15:15). (d) Sensitivity at heterozygous sites in metagenomes from (c). Sensitivity is the
probability of correctly calling the heterozygous genotype of sites that differ between the genomes from
which reads were simulated. (e) False discovery rate of genotypes imputed from tag SNPs based on allele
matching in simulations in (a). Imputation is simply done by selecting the genotype associated with the
observed tag SNP. (f) Sequencing coverage estimated using read counts at GT-Pro genotyped SNPs
correlates with the simulated coverage, even when coverage is <1x. Each observation is the estimate from
metagenomic reads simulated with sequencing error from a single isolate genome. (g) Sequencing
coverage ratio estimates based on read counts for each allele at GT-Pro genotyped heterozygous sites
correlate with the simulated ratios of two isolate genomes, even when one is much less abundant than the
other (<=1:15). The more abundant isolate is at 15x coverage in all simulations.
418
Figure 5. Metagenotyping and gene imputation from gut metagenomes
Comparison of metagenotypes from GT-Pro and alignment with gut microbiome samples from a North
America cohort[22] (HMP project; n=358; Table S8). As an example, we show the species *Bacteroides*
stercoris (species id: 103681). Each point represents a SNP, with color indicating if the genotypes from
the two methods agree (green), both methods return a genotype but the alleles disagree (purple), or only
GT-Pro returns a genotype (black). Disagreements largely occur near 0.5 allele frequency, where small
differences in read counts per allele can "flip" the major and minor alleles. (a) The proportion of samples
in which each SNP is genotyped (prevalence) is similar with both methods. (b) Average allele frequency
across samples varies across SNPs but is highly correlated between the two methods. (c and d)
Comparison similar as (a and b) showing the species GCA_000431835.1 (genus: Succinivibrio, species
id: 100412) from a different Madagascar cohort[16] (n=112; Table S9). Prediction of presence/absence of *C*.
difficile pathogenic gene sets in human gut metagenomes from a mix of cohorts (n=7459) (Table S14) a
random forest classifier built using GT-Pro SNPs from flanking regions in 117 *C*. *difficile* isolates
(Figures S23-S24) with 10-fold cross validation. Heatmaps show the predicted (first column) and
observed (based on alignment, second column) presence (black) or absence (white) in each sample
(rows). Barplots show *C. difficile* relative abundance (left), whole genome sequence coverage (middle),
and number of detected genes from the pathogeneticity locus (right), all estimated by mapping reads from
each sample to a *C. difficile* representative genome. Random Forest predictions correlate with abundance,
coverage, and number of detected pathogenic genes (Figure S25). (l) CdtLoc genes. (m) PaLoc genes.
438
Figure 6. Global genetic structure in 6,452 human gut metagenomes.
(a-d) Gut species differ in the amount of common SNP genetic diversity already present in sequenced
genomes. Metagenomic samples from a North American IBD cohort[24] (n=220; Table S11) (purple) are
visualized in two dimensions alongside the UHGG genomes (green). Each plot is the result of applying
UMAP to a matrix of genotypes at GT-Pro SNPs for one species. Each dot represents a strain of that
species (major allele for heterozygous metagenomes); those closer together in UMAP space have more
similar genotypes. (a) *Anaerostipes hadrus* (species id: 102528) and (b) *Ruminococcus_B faecis* (species
id: 100249) are species where metagenomes lie within the diversity previously captured by genomes. (c)
Blautia_A obeum (species id: 100212) and (d) *Dialister invisus* (species id: 104158) are species where
metagenomes harbor combinations of common SNPs outside the range present in genomes, which may
represent novel subspecies. (e) Heatmap of mean allele sharing scores over all species between
metagenomes from different pairs of countries. Crossed cells indicates missing scores due to insufficient
(< 5000) pairs of samples. (f) Analysis of inter-continental population differentiation (FST) for 78
prevalent species. Each boxplot represents a distribution of inter-continental FST for one species, ordered

by medians. (g) An example of geographic patterns captured by within-species genetic variation in the
GT-Pro metagenotypes of specific species. Each dot is a metagenomic sample, colored by continents.
Dimension reduction and visualization performed with UMAP. The example is from *Agathobacter*
rectalis (species id: 102492). Nearby samples in UMAP space have similar abundance profiles; the
absence of distinct groups indicates that relative abundance does not show strong geographic clustering.
(h) UMAP analysis based on the relative abundances of the 881 GT-Pro species in the same samples as
(i).

Figure 1

Figure 2

472   **Figure 3**
473

Figure 4

Figure 5

482   **Figure 6**

484