Predicting feral cat-reduction targets and costs on large islands using stochastic

population models

Kathryn R. W. Venning[1,*], Frédérik Saltré[1], Corey J. A. Bradshaw[1]

[1]Global Ecology, College of Science and Engineering, GPO Box 2100, Flinders University,

Adelaide, South Australia 5001, Australia

*Corresponding author: Global Ecology, College of Science and Engineering, Flinders

University, GPO Box 2100, Adelaide, South Australia 5001, Australia; tel: +61 8 8201 2090;

e-mail: kathryn.venning@flinders.edu.au

Acknowledgments

The authors acknowledge the traditional owners of the land we work on and pay our respects

to Elders, past, present and emerging. We thank the Kangaroo Island section of the South

Australia Department of Environment and Water for data access and insights.

Data availability

The R code described in this manuscript is available at

https://github.com/KathrynVenning/FeralCatEradication

Declaration of interest

NIL

Predicting feral cat-reduction targets and costs on large islands using stochastic

population models

Abstract

Feral cats are one of the most destructive invasive predators worldwide. Due to the high risk

of pushing native species to extinction in Australia, density-reduction or eradication

campaigns can greatly improve the persistence probability and recovery of native fauna.

Kangaroo Island — Australia's third-largest island — was recently nominated as a complete

cat-eradication site by the federal government. Because most population density-reduction

campaigns are costly and not effective in the long term, mathematical models predicting

optimal culling regimes can guide management plans, especially if they include estimates of

costs under different policy decisions. We constructed a stochastic population model with

cost estimates to test the relative effectiveness and cost-efficiency of two main culling

scenarios for Kangaroo Island to 2030: (1) constant proportional annual cull, and (2) high

initial culling followed by a constant proportional maintenance cull. We also examined the

effectiveness of a trap-neuter-return scenario to compare with the culling outcomes. We

found that an average culling proportion of ≥ 0.3 would reduce the population to $\leq 10\%$ of

the founding population, while a two-phase cull where an initial cull of ≥ 0.6 was followed

by a maintenance cull of ≥ 0.45 would reduce the final population to 1% of its initial size by

2030. Costs estimates varied widely depending on capture techniques used, but a

combination of *Felixer*™ cat-eradication units, conventional traps, and hunting was the most

cost-effective (minimum costs estimated at AU$46.5 million–AU$51.6 million). Our model

results provided direction for the efficient eradication of feral cats on Kangaroo Island and

can be transferred to feral-cat management elsewhere.

Key words: Leslie matrix, matrix population model, feral cats, invasive species, eradication

costs, Kangaroo Island, culling model

Introduction

Since its domestication approximately 10,000 years ago, the common house cat *Felis catus*

has spread throughout the globe and become established in most habitat types (including

most islands) (Fitzgerald et al. 1991, Medina et al. 2011, Woinarski et al. 2015), due to both

accidental and deliberate human facilitation. Because they are generalist predators, feral cats

are today one of the most destructive invasive mammal predators worldwide (Lowe et al.

2000), contributing many or most of the predation-induced terrestrial extinctions recorded

globally (e.g., > 63 species, including 26% of bird, mammal and reptile extinctions) (Doherty

et al. 2016).

Cats are opportunistic hunters and scavengers that are able to adapt easily to fluctuating

prey availability, switching between small invertebrates to mammals as large or even larger

than themselves (Barratt 1997, Bonnaud et al. 2011, Moseby et al. 2015). They can also

easily adapt to arid and hostile environments, surviving with minimal access to fresh water

(Fitzgerald and Turner 2000, Medina et al. 2008, Woinarski et al. 2015). This makes feral

cats particularly destructive to island ecosystems, because they have the added vulnerability

of naïve endemic prey (Close 2005, Bonnaud et al. 2011, Medina et al. 2014, Doherty et al.

2016). Australia in particular is itself a large island, disproportionately represented in

mammal extinctions since European settlement (Abbott 2002, Abbott 2008) so that in

combination with foxes (*Vulpes Vulpes*), cats have become one of the most prolific killers

across the continent (Dickman 1996, Doherty et al. 2016).

The most effective method for reducing feral cat populations and their associated

predation on native species is eradication wherever possible (Andersen et al. 2004, Schmidt

et al. 2009), particularly in insular environments (Bester et al. 2002, Nogales et al. 2004).

Alternative non-lethal approaches (such as trap-neuter-release) exist (Gibson et al. 2002,

Wallace and Levy 2006, Longcore et al. 2009, Miller et al. 2014), but the threat cats pose to

native fauna makes such methods too slow and ineffective for an ecologically meaningful

reductions in cat densities and predation rates (Longcore et al. 2009). While this approach

might appeal to a section of society that does not agree with killing to control (Andersen et al.

2004), the high expense of implementing the approach at sufficiently broad scales, coupled

with its low effectiveness (Longcore et al. 2009), mean that it is not widely used for feral cat

management.

Most density-reduction campaigns based on direct killing have been typically

implemented ad hoc because of the imminent threat cats posed to threatened species and the

requirement to achieve outcomes quickly (Bester et al. 2002, Denny and Dickman 2010). As

such, available funds or resources are often used up quickly without the benefit of long-term

planning based on the projections of empirical density-reduction models (Denny and

Dickman 2010). Unsurprisingly, this unplanned approach typically renders such eradication

attempts unsuccessful (Bester et al. 2002). Campbell et al. (2011) found that inappropriate

methods and poorly timed roll-out were attributed to most island eradication failures.

Custom-designed culling models that take habitat-specific population dynamics into account

while planning the most efficient and cost-effective application of resources are therefore

ideal precursors to any eradication program (Smith et al. 2005, McMahon et al. 2010).

Culling models are effective because of their ability to consider real-time population

dynamics and resources available to recommend feasible density-reduction plans. Many types

of culling models exist depending on the choice of scenario to reduce the population, such as

constant proportional culling (McCarthy et al. 2013), or a high initial proportional culling rate

followed by a constant proportional maintenance thereafter (a two-phase reduction model)

(Campbell et al. 2011). Such models are instrumental in guiding successful eradication by

providing targets and parameters that lead to efficient population reduction of the target

species (Smith et al. 2005). Eradication often employs a three-part strategy: (*i*) an initial high

cull, (*ii*) a consistent maintenance cull to ensure continued population decline, and (*iii*) a final

'clean up' stage. In the clean-up stage, management typically needs to switch eradication

tactics to remove the last remaining individuals in the population as they become more

difficult to detect (Bester et al. 2002, Nogales et al. 2004).

Cat eradication is currently underway on part of Kangaroo Island, Australia's third-largest

island (Figure 1). The initial planning stages of the eradication began in 2016, with a

proposed completion date of 2030. The management strategy for eradication is as follows:

2016-2019: planning and trialling control techniques, 2019-2023: create a safe haven by

eradicating cats from the Dudley Peninsula on the east of the island, and 2023-2030:

eradication on the remainder of island, with full eradication complete by 2030 (Natural

Resources 2013, Department of the Environment and Energy 2015). The program directors

plan to use three main techniques for feral cat eradication: trapping, hunting, and *Felixers™*

(an automated toxin-delivery system that uses rangefinder sensors to distinguish target cats

from non-target species and spray targets with a measured dose of toxic gel; thylation.com).

Although poison baiting (using 1080 or PAPP) has been proposed, the approach is not

currently being considered for Kangaroo Island due to potential risk to domestic animals and

native animal species.

Our aim was to design an ideal set of culling conditions that will most efficiently reduce

feral cat densities on Kangaroo Island following the timeline established above. More

specifically, we (1) constructed a culling model that can be applied to guide cat eradication

on Kangaroo island, (2) identified the relative costs of employing different combinations of

the methods available, and (3) used the culling model to identify a regime that will most

effectively reduce the feral cat population by the 2030 deadline. Specifically, we tested the

efficacy (proportion of the population reduced, and over what time) of two culling methods:

(*i*) constant proportional annual culling, (*ii*) high initial culling followed by a constant

maintenance cull. We hypothesise that a two-phase eradication program, with a high initial

proportional reduction and steady maintenance culling thereafter, will reach the target

population density by 2030 more efficiently than a constant annual culling rate. The two-

phase model will likely be the most effective in reducing feral cat densities because high

initial effort tends to be the cheapest and most effective means of achieving high rates of

reduction (Bester et al. 2002, Nogales et al. 2004, Robertson 2008, Denny and Dickman

2010), but maintenance culling is required thereafter to prevent the population from

recovering.

Methods

Study site

Located approximately 12 km south of the Fleurieu Peninsula (South Australia) at its nearest

point, Kangaroo Island is Australia's third largest island (155 km long and 55 km wide),

covering ~ 4400 km^2 (Masters et al. 2004, Higgins-Desbiolles 2011) (Figure 1). The island

has retained around 53% of its native vegetation, with 35% of the remaining land cover

devoted to dryland agriculture (Willoughby et al. 2018). The island is uniquely absent of

invasive European rabbits (*Oryctolagus cuniculus*) and is one of the few locations in

Australia free of red foxes (*Vulpes Vulpes*); resident cats can therefore feed on a wider range

of native species than elsewhere on mainland Australia (Bonnaud et al. 2011). For our study,

we separated the island into the Dudley Peninsula (east of the narrowest point of Kangaroo

Island, adjacent to Pelican Lagoon; Figure 1) and the western side (west of the narrowest

point), to reflect the management strategy for the entire island.

Feral cat densities on Kangaroo Island are thought to range from 0.15 to 2.97 km^{-2}, with a

median of 0.7 km^{-2} on the western end (Hohnen et al. 2020). Taggart et al. (2019) estimated

that relative feral cat densities in eastern Kangaroo Island were 10 times higher than on the

adjacent mainland.

Model

We constructed Leslie matrices to represent age-specific fertility and survival (Caswell 2001)

for the cat population. Leslie matrices (also known as population demographic matrices)

represent age-specific fertility and survival for a given population (Caswell 2001). We

obtained cat fertility and survival estimates from six studies of domestic, stray and feral cat

population across the USA and Australia (Budke and Slater 2009) and summarised the

population dynamics from a study in Western Australia done in the preliminary stages of cat

eradication (Short and Turner 2005). We calculated mean and standard deviations of the age-

specific demographic rates (i.e., survival, fertility) necessary for stochastic representations of

the model (see below). We only used these fertility and survival estimates for females,

assuming a 1:1 sex ratio (Bloomer and Bester 1991, Budke and Slater 2009).

The maximum age for feral cats on Kangaroo Island is 6 years; however, we split the

initial year into two sections: pre- and post-breeding. The pre-breeding time step incorporates

animals < 10 months of age, because feral cats reach sexual maturity at around 10 months;

this was represented as individuals in this stage having zero fertility. For all resulting

predictions of changing population size, we assumed that survival was the same for males

and females.

We stochastically resampled the deterministic matrix A in all subsequent projections by

resampling based on the standard deviation estimated from minimum and maximum fertility

and survival values (Budke and Slater 2009), which incorporates both measurement error and

inter-annual variability (process error). We assumed a Gaussian distribution around the mean

of both survival and fertility, using the standard deviations for resampling (Table 1).

Untreated population

To simulate how incrementing intensities of reduction change the projected population size,

we first simulated a population not exposed to any culling as a benchmark. We defined the

population's stable age distribution (Caswell 2001), and then multiplied this stable age

structure by a starting population size of 3083 (approximation from density estimates on the

Dudley Peninsula and western side of island) (Taggart et al. 2019, Hohnen et al. 2020). We

then expressed all subsequent projections as a proportion of this founding population size to

avoid the uncertainty in initial population size estimates. Because Kangaroo Island is insular,

we assumed a closed population structure (i.e., no immigration or permanent emigration).

Although cats could be brought to the island or domestic cats can be abandoned, there are

insufficient data to consider the population as demographically open.

We included a logistic compensatory density-feedback function by reducing survival when

the population exceeded double the size of the current population of the form:

$$S_{\text{mod}} = \frac{\kappa}{1 + \left(\frac{N}{\tau}\right)^{\theta}}$$

where, S_{mod} is the proportion of realised survival (survival modifier) as a function of the population's proximity to carrying capacity (twice the founding population size, $K = 6166$), N is the population size, and κ, τ and θ are constants: $\kappa = 0.998$, , $\tau = 14821.78$ and $\theta = 2.835$. We thus assumed survival to reduce as population approaches carrying capacity, with no changes to fertility because landscape managers currently consider the population to be sub-optimal with respect to available food resources (Jones and Coman 1982, Read and Bowen 2001). Most research on feral cat population control does not consider the habitat's carrying capacity (Andersen et al. 2004); however, feral cats seem to maintain consistent fertility regardless of population density, although average survival tends to decrease as the population approaches carrying capacity (Courchamp and Sugihara 1999, Nutter 2006).

Reduction scenarios

1. Trap-neuter-release

To compare the efficacy of all proposed density control and reduction scenarios with fertility-reduction methods, we constructed a model that simulated a trap-neuter-release scenario. In this model, no animals are removed from the population, but fertility is reduced to simulate sterilisation. We ran this model using the same methods for an unculled population, but we ran an incrementing overall reduction in fertility from 50–99%.

2. Culling model

We ran each culling scenario for (*a*) the entire island, (*b*) the Dudley Peninsula, and (*c*) the western side of Kangaroo Island to reflect the current management plan. We built two culling models: (*i*) constant proportional annual culling, and (*ii*) a high initial proportional cull in the first two years, followed by a constant proportional maintenance cull. In each model, we removed individuals from the population vector proportional to the total culling invoked in that time step and the stable age distribution. We ran each model for 10,000 iterations (randomly sampling 10,000 times from the stochastic survival and fertility vectors) to calculate the mean and 95% confidence bounds for minimum proportional population size. Projection intervals varied depending on the geographic scale: ten years (2020–2030) for the entire island, three years (2020–2023) for the Dudley Peninsula, and seven years (2023–2030) for the western side of the island.

For the constant cull, we simulated constant proportional annual culling (*c*) (i.e., we

reduced the population each year by the same proportion for the duration of the projection

interval) from *c* = 0.10 to 0.90. Once we determined the minimum initial culling proportion

that precipitated a decline, we refined the *c* proportions considered from 0.50 to 0.99 in 0.01

intervals. For the two-phase approach, we applied high initial culling only to the first two

years of the eradication project (*c* = 0.50–0.99), with the maintenance culling applied to all

years thereafter (*c* = 0.01–0.50) until the end of the projection interval. For all iterations of

both models, we recorded the minimum projected proportional population size (p*N*) for each

value of *c*, with a culling proportion interval of 0.01.

Cost

Based on previous information regarding the reduction in capture efficiency as population

density declines (Bloomer and Bester 1992, Nogales et al. 2004, Parkes et al. 2014), we

assumed an eradication technique's efficiency (*f*, ranging from 0 to 1) follows a Type III

functional response (i.e., sigmoidal; Nunney 1980, Denno and Lewis 2009) relative to

proportional population size:

236
$$f = \frac{\alpha}{1 + (\beta e^{-\gamma \mathrm{p}N})}$$

where *f* is the relative efficiency of the culling technique, *α, β* and *γ* are constants: *α* = 1.01 (±

0.01), *β* = 85.61 (± 15.40) and *γ* = 8.86 (± 0.41), and p*N* = proportional population size. We

assumed the same efficiency reduction across trapping, hunting and *Felixers™* as a function

of population size, such that the smaller the remaining population of cats, the less efficient

each method was relative to the start of the eradication campaign. We then applied this

reduction to the culling model with pre-set costs for each technique (see below), to estimate

the total cost of eradication.

We sourced trapping and shooting cost data from Holmes et al. (2015), with additional

trapping costs from trapping supplies (traps.com.au) and *Felixer™* data from Hodgens

(2019). We summarised the catch rates and costs for each technique: (*i*) *Felixer™* — each

unit costs AU$13,000. Based on trials on Dudley Peninsula, 6 *Felixer™* units were

successful in killing 13 cats over 38 days, which translates to an annual kill rate of 0.080 (cats

killed unit^{-1} year^{-1}). (*ii*) Traps — each trap costs between AU$157 and AU$297 (sampled

uniformly). Again, based on trials on Dudley Peninsula, 40 traps caught 21 cats over 148

days, which translates to an annual trap rate of 0.198 cats trap^{-1} year^{-1}. (*iii*) Shooting — from

Holmes et al. (2015), we estimated a kill rate per person-hour based on 1044 kills (872 direct

+ 172 from wounds) over 14,725 person-hours (= 0.071 cats killed person^{-1} hour^{-1}).

Ammunition and labour costs equate to AU\$25.92 hour^{-1}.

Assuming the same density of *Felixers™* and traps on Dudley Peninsula applied to the

entire island based on a comparison of relative areas (Dudley Peninsula = 37,500 ha;

Kangaroo Island = 440,500 ha), we tabulated the number of cats killed by these two methods

combined for the incrementing proportional cull, and then calculated the shortfall of cats not

killed to meet the proportional cull required for that year. We considered three different

approaches to meeting this shortfall: (*i*) increasing the number of *Felixers™* only, (*ii*)

increasing the number of traps only, or (*iii*) meeting the shortfall entirely with follow-up

hunting. In each shortfall scenario, we tabulated the total costs across the projection interval

and expressed these as a function of the increments in proportional culling.

Results

Untreated population

An untreated (no-cull) population is expected to increase to a median of 1.8 times the

founding population (i.e., to 5525 individuals when starting with 3083) by 2030 (95%

confidence limits: 1.39–2.21) (Figure 2a). The instantaneous rate of change (*r*) from the

deterministic matrix for the Kangaroo Island population is 0.167. The deterministic (mean)

matrix gave a generation length of 4.7 years. The population is projected to approach

carrying capacity (set arbitrarily at twice the current population size) by 2026.

Trap-neuter-release

We simulated a trap-neuter-release fertility reduction to compare its efficiency to direct

killing methods. Reduced total fertility of 50% and 80% yielded *r* = 0.10 and -0.19,

respectively. A reduction of 80% fertility each year reduced the population to a median

proportion of 0.0754 (0.048–0.12), with 50% fertility control only reducing the population to

0.66 (0.041–0.974) of the founding population.

Culling

Dudley Peninsula 2019–2023: A consistent yearly cull of ≥ 0.52 reduced the population to ≤

0.10 of the founding population, with an annual cull of 0.68 and 0.88 resulting in 0.032 and <

0.01 of the founding population, respectively. A two-phase model reduced the population ≤

0.10 with an initial cull of ≥ 0.70 and a maintenance cull of ≥ 0.45, while a combination of ≥

0.7 (initial) and ≥ 0.65 (maintenance) would reduce the population to ≤ 0.036.

Western Kangaroo Island 2023–2030: An annual proportional cull of ≥ 0.3 would reduce

the population ≤ 10%, while an annual cull of ≥ 0.45 would reduce the population to 0.01 of

its original size. To reduce the population to < 0.10, a two-phase cull of ≥ 0.6 and ≥ 0.2

would be required. An initial cull of ≥ 0.6 followed by a maintenance cull of ≥ 0.45 would

reduce the population to ≤ 0.01.

The entire island 2019–2030: For the constant proportional cull, the population will be

reduced to ≤ 0.10 of the founding population with an annual proportional cull of ≥ 0.3. An

annual cull of ≥ 0.45 would reduce the population to ≤ 0.01 (Figure 2b). A two-phase cull

with ≥ 0.60 (initial) and ≥ 0.2 (maintenance) would reduce the population to ≤ 0.10. To

achieve ≤ 0.01, ≥ 0.60 (initial) and ≥ 0.45 (maintenance) is required (Figure 3).

Cost

To reduce the entire Kangaroo Island population to a 0.10 of its original size using a two-

phase cull, a minimum of AU$6.1 billion (AU$5.1 billion–AU$6.8 billion) would be required

if *Felixer™s* were used to make up the yearly shortfall (using the same densities of

Felixers™ and traps as on Dudley Peninsula) (Figure 4a). In contrast, making up the shortfall

with additional traps would place total costs at AU$53.1 million (AU$40.5 million–AU$69.1

million) for the same target (Figure 4b). Finally, making up the shortfall with hunting would

cost a total of AU$25.0 million (AU$22.7 million–AU$26.5 million) (Figure 4c).

Changing the target population size to 0.01 of the original, the total minimum costs would

increase to AU$16.9 billion (AU$15.8 billion–AU$18.0 billion) if the shortfall was made

with *Felixers™*, AU$129.3 million (AU$94.0 million–AU$167.8 million) if the shortfall was

made with traps, or AU$49.1 million (AU$46.5 million–AU$51.6 million) if the shortfall

was made with hunting (Figure 4).

Discussion

In each cull scenario we considered, a successful reduction of the feral cat population on

Kangaroo Island to below 0.01 of its original size is achievable, but the minimum costs

involved according to the different scenarios we ran could range from AU$46.5 million

(AU$10 ha^{-1}, making up shortfall with hunting; Figure 4c) to AU$18 billion (AU$40,863 ha^{-1},

making up shortfall with *Felixers™*,; Figure 4a) depending on the method used and the

inherent uncertainty in the parameters we estimated. Whether these costs are a true reflection

of the actual costs depends on our untested assumption of the functional response, and the

per-unit outlay of the different methods to meet the annual shortfall for the different

intensities of cull. However, the lower end of our estimates for Kangaroo Island are cheaper

than the costs to remove cats from Macquarie Island (AU\$258 ha^{-1}), likely due to the latter's

remoteness (Robinson and Copson 2014). Nonetheless, our comparisons do illustrate that

high initial culls (> 45%) followed by moderate maintenance culls (20–45%) would be

sufficient to reduce the population to 1-10% of its original size (Figure 3), and that hunting is

the most cost-effective way to meet these targets (especially if other methods are rolled out

simultaneously).

That hunting is cheaper than other methods is unsurprising given that no hardware other

than rifles and ammunition are needed to be purchased outright, in contrast to the higher

overheads associated with traps or *Felixer™* units (Holmes et al. 2015, Hodgens 2019).

Additionally, hunting is considered more humane due to the minimised contact with the

animal and the instant death with a correctly executed headshot (Sharp and Saunders 2011).

However, hunting requires many people working full time, whereas the other techniques are

more passive. Of course, complete eradication would necessarily entail additional costs as the

final individuals were identified, hunted, and destroyed (Bester et al. 2002, Nogales et al.

2004). This would be more expensive than normal eradication efforts and would require a

more strategic isolation of remaining individuals.

Our results identify that fertility-reduction using trap-neuter-release control methods are

an ineffective option for reducing pest densities (Longcore et al. 2009), even though it often

remains attractive for predator control in the public's eye because it does not involve direct

killing (Robertson 2008). Instead, our model outputs show that a realised annual fertility

equivalent to ≤ 25% of expected values (i.e., pre-fertility treatment) would reduce the

population to ≤ 10%. In other words, managers would have to maintain at least 75% of the

female cat population sterilised each year from 2019-2030 for the program to reduce the

population to ≤ 10% of its original size. Whether this is feasible or cost-effective is beyond

the scope of our study, but it does demonstrate that fertility-reduction is a much less efficient

method to eradicate cats than culling. Additionally, sterilised individuals returned to the

population could still continue to eat native fauna until they perished due to natural causes, so

the overall ecological impact would be necessarily greater than a culling-based program

where individuals are removed permanently when treated.

If left unmanaged, cats pose a serious threat to the persistence of native species. Indeed,

cats have been identified as one of the greatest threats to birdlife globally (Butchart et al.

2006), being major contributors or the sole source of predation and extinction of bird species

from New Zealand to Mexico (Nogales et al. 2004, Medina et al. 2011). Further, Smith and

Quin (1996) cite cats as the best predictor of population decline in small conilurine rodents in

Australia. The successful eradication of cats, especially from islands, has shown that in the

absence of cat predation, native fauna is often able to recover to viable population sizes

within a few generations (Bester et al. 2002, Hughes et al. 2008).

We conclude that the most appropriate approach to reduce cat densities on Kangaroo

Island is a two-stage method, with a high initial reduction of at least 60–70% and a

maintenance cull of 50–65%. Although a constant proportional annual cull can be effective, it

is generally less efficient than a two-stage approach. As culling reduces density and drives

the population closer to extinction, it becomes progressively more difficult and expensive to

cull remaining individuals (Nogales et al. 2004, Parkes et al. 2014). This is because most

culling methods are passive and rely on a 'non-negligible probability' of the target animal

encountering *Felixers*™ or traps (Moseby and Hill 2011). Although these techniques can be

accompanied by visual, scent or sound lures, the target animal still needs to be in range to be

enticed by them. Thus, encounters at low densities become increasingly less likely (Veitch

2001), and rising per-capita food abundance as the predator's population dwindles can make

baits or food lures less attractive (Parkes et al. 2014). Cats in particular are intelligent and can

learn to avoid traps. Thus, while trapping is generally considered effective for density

reduction (Nogales et al. 2004), it is most effective in the early stages of eradication programs

(Nogales et al. 2004). Therefore, a two-stage approach allows for the implementation of

widespread control that is effective at high densities, followed by a more targeted approach

through consecutive maintenance as the population continues to decline.

Because the Kangaroo Island trial eradication program is still in its initial stage, many of

the demographic, effectiveness, and cost parameters are not yet available specifically for this

region, lending further uncertainty to our conclusions. For example, we were obliged to

obtain demographic rates from data published for cat populations elsewhere (Budke and

Slater 2009), and we do not yet have access to spatially variable data with which to

parameterise spatially explicit models that could potentially improve realism (McMahon et

al. 2010). A better quantification of carrying capacity, as well as the mechanisms by which

density feedback operates on the demographic rates would also improve predictive accuracy.

Many attempts to eradicate cats have been made, but they vary widely in success (Algar et

al. 2002, Bester et al. 2002, Nogales et al. 2004, Algar et al. 2011, Campbell et al. 2011,

Parkes et al. 2014). For effective eradication to be achieved, culling programs must be based

on empirical data and ideally, directed by models like ours. Our model should allow

practitioners to plan their culling programs better, and to allocate the resources needed to

achieve their targets efficiently and cost-effectively. As more site-specific data become

available, we expect the model's predictions to become ever-more realistic to identify the

most plausible and cheapest pathways to eradication.

References

Abbott, I. 2002. Origin and spread of the cat, *Felis catus*, on mainland Australia, with a
discussion of the magnitude of its early impact on native fauna. Wildlife Research
29:51-74.10.1071/WR01011
Abbott, I. 2008. The spread of the cat, *Felis catus*, in Australia: re-examination of the current
conceptual model with additional information. Conservation Science Western
Australia **7**:1-17
Algar, D., G. Angus, and M. Onus. 2011. Eradication of feral cats on Rottnest Island,
Western Australia. Journal of the Royal Society of Western Australia **94**:439-443
Algar, D., A. Burbidge, and G. Angus. 2002. Cat eradication on Hermite Island, Montebello
Islands, Western Australia. Turning the tide: the eradication of invasive species:14-18
Andersen, M. C., B. J. Martin, and G. W. Roemer. 2004. Use of matrix population models to
estimate the efficacy of euthanasia versus trap-neuter-return for management of free-
roaming cats. Journal of the American Veterinary Medical Association **225**:1871-
1876.10.2460/javma.2004.225.1871
Barratt, D. G. 1997. Predation by house cats, *Felis catus* (L.), in Canberra, Australia. I. prey
composition and preference. Wildlife Research **24**:263-277.10.1071/WR96020
Bester, M., J. Bloomer, R. Van Aarde, B. Erasmus, P. Van Rensburg, J. Skinner, P. Howell,
and T. Naude. 2002. A review of the successful eradication of feral cats from sub-
Antarctic Marion Island, Southern Indian Ocean. South African Journal of Wildlife
Research **32**:65-73
Bloomer, J. P., and M. N. Bester. 1991. Effects of hunting on population characteristics of
feral cats on Marion Island. South African Journal of Wildlife Research **21**:97-102
Bloomer, J. P., and M. N. Bester. 1992. Control of feral cats on sub-Antarctic Marion Island,
Indian Ocean. Biological Conservation **60**:211-219.10.1016/0006-3207(92)91253-O
Bonnaud, E., F. M. Medina, E. Vidal, M. Nogales, B. Tershy, E. Zavaleta, C. J. Donlan, B.
Keitt, M. Le Corre, and S. V. Horwath. 2011. The diet of feral cats on islands: a
review and a call for more studies. Biological Invasions **13**:581-603.10.1007/s10530-
010-9851-3
Budke, C. M., and M. R. Slater. 2009. Utilization of matrix population models to assess a 3-
year single treatment nonsurgical contraception program versus surgical sterilization
in feral cat populations. Journal of Applied Animal Welfare Science **12**:277-
292.10.1080/10888700903163419
Butchart, S. H., A. J. Stattersfield, and N. J. Collar. 2006. How many bird extinctions have
we prevented? Oryx **40**:266-278
Campbell, K., G. Harper, D. Algar, C. Hanson, B. Keitt, and S. Robinson. 2011. Review of
feral cat eradications on islands. Island Invasives: Eradication and Management:37-46

Caswell, H. 2001. Age-Classified Matrix Models. Matrix Population Models, constuction,
analysis, and interpretation. Sinauer Associates, Inc., Sunderland, Massachusetts

Close, R. 2005. Introduced Mammals of the World: Their History, Distribution and
Influence. John Long. A Review by Robert L Close. Australian Mammalogy **27**:109-
110.10.1071/AM05109_BR

Courchamp, F., and G. Sugihara. 1999. Modeling the biological control of an alien predator
to pretect island species from extinciton. Ecological Applications **9**:112-
123.10.1890/1051-0761

Denno, R. F., and D. Lewis. 2009. Predator-Prey Interactions. Pages 202-212 *in* S. Levin A,
editor. The Princeton Guide to Ecology. Princeton University Press, Princeton, New
Jersey

Denny, E. A., and C. Dickman. 2010. Review of cat ecology and management strategies in
Australia. Invasive Animals Cooperative Research Centre, Canberra

Department of the Environment and Energy. 2015. Threatened species strategy. Australian
Government, Canberra

Dickman, C. R. 1996. Impact of exotic generalist predators on the native fauna of Australia.
Wildlife Biology **2**:185-195.10.2981/wlb.1996.018

Doherty, T., C. Dickman, C. Johnson, S. Legge, E. Ritchie, and J. Woinarski. 2016. Impacts
and management of feral cats *Felis catus* in Australia. Mammal Review **47**:83-
97.10.1111/mam.12080

Fitzgerald, B., B. Karl, and C. Veitch. 1991. The diet of feral cats (*Felis catus*) on Raoul
Island, Kermadec Group. New Zealand Journal of Ecology:123-129

Fitzgerald, B. M., and D. C. Turner. 2000. Hunting behaviour of domestic cats and their
impact on prey populations. The domestic cat: the biology of its behaviour **151**

Gibson, K. L., K. Keizer, and C. Golding. 2002. A trap, neuter, and release program for feral
cats on Prince Edward Island. The Canadian Veterinary Journal **43**:695-692

Higgins-Desbiolles, F. 2011. Death by a thousand cuts: governance and environmental trade-
offs in ecotourism development at Kangaroo Island, South Australia. Journal of
Sustainable Tourism **19**:553-570.10.1080/09669582.2011.560942

Hodgens, P. 2019. Felixer V's Felis Report. Kangaroo Island

Hohnen, R., K. Berris, P. Hodgens, J. Mulvaney, B. Florence, B. P. Murphy, S. Legge, C. R.
Dickman, and J. C. Z. Woinarski. 2020. Feral cat density and population size across
Kangaroo Island, South Australia 10.1071/WR19137:10.1071/WR19137

Holmes, N., K. Campbell, B. Keitt, R. Griffiths, J. Beek, C. Donlan, and K. Broome. 2015.
Reporting costs for invasive vertebrate eradications. Biological Invasions **17**:2913-
2925

Hughes, B. J., G. R. Martin, and S. J. Reynolds. 2008. Cats and seabirds: effects of feral
Domestic Cat *Felis silvestris catus* eradication on the population of sooty terns
Onychoprion fuscata on Ascension Island, South Atlantic. Ibis **150**:122-
131.10.1111/j.1474-919x.2008.00838.x

Jones, E., and B. Coman. 1982. Ecology of the feral cat, *Felis catus* (L.), in south-eastern
Australia III. Home ranges and population ecology in semiarid north-west Victoria.
Wildlife Research **9**:409-420.10.1071/WR9820409

Longcore, T., C. Rich, and L. M. Sullivan. 2009. Critical assessment of claims regarding
management of feral cats by trap–neuter–return. Conservation Biology **23**:887-
894.10.1111/j.1523-1739.2009.01174.x

Lowe, S., M. Browne, S. Boudjelas, and M. De Poorter. 2000. 100 of the world's worst
invasive alien species: a selection from the global invasive species database. Invasive
Species Specialist Group Auckland

Masters, P., T. Duka, S. Berris, and G. Moss. 2004. Koalas on Kangaroo Island: from
introduction to pest status in less than a century. Wildlife Research **31**:267-
272.10.1071/WR03007
McCarthy, R. J., S. H. Levine, and J. M. Reed. 2013. Estimation of effectiveness of three
methods of feral cat population control by use of a simulation model. Journal of the
American Veterinary Medical Association **243**:502-511.10.2460/javma.243.4.502
McMahon, C., B. Brook, N. Collier, and C. Bradshaw. 2010. Spatially explicit spreadsheet
modelling for optimising the efficiency of reducing invasive animal density. Methods
in Ecology and Evolution **1**:53-68.10.1111/j.2041-210X.2009.00002.x
Medina, F. M., E. Bonnaud, E. Vidal, and M. Nogales. 2014. Underlying impacts of invasive
cats on islands: not only a question of predation. Biodiversity and Conservation
23:327-342.10.1007/s10531-013-0603-4
Medina, F. M., E. Bonnaud, E. Vidal, B. R. Tershy, E. S. Zavaleta, C. J. Donlan, B. S. Keitt,
M. Corre, S. V. Horwath, and M. Nogales. 2011. A global review of the impacts of
invasive cats on island endangered vertebrates. Global Change Biology **17**:3503-
3510.10.1111/j.1365-2486.2011.02464.x
Medina, F. M., M. López-Darias, M. Nogales, and R. García. 2008. Food habits of feral cats
(*Felis silvestris catus* L.) in insular semiarid environments (Fuerteventura, Canary
Islands). Wildlife Research **35**:162-169.10.1071/WR07108
Miller, P. S., J. D. Boone, J. R. Briggs, D. F. Lawler, J. K. Levy, F. B. Nutter, M. Slater, and
S. Zawistowski. 2014. Simulating free-roaming cat population management options in
open demographic environments. PLOS ONE
9:e113553.10.1371/journal.pone.0113553
Moseby, K. E., and B. M. Hill. 2011. The use of poison baits to control feral cats and red
foxes in arid South Australia I. Aerial baiting trials. Wildlife Research **38**:338-
349.10.1071/WR10235
Moseby, K. E., D. E. Peacock, and J. L. Read. 2015. Catastrophic cat predation: A call for
predator profiling in wildlife protection programs. Biological Conservation **191**:331-
340.10.1016/j.biocon.2015.07.026
Natural Resources. 2013. KI feral cat eradication program: frequently asked questions.
Government of South Australia, Kangaroo Island
Nogales, M., A. Martín, B. R. Tershy, C. J. Donlan, D. Veitch, N. Puerta, B. Wood, and J.
Alonso. 2004. A review of feral cat eradication on islands. Conservation Biology
18:310-319.10.1111/j.1523-1739.2004.00442.x
Nunney, L. 1980. The influence of the type 3 (sigmoid) functional response upon the stability
of predator-prey difference models. Theoretical Population Biology **18**:257-
278.10.1016/0040-5809(80)90052-0
Nutter, F. B. 2006. Evaluation of a trap-neuter-return management program for feral cat
colonies: population dynamics, home ranges, and potentially zoonotic diseases. North
Carolina State University, North Carolina
Parkes, J., P. Fisher, S. Robinson, and A. Aguirre-Muñoz. 2014. Eradication of feral cats
from large islands: an assessment of the effort required for success. New Zealand
Journal of Ecology:307-314
Read, J., and Z. Bowen. 2001. Population dynamics, diet and aspects of the biology of feral
cats and foxes in arid South Australia. Wildlife Research **28**:195-
203.10.1071/WR99065
Robertson, S. A. 2008. A review of feral cat control. Journal of Feline Medicine and Surgery
10:366-375.10.1016/j.jfms.2007.08.003

Robinson, S. A., and G. R. Copson. 2014. Eradication of cats (*Felis catus*) from subantarctic
Macquarie Island. Ecological Management & Restoration **15**:34-
40.10.1111/emr.12073
Schmidt, P. M., T. M. Swannack, R. R. Lopez, and M. R. Slater. 2009. Evaluation of
euthanasia and trap-neuter-return (TNR) programs in managing free-roaming cat
populations. Wildlife Research **36**:117-125.10.1071/WR08018
Sharp, T., and G. Saunders. 2011. A model for assessing the relative humaneness of pest
animal control methods. Department of Agriculture, Fisheries and Forestry Canberra,
Australia
Short, J., and B. Turner. 2005. Control of feral cats for nature conservation. IV. Population
dynamics and morphological attributes of feral cats at Shark Bay, Western Australia.
Wildlife Research **32**:489-501.10.1071/WR04102
Smith, A. P., and D. Quin. 1996. Patterns and causes of extinction and decline in Australian
conilurine rodents. Biological Conservation **77**:243-267
Smith, G. C., H. I. S., and P. A. Robertson. 2005. A model of ruddy duck Oxyura jamaicensis
eradication for the UK. Journal of Applied Ecology **42**:546-555.doi:10.1111/j.1365-
2664.2005.00977.x
Taggart, P. L., B. A. Fancourt, A. J. Bengsen, D. E. Peacock, P. Hodgens, J. L. Read, M. M.
McAllister, and C. G. Caraguel. 2019. Evidence of significantly higher island feral cat
abundance compared with the adjacent mainland. Wildlife Research **46**:378-
385.10.1071/wr18118
Veitch, C. 2001. The eradication of feral cats (*Felis catus*) from little Barrier Island, New
Zealand. New Zealand journal of zoology **28**:1-12.10.1080/03014223.2001.9518252
Wallace, J. L., and J. K. Levy. 2006. Population characteristics of feral cats admitted to seven
trap-neuter-return programs in the United States. Journal of Feline Medicine and
Surgery **8**:279-284.10.1016/j.jfms.2006.02.004
Willoughby, N., D. Thompson, M. Royal, and M. Miles. 2018. South Australian land cover
layers: an introduction and summary statistics, DEW technical report 2018/01.
Technical Report, Government of South Australia, Department for Environment and
Water, Adelaide
Woinarski, J. C. Z., A. A. Burbidge, and P. L. Harrison. 2015. Ongoing unraveling of a
continental fauna: Decline and extinction of Australian mammals since European
settlement. Proceedings of the National Academy of Sciences **112**:4531-
4540.10.1073/pnas.1417301112
561
562

Figure Captions

Figure 1 – Map of Kangaroo island relative to the Australian mainland. The shortest distance from the mainland (southern tip of Fleurieu Peninsula) to Kangaroo Island is approximately 14 km.

Figure 2 – (a) Average proportion of the initial cat population (N_1) on Kangaroo Island projected from 2020–2030 for the unculled scenario. Black line indicates the median value from 10,000 iterations, along with 95% confidence intervals (dashed lines). (b) Minimum proportion of the Kangaroo Island feral cat population remaining after a constant proportional annual cull ranging from 0.5 to 0.9. Solid black line represents median minimum proportion of the initial population (N_1) after 10,000 iterations with 95% confidence intervals indicated as dashed lines.

Figure 3 –Estimated median minimum proportion of the final population remaining (relative to start population N_1) for combinations of initial proportional (i.e., *initial cull*: 0.5–0.9) and maintenance proportional (i.e., *maintenance cull*: 0.1–0.5) culling. Proportion of population remaining after culling scenarios represented by colour bar ranging from lowest (purple) to highest (yellow) remaining proportional population.

Figure 4 – Estimated median total costs of feral cat eradication on Kangaroo Island for combinations of initial proportional (i.e., *initial cull*) and maintenance proportional (i.e., *maintenance cull*) culling, where the shortfall in the number of cats killed from *Felixer*™ units and traps is provided by (a) additional *Felixer*™ units, (b) traps, or (c) hunting. Cost of eradication (in AU$, adjusted for 2020) indicated by colour bar ranging from lowest (purple) to highest (yellow) costs. Contours and white values indicate cost in $AU millions or billions accordingly. Note different *z*-axis (contour) scales in a, b, and c.

Table 1 – Demographic rates (survival probability and fertility — kittens/female) used in the stochastic matrices to project the cat population under different culling scenarios (Budke and Slater 2009).

	minimum	mean	maximum
fertility			
pre-breeding juvenile	0.00	0.00	0.00
subadult	0.35	0.75	1.58
adult	1.98	2.52	3.78
survival			
pre-breeding juvenile	0.27	0.46	0.73
subadult	0.27	0.46	0.73
adult	0.55	0.70	0.78

pre-breeding juvenile: < 10 months old; not sexually mature

Figure 1

Figure 2

Figure 3

Figure 4