Predicting targets and costs for feral-cat reduction on large islands

using stochastic population models

Kathryn R. W. Venning[1,*], Frédérik Saltré[1], Corey J. A. Bradshaw[1]

[1]Global Ecology, College of Science and Engineering, GPO Box 2100, Flinders University,

Adelaide, South Australia 5001, Australia

*Corresponding author: Global Ecology, College of Science and Engineering, Flinders

University, GPO Box 2100, Adelaide, South Australia 5001, Australia; tel: +61 8 8201 2090;

e-mail: kathryn.venning@flinders.edu.au

Acknowledgments

The authors acknowledge the traditional owners of the land we work on and pay our respects

to Elders, past, present and emerging. We thank the Kangaroo Island section of the South

Australia Department of Environment and Water for data access and insights.

Data availability

The R code to create the model simulations is available at

https://github.com/KathrynVenning/FeralCatEradication

Declaration of interest

NIL

Predicting targets and costs for feral-cat reduction on large islands using stochastic population models

Abstract

Feral cats are one of the most destructive invasive predators worldwide. Due to the high risk of feral predators pushing native species to extinction in Australia, density-reduction or eradication campaigns can improve the persistence probability and recovery of native fauna. Kangaroo Island — Australia's third-largest island — was recently nominated as a complete cat-eradication site by the federal government. Because many population density-reduction campaigns are costly and not effective in the long term, mathematical models predicting optimal culling regimes can guide management plans, especially if they include estimates of costs under different policy decisions. We constructed a stochastic population model with cost estimates to test the relative effectiveness and cost-efficiency of two main culling scenarios for Kangaroo Island to 2030: (1) constant proportional annual cull (one-phase), and (2) high initial culling followed by a constant proportional maintenance cull (two-phase). We also examined the effectiveness of a trap-neuter-release scenario to compare with the culling outcomes. We found that an average culling proportion of ≥ 0.3 would reduce the population to ≤ 0.1 of the founding population, while a two-phase cull where an initial cull of ≥ 0.6 was followed by a maintenance cull of ≥ 0.45 would reduce the final population to 0.01 of its initial size by 2030. Costs estimates varied widely depending on capture techniques used, but a combination of *Felixer*™ cat-eradication units, conventional traps, and hunting was the most cost-effective in the two-phase culling scenario when hunting is used to make up culling shortfalls (minimum costs estimated at AU\$46.5 million–AU\$51.6 million with an initial cull of ≥ 0.6, maintenance cull of ≥ 0.45). Trap-neuter-release was an inefficient approach compared to the culling scenarios. Our model results provided direction for the efficient eradication of feral cats on Kangaroo Island and can be transferred to feral-cat management elsewhere.

Key words: Leslie matrix, matrix population model, feral cats, invasive species, eradication costs, Kangaroo Island, culling model

Introduction

Since its domestication approximately 10,000 years ago, the common house cat *Felis catus*

has spread throughout the globe and become established in most habitat types (including on

most islands) (Fitzgerald et al. 1991, Medina et al. 2011, Woinarski et al. 2015), due to both

accidental and deliberate human facilitation (Driscoll et al. 2007). Because they are generalist

predators, feral cats are today one of the most destructive invasive mammal predators

worldwide (Lowe et al. 2000, Doherty et al. 2016), contributing to many or most of the

predation-induced terrestrial extinctions recorded globally (e.g., > 63 species, including 26%

of bird, mammal and reptile extinctions) (Doherty et al. 2016).

Cats are also considered to be opportunistic hunters and scavengers that are able to adapt

to fluctuating prey availability, switching between small invertebrates to mammals as large or

even larger than themselves (Barratt 1997, Bonnaud et al. 2011, Moseby et al. 2015). They

can also easily adapt to arid and hostile environments, surviving with minimal access to fresh

water (Fitzgerald and Turner 2000, Medina et al. 2008, Woinarski et al. 2015). This makes

feral cats particularly destructive to island ecosystems, where their prey have the added

vulnerability of naivety to generalist predators for which they have evolved no or inefficient

anti-predator behaviours (Close 2005, Bonnaud et al. 2011, Medina et al. 2014, Doherty et al.

2016). Australia in particular is a large island, disproportionately represented in mammal

extinctions since European settlement (Abbott 2002, Abbott 2008) so that in combination

with foxes (*Vulpes vulpes*), cats have become one of the most prolific killers of native species

across the continent (Dickman 1996, Doherty et al. 2016).

The most effective method for reducing feral cat populations and their associated

predation on native species is eradication wherever possible (Andersen et al. 2004, Schmidt

et al. 2009), particularly in insular environments (Bester et al. 2002, Nogales et al. 2004,

Doherty et al. 2016). Alternative non-lethal approaches (such as trap-neuter-release) also

exist (Gibson et al. 2002, Wallace and Levy 2006, Longcore et al. 2009, Miller et al. 2014),

and while such an approach might appeal to members of the public that do not agree with

lethal control (Andersen et al. 2004), the high expense of broad-scale implementation,

coupled with its low effectiveness (Longcore et al. 2009) mean it is not widely used for cat

management in Australia. Despite this, the trap-neuter-release management option is

commonly considered in density-control programs, or proposed by communities (Deak et al.

2019).

Most density-reduction campaigns based on direct killing have been typically

implemented *ad hoc* because of the imminent menace cats pose to their prey species and the

requirement to achieve outcomes quickly (Bester et al. 2002, Denny and Dickman 2010). As

such, available funds or resources can be used up quickly without the benefit of long-term

planning based on the projections of empirical density-reduction models (Denny and

Dickman 2010), thus threatening the success of a program. As a result, inappropriate methods

and poorly timed roll-out have been attributed to most island eradication failures (Campbell

et al. 2011). Custom-designed culling models that plan the most efficient and cost-effective

application of resources are therefore ideal precursors to any eradication program (Smith et

al. 2005, McMahon et al. 2010).

Culling models are effective because of their ability to consider real-time population

dynamics and resource availability to recommend feasible density-reduction plans

(McMahon et al. 2010). Multiple types of culling model exist (e.g., spatially explicit, aspatial,

density-driven, area-dependent) depending on the choice of scenario to reduce the population,

such as a single-phase, constant proportional culling (McCarthy et al. 2013), or a two-phase

cull with a high initial proportional culling rate followed by a constant proportional

maintenance thereafter (a two-phase reduction model) (Campbell et al. 2011). Such models

are instrumental in guiding successful eradication by providing targets and parameters that

lead to efficient population reduction of the target species (Smith et al. 2005). Such two-

phase eradication strategies (high initial cull followed by a consistent maintenance 'press'

cull to ensure continued population decline) often still require a final 'clean up' stage where

different (and usually more expensive) strategies are needed to eradicate the last surviving

individuals that are difficult to detect (Bester et al. 2002, Nogales et al. 2004). High initial

culls followed by maintenance culling capitalise on the notion that when densities are high,

culling is more efficient, while the maintenance culling continues to reduce the population

(Nogales et al. 2004, Denny and Dickman 2010).

Cat eradication is currently underway on part of Kangaroo Island, Australia's third-largest

island (Fig. 1). The initial planning stages of the eradication began in 2016, with a proposed

completion date of 2030. The program directors plan to use three main techniques for cat

eradication: trapping, hunting, and *Felixer*™ units (a *Felixer* is an automated toxin-delivery

system that uses rangefinder sensors to distinguish target cats from non-target species and

spray targets with a measured dose of toxic gel; thylation.com). The social licence to apply

lethal population reduction via hunting, baiting, or trapping is largely a function of the

public's perception of the proposed methods (Deak et al. 2019), and relies on the co-

operation of land-owners. This applies to Kangaroo Island given it has > 4200 permanent

residents spread across most of the island. Perception surveys done between 1993 and 2018

showed that > 90% of the Kangaroo Island community supported domestic and feral cat

management (Berris et al. 2019). Additionally, stakeholders (community members, land

owners, relevant governmental departments) are in support and are actively involved in the

development and implementation of the project (Natural Resource Kangaroo Island 2015).

Kangaroo Island has pre-existing community engagement for wildlife-conservation

programs, and local residents are educated about the damage caused by feral cats (Berris et

al. 2019). These public-information programmes improve the local public's perception of

feral-animal management (Deak et al. 2019, Walsh et al. 2019).

Our aim was to design an ideal set of culling conditions that will most efficiently reduce

feral cat densities on Kangaroo Island. More specifically, we (1) constructed stochastic

variants of both culling and fertility-reduction (trap-neuter-release) models under different

application scenarios that can be applied to guide cat eradication on Kangaroo Island, (2)

estimate the relative costs of employing different combinations of the methods available, and

(3) used the culling model to identify a regime that will most effectively reduce the feral cat

population by the 2030 deadline. Specifically, we tested the efficacy (proportion of the

population reduced, and over what time) of two culling scenarios: (*i*) constant proportional

annual culling (one-phase), (*ii*) high initial culling followed by a constant maintenance cull

(two-phase). We hypothesise that the two-phase culling model will reach the target

population density by 2030 more efficiently than the one-phase culling model because initial

effort tends to be the cheapest and most effective means of achieving high rates of reduction

(Bester et al. 2002, Nogales et al. 2004, Robertson 2008, Denny and Dickman 2010). In any

case, maintenance culling is required thereafter to prevent the population from recovering.

Methods

Study site

Located approximately 12 km south of the Fleurieu Peninsula (South Australia) at its nearest

point, Kangaroo Island is Australia's third largest island (155 km long and 55 km wide),

covering ~ 4400 km^2 (Masters et al. 2004, Higgins-Desbiolles 2011) (Fig. 1). The island has

retained around 53% of its native vegetation, with 35% of the remaining land cover devoted

to dryland agriculture (Willoughby et al. 2018). The island is uniquely absent of invasive

European rabbits (*Oryctolagus cuniculus*) and is one of the few locations in Australia free of

red foxes (*Vulpes vulpes*); resident cats can therefore feed on a wider range of native species than elsewhere on mainland Australia (Bonnaud et al. 2011). For our study, we separated the island into the Dudley Peninsula (east of the narrowest point of Kangaroo Island, adjacent to Pelican Lagoon; Fig. 1) and the western side (west of the narrowest point), to reflect the management strategy for the entire island.

Feral cat densities on Kangaroo Island are thought to range from 0.15 to 2.97 cat km^{-2}, with a median of 0.7 cat km^{-2} on the western end (Hohnen et al. 2020). Taggart et al. (2019) estimated that relative feral cat densities in eastern Kangaroo Island were ~ 10 times higher than on the adjacent mainland (Kangaroo Island relative abundance = 14.6 cats camera-trap site^{-1}; mainland = 1.39 cats site^{-1}; 11 sites on both the Island and mainland).

Model

We constructed a Leslie matrix to represent age-specific fertility and survival (Caswell 2001) for the cat population. We obtained cat fertility and survival estimates from six studies of domestic, stray and feral cat population across the USA and Australia (Budke and Slater 2009) and summarised the population dynamics from a study in Western Australia done in the preliminary stages of cat eradication (Short and Turner 2005). We calculated mean and standard deviations of the age-specific demographic rates (i.e., survival, fertility) necessary for stochastic representations of the model (see below). We only used these fertility and survival estimates for females, assuming a 1:1 sex ratio (Bloomer and Bester 1991, Budke and Slater 2009).

According to demographic rates published in the peer-reviewed literature, the maximum age for feral cats ranges from 3 (Budke and Slater 2009) to 9 years (Van Aarde 1983); however, after consulting local wildlife managers, we set the maximum age for feral cats on Kangaroo Island at 6 years. We then split the initial year into two sections: pre- and post-breeding. The pre-breeding time step incorporates animals < 10 months of age, because feral cats reach sexual maturity at around 10 months (Jones and Coman 1982); this was represented as individuals in this stage having zero fertility. For all resulting predictions of changing population size, we assumed that survival was the same for males and females given no evidence to the contrary. We present all parameters and their ranges in Table 1.

We stochastically resampled the deterministic matrix **A** in all subsequent projections based on the standard deviation estimated from minimum and maximum fertility and survival values (Budke and Slater 2009), which incorporates both measurement error and inter-annual variability (process error). We assumed a Gaussian distribution around the mean of fertility

186   and the β distribution for survival probability, using the standard deviations for resampling of

187   each (Table 1).

188   The deterministic matrix **A** is:

$$\mathbf{A} = \begin{bmatrix} f_1 & f_2 & f_3 & f_4 & f_5 & f_6 & f_7 \\ s_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & s_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & s_3 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & s_4 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & s_5 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & s_6 & s_7 \end{bmatrix}$$

where f_x = age (x) -specific fertility and s_x = age-specific survival. See also Table 1 for

parameter values.

Untreated (control) population

To simulate how incrementing intensities of reduction change the projected population size,

we first simulated a population not exposed to any culling to represent a 'control' population.

We calculated the population's stable age distribution from the base matrix **A** (Caswell

2001), and then multiplied this stable age structure by a starting population size of 3083

(approximation from density estimates on the Dudley Peninsula and western side of island)

(Taggart et al. 2019, Hohnen et al. 2020). We then expressed all subsequent projections as a

proportion of this founding population size to avoid the uncertainty in initial population size

estimates. Kangaroo Island is insular, so there are fewer opportunities for migration into the

population compared to the mainland. However, we did account for some 'leakage' into the

population (domestic release or ferry stow-away) by re-running the top-performing culling

scenario (initial proportional cull = 0.6, maintenance cull = 0.45 — see Results) and adding

an incrementing a number of 'leaked' individuals (between 10 and 1000 cats) into the

population annually to test how these additions would affect final population sizes post-

culling (see Supplementary Information, Appendix A, Fig. S1).

We included a logistic compensatory density-feedback function by reducing survival when

the population exceeded double the size of the current population (see below) of the form:

$$S_{\mathrm{mod}} = \frac{\kappa}{1 + \left(\frac{N}{\tau}\right)^{\theta}}$$

where S_{mod} is the proportion of realised survival (survival modifier) as a function of the

population's proximity to carrying capacity (twice the founding population size, $K = 6166$;

see below), N is the population size, and κ, τ and θ are constants: $\kappa = 0.998$, $\tau = 9139.11$, and

$\theta = 2.835$ (see Supplementary Material, Appendix B, Fig. S2). We thus assumed that survival

probability would decline as the population approached carrying capacity (double the size of

the current population based on local landscape managers' recommendations), because

landscape managers currently consider the population to be sub-optimal with respect to

available food resources (Jones and Coman 1982, Read and Bowen 2001). While the

threshold is somewhat arbitrary, it does realistically allow the population to increase if no

additional mortality sources are imposed. Most research on feral cat population control does

not consider the habitat's carrying capacity (Andersen et al. 2004); however, feral cats seem

to maintain consistent fertility regardless of population density, although average survival

tends to decrease as the population approaches carrying capacity (Courchamp and Sugihara

1999, Nutter 2006). We therefore did not adjust fertility relative to population size.

Reduction scenarios

1. Trap-neuter-release

To compare the efficacy of all proposed density control and reduction scenarios with fertility-

reduction methods, we constructed a model that simulated a trap-neuter-release scenario.

Although not widely implemented, the trap-neuter-release method is often suggested by a

certain element of the public as a more ethical alternative to lethal control. We included this

scenario here to compare its efficacy directly to the culling scenarios described below. In this

model, no animals are removed from the population, but fertility is reduced to simulate

sterilisation. We ran this model using the same methods for an unculled population (and over

the same interval), but we reduced fertility for each iteration across a range of values (50–

99%, at 1% intervals). That is, the realised fertility for each iteration varied between 50% and

1% of the 'control' model's values for the duration of the project.

2. Culling model

We built two culling models: (*i*) constant proportional annual culling (one-phase), and (*ii*) a

high initial proportional cull in the first two years, followed by a constant proportional

maintenance cull (two-phase). Here we consider only these phases of a strategy where the last

step likely requires a 'clean-up' — the latter is difficult to consider in a model because the

focus shifts to individuals. We instead defined a threshold at the end of the model, where

moving to the 'clean-up' stage is deemed feasible. In each model, we removed individuals

from the population vector proportional to the total culling invoked in that time step and the

stable age distribution. We ran each model for 10,000 iterations (randomly sampling 10,000

times from the stochastic survival and fertility vectors) to calculate the mean and 95%

confidence bounds for minimum proportional population size. We set the projection interval

to 10 years to represent the management strategy for eradication by 2030 (2020–2030).

For the one-phase scenario, we simulated constant proportional annual culling (c) (i.e., we

reduced the population each year by the same proportion for the duration of the projection

interval) from $c = 0.10$ to 0.90. We first determined the minimum c to provoke a population

decline (this was 0.50 proportional cull), we then incremented this parameter between 0.50

and 0.99 at intervals of 0.01. For the two-phase scenario, we applied high initial culling only

in the first two years of the eradication project ($c = 0.50$–0.99), with maintenance culling

applied to all years thereafter ($c = 0.01$–0.50) until the end of the projection interval. For all

iterations of both models, we recorded the minimum projected proportional population size

(pN) for each value of c, at an incrementing proportional culling of 0.01.

Cost

Based on previous information regarding the reduction in capture efficiency as population

density declines (Bloomer and Bester 1992, Nogales et al. 2004, Parkes et al. 2014), we

assumed an eradication technique's efficiency (f, ranging from 0 to 1) follows a Type III

functional response (i.e., sigmoidal; Nunney 1980, Denno and Lewis 2009) relative to

proportional population size:

269
$$f = \frac{\alpha}{1 + (\beta e^{-\gamma pN})}$$

where f is the relative efficiency of the culling technique, α, β and γ are constants: $\alpha = 1.01$, β

$= 85.61$, and $\gamma = 8.86$, and pN = proportional population size (see Supplementary Material,

Appendix C, Fig. S3). We assumed the same efficiency reduction across trapping, hunting

and *Felixers™* as a function of population size, such that the smaller the remaining

population of cats, the less efficient each method was relative to the start of the eradication

campaign. We then applied this reduction to the culling model with pre-set costs for each

technique (see below), to estimate the total cost of eradication.

We sourced trapping and shooting cost data from Holmes et al. (2015), with additional

trapping costs from trapping supplies (traps.com.au) and *Felixer™* data from Hodgens

(2019). We summarised the catch rates and costs for each technique: (*i*) *Felixer™* — each

unit costs AU$13,000. Based on trials on part of Dudley Peninsula, 6 *Felixer™* units over

approximately 12,000 ha were successful in killing 13 cats over 38 days, which translates to

an annual kill rate of 0.080 (cats killed unit^{-1} year^{-1}). (*ii*) Traps — each trap costs between

AU$157 and AU$297 (sampled uniformly). Again, based on trials on Dudley Peninsula, 40

traps over approximately 12,000 ha caught 21 cats over 148 days, which translates to an

annual trap rate of 0.198 cats trap^{-1} year^{-1}. (*iii*) Shooting — from Holmes et al. (2015), we

estimated a kill rate per person-hour based on 1044 kills (872 direct + 172 from wounds) over

14,725 person-hours (= 0.071 cats killed person^{-1} hour^{-1}). Ammunition and labour costs

equate to AU$25.92 hour^{-1}.

To estimate total costs, we first assumed that the density of *Felixers*™ and traps applied

on Dudley Peninsula (Dudley Peninsula = 37,500 ha) could be extrapolated to the much

larger area of the entire island (440,500 ha). Based on these densities, we calculated the total

number of units and traps required for the entire island, and then tabulated the number of cats

killed by these two methods combined for the incrementing proportional cull. If the total

number of cats killed by these two methods fell short of the proportional cull target in any

given year and iteration, we applied three different scenarios where we varied the method

used to achieve the proportional target beyond the initial roll-out of units and traps. The three

different approaches to meet the shortfalls were: (*i*) increasing the number of *Felixers*™ only,

(*ii*) increasing the number of traps only, or (*iii*) meeting the shortfall entirely with follow-up

hunting. In each shortfall scenario, we tabulated the total costs across the projection interval

and expressed these as a function of the increments in proportional culling.

Of course, this approach assumes a simultaneous roll-out of all *Felixer*™ units and traps

across the entire island, when a more efficient approach might instead be to purchase a

smaller number of units/traps and deploy them in a spatially sequential roll-out (i.e., a

moving 'wave' of units applied to specific regions of the island in sequence). We therefore

also ran a modified scenario to reflect this type of spatial pattern of application by arbitrarily

assuming a smaller number of units/traps across the entire landscape. Reducing the purchase

cost per unit/trap by the same arbitrary value is therefore functionally equivalent to a spatially

sequential roll-out of this smaller sample of units/traps. For this example scenario, we

therefore reduced the purchase cost of both *Felixers*™ and traps by two-thirds unit^{-1} (see

Supplementary Material, Appendix D, Fig. S4).

Results

Untreated population

An untreated (no-cull) 'control' population is expected to increase to a median of 1.8 times

the founding population (i.e., to 5525 individuals when starting with 3083) by 2030 (95%

confidence limits: 1.39–2.21 times) (Fig. 2a). The instantaneous rate of change (r) from the

deterministic matrix for the Kangaroo Island population is 0.167. The deterministic (mean)

matrix gave a generation length of 4.7 years. The population is projected to approach

carrying capacity (set arbitrarily at twice the current population size) by 2026 (Fig. 2a), at

which time the population's $r = 0.0061$ (cf. initial $r = 0.190$).

Trap-neuter-release

The trap-neuter-release scenario with a fertility reduction of 80% each year reduced the

population to a median proportion of 0.0754 (0.048–0.120) of its initial size by 2030 (Fig. 3),

and a 50% fertility decline only reduced the population to 0.66 (0.041–0.974) of the founding

population over the same timeframe.

Culling

Culling timeline 2019–2030: For the one-phase scenario, an annual proportional cull of ≥ 0.3

would reduce the population to ≤ 0.10 of its initial size. An annual cull of ≥ 0.45 would

reduce the population to ≤ 0.01 of its initial size (Fig. 2b). A two-phase cull with ≥ 0.60

(initial) and ≥ 0.2 (maintenance) would reduce the population to ≤ 0.10 of its initial size. To

achieve a population ≤ 0.01 of its initial size, ≥ 0.60 (initial) and ≥ 0.45 (maintenance) are

required (Fig. 4). Stopping the program after the initial culling during the first two years (i.e.,

without any maintenance culling), the population would recover to its initial size in 15 years

(range: 11–21 years; Fig. S4), whereas stopping the maintenance cull in the 9th year (i.e., a

year before termination of the program) would result in population recovery to initial size in

42 years (range: 35–50 years) (Fig. S4).

Cost

To reduce the entire Kangaroo Island population to a 0.10 of its original size ($0.1N_1$) using a

two-phase cull, a minimum of AU\$25.0 million (AU\$22.7 million–AU\$26.5 million) (Fig.

5c), would be required if hunting was used to make up the yearly shortfall. In contrast,

making up the shortfall with additional traps (using the same densities of traps and *Felixers*™

as on Dudley Peninsula) would increase the average costs by 112.4% to AU\$53.1 million

(AU\$40.5 million–AU\$69.1 million) for the same target (Fig. 5b). Finally, making up the

shortfall with additional *Felixer*™ units would increase the average cost relative to the

hunting-shortfall scenario by > 24,000% to AU$6.1 billion (AU$5.1 billion–AU$6.8 billion)

(Fig. 5a). Changing the target population size to 0.01 of the initial (0.01N_1), the total

minimum costs would increase to AU$49.1 million (AU$46.5 million–AU$51.6 million) if

the shortfall was made with hunting (96.4% more than the 0.1N_1 hunting-shortfall scenario),

AU$129.3 million (AU$94.0 million–AU$167.8 million) if the shortfall was made with traps

(163.3% more than the 0.01N_1 hunting-shortfall scenario), or AU$16.9 billion (AU$15.8

billion–AU$18.0 billion) if the shortfall was made with *Felixers™* (> 34,000% more than the

0.01N_1 hunting-shortfall scenario) (Fig. 5).

Assuming a two-thirds reduction in costs for *Felixer™* and trap units (equivalent to

needing only one-third of the units rolled out in a spatially sequential manner), the scenario

where the shortfall is made up of additional *Felixer™* units, the total costs for the 0.01N_1

scenario (initial 0.6, maintenance 0.45), are reduced to AU$3.3 billion (AU$3.0 billion –

AU$3.7 billion) compared to the full-cost scenario for *Felixers*™ (Supplementary Material,

Appendix E, Fig. S5). The total costs for the 0.1N_1 scenario (0.6 initial, 0.2 maintenance), are

reduced to AU$2.0 billion (AU$1.7 billion–AU$2.2 billion) compared to the full-cost

scenario for *Felixers™* (Supplementary Material, Appendix E, Fig. S5). Both scenarios

(0.1N_1 and 0.01N_1) predictably reduce the cost of the culling scenario to ~ 30% of the full-

cost scenarios.

Discussion

In each cull scenario we considered, a successful reduction of the feral cat population on

Kangaroo Island to below 0.01 of its initial size by 2030 is achievable, but the minimum

costs involved according to the different scenarios we ran could range from AU$46.5 million

(AU$10 ha^{-1}, with hunting; Fig. 5c) to AU$18 billion (AU$40,863 ha^{-1}, with *Felixers™*; Fig.

5a) depending on the method used and the inherent uncertainty in the parameters we

estimated. Simulating a spatial roll-out where one-third of the additional *Felixers™* are

required brings the maximum down to AU$3.3 billion (AU$3.0 billion – AU$3.7 billion;

0.01N_1 *Felixer*-shortfall scenario) (Supplementary Material, Appendix E, Fig. S5). Whether

these costs are a true reflection of the actual costs depends on the accurate form of the (as-yet

unmeasured) functional response, and the assumed per-unit outlay of the different methods to

meet the annual shortfall for the different intensities of cull. However, the lower end of our

estimates for Kangaroo Island are cheaper than the costs to remove cats from Macquarie

Island (AU$258 ha^{-1}), likely due to the latter's remoteness (Robinson and Copson 2014).

Cost estimates from other cat eradications on islands (e.g., Marion Island) are currently not

available. Nonetheless, our comparisons do illustrate that high initial culls (> 0.45) followed

by moderate maintenance culls (0.20–0.45) would be sufficient to reduce the population to

0.01–0.10 of its original size (Fig. 4), and that hunting is the most cost-effective way to meet

these targets (especially if other methods are rolled out simultaneously).

That hunting is cheaper than other methods is unsurprising given that no hardware other

than rifles and ammunition are needed to be purchased outright, in contrast to the higher

overheads associated with traps or *Felixer™* units (Holmes et al. 2015, Hodgens 2019).

Indeed, hunting requires many people working full time, whereas the other techniques are

more passive (yet the latter also require set up, monitoring, maintenance, displacement, and

removal by staff). However, approximately 500 person hours are required to equate to the

current cost of a single *Felixer™* unit (hunting ~ AU\$26 per person hour^{-1} vs. AU\$13,000

Felixer™ unit), although assuming a spatial roll-out of fewer units is functionally identical to

a similar reduction in per-unit cost. Additionally, hunting is considered more humane due to

the minimised contact with the animal and the instant death with a correctly executed

headshot (Sharp and Saunders 2011), but access to private land and potential conflict with

private landholders could complicate hunting in non-economic ways. Of course, complete

eradication would necessarily entail additional costs as the final individuals were identified,

hunted, and destroyed (Bester et al. 2002, Nogales et al. 2004). This would be more

expensive than normal eradication efforts and would require a strategic isolation of remaining

individuals, because even a few individuals remaining could conceivable seed a recovery

population that could achieve initial population size in several decades (Supplementary

Material, Appendix D).

Our results identify that fertility-reduction using trap-neuter-release methods are

comparatively ineffective for reducing pest densities (Longcore et al. 2009). Instead, our

model outputs show that a realised annual fertility equivalent to ≤ 25% of expected values

(i.e., pre-fertility treatment) would reduce the population to ≤ 10%. In other words, managers

would have to maintain at least 75% of the female cat population sterilised each year from

2019–2030 for the program to reduce the population to ≤ 10% of its original size. Whether

this is feasible or cost-effective is beyond the scope of our study, but it does demonstrate that

fertility-reduction is a much less efficient method to eradicate cats than culling. Additionally,

sterilised individuals returned to the population could still continue to eat native fauna until

they perished due to natural causes, so the overall ecological impact would be necessarily

greater than a culling-based program where individuals are removed permanently when first

treated.

We conclude that the most appropriate approach to reduce cat densities on Kangaroo

Island is a two-stage method, with a high initial reduction of at least 0.6—0.7 and a

maintenance cull of 0.45—0.65. Although a constant proportional annual cull can be

effective, it is generally less efficient than a two-stage approach, especially if some leakage

of domestic cats occurs. Indeed, our models shows that even as many as 20 cats released into

the population per year would see the final proportional population size increase from 0.005

to 0.015 at the end of the management programme (or, from 20 to ~ 45 individuals;

Supplementary Material, Appendix A). This is because effort is spread equally among

temporal windows in the constant proportional scenario, and therefore must 'catch up'

relative to a large, initial cull given that more surviving individuals are still breeding in the

former. As culling reduces density and drives the population closer to extinction, it becomes

progressively more difficult and expensive to cull remaining individuals (Nogales et al. 2004,

Parkes et al. 2014). This is because most culling methods are passive and rely on a 'non-

negligible probability' of the target animal encountering *Felixers™* or traps (Moseby and Hill

2011). Although these techniques can be accompanied by visual, scent or sound lures, the

target animal still needs to be in range to be enticed by them. Thus, encounters at low

densities become increasingly less likely (Veitch 2001, Campbell et al. 2011), and rising per-

capita food abundance as the predator's population dwindles can make baits or food lures less

attractive (Parkes et al. 2014). Cats in particular are intelligent and can learn to avoid traps.

Thus, while trapping is generally considered effective for density reduction (Nogales et al.

2004), it is most effective in the early stages of eradication programs (Nogales et al. 2004).

Therefore, a two-stage approach allows for the implementation of widespread control that is

effective at high densities, followed by a more targeted approach through consecutive

maintenance as the population continues to decline.

Although our model is focused on Kangaroo Island, the merits of the stochastic framework

we developed imply that the model is transferable to other regions and even other species.

Altering locally measured demographic rates, population sizes, and reduction targets are

feasible with this approach. Our model therefore has applications for other species, including

European red foxes (*Vulpes vulpes*) (Edwards et al. 2004) in Australia, brush-tail possums

(*Trichosurus vulpecula*) and stoats (*Mustela erminea*) in New Zealand (Brown et al. 2015),

and racoons (*Procyon lotor*) in central Europe (Beltrán-Beck et al. 2012).

Because the Kangaroo Island trial eradication program is still in its initial stage, many of

the demographic, effectiveness, and cost parameters are not yet available specifically for this

region, lending further, but as yet unquantifiable, uncertainty to our conclusions. For

example, we were obliged to obtain demographic rates from data published for cat

populations elsewhere (Budke and Slater 2009), and we do not yet have access to spatially

variable data with which to parameterise spatially explicit models that could potentially

improve realism (e.g., such as variation in demographic rates among habitat types — see the

following *Perspective* section) (McMahon et al. 2010). A better quantification of carrying

capacity, as well as the mechanisms by which density feedback operates on the demographic

rates would also improve predictive accuracy.

Many attempts to eradicate cats have been made, but they vary widely in success (Algar et

al. 2002, Bester et al. 2002, Nogales et al. 2004, Algar et al. 2011, Campbell et al. 2011,

Parkes et al. 2014). For effective eradication to be achieved, culling programs must be based

on empirical data and ideally, directed by models like ours. Our model should allow

practitioners to make their culling programs more efficient, and to allocate the resources

needed to achieve their targets efficiently and cost-effectively. As more site-specific data

become available, we expect the model's predictions to become ever-more realistic to

identify the most plausible and cheapest pathways to eradication.

Perspective on the Kangaroo Island cat-eradication program

Poison baits (e.g., 1080 or PAPP) were not initially considered as part of the feral cat control

on Kangaroo Island because of the perceived risk posed to native and domestic animals

(Berris et al. 2019). However, we have confirmed that poison baits will now be used in

concert with the other proposed methods, subject to land-owner consent (Natural Resource

Kangaroo Island 2020). We omitted the use of baits in our model because there are

insufficient data for the use and roll-out in the eradication program. That said, when more

data become available, baits can be included into the model framework, which will likely

decrease the cost of the other control methods we specifically modelled (Nogales et al. 2004).

Due to insufficient data for many model parameters and functions, we were obliged either

to make (arguably defensible) assumptions or use data from other locations/studies (Budke

and Slater 2009). Although we are confident that our results and scenarios are relevant, they

will be undoubtedly be improved by the refinement of locally measured parameters such as

age-specific demographic rates, updated density estimates following the 2020 bushfires, cost

data, efficiency relationships, strength of compensatory density feedback, and probability of

leakage. A more detailed schedule for resource application including budget restrictions,

timeline flexibility, currently available resources (to reduce initial costs via unit purchasing)

and available staff would also assist in improving predicted scenario realism. Further, the

Felixer™ is still in the initial phases of production and is not yet produced on a large

commercial scale. The *Felixer™* has many merits in regards to feral cat control because of its

hazard reduction for baiting non-target species (Read et al. 2019); thus, it is likely to be

increasingly applied in future management projects, especially as costs per unit decline.

References

Abbott, I. 2002. Origin and spread of the cat, *Felis catus*, on mainland Australia, with a
discussion of the magnitude of its early impact on native fauna. Wildlife Research
29:51-74.10.1071/WR01011
Abbott, I. 2008. The spread of the cat, *Felis catus*, in Australia: re-examination of the current
conceptual model with additional information. Conservation Science Western
Australia **7**:1-17
Algar, D., G. Angus, and M. Onus. 2011. Eradication of feral cats on Rottnest Island,
Western Australia. Journal of the Royal Society of Western Australia **94**:439-443
Algar, D., A. Burbidge, and G. Angus. 2002. Cat eradication on Hermite Island, Montebello
Islands, Western Australia. Turning the tide: the eradication of invasive species:14-18
Andersen, M. C., B. J. Martin, and G. W. Roemer. 2004. Use of matrix population models to
estimate the efficacy of euthanasia versus trap-neuter-return for management of free-
roaming cats. Journal of the American Veterinary Medical Association **225**:1871-
1876.10.2460/javma.2004.225.1871
Barratt, D. G. 1997. Predation by house cats, *Felis catus* (L.), in Canberra, Australia. I. prey
composition and preference. Wildlife Research **24**:263-277.10.1071/WR96020
Beltrán-Beck, B., F. J. García, and C. Gortázar. 2012. Raccoons in Europe: disease hazards
due to the establishment of an invasive species. European Journal of Wildlife
Research **58**:5-15
Berris, K., J. Mulvaney, B. Florence, P. Hodgens, and V. Bolwell. 2019. Felixer versus felis
project; community engagement report., Natural Resources Kangaroo Island,
Kangaroo Island, SA
Bester, M., J. Bloomer, R. Van Aarde, B. Erasmus, P. Van Rensburg, J. Skinner, P. Howell,
and T. Naude. 2002. A review of the successful eradication of feral cats from sub-
Antarctic Marion Island, Southern Indian Ocean. South African Journal of Wildlife
Research **32**:65-73
Bloomer, J. P., and M. N. Bester. 1991. Effects of hunting on population characteristics of
feral cats on Marion Island. South African Journal of Wildlife Research **21**:97-102
Bloomer, J. P., and M. N. Bester. 1992. Control of feral cats on sub-Antarctic Marion Island,
Indian Ocean. Biological Conservation **60**:211-219.10.1016/0006-3207(92)91253-O
Bonnaud, E., F. M. Medina, E. Vidal, M. Nogales, B. Tershy, E. Zavaleta, C. J. Donlan, B.
Keitt, M. Le Corre, and S. V. Horwath. 2011. The diet of feral cats on islands: a
review and a call for more studies. Biological Invasions **13**:581-603.10.1007/s10530-
010-9851-3

Brown, K., G. Elliott, J. Innes, and J. Kemp. 2015. Ship rat, stoat and possum control on
mainland New Zealand. An overview of techniques, successes and challenges.
Wellington, Department of Conservation
Budke, C. M., and M. R. Slater. 2009. Utilization of matrix population models to assess a 3-
year single treatment nonsurgical contraception program versus surgical sterilization
in feral cat populations. Journal of Applied Animal Welfare Science **12**:277-
292.10.1080/10888700903163419
Campbell, K., G. Harper, D. Algar, C. Hanson, B. Keitt, and S. Robinson. 2011. Review of
feral cat eradications on islands. Island Invasives: Eradication and Management:37-46
Caswell, H. 2001. Age-Classified Matrix Models. Matrix Population Models, constuction,
analysis, and interpretation. Sinauer Associates, Inc., Sunderland, Massachusetts
Close, R. 2005. Introduced Mammals of the World: Their History, Distribution and
Influence. John Long. A Review by Robert L Close. Australian Mammalogy **27**:109-
110.10.1071/AM05109_BR
Courchamp, F., and G. Sugihara. 1999. Modeling the biological control of an alien predator
to pretect island species from extinciton. Ecological Applications **9**:112-
123.10.1890/1051-0761
Deak, B., B. Ostendorf, D. Taggart, D. E. Peacock, and D. Bardsley. 2019. The significance
of social perceptions in implementing successful feral cat management strategies: a
global review. Animals **9**.10.3390/ani9090617
Denno, R. F., and D. Lewis. 2009. Predator-Prey Interactions. Pages 202-212 *in* S. Levin A,
editor. The Princeton Guide to Ecology. Princeton University Press, Princeton, New
Jersey
Denny, E. A., and C. Dickman. 2010. Review of cat ecology and management strategies in
Australia. Invasive Animals Cooperative Research Centre, Canberra
Dickman, C. R. 1996. Impact of exotic generalist predators on the native fauna of Australia.
Wildlife Biology **2**:185-195.10.2981/wlb.1996.018
Doherty, T., C. Dickman, C. Johnson, S. Legge, E. Ritchie, and J. Woinarski. 2016. Impacts
and management of feral cats *Felis catus* in Australia. Mammal Review **47**:83-
97.10.1111/mam.12080
Driscoll, C. A., M. Menotti-Raymond, A. L. Roca, K. Hupe, W. E. Johnson, E. Geffen, E. H.
Harley, M. Delibes, D. Pontier, A. C. Kitchener, N. Yamaguchi, S. J. O'Brien, and D.
W. Macdonald. 2007. The Near Eastern Origin of Cat Domestication. Science
317:519-523.10.1126/science.1139518
Edwards, G., A. Pople, K. Saalfeld, and P. Caley. 2004. Introduced mammals in Australian
rangelands: Future threats and the role of monitoring programmes in management
strategies. Austral Ecology **29**:40-50.10.1111/j.1442-9993.2004.01361.x
Fitzgerald, B., B. Karl, and C. Veitch. 1991. The diet of feral cats (*Felis catus*) on Raoul
Island, Kermadec Group. New Zealand Journal of Ecology:123-129
Fitzgerald, B. M., and D. C. Turner. 2000. Hunting behaviour of domestic cats and their
impact on prey populations. The domestic cat: the biology of its behaviour **151**
Gibson, K. L., K. Keizer, and C. Golding. 2002. A trap, neuter, and release program for feral
cats on Prince Edward Island. The Canadian Veterinary Journal **43**:695-692
Higgins-Desbiolles, F. 2011. Death by a thousand cuts: governance and environmental trade-
offs in ecotourism development at Kangaroo Island, South Australia. Journal of
Sustainable Tourism **19**:553-570.10.1080/09669582.2011.560942
Hodgens, P. 2019. Felixer V's Felis Report. Kangaroo Island
Hohnen, R., K. Berris, P. Hodgens, J. Mulvaney, B. Florence, B. P. Murphy, S. Legge, C. R.
Dickman, and J. C. Z. Woinarski. 2020. Feral cat density and population size across
Kangaroo Island, South Australia 10.1071/WR19137:10.1071/WR19137

Holmes, N., K. Campbell, B. Keitt, R. Griffiths, J. Beek, C. Donlan, and K. Broome. 2015.
Reporting costs for invasive vertebrate eradications. Biological Invasions **17**:2913-
2925

Jones, E., and B. Coman. 1982. Ecology of the feral cat, *Felis catus* (L.), in south-eastern
Australia III. Home ranges and population ecology in semiarid north-west Victoria.
Wildlife Research **9**:409-420.10.1071/WR9820409

Longcore, T., C. Rich, and L. M. Sullivan. 2009. Critical assessment of claims regarding
management of feral cats by trap–neuter–return. Conservation Biology **23**:887-
894.10.1111/j.1523-1739.2009.01174.x

Lowe, S., M. Browne, S. Boudjelas, and M. De Poorter. 2000. 100 of the world's worst
invasive alien species: a selection from the global invasive species database. Invasive
Species Specialist Group Auckland

Masters, P., T. Duka, S. Berris, and G. Moss. 2004. Koalas on Kangaroo Island: from
introduction to pest status in less than a century. Wildlife Research **31**:267-
272.10.1071/WR03007

McCarthy, R. J., S. H. Levine, and J. M. Reed. 2013. Estimation of effectiveness of three
methods of feral cat population control by use of a simulation model. Journal of the
American Veterinary Medical Association **243**:502-511.10.2460/javma.243.4.502

McMahon, C., B. Brook, N. Collier, and C. Bradshaw. 2010. Spatially explicit spreadsheet
modelling for optimising the efficiency of reducing invasive animal density. Methods
in Ecology and Evolution **1**:53-68.10.1111/j.2041-210X.2009.00002.x

Medina, F. M., E. Bonnaud, E. Vidal, and M. Nogales. 2014. Underlying impacts of invasive
cats on islands: not only a question of predation. Biodiversity and Conservation
23:327-342.10.1007/s10531-013-0603-4

Medina, F. M., E. Bonnaud, E. Vidal, B. R. Tershy, E. S. Zavaleta, C. J. Donlan, B. S. Keitt,
M. Corre, S. V. Horwath, and M. Nogales. 2011. A global review of the impacts of
invasive cats on island endangered vertebrates. Global Change Biology **17**:3503-
3510.10.1111/j.1365-2486.2011.02464.x

Medina, F. M., M. López-Darias, M. Nogales, and R. García. 2008. Food habits of feral cats
(*Felis silvestris catus* L.) in insular semiarid environments (Fuerteventura, Canary
Islands). Wildlife Research **35**:162-169.10.1071/WR07108

Miller, P. S., J. D. Boone, J. R. Briggs, D. F. Lawler, J. K. Levy, F. B. Nutter, M. Slater, and
S. Zawistowski. 2014. Simulating free-roaming cat population management options in
open demographic environments. PLOS ONE
9:e113553.10.1371/journal.pone.0113553

Moseby, K. E., and B. M. Hill. 2011. The use of poison baits to control feral cats and red
foxes in arid South Australia I. Aerial baiting trials. Wildlife Research **38**:338-
349.10.1071/WR10235

Moseby, K. E., D. E. Peacock, and J. L. Read. 2015. Catastrophic cat predation: A call for
predator profiling in wildlife protection programs. Biological Conservation **191**:331-
340.10.1016/j.biocon.2015.07.026

Natural Resource Kangaroo Island. 2015. Feral cat eradicaiton on Kangaroo Island 2015-
2030 PROSPECTUS. Kangaroo Island

Natural Resource Kangaroo Island. 2020. Dudley Peninsula feral cat eradication operations
plan: summary may 2020 - mid 2023.

Nogales, M., A. Martín, B. R. Tershy, C. J. Donlan, D. Veitch, N. Puerta, B. Wood, and J.
Alonso. 2004. A review of feral cat eradication on islands. Conservation Biology
18:310-319.10.1111/j.1523-1739.2004.00442.x

Nunney, L. 1980. The influence of the type 3 (sigmoid) functional response upon the stability
of predator-prey difference models. Theoretical Population Biology **18**:257-
278.10.1016/0040-5809(80)90052-0

Nutter, F. B. 2006. Evaluation of a trap-neuter-return management program for feral cat
colonies: population dynamics, home ranges, and potentially zoonotic diseases. North
Carolina State University, North Carolina

Parkes, J., P. Fisher, S. Robinson, and A. Aguirre-Muñoz. 2014. Eradication of feral cats
from large islands: an assessment of the effort required for success. New Zealand
Journal of Ecology:307-314

Read, J., and Z. Bowen. 2001. Population dynamics, diet and aspects of the biology of feral
cats and foxes in arid South Australia. Wildlife Research **28**:195-
203.10.1071/WR99065

Read, J. L., T. Bowden, P. Hodgens, M. Hess, H. McGregor, and K. Moseby. 2019. Target
specificity of the felixer grooming "trap". Wildlife Society Bulletin **43**:112-
120.10.1002/wsb.942

Robertson, S. A. 2008. A review of feral cat control. Journal of Feline Medicine and Surgery
10:366-375.10.1016/j.jfms.2007.08.003

Robinson, S. A., and G. R. Copson. 2014. Eradication of cats (*Felis catus*) from subantarctic
Macquarie Island. Ecological Management & Restoration **15**:34-
40.10.1111/emr.12073

Schmidt, P. M., T. M. Swannack, R. R. Lopez, and M. R. Slater. 2009. Evaluation of
euthanasia and trap-neuter-return (TNR) programs in managing free-roaming cat
populations. Wildlife Research **36**:117-125.10.1071/WR08018

Sharp, T., and G. Saunders. 2011. A model for assessing the relative humaneness of pest
animal control methods. Department of Agriculture, Fisheries and Forestry Canberra,
Australia

Short, J., and B. Turner. 2005. Control of feral cats for nature conservation. IV. Population
dynamics and morphological attributes of feral cats at Shark Bay, Western Australia.
Wildlife Research **32**:489-501.10.1071/WR04102

Smith, G. C., H. I. S., and P. A. Robertson. 2005. A model of ruddy duck Oxyura jamaicensis
eradication for the UK. Journal of Applied Ecology **42**:546-555.doi:10.1111/j.1365-
2664.2005.00977.x

Taggart, P. L., B. A. Fancourt, A. J. Bengsen, D. E. Peacock, P. Hodgens, J. L. Read, M. M.
McAllister, and C. G. Caraguel. 2019. Evidence of significantly higher island feral cat
abundance compared with the adjacent mainland. Wildlife Research **46**:378-
385.10.1071/wr18118

Van Aarde, R. 1983. Demographic parameters of the feral cat Felis catus population at
Marion Island. South African Journal of Wildlife Research **13**:12-16

Veitch, C. 2001. The eradication of feral cats (*Felis catus*) from little Barrier Island, New
Zealand. New Zealand journal of zoology **28**:1-12.10.1080/03014223.2001.9518252

Wallace, J. L., and J. K. Levy. 2006. Population characteristics of feral cats admitted to seven
trap-neuter-return programs in the United States. Journal of Feline Medicine and
Surgery **8**:279-284.10.1016/j.jfms.2006.02.004

Walsh, A., H. Wilson, H. Bower, P. McClelland, and J. Pearson. 2019. Winning the hearts
and minds–proceeding to implementation of the Lord Howe Island rodent eradication
project: a case study. Island Invasives: Scaling up to Meet the Challenge'.(Eds CR
Veitch, MN Clout, AR Martin, JC Russell and CJ West.) pp:522-530

Willoughby, N., D. Thompson, M. Royal, and M. Miles. 2018. South Australian land cover
layers: an introduction and summary statistics, DEW technical report 2018/01.

671   Technical Report, Government of South Australia, Department for Environment and
672    Water, Adelaide
673 Woinarski, J. C. Z., A. A. Burbidge, and P. L. Harrison. 2015. Ongoing unraveling of a
674    continental fauna: Decline and extinction of Australian mammals since European
675    settlement. Proceedings of the National Academy of Sciences **112**:4531-
676    4540.10.1073/pnas.1417301112
677
678

Table 1 – Mean parameter values and their standard deviations (SD) used in the stochastic model.

parameter	mean	SD
fertility (daughters)		
pre-breeding* juvenile (f_1)	0.000	—
sub-adult (f_2)	0.745	0.307
adult (f_3–f_7)	2.520	0.450
survival		
pre-breeding juvenile (s_1)	0.460	0.115
sub-adult (s_2)	0.460	0.115
adult (s_3–s_6)	0.700	0.058
senescent (s_7)	0.550	0.058

*pre-breeding juvenile: < 10 months old; not sexually mature

Figure Captions

Figure 1 – Map of Kangaroo island relative to the Australian mainland. The shortest distance from the mainland (southern tip of Fleurieu Peninsula) to Kangaroo Island is approximately 14 km.

Figure 2 – (a) Average proportion of the initial cat population (N_1) on Kangaroo Island projected from 2020–2030 for the unculled scenario. Black line indicates the median value from 10,000 iterations, along with 95% confidence intervals (grey-shaded area). (b) Minimum proportion of the Kangaroo Island feral cat population remaining after a constant proportional annual cull ranging from 0.5 to 0.9. Solid black line represents median minimum proportion of the initial population (N_1) after 10,000 iterations with 95% confidence intervals indicated as grey-shaded area.

Figure 3 – Estimated median minimum proportion of founding population remaining (founding population N_1), with decreasing fertility (fertility reduced 50–99%). Solid black line represents median minimum proportion of the initial population (N_1) remaining after fertility reduction scenarios, with 95% confidence intervals indicated as grey-shaded area.

Figure 4 –Estimated median minimum proportion of the final population remaining (relative to start population N_1) for combinations of initial proportional (i.e., *initial cull*: 0.5–0.9) and maintenance proportional (i.e., *maintenance cull*: 0.1–0.5) culling. Proportion of population remaining after culling scenarios represented by colour bar ranging from lowest (purple) to highest (yellow) remaining proportional population.

Figure 5 – Estimated median total costs of feral cat eradication on Kangaroo Island for combinations of initial proportional (i.e., *initial cull*) and maintenance proportional (i.e., *maintenance cull*) culling, where the shortfall in the number of cats killed from *Felixer*™ units and traps is provided by (a) additional *Felixer*™ units, (b) traps, or (c) hunting. Cost of eradication (in AU$, adjusted for 2020) indicated by colour bar ranging from lowest (purple) to highest (yellow) costs. Contours and white values indicate cost in $AU millions or billions accordingly. Note different *z*-axis (contour) scales in a, b, and c.

Figure 1

Figure 2

679 **Figure 3**

Figure 4

Figure 5

SUPPLEMENTARY MATERIAL

Predicting targets and costs for feral-cat reduction on large islands using

stochastic population models

Kathryn R. W. Venning, Frédérik Saltré, Corey J. A. Bradshaw

The following supplementary material includes figures, methods, results and further

contextualisation for information given in the main text.

Table of contents

Appendix A – Leakage into the population

Background

The opportunities for immigration of cats to this island population are low because of strict

controls on cat ownership and importation. However, accidental or deliberate releases, or

domestic cats breeding with feral animals, are still plausible. All domestic cats on Kangaroo

Island must be de-sexed, registered and microchipped by law (Natural Resource Kangaroo

Island 2015), which greatly reduces the possibility of a domestic-feral breeding. Nonetheless,

we considered a hypothetical scenario of 'leakage' of domestic cats into the population on

Kangaroo Island to explore this phenomenon's effects on eradication efficiency.

Approach

We used the optimal-culling scenario (two-phase; initial 0.6, maintenance 0.45) to identify

how leakage would affect the population at the end of the culling scenario. To achieve this,

we added an incrementing number of individuals (between 10 and 100, in intervals of 10)

annually to the population vector, while simultaneously running the culling scenario. We ran

each leakage scenario for 10,000 iterations (randomly sampling 10,000 times from the

stochastic survival and fertility vectors) to calculate the mean and 95% confidence limits for

minimum population size.

Results

If no cats are leaked into the Kangaroo Island population (i.e., baseline expectation), the final

population at the completion of culling would be 16 cats (range: 9–27, 95% confidence

intervals). This would increase by a factor of 1.8 (i.e., 29 cats; range: 21–41) if 10 cats were

added annually (Fig. S1). If 50 cats were added annually, the final population would increase

to 93 cats (79–111) (Fig. S1).

Figure S1. Final population (*N*) at completion of the ideal culling scenario (two-phase; initial 0.60, maintenance
0.45), adding 10–100 cats annually into the population. Solid black line represents mean population after
stochastic resampling. Grey shaded area represents 95% confidence bounds for minimum population size.

Implications

There are currently 244 registered domestic cats on Kangaroo Island, including a single

breeder cat (presuming all cat owners follow the by-laws, this equates to only one domestic

cat able to reproduce; Natural Resource Kangaroo Island 2015). It is therefore highly unlikely

that 100 cats are leaked into the population annually. However, it is possible that a few cats

(1–5) are accidently released annually. Likewise, it is possible some cat owners do not

register their pets or accommodate 'stray' cats (provide food and/or water, but no direct

ownership). Based on our simulation, leakage to this extent is unlikely to reduce overall

eradication efficiency, particularly if owners abide by the by-laws, and with > 90% public

approval for eradication (Natural Resource Kangaroo Island 2015) compliance with existing

by-laws will be high. Regardless, even an annual leakage of 10 cats could increase the final

population after the culling scenario to 1.8 times compared to the baseline scenario (Fig. S1).

As such, wildlife practitioners and council members should continue community engagement

and to enforce cat ownership by-laws.

Appendix B – Survival modifier (compensatory density-feedback mechanism)

Figure S2. Reduction in probability of survival relative to population size (N), with carry capacity (K) set to
twice the size of the founding population (6,166). κ, τ and θ are constants ($\kappa = 0.998$, $\tau = 9139.109$ and $\theta =$
2.835).

$$S_{\text{mod}} = \frac{\kappa}{1 + \left(\dfrac{N}{\tau}\right)^{\theta}}$$

74
75

Appendix C – Type III functional response

To force the model to adjust the efficacy of the different culling methods according to the number of individuals in the population, we constructed an arbitrary Type III functional response of the form:

$$f = \frac{\alpha}{1 + (\beta e^{-\gamma pN})}$$

where f is the relative efficiency of the culling technique, α, β and γ are constants: $\alpha = 1.01$, $\beta = 85.61$, and $\gamma = 8.86$, and pN = proportional population size (Fig. S3). There is no empirical basis for these particular parameter values; instead, we wanted to demonstrate a generic sigmoidal function to alter the efficiency of each culling method.

Figure S3. Reduction in the efficiency of a culling method (f) as the proportion of the remaining population size declines (pN). Here, α, β and γ are constants ($\alpha = 1.01$, $\beta = 85.61$ and $\gamma = 8.86$).

Appendix D – Stopping culling prematurely

Background

Following eradications to completion are essential to achieve density-reduction or eradication goals, but due to funding limitations or unforeseen circumstances, such programs are sometimes cancelled or stopped prematurely (Keitt et al. 2011, Capizzi 2020). Depending on at what stage of eradication a program is stopped, the remaining population can rebuild to its pre-management densities or beyond (Armstrong et al. 2001, Bradshaw et al. 2007).

Approach

To examine how quickly the Kangaroo Island cat population would recover to initial population size ($N_1 = 3083$) relative to variation in the premature stopping of the culling program, we stopped the optimal management cull (0.45 proportionate annual cull) at yearly values following the initial cull (0.60 proportionate cull for first two years). We then allowed the population to rebuild using the same methods as in the untreated population (see main text *Methods*). We then recorded the time it took for the rebuilding population to equate to N_1. We ran each stoppage scenario for 10,000 iterations (randomly sampling 10,000 times from the stochastic survival and fertility vectors) to calculate the mean and 95% confidence limits for population size.

Results

Without a maintenance cull (i.e., only two years with 0.6 proportional cull applied), the population recovers to its initial size in 15 years (range: 11–21 years) (Fig. S4). Stopping culling prior to the end of the targeted duration of the 10-year culling program (i.e., 9[th] year) results in population recovery to initial size in 42 years (range: 35–50 years) (Fig. S4).

116 **Figure S4**. Years required for population to rebuild to founding population size (N_1) when the maintenance cull
117 is stopped prematurely (year maintenance cull stopped). Initial culling of 0.60 for two years is followed by
118 maintenance culling (0.45) for the years indicated. Solid black line represents the median time to rebuild to
119 founding population (years), and the grey area represents the 95% confidence interval.
120

122   **Appendix E – Reduced initial cost of *Felixer*™ units**

123
124   **Figure S5**. Estimated median total cost of feral cat eradication on Kangaroo Island for combinations of two-
125   phase cull (high initial proportional cull followed by steady maintenance proportional cull), where shortfall is
126   made up by *Felixer*™ units. Here, we reduced the initial cost of *Felixer*™ units by 2/3 to simulated resource
127   sharing and/or reduced initial costs. Costs of eradication (in AU$, adjusted for 2020) indicated by colour bar
128   ranging from lowest (purple) to highest (yellow). Contours and white values indicate cost in $AU billions.

References

Armstrong, D. P., J. K. Perrott, and I. Castro. 2001. Estimating impacts of poison operations using mark-recapture analysis: hihi (Notiomystis cincta) on Mokoia Island. New Zealand Journal of Ecology:49-54

Bradshaw, C. J. A., I. C. Field, D. M. J. S. Bowman, C. Haynes, and B. W. Brook. 2007. Current and future threats from non-indigenous animal species in northern Australia: a spotlight on World Heritage Area Kakadu National Park. Wildlife Research **34**:419-436.WR06056

Capizzi, D. 2020. A review of mammal eradications on Mediterranean islands. Mammal Review **50**:124-135.10.1111/mam.12190

Keitt, B., K. Campbell, A. Saunders, M. Clout, Y. Wang, R. Heinz, K. Newton, and B. Tershy. 2011. The global islands invasive vertebrate eradication database: a tool to improve and facilitate restoration of island ecosystems. Island invasives: eradication and management. IUCN, Gland, Switzerland:74-77.10.1071/WR06056

Natural Resource Kangaroo Island. 2015. Feral cat eradicaiton on Kangaroo Island 2015-2030 PROSPECTUS. Kangaroo Island