PSEUDOMONAS AERUGINOSA VOLATILOME CHARACTERISTICS AND ADAPTATIONS

IN CHRONIC CYSTIC FIBROSIS LUNG INFECTIONS

Trenton J. Davis[a,b], Ava V. Karanjia[a,c], Charity N. Bhebhe[a], Sarah B. West[a], Matthew

Richardson[d,e], and Heather D. Bean[a,b,#]

[a]School of Life Sciences, Arizona State University, Tempe, AZ 85287, USA

[b]Center for Fundamental and Applied Microbiomics, The Biodesign Institute, Tempe, AZ, USA

[c]School of Engineering of Matter, Transport, and Energy, Arizona State University, Tempe, AZ,

USA

[d]Department of Respiratory Sciences, College of Life Sciences, University of Leicester,

Leicester, UK

[e]NIHR Biomedical Research Centre (Respiratory theme), Institute for Lung Health, Leicester,

UK

[#]Corresponding author: Heather D. Bean

17   Heather.D.Bean@asu.edu

ABSTRACT

Pseudomonas aeruginosa chronic lung infections in individuals with cystic fibrosis (CF) significantly reduce quality of life and increase morbidity and mortality. Tracking these infections is critical for monitoring patient health and informing treatments. We are working toward the development of novel breath-based biomarkers to track chronic *P. aeruginosa* lung infections *in situ*. Using comprehensive two-dimensional gas chromatography coupled to time-of-flight mass spectrometry (GC×GC-TOFMS), we characterized the *in vitro* volatile metabolomes (or volatilomes) of 81 *P. aeruginosa* isolates collected from 17 CF patients over at least a five-year period of their chronic lung infections. We detected 539 volatiles produced by the *P. aeruginosa* isolates, 69 of which were core volatiles that were highly conserved. We found that each early infection isolate has a unique volatilome, and as infection progresses, the volatilomes of isolates from the same patient become increasingly dissimilar, to the point that these intra-patient isolates are no more similar to one another than to isolates from other patients. We observed that the size and chemical diversity of *P. aeruginosa* volatilomes do not change over the course of chronic infections; however, the relative abundances of core hydrocarbons, alcohols, and aldehydes do change, and are correlated to changes in phenotypes associated with chronic infections. This study indicates that it may be feasible to track *P. aeruginosa* chronic lung infections by measuring changes to the infection volatilome, and lays the groundwork for exploring the translatability of this approach to direct measurement using patient breath.

IMPORTANCE

Pseudomonas aeruginosa is a leading cause of chronic lung infections in cystic fibrosis (CF), and are correlated with lung function declines. Significant clinical efforts are, therefore, aimed at detecting infections and tracking them for phenotypic changes, such as mucoidy and antibiotic resistance. Both the detection and tracking of lung infections relies on sputum cultures, but due to improvements in CF therapies, sputum production is declining though risks for lung infections

persist. Therefore, we are working toward the development of breath-based diagnostics for CF

lung infections. In this study we characterized of the volatile metabolomes of 81 *P. aeruginosa*

clinical isolates collected from 17 CF patients over a duration of at least five years of a chronic

lung infection. We found that the volatilome of *P. aeruginosa* adapts over time, and correlates to

infection phenotype changes, suggesting it may be possible to track chronic CF lung infections

with a breath test.

KEYWORDS

Pseudomonas aeruginosa; cystic fibrosis; metabolomics; volatile organic compounds; chronic

infection; adaptation

RUNNING TITLE

P. aeruginosa cystic fibrosis volatilome

INTRODUCTION

Cystic fibrosis (CF) is an autosomal recessive disease caused by a mutation in the CFTR protein that regulates ion transport across epithelia. In the lungs, reduced or lost CFTR function leads to defective mucociliary transport, facilitating infection and colonization by a plethora of microorganisms (1). *Pseudomonas aeruginosa* is one of the most prevalent lung pathogens in CF – especially after adolescence (2) – and is able to establish chronic infections that can last for years to decades (3, 4). *P. aeruginosa* lung infection is associated with more rapid lung function decline, increased risk of hospitalization, and increased risk of death (5, 6). Significant clinical efforts are, therefore, aimed at diagnosing and treating new infections to delay the onset of chronic infection (7).

As *P. aeruginosa* transitions from an acute infection to chronicity, it undergoes a variety of genotypic, phenotypic, and metabolic changes (8, 9). It has been well-established that several phenotypes are correlated with chronic infection, including reduced motility and increased mucoidy, antibiotic resistance, and biofilm formation (9-11), and some of these phenotypes are also correlated with poorer patient outcomes (6, 12, 13). In light of this, *P. aeruginosa* phenotypes could be used to evaluate infection stage, CF disease progression, and patient morbidity risk (10, 12, 14, 15). Poor access to infection sites in the lower airways, however, delays the detection of new *P. aeruginosa* infections and reduces the feasibility and accuracy of disease state tracking via bacterial phenotypes, whether by culture-dependent or culture-independent methods (16).

We are developing breath-based diagnostics to detect new *P. aeruginosa* infections and track chronic infection phenotypes in the CF lung. Breath consists of thousands of volatile organic compounds containing a wealth of information about human health and disease (17, 18), and is being leveraged for the development of diagnostics for wide-ranging conditions (19, 20). In the context of chronic lung infections, breath-based diagnostics possess significant advantages over other diagnostic modalities. First, breath provides a non-invasive way of

sampling the entire ventilated lung, a limitation of both sputum and bronchoalveolar lavage fluid

that leads to delayed diagnosis of new infections, or the detection of infection phenotype

changes (16, 21-24). Second, breath sampling captures metabolic information about the

pathogens *in situ*, eliminating the need for microbial enrichment steps and speeding the time-to-

diagnosis by days to weeks (25, 26). Additionally, *in situ* metabolic measurements provide

information on the physiology of the pathogens in the context of disease versus the context of

the lab. We propose that a breath-based diagnostic can be created for detecting and tracking

CF *P. aeruginosa* lung infections by correlating the microbial genotypic and phenotypic changes

that occur during chronic infections to changes in the infection volatile metabolome, or

"volatilome".

In order to develop a set of biomarkers for diagnosing and tracking *P. aeruginosa* lung

infections in CF, we must consider the genomic, phenomic, and volatilomic diversity of the

species, and how that diversity may be altered via adaptation to the CF lung environment. A

recent effort to sequence more than 1300 *P. aeruginosa* isolates demonstrates that the

genomes are highly flexible, with only 1% of the pan-genome conserved across all isolates (27).

An untargeted analysis of the volatile metabolites produced *in vitro* by 24 *P. aeruginosa* clinical

isolates shows that the volatilomes are also diverse, with 18% of the pan-volatilome conserved

in this comparatively small study (28). In the context of CF, most early *P. aeruginosa* infections

in young patients are caused by unique strains that come from the patients' environments (29-

32). We therefore expect the volatilomes early in infection to have high inter-patient dissimilarity,

reflecting the genomic diversity of *P. aeruginosa* CF isolates (3, 33-35), and to be chemically

diverse (*i.e.*, volatiles from a wide range of chemical classes). Once chronic infections are

established, however, the "founder" *P. aeruginosa* strain evolves into a population of clonal sub-

strains (3, 8, 36-40), many of which harbor CF-typical loss-of-function mutations in regulatory

genes and sigma factors (*e.g.*, *lasR* and *mucA*) (3, 41, 42). From this, we posit that the size of

the *P. aeruginosa* volatilome will shrink during chronic infection, with a concomitant decrease in

chemical diversity and a reduction in inter-patient dissimilarity.

The primary goal of this study is to build a foundation for using volatilomes to diagnose

and track *P. aeruginosa* CF lung infections by exploring how the *in vitro* volatilome of *P.*

aeruginosa CF isolates change over the course of chronic infection. We used comprehensive

two-dimensional gas chromatography coupled with time-of-flight mass spectrometry (GC×GC-

TOFMS) to characterize the volatilomes of 81 *P. aeruginosa* isolates from early and late chronic

lung infections from 17 persons with CF. Using an untargeted metabolomics approach, we

characterized the size and chemical composition of the CF *P. aeruginosa* volatilome, and

identified core volatiles that would be a primary source of breath biomarkers for diagnosing

infections. We also investigated how the *P. aeruginosa* volatilome is shaped by the CF lung

environment, characterizing changes in volatilome sizes and compositions over time, and how

these changes relate to intra-patient and inter-patient volatilome dissimilarities. A secondary

goal of this study is to provide further data on the *P. aeruginosa* volatilome via the largest single

analysis of *P. aeruginosa* headspace volatiles, to-date.

METHODS

Bacterial isolates

Eighty one *P. aeruginosa* isolates from 17 individuals with CF were acquired from the

Cystic Fibrosis Isolate Core at Seattle Children's Center for Global Infectious Disease

Research. From the majority of patients we obtained three isolates: one early isolate, defined as

the first cultured *P. aeruginosa* isolate, and two late isolates collected a minimum of five years

after the first isolate. For one patient (Patient 23), 32 additional isolates were collected at

intervals over 7.5 years between the collections of the early and two late isolates. For one

patient, only one isolate was included in this study. The full strain names of the isolates are

provided in Table S1, and all isolates are available upon request to the CF Isolate Core

(https://www.seattlechildrens.org/research/resources/cystic-fibrosis-isolate/).

We quantified five phenotypes that are correlated with chronic infections: proteases,

pyocyanin, rhamnolipids, twitching motility, and mucoidy. *P. aeruginosa* was cultured from

glycerol stocks on LB agar for 24 h, then a single colony was cultured aerobically to stationary

phase in LB Lennox broth (10 g tryptone, 5 g yeast extract, 5 g NaCl per liter) at 37 °C with

shaking at 200 RPM, unless otherwise noted. Results of phenotype assays are reported in

Table S1 as means of biological triplicates, except for mucoidy for which five replicates were

measured. Pyocyanin production was evaluated by methods adapted from (43). Cell-free *P.*

aeruginosa culture supernatant (7.5 mL) was extracted with 4.5 mL of chloroform, inverted for 2

min and centrifuged at $4,122 \times g$ for 15 min. Three milliliters of the organic phase was removed

and extracted with 750 μL of 0.2 *N* hydrochloric acid and vortexed for 2 min. The aqueous

phase was aliquoted into a 96-well plate and absorbance was measured at 520 nm. Protease

production was evaluated by methods adapted from (15). A sterile wooden inoculation stick was

dipped into a culture of *P. aeruginosa* and then gently touched to the surface of a brain-heart

infusion-skim milk agar (1.5%) plate. Plates were incubated upright at 37 °C and zones of

clearance were measured at 48 h. Rhamnolipids production was evaluated by methods adapted

from (44). Proteose peptone-glucose-ammonium salts (PPGAS) medium was inoculated from

an LB pre-culture of *P. aeruginosa* and cultured at 37 °C to stationary phase with shaking at 200

RPM. Cultures were centrifuged at $21,694 \times g$ for 1 min to pellet cells. The supernatant was

serially diluted two-fold in PPGAS and 20 μL of each dilution was spotted onto a microtiter plate

lid. Spots were classified as drops or collapsed drops and assigned numeric scores

corresponding to the number of dilutions needed to obtain a drop. Twitching motility was

evaluated by methods adapted from (45, 46). The pointed end of a sterile toothpick was touched

to the edge of a single *P. aeruginosa* colony and then stabbed to the bottom of a twitching

motility agar plate (per 1 L: 10 g tryptone, 5 g NaCl, 5 g yeast extract, 10 g agar). Plates were

incubated at 37 °C for 24 h and the radius of the interstitial biofilm was measured. For mucoidy,

frozen glycerol stocks of *P. aeruginosa* were streaked onto LB Lennox agar (1.5%) plates and

grown for 48 hrs. Colonies were visually inspected and the degree of mucoid morphology was

scored from 0 to 2, in 0.5 increments (0 = highly mucoid, 2 = non-mucoid). Twenty pairs of

isolates (one Early and one Late) from ten patients were selected for additional statistical

analyses, described below, based on changes to their phenotypes during chronic infections.

Sample preparation

Isolates were cultured as previously described (47). Briefly, isolates were cultured

aerobically for 16 h at 37 °C in 5 mL of lysogeny broth-Lennox (LB), and then diluted 1000-fold

into 25 mL of fresh LB and grown for 24 h under the same conditions. For metabolomics

analyses, cells were removed via centrifugation through a 0.2 μm filter and 2 mL of each filtrate

was transferred to a 10 mL GC headspace vial with screw caps. Samples were prepared in

biological triplicate, with LB media controls prepared in parallel, and stored at –20 °C prior to

analysis.

GC×GC analysis and data processing

Culture filtrates were thawed and maintained at 4 °C until analyzed, as previously

described (47). Headspace volatiles were characterized using a Pegasus® 4D GC×GC-TOFMS

(LECO® Corporation, St. Joseph, MI) equipped with a MPS Pro® rail autosampler (Gerstel®

Inc, Linthicum Heights, MD). Column set configuration, and GC×GC, MS, and data processing

method parameters are previously reported (47), and summarized in Miscellaneous Table 1

(https://doi.org/10.1101/2020.06.13.126698).

Data processing steps are outlined in Miscellaneous Figure 1. Chromatographic artifacts

and suspected contaminants based on peak names and comparisons to blanks were removed

(Miscellaneous Table 2), as well as poorly modulated peaks eluting prior to 358 s (acetone

retention time) in the first dimension. Missing values were handled as follows: if detected in one

of three replicates, the measured value was permuted to 0; if detected in two of three replicates,

the missing value was imputed as half of the minimum detected value for that compound across

all samples.

Probabilistic quotient normalization (PQN) was applied to account for differences in peak

abundance due to variations in culture cell density, followed by a $\log_{10}$ transformation (48).

Peaks were further filtered out based on the following criteria: (1) within-triplicate intraclass

correlation coefficient (ICC) < 0.75 (definition: absolute agreement; model: two-way mixed

effects; type: mean of k measurements) (49); and (2) only detected in sterile media, or not

significantly greater in abundance compared to sterile media using the Wilcoxon rank-sum test

with Benjamini-Hochberg adjustment (significance threshold of .05) (50).

For statistical analyses beyond the reporting of detected peaks, peaks that showed

significant correlations to run order were identified using Kendall's tau with Benjamini-Hochberg

adjustment (-0.6 ≤ τ ≥ 0.6, significance threshold of .05) and removed. Principal component

analysis (PCA) revealed an apparent batch effect that was attributed to, at least in part, a non-

biological phenomenon and was described previously (47). This batch effect was corrected

using an empirical Bayes approach (51). The geometric means of triplicates of the batch-

corrected data were used for further statistical analyses. For analysis of the ten paired early and

late chronic infection isolates, batch-corrected data was not used as all of these isolates were

contained within the same batch.

Peaks were putatively identified using published reporting standards (52). Level 2

identifications were determined based on the following criteria: (1) ≥ 80% mass spectral forward

match using the NIST 2011 library; (2a) first dimension retention times that possess a strong

linear fit with carbon number in homologous series ($r^2 \geq 0.995$, for the identification of select 2-,

3-, and 4-ketones), or (2b) experimentally determined linear retention indices (LRI) consistent

with published LRIs, as determined by the following acceptance criteria based on the

differences between experimental and published non-polar and polar column RIs for Grob's test

mix:

$$\left(\frac{\mathrm{RI}_{\mathrm{experimental}} - \mathrm{RI}_{\mathrm{non-polar}}}{\mathrm{RI}_{\mathrm{polar}} - \mathrm{RI}_{\mathrm{non-polar}}} \right) \times 100 = 0 - 43\%$$

Level 3 identifications were based on at least an 80% mass spectral forward match score.

Chemical classifications (alcohols, aldehydes, aromatics, carboxylic acids, esters,

hydrocarbons, ketones, thiols, or other) were assigned based primarily on their mass spectral

identity, and secondarily by their chromatographic characteristics. Using polar second

dimension columns, compounds of differing functional classes elute at easily discernible

second-dimension retention times in a stratified manner, typically (from smallest ^2t_R to largest):

hydrocarbons < ketones < aromatics (53-57). In this study, compounds identified as

hydrocarbon, ketones, and aromatics eluted at approximately 0.71 s ± 0.05 s, 0.88 s ± 0.22 s,

and 1.12 s ± 0.26 s, respectively. Level 4 identifications are those that meet none of the above-

mentioned criteria.

Statistical analyses

A Phenotype Score was determined, for each replicate, by scaling the data for each

phenotype to a range of 0 to 1, where 1 is the maximum value and 0 is the minimum value

(excluding outliers). Scaled phenotype data were then summed to yield the Score, and

averaged across replicates. Patient age was not included in calculation of Scores. Two-tailed

Wilcoxon signed-rank tests (with continuity correction, where appropriate; significant threshold

of .05) were used to test for significant differences between Late and Early isolate Scores.

Volatile compounds that were detected in at least 95% of all samples were classified as

"core" volatiles. Significant differences between inter-isolate phenotypes were tested using the

Wilcoxon signed-rank test (significance threshold of .05). Significant differences between intra-

and inter-patient volatilomes were tested using one-way analysis of variance (ANOVA)

(significance threshold of .05), and Tukey's honestly significant differences (HSD) multiple

comparison procedure (significance threshold of .05). The relatedness of isolates based on their

volatilomes were assessed using agglomerative hierarchical clustering analysis (HCA), non-

metric multidimensional scaling (NMDS), and permutational multivariate analysis of variances

(PERMANOVA) on the Euclidean distances between isolates. Linear regression and Pearson's

correlation of Euclidean distances of Patient 23 isolates were used to assess intra-patient

isolate dissimilarity over time. All statistical analyses were performed using R version 3.3.2 (The

R Foundation for Statistical Computing) with the following packages (version): *ICC* (2.3.0),

pairwiseAdonis (0.0.1), *pheatmap* (1.0.8), *stats* (3.3.2), and *sva* (3.22.0).

Data availability

Metabolomic data (chemical feature peak areas and retention time information) included

in this study are available at the NIH Common Fund's National Metabolomics Data Repository

(NMDR) website, the Metabolomics Workbench, at www.metabolomicsworkbench.org, where it

has been assigned Project ID PR000970, Study ID ST001414 (DOI:

http://dx.doi.org/10.21228/M89Q4F). Additional Miscellaneous Information mentioned can be

found alongside Supplementary Information at bioRxiv (DOI:

https://doi.org/10.1101/2020.06.13.126698)

RESULTS

Characteristics of *P. aeruginosa* CF isolates

We obtained from a biorepository 81 *P. aeruginosa* chronic infection isolates, which had

been collected from 17 individuals with CF. From 14 patients, two or three *P. aeruginosa*

isolates were obtained: one isolate that was the first cultured *P. aeruginosa*, and one or two

isolates that were collected at least five years after the first. We refer to these isolates as Early

and Late, respectively. For one patient (Patient 75), four isolates were collected: two Early

isolates collected one month apart and two Late isolates, collected ten and 16 years after the

first. For one patient (Patient 23), 32 additional isolates were collected over the course of a 7.5 y

infection period, which we have termed Intermediate isolates. For one patient, only one isolate

was collected. These isolates were genetically characterized in a study by Smith *et al*., who

determined that the intra-patient replicate isolates are all clonally related, with the exception of

the four isolates from patient 75, which are actually two clonally-related Early/Late pairs

(isolates 75E-1 and 75L-2 are clonal, and isolates 75E-2 and 75L-1 are clonal) (3).

For all isolates, we characterized five clinically relevant phenotypes *in vitro*: mucoidy,

pyocyanin production, rhamnolipids production, protease production, and twitching motility (Fig.

1; Table S1). These phenotypes are commonly altered during the course of chronic CF lung

infections (15). There were wide ranges in the expression of these phenotypes across all

isolates, as expected. Several isolate sets exhibited phenotypes consistent with age of

collection in that Early isolates possessed higher degrees of motility, higher quantities of

pyocyanin, rhamnolipids, and proteases (indicating intact quorum regulation), and lower

mucoidy compared to their cognate Late isolates (*e.g.*, Patients 23, 31, 33, 36, and 76). The

associations between relative patient age at isolate collection and *P. aeruginosa* phenotypes

were not perfect, however. For example, Early isolate 62E had no detectable pyocyanin,

proteases, rhamnolipids, or twitching motility, and many of the Late isolates retained more early-

like phenotypes (e.g., both Late isolates of 41, 66, 71, 75, and 101, and 60L-1, 74L-2, and 100L-

2). Despite observable changes in isolate phenotype within patient, no phenotypes were

significantly different when comparing Late isolate(s) to the Early isolate. Collectively, the 81

isolates were highly varied, and represented the array of phenotypes we expect to observe in

the span of *P. aeruginosa* CF lung infections from initial infection to long-established chronic

infections.

The *in vitro* volatilome of *P. aeruginosa* CF isolates

Taking an untargeted metabolomics approach, we cultured the clinical *P. aeruginosa*

isolates *in vitro* and characterized their volatilomes using GC×GC-TOFMS. Following extensive

data processing, including removal of analytically biased chromatographic features, we

conservatively attribute 539 non-redundant volatile compounds to the growth and metabolism of

the *P. aeruginosa* isolates (Table S2). Among these, 69 compounds are shared by at least 95%

($n \geq 77$) of the isolates, representing core volatiles. Using minimum metabolomics reporting

standards, we assigned compound identification levels 1-4 (1 high, with 2 being the highest in

this study) based on a combination of mass spectral and chromatographic characteristics. We

were able to assign putative names to 14 core and 33 non-core compounds with an ID level 2

(Table 1), many of which have been previously reported as *P. aeruginosa* volatiles (Table 1).

While we detected 2-aminoacetophenone (the volatile compound responsible for the grape-like

odor of *P. aeruginosa*), it possessed an intraclass correlation slightly lower than the applied

threshold of 0.75 and was ultimately filtered out of our peak list. We also identified several

compounds of a variety of chemical classes that have not been previously reported for *P.*

aeruginosa, including: alcohols (3-methyl-1-buten-1-ol, 1-butoxy-2-propanol, 2-butyl-1-octanol),

esters (butanoic acids, methyl isovalerate), aromatics (tetrahydrofuran, tetramethylpyrazine, 2-

butylfuran, 3-cyano-2,5-dimethylpyrazine, 2,4-(1,1-dimethyl)-phenol), unsaturated ketones (3-

hydroxy-butan-2-one, 4-undecanone, 1-octen-3-one, 3-penten-2-one, 6-methyl-2-heptanone, 1-

(4-ethylphenyl)-ethanone), and an aldehyde (decanal).

Chemical classifications were assigned to ID level 2 and 3 compounds (187 compounds)

using mass spectral match and retention time characteristics (Fig. 2). Among Level 2 and 3 core

volatiles, the majority of classified compounds were hydrocarbons (41%), followed by ketones

(16%). Other oxidized compounds, including alcohols, aromatics, acids, and thiols, accounted

for an additional 43%. The non-core volatiles had similar chemical compositions, which are

consistent with previous studies of the *P. aeruginosa* volatilome (25, 28, 53, 58, 59). Of the

entire volatilome, 65% of the volatiles (*n* = 352) possessed less than an 80% mass spectral

match to the 2011 NIST MS library, and as such were classified as unknowns.

Table 1 Putatively identified volatile compounds metabolized by *P. aeruginosa* CF isolates.

Compound Name[a]	Formula	Chemical Class	Samples (*n* = 81)	Refs.[b]
Core volatile compounds				
3-methylbutanal	$C_5H_{10}O$	Aldehyde	79	
thiocyanic acid, methyl ester	C_2H_3NS	Ester	77	(53)
3-methyl-1-butanol	$C_5H_{12}O$	Alcohol	77	(58, 60, 61)
pyridine	C_5H_5N	Aromatic	77	
3-penten-2-one	C_5H_8O	Ketone	77	
3-methyl-1-buten-1-ol	$C_5H_{10}O$	Alcohol	78	
4-methyl-3-penten-2-one	$C_6H_{10}O$	Ketone	81	(28, 62)
hexanal	$C_6H_{12}O$	Aldehyde	81	(58, 60, 62)
heptanal	$C_7H_{14}O$	Aldehyde	80	
2,5-dimethylpyrazine	$C_6H_8N_2$	Aromatic	81	(53)
2-nonanone	$C_9H_{18}O$	Ketone	81	(28, 53, 58, 59, 61-67)
4-ethyl-1,2-dimethylbenzene	$C_{10}H_{14}$	Aromatic	77	
1,3,3-trimethyl-(bicyclo)-heptan-2-ol	$C_{10}H_{18}O$	Alcohol	78	
1-(4-ethylphenyl)-ethanone	$C_{10}H_{12}O$	Ketone	79	
Non-core volatile compounds				
dimethyl sulfide	C_2H_6S	Other	29	(28, 53, 58, 60, 61, 64, 68, 69)
2-methyl-3-buten-2-ol	$C_5H_{10}O$	Alcohol	35	
2-butanone	C_4H_8O	Ketone	1	(28, 53, 58, 59, 61, 63, 66, 70)
tetrahydrofuran	C_4H_8O	Aromatic	1	
acetic acid	$C_2H_4O_2$	Carboxylic acid	55	
methyl thioacetate	C_3H_6OS	Thiol	3	(58, 60, 62)

2,4-dimethylfuran	C_6H_8O	Aromatic	66	(28)
butanoic acid, methyl ester	$C_5H_{10}O_2$	Ester	3	
3-hydroxybutan-2-one	$C_4H_8O_2$	Ketone	18	
dimethyl disulfide	$C_2H_2S_2$	Other	1	(28, 58-62, 66, 68-73)
2-methyl-butanoic acid, methyl ester	$C_6H_{12}O_2$	Ester	3	
3-methyl-butanoic acid, methyl ester	$C_6H_{12}O_2$	Ester	3	
furfural	$C_5H_4O_2$	Aromatic	67	(60)
2-butylfuran	$C_8H_{12}O$	Aromatic	6	
2-hexen-1-ol	$C_6H_{12}O$	Alcohol	51	(28)
1-butoxy-2-propanol	$C_7H_{16}O_2$	Alcohol	61	
6-methyl-2-heptanone	$C_8H_{16}O$	Ketone	33	
1-methylethylbenzene	C_9H_{12}	Aromatic	57	
3-octanone	$C_8H_{16}O$	Ketone	56	(28, 61)
4,6-dimethyl-2-heptanone	$C_9H_{18}O$	Ketone	47	
2-ethyl-5-methylpyrazine	$C_7H_{10}N_2$	Aromatic	58	
4-nonanone	$C_9H_{18}O$	Ketone	49	(28)
tetramethylpyrazine	$C_8H_{12}N_2$	Aromatic	5	
2-octenal	$C_8H_{14}O$	Aldehyde	48	(53)
octanenitrile	$C_8H_{15}N$	Other	73	(28)
2,4-octadienal	$C_8H_{12}O$	Aldehyde	9	(28)
3-decanone	$C_{10}H_{20}O$	Ketone	41	(53)
decanal	$C_{10}H_{20}O$	Aldehyde	65	
2-decanone	$C_{10}H_{20}O$	Ketone	13	(28)
4-undecanone	$C_{11}H_{22}O$	Ketone	30	
2-dodecanone	$C_{12}H_{24}O$	Ketone	18	(28, 53)
2-butyl-1-octanol	$C_{12}H_{26}O$	Alcohol	48	
2,4-bis(1,1-dimethyl)-phenol	$C_{14}H_{22}O$	Aromatic	26	

[a]Bolded compounds have not been previously reported as *P. aeruginosa* volatile compounds.
[b]Previous references for these volatiles in *P. aeruginosa* cultures are provided.

Relationships in the *P. aeruginosa* CF volatilome within and between patients

Because every patient is infected by a genetically-unique strain, we hypothesized that

the dissimilarities of intra-patient volatilomes would be lower than inter-patient. Additionally, due

to the greater metabolic potential of early-infection isolates with intact regulatory networks, we

hypothesized that the volatilomes of inter-patient Early isolates would be more dissimilar than

those of inter-patient Late isolates. We calculated the pairwise Euclidean distances as a

measure of dissimilarity between all Early isolate volatilomes ("inter-Early dissimilarity"), all Late

isolate volatilomes ("inter-Late dissimilarity"), and between Early and Late isolate volatilomes

from the same patient ("intra-patient dissimilarity"). Significant differences between means were

tested using one-way ANOVA with Tukey's HSD multiple comparison procedure. Interestingly,

the inter-Early isolate dissimilarity was the lowest (mean 35.2, median 33.8) and the inter-Late

isolate dissimilarity was the highest (mean 39.1, median 39.1), and these differences were

significant (95% CI: 3.05 – 4.86; P = 0+) (Fig. 3). Similarly, mean inter-Late volatilome

dissimilarity was significantly greater than that of intra-Patient volatilomes (mean 36.4, median

35.6) (95% CI: 0.41 – 4.97; P = .013). Even more intriguing was that the inter-Early dissimilarity

was lower than even the intra-patient dissimilarity, though this difference was not significant

(95% CI: -1.11 – 3.66; P = .515). For comparison, we also calculated the pairwise Euclidean

distances of all isolates in this study (Pooled; mean 38.0, median 37.9); the mean inter-Early

and mean inter-Late dissimilarities were both significantly lower than the mean dissimilarity of

the Pooled isolates (95% CIs: 1.99 – 3.77, -1.65 – 0.51; P = 0+).

Visualizing the Euclidean distances between isolate volatilomes via non-metric

multidimensional scaling (NMDS) and ordered dissimilarity images (ODI), we looked for

clustering of clonal isolates, indicating intra-patient similarities. The NMDS and ODI of the

Euclidean distances of all 81 isolate volatilomes show a primary cluster comprised mostly of

intermediate isolates of Patient 23 (Fig. S1). To reduce the influence of the over-represented

Patient 23 isolates in data visualization, we truncated our data set to 48 isolates by removing

the Intermediate isolates of Patient 23, and the lone Patient 21 isolate. The ODI plot and NMDS

of the truncated set revealed three clusters of similarity, none of which include all isolates from a

single patient (Fig. 4; Fig S2a,b). This refutes our hypothesis that isolates from the same patient

maintain a similar volatilome over time. To determine if any inter-patient or intra-patient isolate

pairs are significantly different, we performed permutational multivariate analysis of variance

(PERMANOVA) on the Euclidean distances for all possible pairs. Following Benjamini-Hochberg

adjustment of *P* values, no pairs of isolates were identified as having significantly different

volatilomes ($q \geq .1$ for all isolate pairs). However, PERMANOVA conducted on all isolates

indicated strong significant differences between the 81 volatilomes (*pseudo-F*$_{80,161}$ = 28.0, *P*

< .01), as well as between the volatilomes of the truncated set of 48 isolates (*pseudo-F*$_{47,96}$ =

29.2, *P* = 0+), reinforcing the inference from the ODI and NMDS that the CF *P. aeruginosa*

volatilome is heterogeneous.

Though we do not find that patient isolates maintain similar volatilomes over the duration

of chronic infection, it is noteworthy that the majority of Patient 23's isolates have strong

similarities (Fig. S1). Taking a closer look at the 35 isolates from Patient 23 (Fig. 5; Fig. S2c),

the ODI and NMDS revealed four distinct similarity neighborhoods: one neighborhood consisting

of only the Early isolate 23E, which stands alone; a second consisting of the four latest-collected

isolates 23I-32, 23I-33, 23L-1, 23L-2; a third consisting of isolates 23I-7, 23I-9, 23I-12, 23I-14,

23I-29; and a fourth encompassing the remainder of the isolates. The four latest-collected

isolates are less defined by their similarity to each other than their dissimilarity to all other

isolates collected during the infection. A linear regression of the Euclidean distances between

the Early isolate and the Intermediate and Late isolates as a function of patient age showed that

isolates became increasingly dissimilar to the Early isolate over time (Pearson's r = 0.70, *P* =

0+) (Fig. 6). These observations are underpinned by the phenotypic differences of the four

latest isolates, and the number and types of known mutations that were accumulated in the 23L

versus 23E isolates (3). As described by Smith *et al.*, the *P. aeruginosa* infection sampled from

Patient 23 (referred to as Patient 1 in the referenced publication) diversified from patient ages

1.5 to 3 y into a population of isolates, with mutations in virulence, motility, quorum sensing, iron

transport, efflux, and transcription and translation genes. The volatilome dissimilarities we

measured reflect the genotypic diversification of the infection as a function of time (Fig. 6).

Chemical characteristics of the volatilomes of Early and Late isolates

We compared the chemical characteristics of the Early and Late volatilomes, positing that the size and chemical diversity of the volatilomes would decrease from Early to Late infection due to loss-of-function mutations. To perform the comparison, we needed to balance the size of the isolate groups, and did so by selecting ten pairs of *P. aeruginosa* isolates that had large changes in phenotypes over the duration of infection, thereby enhancing the differences we may find in the Early vs. Late volatilomes (Fig. 1; Table S1). Visual inspection of the ten pairs of GC×GC chromatograms suggests reductions in the number and variety of volatile compounds produced by Late isolates (Fig. 7, Miscellaneous Figure 2). Contrary to the appearance of the chromatograms, however, the overall size and chemical compositions of the Early and Late volatilomes were similar to each other (Fig. S3a-d), with hydrocarbons representing approximately 50% and alcohols and ketones together representing approximately 30% of the 410 Early and 441 Late volatiles. We quantified the chemical richness and diversity of the pooled Early and pooled Late isolates using the Shannon-Wiener diversity index (Table S3) and found the volatilomes of Early and Late isolates were similar, whether using all volatiles or only the non-core volatiles.

The uniformity in the Early versus Late volatilomes is not a function of data aggregation, as the volatilomes of the individual isolates are highly similar in number and chemical composition, with the exception of one Late isolate (76L) that has a slow-growth phenotype (Fig. S3e-h). Furthermore, when looking at individual isolate pairs, for the vast majority there were no differences in the richness and diversity of their volatilomes (Table S3). Together, these results suggest that there are not major differences between the number and diversity of chemical compounds when comparing Early and Late infection isolates. Hierarchical clustering analysis (HCA) of the *P. aeruginosa* isolates by their volatilomes underscores this observation; we

observed no discernible clustering of the Early versus Late isolates based on the presence and

absence of volatiles (Fig. S4a).

Abundance of volatile compounds in Early and Late isolates

Though we did not observe any differences in the Early and Late isolate volatilomes

based on presence and absence of metabolites, the chromatograms suggest a reduced

volatilome in late infection. We therefore posited that while the numbers of volatiles do not

significantly change from Early to Late infection isolates, the relative concentrations of volatiles

do. Using volatile abundances, we performed HCA on the 20 selected Early and Late isolates

and found clustering by the time of isolate collection and phenotype score (Fig. S4b). We were

interested in whether or not clustering still occurs in the larger, less curated set of 48 isolates

(*i.e.*, the truncated set), which has more discordance between phenotype and time of collection

(Fig. 1). HCA of the truncated set of isolates shows that Early isolates are generally more similar

to one another than Late isolates are (Fig. 8), reinforcing observations made by calculating

Euclidean distances; however, significant proportions of the Early and the Late isolates are mis-

clustered. Interestingly, the majority of the Late isolates clustering with the Early isolates in the

upper clade have more early-like phenotypes, and several Early isolates that have more late-

like phenotypes (*e.g.*, 62E, 73E, 70E) cluster in the lower clade with Late isolates, indicating a

relationship between the isolate phenomes and volatilomes. We also observed clustering by

phenotype when the volatilomes of all 81 isolates were analyzed (Fig. S5). Similar phenomena

are observed when clustering the truncated set or all 81 isolates by the core volatilome (Fig.

S6), and even when using only the 23 core alcohols, aldehydes, and hydrocarbons (Fig. 9; Fig.

S7), suggesting that a set of conserved volatiles could be identified and used as biomarkers for

detecting phenotypic changes in chronic infections.

DISCUSSION

P. aeruginosa isolates from cystic fibrosis lung infections have a large and diverse

volatilome. We conservatively attribute 539 detected volatiles to *P. aeruginosa* growth and

metabolism, only 13% of which were core volatiles. For volatiles that we could assign to a

chemical class, hydrocarbons represented the highest proportion, followed by ketones, which

we observed both in the pan-volatilome and for individual patient isolates. Approximately 40% of

core volatiles have been identified in other *P. aeruginosa* metabolomic studies, including the

most frequently detected *in vitro* volatiles 2-butanone, 2-nonanone, dimethyl sulfide, and

dimethyl disulfide (see Table 1 for references). The majority of the core volatiles we detected,

however, are reported here for the first time. This opens up speculation that there may be a

unique core volatile profile for strains colonizing the CF lung, though due to the inter-strain

variation in the *P. aeruginosa* volatilome, many more CF and non-CF isolates will need to be

characterized via untargeted GC×GC analyses to determine if this may be the case. To date, a

number of studies have attempted to identify collections of volatiles to differentiate *P.*

aeruginosa-positive or -negative subjects from breath (58, 74-81) and lavage (25) with

promising results. Compared to this study, however, the referenced analyses were either

targeted at single compounds (e.g., hydrogen cyanide or 2-aminoacetophenone), did not

characterize the volatilome of *P. aeruginosa* in detail, and/or did not draw inferences to infection

stage or changes across time. This study represents the first untargeted comparative analysis of

P. aeruginosa volatilomes over long-term chronic infections, expanding the body of knowledge

on *P. aeruginosa* metabolism and broadening the potential applications for breath-based

diagnostics.

Despite the trend toward reduced or loss-of-phenotype expression in late infection

isolates – which we hypothesized would correspond to a reduced metabolome size – we

observed that Late isolates produce the same number and variety of volatile compounds as

Early isolates. However, the relative abundance of volatiles produced by the Late isolates is

significantly reduced. Our data suggest that mutations that arise in *P. aeruginosa* chronic

infection isolates do not result in complete inhibition of metabolic pathways, but may instead

reduce metabolic flux. This observation has implications for diagnostics, signifying that presence

and absence of *P. aeruginosa* volatile metabolites would provide little diagnostic value for

tracking chronic lung infections. Rather, a breath-based diagnostic for monitoring *P. aeruginosa*

adaptations in CF will likely need to measure the relative abundance of metabolites. Importantly,

a subset of core volatiles can be used to cluster isolates based on commonalities in

phenotypes, indicating that a diagnostic based upon the relative abundances of conserved

hydrocarbons, alcohols, and aldehydes (or a few selected volatiles from these classes) may be

sufficient to track *P. aeruginosa* infection in most patients. It should be noted that core *P.*

aeruginosa volatiles are not necessarily unique to *P. aeruginosa*. In fact, a review of the mVOC

2.0 microbial volatile organic compound database (82) shows nearly half of the core volatiles

(and a third of named compounds overall) have been reported as being produced by other CF

lung bacterial pathogens, including *Staphylococcus aureus*, *Klebsiella pneumoniae*,

Haemophilus influenzae, *Stenotrophomonas maltophilia*, and *Burkholderia spp*. This further

highlights the necessity of developing a breath-based diagnostic based on the relative

abundances of sets of volatiles, which has been shown to accurately discriminate between

human pathogens (distantly or closely related to *P. aeruginosa*) in numerous *in vitro*, *ex vivo*,

and *in vivo* analyses (25, 62, 83-85).

Based on the genomic heterogeneity of *P. aeruginosa* (3, 33-35), we hypothesized that

every patient would have a unique *P. aeruginosa* volatilome, and that intra-patient isolates

would have more similar volatilomes than inter-patient isolates. To the contrary, we observed

that the volatilomes of Early and Late isolates from a single patient are no more similar to one

another than they are to any other isolate (intra-patient vs. pooled data, Fig. 3). Interestingly, the

volatilomes of inter-patient Early isolates are even more similar than intra-patient isolates, while

the volatilomes of inter-patient Late isolates have the highest dissimilarity. These data indicate

that the genomic similarity within patient is less influential on the volatilome than the post-

transcriptional diversity that evolves within a chronic *P. aeruginosa* infection population. The

data from Patient 23, however, suggest that intra-patient isolates are more likely to have similar

volatilomes when there are fewer mutational and phenotypic differences between them. From

these results, we propose that a breath-based diagnostic to track *P. aeruginosa* chronic lung

infections would be personalized, with each patient's breathprint at initial infection serving as a

baseline to which subsequent breathprints are compared. Sustained, significant deviations from

the patient's baseline breathprint would indicate that the *P. aeruginosa* population phenotype

has changed, and therefore the patient may be at higher risk for associated poor clinical

outcomes (*e.g.*, higher risk of exacerbation) (15) and at higher risk of infection eradication failure

after antibiotic treatment (12). In a longitudinal case study of a single CF patient, we have

shown that changes in the dominant species colonizing the lung correlate to changes in the

volatile metabolites detected in sputum (86), and therefore monitoring for breathprint deviations

may also work for *P. aeruginosa* infections, as these data suggest. Despite this, the issue of

specificity becomes a concern unless biomarkers for individual *P. aeruginosa* phenotypes are

identified and used for tracking; this is the subject of ongoing work.

We highlight some limitations in this study. Although we make reference to Early isolates

and those phenotypes associated with early infection, it is important to recognize that these

isolates were merely the first culturable *P. aeruginosa* from the patients included in this study. It

is likely that some patients had *P. aeruginosa* in the lung for some years in spite of persistent

negative cultures into late childhood or adolescence (15), exemplified by isolates 62E and 73E

exhibiting late-like phenotypes. Phenotypes discordant with collection time or apparent infection

stage underscores a major drawback of culture-based methods for infection staging (*e.g.*, *in*

vitro phenotyping) in that the respiratory microbiome is prone to under-sampling, especially

during initial colonization, due to the complex structure of the lung. The increased use of new

therapeutics (*e.g.*, CFTR modulators) also impacts culture-based diagnostics as they reduce

sputum production, limiting the opportunity for clinical cultures. Utilizing breath-based

biomarkers, on the other hand, can overcome these challenges as breath represents a sample

of the entire ventilated lung environment.

P. aeruginosa chronic infection phenotypes are positively correlated with advanced

patient age (15, 87-89), and we observe a moderate, but significant correlation (Kendall's tau –

0.41, *P* < .001) between subject age and late-infection phenotype scores in this study. The

variability of isolate volatilomes we observed might, therefore, be explained by the age of their

corresponding patients. Testing this would require analyzing isolate volatilomes from a larger

number of subjects that were diagnosed with their first *P. aeruginosa* infections at younger and

older ages. In either case, the issue of the potential covariance between age and volatilome on

P. aeruginosa diagnostics could be overcome by taking a multifactorial approach as opposed to

volatile biomarkers only. We may find that using a combination of several clinical predictors

such as age, *P. aeruginosa* phenotype, and volatilome together provides more diagnostic power

than any individual metric. It will be important to explore this more closely in future work.

Additionally, while we only presented volatilome trends in relation to five clinically-relevant

phenotypes in this study (production of pyocyanin, rhamnolipids, and proteases, twitching

mobility, and mucoidy), there are additional clinically-relevant phenotypes, including antibiotic

resistance, that should be included in future analyses.

In summary, current methods for tracking *P. aeruginosa* lung infection progression are

predominantly culture-based (16, 21), and our results suggest a potential role for *in vitro* volatile

metabolomics in staging chronic infections in conjunction with established clinical laboratory

methods. The clinically-correlated *P. aeruginosa* phenotypes we measured in this study –

mucoidy, motility, and quorum-regulated traits – are sensitive to environment (90-93); as such,

the metabolomes we characterized from *in vitro* cultures are unlikely to directly reflect the

metabolomes these isolates produced in the CF lung. For the development of a breath test for

tracking infections *in situ*, it will be necessary to utilize culture models that are more reflective of

the lung (94-97) and to collect *ex vivo* and *in vivo* infection volatilomes (25, 75, 98). The utility of

breath for tracking chronic lung infections is being demonstrated in tuberculosis lung infection

models (26), and therefore we anticipate that the development of a breath-based diagnostic for

tracking *P. aeruginosa* chronic lung infections is feasible. It will also be important to ensure that

the *P. aeruginosa* volatiles we have identified are detectable in breath. This is the primary goal

of the observational clinical study IMproving *P. Aeruginosa* deteCTion using Breath (IMPACT-

Breath) that is underway. Despite the limitations of *in vitro* analyses, this study presents the first

comprehensive analysis of the *P. aeruginosa* volatilome from chronic CF lung infections, and

lays the groundwork for the application of volatile metabolomics in tracking CF lung disease.

ACKNOWLEDGEMENTS

Funding to support this work was provided by the Cystic Fibrosis Foundation (Hill17P0,

Hill18A0-CI), the National Institutes of Health (R56HL139846), and the ASU School of Life

Sciences Undergraduate Research (SOLUR) program. Isolates were obtained from the Cystic

Fibrosis Isolate Core at Seattle Children's Research Institute, funded by the National Institutes

of Health (P30 DK089507) and the Cystic Fibrosis Foundation (HOFFMA20Y2-OUT).

AUTHOR CONTRIBUTIONS

HB conceived of the study; HB, TD designed the experiments; AK, CB, SW, and TD collected

phenotype data; TD collected metabolomics data; TD and MR processed and analyzed the

metabolomics data; HB and TD wrote and edited the manuscript; all authors approved the final

version.

CONFLICTS OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

REFERENCES

1. Gibson RL, Burns JL, Ramsey BW. 2003. Pathophysiology and management of

pulmonary infections in cystic fibrosis. Am J Respir Crit Care Med 168:918-951.

2. Marshall B, Faro A, Fink A, Loeffler D, Elbert A, O'Neil T, Rush T, Rizvi S. 2018. Cystic

Fibrosis Foundation patient registry: 2017 annual data report. Cystic Fibrosis

Foundation, Foundation CF, Bethesda, MD.

3. Smith EE, Buckley DG, Wu ZN, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller

SI, Ramsey BW, Speert DP, Moskowitz SM, Burns JL, Kaul R, Olson MV. 2006. Genetic

adaptation by *Pseudomonas aeruginosa* to the airways of cystic fibrosis patients. Proc

Natl Acad Sci U S A 103:8487-8492.

4. Li ZH, Kosorok MR, Farrell PM, Laxova A, West SEH, Green CG, Collins J, Rock MJ,

Splaingard ML. 2005. Longitudinal development of mucoid *Pseudomonas aeruginosa*

infection and lung disease progression in children with cystic fibrosis. JAMA 293:581-

588.

5. Emerson J, Rosenfeld M, McNamara S, Ramsey B, Gibson RL. 2002. *Pseudomonas

aeruginosa* and other predictors of mortality and morbidity in young children with cystic

fibrosis. Pediatr Pulmonol 34:91-100.

6. Kosorok MR, Zeng L, West SEH, Rock MJ, Splaingard ML, Laxova A, Green CG, Collins

J, Farrell PM. 2001. Acceleration of lung disease in children with cystic fibrosis after

Pseudomonas aeruginosa acquisition. Pediatr Pulmonol 32:277-287.

7. Davidson AG, Chilvers MA, Lillquist YP. 2012. Effects of a *Pseudomonas aeruginosa*

eradication policy in a cystic fibrosis clinic. Curr Opin Pulm Med 18:615-21.

8.   Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* evolutionary

adaptation and diversification in cystic fibrosis chronic lung infections. Trends Microbiol

24:327-337.

9.   Hogardt M, Heesemann J. 2013. Microevolution of *Pseudomonas aeruginosa* to a

chronic pathogen of the cystic fibrosis lung, p 91-118. *In* Dobrindt U, Hacker JH,

Svanborg C (ed), Between pathogenicity and commensalism, vol 358. Springer-Verlag

Berlin, Berlin.

10.   Sanders DB. 2014. Phenotypes that matter: *Pseudomonas aeruginosa* and progression

of cystic fibrosis lung disease. Am J Respir Crit Care Med 190:245-246.

11.   Pritt B, O'Brien L, Winn W. 2007. Mucoid *Pseudomonas* in cystic fibrosis. Am J Clin

Pathol 128:32-34.

12.   Mayer-Hamblett N, Ramsey BW, Kulasekara HD, Wolter DJ, Houston LS, Pope CE,

Kulasekara BR, Armbruster CR, Burns JL, Retsch-Bogart G, Rosenfeld M, Gibson RL,

Miller SI, Khan U, Hoffman LR. 2014. *Pseudomonas aeruginosa* phenotypes associated

with eradication failure in children with cystic fibrosis. Clin Infect Dis 59:624-631.

13.   Courtney JM, Bradley J, McCaughan J, O'Connor TM, Shortt C, Bredin CP, Bradbury I,

Elborn JS. 2007. Predictors of mortality in adults with cystic fibrosis. Pediatr Pulmonol

42:525-532.

14.   Hauser AR, Jain M, Bar-Meir M, McColley SA. 2011. Clinical significance of microbial

infection and adaptation in cystic fibrosis. Clin Microbiol Rev 24:29-70.

15.   Mayer-Hamblett N, Rosenfeld M, Gibson RL. 2014. *Pseudomonas aeruginosa in vitro*

phenotypes distinguish cystic fibrosis infection stages and outcomes. Am J Respir Crit

Care Med 190:289-297.

16.   Héry-Arnaud G, Nowak E, Caillon J, David V, Dirou A, Revert K, Munck MR, Frachon I,

Haloun A, Horeau-Langlard D, Le Bihan J, Danner-Boucher I, Ramel S, Pelletier MP,

Rosec S, Gouriou S, Poulhazan E, Payan C, Ferec C, Rault G, Le Gal G, Le Berre R.

2017. Evaluation of quantitative PCR for early diagnosis of *Pseudomonas aeruginosa*

infection in cystic fibrosis: A prospective cohort study. Clin Microbiol Infect 23:203-207.

17.   Kim K-H, Jahan SA, Kabir E. 2012. A review of breath analysis for diagnosis of human

health. Trends Anal Chem 33:1-8.

18.   Ahmed WM, Lawal O, Nilsen TM, Goodacre R, Fowler SJ. 2017. Exhaled volatile

organic compounds of infection: A systematic review. ACS Infect Dis 3:695-710.

19.   Oakley-Girvan I, Davis SW. 2018. Breath based volatile organic compounds in the

detection of breast, lung, and colorectal cancers: A systematic review. Cancer Biomark

21:29-39.

20.   Amann A, Miekisch W, Schubert J, Buszewski B, Ligor T, Jezierski T, Pleil J, Risby T.

2014. Analysis of exhaled breath for disease detection. Annu Rev Anal Chem, Vol 7

7:455-482.

21.   Smith WD, Bardin E, Cameron L, Edmondson CL, Farrant KV, Martin I, Murphy RA,

Soren O, Turnbull AR, Wierre-Gore N, Alton EW, Bundy JG, Bush A, Connett GJ, Faust

SN, Filloux A, Freemont PS, Jones AL, Takats Z, Webb JS, Williams HD, Davies JC.

2017. Current and future therapies for *Pseudomonas aeruginosa* infection in patients

with cystic fibrosis. FEMS Microbiol Lett 364.

22.   Burns JL, Gibson RL, McNamara S, Yim D, Emerson J, Rosenfeld M, Hiatt P, McCcoy

K, Castile R, Smith AL, Ramsey BW. 2001. Longitudinal assessment of *Pseudomonas*

aeruginosa in young children with cystic fibrosis. J Infect Dis 183:444-452.

23.   Hoppe JE, Towler EE, Wagner BD, Accurso FJ, Sagel SD, Zemanick ET. 2015. Sputum

induction improves detection of pathogens in children with cystic fibrosis. Pediatr

Pulmonol 50:638-646.

24.   West SEH, Zeng L, Lee BL, Kosorok MR, Laxova A, Rock MJ, Splaingard MJ, Farrell

PM. 2002. Respiratory infections with *Pseudomonas aeruginosa* in children with cystic

fibrosis - early detection by serology and assessment of risk factors. JAMA 287:2958-

2967.

25.   Nasir M, Bean HD, Smolinska A, Rees CA, Zemanick ET, Hill JE. 2018. Volatile

molecules from bronchoalveolar lavage fluid can 'rule-in' *Pseudomonas aeruginosa* and

'rule-out' *Staphylococcus aureus* infections in cystic fibrosis patients. Sci Rep 8.

26.   Beccaria M, Bobak C, Maitshotlo B, Mellors TR, Purcaro G, Franchina FA, Rees CA,

Nasir M, Black A, Hill JE. 2019. Exhaled human breath analysis in active pulmonary

tuberculosis diagnostics by comprehensive gas chromatography-mass spectrometry and

chemometric techniques. J Breath Res 13.

27.   Freschi L, Vincent AT, Jeukens J, Emond-Rheault JG, Kukavica-Ibrulj I, Dupont MJ,

Charette SJ, Boyle B, Levesque RC. 2019. The *Pseudomonas aeruginosa* pan-genome

provides new insights on its population structure, horizontal gene transfer, and

pathogenicity. Genome Biol Evol 11:109-120.

28.   Bean HD, Rees CA, Hill JE. 2016. Comparative analysis of the volatile metabolomes of

Pseudomonas aeruginosa clinical isolates. J Breath Res 10.

29.   Kelly NM, Fitzgerald MX, Tempany E, O'Boyle C, Falkiner FR, Keane CT. 1982. Does
Pseudomonas cross-infection occur between cystic-fibrosis patients. Lancet 2:688-90.

30.   Grothues D, Koopmann U, von der Hardt H, Tummler B. 1988. Genome fingerprinting of
Pseudomonas aeruginosa indicates colonization of cystic fibrosis siblings with closely
related strains. J Clin Microbiol 26:1973-7.

31.   Thomassen MJ, Demko CA, Doershuk CF, Root JM. 1985. *Pseudomonas aeruginosa*
isolates: Comparisons of isolates from campers and from sibling pairs with cystic
fibrosis. Pediatr Pulmonol 1:40-5.

32.   Wolz C, Kiosz G, Ogle JW, Vasil ML, Schaad U, Botzenhart K, Doring G. 1989.
Pseudomonas aeruginosa cross-colonization and persistence in patients with cystic
fibrosis. Use of a DNA probe. Epidemiol Infect 102:205-14.

33.   Ashish A, Paterson S, Mowat E, Fothergill JL, Walshaw MJ, Winstanley C. 2013.
Extensive diversification is a common feature of *Pseudomonas aeruginosa* populations
during respiratory infections in cystic fibrosis. J Cyst Fibros 12:790-3.

34.   Cramer N, Klockgether J, Wrasman K, Schmidt M, Davenport CF, Tummler B. 2011.
Microevolution of the major common *Pseudomonas aeruginosa* clones C and PA14 in
cystic fibrosis lungs. Environ Microbiol 13:1690-704.

35.   Mowat E, Paterson S, Fothergill JL, Wright EA, Ledson MJ, Walshaw MJ, Brockhurst
MA, Winstanley C. 2011. *Pseudomonas aeruginosa* population diversity and turnover in
cystic fibrosis chronic infections. Am J Respir Crit Care Med 183:1674-1679.

36.   Bragonzi A, Paroni M, Nonis A, Cramer N, Montanari S, Rejman J, Di Serio C, Doring G,
Tummler B. 2009. *Pseudomonas aeruginosa* microevolution during cystic fibrosis lung

infection establishes clones with adapted virulence. Am J Respir Crit Care Med 180:138-

45.

37.   Kordes A, Preusse M, Willger SD, Braubach P, Jonigk D, Haverich A, Warnecke G,

Haussler S. 2019. Genetically diverse *Pseudomonas aeruginosa* populations display

similar transcriptomic profiles in a cystic fibrosis explanted lung. Nat Commun 10:3397.

38.   Clark ST, Guttman DS, Hwang DM. 2018. Diversification of *Pseudomonas aeruginosa*

within the cystic fibrosis lung and its effects on antibiotic resistance. FEMS Microbiol Lett

365.

39.   Schick A, Kassen R. 2018. Rapid diversification of *Pseudomonas aeruginosa* in cystic

fibrosis lung-like conditions. Proc Natl Acad Sci U S A 115:10714-10719.

40.   Wilder CN, Allada G, Schuster M. 2009. Instantaneous within-patient diversity of

Pseudomonas aeruginosa quorum-sensing populations from cystic fibrosis lung

infections. Infect Immun 77:5631-9.

41.   Smania AM, Segura I, Pezza RJ, Becerra C, Albesa I, Argarana CE. 2004. Emergence

of phenotypic variants upon mismatch repair disruption in *Pseudomonas aeruginosa*.

Microbiology 150:1327-1338.

42.   Ciofu O, Mandsberg LF, Bjarnsholt T, Wassermann T, Hoiby N. 2010. Genetic

adaptation of *Pseudomonas aeruginosa* during chronic lung infection of patients with

cystic fibrosis: Strong and weak mutators with heterogeneous genetic backgrounds

emerge in *mucA* and/or *lasR* mutants. Microbiology 156:1108-1119.

43.   Essar DW, Eberly L, Hadero A, Crawford IP. 1990. Identification and characterization of

genes for a second anthranilate aynthase in *Pseudomonas aeruginosa*:

Interchangeability of the two anthranilate synthases and evolutionary implications. J

Bacteriol 172:884-900.

44.   Caiazza NC, Shanks RMQ, O'Toole GA. 2005. Rhamnolipids modulate swarming

motility patterns of *Pseudomonas aeruginosa*. J Bacteriol 187:7351-7361.

45.   Turnbull L, Whitchurch CB. 2014. Motility assay: Twitching motility. Methods Mol Biol

1149:73-86.

46.   Glessner A, Smith RS, Iglewski BH, Robinson JB. 1999. Roles of *Pseudomonas*

aeruginosa las and rhl quorum-sensing systems in control of twitching motility. J

Bacteriol 181:1623-9.

47.   Higgins Keppler EA, Jenkins CL, Davis TJ, Bean HD. 2018. Advances in the application

of comprehensive two-dimensional gas chromatography in metabolomics. Trends Anal

Chem 109:275-286.

48.   Dieterle F, Ross A, Schlotterbeck G, Senn H. 2006. Probabilistic quotient normalization

as robust method to account for dilution of complex biological mixtures. Application in ^{1}H

NMR metabonomics. Anal Chem 78:4281-4290.

49.   Koo TK, Li MY. 2016. A guideline of selecting and reporting intraclass correlation

coefficients for reliability research. J Chiropr Med 15:155-163.

50.   Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and

powerful approach to multiple testing. J R Stat Soc Series B Methodol 57:289-300.

51.   Johnson WE, Li C, Rabinovic A. 2007. Adjusting batch effects in microarray expression

data using empirical bayes methods. Biostatistics 8:118-27.

52.   Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, Fan TWM, Fiehn O,

721   Goodacre R, Griffin JL, Hankemeier T, Hardy N, Harnly J, Higashi R, Kopka J, Lane AN,

722   Lindon JC, Marriott P, Nicholls AW, Reily MD, Thaden JJ, Viant MR. 2007. Proposed

723   minimum reporting standards for chemical analysis Chemical Analysis Working Group

724   (CAWG) Metabolomics Standards Initiative (MSI). Metabolomics 3:211-221.

725   53.   Bean HD, Dimandja J-MD, Hill JE. 2012. Bacterial volatile discovery using solid-phase

726   microextraction and comprehensive two-dimensional gas chromatography–time-of-flight

727   mass spectrometry. J Chromatogr B 901:41-46.

728   54.   Vogt L, Gröger T, Zimmermann R. 2007. Automated compound classification for ambient

729   aerosol sample separations using comprehensive two-dimensional gas

730   chromatography–time-of-flight mass spectrometry. J Chromatogr A 1150:2-12.

731   55.   Seeley JV, Kramp F, Hicks CJ. 2000. Comprehensive two-dimensional gas

732   chromatography via differential flow modulation. Anal Chem 72:4346-4352.

733   56.   Adahchour M, Beens J, Vreuls RJJ, Brinkman UAT. 2006. Recent developments in

734   comprehensive two-dimensional gas chromatography (GC x GC): IV. Further

735   applications, conclusions and perspectives. Trends Anal Chem 25:821-840.

736   57.   Baptista I, Santos M, Rudnitskaya A, Saraiva JA, Almeida A. 2019. A comprehensive

737   look into the volatile exometabolome of enteroxic and non-enteroxic *Staphylococcus*

738   *aureus* strains. Int J Biochem Cell Biol 108:40-50.

739   58.   Shestivska V, Španêl P, Dryahina K, Sovová K. 2012. Variability in the concentrations of

740   volatile metabolites emitted by genotypically different strains of *Pseudomonas*

741   *aeruginosa*. Journal of Applied Microbiology.

59. Labows JN, McGinley KJ, Webster GF, Leyden JJ. 1980. Headspace analysis of volatile

metabolites of *Pseudomonas aeruginosa* and related species by gas chromatography-

mass spectrometry. J Clin Microbiol 12:521-6.

60. Lawal O, Knobel H, Weda H, Nijsen TME, Goodacre R, Fowler SJ, BreathDx c. 2018.

TD/GC-MS analysis of volatile markers emitted from mono- and co-cultures of

Enterobacter cloacae and *Pseudomonas aeruginosa* in artificial sputum. Metabolomics

14:66.

61. Filipiak W, Sponring A, Baur MM, Filipiak A, Ager C, Wiesenhofer H, Nagl M, Troppmair

J, Amann A. 2012. Molecular analysis of volatile metabolites released specifically by

Staphylococcus aureus and *Pseudomonas aeruginosa*. BMC Microbiol 12:113.

62. Neerincx AH, Geurts BP, Habets MFJ, Booij JA, van Loon J, Jansen JJ, Buydens LMC,

van Ingen J, Mouton JW, Harren FJM, Wevers RA, Merkus P, Cristescu SM, Kluijtmans

LAJ. 2016. Identification of *Pseudomonas aeruginosa* and *Aspergillus fumigatus* mono-

and co-cultures based on volatile biomarker combinations. J Breath Res 10:9.

63. Boots AW, Smolinska A, van Berkel J, Fijten RRR, Stobberingh EE, Boumans MLL,

Moonen EJ, Wouters EFM, Dallinga JW, Van Schooten FJ. 2014. Identification of

microorganisms based on headspace analysis of volatile organic compounds by gas

chromatography-mass spectrometry. J Breath Res 8:12.

64. Zscheppank C, Wiegand HL, Lenzen C, Wingender J, Telgheder U. 2014. Investigation

of volatile metabolites during growth of *Escherichia coli* and *Pseudomonas aeruginosa*

by needle trap-GC-MS. Anal Bioanal Chem 406:6617-28.

65. Senecal AG, Magnone J, Yeomans W, Powers EM. 2002. Rapid detection of pathogenic

bacteria by volatile organic compound (VOC) analysis. Proc SPIE 4575:610-613.

66.   Zechman JM, Aldinger S, Labows JN, Jr. 1986. Characterization of pathogenic bacteria

by automated headspace concentration-gas chromatography. J Chromatogr 377:49-57.

67.   Briard B, Heddergott C, Latge JP. 2016. Volatile compounds emitted by *Pseudomonas*

aeruginosa stimulate growth of the fungal pathogen *Aspergillus fumigatus*. mBio

7:e00219.

68.   Preti G, Thaler E, Hanson CW, Troy M, Eades J, Gelperin A. 2009. Volatile compounds

characteristic of sinus-related bacteria and infected sinus mucus: Analysis by solid-

phase microextraction and gas chromatography-mass spectrometry. J Chromatogr B

Analyt Technol Biomed Life Sci 877:2011-8.

69.   Allardyce RA, Langford VS, Hill AL, Murdoch DR. 2006. Detection of volatile metabolites

produced by bacterial growth in blood culture media by selected ion flow tube mass

spectrometry (SIFT-MS). J Microbiol Methods 65:361-5.

70.   Rudzinski CM, Herzig-Marx R, Lin J, Szpiro A, Johnson B. 2004. Pathogen detection

using headspace analysis abstr Scientific Conference on Chemical and Biological

Defense Research, Hunt Valley, MD, MIT Lincoln Laboratory,

71.   Carroll W, Lenney W, Wang T, Spanel P, Alcock A, Smith D. 2005. Detection of volatile

compounds emitted by *Pseudomonas aeruginosa* using selected ion flow tube mass

spectrometry. Pediatr Pulmonol 39:452-6.

72.   Thorn RM, Reynolds DM, Greenman J. 2011. Multivariate analysis of bacterial volatile

compound profiles for discrimination between selected species and strains *in vitro*. J

Microbiol Methods 84:258-64.

73.   Schöller C, Molin S, Wilkins K. 1997. Volatile metabolites from some gram-negative
787   bacteria. Chemosphere 35:1487-95.

788   74.   Lammers M, Kos R, De Vries R, Brinkman P, Dagelet JWF, Rutjes NW, Majoor CJ,
789   Terheggen-Lagro SWJ, Weersink EJM, Bos LDJ, Sterk PJ, Maitland-Van Der Zee A-H,
790   Neerincx AH, Group TACR. 2018. Detection of *Pseudomonas aeruginosa* infections in
791   cystic fibrosis patients by enose technology. 52:PA3412.

792   75.   Neerincx AH, Brinkman P, Van der Gouwe P, Paff T, Fens N, Gerritsen MG, Haarman E,
793   Van Aalderen WMC, Majoor CJ, Weersink EJM, Bos LD, Sterk PJ. 2018. Detection of
794   *Pseudomonas aeruginosa* in exhaled breath of cystic fibrosis patients. Eur Respir J 52.

795   76.   Pabary R, Huang J, Kumar S, Alton EW, Bush A, Hanna GB, Davies JC. 2016. Does
796   mass spectrometric breath analysis detect *Pseudomonas aeruginosa* in cystic fibrosis?
797   Eur Respir J 47:994-7.

798   77.   Gilchrist FJ, Belcher J, Jones AM, Smith D, Smyth AR, Southern KW, Spanel P, Webb
799   AK, Lenney W. 2015. Exhaled breath hydrogen cyanide as a marker of early
800   *Pseudomonas aeruginosa* infection in children with cystic fibrosis. ERJ Open Res 1.

801   78.   Gilchrist FJ, Bright-Thomas RJ, Jones AM, Smith D, Spanel P, Webb AK, Lenney W.
802   2013. Hydrogen cyanide concentrations in the breath of adult cystic fibrosis patients with
803   and without *Pseudomonas aeruginosa* infection. J Breath Res 7:026010.

804   79.   Robroeks C, van Berkel J, Dallinga JW, Jobsis Q, Zimmermann LJI, Hendriks HJE,
805   Wouters MFM, van der Grinten CPM, van de Kant KDG, van Schooten FJ, Dompeling E.
806   2010. Metabolomics of volatile organic compounds in cystic fibrosis patients and
807   controls. Pediatric Research 68:75-80.

808   80.   Scott-Thomas AJ, Syhre M, Pattemore PK, Epton M, Laing R, Pearson J, Chambers ST.

809   2010. 2-aminoacetophenone as a potential breath biomarker for *Pseudomonas*

810   *aeruginosa* in the cystic fibrosis lung. BMC Pulmonary Medicine 10:10.

811   81.   Enderby B, Smith D, Carroll W, Lenney W. 2008. Hydrogen cyanide as a biomarker for

812   *Pseudomonas aeruginosa* in the breath of children with cystic fibrosis. Pediatr Pulmonol

813   44:142-147.

814   82.   Lemfack MC, Gohlke BO, Toguem SMT, Preissner S, Piechulla B, Preissner R. 2018.

815   mVOC 2.0: A database of microbial volatiles. Nucleic Acids Res 46:D1261-D1265.

816   83.   Dryahina K, Sovová K, Nemec A, Španěl P. 2016. Differentiation of pulmonary bacterial

817   pathogens in cystic fibrosis by volatile metabolites emitted by their *in vitro* cultures:

818   *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Stenotrophomonas maltophilia* and

819   the *Burkholderia cepacia* complex. J Breath Res 10:037102.

820   84.   Zhu J, Bean HD, Jiménez-Díaz J, Hill JE. 2013. Secondary electrospray ionization-mass

821   spectrometry (SESI-MS) breathprinting of multiple bacterial lung pathogens, a mouse

822   model study. J Appl Physiol 114:1544-1549.

823   85.   Zhu J, Jiménez-Díaz J, Bean HD. 2013. Robust detection of *P. aeruginosa* and *S.*

824   *aureus* acute lung infections by secondary electrospray ionization-mass spectrometry

825   (SESI-MS) breathprinting: From initial infection to clearance. J Breath Res 7.

826   86.   Hahn A, Whiteson K, Davis TJ, Phan J, Sami I, Koumbourlis AC, Freishtat RJ, Crandall

827   KA, Bean HD. 2020. Longitudinal associations of the cystic fibrosis airway microbiome

828   and volatile metabolites: A case study. Front Cell Infect Microbiol 10.

87.   Hoffman LR, Kulasekara HD, Emerson J, Houston LS, Burns JL, Ramsey BW, Miller SI.

2009. *Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis lung

disease progression. J Cyst Fibros 8:66-70.

88.   Ren CL, Konstan MW, Yegin A, Rasouliyan L, Trzaskoma B, Morgan WJ, Regelmann

W, Scientific Advisory Group I, Coordinators of the Epidemiologic Study of Cystic F.

2012. Multiple antibiotic-resistant *Pseudomonas aeruginosa* and lung function decline in

patients with cystic fibrosis. J Cyst Fibros 11:293-9.

89.   Hunter RC, Klepac-Ceraj V, Lorenzi MM, Grotzinger H, Martin TR, Newman DK. 2012.

Phenazine content in the cystic fibrosis respiratory tract negatively correlates with lung

function and microbial complexity. Am J Respir Cell Mol Biol 47:738-745.

90.   Grosso-Becerra MV, Santos-Medellín C, González-Valdez A, Méndez JL, Delgado G,

Morales-Espinosa R, Servín-González L, Alcaraz LD, Soberón-Chávez G. 2014.

Pseudomonas aeruginosa clinical and environmental isolates constitute a single

population with high phenotypic diversity. BMC Genomics 15.

91.   Bragonzi A, Worlitzsch D, Pier GB, Timpert P, Ulrich M, Hentzer M, Andersen JB,

Givskov M, Conese M, Doring G. 2005. Nonmucoid *Pseudomonas aeruginosa*

expresses alginate in the lungs of patients with cystic fibrosis and in a mouse model. J

Infect Dis 192:410-419.

92.   Moskowitz SM, Emerson JC, McNamara S, Shell RD, Orenstein DM, Rosenbluth D,

Katz MF, Ahrens R, Hornick D, Joseph PM, Gibson RL, Aitken ML, Benton WW, Burns

JL. 2011. Randomized trial of biofilm testing to select antibiotics for cystic fibrosis airway

infection. Pediatr Pulmonol 46:184-192.

93.   Cornforth DM, Dees JL, Ibberson CB, Huse HK, Mathiesen IH, Kirketerp-Moller K,

Wolcott RD, Rumbaugh KP, Bjarnsholt T, Whiteley M. 2018. *Pseudomonas aeruginosa*

transcriptome during human infection. Proc Natl Acad Sci U S A 115:E5125-E5134.

94.   Harrison F, Muruli A, Higgins S, Diggle SP. 2014. Development of an *ex vivo* porcine

lung model for studying growth, virulence, and signaling of *Pseudomonas aeruginosa*.

Infect Immun 82:3312-3323.

95.   Bayes HK, Ritchie N, Irvine S, Evans TJ. 2016. A murine model of early *Pseudomonas*

aeruginosa lung disease with transition to chronic infection. Sci Rep 6.

96.   Crabbé A, Ledesma MA, Nickerson CA. 2014. Mimicking the host and its

microenvironment *in vitro* for studying mucosal infections by *Pseudomonas aeruginosa*.

Pathog Dis 71:1-19.

97.   Sønderholm M, Kragh KN, Koren K, Jakobsen TH, Darch SE, Alhede M, Jensen PO,

Whiteley M, Kühl M, Bjarnsholt T. 2017. *Pseudomonas aeruginosa* aggregate formation

in an alginate bead model system exhibits *in vivo*-like characteristics. Appl Environ

Microbiol 83.

98.   Bean HD, Jiménez-Díaz J, Zhu J, Hill JE. 2015. Breathprints of model murine bacterial

lung infections are linked with immune response. Eur Respir J 45:21-24.

870 **FIG 1** Relative phenotype production of 81 *P. aeruginosa* CF chronic lung infection isolates.

871 Each phenotype was independently scaled to a range of 0 to 1, with 0 representing the

872 minimum data point and 1 representing the maximum data point (excluding outliers), and the

873 relative productions are depicted in the heat map as standardized values (mean-centered and

874 scaled to unit variance) across isolates. Phenotype Score ("Score") is the sum of the scaled,

875 non-standardized phenotype values, rounded to the nearest integer.

877 **FIG 2** Size and chemical composition of volatile compounds metabolized by 81 *P. aeruginosa*

878 CF chronic infection isolates. **(a)** Identified compounds; **(b)** Identified and unknown compounds;

879 **(c)** Identified compounds, scaled to 100%; **(d)** Identified and unknown compounds, scaled to

880 100%. C = Core compounds, defined as detected in 95% or more of samples; NC = Non-core

881 compounds; C+NC = sum of Core and Non-core compounds.

FIG 3 Violin plots and box plots of pairwise dissimilarities (defined as Euclidean distance) of *P.*

aeruginosa chronic infection isolate volatilomes. Inter-Early are between-patient Early-Early

isolate dissimilarities. Inter-Late are between-patient Late-Late isolate dissimilarities. Intra-

Patient are within-patient Early-Late isolate dissimilarities. Pooled represents all pairwise

volatilome dissimilarities. Red circles indicate the mean. Significant differences between means

were identified using one-way ANOVA with Tukey's HSD. *P < .05, **P < .01, ***P < .001.

FIG 4 Ordered dissimilarity image (ODI) of the truncated set of 48 *P. aeruginosa* clinical isolates

depicting volatilome dissimilarity defined by Euclidean distance.

FIG 5 Ordered dissimilarity image (ODI) of the 35 *P. aeruginosa* clinical isolates from Patient 23

depicting volatilome dissimilarity defined by Euclidean distance. For visual clarity, the isolate

labels do not include patient number (*e.g.*, 23).

FIG 6 Dissimilarity of Patient 23 volatilomes over time, defined as Euclidean distance from the

Early isolate (not shown). For visual clarity, the isolate labels do not include patient number

(*e.g.*, 23) and intermediate isolate labels have been truncated to only include the numeral that

signifies collection order (*e.g.*, I-1 is represented as 1). The black line represents the linear

regression fit (r = 0.70, *P* = 0+), and the shaded region represents the standard error of the

regression line.

FIG 7 Representative GC×GC chromatograms of Early (top) and Late (bottom) chronic infection

isolates (41E and 41L-2 are depicted). Dark blue represents the baseline, and peak intensity is

depicted using a color gradient from light blue (low) to dark red (high). Chromatographic regions

of ^1t_R < 200 s and ^2t_R < 0.5 s were excluded for visual clarity. All chromatograms for the ten

selected Early and Late isolate pairs are provided in Supplementary Fig. 7.

FIG 8 Hierarchical clustering analysis (HCA) of the truncated set of 48 *P. aeruginosa* clinical CF

isolates, based on the relative abundance of 539 volatile compounds. Volatiles are in columns

(standardized relative abundance). Clustering is based on rows (isolates), which are color-

coded by their phenotype score (left color block) and relative time of collection (right color

block). ALC = alcohols, ALD = aldehydes, ARO = aromatics, CA/EST = carboxylic acids and

esters, HC = hydrocarbons, KET = ketones, THI = thiols, OTH = other compounds, UNK =

unknown identity.

FIG 9 Hierarchical clustering analysis (HCA) of the truncated set of 48 *P. aeruginosa* clinical CF

isolates, based on the relative abundance of 23 core alcohols, aldehydes, and hydrocarbons.

Volatiles are in columns (standardized relative abundance). Clustering is based on rows

(isolates), which are color-coded by their phenotype score (left color block) and relative time of

collection (right color block). ALC = alcohols, ALD = aldehydes, HC = hydrocarbons.

FIG S1. Non-metric multidimensional scaling (NMDS) plots of 81 *P. aeruginosa* clinical

isolates depicting volatilome dissimilarity defined by Euclidean distance **(a)** colored by

collection time and **(b)** colored by patient; **(c)** Ordered dissimilarity image (ODI) of 81 *P.*

aeruginosa clinical isolates depicting volatilome dissimilarity defined by Euclidean

distance.

FIG S2. Non-metric multidimensional scaling (NMDS) plots of the truncated set of 48 *P.*

aeruginosa clinical isolates depicting volatilome dissimilarity defined by Euclidean

distance **(a)** colored by collection time and **(b)** colored by patient; **(c)** NMDS plot of the

35 *P. aeruginosa* clinical isolate volatilomes from Patient 23 depicting volatilome

dissimilarity defined by Euclidean distance.

FIG S3. Size and chemical composition of volatile compounds metabolized by the

selected ten Early-Late pairs of *P. aeruginosa* chronic infection CF isolates. **(a)**

Identified compounds; **(b)** Identified and unknown compounds; **(c)** Identified

compounds, scaled to 100%; **(d)** Identified and unknown compounds, scaled to 100%.

Size and chemical composition of volatile compounds metabolized by each of the ten

Early-Late pairs of *P. aeruginosa* chronic infection CF isolates. **(e)** Identified

compounds; **(f)** Identified and unknown compounds; **(g)** Identified compounds, scaled to

100%; **(h)** Identified and unknown compounds, scaled to 100% C = Core compounds,

defined as detected in 95% or more of samples; NC = Non-core compounds; C+NC =

sum of Core and Non-core compounds.

FIG S4. Hierarchical clustering analysis (HCA) of the ten Early-Late pairs of *P.*

aeruginosa clinical CF isolates, based on **(a)** presence and absence and **(b)** relative

abundance of 472 volatile compounds. Core compounds are present in at least 95% of

isolates; Non-core compounds are present in less than 95%. Volatiles are in columns.

Clustering is based on rows (isolates), which are color-coded by their phenotype score

(left color block) and relative time of collection (right color block).

FIG S5. Hierarchical clustering analysis (HCA) of the 81 *P. aeruginosa* clinical CF

isolates, based on the relative abundance of 539 volatile compounds. Core compounds

are present in at least 95% of isolates; Non-core compounds are present in less than

95%. Volatiles are in columns (standardized relative abundance). Clustering is based on

rows (isolates), which are color-coded by their phenotype score (left color block) and

relative time of collection (right color block).

FIG S6. Hierarchical clustering analysis (HCA) of **(a)** the truncated set of 48 and **(b)** full

set of 81 *P. aeruginosa* clinical CF isolates, based on the relative abundance of the 69

core volatile compounds. Volatiles are in columns (standardized relative abundance).

Clustering is based on rows (isolates), which are color-coded by their phenotype score

(left color block) and relative time of collection (right color block).

FIG S7. Hierarchical clustering analysis (HCA) of the 81 *P. aeruginosa* clinical CF

isolates, based on the relative abundance of 23 core alcohols, aldehydes, and

hydrocarbons. Volatiles are in columns (standardized relative abundance). Clustering is

based on rows (isolates), which are color-coded by their phenotype score (left color

block) and relative time of collection (right color block).