MAGI1 inhibits the AMOTL2/p38 stress pathway and prevents luminal breast tumorigenesis

Diala Kantar [1], Emilie Bousquet Mur [1,+], Vera Slaninova [1,+], Yezza Ben Salah [1], Luca Costa [3], Elodie Forest

[1], Patrice Lassus [1], Charles Géminard [1], Marion Larroque [1,2], Florence Boissière-Michot [1,2], Béatrice

Orsetti [1], Charles Theillet [1], Jacques Colinge [1], Christine Benistant [3], Antonio Maraver [1], Lisa Heron-

Milhavet [1,*], and Alexandre Djiane [1,*]

[1] IRCM, Univ Montpellier, Inserm, ICM, Montpellier, France. [2] Translational Research Unit, ICM,

Montpellier, France. [3] CBS, Univ Montpellier, CNRS, Inserm, Montpellier, France.

[+] Co-second authors: E.B. and V.S.

[*] Co-last authors and co-corresponding authors: L.H.M. and A.D.

[*] Authors to whom all correspondence and material requests should be addressed: L.H.M. and A.D.

(Emails: lisa.heron-milhavet@inserm.fr / alexandre.djiane@inserm.fr)

ABSTRACT

Alterations to cell polarization or to intercellular junctions are often associated with epithelial cancer

progression, including breast cancers (BCa). We show here that the loss of the junctional scaffold

protein MAGI1 is associated with bad prognosis in luminal BCa, and promotes tumorigenesis. E-

cadherin and the actin binding scaffold AMOTL2 accumulate in *MAGI1* deficient cells which are

subjected to increased stiffness. These alterations are associated with low YAP activity, the terminal

Hippo-pathway effector, but with an elevated ROCK and p38 Stress Activated Protein Kinase activities.

Blocking ROCK prevented p38 activation, suggesting that MAGI1 limits p38 activity in part through

releasing actin strength. Importantly, the increased tumorigenicity of *MAGI1* deficient cells is rescued

in the absence of AMOTL2 or after inhibition of p38, demonstrating that MAGI1 acts as a tumor-

suppressor in luminal BCa by inhibiting an AMOTL2/p38 stress pathway.

1   INTRODUCTION

Breast cancers (BCa) arise from mammary epithelial cells, and show tremendous diversity.

Based on transcriptome, BCa are typically divided into five main subtypes: luminal A, luminal B, HER2-

enriched, normal-like, and basal (reviewed in [1] [2]), partly overlapping with previous classifications (such

as ER/PR/HER2 receptors expression status). Basal tumors, overlapping strongly with triple-negative

breast cancers, are the most aggressive and present the worst prognosis. However, even though

luminal A and B subtypes (typically ER/PR positives) usually respond well to current treatments, when

"luminal" tumors relapse, they become very difficult to treat with very poor outcome.

The apico-basal (A/B) polarity of mammary epithelial cells is established and maintained by the

asymmetric segregation of evolutionarily conserved protein complexes [3]. A/B polarity governs the

correct position of different intercellular junctions that ensure the integrity of the epithelial sheet.

Adherens Junctions (AJs) constitute apical adhesive structures mediated by the homophilic trans-

interactions of E-cadherin/catenin complexes [4] [5]. At the very apical part of epithelial cells, Tight

junctions (TJs) ensure the impermeability of tissues in part through the action of Claudins/Zonula

Occludens proteins complexes [6]. The transmembrane proteins of these different junctions (E-cadherin,

Claudins…) are coupled and anchored to the actin cytoskeleton via adaptor scaffold proteins, thus

coupling adhesion and cell contacts to actin dynamics. As such AJs represent mechano-sensory

structures, where tensile or compressive forces, arising from internal or external sources, will be

sensed to promote short term responses such as changes in adhesion and actomyosin dynamics, but

also long term responses to control cell density, in particular through the mechano-sensitive Hippo

pathway [7,8]. In epithelial cells, the compressive forces exerted by the cortical actin cytoskeleton or by

cell packing, sensed at AJs, antagonize the proliferative and anti-apoptotic effects of YAP/TAZ in part

through α-catenin conformational change, or through the release of Merlin/NF2 [9–14]. Other signaling

pathways such as β-catenin, ERK, or JNK/SAPK stress pathways can also be influenced by polarity and

forces sensed at junctions [11,15–18]. BCa are derived from transformed mammary epithelia where

frequent alterations in TJs and/or AJs composition and structure, and the associated abnormal

mechano-sensory responses have been linked to cell dysfunction such as uncontrolled cell

proliferation and metastasis [19,20].

At apical junctions, cortical scaffold proteins such as MAGI (MAGUKs with Inverted domain

structure), assemble large and dynamic molecular complexes. In humans, among the three MAGI

proteins (MAGI1-3), MAGI1 is the most widely expressed and contains several protein-protein

interaction domains including six PDZ and two WW domains [21] mediating interactions with TJ proteins

[22]. In *Drosophila*, we have demonstrated that Magi, the sole fly MAGI homolog, is required for E-

cadherin belt integrity and AJ dynamics ultimately restricting cell numbers in the developing eye [23,24]. The regulation of Cadherin complexes by MAGIs, is evolutionarily conserved from C. elegans [25], to vertebrates, and in particular vertebrate endothelial cells where MAGI1 links VE-cadherin complexes with the underlying cortical actin cytoskeleton [26]. A tumor suppressive role of MAGI1 (reviewed by Feng et al. [27]) is suggested by the correlation between low *MAGI1* expression and poor prognosis in various cancers, such as hepatocellular carcinoma [28]. The anti-tumoral activity of MAGI1 is further supported by its ability to bind the tumor-suppressor PTEN [29–31], the fact that it is often targeted by viral oncoproteins [32,33], and by the observation that in colorectal cancer cells, MAGI1 was upregulated in response to cyclooxygenase-2 inhibitor and prevented metastasis [34]. Thus it appeared that the apical junction-localized MAGI1 scaffold protein participates in multiple complexes to fine-tune adhesion and signaling, and may therefore be considered as a tumor suppressor.

The angiomotin (AMOT) family of membrane-associated scaffold proteins is composed of three members: AMOT, AMOTL1, and AMOTL2, playing important roles in the regulation of intercellular junctions [35]. In endothelial cells and in Zebrafish developing embryos, AMOTL2 has been shown to link cadherins and the actin cytoskeleton. In particular, AMOTL2 is required for the maintenance of tension (mainly arising from compressive forces) at the level of cadherin junctions to properly shape strong blood vessels [26,36]. Moreover, AMOTs are also known to control the Hippo tumor-suppressor signaling pathway. Canonical Hippo pathway, through sequential activation of the Hippo/MST and Wts/LATS kinases, promotes the phosphorylation and cytoplasmic retention of the transcriptional co-activators Yki/YAP/TAZ (reviewed by [37]). Specific PPxY motifs found in AMOTs interact with the WW domains of YAP and TAZ [38] and AMOTs act thus as negative regulators of YAP/TAZ mediated transcription by trapping YAP [38,39] and TAZ [40] in the cytoplasm. Recent studies identified interactions between MAGI1 and AMOTs, in particular in the regulation of intercellular junctions [41] [42,43].

We report here that low levels of *MAGI1* are associated with bad prognosis in luminal ER+ BCa. Impairing MAGI1 expression in luminal BCa cells promoted their growth in 2D, and their ability to grow in low attachment conditions in-vitro and in subcutaneously grafted nude mice. We further document that *MAGI1* deficient cells accumulate specific junctional proteins, including E-cadherin and AMOTL2. Mechanistically, we show that all *MAGI1* deficient phenotypes are suppressed by down-regulating *AMOTL2* suggesting that AMOTL2 and its effects on junctions is the primary cause of the increased tumorigenicity after MAGI1 loss. Consistent with these observations, we could further show that *MAGI1* deficient luminal BCa cells experience higher stiffness as evidenced by increased Young's modulus, and ROCK activity as evidenced by increased ROCK-specific Ser19 phosphorylation of the regulatory Myosin Light Chain 2. They activate the p38 stress signaling pathway while YAP activity is

1   antagonized. Finally, we provide evidence that releasing actin cytoskeletal strength, or impairing p38

2   activity can revert the effects of MAGI1 loss, supporting a model by which, in response to MAGI1 loss,

3   elevated AMOTL2/E-cadherin and junction dysfunction, together with actin cytoskeletal tension

4   activate the p38 stress pathway fueling tumorigenicity.

MATERIALS & METHODS

Plasmids, mutant constructs and shRNA cloning

All expression plasmids were generated with the Gateway system (Invitrogen). The Entry vector

pDONR223-MAGI1 was a gift from William Hahn (Addgene plasmid #23523) [44] and it was used to

introduce the different point mutations in *MAGI1* by mutagenesis PCR using PfuTurbo (Agilent). All the

Gateway destination vectors that were generated are listed as follows : *pCMV10 3xFlag RfB-MAGI1,*

pCMV10 3xFlag RfB-MAGI1 P331A, pCMV10 3xFlag RfB-MAGI1 P390A and *pCMV10 3xFlag RfB-MAGI1-*

P331/390A. *pEZY-EGFP-MAGI1* was constructed by Gateway recombination between the *pDONR223-*

MAGI1 and the *pEZY-EGFP* destination vector that was a gift from Zu-Zhu Zhang (Addgene plasmid

#18671) [45]. *pcDNA3 HA-AMOT, pcDNA3 HA-AMOT Y242A, pcDNA3 HA-AMOT Y287A* and *pcDNA3 HA-*

AMOT Y242/287A were gifts from Kunliang Guan (Addgene plasmids #32821, #32823, #32824 and

#32822 respectively) [38].

shRNA directed against human *MAGI1* and *AMOTL2* were constructed and cloned in the pSIREN-

RetroQ vector (TaKaRa) according to the manufacturer's conditions between BamHI and EcoRI cloning

sites as previously described for other genes [46]. Targeted sequences were:

shRNA-MAGI1(1-1): CACCTATGAAGGAAACTATT

shRNA-MAGI1(3-1): GATCTCATAGTGGAAGTTAA

shRNA-AMOTL2(1416): GGAACAAGATGGACAGTGA

shRNA-AMOTL2(3667): GAGATGTCTTGTTAGCATA

All hairpins were validated (http://cancan.cshl.edu/cgi-bin/Codex/Codex.cgi). The shRNA targeting

Luciferase as a control was kindly provided by Celine Gongora (IRCM). Retroviral particles were

produced in HEK293 cells and used to infect MCF7 and T47D (and HCT116) cell lines that were then

selected with 1 µg/ml puromycin.

Cell culture and cell transfection

MCF7, T47D and HCT116 human cell lines originated from the *TumoroteK* bank (SIRIC Montpellier

Cancer) and have been authenticated by STR (Short-tandem repeat) profiling and certified mycoplasm-

free. MCF7 cell lines were grown respectively in DMEM:HAM F12 or DMEM medium supplemented

with penicillin and streptomycin antibiotics (1%) and fetal calf serum (2% or 10%). T47D and HCT116

were grown in RPMI medium supplemented with antibiotics (1%) and fetal calf serum (2 % or 10%).

MCF7, T47D and HCT116 shRNA cell lines were grown in their respective medium in the presence of puromycin (1 µg/mL) for selection of cells expressing shRNAs.

Plasmid transfections were performed in MCF7 cells using Lipofectamine2000 reagent (Invitrogen) according to the manufacturer's directions. Cell extracts were assayed for protein expression 24-48 hours post-transfection. Concerning *AMOTL2*siRNA transfection (Dharmacon siGENOME #M-01323200-0005), a final concentration of 100 nM was used using Lipofectamine2000 reagent as well.

Western blotting

Proteins issued from transfected MCF7, MCF7shRNA, T47DshRNA or HCT116shRNA cell lines were extracted, analyzed by SDS-PAGE. Antibodies used were mouse anti-MAG1 (1/250; Santa Cruz #sc100326), rabbit anti-MAGI1 (1/1000; Sigma #HPA031853), rabbit anti-MAGI2 (1/250; Santa Cruz #sc25664), rabbit anti-MAGI3 (1/250; Santa Cruz discontinued), mouse anti-tubulin (1/10000; Sigma-Aldrich #T6074), mouse anti-AMOT (1/250; Santa Cruz #sc166924), rabbit AMOTL2 (1/600; Proteintech #23351-1-AP), mouse M2 anti-Flag (1/2000; Sigma-Aldrich #F1804), mouse anti-HA (1/2000; BioLegend #901501), rabbit anti-GFP (1/2000; Torrey Pines Laboratories #TP401), rabbit anti-YAP/TAZ (1/1000; Cell Signaling Technology #8418), rabbit anti YAP (1/1000; Cell Signaling Technology #14074), p-YAP (S127) (1/1000; Cell Signaling Technology #13008) and p-YAP (S397) (1/1000; Cell Signaling Technology #13619), rabbit anti LATS1 (1/1000; Cell Signaling Technology #3477), p-LATS1 (1/1000; Cell Signaling Technology #8654), p-Mob1 (1/1000; Cell Signaling Technology #8699) mouse anti-actin (1/250; Developmental Studies Hybridoma Bank #JLA20), goat anti-SCRIB (1/250; Santa Cruz #sc11049), rabbit anti-Claudin-1 (1/1000; Cell Signaling Technology #13255), anti-Claudin-3 (1/1000; Genetex #15102), anti-ZO-1 (1/1000; Cell Signaling Technology #8193), anti-β-catenin (1/1000; Cell Signaling Technology #8480), anti-PARD3 (1/1000; Millipore #07-330), anti-E-cadherin (1/1000; Cell Signaling Technology #3195), anti-p-p38 (Cell Signaling Technology #4511), anti-p38 (Cell Signaling Technology #8690), anti-p-JNK/SAPK (Cell Signaling Technology #9251), anti-JNK/SAPK (Cell Signaling Technology #9252), anti-p-MLC Ser19 (Cell Signaling Technology #3675), anti-MLC (Cell Signaling Technology #3672), anti-p-Akt (Cell Signaling Technology #4060), anti-Akt (Cell Signaling Technology #4691), anti-p-ERK1/2 (Cell Signaling Technology #4370) and anti-ERK1/2 (Cell Signaling Technology #4695).

RNA extraction, reverse transcription and real time RT-qPCR

RNAs were extracted from cells using RNeasy plus kit (Qiagen) and cDNAs were prepared starting from 1 µg of RNAs using the Superscript III reverse transcriptase (Invitrogen) and following manufacturer's directions. Real time quantitative PCR was then performed on cDNAs using SYBR Green I Mastermix (Roche) according to the manufacturer's conditions on a Light Cycler 480 device (Roche). All primers used are listed as follows, 5' to 3':

MAGI1_For: CGTAAAGTGGTTTTTGCGGTGC

MAGI1_Rev: TCTCCACGTCGTAGGGCTGC

MAGI2_For: ATCATTGGTGGAGACGAGCC

MAGI2-Rev: TAGCCACGACACAACACCAG

MAGI3_For: CTGCACTTTTCAGTCTTCTTTTGAC

MAGI3_Rev: CTGAACCAAATTACGTGGCCC

AMOTL2_For: CCAAGTCGGTGCCATCTGTT

AMOTL2_Rev: CCATCTCTGCTCCCGTGTTT

CTGF_For: TTCCAAGACCTGTGGGAT

CTGF_Rev: GTGCAGCCAGAAAGCTC

CYR61_For: ACCAAGAAATCCCCCGAACC

CYR61_Rev: CGGGCAGTTGTAGTTGCATT

BIRC2_For: GTCAGAACACCGGAGGCATT

BIRC2_Rev: TGACATCATCATTGCGACCCA

AREG_For: CGAAGGACCAATGAGAGCCC

AREG_Rev: AGGCATTTCACTCACAGGGG

HPRT_For: CTGACCTGCTGGATTACA

HPRT_Rev: GCGACCTTGACCATCTTT

Patients and TMA construction

Breast cancer samples were retrospectively selected from the Institut régional du Cancer de Montpellier (ICM) pathology database using the following inclusion criteria: chemotherapy-naïve at the time of surgery and estrogen (ER), progesterone (PR) receptor and HER2 status available. The tissue micro-array (TMA) was constructed to encompass the four subtypes of breast cancer. Fifty samples were identified: 14 with hormone receptor positive expression (>10% of tumor cells expressing ER and PR) and HER2-negative (scored 0 by immunohistochemistry), 12 with hormone receptor positive but HER2-positive (scored 3+ by immunohistochemistry), 10 hormone receptor negative HER2 positive and 14 triple negative samples (ER-,PR- and HER2-negative). Tumor samples were collected following French laws under the supervision of an investigator and declared to the French Ministry of Higher

Education and Research (Biobanque BB-0033-00059; declaration number DC-2008–695). All patients

were informed about the use of their tissue samples for biological research and the study was

approved by the local translational research committee.

Tissue blocks with enough material upon gross inspection were initially selected and then the presence

of breast carcinoma was evaluated on hematoxylin–eosin-stained sections. The areas to be used for

the construction of the TMA were marked on the slide and on the donor block. Samples corresponding

to the selected areas were extracted using a manual arraying instrument (Manual Tissue Arrayer 1,

Beecher Instruments, Sun Prairie, WI, USA). To take into account the tumor heterogeneity, tumor

sampling consisted of two cores (1 mm in diameter) from different tumor areas from a single specimen,

and placed at the specified coordinates.

Immunohistochemistry

Four µm thin sections of the TMA were mounted on Flex microscope slides (Dako) and allowed to dry

overnight at room temperature before immunohistochemistry (IHC) processing.PT-Link® system

(Dako) was used for pre-treatment, allowing simultaneous de-paraffinization and antigen retrieval.

Heat-induced antigen retrieval was executed for 15 minutes in High pH Buffer (Dako) at 95°C.

Immunohistochemistry procedure was performed using the Dako Autostainer Link48 platform. Briefly,

endogeneous peroxidase was quenched using Flex Peroxidase Block (Dako) for 5 min at room

temperature. TMA sections were incubated with rabbit polyclonal antibodies against MAGI1 (1/50;

Sigma Aldrich #HPA0311853) and YAP (1/100; Santa Cruz #15407) at room temperature. Following an

amplification step with a rabbit linker (Dako) and two rinses in buffer, the slides were incubated with

a horseradish peroxidase-labeled polymer coupled to secondary anti-mouse and anti-rabbit antibodies

for 20 min, followed by appliance of 3,3'-Diaminobenzidine for 10 min as substrate. Counterstaining

was performed using Flex Hematoxylin (Dako) followed by washing the slides under tap water for 5

min. Finally, slides were mounted with a coverslip after dehydration. MCF7 cell line transfected with

Flag-MAGI1 expression vector was used as positive control for MAGI1 IHC.

Staining intensity of each marker was performed in a blinded fashion by two trained observers with

the following criteria: 0 (negative), 1 (weak), 2 (moderate), 3 (strong). When both spots were

assessable, data were consolidated into a single score as the mean of the duplicate.

Immunofluorescence

Cells seeded on glass coverslips were fixed 10 min in paraformaldehyde (4 %), before being

permeabilized in PBS / 0.1% TritonX-100 for 10 min. After blocking in PBS / 0.5% BSA, cells were

incubated with primary antibodies overnight at 4C. Antibodies used were mouse anti-MAGI1 (1/100;

Sigma Aldrich #HPA031853), rabbit anti-AMOT (1/100; Proteintech #24550-I-AP), rabbit anti AMOTL2

(1/100; Atlas Antibodies #HPA063027), rabbit anti-E-cadherin (1/100; Cell Signaling Technology

#3195), mouse anti-E-cadherin (1/200; BD Transduction Lab #610182), phalloidin (1/200; Sigma-

Aldrich #P1951), rabbit anti-β-catenin (1/1000; Cell Signaling Technology #8480), rabbit anti-Claudin-

3 (1/100; Genetex #GTX15102), anti-PARD3 (1/1000; Millipore #07-330) and anti-ZO-1 (1/100; Cell

Signaling Technology #8193). Secondary Alexa Fluor Antibodies (1/200; Invitrogen) were used as

described previously [24] for 1 hour at room temperature before mounting the coverslips with

Vectashield (Vector Laboratories #H-1200) and imaging on a Zeiss Apotome microscope.

Atomic Force Microscopy (AFM) measurements

Cells were plated on glass 35mm fluorodish at intermediate (1/3) dilution density, grown overnight,

and placed in 2% serum medium before AFM experiments. Cells grew as clusters of cells. AFM Force

Spectroscopy (AFM-FS) experiments [47] on living cell clusters were performed on a JPK Nanowizard 4

microscope equipped with a CellHesion stage (JPK). The AFM is mounted on an inverted Axiovert 200M

microscope system (Carl Zeiss) equipped with a 100x (1.5NA, plan-Apochromat) objective lens (Carl

Zeiss). We employed a heating module (JPK) placed at the sample level to maintain cells at the

physiological temperature of 37°C during measurements. We used CP-CONT-SIO-D cantilevers with

10μm colloidal beads as tip (NanoAndMore). Cantilever stiffness and optical lever sensitivities were

both calibrated in liquid environment using the Contact-Free-Method provided by JPK AFM, and based

on a mix of a thermal and Sader [48] calibration methods. Calibrated springs constant for cantilevers

were evaluated in the range of 0.15-0.24 N/m. AFM-FS indentation cycles were performed using a 5-

10nN force threshold to induce 2-3μm (approx. 20% of cell height) maximal indentation lengths. A

squared grid of 4x4 pixels covering a region of 50μm X 50μm was fixed at the center of cell cluster

monolayers, defining a force map constituted by 16 indentation curves. At least 5 force maps were

acquired in each experiment on each cell category and 3 separated experiments were performed.

Analyses were carried out using the JPK AFM data processing software. The elastic Young's modulus

(E; Pa) was evaluated by fitting each force versus tip-cell distance curve with the Hertz contact model

for indenting an infinite isotropic elastic half space with a solid sphere as described in [49].

Cell cycle analysis

MCF7shRNA cell lines were fixed with ethanol and stained with propidium iodide for cell cycle analysis according to the on-line technical protocol (BioLegend http://www.biolegend.com/media_assets/support_protocol) and analyzed using a Gallios Flow cytometer (Beckman Coulter).

Mammosphere culture

MCF7*shLuc* and MCF7*shMAGI1* cell suspensions were obtained by trypsinization and then filtered using 30μm cell strainer (Miltenyi). Filtered cells were immediately seeded in 96 well plates coated with 1% Poly (2-hydroxyethyl methacrylate; Sigma Aldrich) at a density of 200 or 50 cells per well. Cells were grown in MEBM basal medium supplemented with B27 without vitamin A (50X), 20 ng/mL EGF, 20 ng/mL bFGF and 4 μg/mL heparin. Cells were incubated at 37 °C with 5% CO_2 during 5 days.

Mammosphere acquisition and analysis

At day 5, mammospheres were labeled using 0.2 nM Calcein-AM (PromoCell GmbH) and incubated 30 minutes at room temperature. Data acquisition was done using Incucyte® (Essen Bioscience, Sartorius) for brigthfield and green fluorescence. Image analysis was processed with IncuCyteS3 ®Software using the following parameters: Segmentation [Top Hat, Radius (100μm), Threshold (2 GCU)], Edge split [on, edge sensitivity -55], Hole fill [100μm^2].

Immunoprecipitation and co-immunoprecipitation

Protein extracts were prepared in lysis buffer (NaCl 150 mM, Tris pH 7.4 10 mM, EDTA 1 mM, Triton X-100 1%, NP-40 0.5%, cOmplete, EDTA-free protease inhibitors (Roche #11873580001) for 30 min on ice before centrifugation. Immunoprecipitations were performed overnight at 4°C on a rocking wheel using either mouse anti-MAGI1 antibody for endogenous immunoprecipitations or anti-HA antibody and EZview Red anti-FLAG M2 affinity gel (Sigma-Aldrich #F1804) for co-immunoprecipitations. Protein G sepharose was then added to the MAGI1- or HA- immunoprecipitates for 1 hour at 4°C before extensive washes. Concerning the FLAG immunoprecipitation, washes were performed followed by protein elution by competition with 3XFLAG peptide (150 ng/μL final concentration) during 1 hour at

4°C. The different immunoprecipitates were then submitted to Western blotting for detection of

protein complexes.

Subcellular fractionation - Isolation of cytoplasmic, nuclear and membrane protein fractions.

Subcellular fractionation of cultured human cell lines was performed as previously described; samples,

buffers and centrifugations were kept and performed at 4°C at all time. For more details, the protocol

is available on-line (http://www.bio-protocol.org/e754).

Cell growth assay

2D cell growth assay was analyzed using MTT (Thiazolyl Blue Tetrazolium Bromide (Sigma Aldrich

#M2128)). Briefly, cells seeded in 96 well plates were stopped from day 0 to day 7 by incubation with

MTT solution (0.5 mg/mL final concentration) during 4 hours at 37 °C. Before reading OD at 570 nm,

cells were incubated with DMSO to solubilize the formazan crystals formed in the presence of MTT.

3D Spheroid formation and measures

10 000 cells were seeded in Ultra-Low Attachment 96 well plates (Costar) cultured in their regular

medium supplemented with 15 % FCS. After 3 days, all cells formed 3D spheroids that can be further

analyzed by perimeter and circularity measurements. Pictures of spheroids were taken 3 days after

seeding cells to assay the ability of cell lines to form spheroids and both the perimeters and the

circularity of spheroids were calculated. Circularity was calculated (4px(area/(perimeter)^2) where 1 is

considered as a perfect circle.

p38 MAPK inhibitor treatment

The different MCF7*shRNA* cell lines were plated at $3x10^6$ cells per 60mm plates. The day after, 10 µM

of Ralimetinib (LY2228020; Selleckchem #S1494) was added and 24 h later cells were collected and

counted to be plated for the different experiments (Ralimetinib treatment is maintained throughout

the experiments). For proliferation assay, 3000 cells were plated in 96-well plates and plates were

stopped at Day 0, Day 2, Day 5 and Day 8 using 10 % of Trichloroacetic acid (TCA; Sigma Aldrich #T4885)

for 10 min, then washed 3 times with PBS and stored at 4°C until the end of the experiment. Once the

experiment is finished, 50 µl of Sulforhodamine B (SRB; Sigma Aldrich #230162) 0.04% diluted in acetic acid 1% were added for 30 min before extensive washes with acetic acid 1% and drying step overnight at room temperature. Before reading OD at 540 nm, cells were incubated with Tris 10 mM pH 10.5. For Soft agar assay, 10 000 cells were plated in 12-well plates before proceeding to the assay detailed below. The same experiments were conducted with another p38 inhibitor using 10 µM of Pexmetinib (ARRY-614; Selleckchem #S7799).

ROCK inhibitors treatment

The treatments of MCF7*shRNA* cell lines with either Blebbistatin (Selleckchem #7099) or Y-27632 (Selleckchem #S1049) were both performed at a final concentration of 10 µM and cells were treated during 2 hours before being lysed for Western blot analyses.

Soft Agar assay

To assess anchorage independent growth, soft agar assays were performed. A first layer of 0.8% Noble agar (Sigma Aldrich #A5431) was prepared in 12-well plates and left at 4C during 1 hour for the agar to solidify gently. The plate was kept at 37C before the cell-containing agar layer was prepared. 5,000 or 10,000 cells were imbedded in the second layer containing 0.4% Noble agar on top of the first layer. When the second layer was solidified, fresh medium was added on top of the cells. MCF7 cell lines were cultured during 14 to 21 days before Crystal violet coloration (0.01% final concentration; Sigma Aldrich #C0775) or MTT assay (1 mg/mL final concentration).

Primary tumor growth assay

MCF7shRNA and HCT116shRNA cell lines were injected subcutaneously in six-week-old female athymic Nude-Foxn1 mice (Envigo). Both flanks of each mouse were injected with cells mixed with Matrigel (10^6 cells injected for HCT116; 10×10^6 cells injected for MCF7). Mice injected with MCF7 cells were subjected to neck dropping with 50µl of β-estradiol (1.5 mg/mL diluted in ethanol) to stimulate MCF7 cell growth. Animal procedures were performed according to protocols approved by the French national committee on animal care.

RESULTS

The loss of MAGI1 enhances tumorigenicity of luminal breast cancer (BCa) cells

In order to study the role of the apical scaffold MAGI1 during breast carcinogenesis, we first

analyzed the expression of MAGI1 by western blot in both luminal and basal BCa cell lines: MAGI1

expression was restricted to luminal ER+ lineages (T47D and MCF7 luminal A, and to a lesser extent in

ZR75 luminal B), while no expression could be detected in basal ER- lineages (immortalized MCF10A

and triple negative MDA-MB-468 and BT549; Supplemental Fig. 1A). Moreover, public database mining

(http://www.kmplot) [50], indicated that low *MAGI1* RNA expression levels were associated with worse

prognosis in relapse-free survival for BCa patients, but only in ER+ (mainly luminal) molecular BCa

subtypes (Kaplan Meier curve; Supplemental Fig. 1B).

We thus decided to investigate the functional role of MAGI1 and the consequence of MAGI1

knockdown in luminal BCa, and generated MCF7 and T47D cell lines in which *MAGI1* was targeted by

constitutive shRNA. Two independent shRNA constructs targeting different parts of the *MAGI1*

transcript were used, *shMAGI1(3-1)* and *shMAGI1(1-1)*, which led respectively to more than 90% and

around 50% MAGI1 knockdown at the protein level as shown by western blot and immunofluorescence

analyses (Supplemental Fig. 1D&E). The knockdown was specific for MAGI1 and did not induce

compensatory up-regulation of MAGI2 and MAGI3, the two remaining MAGI family members

(Supplemental Fig. 1D&F). *shMAGI1(3-1)* was the most potent resulting in almost knock-out like down-

regulation, and was therefore used in all subsequent studies and referred to as *shMAGI1* thereafter.

First, *MAGI1* knockdown increased 2D cell growth of MCF7 cells (25% increase at day 7) as

assayed by MTT cell growth assay (Fig. 1A). It was associated with a weak increase in the proportion of

cells in the S phase of the cell cycle from 21% in control MCF7*shLuc* to 29% in MCF7*shMAGI1* cells (Fig.

1B). More importantly, in anchorage-independent soft agar cell growth conditions, MCF7*shMAGI1* cell

lines formed circa twice as many colonies compared to *shLuc* controls (Fig. 1C). Similarly,

MCF7*shMAGI1* cells showed enhanced clonal mammosphere capacities which were correlated with an

expanded area occupied by the spheres (60% increase; Fig. 1D). Significantly, when MCF7*shMAGI1*

cells were injected subcutaneously in nude mice, they grew as a tumor mass much more rapidly and

extensively than control MCF7*shLuc* cells (Fig. 1E). Together, these results show that the loss of MAGI1

in MCF7 cells promotes tumorigenicity (Fig. 1A-E). Similar results were obtained in T47D, a second

luminal A BCa cell line, in which MAGI1 knockdown (*shMAGI1*) led to elevated 2D and anchorage-

independent cell growth (Supplemental Fig. 2A&B). Even though BCa was the prime focus of this study,

the effects of *MAGI1* knockdown appeared not restricted to mammary cells, and similar observations

on anchorage-free growth in soft agar, and orthotopic tumor growth in nude mice were also observed

using HCT116 colon cancer cells (Supplemental Fig. 2E&F) extending earlier reports [34]. Altogether, our

results show that the loss of MAGI1 in luminal A BCa cells enhances their tumorigenic behaviors.

The loss of MAGI1 promotes the accumulation of epithelial junctions components

In order to further investigate the role of MAGI1 in breast tissue, we sought to determine its

localization in mammary epithelial cells. Immunohistochemistry on BCa patient tissue micro-array

revealed that in normal breast cells, MAGI1 was expressed only in the luminal epithelial cells and not

in the underlying basal myo-epithelial layer, consistent with MAGI1 expression exclusively in luminal-

type BCa cell lines (Supplemental Fig. 1A). At the sub-cellular level, MAGI1 was localized at the apical

pole of luminal breast cells (Supplemental Fig. 1C, arrows). Using MCF7 cancer cells, immortalized

hMEC (human mammary epithelial cells), and polarized canine MDCK cells, we further confirmed by

immunofluorescence that MAGI1 localized near the plasma membrane, overlapping with junction

components such as E-cadherin (AJ), ZO1, and Claudin3 (TJ), suggesting that MAGI1 is a TJ and/or AJ

resident protein (Fig. 2A, arrows and data not shown).

The localization of MAGI1 at apical junctions prompted us to explore whether MAGI1 could

control their biology. Performing western-blot analyses on whole protein extracts, the loss of MAGI1

did not affect the total protein abundance of the typical TJs component ZO-1 nor of the AJs component

PARD3 (Fig. 2B). However, in MCF7*shMAGI1* we observed increases in the junctional proteins β-catenin

(x1.25), E-cadherin (x1.6) and AMOTL2 (x1.8) (Fig. 2B). Consistently, performing immunofluorescence

on fixed MCF7*shMAGI1* cells, the membrane levels and localization of ZO-1 and PARD3 were

unchanged, while E-cadherin staining was increased, and less tightly restricted to the plasma

membrane (Fig. 2C). The altered distribution of E-cadherin in *shMAGI1* cells is reminiscent of the role

of Magi in *Drosophil*a where we showed that it controls E-cadherin belt at AJs in epithelial cells of the

developing eye [24].

In epithelial cells, the increase in E-cadherin material is often associated with increased

strength of the junctions, and through the linkage to the underlying actin to increases in tension and

overall compressive forces (reviewed by [7]). While we did not observe obvious changes in the intensity

or morphology of F-actin upon MAGI1 knockdown (Supplemental Fig. 3A&B), MCF7*shMAGI1* cells

exhibited behaviors compatible with increased compressive forces. First, when cultured as spheroids,

MCF7*shMAGI1* cells grew as round masses of cells, rounder than MCF7*shLuc* controls with circa 30%

smaller perimeter (Fig. 3A&B) and better circularity (0.95 vs 0.62 respectively; Fig. 3C), a sign that

MCF7*shMAGI1* cells aggregates were more compact, even though composed of the same number of

cells as MCF7*shLuc* controls. This compaction was also observed in T47D *shMAGI1* cells (Supplemental

Fig. 2C&D). Second, using Atomic Force Microscopy (AFM), we showed that MCF7*shMAGI1* cells were

stiffer than controls (higher Young's modulus factor; Fig. 3D), strongly suggesting that MCF7*shMAGI1*

cells may have a stronger cytoskeleton and might experience higher internal pressure. Since AFM was

performed on small MCF7 cells clusters, our data indicate that the increased compressive forces were

generated from within the cells, likely through increased actin contractility. Indeed, elevated ROCK

activity was detected in MAGI1 deficient cells as shown by the increase in ROCK-specific serine 19

phosphorylation of the regulatory myosin light chain MLC2 (Fig. 3E&F), suggesting that the actin

cytoskeleton was under higher tension. Furthermore, treating MCF7*shMAGI1* cells with ROCK

inhibitors completely abolished the increased cell stiffness and the Young's modulus factors measured

by AFM after inhibitors treatment reached similarly low levels in *shMAGI1* and *shLuc* controls

(Supplemental Fig. 3C).

MAGI1 loss does not promote YAP nor β-catenin signaling

In epithelial cells, E-cadherin-based AJs regulate the activity of mechanosensitive pathways,

and in particular the Hippo and Wnt/β-catenin pathways, two major pathways controlling cell

proliferation and frequently mis-regulated during carcinogenesis [8,51]. We thus explored whether these

two pathways could be affected following MAGI1 depletion.

Monitoring Hippo pathway and YAP activity in stable MCF7*shMAGI1* showed that in the

absence of *MAGI1*, total YAP and phosphorylated YAP (on both serine residues S127 and S397)

accumulated in the cells while TAZ levels remained unchanged (Western blot analyses; Fig. 4A). By

immunofluorescence, we then showed that unlike what was observed for control cells grown at similar

confluence, YAP did not accumulate in the nuclei of MCF7*shMAGI1* cells (Fig. 4B). These findings were

further confirmed in luminal BCa patients TMA, where we observed a strong correlation in ER+ tumors

between MAGI1 apical membrane localization and YAP nuclear localization: only 27 % of the patients

with no MAGI1 expression or mis-localized MAGI1 (no membrane staining) showed YAP nuclear

localization while 92% of patients with membranous MAGI1 showed nuclear YAP (Figure 4C).

Accordingly, *MAGI1* knockdown in MCF7 cells resulted in impaired YAP transcriptional activity with

decreased expression of canonical YAP target genes such as *CTGF, CYR61, BIRC2,* and *AREG* (Figure 4D).

Surprisingly, even though YAP phosphorylations on S127 and S397 are mediated by LATS kinases

(Reviewed in [52,53]), the loss of MAGI1 did not promote a higher activity of the Hippo pathway and the

levels of phosphorylated activated LATS1 kinase and of phosphorylated MOB1 were unaffected (Figure 4A). This suggests that the elevated phosphorylated YAP were not the consequence of increased upstream Hippo pathway activity (MST kinases), but might reflect higher p-YAP stability, either because it is protected from destruction, or because it is not dephosphorylated as efficiently, or a combination of the two. These results are in agreement with earlier studies showing that E-cadherin junctional tension in epithelial cells promote YAP/TAZ nuclear exclusion either through α-catenin conformational change [12–14], or through Merlin/NF2-mediated cytoplasmic retention [10]. Together, these results show that the junctional MAGI1 scaffold prevents the accumulation of phosphorylated YAP in the cytoplasm of luminal BCa cells.

Similarly, even though higher overall protein levels of β-catenin could be observed in MCF7*shMAGI1* cells compared to controls, this was not reflected by an increase in the active unphosphorylated β-catenin pool (Figure 4A). These results are in agreement with studies showing that increased compression transduced by apical junctions in epithelial cells prevent β-catenin nuclear accumulation [11,16]. These results thus show that the loss of MAGI1 does not promote YAP nor β-catenin signaling, and suggest that these pathways do not mediate the increased tumorigenicity of *MAGI1* deficient cells.

p38 Stress Activated Protein Kinase mediates tumorigenicity of MAGI1 deficient cells

We thus asked which mechanisms mediate the increase in 2D cell proliferation and in anchorage-independent cell growth observed upon MAGI1 loss (see Fig. 1), focusing on other known oncogenic pathways. The ERK/MAP Kinase and JNK stress kinase pathways could be influenced by junctions and the forces they sense [15,17,18]. In response to MAGI1 loss, we did not detect any alterations of the main oncogenic pathways: Akt, ERK1/2, and JNK pathways (Fig. 5A&B). Interestingly, the p38 stress kinase was strongly activated in MCF7*shMAGI1* compared to MCF7*shLuc*, and the amount of phosphorylated p38, normalized to total p38 protein level, was increased 2 fold (Fig. 5A&4B). The p38 pathway is activated in response to a wide variety of cellular stresses and has been implicated either as a tumor-suppressor or as an oncogene in various cancers, including BCa [54–58]. Using p38 specific inhibitors, we then tested whether the p38 pathway could mediate, at least in part, the increased tumorigenicity of MCF7 luminal cells. In 2D cell growth, MCF7*shMAGI1* treated with LY2228820 p38α/β inhibitor (also named Ralimetinib) grew less than untreated cells (2 fold suppression at 8 days of growth; Figure 5C). It should be noted that p38 inhibition also slowed the growth of MCF7*shLuc* control cells, but the effect was far less pronounced than for MCF7*shMAGI1* cells. In soft agar,

LY2228820 treatment suppressed strongly the anchorage-independent cell growth of MCF7*shMAGI1* (2.5 fold decrease) as compared to its effect on MCF7*shLuc* control cells (Figure 5D). Similar effects were obtained using a second independent p38 inhibitor, ARRY-614 (p38 and Tie2 inhibitor), even though we observed higher toxicity (data not shown), strongly supporting that the effects observed were the consequence of p38 inhibition and not any unspecific action of the drugs. Taken together, our results demonstrate the essential role of the p38 signaling pathway for MAGI1-deficient MCF7 luminal BCa cells tumorigenicity, consistent with previous report identifying the critical role of p38α in mammary tumorigenesis using mouse BCa models [54].

Strikingly, treating MCF7*shMAGI1* cells with the Rho Kinase (ROCK) inhibitors Y-27632 or Blebbistatin that prevent Myosin-mediated F-actin tension, abolished p38 activation (Fig. 5E&F) and largely decreased elastic cell Young's modulus (Supplemental Fig. 3C), showing that increased ROCK activity and actin strength represent critical events to activate p38 signaling. However, due to their cell toxicity, these ROCK inhibitors treatments could only be maintained for short periods, preventing us to assay formally, whether releasing actin tension in MAGI1 deficient cells could suppress their tumorigenicity as would be expected from their effects on p38.

MAGI1 interacts with the angiomotin family members AMOT and AMOTL2 at the junctions

In order to better understand how the loss of MAGI1 and the associated alterations in ROCK activity and in E-cadherin-based junctions could lead to p38 stress-kinase activation, we decided to focus our studies on the junctional MAGI1 molecular complexes. Several studies, including proteomic approaches, have described a physical association between MAGI1 and AMOT or AMOTL2, two members of the angiomotin family of apical scaffolds, acting both as junctions/actin linkers as well as negative regulators of YAP transcriptional activity [41–43,59,60]. We thus, investigated whether AMOTs could mediate the effects of MAGI1 to restrict tumorigenesis in breast luminal cells.

First, using co-immunoprecipitation experiments, we confirmed that overexpressed Flag-tagged MAGI1 or endogenous MAGI1 interacted with endogenous AMOT and AMOTL2 in MCF7 luminal BCa cells (Fig. 6A&B). Building on the availability of numerous constructs generated by the group of KL. Guan [61], we have then used AMOT as a paradigm to study the interaction between MAGI1 and AMOTs through co-immunoprecipitation experiments between Flag-tagged MAGI1 and HA-tagged AMOT constructs. Using point mutations affecting key structural Proline residues in the first and second WW motifs of MAGI1, we showed that the second WW motif was required for the interaction with AMOT, since co-immunoprecipitation was abolished by a point mutation on Proline residue 390

(mutated to Alanine MAGI1 P390A; Fig. 6C&D). The first WW domain appeared dispensable as no change in binding was observed when mutating Proline 331 (MAGI1 P331A). Similar results were obtained performing co-immunoprecipitations on endogenous AMOTL2 (Fig. 6E), showing that as for AMOT, the second WW domain of MAGI1 was required for AMOTL2 binding, confirming earlier findings [43]. The N-terminus part of AMOT contains two PPXY motifs, which represent canonical WW domains interactors. Using HA-tagged AMOT point mutants, we then determined that the interaction between MAGI1 and AMOT occurred through the second PPXY motif (Fig. 6F&G) since co-immunoprecipitation was lost in AMOT Y287A (Tyrosine 287 mutated to Alanine; the mutation AMOT Y242A did not show any effect). Together these results confirm using precise point mutations that MAGI1 binds to AMOT and AMOTL2 through its second WW domain, and at the level of the conserved second PPXY motifs of the N-terminal domains of AMOT family scaffolds.

Then, using immunofluorescence on MCF7 fixed cells, MAGI1, AMOT, and AMOTL2 localized in overlapping membrane domains corresponding to cellular junctions (Fig. 6H; the AMOT nuclear staining is non-specific). Even though, MAGI1 colocalized with AMOTL2 and AMOT (Fig. 6H, arrows), it was not required for their proper membrane localizations as they were unaffected in *shMAGI1,* suggesting that even though they physically interact once in proximity, they must reach their membrane localization independently (Fig. 6I). Nevertheless, even though MAGI1 was not required for AMOTL2 localization, it did however control AMOTL2 levels, and AMOTL2 accumulated 1.8 fold in MCF7*shMAGI1* compared to controls as measured by western blots on whole protein extracts (Supplemental Fig. 3A&B). These results suggest that MAGI1 regulates the stability and/or degradation of AMOTL2, but using classic drugging approaches, we were unable to identify the mechanisms involved.

Together our results confirm the physical interaction between MAGI1 and AMOTs and show that MAGI1 restricts the total AMOTL2 protein levels, through a mechanism that remains to be established.

AMOTL2 mediates the increased tumorigenicity of MAGI1-deficient luminal BCa cells

We observed that in luminal BCa cells, the loss of MAGI1 triggered both i) an accumulation of AJs material including its binding partner AMOTL2, and ii) an activation of the p38 pathway driving an increase in tumorigenicity. We thus studied next whether these two different aspects were linked or independent consequences of MAGI1 loss. Generating double knockdown MCF7 cells for *MAGI1* and *AMOTL2*, we could show that the increased 2D growth and anchorage-independent growth after MAGI1 loss were suppressed by the knockdown of *AMOTL2* (Fig. 7A&B). Consistently, *AMOTL2*

knockdown abolished the accumulation of E-cadherin and the activation of the p38 stress kinase (phospho-p38 levels; Fig. 7C&D). These results identify AMOTL2 accumulation as a critical mediator in the p38 activation and tumorigenesis induced by impaired *MAGI1*.

Together, these results show that both AMOTL2 and ROCK activity are required for p38 activation in MAGI1 deficient luminal BCa cells. They support a model in which, the AMOTL2 accumulation in MAGI1 deficient cells governs an increase of E-cadherin-based junctions, which together with ROCK activity and cortical actin tension, activates the tumorigenic activity of the p38 signaling pathway (Fig.7E).

DISCUSSION

In this study, we explored the function of the junctional component MAGI1 in breast carcinogenesis and showed that the loss of MAGI1 in luminal BCa cell lines promoted tumorigenesis. At the cellular level, we showed that the loss of MAGI1 induced an accumulation of AJ components and increased cell compression behaviors. Importantly, we showed that the increased tumorigenicity of MAGI1 deficient luminal BCa cells was mediated by p38 stress signaling. Molecularly, we confirmed that MAGI1 interacted with the AMOT family of apical scaffolds and actin linkers, and showed that decreasing *AMOTL2* levels, completely suppressed the effect of *MAGI1* loss on the activation of p38 and on tumorigenicity. Finally, we provided evidence that inhibiting ROCK activity could alleviate the p38 activation, supporting a model in which MAGI1 suppresses luminal BCa by preventing both ROCK and AMOTL2-mediated Junction dysfunction, and subsequent p38 stress signaling.

The loss of MAGI1 led to an increase in tumorigenicity, and an accumulation of AJ material, in particular E-cadherin, reminiscent to the role we described previously for the unique MAGI homologue in *Drosophila* [24]. This correlation could appear counterintuitive as E-cadherin knockdown promotes invasion and migration in cultured cells [62], and is critical during EMT and for aggressiveness (recently reviewed in [63]). However, in several cancers including some BCa, E-cadherin expression is maintained with a proposed tumor supporting role [64,65] and recent mouse models of luminal and basal invasive ductal carcinomas have demonstrated that mammary cancer cells proliferation, resistance to apoptosis, and distant metastatic seeding potential, require E-cadherin [66]. These observations reconcile the high E-cadherin levels and increased tumorigenicity observed upon *MAGI1* loss, and suggest that at least in specific phases of tumorigenesis (e.g. mass proliferation, colony formation), E-cadherin could play a positive role during BCa.

One striking feature of MAGI1 depleted cells is the protein accumulation of AMOTL2. Even though the exact mechanisms remain to be described, the stability of the related AMOT is controlled by the activity of the RNF146/Tankyrase pathway [67,68], and by phosphorylation by the Hippo pathway core kinase LATS1 (reviewed by [37]). Importantly AMOTL2, a binding partner of MAGI1, is required for the increased tumorigenicity of MAGI1-depleted luminal BCa cells, showing that AMOTL2 is a critical mediator in this context. AMOTs are apical scaffolds playing important roles in cadherin-based junction regulations and its linkage to the actin cytoskeleton [26,35,36]. Previous reports have shown that AMOTL1 is overexpressed in invasive ductal carcinomas, leading to invasive behaviors. However, this effect of AMOTL1 appears specific to ER-negative subtypes [69]. AMOTL1 is not expressed in breast luminal lineages, in which AMOTL2 is the most abundant (twice more abundant than AMOT in MCF7 cells; our

unpublished observation). We propose thus that in ER-positive luminal breast lineages, AMOTL2 overexpression could play similar roles as marker and mediator of tumorigenesis. Interestingly, a short isoform of AMOTL (AMOTL2 p60 in contrast to the long AMOTL2 p100) was reported overexpressed in *in situ* and invasive ductal carcinoma cells [70]. The AMOTL2 p60 isoform lacks the PPxY domain critical for its interaction with MAGI1, and it is possible this isoform is thus insensitive to MAGI1-dependent destabilization, even though this remains to be studied. Hypoxia and Fos signaling govern the specific expression of AMOTL2 p60 [70], and the accumulation of AMOTL2 in the context of *MAGI1* mutation could thus represent an alternative mechanism for AMOTL2 expression, extending the relevance of AMOTL2 up-regulation during carcinogenesis.

AMOTL2 couples adhesion complexes to the actin cytoskeleton to allow F-actin tension and thus morphogenesis in endothelial cells (coupling with VE-cadherin [71]) and in epithelial cell during embryogenesis (E-cadherin [59]). Importantly, the specific cellular morphology defects in AMOTL2 knockdown could be mimicked by inhibiting ROCK kinase and Myosin phosphorylation showing that AMOTL2 and actin tension are linked [35]. Relaxing actin tension using two ROCK inhibitors (Y-27632 and Blebbistatin), or preventing AMOTL2 accumulation (siRNA), both robustly suppressed p38 activation, a key feature mediating tumorigenesis after MAGI1 loss. So how could AMOTL2 and ROCK interact to mediate the effects of *MAGI1* loss? They could either (i) be two independent consequences of MAGI1 loss or (ii) they could be linked causally. Due to the toxicity of long term exposures to ROCK inhibitors, we were unable to study whether AMOTL2 stability could be a consequence of ROCK activity. Preliminary results would indicate that AMOTL2 does not control ROCK activity (data not shown), suggesting that ROCK might act upstream, or in parallel to AMOTL2, but more experiments are needed to better determine how MAGI1 controls ROCK activity, and how ROCK and AMOTL2 regulate p38 activity.

p38 signaling is activated by many upstream signals including a wide variety of environmental stresses such as UV, genotoxic agents, heat shock, or hyperosmotic conditions [18]. During osmotic shock, the cellular cortex is subjected to external pressure reminiscent to compressive forces, and we propose therefore that E-cadherin and AMOTL2 enrichment, together with increased ROCK activity and actin tension result in similar compressive forces that could represent a new stress signal activating p38. Amongst the four different p38 kinases (p38$\alpha,\beta,\gamma,\delta$), P38$\alpha$ is the most ubiquitously expressed. p38 signaling has been implicated either as a tumor suppressor or as an oncogene depending on cancer type and stage [18,56–58]. In BCa, targeting p38δ resulted in a reduced rate of cell proliferation and enhanced cell-matrix adhesion [72]. More importantly, BCa mouse models, demonstrated the critical role of p38α (Mapk14) during the initiation, and proliferation of mammary tumors. The tumor promoting role of p38α involves a protective role against deleterious DNA damage in the mammary epithelial

cancer cells [54]. We observed in *MAGI1*-deficient cells an increased proportion of cells in S-phase, indicating that these cells might have a slower S-phase. Slow S-phase is classically observed when cells have to overcome a high level of DNA damage. These results highlight the need for further studies to better understand the link between MAGI1 loss, p38 activation, DNA damage, and breast tumorigenesis.

Besides elevated p38 signaling, MAGI1-deficient cells exhibited low YAP nuclear activity, consistent with earlier reports showing that in epithelial cells with high E-cadherin and/or under compressive forces α-Catenin and NF2/Merlin act to exclude YAP from the nucleus [10]. Here we demonstrated the critical role of AMOTL2 downstream of MAGI1 to mediate E-cadherin accumulation. Since AMOTs have been shown to trap YAP in the cytoplasm [61], further studies are thus required to better understand the interactions between MAGI1/AMOTL2 and α-Cat/NF2 in the control of YAP localization and activity. This low YAP signaling could appear surprising since the oncogenic role of nuclear YAP is well established [73]. However, the oncogenic role of YAP in luminal BCa remains debated. While elevated YAP/TAZ activity gene signatures have been reported to correlate with more aggressive BCa [74–76], aggressive BCa are enriched in basal/triple negative sub-types. Furthermore, several studies report conflicting or no correlation between YAP staining levels and clinical outcomes in BCa patients (reviewed in [73]) suggesting that YAP/TAZ levels and nucleo/cytoplasmic distributions in luminal BCa could be re-examined.

REFERENCES

1. Holliday, D. L. & Speirs, V. Choosing the right cell line for breast cancer research. *Breast Cancer Res. BCR* **13**, 215 (2011).

2. Guedj, M. *et al.* A refined molecular taxonomy of breast cancer. *Oncogene* **31**, 1196–1206 (2012).

3. Knust, E. & Bossinger, O. Composition and formation of intercellular junctions in epithelial cells. *Science* **298**, 1955–1959 (2002).

4. Coopman, P. & Djiane, A. Adherens Junction and E-Cadherin complex regulation by epithelial polarity. *Cell. Mol. Life Sci. CMLS* **73**, 3535–3553 (2016).

5. Jeanes, A., Gottardi, C. J. & Yap, A. S. Cadherins and cancer: how does cadherin dysfunction promote tumor progression? *Oncogene* **27**, 6920–6929 (2008).

6. Balda, M. S. & Matter, K. Tight junctions as regulators of tissue remodelling. *Curr. Opin. Cell Biol.* **42**, 94–101 (2016).

7. Charras, G. & Yap, A. S. Tensile Forces and Mechanotransduction at Cell-Cell Junctions. *Curr. Biol. CB* **28**, R445–R457 (2018).

8. Pinheiro, D. & Bellaïche, Y. Mechanical Force-Driven Adherens Junction Remodeling and Epithelial Dynamics. *Dev. Cell* **47**, 3–19 (2018).

9. Benham-Pyle, B. W., Pruitt, B. L. & Nelson, W. J. Cell adhesion. Mechanical strain induces E-cadherin-dependent Yap1 and β-catenin activation to drive cell cycle entry. *Science* **348**, 1024–1027 (2015).

10. Furukawa, K. T., Yamashita, K., Sakurai, N. & Ohno, S. The Epithelial Circumferential Actin Belt Regulates YAP/TAZ through Nucleocytoplasmic Shuttling of Merlin. *Cell Rep.* **20**, 1435–1447 (2017).

11. Hirata, H., Samsonov, M. & Sokabe, M. Actomyosin contractility provokes contact inhibition in E-cadherin-ligated keratinocytes. *Sci. Rep.* **7**, 46326 (2017).

12. Kim, N.-G., Koh, E., Chen, X. & Gumbiner, B. M. E-cadherin mediates contact inhibition of proliferation through Hippo signaling-pathway components. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 11930–11935 (2011).

13. Schlegelmilch, K. *et al.* Yap1 acts downstream of α-catenin to control epidermal proliferation. *Cell* **144**, 782–795 (2011).

14. Silvis, M. R. *et al.* α-catenin is a tumor suppressor that controls cell accumulation by regulating the localization and activity of the transcriptional coactivator Yap1. *Sci. Signal.* **4**, ra33 (2011).

15. Aoki, K. *et al.* Propagating Wave of ERK Activation Orients Collective Cell Migration. *Dev. Cell* **43**, 305-317.e5 (2017).

16. Hall, E. T., Hoesing, E., Sinkovics, E. & Verheyen, E. M. Actomyosin contractility modulates Wnt signaling through adherens junction stability. *Mol. Biol. Cell* **30**, 411–426 (2019).

17. Pereira, A. M., Tudor, C., Kanger, J. S., Subramaniam, V. & Martin-Blanco, E. Integrin-dependent activation of the JNK signaling pathway by mechanical stress. *PloS One* **6**, e26182 (2011).

18. Wagner, E. F. & Nebreda, A. R. Signal integration by JNK and p38 MAPK pathways in cancer development. *Nat. Rev. Cancer* **9**, 537–549 (2009).

19. Beavon, I. R. The E-cadherin-catenin complex in tumour metastasis: structure, function and regulation. *Eur. J. Cancer Oxf. Engl. 1990* **36**, 1607–1620 (2000).

20. Bosch-Fortea, M. & Martín-Belmonte, F. Mechanosensitive adhesion complexes in epithelial architecture and cancer onset. *Curr. Opin. Cell Biol.* **50**, 42–49 (2018).

21. Laura, R. P., Ross, S., Koeppen, H. & Lasky, L. A. MAGI-1: a widely expressed, alternatively spliced tight junction protein. *Exp. Cell Res.* **275**, 155–170 (2002).

22. Hirabayashi, S. *et al.* JAM4, a junctional cell adhesion molecule interacting with a tight junction protein, MAGI-1. *Mol. Cell. Biol.* **23**, 4267–4282 (2003).

23. Padash Barmchi, M., Samarasekera, G., Gilbert, M., Auld, V. J. & Zhang, B. Magi Is Associated with the Par Complex and Functions Antagonistically with Bazooka to Regulate the Apical Polarity Complex. *PloS One* **11**, e0153259 (2016).

24. Zaessinger, S., Zhou, Y., Bray, S. J., Tapon, N. & Djiane, A. Drosophila MAGI interacts with RASSF8 to regulate E-Cadherin-based adherens junctions in the developing eye. *Dev. Camb. Engl.* **142**, 1102–1112 (2015).

25. Lynch, A. M. *et al.* A genome-wide functional screen shows MAGI-1 is an L1CAM-dependent stabilizer of apical junctions in C. elegans. *Curr. Biol. CB* **22**, 1891–1899 (2012).

26. Hultin, S. *et al.* AmotL2 links VE-cadherin to contractile actin fibres necessary for aortic lumen expansion. *Nat. Commun.* **5**, 3743 (2014).

27. Feng, X., Jia, S., Martin, T. A. & Jiang, W. G. Regulation and involvement in cancer and pathological conditions of MAGI1, a tight junction protein. *Anticancer Res.* **34**, 3251–3256 (2014).

28. Zhang, G., Liu, T. & Wang, Z. Downregulation of MAGI1 associates with poor prognosis of hepatocellular carcinoma. *J. Investig. Surg. Off. J. Acad. Surg. Res.* **25**, 93–99 (2012).

29. Kozakai, T. *et al.* MAGI-1 expression is decreased in several types of human T-cell leukemia cell lines, including adult T-cell leukemia. *Int. J. Hematol.* **107**, 337–344 (2018).

30. Zhang, G. & Wang, Z. MAGI1 inhibits cancer cell migration and invasion of hepatocellular carcinoma via regulating PTEN. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* **36**, 381–385 (2011).

31. Zmajkovicova, K. *et al.* MEK1 is required for PTEN membrane recruitment, AKT regulation, and the maintenance of peripheral tolerance. *Mol. Cell* **50**, 43–55 (2013).

32. Kranjec, C., Massimi, P. & Banks, L. Restoration of MAGI-1 expression in human papillomavirus-positive tumor cells induces cell growth arrest and apoptosis. *J. Virol.* **88**, 7155–7169 (2014).

33. Makokha, G. N. *et al.* Human T-cell leukemia virus type 1 Tax protein interacts with and mislocalizes the PDZ domain protein MAGI-1. *Cancer Sci.* **104**, 313–320 (2013).

34. Zaric, J. *et al.* Identification of MAGI1 as a tumor-suppressor protein induced by cyclooxygenase-2 inhibitors in colorectal cancer cells. *Oncogene* **31**, 48–59 (2012).

35. Hildebrand, S. *et al.* The E-cadherin/AmotL2 complex organizes actin filaments required for epithelial hexagonal packing and blastocyst hatching. *Sci. Rep.* **7**, 9540 (2017).

36. Hultin, S. *et al.* AmotL2 integrates polarity and junctional cues to modulate cell shape. *Sci. Rep.* **7**, 7548 (2017).

37. Maugeri-Saccà, M. & De Maria, R. The Hippo pathway in normal development and cancer. *Pharmacol. Ther.* **186**, 60–72 (2018).

38. Zhao, B. *et al.* Angiomotin is a novel Hippo pathway component that inhibits YAP oncoprotein. *Genes Dev.* **25**, 51–63 (2011).

39. Yu, F.-X. & Guan, K.-L. The Hippo pathway: regulators and regulations. *Genes Dev.* **27**, 355–371 (2013).

40. Chan, S. W. *et al.* Hippo pathway-independent restriction of TAZ and YAP by angiomotin. *J. Biol. Chem.* **286**, 7018–7026 (2011).

41. Wang, W. *et al.* Defining the protein-protein interaction network of the human hippo pathway. *Mol. Cell. Proteomics MCP* **13**, 119–131 (2014).

42. Bratt, A. *et al.* Angiomotin regulates endothelial cell-cell junctions and cell motility. *J. Biol. Chem.* **280**, 34859–34869 (2005).

43. Patrie, K. M. Identification and characterization of a novel tight junction-associated family of proteins that interacts with a WW domain of MAGI-1. *Biochim. Biophys. Acta* **1745**, 131–144 (2005).

44. Johannessen, C. M. *et al.* COT drives resistance to RAF inhibition through MAP kinase pathway reactivation. *Nature* **468**, 968–972 (2010).

45. Guo, F., Chiang, M.-Y., Wang, Y. & Zhang, Y.-Z. An in vitro recombination method to convert restriction- and ligation-independent expression vectors. *Biotechnol. J.* **3**, 370–377 (2008).

46. Marzi, L. *et al.* FOXO3a and the MAPK p38 are activated by cetuximab to induce cell death and inhibit cell proliferation and their expression predicts cetuximab efficacy in colorectal cancer. *Br. J. Cancer* **115**, 1223–1233 (2016).

47. Saavedra V, O., Fernandes, T. F. D., Milhiet, P.-E. & Costa, L. Compression, Rupture, and Puncture of Model Membranes at the Molecular Scale. *Langmuir ACS J. Surf. Colloids* **36**, 5709–5716 (2020).

48. Fernandes, T. F. D., Saavedra, O., Margeat, E., Milhiet, P.-E. & Costa, L. Synchronous, Crosstalk-free Correlative AFM and Confocal Microscopies/Spectroscopies. *Sci. Rep.* **10**, 7098 (2020).

49. Cartagena-Rivera, A. X., Van Itallie, C. M., Anderson, J. M. & Chadwick, R. S. Apical surface supracellular mechanical properties in polarized epithelium using noninvasive acoustic force spectroscopy. *Nat. Commun.* **8**, 1030 (2017).

50. Györffy, B. *et al.* An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast Cancer Res. Treat.* **123**, 725–731 (2010).

51. Nusse, R. & Clevers, H. Wnt/β-Catenin Signaling, Disease, and Emerging Therapeutic Modalities. *Cell* **169**, 985–999 (2017).

52. Furth, N. & Aylon, Y. The LATS1 and LATS2 tumor suppressors: beyond the Hippo pathway. *Cell Death Differ.* **24**, 1488–1501 (2017).

53. Misra, J. R. & Irvine, K. D. The Hippo Signaling Network and Its Biological Functions. *Annu. Rev. Genet.* **52**, 65–87 (2018).

54. Cánovas, B. *et al.* Targeting p38α Increases DNA Damage, Chromosome Instability, and the Anti-tumoral Response to Taxanes in Breast Cancer Cells. *Cancer Cell* **33**, 1094-1110.e8 (2018).

55. Gupta, J. & Nebreda, A. R. Roles of p38α mitogen-activated protein kinase in mouse models of inflammatory diseases and cancer. *FEBS J.* **282**, 1841–1857 (2015).

56. Gupta, J. *et al.* Dual function of p38α MAPK in colon cancer: suppression of colitis-associated tumor initiation but requirement for cancer cell survival. *Cancer Cell* **25**, 484–500 (2014).

57. Hardwick, J. C., van den Brink, G. R., Offerhaus, G. J., van Deventer, S. J. & Peppelenbosch, M. P. NF-kappaB, p38 MAPK and JNK are highly expressed and active in the stroma of human colonic adenomatous polyps. *Oncogene* **20**, 819–827 (2001).

58. Maik-Rachline, G., Zehorai, E., Hanoch, T., Blenis, J. & Seger, R. The nuclear translocation of the kinases p38 and JNK promotes inflammation-induced cancer. *Sci. Signal.* **11**, (2018).

59. Hildebrand, S. *et al.* The E-cadherin/AmotL2 complex organizes actin filaments required for epithelial hexagonal packing and blastocyst hatching. *Sci. Rep.* **7**, 9540 (2017).

60. Couzens, A. L. *et al.* Protein interaction network of the mammalian Hippo pathway reveals mechanisms of kinase-phosphatase interactions. *Sci. Signal.* **6**, rs15 (2013).

61. Zhao, B. *et al.* Angiomotin is a novel Hippo pathway component that inhibits YAP oncoprotein. *Genes Dev.* **25**, 51–63 (2011).

62. Frixen, U. H. *et al.* E-cadherin-mediated cell-cell adhesion prevents invasiveness of human carcinoma cells. *J. Cell Biol.* **113**, 173–185 (1991).

63. Yang, J. *et al.* Guidelines and definitions for research on epithelial-mesenchymal transition. *Nat. Rev. Mol. Cell Biol.* (2020) doi:10.1038/s41580-020-0237-9.

64. Kleer, C. G., van Golen, K. L., Braun, T. & Merajver, S. D. Persistent E-cadherin expression in inflammatory breast cancer. *Mod. Pathol. Off. J. U. S. Can. Acad. Pathol. Inc* **14**, 458–464 (2001).

65. Rodriguez, F. J., Lewis-Tuffin, L. J. & Anastasiadis, P. Z. E-cadherin's dark side: possible role in tumor progression. *Biochim. Biophys. Acta* **1826**, 23–31 (2012).

66. Padmanaban, V. *et al.* E-cadherin is required for metastasis in multiple models of breast cancer. *Nature* **573**, 439–444 (2019).

67. Campbell, C. I. *et al.* The RNF146 and tankyrase pathway maintains the junctional Crumbs complex through regulation of angiomotin. *J. Cell Sci.* **129**, 3396–3411 (2016).

68. Wang, W. *et al.* Tankyrase Inhibitors Target YAP by Stabilizing Angiomotin Family Proteins. *Cell Rep.* **13**, 524–532 (2015).

69. Couderc, C. *et al.* AMOTL1 Promotes Breast Cancer Progression and Is Antagonized by Merlin. *Neoplasia N. Y. N* **18**, 10–24 (2016).

70. Mojallal, M. *et al.* AmotL2 disrupts apical-basal cell polarity and promotes tumour invasion. *Nat. Commun.* **5**, 4557 (2014).

71. Hultin, S. *et al.* AmotL2 links VE-cadherin to contractile actin fibres necessary for aortic lumen expansion. *Nat. Commun.* **5**, 3743 (2014).

72. Wada, M. *et al.* P38 delta MAPK promotes breast cancer progression and lung metastasis by enhancing cell proliferation and cell detachment. *Oncogene* **36**, 6649–6657 (2017).

73. Zanconato, F., Cordenonsi, M. & Piccolo, S. YAP/TAZ at the Roots of Cancer. *Cancer Cell* **29**, 783–803 (2016).

74. Bartucci, M. *et al.* TAZ is required for metastatic activity and chemoresistance of breast cancer stem cells. *Oncogene* **34**, 681–690 (2015).

75. Cordenonsi, M. *et al.* The Hippo transducer TAZ confers cancer stem cell-related traits on breast cancer cells. *Cell* **147**, 759–772 (2011).

76. Di Agostino, S. *et al.* YAP enhances the pro-proliferative transcriptional activity of mutant p53 proteins. *EMBO Rep.* **17**, 188–201 (2016).

ACKNOWLEDGEMENTS

We acknowledge the contribution of G. Froment, D. Nègre and C. Costa from lentivectors' production facility of SFR Biosciences (UMS3444/CNRS, US8/Inserm, ENS Lyon, UCBL, France). We thank the ABIC (IRCM), MRI (IRCM), MGC (IGMM) platforms, and the URT (IRCM, Montpellier, France). We thank V. Chambon and T. Soirat for technical help with the mammosphere formation assay and PE Milhiet (CBS, Montpellier, France) for helpful discussion and AFM set up. All plasmids obtained from Addgene are detailed in the Material and Methods section and all references have been included. We also thank Drs C. Gongora, L. Holmgren, N. Tapon, and B. Zhao for discussion, tools and advice. This work was supported by grants from Fondation ARC and Ligue Régionale Contre le Cancer (34) to LHM and AD. DK is supported by Ligue Nationale Contre le Cancer. VS and EB were supported by Fondation de France.

AUTHOR CONTRIBUTIONS

DK, VS, EB, YBS, EF, ML, FBM, CB, LC, BO, and LHM performed experiments. LHM and AD designed the experiments. DK, JC, BO, LHM, and AD analyzed the data. DK, PL, CG, JC, CT, AM, LHM, and AD interpreted the data. LHM and AD wrote the manuscript.

FIGURE LEGENDS

Figure 1. MAGI1 impairment induces tumorigenic phenotypes in epithelial cells

(A) MTT assay (OD 560 nm) representing 2D cell growth of MCF7*shMAGI1* as compared to MCF7*shLuc*

cells. Bars represent mean ± Standard Deviation (SD; n=10 wells as replicates) of a representative

experiment (out of 3). Unpaired two-tailed Student's t-test; *** $p < 0.001$.

(B) Cell cycle phases of MCF7*shLuc* (a) and MCF7*shMAGI1* (b) cells labeled with propidium iodide and

analyzed by Flow cytometry, showing a slight increased proportion of cells in S phase at the expense

of G0/G1 after *MAGI1* invalidation.

(C) Top: quantification of colony numbers of MCF7*shMAGI1* cells grown in anchorage independent

conditions (soft agar assay) and represented as fold increase compared to MCF7*shLuc* cells. Data are

presented as the means ±SD (n=3). Unpaired two-tailed Student's t-test; *** $p < 0.001$. Bottom:

western blot analysis of whole protein extracts issued from the cells used in the soft agar assays and

showing the relative amounts of MAGI1 in the different cell lines; Tubulin was used as a loading control.

(D) Top: quantification of mammospheres represented as total area issued from MCF7*shMAGI1* cells

and shown as fold increase compared to MCF7*shLuc* cells, starting from 200 cells. Data are presented

as the means ±SD (n=4). Unpaired two-tailed Student's t-test; *** $p < 0.001$. Bottom: images of

mammosphere formation assay showing a representative experiment.

(E) Primary tumor growth of MCF7*shLuc* and MCF7*shMAGI1* cells injected subcutaneously in nude

mice. Primary tumor growth was assessed by measuring the tumor volume over time until the tumors

were too big and the mice had to be euthanized. Bars correspond to the mean ± Standard Error to the

Mean (n=8 mice per group). Unpaired two-tailed Student's t-test; * $p < 0.05$.

Figure 2. The loss of MAGI1 affects E-cadherin levels and localization in MCF7 cells

(A) Representative immunofluorescence experiments performed in MCF7 cells monitoring MAGI1

localization (red) compared to E-cadherin, ZO1, and Claudin3 (green). DAPI (blue) was used to stain

DNA and the nuclei. White arrows indicate staining overlap. Scale bar=10 μm.

(B) Western blot analysis of whole protein extracts (n=4) monitoring the expression the junctional

components Claudin-1, ZO-1, β-catenin, PARD3, E-cadherin and AMOTL2 in MCF7*shMAGI1* cell lines

compared to MCF7*shLuc*. Tubulin was used as a loading control. Note the increase in E-cadherin levels. Western blot experiments have been repeated at least four times. Protein expression levels were quantified as compared to Tubulin (not shown).

(C) Representative immunofluorescence experiments (n=3) were performed on MCF7*shMAGI1* compared to MCF7*shLuc* cells monitoring the localization of β-catenin, PARD3, Claudin-3, ZO-1 and E-cadherin (green). DAPI (blue) was used to stain DNA and the nuclei. Scale bar=10 μm. Note the slight sub-cortical accumulation of E-cadherin shown in the high magnification images.

Figure 3. The loss of MAGI1 affects MCF7 cell compaction, ROCK activity, and compressive forces

(A) Representative phase contrast and fluorescence images of MCF7shLuc and MCF7shMAGI1 cells grown in 3D spheroid cultures and stained with E-cadherin (red) and DAPI (blue).

(B) Calculated perimeters for 3D spheroid cultures of MCF7shMAGI1 normalized by the perimeter of MCF7shLuc cells. Bars represent mean ±SD (n=10 spheroids) of five independent experiments. Unpaired two-tailed Student's t-test; * $p < 0.05$.

(C) Calculated circularity for 3D spheroid cultures of MCF7shLuc and MCF7shMAGI1 (calculations were done with the ImageJ software where a value of 1 is considered as a perfect circle). Bars represent mean ±SD (n=10 spheroids) of five independent experiments. Unpaired two-tailed Student's t-test; * $p < 0.05$.

(D) Elastic Young's modulus (EYM) of cells: Hertz contact mechanics model for spherical indenters was used. In MCF7shMAGI1 cells, the apical surface EYM is significantly elevated when compared to control MCF7shLuc cells. Data are represented as mean +/- standard deviation: shLuc EYM= 258.8 +/- 101, shMAGI1 EYM= 302.4 +/- 91. Unpaired two-tailed Student's t-test; * $p < 0.05$ (n=132 and 94 for shLuc and shMAGI1 respectively).

(E) Western blot analysis on whole protein extracts (n=3) of ROCK-specific ser19 phosphorylation of Myosin Light Chain 2, total MLC2 and MAGI1 in MCF7*shMAGI1* cell lines compared to MCF7*shLuc*. GAPDH was used as a loading control.

(F) Quantification of the representative Western blot showing protein expression represented in panel E. Phosphorylated proteins were quantified as compared to their total protein counterparts to evaluate their activation.

Figure 4. The loss of MAGI1 tumorigenic phenotypes are not associated with active YAP

(A) Western blot analysis on whole protein extracts (n=3) of phosphorylated YAP (S397 & S127), total YAP and TAZ, phosphorylated LATS1, total LATS1, phosphorylated MOB1, non-phosphorylated (active) β-catenin, total β-catenin and MAGI1 in MCF7*shMAGI1* cell lines compared to MCF7*shLuc*. Tubulin was used as a loading control.

(B) Representative immunofluorescence images (n=3) of endogenous MAGI1 and YAP protein expression in MCF7*shLuc* and MCF7*shMAGI1* cells cultured to similar confluence (60.4 % and 63.2 % respectively). Scale bar=10 μm.

(C) Quantification of MAGI1 and YAP staining in a tissue micro-array (TMA) from luminal breast cancer patients in a cohort of 28 patients, showing the strong correlation between MAGI1 membranous and YAP nuclear localizations; Fisher exact test, *** $p < 0.001$. Histogram shows the percentages of patients having a nuclear or a cytoplasmic YAP staining according to the absence or the presence of MAGI1 membranous staining.

(D) Real-time qRT-PCR showing the expression of canonical Hippo pathway target genes (CTGF, CYR61, BIRC2 and AREG) in MCF7*shMAGI1* compared to the control MCF7*shLuc* cells and normalized with GAPDH expression. Data are presented as the means ±SD (n=3). Unpaired two-tailed Student's t-test; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.

Figure 5. The loss of MAGI1 tumorigenic phenotypes are associated with p38 stress signaling pathway activation

(A) Western blot analysis showing protein expression and/or activation of Akt and MAPK proliferation signaling pathways as well as p38 and JNK stress signaling pathways in MCF7*shMAGI1 compared* to MCF7*shLuc* cells. Tubulin was used as a loading control. Western blot experiments have been repeated at least three times.

(B) Quantification of the representative Western blot showing protein expression represented in panel A. Phosphorylated proteins were quantified as compared to their total protein counterparts to evaluate their activation.

(C) 2D cell growth assay of MCF7*shMAGI1* treated, or not, with the p38 MAPK inhibitor LY2228820 (LY in the figure) compared to MCF7*shLuc*. Bars represent mean ±SD (n=6 wells as replicates) of a

representative experiment (n=3). Unpaired two-tailed Student's t-test; * p<0.05. Note the increased

percentage of drop for MCF7*shMAGI1* cells at day 8.

(D) Quantification of colony numbers of MCF7*shMAGI1* cells non treated or treated with LY2228820

grown in anchorage independent conditions (soft agar assay) and represented as fold increase

compared to MCF7*shLuc* cells non treated or treated with the p38 inhibitor (LY in the figure). Data are

presented as the means ±SD (n=3). Unpaired two-tailed Student's t-test; *** p < 0.001.

(E) Western blot analysis showing protein levels of phosphorylated p38 and total p38 in MCF7*shMAGI1*

cells treated or non-treated (NT) with Blebbistatin (Blebb) or Y-27632. Both inhibitors were used at 10

μM during 2 h and Tubulin that was used as a loading control.

(F) Quantification of the representative Western blot showing protein expression and represented in

panel E. Phosphorylated p38 was quantified as compared to total p38 to evaluate its activation.

Figure 6. MAGI1 interacts with AMOT and AMOTL2

(A) Western blot of Flag-MAGI1 immunoprecipitates: Flag-MAGI1 transfections were performed in

MCF7 cells and endogenous AMOT (upper panels) and AMOTL2 (lower panels) were revealed; Flag

blotting was used as the immunoprecipitation control. Note that we could hardly detect Flag-MAGI1

by Western blot in the inputs, even though MAGI1 was well immunoprecipitated, due to sensitivity

issues associated with single Flag tag in Flag-MAGI1 used in this typical experiment.

(B) Western blot analysis of endogenous MAGI1 immunoprecipitates revealing the presence of

endogenous AMOT (upper panels) and AMOTL2 (lower panels). MAGI1 blotting was used as the

immunoprecipitation control (*shows non-specific bands with the anti-MAGI1 antibody).

(C) Schematic of full length MAGI1 and of the point MAGI1 WW domain mutants used (P331A &

P390A).

(D) Western blot analysis of Flag-MAGI1 immunoprecipitates on protein extracts from MCF7 cells

transfected with HA-AMOT and different Flag tagged MAGI1 constructs, showing that the interaction

with AMOT occurred through the second WW domain of MAGI1 (interaction lost in P390A; red).

(E) Western blot analysis of Flag-MAGI1 immunoprecipitates on protein extracts from MCF7 cells

transfected with Flag-MAGI1 or Flag-MAGI1 P390A mutant, revealing the interaction of endogenous

AMOTL2 with Flag-MAGI1 but not Flag-MAGI1 P390A (red).

(F) Schematic of full-length AMOT and of the point AMOT PPXY mutants used (Y242A & Y287A).

(G) Western-blot analysis of Flag-MAGI1 immunoprecipitates on protein extracts from MCF7 cells transfected with Flag-MAGI1 and different HA-AMOT constructs, showing that the interaction occurs through the second PPXY domain of AMOT (interaction lost in Y287A; red). Similar levels of interaction were obtained for HA-AMOT/Flag-MAGI1 and HA-AMOT Y242A/Flag-MAGI1 complexes (quantified using the ImageJ software).

(H) Representative immunofluorescence images of endogenous staining for MAGI1 (red; top panels) co-localizing at the plasma membrane (white arrows) with AMOT (green; left) or AMOTL2 (green; right) in MCF7 cells. Note the non-specific nuclear staining for AMOT. Scale bar=10 µm.

(I) Western blot analysis after subcellular fractionation showing the relative amount of AMOTL2 protein in the cytoplasmic and membrane fractions in MCF7*shMAGI1* compared to MCF7*shLuc* cells. Tubulin was used as a cytoplasmic control for the fractionation and for normalization.

Figure 7. AMOTL2 mediates the effects of MAGI1 on junctions and on p38 signaling

(A) MTT assay (OD 560 nm) representing 2D cell growth of MCF7*shMAGI, MCF7shAMOTL2* and MCF7*shMAGI1/shAMOTL2* cells as compared to MCF7*shLuc.* Bars represent mean ±SD (n=10 wells as replicates) of a representative experiment (n=3). Unpaired two-tailed Student's t-test; ** $p<0.01$; *** $p < 0.001$.

(B) Quantification of colony numbers of MCF7 cells (*shMAGI1, shAMOTL2* & *shMAGI1/shAMOTL2*) grown in anchorage independent conditions (soft agar assay) and represented as fold increase compared to MCF7*shLuc* cells. Data are presented as the means ±SD (n=3). Unpaired two-tailed Student's t-test; * $p < 0.05$. Western blot analyses confirmed the 90 % knockdown for MAGI1 in MCF7*shMAGI1* and MCF7*shMAGI1/shAMOTL2* and QPCR data confirmed the 50 to 60 % knockdown for AMOTL2 in MCF7*shAMOTL2* and MCF7*shMAGI1/shAMOTL2* respectively (determined by RT-qPCR and/or Western Blot ; Data not shown).

(C) Western blot analysis showing protein expression and/or phosphorylation (activation) of p38 and JNK stress signaling pathways as well as junctions' components in MCF7*shMAGI1* as compared to MCF7*shLuc* cells when *AMOTL2*siRNA were transfected. Tubulin was used as a loading control. Western blot experiments have been repeated at least three times.

(D) Quantification of the representative Western blot showing protein expression and represented in panel C. Phosphorylated proteins were quantified as compared to their total protein counterparts to evaluate their activation and junctions' proteins were quantified as compared to tubulin that was used as a loading control. Upper: Phosphorylated p38 / Total p38 protein quantification and Lower: protein / Tubulin quantification.

(E) Model for the role of MAGI1 during Luminal BCa. MAGI1 prevents the accumulation of junctional AMOTL2 and E-cadherin as well as ROCK activity thus releasing cellular stiffness. Increased AMOTL2 and ROCK then activate p38 stress signaling responsible for the increased tumorigenicity of MAGI1-deficient cells. Anti and Pro tumorigenic events are highlighted in green and red respectively.

SUPPLEMENTAL MATERIAL

SUPPLEMENTAL FIGURE LEGENDS

Suppl Figure 1

(A) Western blot showing MAGI1 expression in different breast cancer cell lines (T47D and MCF7 from

the luminal A sub-type, MDA-MB-468 and BT549 from basal sub-type and ZR75 from luminal B sub-

type. Tubulin was used as loading control.

(B) Kaplan-Meier survival curves and log-rank analysis. Relapse-free survival (RFS) curves for 841 ER

positive breast cancer patients as a function of MAGI1 expression. Patients were split according to

median expression. Data for MAGI1 (225474_at) were obtained using the KM plotter website at

http://kmplot.com (Gyorffy B, Lanczky A, et al, Breast Cancer Res Treatment, 2010).

(C) Representative immunohistochemical staining of MAGI1 performed in normal human breast

samples. Brown staining indicates positive immunoreactivity and arrows show MAGI1 staining.

(D) Western blot analysis of MAGI1, MAGI2 and MAGI3 expression in MCF7*shMAGI1(1-1)* and

MCF7*shMAGI1(3-1)* showing respectively 50% and 100% knockdown as compared to MCF7*shLuc*

control cell line. Tubulin was used as a loading control.

(E) Representative immunofluorescence images showing MAGI1 staining (red) in the *MCF7shMAGI1*

(1-1) and *(3-1)* and in MCF7*shLu*c cell lines. Scale bar=10 μm.

(F) Real-time qRT-PCR showing the expression of *MAGI1*, *MAGI2* and *MAGI3* mRNA in

MCF7*shMAGI1(1-1)* and in MCF7*shMAGI1(3-1)* compared to the control MCF7*shLuc* cell line. Data are

presented as the means ± SD. Three independent experiments; unpaired two-tailed Student's t-test (*

$p < 0.05$; n.s. not significant).

Suppl Figure 2

(A) MTT assay (OD 560 nm) representing 2D cell growth of T47D*shMAGI1* as compared to T47D*shLuc*

cells. Bars represent mean ± SD (n=10 wells as replicates) of a representative experiment (n=3),

showing similar increased cell growth as observed for MCF7 cells. Unpaired two-tailed Student's t-test;

*** p < 0.001.

(B) Upper: quantification of the number of colonies of T47D*shMAGI1* grown in anchorage independent

conditions (soft agar assay) represented as fold increase compared to T47D*shLuc* cells. Data are

presented as the means (n=3) ± Standard Deviation. Unpaired two-tailed Student's t-test; ** p < 0.01.

Below: western blot analysis on whole protein extracts from cells used on the left showing the efficient

MAGI1 protein knockdown; Tubulin was used as a loading control.

(C) Calculated perimeters for 3D spheroids cultures of T47D*shMAGI1* (shMAGI1(3-1) cell line only)

normalized by the perimeter of T47D*shLuc* cells. Bars represent mean ± SD (n=10 spheroids) of three

independent experiments. Unpaired two-tailed Student's t-test; * p < 0.05.

(D) Calculated circularity for 3D spheroid cultures of T47D*shLuc* and T47D*shMAGI1* (shMAGI1(3-1) cell

line only). Calculations were done with the ImageJ software where a value of 1 is considered as a

perfect circle. Bars represent mean ± SD (n=10 spheroids) of three independent experiments. Unpaired

two-tailed Student's t-test; * p < 0.05.

(E) Upper: quantification of the number of colonies of HCT116*shMAGI1* grown in anchorage

independent conditions (soft agar assay) represented as fold increase compared to HCT116*shLuc*. Data

are presented as the means (n=3) ± Standard Deviation. Unpaired two-tailed Student's t-test; * p <

0.05. Below: analysis on whole protein extracts from cells used on the left showing the efficient MAGI1

protein knockdown; Tubulin was used as a loading control.

(F) Primary tumor growth of HCT116*shLuc* and HCT116*shMAGI1* cells injected subcutaneously in nude

mice. Primary tumor growth was assessed by measuring the tumor volume over time until the tumors

were too big and the mice had to be euthanized. Bars correspond to the average ± SEM (n=8 mice).

Unpaired two-tailed Student's t-test (* p < 0.05).

Suppl Figure 3

(A&B) Representative immunofluorescence images of endogenous AMOTL2 and actin (phalloidin red

staining) in MCF7*shLuc* (A) and MCF7*shMAGI1* (B) showing unaffected localisation at the plasma

membrane. Scale bar=10 µm.

(C) Elastic Young's modulus (EYM) of cells treated, or not, with Y-27632 (10µM for 45 min): Hertz

contact mechanics model for spherical indenters was used. In treated cells, the apical surface EYM is

significantly decreased when compared to untreated cells. Data are represented as mean +/- standard

deviation: shLuc untreated EYM= 248.4 +/- 133.5, treated with Y-27632 EYM= 112.0 +/- 81.5, shMAGI1

untreated EYM= 321.1 +/- 107.1 and shMAGI1 treated with Y-27632 EYM= 103.3 +/-79.9. Anova and

unpaired two-tailed student's *t*-test; * $p < 0.05$ (n=67, 49, 57 and 63 for untreated and treated control

shLuc and shMAGI1, respectively).

A

2D cell growth assay (MCF7)

OD (560 nm)

■ MCF7shLuc
■ MCF7shMAGI1

***

Day0 Day1 Day2 Day3 Day4 Day7

B

a

G2/M 22%   SubG1 3%
                      G2/M 22%
S 21%   G0/G1 54%

MCF7shLuc

b

SubG1 2%
S 29%   G0/G1 47%

MCF7shMAGI1

C

Soft agar assy(MCF7)

Colony numbers (Fold Increase)

***

shLuc   shMAGI1

shLuc shMAGI1

MAGI1

Tubulin

D

Mammosphere formation assay (MCF7)

Green total area (Fold increase)

***

shLuc   shMAGI1

shLuc   shMAGI1

E

Primary tumor growth (MCF7)

Tumor volume (mm³)

● shLuc
■ shMAGI1

*
(n=8)

Days post implantation

Figure 2

MCF7shLuc MCF7shMAGI1

Phase contrast (2X)

Ecadh/DAPI (10X)

A

B

Spheroid perimeter
(MCF7)

C

Spheroid circularity
(MCF7)

D

E

shLuc shMAGI1

MLC2

P-MLC2

MAGI1

GAPDH

F

P-MLC2

Figure 4

Figure 5

Figure 6

Figure 7

A

2D cell growth assay (T47D)

B   Suppl Figure 2

Soft agar assay (T47D)

C

Spheroid perimeter (T47D)

D

Spheroid circularity (T47D)

E

Soft agar assay (HCT116)

F

Primary tumor growth (HCT116)

A

MCF7shLuc

B

MCF7shMAGI1

C