Research Article

Animal Behavior

Title:

Sex differences in vocalizations to familiar or unfamiliar females in mice

Eri Sasaki*, Yuiri Tomita*, and Kouta Kanno.

Affiliations

Lab. of Neuroscience, Course of psychology, Faculty of Law, Economics and

Humanities, Kagoshima University

* These four authors contributed equally to this work.

Correspondence: Kouta Kanno, Ph.D.

Lab. of Neuroscience, Course of Psychology

Department of Humanities,

Faculty of Law, Economics, and Humanities,

Kagoshima University

Korimoto 1-21-30, Kagoshima City, Kagoshima 890-0065, Japan

E-mail: canno@leh.kagoshima-u.ac.jp

Tel: +81-99-285-7624

Fax: +81-99-285-7609

Abstract

Mice, both wild and laboratory strains, emit ultrasounds to communicate. The

sex differences between male to female (male–female) and female to female (female–

female) ultrasonic vocalizations (USVs) have been discussed for decades. In the present

study, we compared the number of USVs emitted to familiar and unfamiliar female by

both males (male–female USVs) and females (female–female USVs). We found that

females vocalized more to unfamiliar than to familiar females. In contrast, males

exhibited more USVs to familiar partners. This sexually dimorphic behavior suggests

that mice change their vocal behavior in response to the social context, and their

perception of the context is based on social cognition and memory. In addition, because

males vocalized more to familiar females, USVs appear to be not just a response to

novel things or individuals, but also to be a social response.

Keywords: ultrasonic vocalizations, social behavior, affiliation, social cognition, and

memory

1. Introduction

Social animals utilize signals for some modalities to communicate with other individuals. Among such signals, vocal communications are widely observed in animals and are superior in many aspects [1]. Recently, vocal communication in mice has attracted considerable academic attention because of its power and utility in the investigation of molecular and neural mechanisms for social behaviors and their deficits [2–4]. In mice, ultrasonic vocalizations (USVs) are used for communication, and they can be widely observed in both wild and laboratory strains [5]. There are two types of USVs commonly known in mice: pupUSVs and courtship vocalizations [4]. Mice pups in the first several days after birth emit ultrasounds (pupUSVs), mainly when they are outside their nest. A state of arousal due to the perception of cold, perception of unusual tactile stimulation, or loss of social contact triggers the emission of pupUSVs, which enhance the mother's approach to the sound source [6]. The mother's attention and movement lead to the expression of maternal behavior. On the other hand, adult male mice emit ultrasounds in the presence of adult females or her urine [7]; these are known as courtship vocalizations or courtship songs [8]. Activational effects of sex hormones

on male vocalizations have been reported [9,10]. Furthermore, the number of male

USVs is enhanced by sociosexual experience with females [11], and usage of syllable

types (call patterns) is modulated according to the progressive phase of sexual behaviors

[12]. Thus, USVs from males to females are thought to be male-specific precopulatory

behaviors [11,13].

In this regard, vocalizations in mice involved in these two contexts have been

well investigated since they were first reported about fifty years ago [14–16]. In the

process of such research history, adult females were initially not thought to vocalize

either to males or females [17]. However, using several strains, Maggio and Whitney

thoroughly revealed that adult females emit USVs to females (female–female), where

they rarely emit to males (female–male) and that the number of female–female USVs is

comparable to male–female courtship USVs [18]. Thereafter, a study observing female–

female USVs in albino NMRI female mice indicated that females vocalize more to

unfamiliar females than to familiar ones [19]. This finding suggests that female–female

USVs reflect social cognition and memory. In fact, administration of a classical

cholinergic antagonist (scopolamine) disrupted this social memory [19]. Although

female–female USVs are known to be related to this social memory, the functions of these USVs is poorly understood.

In the study by D'Amat et al. [19] described above, familiar individuals were those that were previously presented only several minutes in advance (while simultaneously recording ultrasound). Therefore, we decided to use female individuals that were reared in the same cage after weaning as familiar individuals. In other words, we hypothesized that female–female USVs reflect social affiliation if females emit USVs more to familiar females than to unfamiliar ones under such conditions. If this hypothesis is correct, the results of the study by D'Amat et al. and our expected results could be explained as follows: females express high levels USVs to establish an affiliative relationship with a stranger of the same sex first, and, at the same time, females also emit more USVs to familiar social members. In the present study, we also compared female–female vocalizations and male–female vocalizations using the B6 strain, which is currently the most commonly used laboratory strain. Social cognition and memory in male mice has not yet been measured in this context, but it is important to measure the sociality of both sexes using the same methods because mouse models

and USVs have been recently investigated to clarify the basic mechanisms of

developmental disorders [2–4].

2. Materials and Methods

(a) Animals

We used an inbred strain of B6 (C57BL/6J) mice. Mice were purchased from

Japan SLC (Hamamatsu, Japan). Subject animals were used when they were 8–10

weeks-old for recording tests.

For recordings of male–female USVs, a male and an age-matched unfamiliar

female mouse were cohoused in a standard cage (182 × 260 × 128 mm, CREA Japan)

immediately following arrival from the supplier until experiments began.

For recordings of female–female USVs, females that were bred together in the

same cage after weaning were purchased, and two individuals were housed in each cage

immediately after arrival from the supplier until experiments began. One was used as

the subject, and the other was used as the familiar partner (see Fig. 1A).

For both recordings, age-matched, unfamiliar adult females were also

purchased, and 3–4 animals were housed in the same cage. These animals were used

repeatedly as an unfamiliar intruder for recordings (see below and Fig. 1A).

Food and water were supplied *ad libitum*, and the animals were kept under a

standard 12-h:12-h light-dark cycle. The environment was maintained at a constant

temperature (22–25°C) and humidity (50 ± 5%). All experimental procedures were

approved by the Institutional Animal Use Committee of Kagoshima University

(#L18004 and #L19003).

(b) Experimental design and general procedures.

The subject male was cohoused with an unfamiliar female, and the subject

female was cohoused with a female that was bred together with the subject female in the

same cage after weaning, as described above. These male–female and female–female

were cohoused for 7 days without exchange of beddings. At the 7th day (Day 7 shown in

Fig. 1A) after the pairing of male–female and female–female began (Day 0 shown in

Fig. 1A), the partner female was removed. Recordings of USVs were conducted twice

(recording 1 and 2) for all subjects, and two experimental conditions were tested in both

male–female and female–female contexts; some subjects were used for recording 1 30

min after the removal of the partner, and the other subjects were used 1 day (24–25 h)

after the removal. Recording 2 was conducted 30 minutes after recording 1 for all

subjects. USVs were induced by introducing a female mouse. The subjects were

exposed to the familiar partner and an unfamiliar female once each, and the order of

encountering the familiar or unfamiliar partner was counter-balanced between

recordings 1 and 2.

(c) Settings and procedures for ultrasound recordings

The microphone was hung 16 cm above the floor of a sound-attenuating

chamber. Immediately prior to recording, the home cage of the subjects was moved into

the chamber where a red dim light was placed. The ultrasound recording was performed

for 100 s after the intruder was introduced into the home cage.

According to previous studies [12,20,21], we conducted recordings without

the devocalization of female encounters under natural conditions. Previous experiments

suggested that most vocalizations are produced by the resident subjects and that the

vocal contribution of female encounters is very limited in the male–female and female–

female contexts [22–24].

(d) Ultrasound analysis

Most of the settings for the USV recordings followed those of our previous

study [11], with some modifications. Ultrasonic vocalizations were recorded using a

CM16/CMPA condenser microphone (Avisoft Bioacoustics, Berlin, Germany),

ultrahigh-speed ADDA converter BSA768AD-KUKK1710 and its software

SpectoLibellus2D (Katou Acoustics Consultant Office, Kanagawa, Japan) with a

sampling rate of 384 kHz (to measure 20-192 kHz).

Recorded sounds were saved on a computer as .wav files using

SpectoLibellus2D. These sound files were analyzed with the GUI-based software

USVSEG (implemented as MATLAB scripts) recently developed by us [25].

Continuous sound signals with frequencies ranging from to 40–160 kHz and durations

ranging from 3 to 300 ms were analyzed and detected as syllables. Each syllable was

segmented and saved as a jpeg image file, and the noise data (false positives) were

manually excluded. Finally, the number of vocalizations (syllables) was quantified.

(e) Statistics

Statistical analyses were performed using JMP 14 software (SAS Institute,

Cary, NC, USA). In the present study, most of the data seemed not to be distributed

normally; therefore, nonparametric tests were used. To compare two conditions within

an individual, Wilcoxon signed-rank tests were applied (see Fig. 1), and Kruskal–Wallis

tests were used for comparison among groups, followed by post-hoc Steel–Dwass tests

(see Fig. 2). Values are reported as dot plots with boxplots. Differences with p values P

< 0.05 were considered significant.

3. Results

(a) Female–female vocalization

Under the 30 min isolation condition, the number of USVs emitted from

subject females to unfamiliar female was significantly higher than to the familiar

partner (Fig. 1B; Wilcoxon signed-rank test, $W_{(14, 14)} = 39.0$, $P = 0.02$). On the other

hand, there was no significant difference between the number of USVs emitted to the

unfamiliar and familiar partner after 1 day isolation (Fig. 1C; Wilcoxon signed-rank test,

$W_{(11, 11)} = 3.0$, $P = 0.9063$).

(b) Male–female vocalization

Under the 30 min isolation condition, there was no significant difference

between the number of USVs emitted from males to unfamiliar and familiar partners

(Fig. 1D; Wilcoxon signed-rank test, $W_{(15, 15)} = 26.0$, $P = 0.1758$). On the other hand, the

number of USVs emitted to the familiar partner was significantly higher than that of the

unfamiliar female after 1 day isolation (Fig. 1E; Wilcoxon signed-rank test, $W_{(15, 15)} =$

-47.5, $P = 0.0067$).

(c) Total comparison among groups

Differences between the number of USVs emitted to familiar and unfamiliar

partners ([#. familiar] - [#. unfamiliar]) were calculated from the data shown in Fig.1

and were compared among the 4 groups (Fig. 2). The Kruskal–Wallis test indicated

significant differences between groups (χ^2 = 19.4288, df = 3, P = 0.0002) and the Steel–

Dwass test (performed *post-hoc*) showed that the value of males under the 1 day

isolation condition was significantly higher than that of males or females under the 30

min isolation condition (P = 0.0085 and P = 0.0025, respectively), or females under the

1day isolation condition (P = 0.0337).

4. Discussion

As described in the Introduction, we prepared female individuals with

affiliative relations. Nevertheless, contrary to our expectation, the result of the previous

study of D'Amat et al. [19] was replicated in our 30 min isolation condition (Fig. 1B);

here we found that females vocalized more to an unfamiliar female than to familiar

females. It is unclear as to why females rarely vocalized under the 1day isolation

condition, but no difference was found between the number of USVs to the unfamiliar

and familiar females under such conditions, as was the case in D'Amat et al's studies.

We believe that it is safe to conclude that B6 females vocalize more to unfamiliar

females.

On the other hand, it was surprising that males exhibited an opposite tendency

toward females. It has been reported that male courtship USVs are enhanced by

sociosexual experience [11] and that vocal usage is altered according to the progressive

phase of sexual behavior [12]. These results indicate that male USVs directed towards

females express sexual motivation. We expected males to show more vocalization to

unfamiliar females; however, this was not a "flirtatious"' tendency, as males made many

vocalizations to the familiar partner. Even though different experimental conditions,

such as length of cohousing, could possibly have led to different results, it appears that

males exhibited more sexual motivation toward the familiar partner (cohoused for 1

week) under the present conditions.

Although it is not clear why the optimal conditions for the observation of

differential vocal responses to the familiar or unfamiliar females differ between males

and females, the important finding of this study is that there is a sex difference in such

vocal responses. For decades, the sex differences between male–female and female–

female USVs have been discussed. The amount of USVs between male–female and

female–female is comparable [18]. In addition, the vocal repertoire between sexes is

also reported to be comparable [23]. In contrast, the sex differences shown in the

present study indicate that mice change their vocal behavior in response to the context

of whom they are dealing with; their perception of the context is based on social

cognition and memory. In addition, because males vocalized more to familiar females,

USVs are thought to be not just a response to novel things or individuals but also to be a

social response. These findings with wild-type mice provide a practical behavioral assay

for the disease model mice, focusing on their social impairment.

Although such sex differences in USVs have been revealed, the past and

present studies have yet to determine the biological significance and function of

female–female USVs. Since Maggio and Whitney [18] hypothesized that female

vocalization contributes to the formation of social hierarchy among females, this

hypothesis has been referred to in some studies [19,23]. However, to our knowledge,

this hypothesis has never been investigated experimentally in mice. A recent study made

an important point about sex differences in acoustic features, with more diversity

(inter-individual difference) observed in the vocal repertoire in males than in females

[26]. Nonetheless, the biological significance and function of female USVs remain

unclear, and further studies, including investigation of the associated neural mechanisms,

are needed.

Ethics. All experimental procedures were approved by the Institutional Animal Use

Committee of Kagoshima University (#L18004 and #L19003).

Data accessibility. The dataset supporting this article are available from the figshare

Digital Repository: XXXXXXXXX

Authors' contributions K.K. contributed to the experimental design. E.S. and Y.T.

conducted the experiments. E.S., Y.T., and K.K. contributed to the interpretation of data

and writing of the manuscript.

Competing interests. There are no conflicts of interest to declare.

Funding. This work was supported by JSPS KAKENHI grant numbers 18K13371 and

19H04912.

Acknowledgments. This work was supported by a Grant-in-Aid for Scientific Research

on Innovation Areas "Integrative Research toward Elucidation of Generative Brain

Systems for Individuality" (19H04912) from MEXT to K.K. We thank various supports,

including daily discussion, of this project. We would like to thank Editage

(www.editage.com) for English language editing.

References

1. Brudzynski SM. 2010 Vocalization as an ethotransmitter: Introduction to the
handbook of mammalian vocalization. In *Handbook of Mammalian Vocalization* (ed
SM Brudzynski), pp. 3–9. ELSEVIER. (doi:10.1016/B978-0-12-374593-4.00001-2)

2. Fischer J, Hammerschmidt K. 2011 Ultrasonic vocalizations in mouse models for
speech and socio-cognitive disorders: insights into the evolution of vocal
communication. *Genes Brain Behav* **10**, 17–27.
(doi:10.1111/j.1601-183X.2010.00610.x)

3. Lahvis GP, Alleva E, Scattoni ML. 2011 Translating mouse vocalizations: prosody
and frequency modulation. *Genes Brain Behav* **10**, 4–16.
(doi:10.1111/j.1601-183X.2010.00603.x)

4. Konopka G, Roberts TF. 2016 Animal models of speech and vocal communication
deficits associated with psychiatric disorders *Biol Psychiatry* **79**, 53–61.
(doi:10.1016/j.biopsych.2015.07.001)

5. Sales GD. 2010 Ultrasonic calls of wild and wild-type rodents. In *Handbook of*

Mammalian Vocalization: An Integrative Neuroscience Approach (ed SM Brudzynski), pp. 77–88. ELSEVIER. (doi:10.1016/B978-0-12-374593-4.00009-7)

6. Ehret G. 2005 Infant rodent ultrasounds - A gate to the understanding of sound communication. *Behav Genet* **35**, 19–29. (doi:10.1007/s10519-004-0853-8)

7. Nyby J, Wysocki CJ, Whitney G, Dizinno G, Schneider J. 1979 Elicitation of male mouse (Mus musculus) ultrasonic vocalizations: I. Urinary cues. *J Comp Physiol Psychol* **93**, 957–975. (doi:10.1037/h0077623)

8. Holy TE, Guo Z. 2005 Ultrasonic songs of male mice. *PLoS Biol* **3**, e386. (doi:10.1371/journal.pbio.0030386)

9. Nyby J, Matochik JA, Barfield RJ. 1992 Intracranial androgenic and estrogenic stimulation of male-typical behaviors in house mice (Mus domesticus). *Horm Behav* **26**, 24–45.

10. Sipos ML, Nyby JG. 1996 Concurrent androgenic stimulation of the ventral tegmental area and medial preoptic area: synergistic effects on male-typical reproductive behaviors in house mice. *Brain Res* **729**, 29–44.

11. Kanno K, Kikusui T. 2018 Effect of sociosexual experience and aging on number of courtship ultrasonic vocalizations in male mice. *Zoolog Sci* **35**, 208–214. (doi:10.2108/zs170175)

12. Matsumoto YK, Okanoya K. 2016 Phase-specific vocalizations of male mice at the initial encounter during the courtship sequence. *PLoS One* **11**, e0147102. (doi:10.1371/journal.pone.0147102)

13. Burns-Cusato M, Scordalakes EM, Rissman EF. 2004 Of mice and missing data: what we know (and need to learn) about male sexual behavior. *Physiol Behav* **83**, 217–232. (doi:10.1016/j.physbeh.2004.08.015)

14. Zippelius HM, Schleidt WM. 1956 Ultraschall-Laute bei jungen Mäusen. *Naturwissenschaften* **43**, 502. (doi:10.1007/BF00632534)

15. Noirot E. 1966 Ultra-sounds in young rodents. I. Changes with age in albino mice.

Anim Behav **14**, 459–462. (doi:10.1016/S0003-3472(66)80045-3)

16. Sewell GD. 1967 Ultrasound in adult rodents. *Nature* **215**, 512. (doi:10.1038/215512a0)

17. Nyby J, Dizinno G, Whitney G. 1977 Sexual dimorphism in ultrasonic vocalizations of mice (Mus musculus): gonadal hormone regulation. *J Comp Physiol Psychol* **91**, 1424–1431.

18. Maggio JC, Whitney G. 1985 Ultrasonic vocalizing by adult female mice (Mus musculus). *J Comp Psychol Wash DC 1983* **99**, 420–436. (doi:10.1037/0735-7036.99.4.420)

19. D'Amato FR, Moles A. 2001 Ultrasonic vocalizations as an index of social memory in female mice. *Behav Neurosci* **115**, 834–840. (doi:10.1037/0735-7044.115.4.834)

20. Ey E, Torquet N, Le Sourd AM, Leblond CS, Boeckers TM, Faure P, Bourgeron T. 2013 The autism ProSAP1/Shank2 mouse model displays quantitative and structural abnormalities in ultrasonic vocalisations. *Behav Brain Res.* **256**, 677–689. (doi:10.1016/j.bbr.2013.08.031)

21. Ferhat A-T, Le Sourd A-M, de Chaumont F, Olivo-Marin J-C, Bourgeron T, Ey E. 2015 Social communication in mice–are there optimal cage conditions? *PloS One* **10**, e0121802. (doi:10.1371/journal.pone.0121802)

22. White NR, Prasad M, Barfield RJ, Nyby JG. 1998 40- and 70-kHz vocalizations of mice (Mus musculus) during copulation. *Physiol Behav* **63**, 467–473. (doi:10.1016/S0031-9384(97)00484-8)

23. Hammerschmidt K, Radyushkin K, Ehrenreich H, Fischer J. 2012 The structure and usage of female and male mouse ultrasonic vocalizations reveal only minor differences. *PLoS One* **7**, e41133. (doi:10.1371/journal.pone.0041133)

24. Neunuebel JP, Taylor AL, Arthur BJ, Egnor SR. 2015 Female mice ultrasonically interact with males during courtship displays. *eLife* **4**, 1–24. (doi:10.7554/eLife.06203)

25. Tachibana RO, Kanno K, Okabe S, Kobayasi KI, Okanoya K. 2020 USVSEG: A

robust method for segmentation of ultrasonic vocalizations in rodents. *PLoS ONE*

15, 1–16. (doi:10.1371/journal.pone.0228907)

26. Matsumoto YK, Okanoya K. 2018 Mice modulate ultrasonic calling bouts

according to sociosexual context. *R Soc Open Sci* **5**, 1–15.

(doi:10.1098/rsos.180378)

Legends for figures

Fig. 1. Male and female USVs to familiar or unfamiliar females. (A) Schematic

diagram of the experimental procedure. (B) Female USVs to the familiar or unfamiliar

females under the condition of 30 min isolation before recording 1. (C) Females' USVs

to the familiar or unfamiliar females under the condition of 1day isolation before

recording 1. (D) Male USVs to the familiar or unfamiliar females under the condition of

30 min isolation before recording 1. (E) Male USVs to familiar or unfamiliar females

under the condition of 1day isolation before recording 1. *$P < 0.05$, Wilcoxon

signed-rank test; B, n = 11; C, n = 14, D and E, n = 15.

Fig. 2. Sex differences in vocal response to familiar or unfamiliar females. Total

comparison among groups is shown in Fig. 1. *P < 0.05, Steel–Dwass test after

Kruskal–Wallis test; male 30 min and 1day isolation, n = 15; female 30 min isolation, n

= 11, female 1day isolation, n = 14.