Global epistasis emerges from a generic model of a complex trait

Gautam Reddy*

NSF-Simons Center for Mathematical & Statistical Analysis of Biology, Harvard University, Cambridge, MA 02138

Michael M. Desai†

NSF-Simons Center for Mathematical and Statistical Analysis of Biology, Harvard University, Cambridge, MA 02138
Department of Organismic and Evolutionary Biology,
Harvard University, Cambridge, MA 02138, USA
Quantitative Biology Initiative, Harvard University, Cambridge, MA 02138, USA and
Department of Physics, Harvard University, Cambridge, MA 02138, USA.
(Dated: June 14, 2020)

Epistasis between mutations can make adaptation contingent on evolutionary history. Yet despite widespread "microscopic" epistasis between the mutations involved, microbial evolution experiments show consistent patterns of fitness increases during laboratory adaptation. Recent work has found that this consistency is driven in part by global patterns of diminishing-returns and increasing-costs epistasis, which make mutations systematically less beneficial (or more deleterious) on more-fit genetic backgrounds. This global "macroscopic" epistasis is thought to make phenotypic evolution repeatable, even while the genetic basis of this evolution is highly stochastic. However, the mechanistic basis of consistent macroscopic epistasis remains unknown. Here we show, using a generic model of a complex trait, that global diminishing-returns and increasing-costs epistasis arise naturally as a consequence of pervasive microscopic epistasis, emerging simply as a statistical trend due to an imbalance between positive and negative contributions to the fitness. Our model predicts a specific quantitative relationship between the magnitude of global epistasis and the stochastic effects of microscopic epistasis, which we confirm by re-analyzing existing data. We also describe the predictions of this model for the patterns of fitness evolution and for how the distribution of fitness effects shifts as a population adapts, and propose additional experimental tests of these results.

I. INTRODUCTION

Despite the idiosyncrasies of epistasis, a number of laboratory microbial evolution experiments show systematic patterns of convergent phenotypic evolution and declining adaptability. A striking example is provided by the *E.coli* long-term evolution experiment (LTEE) (Figure 1a): 12 replicate populations that adapt in parallel show remarkably similar trajectories of fitness increase over time [1, 2], despite stochasticity in the identity of fixed mutations and the underlying dynamics of molecular evolution [3, 4]. Similar consistent patterns of fitness evolution characterized by declining adaptability over time have also been observed in parallel yeast populations evolved from different genetic backgrounds and initial fitnesses [5] (Figure 1b) and in other organisms [6–12]. Declining adaptability is thought to arise from diminishing-returns epistasis [5, 13, 14], where a global coupling induced by epistatic interactions systematically reduces the effect size of individual beneficial mutations on fitter backgrounds. Diminishing-returns manifests as a striking linear dependence of the fitness effect of a mutation on background fitness (Figure 1c). While diminishing-returns can be rationalized as the saturation of a trait close to a fitness peak, recent work shows a similar dependence on background fitness even for dele-

terious mutations, which become more costly on higher fitness backgrounds [15]. This suggests that fitter backgrounds are also less robust to deleterious effects (Figure 1d), a phenomenon that has been termed increasing-costs epistasis. The mechanistic basis for the global coupling that results in these effects is unknown.

Put together, these observations suggest that the contributions to the fitness effect, s_i, of a mutation at a locus i in a given genetic background can be written as

$$s_i = s_{\text{additive},i} + s_{\text{genotype},i} - c_i y, \qquad (1)$$

where $s_{\text{additive},i}$ is the additive effect of the mutation, $s_{\text{genotype},i}$ is its genotype-dependent epistatic contribution independent of the background fitness y (i.e. idiosyncratic epistasis), and c_i quantifies the magnitude of global epistasis for locus i. Eq. (1) reflects the observation that the strength of global epistasis depends on the specific mutation and applies independently of whether its additive effect is deleterious (increasing-cost) or beneficial (diminishing-returns). Over the course of adaptation in a fixed environment, global epistatic feedback on mutational effects can lead to a long-term decrease in adaptability. If this feedback dominates, Eq. (1) suggests that the dependence of the fitness effect on evolutionary history is summarized entirely by the current fitness, and therefore results in predictable fitness evolution.

———————
*gautam_nallamala@fas.harvard.edu
†mdesai@oeb.harvard.edu

FIG. 1: Declining adaptability and global epistasis in microbial evolution experiments. (a) Convergent phenotypic evolution in the *E.coli* long-term evolution experiment: the fitness relative to the common ancestor of 11 independently adapting populations over 50,000 generations is shown (data from [1]). The 12th population, Ara+6, has limited data and is not shown. (b) Yeast strains with lower initial fitness adapt faster (data from [5]). The fitness gain after 250 (green) and 500 (orange) generations of 640 independently adapting populations with 64 different founders and 10 replicates of each founder. Mean and SE are computed over replicates. (c) Diminishing returns of specific beneficial mutations on fitter backgrounds for three knocked out genes (green, orange and purple) (data from [5]). Control in pink. (d) Increasing costs of specific deleterious mutations on fitter backgrounds (data from [15]). The fitness effect relative to the least fit background for the mean over 91 mutations (in red) and five of the 91 mutations are shown. Linear fits for the five specific mutations and the mean using dashed and solid lines respectively are shown.

II. RESULTS

A. Diminishing-returns and increasing-costs epistasis

We begin by examining the most general form for expressing the relationship between genotype and fitness (i.e. to describe the fitness landscape). Consider first the case of a single locus, x_1, that has two possible variants, which we denote as $x_1 = \pm 1$. We note that this convention is non-standard in much of the population and quantitative genetics literature (which would instead often take $x_1 = 0$ and $x_1 = 1$ as the two possible genotypes), but this is an equivalent formulation [16–18] which we employ here because it will prove more convenient for our purposes. Given this formulation, we can write the fitness, y, of the two possible genotypes as

$$y = \bar{y} + f_1 x_1, \qquad (2)$$

where $\bar{y}$ is a constant that sets the overall scale of fitness. A mutation at the locus is simply a flip of sign, $x_1 \to -x_1$,

so the fitness effect of the mutation at this locus (from, say, $x_1 = -1$ to $x_1 = 1$) is given by $s = (\bar{y} + f_1) - (\bar{y} - f_1) = 2f_1$.

If there are two loci that contribute to fitness, we can extend Eq. (2) to this case as:

$$y = \bar{y} + f_1 x_1 + f_2 x_2 + f_{12} x_1 x_2, \qquad (3)$$

where again $x_1, x_2 = \pm 1$. It is straightforward to compute the fitness effect of each mutation and the epistatic interaction between them from this landscape (taking $(x_1, x_2) = (-1, -1)$ as the wild-type, we have $s_1 = 2f_1 - 2f_{12}$, $s_2 = 2f_2 - 2f_{12}$, and $\epsilon \equiv y_{1,1} - y_{-1,-1} - s_1 - s_2 = 4f_{12}$). The advantage of our formulation of the landscape is that the additive and epistatic contributions to the overall genetic variance are clearly delineated. In particular, if in a population, the two loci are unlinked (i.e., uncorrelated) and variants are equiprobable, then the mean fitness is simply $\bar{y}$ and the genetic variance is $f_1^2 + f_2^2 + f_{12}^2$. Therefore, f_1 and f_2 correspond to the additive contributions to the variance and f_{12} quantifies the epistatic contribution. If we assume fitness changes from mutations are due to the modification of parameters such as rate constants within a cellular network, then f_1, f_2 and f_{12} are related to the first and second-order terms respectively in a Taylor expansion of this network's contribution to the fitness with respect to the modified parameters.

Generalizing Eq. (3) further to an arbitrary number of loci, a map between a quantitative trait (such as fitness), y, and the underlying genotype can be expressed as a sum of combinations of L biallelic loci $x_1, x_2, \ldots, x_L$ that take on values $x_i = \pm 1$:

$$y = \bar{y} + \sum_i f_i x_i + \sum_{i>j} f_{ij} x_i x_j + \sum_{i>j>k} f_{ijk} x_i x_j x_k + \ldots \qquad (4)$$

We emphasize that this is a completely general framework; any fitness landscape can be represented in the form described in Eq. (4). As before, the coefficients of terms linear in x_i represent the additive contributions of the contributing loci to the fitness, whereas the higher-order terms quantify epistatic interactions of all orders, and $\bar{y}$ is the average fitness across all possible genotypes. Importantly, Eq. (4) makes apparent the idiosyncrasies induced by epistasis: a mutation at a locus with l interacting partners has an effect composed of 2^{l-1} independent contributions. To explicitly compute the fitness effect of a mutation at locus i in a particular genetic background, we simply flip the sign of x_i, keeping all other x_j constant, and write down the difference in fitness that results. This fitness effect will generically involve a sum over a large number of terms involving the f's in Eq. (4). While this may suggest that an analysis of fitness effects via Eq. (4) is impossible, it in fact lends itself to a drastic simplification: we generically expect the fitness effects of individual mutations to decrease linearly with background fitness if the locus has a significant number

of independent interactions that contribute to the fitness (i.e., provided that the number of nonzero epistatic terms is large). In other words, Eq. (4) implies that widespread independent idiosyncratic epistatic interactions lead to global diminishing-returns and increasing-costs epistasis.

We present a full derivation of this result in the Appendix. Here we explain the key intuition using a heuristic argument. The argument is based on a simple idea: for a well-adapted organism (y large) with complex epistatic interactions, a mutation is more likely to disrupt rather than enhance fitness. To be quantitative, consider a highly simplified scenario where some number N of the f's in Eq. (4) are ± 1 at random, and the others are 0. In this case, the fitness of a given genotype is a sum of N_+ and N_- interactions that contribute positively and negatively to the trait respectively, each with unit magnitude, so that $y = N_+ - N_-$. When positive and negative interactions balance, the organism is in a "neutrally-adapted" state ($y \approx \bar{y}$). By selecting for positive interactions, adaptation generates a bias so that $N_+ > N_-$ and $y > \bar{y}$. If locus i involved in a fraction v_i of all of $N = N_+ + N_-$ interactions is mutated, the effect of the mutation, on average, is to flip the sign of $N_+ v_i$ positive interactions and $N_- v_i$ negative interactions. The new fitness is then $y_i = N_+ - N_- - 2N_+ v_i + 2N_- v_i = (1 - 2v_i)y$. The linear negative feedback relation is immediately apparent and emerges as a systematic trend simply due to a sampling bias towards positive interactions. Of course, while this relation is true on average, it is possible that locus i affects more or less positive interactions due to sampling fluctuations. Provided only that N is large, these fluctuations are approximately Gaussian with magnitude $[Nv_i(1 - v_i)]^{1/2}$.

This basic argument holds beyond the simple model with unit interactions. In the more general case, we find that the v_i described above is replaced by the expression

$$v_i \equiv \frac{V_i}{V} = \frac{f_i^2 + \sum_{j \neq i} f_{ij}^2 + \cdots}{\sum_k f_k^2 + \sum_{k>l} f_{kl}^2 + \cdots}. \tag{5}$$

Note that V corresponds to the total genetic variance due to all loci (i.e. the variance in fitness across all possible genotypes) while V_i corresponds to the variance due to locus i (i.e. the amount by which the total genetic variance would decrease if we fixed the allele at locus i). We therefore refer to v_i as the *variance fraction* of locus i, since it quantifies the magnitude of contributions to the variance in the trait from locus i relative to the contributions from all loci. However, we emphasize that the v_i's do not sum to one unless there is no epistasis (with epistasis, typically $\sum_i v_i > 1$, reflecting the fact that the variance contributed by different loci overlap).

In the Appendix, we use a generalization of the heuristic argument above to show that, given this definition of v_i, the fitness effect of a mutation at locus i on a background of fitness y can be written as

$$s_i = -2v_i(y - \bar{y}) + 2\sqrt{v_i(1 - v_i)}\eta_i. \tag{6}$$

Here the first term corresponds to the global pattern of diminishing returns, while the second corresponds to idiosyncratic fluctuations away from this trend (where η_i is a background genotype and locus dependent stochastic term which is normally distributed with zero mean and a variance equal to the genetic variance over the 2^L equally likely background genotypes; see SI Appendix for further comments). We note that Eq. (6) implies that v_i plays a central role in our theory. It determines the slope of the negative relationship between fitness effect and background fitness (i.e. the strength of global epistasis for locus i); this corresponds to the first term in Eq. (6). At the same time, it determines the size of the idiosyncratic fluctuations away from this trend; this corresponds to the second term in Eq. (6).

We emphasize that Eq. (6) does not depend on the specific details of the genetic architecture or the fitness landscape and is true as long as there are sufficient non-zero and independent epistatic terms contributing to the fitness such that the central limit theorem holds. Since the total number of possible epistatic terms grows exponentially with the number of loci, it is therefore natural to expect Eq. (6) to apply generically for a complex trait. From a phenomenological perspective, Eq. (6) is sufficient to describe the core observations of global diminishing-returns and increasing-costs epistasis, with a linear dependence of fitness effect on background fitness and a locus-specific dependence of the magnitude of this global epistasis.

Essentially, our analysis captures what the fitness effect of a mutation looks like when regressed against parental fitness, with the dependence of the fitness effect on background genotype (i.e., the 'microscopic' epistasis term) appearing as the residuals from the regression. Eq. (6) has important implications which immediately lead to two testable predictions. First, the *variance* of the residuals is proportional to the *slope* of the regression. In other words, the magnitude of microscopic epistasis of a locus is related to its global coupling to the trait, Second, η_i is normally distributed and thus the full distribution of the residuals can be predicted from the slope. We also note that the slope can be used to probe the contribution of a locus to the trait (i.e. its variance fraction), taking into account all orders of epistasis.

We note that in a purely additive model with no epistasis, we find from Eq. (5) that the sum of variances of loci should add up to the total genetic variance. Each variance fraction is thus $O(L^{-1})$. Thus in this additive case, declining adaptability does not involve diminishing-returns, but instead occurs only as an effect of running out of a finite supply of available mutations. However, this is no longer true when epistasis comes into play. While strong microscopic epistasis is not strictly required for diminishing-returns to apply, epistatic interactions allow for a larger number of significant loci and a wide range of variance fractions across loci. Importantly, while fitness effects of individual mutations (and hence the distribution of fitness effects) may change over the course

FIG. 2: Global epistasis is recapitulated in a generic model of a complex trait and leads to testable predictions. (a) The distribution of variance fractions (DVF) over 400 loci for the simulated genotype-phenotype map. (b) The predicted linear relationship between fitness effect (relative to the fitness effect on the least fit background) and background fitness for the mean over 30 randomly chosen loci (red, solid line) and five loci (dashed lines in colors) is recapitulated. The slope of the linear fit for each locus is proportional to its variance fraction, v (slope $= -2v$). Mean and SE are over backgrounds of approximately equal fitness. See Methods for more details. (c) The mean fitness gain after 25 (green) and 50 (orange) generations of simulated evolution of 768 independently adapting populations with 64 unique founders and 12 replicates each. Means and SEs are computed over the 12 replicates. Error bars are s.e.m. (d) The relationship predicted from theory between the residual variance from the linear fit for each locus and its slope is confirmed in simulations. (e) The mean fitness effect for single mutants at 30 loci and double mutants from all possible pairs of the 30 loci. The slope for the double mutants is predicted to be roughly twice that of single mutants. (f) The estimated variance fraction of a double mutant with mutations at two loci is predicted from theory and confirmed in simulations to be approximately the sum of the variance fractions for single mutations at the two loci. Sub-additivity is due to epistasis between the two loci. See Methods for more details.

of evolution due to epistasis, the distribution of variance fractions (DVF), $P(v)$, is an invariant measure of the effect-sizes of mutations available to the organism during adaptation. As we will see, this means that the DVF plays an important role in determining long-term adaptability.

B. Numerical results and experimental tests

To test our analytical results, we first demonstrate that the effects described are reproduced in numerical simulations and then analyze existing experimental data. We numerically generated a genotype-phenotype map of the form in Eq. (4) with $L = 400$ loci and an exponential DVF, $P(v) = \bar{v}^{-1}e^{-v/\bar{v}}$, where $\bar{v} = 0.02$ (Methods). This DVF is shown in Figure 2a. The fitness effect of mutations at 30 loci was measured for 640 genotypes with a range of fitnesses (Figure 2b), which recapitulated the linear dependence on background fitness (Figure 1c,d) and a negative slope equal to twice the variance fraction predicted from Eq. (6). We further simulated the evolution of randomly generated genotypes similar to the experimental procedure used in Kryazhimskiy et al. [5] (Figure 2c), which reproduced the patterns of declining adaptability observed in experiments (Figure 1b). Note

that ~10 mutations are fixed during this simulated evolution; declining adaptability here is not due to a finite-sites effect.

As mentioned previously, a non-trivial prediction from Eq. (6) is the proportional relationship between the magnitude of global epistasis (quantified by the negative slope) and the magnitude of microscopic epistasis (quantified by the residual variance); see also Figure 3a. We verify this relationship in simulations (Figure 2d). The theory additionally predicts that the negative slope of a double mutant at loci i, j is $v_{ij} = v_i + v_j - 2e_{ij}$, where e_{ij} quantifies the magnitude of epistatic interactions between i and j. We further assume that $e_{ij} = O(\bar{v}^2)$ (specifically, $e_{ij} = v_i v_j$ for the genotype-phenotype map used for numerics). Since v_i and v_j are typically small for a complex trait, we expect near-additivity $v_{ij} \approx v_i + v_j$ and that any deviations are sub-additive, which is confirmed in simulations (Figure 2e,f).

While testing the latter prediction on double mutants requires further experiments, the proportional relationship between slope and residual variance can be immediately tested. To do so, we re-analyzed the data from Johnson et al. [15], which measured the fitness effect of 91 insertion mutants on ~145 backgrounds. The estimated variance fraction of each of the 91 mutations is shown in Figure 3b. These mutations were selected after

FIG. 3: Experimental observations from Johnson *et al.* [15] are consistent with theoretical predictions. (a) The fitness effect of one of the 91 mutations from [15] plotted against background fitness. The quantities relevant for the theoretical prediction are shown. (b) The distribution of the measured variance fractions for the 91 insertion mutations. (c) The measured variance of the residuals (obtained from subtracting the background-fitness-dependent linear terms from the fitness effect) is proportional to the measured variance fraction, v (specifically $v(1-v)$), shown here for the 91 mutations. (d) The shape of the distribution of residuals pooled from all 91 mutations aligns well with the prediction from Eq. (6). The variances of the two distributions are matched. Inset: same plot in log-linear scale. See Methods for more details.

screening for nonzero effect, and thus the DVF is biased upwards. The mean variance fraction is $\bar{v} \approx 0.06$. The data shows a remarkable confirmation of the relationship between variance fraction and residual variance (Figure 3c, $R^2 = 0.8$). The Gaussian-distributed term in Eq. (6) also predicts the shape of the distribution of the residuals given the variance fractions, which aligns well with the empirical distribution of the residuals (Figure 3d).

In short, we have a developed a theoretical framework to show that linear patterns of global diminishing-returns and increasing-costs epistasis are a simple consequence of widespread microscopic epistasis combined with an imbalance between positive and negative contributions to the fitness. In what follows, we extend this basic argument to analyze how global epistasis influences the distribution of fitness effects and consequently long-term adaptation.

C. The distribution of fitness effects

Long-term adaptation is determined by the distribution of fitness effects (DFE) over possible mutations and the stochastic dynamical processes that lead to fixation.

While Eq. (6) represents the distribution of the fitness effects of a specific mutation at locus i over *all* genotypes in the population that have fitness y, we are now instead interested in the DFE, where fitness effects are measured for all the mutations of a *particular* genotype that has fitness y. For now we ignore the influence of evolutionary history on the DFE; we expand on that complication in the following Section.

Examining the DFE over L loci for a randomly chosen genotype of fitness y can be thought of as sampling the fitness effects $s_1, s_2, \ldots, s_L$ from the conditional joint distribution $P(s_1, s_2, \ldots, s_L | y)$, which generally depends on epistasis. If the number of nonzero epistatic terms is large, $P(s_1, s_2, \ldots, s_L | y)$ is a multivariate normal distribution defined entirely by the means and covariances of the $L+1$ variables $y, s_1, s_2, \ldots, s_L$, which in turn can be computed in terms of the f's from Eq. (4). In particular, the conditional means and covariances are $\text{Mean}_y(s_i) = -2v_i(y - \bar{y})$, $\text{Cov}_y(s_i, s_j) = 4V(e_{ij} - v_i v_j)$, where e_{ij} is the epistatic variance fraction between loci i and j and $e_{ii} = v_i$. This implies that the conditional correlation between fitness effects is $(e_{ij} - v_i v_j)/\sqrt{v_i v_j (1 - v_i)(1 - v_j)}$.

The DFE simplifies considerably if we make certain additional assumptions on the magnitude of epistatic interactions. If we assume the typical variance fraction $\bar{v}$ is small (i.e., $\bar{v} \ll 1$) and also that e_{ij} is $O(\bar{v}^2)$, then correlations are $O(\bar{v})$ and thus negligible. Then, in a particular sample $s_1, s_2, \ldots, s_L$, we can think of each s_i as being drawn independently with mean $-2v_i(y - \bar{y})$ and variance $4v_i V$. If L is large, this leads to a DFE

$$\rho(s|y) = \int_0^1 dv (2\sqrt{vV})^{-1} P(v) \varphi\left(\frac{s + 2v(y - \bar{y})}{2\sqrt{vV}}\right), \quad (7)$$

where $P(v)$ is the DVF across loci and φ is the standard normal pdf. Curiously, the correlations between s_i's vanish when $e_{ij} = v_i v_j$ and the DFE is determined entirely by the DVF. Further below, we introduce a fitness landscape model for which this relation does hold. Diminishing-returns is naturally incorporated in Eq. (7): the mean of s is $-2\bar{v}(y - \bar{y})$, i.e., the DFE shifts progressively towards deleterious values with increasing fitness.

D. Historical contingency in adaptive trajectories

A key unresolved question is the extent to which evolutionary history influences the DFE and the dynamics of adaptation [19]. That is, what does our theory say about historical contingency?

Suppose a clonal population of fitness y_0 accumulates k successive mutations resulting in fitnesses $y_1, y_2, \ldots, y_k$. By virtue of arising on the same ancestral background, the fitness gain of a new mutation, s_{k+1}, is in general correlated with the full sequence of past fitnesses and the identity of the k mutations through its epistatic interactions with them. Based on these correlations, we

can use linear response theory to write

$$s_{k+1} = \sum_{i=0}^{k} w_{k+1,i} y_i + \epsilon, \qquad (8)$$

where the weights $w_{k+1,i}$ depend on the variance fraction (v_{k+1}) of the new mutation and its epistatic interactions with past mutations. Here ϵ is the stochastic residual that depends on the initial genotype and the weights (SI Appendix). Eq. (8) is a generalization to a sequence of mutations of Eq. (6), i.e., when $k = 0$.

To gain intuition, it is useful to first analyze Eq. (8) when $k = 1$ (i.e. to compute the effect of a second mutation conditional on the first). In this case, we show that (SI Appendix)

$$s_2 \simeq -2v_2(y_1 - \bar{y}) + \frac{v_1 v_2 - e_{12}}{v_1} s_1 + \epsilon, \qquad (9)$$

where $s_1 = y_1 - y_0$ is the fitness effect due to mutation 1. The first term on the right hand side is the dependence on the fitness of the immediate ancestor, similar to the corresponding term in Eq. (6). The second term quantifies the influence of epistasis between loci 1 and 2 on s_2. When $e_{12} = v_1 v_2$, dependence on s_1 vanishes entirely and s_2 depends only on y_1. In contrast, if loci 1 and 2 do not interact, $e_{12} = 0$, and s_2 is, on average, larger *if* the mutation at 1 is beneficial compared to when it is deleterious. This has an intuitive interpretation: diminishing-returns applies to the overall fitness and the mechanism through which it acts is epistasis. However, if mutations 1 and 2 do not interact, then the increase in fitness corresponding to mutation 1 does not actually reduce the effect of mutation 2 (as expected by diminishing-returns) so the expected effect of mutation 2 is larger. This analysis suggests that during adaptation, since selection favors mutations with stronger fitness effects on the current background, a mutation that interacts less with previous mutations is more likely to be selected.

To identify the conditions under which history plays a minimal role, we would like to examine when s_{k+1} depends only on the current fitness, y_k, and is independent of both the past fitnesses and idiosyncratic epistasis. If this were true, then Eq. (6) would apply for new mutations that arise through the course of a single evolutionary path (i.e. the fitness effect of a new mutation is "memoryless" and depends only on its variance fraction and the current fitness). Surprisingly, such a condition does exist. We show that this occurs when the magnitude of epistatic interactions between the new mutation and the k previous mutations, $e_{k+1,1:k}$, satisfies a specific relation: $e_{k+1,1:k} = v_{k+1} v_{1:k}$, where $v_{1:k}$ is the combined variance fraction of the k previous mutations (SI Appendix). In general, this condition is not satisfied, implying that there will be historical contingency which can be analyzed using the framework above. Remarkably, it turns out that a fitness landscape model for which the condition is satisfied does exist and arises from certain

intuitive assumptions on the organization of biological pathways and cellular processes. This fitness landscape model is described below.

E. The connectedness model

We introduce the "connectedness" model (CN model, for short). In this model, each locus i is involved in a fraction μ_i of independent "pathways", where each pathway has epistatic interactions between all loci involved in that pathway (Figure 4a). The probability of an epistatic interaction between three loci (i, j, k) is then proportional to $\mu_i \mu_j \mu_k$, since this is the probability that these loci are involved in the same pathway. When the number of loci L is large, we show that in this model, $v_i = \mu_i/(1 + \bar{\mu})$, and when L is small, we get $v_i = \mu_i/\bar{\mu}L$, where $\bar{\mu}$ is the average over all loci (SI Appendix). The CN model therefore has a specific interpretation: the outsized contribution to the fitness from certain loci (large v_i) is due to their involvement in many different complex pathways (large μ_i) and not from an unusually large perturbative effect on a few pathways. The distribution, $P(\mu)$, across loci determines the DVF.

Statistical fitness landscapes such as the NK model and the Rough Mt. Fuji model [19–23] are related to the CN model, with one important difference that distinguishes them and gives qualitatively different dynamics of adaptation (shown further below). Specifically, in contrast to the CN model, the classical fitness landscape picture corresponds to a mean-field theory where the variance fraction of every locus is assumed to be the same. In the special case of all v_i being equal, the 'homogeneous' CN model is equivalent to a Gaussian fitness landscape with exponentially-decaying correlations (SI Appendix). The homogeneous CN model has tunable ruggedness, i.e., the landscape transitions from additivity to maximal epistasis with increasing $\bar{\mu}$. Maximal epistasis corresponds to $\mu_i = 1$ (and hence $v_i = 1/2$) for all loci. From Eq. (6), this implies that the new fitness after a mutation occurs is independent of the previous fitness, consistent with the expectation from a House-of-Cards model.

Classical fitness landscape models with exponentially-decaying correlations have memoryless fitness effects under the restrictive assumption that every locus is equivalent [19]. We show that the dynamics of adaptation of the more general CN model are also memoryless, i.e., the condition detailed in the previous section holds true (SI Appendix). Yet, as we show below, the predicted dynamics for the CN model are very different to those from a classical fitness landscape model.

We emphasize that the well-connectedness assumed for the CN model is not a requirement for Eq. (6) to hold. However, how diminishing-returns influences the long-term dynamics of adaptation depends on the specific genetic architecture and the corresponding fitness landscape. Consider for example an alternative model of genetic networks organized in a modular structure

FIG. 4: The DFE and long-term adaptation dynamics predicted for the connectedness model. (a) Schematic of the connectedness (CN) model, where each locus is associated with a fraction μ of pathways that contribute to the organism's fitness. (b) An alternative model with modular organization, where sets of loci interact only within the pathways specific to a single module. (c) The DFE predicted from Eq. (12) matches those obtained from simulated evolution of genotypes from the CN model. 128 randomly drawn genotypes (400 loci) with initial fitness y close to zero are evolved to $y = 2.5$ and $y = 5$ and the DFE is measured across loci and genotypes. We chose $\bar{y} = 0$ and $V = 1$ so that y represents adaptedness. Insets: same plots in log-linear scale. Note that the number of beneficial mutations acquired during the simulated evolution ($\sim$10-20) is much less than the total number of loci (400). (d) For a neutrally-adapted organism, the theory predicts quick adaptation to a well-adapted state beyond which the adaptation dynamics are independent of the specific details of the genotype-fitness map. Shown here is the mean adaptation curve predicted under strong-selection-weak-mutation (SSWM) assumptions, which leads to a power-law growth of fitness with exponent 1/5 in the well-adapted regime (inset). (e) The number of fixed beneficial mutations under SSWM, which grows as a power-law with exponent 2/5 in the well-adapted regime (inset). The shaded region is the 95% confidence interval around the mean for (c) and (d). See Methods and SI Appendix for more details.

(Figure 4b). In this model, each locus is part of a single module, and interacts epistatically with other loci in that module to determine the fitness of that module; overall fitness is then determined as a function of the module fitnesses. In this case, the variance contributed by a locus is due to its additive contribution and from epistasis between loci restricted to its module. While the argument for diminishing-returns still applies to the fitness as a whole, it follows from the same argument that diminishing-returns should also apply to each module separately. Consequently, the dynamics of adaptation for the modular model are different from the CN model. For simplicity, we analyze the dynamics of adaptation for the CN model and postpone a discussion of how the dynamics differ for different models to subsequent work.

F. The dynamics of adaptation

We now examine the DFE that follows from Eq. (6) and what that implies for long-term adaptation under the conditions for memoryless fitness effects. We henceforth assume a large number of loci with sparse epistasis (though the total number of nonzero epistatic terms is still large). This implies that $L \gg 1$, $v_i \ll 1$ and $\bar{v}L \gg 1$; we also assume strong-selection-weak-mutation (SSWM) selection dynamics (i.e., a mutation sweeps and fixes in

a population before another one arises). It is convenient to rescale fitnesses based on the total variance in fitness across all possible genotypes by defining $z = V^{-1/2}(y - \bar{y})$, $\sigma = V^{-1/2}s$, $\nu = V^{-1/2}\eta$. Note that ν is normally-distributed with zero mean and unit variance. Here z has an intuitive interpretation as the "adaptedness" of the organism. When the organism is neutrally-adapted ($|z| \ll 1$), positive and negative epistatic contributions to the fitness are balanced and global diminishing-returns is negligible. Global diminishing-returns is relevant when the organism is well-adapted ($z \gg 1$). Below, we give the intuition behind our analysis, which is presented in full detail in the SI.

In the neutrally-adapted regime, the linear negative feedback in Eq. (6) is negligible and the DFE is determined by the distribution of $\simeq v^{1/2}\nu$. Loci with large v can lead to a DFE with a long tail. In this regime, beneficial mutations fix linearly with time. If $\bar{v}$ is the typical variance fraction of a locus, the fitness increases linearly with time as $z \sim n_s \bar{v}^{1/2}$, where n_s is the number of substitutions. Since $\bar{v}$ is a measure of overall epistasis, this implies that epistasis speeds adaptation in the neutrally-adapted regime by allowing access to more influential beneficial mutations.

Fitness increases rapidly until the effect of the negative feedback cannot be neglected. From Eq. (6), this happens when $\bar{v}z \sim \bar{v}^{1/2}\nu$ (i.e., when $z^2 \sim \bar{v}^{-1}$). Intuitively, fitness begins to plateau when its accumulated

benefit from substitutions is comparable to the scale of the total genetic variance ($n_s \bar{v} \sim 1$) and further improvements are due to rare positive fluctuations. In this well-adapted regime, diminishing-returns and increasing-costs epistasis strongly constrain the availability of beneficial mutations, whose effects can be quantified in this model: for a mutation to have a fitness effect σ, we require from Eq. (6) that $\nu \simeq \sigma/2v^{1/2} + v^{1/2}z$, which has probability $\sim e^{-\nu^2/2}$. Beneficial effects of large σ arise when ν has a large positive deviation. The most likely v that leads to a particular σ is when ν is smallest (i.e. at $v^* \simeq \sigma/2z$), in which case $\nu \simeq \sqrt{2\sigma z}$, yielding a tail probability $\sim e^{-\sigma z}$. Remarkably, the beneficial DFE in the well-adapted regime is quite generally an exponential distribution independent of the precise form of the DVF (unless it is singular). In particular, we show that for the DFE, $\rho(\sigma|z)$,

$$\frac{\rho(\sigma|z)}{\rho(-\sigma|z)} = e^{-\sigma z}, \tag{10}$$

which depends solely on the adaptedness of the organism. The exponential form arises because of the Gaussianity of ν, but the argument can be easily extended to ν with non-Gaussian tails. An exponential beneficial DFE has been previously proposed by Orr [24] but arises here due to a qualitatively different argument (see Discussion).

Under SSWM assumptions, from Eq. (10), the typical effect size of a fixed mutation is $\sigma_{\text{fix}} \sim z^{-1}$, which typically has a variance fraction,

$$v_{\text{fix}}^* \simeq \sigma_{\text{fix}}/2z \sim 1/2z^2. \tag{11}$$

The above relation makes precise the effects of increasing-costs epistasis on adaptation. As adaptation proceeds, the delicate balance of high fitness configurations constrains fixed beneficial mutations to have *moderate* variance fractions. A mutation of small variance fraction is likely to confer small benefit and is lost to genetic drift, while one with a large variance fraction is more likely to disrupt an established high fitness configuration.

This intuition is not captured in classical fitness landscape models, which assume statistically equivalent loci, i.e., $v_i = \bar{v}$ for all i and $P(v) = \delta(v - \bar{v})$ is singular. From Eq. (7), we see that this leads to a Gaussian DFE whose mean decreases linearly with increasing fitness, in contrast to the exponential DFE in our theory. The key difference is the lack of loci with intermediate effect, which drive adaptation in the well-adapted regime. As a consequence, the rate of beneficial mutations declines exponentially ($U_b \sim e^{-\bar{v}z^2/2}$) and the fitness thus sharply plateaus at $z \sim \bar{v}^{-1/2}$. In contrast, our theory predicts a much slower depletion of beneficial mutations, $U_b \sim z^{-2}$ (SI Appendix). The rate of adaptation is $dz/dt \sim U_b p_{\text{fix}} \sigma_{\text{fix}} \sim z^{-4}$ (since $p_{\text{fix}} \sim \sigma_{\text{fix}}$), which leads to a slow but steady power-law gain in fitness, $z \sim t^{1/5}$. The rate of fixation of beneficial mutations is $dn_s/dt \sim U_b p_{\text{fix}} \sim z^{-3} \sim t^{-3/5}$, which gives $n_s \sim t^{2/5}$.

We verify our analytical results using numerics. As before, we generated a genotype-phenotype map using the CN model with an exponential DVF, $P(v) = \bar{v}^{-1} e^{-v/\bar{v}}$ and $L = 400$ loci. The DFE in this case can be calculated exactly from Eq. (7):

$$\rho(\sigma|z) = \frac{\bar{v}^{-1}}{2\sqrt{2\bar{v}^{-1} + z^2}} e^{-\sigma z/2 - |\sigma|\sqrt{2\bar{v}^{-1}+z^2}/2}. \tag{12}$$

We simulated the evolution of randomly generated genotypes from $z = 0$ to $z = 2.5$ and $z = 5$ and the DFE across all loci was measured (we chose $\bar{y} = 0, V = 1$ so that $y = z, s = \sigma$). The theoretical prediction for the DFE, Eq. (12), closely aligns with the numerical results (Figure 4c).

Due to computational constraints, it is difficult to simulate evolution deep into the well-adapted regime. To compute the shape of adaptive trajectories and their variability, we instead simulated SSWM dynamics using the DFE directly from Eq. (12), beginning from a neutrally-adapted fitness ($z = 0$). Typical trajectories (Figure 4d) show rapid adaptation to the well-adapted regime beyond which the fitness grows slowly as $t^{1/5}$, as predicted from theory. The predictions for the number of fixed beneficial mutation are also re-capitulated (Figure 4e).

III. DISCUSSION

Recent empirical studies have observed consistent patterns of diminishing-returns and increasing-costs epistasis. Our model gives a simple mechanistic explanation for these observations. In particular, we showed that these patterns are generic consequences of widespread microscopic epistatic interactions. The key intuition underlying this result is that a random mutation typically has a larger disruptive effect on the delicate balance of microscopic epistasis that underpins a fitter background. Our model predicts a specific, quantitative relationship between the magnitudes of global epistasis (i.e., the negative slope of diminishing-returns and increasing-costs epistasis) and microscopic epistasis, which we confirmed using existing data (Figure 3).

Our model leads to other experimentally testable predictions. The most direct and accessible test of our theory is to measure the fitness for all possible combinations of mutations at $\sim$10-15 significant loci and compare (using (5)) the magnitude of global epistasis to the measured fitness coefficients (the f's). Additionally, we predict that the magnitude of global epistasis of a double mutant should be nearly the sum of magnitudes of the corresponding single mutants, and any deviations should be biased towards sub-additivity. Since the predictions involve measuring residual variance, experimental noise can be an important confounding factor.

We emphasize that our key result, Eq. (6), is a general statistical relation that holds irrespective of the specific genetic architecture and the corresponding fitness landscape. Weak epistasis with many loci is sufficient to observe noticeable patterns of global epistasis. However, the argument fails if the contribution of a locus is purely

additive or when epistasis is limited to one or a handful of other loci. In the latter case, we expect the fitness effect of a mutation to be dominated by the allelic states of its partner loci, and thus take on a few discrete values. A few examples from Johnson *et al.* [15] indeed exhibit this pattern, (e.g. cases where the fitness effect of a specific mutation depends primarily on the allelic state at a single other locus).

A surprising empirical observation is that the negative linear relationship between fitness effect and ancestral fitness characteristic of global epistasis has different slopes for different loci. Our model identifies the slope as twice the fraction of variance contributed by a locus to the trait. To explain the wide range of variance fractions observed in data, we developed the connectedness (CN) model, a framework to think about the organization of cellular processes that can lead to loci of widely varying variance fractions. In the CN model, loci have a large variance fraction (VF) due to their involvement in many different pathways rather than due to a large effect on a single pathway. The CN model can be viewed as a generalization of statistical fitness landscapes, where rather than each locus having the same VF, loci have a range of VF, quantified by the distribution of variance fractions (DVF). In the special case of each locus having the same VF, the CN model corresponds to a Gaussian fitness landscape with tunable ruggedness and exponentially-decaying correlations.

Extending our framework to incorporate adaptation, we showed that the distribution of fitness effects (DFE) depends only on the current fitness, rather than the entire evolutionary history, under the intuitive assumptions behind the CN model. The theory therefore gives a simple explanation for why phenotypic evolution can be predictable, even while the specific mutations that underlie this evolution are highly stochastic.

Our framework has an implicit notion of 'adaptedness' without referencing a Gaussian-shaped phenotypic optimum, often assumed in models of adaptation (e.g. Fisher's geometric model) [25–27]. Over the course of adaptation, the DFE shifts towards deleterious values, reflecting diminishing-returns, which naturally arises from our basic arguments. For a well-adapted organism, we show that the DFE for beneficial mutations takes on an exponential form, and leads to universal adaptive dynamics. While an exponential DFE for beneficial mutations has been proposed previously based on extreme value theory [24], our result arises due to an entirely different argument: the tail of the beneficial DFE is determined by loci of intermediate size whose disruptive effect due to increasing-costs is small, yet whose effect size is large enough not to be lost due to genetic drift. Our theory further predicts declining adaptability, with rapid adaptation in a neutrally-adapted regime followed by much slower increases in fitness, resulting in power-law adaptive trajectories when the organism is well-adapted. This is consistent with observations from the *E.coli* LTEE [1, 2]. Our model predicts a quicker decline in the number of substitutions ($n_s \sim t^{2/5}$) compared to the near linear trend observed in the LTEE data [4]; however, the dynamics of fixation in the LTEE deviate strongly from SSWM assumptions, which may explain the discrepancy.

IV. METHODS

The code and data to generate the figures are available at [28].

A. Simulations

We consider a quantitative trait model with L loci. Each locus is assigned a sparsity μ from $P(\mu)$, which is an exponential distribution with mean $\bar{\mu}$. Each of M independent pathways sample loci with each locus i having probability μ_i of being selected to a pathway. We choose $L = 400, \bar{\mu} = 0.02, M = 500$ so that $\bar{\mu}L = 8$ ensures signficant epistasis. All loci in a pathway interact with each other, where additive and higher-order coefficient terms of all orders were drawn independently from a standard normal distribution. The total fitness is the sum of contributions from the M pathways. We normalize the coefficients so that the sum of squares of all coefficients is 1, i.e., the total variance across genotypes is 1. The mean, $\bar{y}$ is close to zero from our sampling procedure. The above procedure is a simple and efficient way to generate epistatic terms to order ~ 20, beyond which the computational requirements are limited by the exponentially increasing demand. Note that the effects described in the paper were also observed with only pairwise and cubic epistatic terms.

The variance fractions shown in Figure 2a can be calculated numerically from the definition. From the theory, given our choice of $P(\mu)$, these should follow an exponential distribution with mean $\bar{v} \approx \bar{\mu}/(1 + \bar{\mu})$. There may be deviations because M is finite whereas the calculations assume $M \to \infty$. To generate Figure 2b, in order to get a range of background fitnesses, we first sample 128 random genotypes. These have fitnesses close to zero; in order to obtain a range of fitness values, we simulated the evolution of these 128 genotypes up to $y = 1, 2, 3, 4, 5$ under strong-selection-weak-mutation (SSWM) assumptions to get $128 \times 5 = 640$ genotypes at roughly five fitness values. The fitness effect of applying a mutation (i.e., flipping its sign) is measured for 30 randomly chosen loci (which are kept fixed) over each of the 640 genotypes. This is shown for five of the 30 and for the mean over the 30 loci in Figure 2b.

To generate Figure 2c, we sampled 64 random genotypes and 12 replicates of each. The evolution of these 768 genotypes was simulated for a total of 50 generations with a mutation rate of 1 per generation. The mean fitness gain over the 12 replicates is plotted for each of the 64 founders against their initial fitness.

To generate Figure 2d, the residuals are measured using the same procedure as for the experimental data analysis (see Sec. IV B) for the initial 128 genotypes at $y \approx 0$ and the 30 loci with the largest variance fraction.

Double mutants were created by mutating all pairs of the 30 randomly chosen loci on the 640 evolved genotypes. Their mean fitness effect was computed and plotted along with the mean fitness effect for single mutants, shown in Figure 2e. The variance fraction of the pair of loci for the double mutant was estimated as before and compared to the sum of the estimated variance fractions of the corresponding single mutants. This is shown in Figure 2f.

To generate the plots in Figure 4c, we simulated the evolution of 128 randomly sampled genotypes to $y = 2.5$ and $y = 5$. The fitness effect of 200 randomly sampled loci was measured and the distribution is plotted.

B. Analysis of the data from Johnson *et al.*

The data from Johnson *et al.* [15] consists of the fitness after the addition of 91 insertion mutations on each of 145 background genotypes. The fitness of a particular mutation at locus i can be modeled as

$$y_i = -c_i y + b_i + s_i(g), \tag{13}$$

where y_i, y are the mutant and background fitnesses respectively, c_i, b_i are constants for each locus and $s_i(g)$ depends on the background genotype g. According to our theory, each locus i has an associated variance fraction v_i and the mutant fitness can be written as

$$y_i - \bar{y} = (1 - 2v_i)(y - \bar{y}) + \underbrace{2\sqrt{v_i(1 - v_i)}\eta_i(g)}_{\text{residual}}, \tag{14}$$

where $\bar{y}$ is an unknown constant and $\eta_i(g)$ highlights the dependence of the fitness effect on the background genotype.

To estimate v_i, we first compute the Pearson correlation $\hat{\rho}_i = \text{Corr}(y_i, y)$ across backgrounds and then use $\hat{v}_i = (1 - \hat{\rho}_i)/2$ to estimate the variance fraction. The Pearson correlation gives a better estimate of $\hat{v}_i$ (on simulated data) compared to the slope from a linear fit while also ensuring that the estimates are positive. The residuals for each of the 145 genotypes for each of the 91 mutations are then estimated as

$$\text{Residual}_i(g) = (y_i - (1 - 2\hat{v}_i)y) - \overline{(y_i - (1 - 2\hat{v}_i)y)}, \tag{15}$$

where the overline represents an average over the 145 genotypes, which is used as an estimate of the constant term. In Figure 3b, we plot the distribution of estimated variance fractions. In Figure 3c, we compute the variance of the residuals across the 145 genotypes for each locus and plot it against the locus' estimated variance fraction. In Figure 3d, we plot the distribution of residuals over all genotypes and loci. The prediction is that the distribution of residuals is determined by $2\sqrt{v_i(1 - v_i)}\eta$, where η is a Gaussian random variable. We multiply $\sqrt{\hat{v}_i(1 - \hat{v}_i)}$ for each locus with 10,000 i.i.d standard normal RVs, pool the resulting numbers for all loci and plot the predicted distribution in Figure 3d. The distributions are variance-matched. While Figure 3c shows that the variance of the residuals aligns with the theoretical prediction of being proportional to slope, Figure 3d shows that the data is also consistent with the predicted Gaussianity of the background-genotype-dependent contribution.

V. ACKNOWLEDGMENTS

We thank Sergey Kryazhimskiy, Andrew Murray, Milo Johnson, and members of the Desai lab for comments on the manuscript. G.R was supported by the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number #1764269) and the Harvard FAS Quantitative Biology Initiative. M.M.D. acknowledges support from the Simons Foundation (Grant 376196) and grant GM104239 from the NIH.

---

[1] Michael J Wiser, Noah Ribeck, and Richard E Lenski. Long-term dynamics of adaptation in asexual populations. *Science*, 342(6164):1364–1367, 2013.

[2] Richard E Lenski, Michael J Wiser, Noah Ribeck, Zachary D Blount, Joshua R Nahum, J Jeffrey Morris, Luis Zaman, Caroline B Turner, Brian D Wade, Rohan Maddamsetti, et al. Sustained fitness gains and variability in fitness trajectories in the long-term evolution experiment with escherichia coli. *Proceedings of the Royal Society B: Biological Sciences*, 282(1821):20152292, 2015.

[3] Olivier Tenaillon, Jeffrey E Barrick, Noah Ribeck, Daniel E Deatherage, Jeffrey L Blanchard, Aurko Dasgupta, Gabriel C Wu, Sebastien Wielgoss, Stephane Cruveiller, Claudine Médigue, et al. Tempo and mode of genome evolution in a 50,000-generation experiment. *Nature*, 536(7615):165–170, 2016.

[4] Benjamin H Good, Michael J McDonald, Jeffrey E Barrick, Richard E Lenski, and Michael M Desai. The dynamics of molecular evolution over 60,000 generations. *Nature*, 551(7678):45–50, 2017.

[5] Sergey Kryazhimskiy, Daniel P Rice, Elizabeth R Jerison, and Michael M Desai. Global epistasis makes adaptation predictable despite sequence-level stochasticity. *Science*, 344(6191):1519–1522, 2014.

[6] Santiago F Elena and Richard E Lenski. Evolution experiments with microorganisms: the dynamics and genetic bases of adaptation. *Nature Reviews Genetics*, 4(6):457–469, 2003.

[7] L Perfeito, A Sousa, T Bataillon, and I Gordo. Rates of fitness decline and rebound suggest pervasive epistasis. *Evolution*, 68(1):150–162, 2014.

[8] Andrea Wünsche, Duy M Dinh, Rebecca S Satterwhite, Carolina Diaz Arenas, Daniel M Stoebel, and Tim F Cooper. Diminishing-returns epistasis decreases adaptability along an evolutionary trajectory. *Nature Ecology & Evolution*, 1(4):1–6, 2017.

[9] Rafael Sanjuán, José M Cuevas, Andrés Moya, and Santiago F Elena. Epistasis and the adaptability of an rna virus. *Genetics*, 170(3):1001–1008, 2005.

[10] Alejandro Couce and Olivier A Tenaillon. The rule of declining adaptability in microbial evolution experiments. *Frontiers in genetics*, 6:99, 2015.

[11] Elizabeth R Jerison, Sergey Kryazhimskiy, James Kameron Mitchell, Joshua S Bloom, Leonid Kruglyak, and Michael M Desai. Genetic variation in adaptability and pleiotropy in budding yeast. *Elife*, 6:e27167, 2017.

[12] Martijn F Schenk, Ivan G Szendro, Merijn LM Salverda, Joachim Krug, and J Arjan GM De Visser. Patterns of epistasis between beneficial mutations in an antibiotic resistance gene. *Molecular biology and evolution*, 30(8):1779–1787, 2013.

[13] Aisha I Khan, Duy M Dinh, Dominique Schneider, Richard E Lenski, and Tim F Cooper. Negative epistasis between beneficial mutations in an evolving bacterial population. *Science*, 332(6034):1193–1196, 2011.

[14] Hsin-Hung Chou, Hsuan-Chao Chiu, Nigel F Delaney, Daniel Segrè, and Christopher J Marx. Diminishing returns epistasis among beneficial mutations decelerates adaptation. *Science*, 332(6034):1190–1192, 2011.

[15] Milo S Johnson, Alena Martsul, Sergey Kryazhimskiy, and Michael M Desai. Higher-fitness yeast genotypes are less robust to deleterious mutations. *Science*, 366(6464):490–493, 2019.

[16] Wim Hordijk and Peter F Stadler. Amplitude spectra of fitness landscapes. *Advances in Complex Systems*, 1(01):39–66, 1998.

[17] Richard A Neher and Boris I Shraiman. Statistical genetics and evolution of quantitative traits. *Reviews of Modern Physics*, 83(4):1283, 2011.

[18] Edward D Weinberger. Fourier and taylor series on fitness landscapes. *Biological cybernetics*, 65(5):321–330, 1991.

[19] Atish Agarwala and Daniel S Fisher. Adaptive walks on high-dimensional fitness landscapes and seascapes with distance-dependent statistics. *Theoretical population biology*, 130:13–49, 2019.

[20] Johannes Neidhart, Ivan G Szendro, and Joachim Krug. Exact results for amplitude spectra of fitness landscapes. *Journal of theoretical biology*, 332:218–227, 2013.

[21] Stuart A Kauffman and Edward D Weinberger. The nk model of rugged fitness landscapes and its application to maturation of the immune response. *Journal of theoretical biology*, 141(2):211–245, 1989.

[22] Peter F Stadler and Robert Happel. Random field models for fitness landscapes. *Journal of Mathematical Biology*, 38(5):435–478, 1999.

[23] Takuyo Aita, Hidefumi Uchiyama, Tetsuya Inaoka, Motowo Nakajima, Toshio Kokubo, and Yuzuru Husimi. Analysis of a local fitness landscape with a model of the rough mt. fuji-type landscape: Application to prolyl endopeptidase and thermolysin. *Biopolymers: Original Research on Biomolecules*, 54(1):64–79, 2000.

[24] H Allen Orr. The distribution of fitness effects among beneficial mutations. *Genetics*, 163(4):1519–1526, 2003.

[25] RA Fisher. *The genetical theory of natural selection.* Clarendon Press, 1930.

[26] H Allen Orr. The genetic theory of adaptation: a brief history. *Nature Reviews Genetics*, 6(2):119–127, 2005.

[27] Guillaume Martin and Thomas Lenormand. A general multivariate extension of fisher's geometrical model and the distribution of mutation fitness effects across species. *Evolution*, 60(5):893–907, 2006.

[28] Gautam Reddy. github:greddy992/global_epistasis, 2020.

Supplementary information for 'Global epistasis in a generic model of a complex trait'

Gautam Reddy and Michael M. Desai

June 14, 2020

1 Diminishing-returns and increasing-costs epistasis in a model of a complex trait

In the main text, we propose that the fitness effect after a mutation at locus i can be written as

$$s_i = -(1 - \rho_i)(y - \bar{y}) + \sqrt{1 - \rho_i^2}\,\eta_i, \quad |\rho_i| \le 1. \tag{1}$$

Here η_i is a genotype and locus dependent contribution, which is distributed as a mean-zero Gaussian with variance equal to the total genetic variance, $\bar{y}$ is a constant, and $1 - \rho_i$ can be viewed as the fraction of fitness "forgotten" due to the mutation.

We will now show that Eq. (1) arises naturally in a generic model of a trait with L biallelic loci. Essentially, we would like to compute the distribution of the fitness across genotypes after a mutation at locus i, y_i, given the fitness y and the set of all parameters Θ of the model (i.e. $P(y_i|y, \Theta)$, with $s_i = y_i - y$). Using the chain rule of probability, we can write

$$P(y_i|y, \Theta) = \sum_g P(y_i|g, \Theta)P(g|y, \Theta), \tag{2}$$

where the sum is over all possible genotypes. While $P(y_i|g, \Theta)$ is determined by the model, $P(g|y, \Theta)$ is the crucial factor that gives weight only to the genotypes that yield the current fitness y. If the fitness is much larger than the mean fitness over all possible genotypes, Eq. (2) implicitly ensures that weight is given to only those genotypic configurations that lead to such an unusually large fitness.

For simplicity, we will analyze the case of a "microcanonical ensemble" where every genotypic configuration that leads to a particular fitness is equally likely (with no linkage). This approximates the experimental setup from Johnson et al [1] against which our predictions will be tested. For a population with different statistics, certain terms below will differ but the qualitative aspects of the result still hold.

It is difficult to directly evaluate the sum in Eq. (2). Below, we give a simple derivation but elaborated so as to make clear the key assumption that leads to Eq. (1).

1.1 Fourier representation of the fitness function

The fitness $y(g)$ for a genotype of length L, $g = \{x_1, x_2, \ldots, x_L\}$, where $x_i = \pm 1$, can be generally written as

$$y(g) = \bar{y} + \sum_i f_i x_i + \sum_{i>j} f_{ij} x_i x_j + \sum_{i>j>k} f_{ijk} x_i x_j x_k + \ldots \tag{3}$$

The symmetric choice of $x_i = \pm 1$ is chosen for mathematical convenience. In this form, the total variance contribution from the qth order epistatic terms is $\sum_{i_1 > i_2 > \ldots i_q} f_{i_1 i_2 \ldots i_q}^2$ and the total genetic variance, defined as

$$V \equiv \frac{1}{2^L} \sum_g (y(g) - \bar{y})^2, \tag{4}$$

is the sum of variance contributions from orders $q = 1$ to L. The sum over the 2^L genotypes is an expectation value assuming all genotypes are equally likely; we will denote this expectation using an overline hereafter. We use this expectation value as a proxy for empirical averages over the "background" genotypes in a population. With a sufficient number of background genotypes, the empirical average should converge to this expectation value.

The representation in Eq. (3) is in fact a Fourier representation of the fitness function on the L-dimensional hypercube, and makes calculations much simpler. For instance, to get the terms from each order, we have

$$\bar{y} = \overline{y(g)}, \quad f_i = \overline{x_i y(g)}, \quad f_{ij} = \overline{x_i x_j y(g)}, \quad \ldots \tag{5}$$

It is useful to define the variance contribution due to a particular locus i as (symmetry w.r.t interchanging indices is used for each term)

$$V_i \equiv f_i^2 + \sum_{j \neq i} f_{ij}^2 + \sum_{k < j \neq i} f_{ijk}^2 + \ldots. \tag{6}$$

The *variance fractions* for each locus,

$$v_i \equiv \frac{V_i}{V}, \tag{7}$$

plays a key role in the model. In particular, we will show that $\rho_i = 1 - 2v_i$ in Eq. (1) when the mutation is the sign flip $x_i \to -x_i$.

1.2 Derivation of Eq. (1)

We would like to relate the fitness before and after locus i is flipped (i.e. $x_i \to -x_i$), denoted by y and y_i respectively. We have from Eq. (3),

$$y - \bar{y} = y_{\bar{i}} + \xi_i, \tag{8}$$
$$y_i - \bar{y} = y_{\bar{i}} - \xi_i, \tag{9}$$

where $y_{\bar{i}}$ and ξ_i respectively contain all the terms not containing and containing locus i, i.e.,

$$\xi_i = f_i x_i + \sum_{j \neq i} f_{ij} x_i x_j + \sum_{k < j \neq i} f_{ijk} x_i x_j x_k + \ldots. \tag{10}$$

We have $y_{\bar{i}} = (y + y_i)/2 - \bar{y}$ and $\xi_i = (y - y_i)/2$. Since $y_{\bar{i}}$ and ξ_i contain all the genotype-dependence, we can write

$$P(y_i | y, \Theta) \propto P(y_i, y | \Theta) \tag{11}$$

$$= \int P(y_i, y | y_{\bar{i}}, \xi_i) P(y_{\bar{i}}, \xi_i | \Theta) dy_{\bar{i}} \xi_i \tag{12}$$

$$= P\left(y_{\bar{i}} = \frac{y + y_i}{2} - \bar{y}, \xi_i = \frac{y - y_i}{2} \middle| \Theta\right). \tag{13}$$

From the properties of the Fourier representation in Eq. (5), it is easy to see that the means are $\overline{y_{\bar{i}}} = 0, \overline{\xi_i} = 0$, the variances are $\overline{y_{\bar{i}}^2} = V - V_i$, $\overline{\xi_i^2} = V_i$, and the covariance is $\overline{y_{\bar{i}}\xi_i} = 0$.

The calculations so far have been exact. We now make the key assumption that $y_{\bar{i}}$ and ξ_i are both normal-distributed across genotypes. This assumption is similar to that of Fisher's infinitesimal model, where the distribution of trait values across strains for a complex trait is argued to be normal-distributed since the trait value is due to infinitesimal independent contributions from many loci. While $y_{\bar{i}}$ is easily seen to be normal-distributed for large L, an argument can be made for ξ_i only if locus i has a large number of epistatic terms. However, even weak epistasis can lead to normality. For instance, for loci with only pairwise epistasis, the number of pairwise epistatic terms is $\propto L$, which justifies the normality assumption for large L.

Under these assumptions and since $y_{\bar{i}}$ and ξ_i are linearly independent, we have

$$P(y_i|y, \Theta) \propto \varphi\left(\frac{y + y_i - 2\bar{y}}{2\sqrt{V - V_i}}\right) \varphi\left(\frac{y - y_i}{2\sqrt{V_i}}\right), \tag{14}$$

where φ is the standard normal pdf. Therefore, y_i is normal-distributed across genotypes and from the above equation can be written as

$$y_i - \bar{y} = (1 - 2v_i)(y - \bar{y}) + 2\sqrt{v_i(1 - v_i)}\eta_i, \tag{15}$$

where the variance fraction $v_i = V_i/V$ was defined previously and $\overline{\eta_i} = 0, \overline{\eta_i^2} = V$. This leads to the form in Eq. (1),

$$s_i = -2v_i(y - \bar{y}) + 2\sqrt{v_i(1 - v_i)}\eta_i. \tag{16}$$

1.3   Comments on the result

The above derivation was presented to clarify the basic assumptions. Eq. (16) can also be derived by simply computing the correlation between y and y_i in Eq. (8) and Eq. (9). We then have $\overline{(y - \bar{y})(y_i - \bar{y})} = \overline{y_{\bar{i}}^2} - \overline{\xi_i^2} = V - 2V_i$. The correlation is then $1 - 2v_i$ and Eq. (16) follows if y_i is also Gaussian.

The expectation values are averages over all the genotypes assuming that every genotype of a particular fitness is equally likely. This is analogous to ensemble averages over a microcanonical ensemble in statistical physics, where one assumes that all the particle configurations that have a particular energy are equally likely. The validity of this assumption depends on the experiment and how the various strains are generated. For instance, in Johnson et al [1], the background genotypes are generated from a cross between two strains, which due to recombination makes each locus have equal probability of being one of the two alleles. Closely-linked loci may be considered together as blocks. However, some of the loci are partially linked, which may lead to deviations from the predictions.

In addition, the sums in Eq. (3) are over L loci involved in a trait. In reality, we may vary a subset P of all loci and take averages over only this subset. The derivation still follows through in this case; we can simply write the fitness in terms of effective parameters as

$$y(g) = \tilde{y} + \sum_{i \in P} \tilde{f}_i x_i + \sum_{i,j \in P} \tilde{f}_{ij} x_i x_j + \dots, \quad \text{where} \tag{17}$$

$$\tilde{y} = \bar{y} + \sum_{i \notin P} f_i x_i + \sum_{i,j \notin P} f_{ij} x_i x_j + \dots, \tag{18}$$

$$\tilde{f}_i = f_i + \sum_{j \notin P} f_{ij} x_j + \sum_{j,k \notin P} f_{ijk} x_j x_k + \dots, \quad \text{and so on.} \tag{19}$$

Here we are abusing notation — it is to be assumed that a coefficient with a particular combination of indices appears only once in the sums. The results from the previous section still apply w.r.t these

new effective parameters. For instance, the total variance and the variance due to locus $i \in P$ are

$$\tilde{V} = \sum_{i \in P} \tilde{f}_i^2 + \sum_{i,j \in P} \tilde{f}_{ij}^2 + \sum_{i,j,k \in P} \tilde{f}_{ijk}^2 + \dots, \tag{20}$$

$$\tilde{V}_i = \tilde{f}_i^2 + \sum_{j \in P} \tilde{f}_{ij}^2 + \sum_{j,k \in P} \tilde{f}_{ijk}^2 + \dots, \tag{21}$$

and the effective variance fraction is $\tilde{v}_i = \tilde{V}_i / \tilde{V}$.

Eq. (16) captures both diminishing-returns and increasing-costs epistasis. Increasing-costs epistasis in this model is due to the delicate stability of configurations that lead to very large fitness values. A mutation with strong epistasis and which has a significant contribution to the variance is likely to disrupt this delicate configuration and lead to a steep drop in fitness. Diminishing-returns is due to a lowering ratio of beneficial to deleterious mutations as adaptation proceeds and arises ultimately due to the conservation of genetic variance. This is implicit in the model since a mutation flips the sign $x \to -x$, which does not change the total genetic variance. Conservation of genetic variance results in Eq. (1) if in addition the randomness due to genetic factors is normal-distributed. This should be true if the loci of non-zero effect have epistatic interactions, since their effect is then a sum of many independent factors.

Eq. (16) is analogous to the fluctuation-dissipation theorem in statistical physics, which relates the response of a thermodynamic system to a perturbation. The relationship between the magnitude of macroscopic epistasis and the variance due to the background genotypes is analogous to the relationship between viscous drag and Brownian motion. The external perturbation in this case is provided by natural selection, which pushes the system away from equilibrium (here $\bar{y}$). Diminishing-returns naturally arises as the tendency of the system to revert to its entropically-favored equilibrium state.

Using the arguments from the previous section, it is easy to show that the variance of a double mutant at loci i and j, V_{ij}, is necessarily sub-additive. In particular, we have

$$V_{ij} = V_i + V_j - 2I_{ij}, \quad \text{where} \tag{22}$$

$$I_{ij} = f_{ij}^2 + \sum_{k \neq i,j} f_{ijk}^2 + \dots \tag{23}$$

is the total *epistatic variance* between loci i and j. The correlation between the new fitness after a double mutation and the previous fitness is $1 - 2V_{ij}/V \equiv 1 - 2v_{ij} = 1 - 2v_i - 2v_j + 2e_{ij}$, where $e_{ij} \equiv I_{ij}/V$ is the *epistatic variance fraction* between i and j.

2 Connectedness model

We introduce a "connectedness" model (the CN model, for short), where each locus has a probability μ_i of being involved in any particular interaction. We can interpret μ_i as the fraction of independent pathways that involve locus i, where each pathway has epistatic interactions between all loci involved in that pathway. The number of independent pathways, M, is assumed to be very large. The probability of an epistatic interaction between, say loci i, j, k, is proportional to $\mu_i \mu_j \mu_k$, since this is the probability that these loci are involved in the same pathway. The magnitude of the interaction term is $f_{ijk}^2 \sim \sigma^2 \mu_i \mu_j \mu_k$, where σ is the magnitude of the perturbative effect of the mutations, which is assumed to be constant for all orders of interaction ($\sigma^2 \propto M^{-1}$ to ensure finite variance for large M). The CN model leads to a specific interpretation (and hence its name): the outsized contribution to the variance from a particular locus is due to its involvement in many different complex pathways and not from an unusually large perturbative effect on a few pathways. For large L, the CN model is specified by the distribution, $P(\mu)$, across loci. In particular, given $P(\mu)$, we can calculate the total genetic variance, V, and the variance due to locus i, V_i. We define $\bar{\mu} \equiv \int_0^1 \mu P(\mu) d\mu$.

We first calculate the *expected* total variance across statistical ensembles. Note that here the expectations are averages over ensembles where the parameters of the model are re-sampled, in contrast

to the derivations presented in Section 1, which were microcanonical ensemble averages over equally likely genotypes. Since each pathway is independently sampled, expectations approximate the values in a single realization as $M \to \infty$. All expectations are denoted $\langle . \rangle$. We calculate the expected variance contribution from one pathway: since all pathways are statistically identical, the total variance from M pathways is simply M times the expected contribution from a single pathway. Since the μ_i are independent, we can then replace μ_i with $\bar{\mu}$ throughout unless explicitly conditioned when computing expectations. The contribution from the qth order interaction between loci $i_1, i_2, \ldots, i_q$ is $\langle f_{i_1 i_2 \ldots i_q}^2 \rangle = \sigma^2 \bar{\mu}^q$, where $\bar{\mu}$ is the mean probability that any locus is selected into the pathway. Since there are $\binom{L}{q}$ terms at order q, the total variance is

$$V = \sigma^2 \left\{ \binom{L}{1} \bar{\mu} + \binom{L}{2} \bar{\mu}^2 + \binom{L}{3} \bar{\mu}^3 + \ldots \right\} \tag{24}$$

$$= \sigma^2 \left((1 + \bar{\mu})^L - 1 \right). \tag{25}$$

The variance due to locus i *conditional* on μ_i is

$$V_i = \sigma^2 \left\{ \mu_i + \binom{L-1}{1} \mu_i \bar{\mu} + \binom{L-1}{2} \mu_i \bar{\mu}^2 + \ldots \right\} \tag{26}$$

$$= \sigma^2 \mu_i \left(1 + \bar{\mu} \right)^{L-1}. \tag{27}$$

Therefore, we have

$$v_i = V_i / V = \frac{\mu_i \left(1 + \bar{\mu} \right)^{L-1}}{(1 + \bar{\mu})^L - 1}. \tag{28}$$

There are two qualitatively different regimes, $\bar{\mu} L \ll 1$ and $\bar{\mu} L \gg 1$. When $\bar{\mu} L \ll 1$, each pathway typically contains a single locus and therefore should lead to an additive model. In this limit, we can write from Eq. (28),

$$v_i \approx \frac{\mu_i}{L \bar{\mu}}, \quad \bar{\mu} L \ll 1, \tag{29}$$

which is consistent with the expectation that $\sum_i v_i = 1$ and $v_i \sim L^{-1}$ for an additive model. In the opposite limit, $\bar{\mu} L \gg 1$, we have

$$v_i \approx \frac{\mu_i}{1 + \bar{\mu}}, \quad \bar{\mu} L \gg 1. \tag{30}$$

The CN model therefore leads to an intuitive interpretation of the variance fraction as being determined by the sparsity of interactions of a locus. Eq. (30) yields an upper bound at $v_i = 0.5$, which occurs when $\mu = 1$ for all loci. From Eq. (15), we see that this case is equivalent to the House-of-Cards model of maximal epistasis where every locus interacts with every other locus. The fitness after a mutation is independent from its previous fitness. The DVF $P(v)$ is directly determined by $P(\mu)$; the CN model can therefore be used as a generative model to generate arbitrary DVFs.

We can further calculate the epistatic variance between two loci in the CN model. The total epistatic variance I_{ij} between loci i and j is

$$I_{ij} = \langle f_{ij}^2 \rangle + \sum_{k \neq i,j} \langle f_{ijk}^2 \rangle + \ldots \tag{31}$$

$$= \sigma^2 \mu_i \mu_j + \sigma^2 \binom{L-2}{1} \mu_i \mu_j \bar{\mu} + \sigma^2 \binom{L-2}{2} \mu_i \mu_j \bar{\mu}^2 + \ldots \tag{32}$$

$$= \sigma^2 \mu_i \mu_j (1 + \bar{\mu})^{L-2}. \tag{33}$$

In the limit $\bar{\mu}L \gg 1$, the epistatic variance fraction after dividing by V is then simply $\mu_i\mu_j/(1+\bar{\mu})^2 = v_iv_j$. Using Eq. (22), we therefore have

$$v_{ij} = v_i + v_j - 2v_iv_j. \tag{34}$$

If v_i's are small, the CN model predicts near-additivity between the effects of two loci. This is *not* inconsistent with the strong epistasis assumption implicit in the limit $\bar{\mu}L \gg 1$: though the total contribution of epistatic interactions to the genetic variance may be large, the epistatic variance between two specific loci can still be negligible. This is because the majority of epistatic variance is due to the combinatorially large number of higher-order epistatic terms whose individual effects themselves can be weak. This has implications for experimental estimates of epistasis, which typically consider at most ~ 10 loci. If one wants to extrapolate the total pairwise variance of $L \gg 10$ loci from the pairwise variance estimates of a subset of $l = 10$ loci, the pairwise contributions should be scaled with a factor $(L/l)^2$ in contrast to L/l for the additive variances.

2.1 Relationship to statistical fitness landscapes

Statistical fitness landscapes such as the NK model and the Rough Mt. Fuji model are closely related to the CN model described above, with one key difference that gives qualitatively different adaptation properties. Specifically, in contrast to the CN model, the classical fitness landscape picture corresponds to a mean-field theory where the effect of each locus is assumed to be a constant. The DFE, which is determined by the DVF in our picture, instead comes from a single constant value $v = \bar{v}$. The fluctuations of the fitness effects are solely due to the genotype-dependent term η, which is a Gaussian. From Eq. (16), the DFE then corresponds to a Gaussian with mean $-2v(y - \bar{y})$. As adaptation proceeds and y increases, the DFE shifts to the deleterious side but retains its Gaussian shape. As we argue in the main text, the adaptation properties that result from this DFE are quite different from those arising from our theory. Further, while the fitness landscape picture may lead to a good approximation to our results when the number of mutations is large, it does not capture the different magnitudes of diminishing-returns for different loci observed in experiments.

The CN model for the special case of homogeneous loci, $\mu_i = \bar{\mu}$ for all i, is similar to the NK model but with one major difference: the number of loci in each pathway in the NK model is fixed at K loci, whereas here the typical pathway size is controlled by a continuous parameter $\bar{\mu}$. This introduces contributions at every epistatic order while effectively imposing sparsity on the contributions from higher-order interactions.

The statistical fitness landscapes often discussed are stationary Gaussian processes on a L-dimensional hypercube. The homogeneous CN model also falls into this class as $M \to \infty$. The key quantity that defines such a fitness landscape is the covariance function between the fitnesses of genotypes g and g', $C(g, g') = C(d)$, where $d \equiv |g - g'|$ is the Hamming distance between two genotypes. We now compute the covariance $C(d)$ for the homogeneous CN model. Each order term in each pathway is independent and the covariances for each order add up over all pathways. We first calculate the expectation of the first order term in a single pathway, which is

$$\sum_{i,i'} \langle f_i x_i f_{i'} x'_{i'} \rangle = \sigma^2 \bar{\mu} \sum_{i,i'} \delta_{ii'} x_i x'_{i'} \tag{35}$$

$$= \sigma^2 \bar{\mu} g.g' \tag{36}$$

$$= \sigma^2 \bar{\mu} \left(L - 2d \right). \tag{37}$$

Here we have used $\langle f_i^2 \rangle = \bar{\mu}\sigma^2$, where the $\bar{\mu}$ comes from the probability of locus i being selected for the module as argued previously. The covariance is linear in the distance between genotypes, as one would expect from an additive model. Directly calculating the higher order terms is more complicated because of the ordering restriction $i > j$ for the second order term and for higher orders. As has been noted previously [2, 3, 4], these can instead be calculated using combinatorics, which we will

demonstrate with the second-order term. We have for the second-order covariance

$$\sum_{i>j,i'>j'} \langle f_{ij} x_i x_j f_{i'j'} x'_{i'} x'_{j'} \rangle = \sigma^2 \bar{\mu}^2 \sum_{i>j,i'>j'} \delta_{ii'} \delta_{jj'} x_i x'_{i'} x_j x'_{j'} \tag{38}$$

$$= \sigma^2 \bar{\mu}^2 \sum_{i>j} (gg')_i (gg')_j, \tag{39}$$

where the element-wise product $(gg')_i \equiv x_i x'_i$ is 1 if $x_i = x'_i$ match and -1 otherwise. If d is the distance between g and g' then the element-wise product gg' has d -1 terms and $L-d$ 1 terms. The term in the summation above is 1 if both $(gg')_i$ and $(gg')_j$ are chosen from the $L-d$ subset or both are chosen from d subset. This term is -1 if one of the two is chosen from the $L-d$ subset and the other from the d subset. The number of terms that are 1s are therefore $\binom{L-d}{2}\binom{d}{0} + \binom{L-d}{0}\binom{d}{2}$ and the number of terms that are -1s are $\binom{L-d}{1}\binom{d}{1}$. This argument is easily extended to higher orders. The general qth order contribution to the covariance is

$$\sigma^2 \bar{\mu}^q \sum_{k=0}^{\min(q,d)} (-1)^k \binom{L-d}{q-k}\binom{d}{k}. \tag{40}$$

It is easily verified that the first-order term matches. When $d = 0$, we recover the binomial coefficients, as expected. The summation above is precisely the Krawtchouk polynomial $\mathcal{K}_q(d; L, 2)$, which we will denote by $\mathcal{K}_q(L, d)$. We therefore have

$$C(d) = \sigma^2 \sum_{q=1}^{d} \bar{\mu}^q \mathcal{K}_q(L, d). \tag{41}$$

The generating function of the Krawtchouk polynomials yields

$$C(d) = \sigma^2 \left((1 + \bar{\mu})^{L-d} (1 - \bar{\mu})^d - 1 \right). \tag{42}$$

The above expression is consistent with intuition: when $\bar{\mu}L \ll 1$, the covariance is linear in d, as expected for an additive model. In the opposite limit of $\bar{\mu}L \gg 1$ and $0 < d < L/2$, the constant term 1 can be ignored, and we see that the covariance is proportional to $e^{-\lambda d}$, where $\lambda = \ln\left(\frac{1+\bar{\mu}}{1-\bar{\mu}}\right)$. Epistasis is maximal when $\bar{\mu} \to 1$, in which case $\lambda \to \infty$ and the covariance rapidly goes to zero with d. This is the House-of-Cards model of maximal epistasis.

3 Adaptation

3.1 The distribution of fitness effects

Long-term adaptation is determined by the distribution of fitness effects (DFE) and the stochastic dynamical processes that lead to fixation. Before analyzing properties of adaptive walks, we clarify what our previous analysis, which ultimately led to Eq. (16), means for the DFE.

Eq. (16) represents the distribution of the fitness effect of a specific mutation at locus i over *all* genotypes in the population that have fitness y. We are instead interested in the DFE, where fitness effects are measured for all the mutations of a *particular* genotype that has fitness y. The DFE for a particular genotype generally depends on the idiosyncratic epistatic interactions between loci. In order to make this explicit, we turn to our analysis framework from Section 1. For notational convenience, we will assume $\bar{y} = 0$ and $V = 1$ in this sub-section and the next.

The DFE over L loci for a particular genotype of a fitness y can be thought of as a *sample*, $s_1, s_2, \ldots, s_L$, from the conditional joint distribution $P(s_1, s_2, \ldots, s_L | y)$. From our Gaussianity assumption, this joint distribution can be defined entirely in terms of the means and covariances of the

$L+1$ variables $y, s_1, s_2, \ldots, s_L$. Recall that the fitness effect of a mutation at locus i is $s_i = y_i - y = -2\xi_i$ as defined in Eq. (10). As shown previously, we have $\overline{s_i y} = -2v_i$. We also have $\overline{s_i s_j} = 4e_{ij}$ with $e_{ii} = v_i$. The means of all the variables are zero. Based off this covariance structure, we can compute $P(s_1, s_2, \ldots, s_L | y)$ using the properties of the conditional distribution of a multivariate normal distribution. It is straightforward to show that conditional on the fitness y, the conditional means are given by $\text{Mean}_y(s_i) = -2v_i y$ and the conditional covariances are $\text{Cov}_y(s_i, s_j) = 4(e_{ij} - v_i v_j)$. This relation makes clear that in general the DFE from a sample $s_1, s_2, \ldots, s_L \sim P(s_1, s_2, \ldots, s_L | y)$ depends on the epistatic interactions between all pairs of loci via e_{ij}. Note that $\text{Var}_y(s_i) = 4v_i(1 - v_i)$, which leads to Eq. (16) for the marginal distribution of a particular locus.

The DFE simplifies considerably if we make certain additional assumptions on the nature of epistatic interactions. In particular, the conditional correlation between fitness effects is

$$\text{Corr}_y(s_i, s_j) = \frac{e_{ij} - v_i v_j}{\sqrt{v_i v_j (1 - v_i)(1 - v_j)}}. \tag{43}$$

If we assume the typical variance fraction, $\bar{v}$ is small (i.e., $\bar{v} \ll 1$) and also that e_{ij} is $O(\bar{v}^2)$, then correlations are $O(\bar{v})$ and thus negligible. Then, in a particular sample $s_1, s_2, \ldots, s_L$, we can think of each s_i as being drawn independently with mean $-2v_i y$ and variance $4v_i(1 - v_i)$. If L is large, then this leads to a DFE

$$\rho(s|y) = \int_0^1 dv (2\sqrt{v(1-v)})^{-1} P(v) \varphi\left(\frac{s + 2vy}{2\sqrt{v(1-v)}}\right), \tag{44}$$

where $P(v)$ is the distribution of variance fractions (DVF) across the loci and φ is the standard normal pdf. Surprisingly, this relation becomes exact for the CN model, where we have shown that $e_{ij} = v_i v_j$ (Eq. (34)) and therefore the correlations between s_i's vanish. In this case, the DFE is thus determined entirely by the DVF, but we will show later that it has certain universal properties independent of even the DVF.

Above, we have measured the DFE for the subset of genotypes that have fitness y without regard to their evolutionary history. Over the course of adaptation, mutations are fixed and certain fitness changes are observed. We would then like to measure the DFE for those genotypes that have undergone a particular adaptive trajectory. As we show below, the DFE again simplifies considerably if certain relations hold.

3.2   History-dependence of the DFE

Using an analysis similar to the one in the previous section, we quantify history-dependence by calculating the correlations between the new fitness and the adaptive history conditional on the genotypes that have undergone a specific sequence of events in the past.

To be precise, suppose an initial clonal population of fitness y_0 gains k successive mutations and the corresponding sequence of fitnesses is $y_1, y_2, \ldots, y_k$. We would like to understand how the fitness of a new mutation at locus $k + 1$, y_{k+1}, depends on past fitnesses and the idiosyncratic epistatic interactions between the previous k mutations and the new mutation. The correlation between any two fitnesses y_i and y_j $(i < j)$ is given by $1 - 2v_{i+1:j}$, where $v_{i+1:j}$ is the variance fraction of the loci $i + 1, i + 2, \ldots, j$ (the subscript notation will be used throughout). In general, $v_{i+1:j}$ accounts for the epistatic interactions of all orders between these $j - i$ loci and is expressible in terms of the coefficients of our original complex trait model in Eq. (3). One can then write the covariance matrix between $y_0, y_1, \ldots, y_{k+1}$. In block form, this is

$$\Sigma = \begin{pmatrix} \Sigma_{0:k,0:k} & \Sigma_{0:k,k+1} \\ \Sigma_{k+1,0:k} & \Sigma_{k+1,k+1} \end{pmatrix}. \tag{45}$$

The mean of y_{k+1} conditional on $y_0, y_1, \ldots, y_k$ is a linear weighted sum of the past fitnesses:

$$\overline{y_{k+1}} = \Sigma_{k+1,0:k} \Sigma_{0:k,0:k}^{-1} \boldsymbol{y}, \tag{46}$$

where $\boldsymbol{y}$ is a vector with elements $y_0, y_1, \ldots, y_k$. In other words, y_{k+1} can be written as

$$y_{k+1} = \sum_{i=0}^{k} w_{k+1,i} y_i + \epsilon, \tag{47}$$

where ϵ is a mean-zero stochastic term that depends on the genotype of the initial population and whose variance can be calculated from Σ. To gain intuition, it is useful to explicitly calculate the case of $k = 1$. In this case, the covariance matrix is

$$\Sigma = \begin{pmatrix} 1 & 1 - 2v_1 & 1 - 2v_{12} \\ 1 - 2v_1 & 1 & 1 - 2v_2 \\ 1 - 2v_{12} & 1 - 2v_2 & 1 \end{pmatrix}. \tag{48}$$

We have

$$\Sigma_{2,01} \Sigma_{01,01}^{-1} = \frac{1}{4v_1(1 - v_1)} \begin{pmatrix} 1 - 2v_{12} & 1 - 2v_1 \end{pmatrix} \begin{pmatrix} 1 & 2v_1 - 1 \\ 2v_1 - 1 & 1 \end{pmatrix} \tag{49}$$

$$= \frac{1}{4v_1(1 - v_1)} \begin{pmatrix} 1 - 2v_{12} - (1 - 2v_1)(1 - 2v_2) & (1 - 2v_2) - (1 - 2v_1)(1 - 2v_{12}) \end{pmatrix}. \tag{50}$$

We therefore have

$$\overline{y_2} = \frac{v_1 - v_2 + v_{12} - 2v_1 v_{12}}{2v_1(1 - v_1)} y_1 + \frac{v_1 + v_2 - 2v_1 v_2 - v_{12}}{2v_1(1 - v_1)} y_0. \tag{51}$$

The dependence on the past has complex dependencies on epistasis between loci 1 and 2 even in this highly simplified case. To identify what contributes to history dependence beyond just the most recent fitness, we re-write Eq. (51) as

$$\overline{y_2} = (1 - 2v_2)y_1 + \frac{(1 - 2v_1)(v_{12} - v_1 - v_2 + 2v_1 v_2)}{2v_1(1 - v_1)} y_1 + \frac{v_1 + v_2 - 2v_1 v_2 - v_{12}}{2v_1(1 - v_1)} y_0. \tag{52}$$

From here, we observe that the dependence on v_1 and y_0 vanishes precisely when $v_{12} = v_1 + v_2 - 2v_1 v_2$. This is not true for all landscapes. However, as noted previously, this is in fact true for the CN model (Eq. (34)) when $\bar{\mu}L \gg 1$. In this case, we get the simple relation

$$y_2 = (1 - 2v_2)y_1 + 2\sqrt{v_2(1 - v_2)}\eta, \tag{53}$$

where the pre-factor in the second term comes from normalization and η is a Gaussian random variable.

From Eq. (22), we have $v_{12} = v_1 + v_2 - 2e_{12}$, where e_{12} is the epistatic variance fraction between loci 1 and 2. We can then re-write Eq. (52) as

$$\overline{y_2} \approx (1 - 2v_2)y_1 + \frac{v_1 v_2 - e_{12}}{v_1} s_1, \tag{54}$$

for $v_1 \ll 1$ and $s_1 = y_1 - y_0$. The equation gives us some intuition about how modular epistasis comes into play. As noted previously, the case $e_{12} = v_1 v_2$ corresponds to a special case when the history dependence vanishes. If instead $e_{12} > v_1 v_2$, the effect of epistasis between loci 1 and 2 is to *decrease* the fitness effect of locus 2 if locus 1 is beneficial ($s_1 > 0$) and increase it if locus 1 is deleterious. If $e_{12} < v_1 v_2$, then the opposite effect occurs: the fitness effect of locus 2 is larger if locus 1 is beneficial. This can be intuited as follows: Suppose instead of the CN model, fitness is determined by a linear sum of contributions from independent modules that contain distinct, non-overlapping sets of loci. Suppose also that loci 1 and 2 belong to two different modules Δ_1, Δ_2 respectively and the mutation at locus 1 is beneficial. Then, conditional on the current fitness y_1 and conditional on a beneficial mutation in Δ_1, it is likely that Δ_2 is less adapted relative to Δ_1. Statistically speaking, diminishing-returns should then be stronger for Δ_1 than any other module. Therefore, the fitness effect of a mutation in another module is larger relative to one in Δ_1 after taking the overall effects of y_1 into account. This analysis suggests that during adaptation, fixed beneficial mutations are more likely to be spread out over different modules.

3.3 Sufficient condition for memoryless fitness gains

The $k = 1$ case suggests that the relation for memoryless fitness gains ($e_{12} = v_1 v_2$) could in fact be true for all k under the CN model, which indeed turns out to be the case, as we show below. Motivated by the $k = 1$ case, we would like to have $w_{k+1,k} = 1 - 2v_{k+1}$ in Eq. (47) and the rest of the weights equal to zero. If this is the case,

$$y_{k+1} = (1 - 2v_{k+1})y_k + \epsilon. \tag{55}$$

Multiplying both sides by y_j and computing the correlations, we get the condition $1 - 2v_{j+1:k+1} = (1 - 2v_{j+1:k})(1 - 2v_{k+1})$ for all j. Therefore, a sufficient condition for memoryless fitness gain is

$$v_{j+1:k+1} = v_{k+1} + v_{j+1:k} - 2v_{k+1}v_{j+1:k} \tag{56}$$

for all k and for all $j < k$. We will now show that this is true for the CN model for $j = 0$, i.e., $v_{1:k+1} = v_{k+1} + v_{1:k} - 2v_{k+1}v_{1:k}$; the rest trivially follows. Let us first analyze what terms contribute towards $V_{1:k}$ (dividing by V gives $v_{1:k}$). When loci 1 through k are mutated, their effect is to flip the signs of their coefficients in Eq. (3). The ones which have changed sign contribute to the de-correlation between the fitnesses before and after the set of mutations, but the epistatic terms which have an even number of flips do not change and therefore their contribution has to be subtracted. $V_{1:k}$ therefore is the sum of squares of all the coefficients in Eq. (3) which have flipped an odd number of times. Suppose $i_1, i_2, \ldots, i_k$ are the indices of the k loci. Then,

$$V_{1:k} = \sum_{i_1=1}^{k} F_{i_1}^2 + \sum_{i_1>i_2>i_3}^{k} F_{i_1 i_2 i_3}^2 + \ldots, \quad \text{where} \tag{57}$$

$$F_{i_1}^2 = f_{i_1}^2 + \sum_{j_1 \neq 1:k} f_{i_1 j_1}^2 + \sum_{j_1 > j_2 \neq 1:k} f_{i_1 j_1 j_2}^2 + \ldots,$$

$$F_{i_1 i_2 i_3}^2 = f_{i_1 i_2 i_3}^2 + \sum_{j_1 \neq 1:k} f_{i_1 i_2 i_3 j_1}^2 + \sum_{j_1 > j_2 \neq 1:k} f_{i_1 i_2 i_3 j_1 j_2}^2 + \ldots,$$

and so on. Now, when the $k + 1$th locus is also flipped, to compute $V_{1:k+1}$, we can add up the two variances $V_{1:k}$ and V_{k+1} except for the cross terms which have an even number of sign flips and which include both the $k + 1$th locus and the other k loci. These have to be subtracted twice because they appear both in $V_{1:k}$ and V_{k+1}. These terms are

$$I_{1:k,k+1} = \sum_{i_1=1}^{k} F_{i_1 k+1}^2 + \sum_{i_1>i_2>i_3}^{k} F_{i_1 i_2 i_3 k+1}^2 + \ldots, \tag{58}$$

where the Fs are defined in a similar fashion as in Eq. (57) except the sums over js run from $k+2$ to L instead of $k+1$ to L. We get the general relation

$$V_{1:k+1} = V_{k+1} + V_{1:k} - 2I_{1:k,k+1}. \tag{59}$$

For the CN model specifically, we have $\langle f_{i_1 i_2 \ldots i_k}^2 \rangle = \sigma^2 \prod_{j=1}^{k} \mu_{i_j}$. This implies

$$\langle I_{1:k,k+1} \rangle = \sum_{i_1=1}^{k} \langle F_{i_1 k+1}^2 \rangle + \sum_{i_1>i_2>i_3}^{k} \langle F_{i_1 i_2 i_3 k+1}^2 \rangle + \ldots, \tag{60}$$

$$= \sigma^2 \mu_{k+1}(1 + \bar{\mu})^{L-k-1} \left(\sum_{i_1=1}^{k} \mu_{i_1} + \sum_{i_1>i_2>i_3}^{k} \mu_{i_1} \mu_{i_2} \mu_{i_3} + \ldots \right).$$

Performing a similar calculation for $\langle V_{1:k} \rangle$, we find

$$\langle V_{1:k} \rangle = \sigma^2 (1 + \bar{\mu})^{L-k} \left(\sum_{i_1=1}^{k} \mu_{i_1} + \sum_{i_1 > i_2 > i_3}^{k} \mu_{i_1} \mu_{i_2} \mu_{i_3} + \dots \right), \tag{61}$$

which gives

$$\langle I_{1:k,k+1} \rangle = v_{k+1} \langle V_{1:k} \rangle. \tag{62}$$

Dividing Eq. (59) by V throughout concludes the derivation.

3.4 Adaptedness

Under the conditions of memoryless fitness gains from the previous section, we can write

$$\sigma = -2vz + 2\sqrt{v(1-v)}\nu, \tag{63}$$

where variables have been re-scaled as $z = V^{-1/2}(y - \bar{y}), \sigma = V^{-1/2}s, \nu = V^{-1/2}\eta$. This equation suggests various forms for the DFE, $\rho(\sigma|z)$, depending on the DVF, $P(v)$. Note that our theory allows us to derive the DFE starting from rather general assumptions on the organization of the genotype-phenotype map, in contrast to past models which assume the DFE as a starting point.

z is a simple measure of "adaptedness": (1) When z is negative, the organism is in the unlikely situation of being "negatively-adapted" to the environment. Beneficial mutations are much more likely than deleterious mutations and adaptation is dominated by loci that have a large v. (2) When $|z| \ll 1$, the organism is "neutrally-adapted". The number of beneficial and deleterious mutations are balanced. (3) When $z \gg 1$, the organism is "well-adapted", where the DFE is strongly skewed towards deleterious mutations.

We will analyze adaptation in the neutrally-adapted and the well-adapted regimes. When the organism is negatively-adapted, Eq. (63) predicts that a few substitutions are sufficient to quickly reach the neutrally-adapted state. In addition, we assume the variance fraction of each locus is small, i.e., $v \ll 1$ but the number of loci L is large enough so that overall epistasis $\approx \bar{v}L \gg 1$ ($\bar{v}$ is the mean variance fraction). We can then ignore the $1 - v$ factor and re-write Eq. (63) as

$$\sigma = -2vz + 2\sqrt{v}\nu. \tag{64}$$

The intuition behind the analytical results is discussed in the main text. We present the formal calculations here.

3.5 Analytical results for an exponential DVF

We begin by calculating the DFE, $\rho(\sigma|z)$, for the specific case when the DVF is an exponential,

$$P(v) = \bar{v}^{-1} e^{-v/\bar{v}}, \tag{65}$$

where $\bar{v} \ll 1$ is the mean variance fraction across loci. The true DVF likely has a large fraction of loci that have no effect; accounting for these loci simply scales the mutation rate and will be ignored in this analysis. The exponential DVF leads to analytical predictions for the shape of the average fitness trajectories under certain simplifying assumptions about the underlying selective forces.

From Eq. (64), we have

$$\rho(\sigma|z) = \int \int dv d\nu P(v) P(\nu) \delta(\sigma + 2vz - 2\sqrt{v}\nu). \tag{66}$$

Here ν is a standard normal random variable, which we integrate out, giving

$$\rho(\sigma|z) = \int_0^1 dv (2\sqrt{v})^{-1} P(v) \varphi\left(\frac{\sigma + 2vz}{2\sqrt{v}}\right), \tag{67}$$

where φ is the normal pdf. For $P(v)$ given in Eq. (65), this integral can be calculated exactly:

$$\rho(\sigma|z) = \frac{\bar{v}^{-1}}{2\sqrt{2\bar{v}^{-1} + z^2}} e^{-\sigma z/2 - |\sigma|\sqrt{2\bar{v}^{-1} + z^2}/2}. \tag{68}$$

The resulting DFE is a double exponential with scale $\sim (z/2 \pm \sqrt{2\bar{v}^{-1} + z^2}/2)^{-1}$. For $|z| \ll 1$ small, the DFE is symmetric around the origin, as expected, with a scale determined by the DVF. As z increases, the DFE skews towards deleterious effects. The typical magnitudes of beneficial and deleterious effects are not independent from the overall ratio of beneficial to deleterious mutations. The well-adapted regime is reached when z^2 is comparable to $\bar{v}^{-1}$. To clearly delineate the two regimes, it is useful to define new variables $x = z\sqrt{\bar{v}/2}$ and $\lambda = \sigma\sqrt{2/\bar{v}}$. In the new variables, the DFE is

$$\rho(\lambda|x) = \frac{1}{4\sqrt{1+x^2}} e^{-\frac{\lambda x + |\lambda|\sqrt{1+x^2}}{2}}. \tag{69}$$

The neutrally-adapted and well-adapted regimes then correspond to $x \lesssim 1$ and $x \gtrsim 1$ respectively. The mean rate of adaptation in units of generations on the x scale is $\langle dx \rangle = \langle dz \rangle \sqrt{\bar{v}/2} = \langle \sigma \rangle \sqrt{\bar{v}/2} = \langle \lambda \rangle \bar{v}/2$, where the expectation is taken over fixation probabilities and the DFE. We assume strong-selection-weak-mutation (SSWM) so that $p_{\text{fix}}(\sigma) \sim 2\sigma$ or $p_{\text{fix}}(\lambda) \sim 2\lambda\sqrt{\bar{v}/2}$ for positive λ and 0 otherwise. We find

$$\langle dx \rangle = NU\bar{v}/2 \int_{-\infty}^{\infty} d\lambda \, p_{\text{fix}}(\lambda)\rho(\lambda|x)\lambda \tag{70}$$

$$= NU\bar{v}\sqrt{\bar{v}/2} \int_0^{\infty} d\lambda \, \lambda^2 \rho(\lambda|x), \tag{71}$$

where N is population size and U is the effective mutation rate for loci with non-zero effect. Integrating over λ, we get

$$\langle dx \rangle = \frac{NU(2\bar{v})^{3/2}}{\sqrt{1+x^2}(x + \sqrt{1+x^2})^3}. \tag{72}$$

Integrating over dx starting from an initial x_0, we therefore obtain an approximation to the mean fitness trajectory

$$NU(2\bar{v})^{3/2} n_{\text{gen}} = \int_{x_0}^{x} dx \sqrt{1+x^2}(x + \sqrt{1+x^2})^3 = T(x) - T(x_0), \tag{73}$$

$$\text{where } T(x) = \frac{4x^5}{5} + \frac{5x^3}{3} + \frac{\sqrt{x^2+1}}{15}\left(12x^4 + 19x^2 + 7\right) + x. \tag{74}$$

Note that this is only an approximation for small x since the typical fixed beneficial effect is a discrete jump. We show the result from Eq. (73) in Figure 1a, where the dependencies on the equilibrium fitness $\bar{y}$ and genetic variance V are highlighted. There are two independent parameters that determine the scale ($\sqrt{\bar{v}/2V}$) and location ($\bar{y}$) of the fitness, and one parameter (NUn_{gen}) that determines the time scale. From Eq. (73), the $x \sim t$ and $x \sim t^{1/5}$ scalings in the neutrally-adapted ($x \ll 1$) and well-adapted regime ($x \gg 1$) respectively are apparent.

The number of substitutions, n_s, as a function of x can also be calculated under the SSWM assumption, which gives

$$\bar{v}n_s = N(x) - N(x_0), \tag{75}$$

$$\text{where } N(x) = x(x + \sqrt{x^2+1})/4 + \sinh^{-1}(x)/4, \tag{76}$$

which can be mapped onto time using Eq. (73). The scalings are $n_s \sim x \sim t$ in the neutrally-adapted regime and $n_s \sim x^2 \sim t^{2/5}$ in the well-adapted regime.

3.6 Asymptotics and general scaling results

We now derive the asymptotic properties of the DFE in the well-adapted regime $(\bar{v}^{1/2}z \gg 1)$. Writing out the Gaussian pdf in Eq. (44), we have

$$\rho(\sigma|z) = \int_0^1 dv (2\sqrt{2\pi v})^{-1} P(v) e^{-\frac{(\sigma+2vz)^2}{8v}} \tag{77}$$

$$= e^{-\sigma z/2} \int_0^1 dv (2\sqrt{2\pi v})^{-1} P(v) e^{-\frac{\sigma^2}{8v} - \frac{vz^2}{2}} \tag{78}$$

$$= e^{-\sigma z/2} \int_0^1 du (2\sqrt{2\pi u y})^{-1} P(u/z) e^{-\frac{z}{2}\left(\frac{\sigma^2}{4u}+u\right)}, \tag{79}$$

where the change of variables $v = u/z$ is used. When $z \gg 1$, Laplace's method can be used. The exponent is minimized when $u = |\sigma|/2$, which gives

$$\rho(\sigma|z) \approx (2z)^{-1} P(|\sigma|/2z) e^{-\frac{z}{2}(\sigma+|\sigma|)}. \tag{80}$$

The contribution towards a fitness effect σ at large z comes largely from loci with variance fraction $v \sim |\sigma|/2z$. Surprisingly, the exponential form of the beneficial DFE is determined entirely by the Gaussian tails of the genotype-dependent term. The argument can be easily generalized to non-Gaussian tail probabilities using a similar calculation. Eq. (80) implies that the ratio of the probabilities of beneficial and deleterious mutations is completely independent of the DVF:

$$\frac{\rho(\sigma|z)}{\rho(-\sigma|z)} = e^{-\sigma z}. \tag{81}$$

Such a relationship has been hypothesized previously based on simulations of an additive finite-sites model and the form of p_{fix} close to the high-interference limit that could result in a fitness plateau [5]. Using our theory, we have shown that this result is indeed true independent of the DVF and under our core hypotheses of normality and memoryless fitness gains. This result closely resembles the fluctuation theorem in statistical physics. If $p_{\text{fix}} \sim e^{\sqrt{V}T_c\sigma}$ in the high-interference limit, where T_c is the coalescent time, fitness should plateau when $p_{\text{fix}}(\sigma)\rho(\sigma|z) = p_{\text{fix}}(-\sigma)\rho(-\sigma|z)$, which is at

$$z_{\text{plateau}} = 2\sqrt{V}T_c. \tag{82}$$

The $\sqrt{V}$ appears to account for the re-scaling $\sigma = s/\sqrt{V}$.

We get Eq. (80) only if $P(v)$ has probability mass at $v = |\sigma|/2z$. This is not the case in the exponentially-correlated fitness landscape model with homogeneous loci, i.e., if for instance $P(v) = \delta(v - \bar{v})$, we instead get from Eq. (44)

$$\rho(\sigma|z) = \varphi\left(\frac{\sigma + 2\bar{v}z}{2\sqrt{\bar{v}}}\right). \tag{83}$$

The DFE in this case is therefore a Gaussian with mean shifting towards the deleterious side. The ratio of beneficial to deleterious mutations goes rapidly to zero as $\simeq \varphi(\sqrt{\bar{v}}z)/\sqrt{\bar{v}}z$ and adaptation sharply plateaus beyond $\sqrt{\bar{v}}z \simeq 1$.

From Eq. (80) various scaling results can be derived that apply independent of $P(v)$ *and* the Gaussian assumption on ν. We retain the exponential form for convenience. The rate of beneficial mutations is

$$U_b = U \int_0^\infty d\sigma (2z)^{-1} P(|\sigma|/2z) e^{-\sigma z} \tag{84}$$

$$= \frac{U}{2z^2} \int_0^\infty dw P(w/2z^2) e^{-w}. \tag{85}$$

Figure 1: The mean fitness trajectory and the mean number of substitutions predicted by the model in the strong-selection-weak-mutation regime. (a) The scaled fitness vs scaled time. Shown below is the fitness in log-scale to highlight the different scalings in the neutrally-adapted and well-adapted regimes. The slopes of the dashed lines are shown. (b) The scaled number of substitutions vs scaled time as in panel (a).

The integral has an effective upper cutoff at $w \sim O(1)$ and can be approximated as $U_b \approx UP(0)/2z^2 \sim z^{-2}$ under certain assumptions for $P(v)$ for small v. The pre-factor $P(0)$ suggests that the number of beneficial mutations depends on the number of small-effect loci. While strong epistatic loci drive adaptation in the neutrally-adapted regime, adaptation in the well-adapted regime is instead driven by weakly epistatic loci.

From Eq. (80), the typical size of a beneficial mutation is $\sigma \sim z^{-1}$. Under SSWM, $p_{\text{fix}} \sim \sigma \sim z^{-1}$. As argued previously, since $U_b \sim z^{-2}$, we get $dz/dt \sim z^{-4}$ and therefore we obtain the two scaling relations

$$z \sim t^{1/5}, n_s \sim t^{2/5}, \tag{86}$$

which apply independently of the form of the DVF in the well-adapted regime.

References

[1] Milo S Johnson, Alena Martsul, Sergey Kryazhimskiy, and Michael M Desai. Higher-fitness yeast genotypes are less robust to deleterious mutations. *Science*, 366(6464):490–493, 2019.

[2] Peter F Stadler and Robert Happel. Random field models for fitness landscapes. *Journal of Mathematical Biology*, 38(5):435–478, 1999.

[3] Johannes Neidhart, Ivan G Szendro, and Joachim Krug. Exact results for amplitude spectra of fitness landscapes. *Journal of theoretical biology*, 332:218–227, 2013.

[4] Atish Agarwala and Daniel S Fisher. Adaptive walks on high-dimensional fitness landscapes and seascapes with distance-dependent statistics. *Theoretical population biology*, 130:13–49, 2019.

[5] Daniel P Rice, Benjamin H Good, and Michael M Desai. The evolutionarily stable distribution of fitness effects. *Genetics*, 200(1):321–329, 2015.