Sterol 14-α-demethylase is vital for mitochondrial functions and stress tolerance in *Leishmania major*

Sumit Mukherjee[a*], Samrat Moitra[a], Wei Xu[a*], Veronica Hernandez[a] and Kai Zhang[a]#

[a]Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409, USA.

*Current address: Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO 63110, USA.

#Corresponding author: Kai Zhang, Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409, USA. Tel: 1-806-834-0550. E-mail: kai.zhang@ttu.edu.

Short title: Sterol synthesis affects mitochondria in *Leishmania*

ABSTRACT

Sterol 14-α-demethylase (C14DM) is a key enzyme in the biosynthesis of sterols and the primary target of azoles. In *Leishmania major*, genetic or chemical inactivation of C14DM leads to accumulation of 14-methylated sterol intermediates and profound plasma membrane abnormalities including increased fluidity and failure to maintain ordered membrane microdomains. These defects likely contribute to the hypersensitivity to heat and severely reduced virulence displayed by the C14DM-null mutants (*c14dm⁻*). In addition to plasma membrane, sterols are present in intracellular organelles. In this study, we investigated the impact of C14DM ablation on mitochondria. Our results demonstrate that *c14dm⁻* mutants have significantly higher mitochondrial membrane potential than wild type parasites. Such high potential leads to the buildup of reactive oxygen species in the mitochondria, especially under nutrient-limiting conditions. Consistent with these mitochondrial alterations, c*14dm⁻* mutants show impairment in respiration and are heavily dependent on glucose uptake and glycolysis to generate energy. Consequently, these mutants are extremely sensitive to glucose deprivation and such vulnerability can be rescued through the supplementation of glucose or glycerol. In addition, the accumulation of oxidants may also contribute to the heat sensitivity exhibited by *c14dm⁻*. Finally, genetic or chemical ablation of C14DM causes increased susceptibility to pentamidine, an antimicrobial agent with activity against trypanosomatids. In summary, our investigation reveals that alteration of sterol synthesis can negatively affect multiple cellular processes in *Leishmania* parasites and make them vulnerable to clinically relevant stress conditions.

AUTHOR SUMMARY

Sterols are well recognized for their stabilizing effects on the plasma membrane, but their functions in intracellular organelles are under explored, which hampers the development of sterol synthesis inhibitors as drugs. Our previous studies have demonstrated significant plasma membrane instability in the sterol biosynthetic mutant *c14dm⁻* in *Leishmania major*, a pathogenic protozoan responsible for cutaneous leishmaniasis causing 1-1.5 million infections a year. While the plasma membrane defects have undoubtedly contributed to the reduced virulence exhibited by *c14dm⁻* mutants, it was not clear whether other cellular processes were also affected. In this study, we revealed profound mitochondrial dysfunctions and elevated level of reactive oxygen species in *c14dm⁻* mutants. These sterol mutants rely heavily on glycolysis to generate energy and are extremely sensitive to glucose restriction. In addition, the accumulation of oxidants appears to be responsible (at least in part) for the previously observed heat sensitivity in *c14dm⁻* mutants. Thus, genetic or chemical inactivation of C14DM can influence the functions of cellular organelles beyond the plasma membrane. These findings shed light on the mechanism of action for azole compounds and provide new insight into the roles of sterol biosynthesis in *Leishmania* parasites.

INTRODUCTION

Parasitic protozoa of the genus *Leishmania* are the causative agents for a group of diseases in humans and animals known as leishmaniasis [1, 2]. During their life cycle, *Leishmania* parasites alternate between flagellated, extracellular promastigotes in sandfly vectors and non-flagellated, intracellular amastigotes in vertebrates. Current drugs for leishmaniasis are plagued with high toxicity, low efficacy and resistance is on the rise [3]. Therefore, research involving the identification and characterization of metabolic pathways that could provide new drug targets is of considerable value.

While mammalian cells synthesize cholesterol as the main sterol, *Leishmania* promastigotes primarily make ergostane-based sterols such as ergosterol and 5-dehydroepisterol [4-6]. Ergosterol is distinguished from cholesterol by the presence of an unsaturated side chain with a methyl group at the C-24 position and an additional double bond at C-7 to C-8 in the B ring of the sterol core. Differing from promastigotes, *Leishmania* amastigotes (which reside in macrophages) mainly salvage cholesterol from the host, while downregulating the de novo ergosterol synthesis (supplemental Figure S14 and S15 in [7]). Despite the low abundance of ergostane-based sterols in amastigotes, sterol synthesis inhibitors and amphotericin B have shown promise as anti-leishmaniasis drugs [5, 8-10], suggesting that the residue amount of endogenous sterols are crucial for amastigotes and/or for the binding of amphotericin B. Among these sterol synthesis inhibitors are azole compounds which primarily target the cytochrome P450 enzyme sterol 14-α-demethylase (C14DM) [5, 11, 12]. Contribution of sterol biosynthesis to *Leishmania* biology is not fully understood, which hinders the development and improvement of azoles as anti-leishmaniasis drugs.

C14DM was found to be refractory to genetic deletion in *Leishmania donovani*, the

causative agent for visceral leishmaniasis [13]. In contrast, our earlier study showed that this

enzyme was dispensable in *Leishmania major* (responsible for cutaneous leishmaniasis in the old

world) [7]. Thus, different *Leishmania* species may possess distinct requirements for sterol

synthesis. Importantly, although *L. major C14DM*-null mutants (*c14dm⁻*) are viable as

promastigotes under normal tissue culture conditions, they show increased plasma membrane

fluidity, disruption in lipid raft formation, and deficiency in lipophosphoglycan (LPG, a major

surface glycoconjugate and virulence factor [14]) synthesis [7]. These plasma membrane defects

probably contribute to the mutants' hypersensitivity to heat and their severely attenuated

virulence in mice, although other factors may be involved as well [7]. A later study on the sterol

C-24-methyltransferase indicate that the accumulation of aberrant, C-14-methylated sterol

intermediates, not the lack of ergostane-based sterols, is mainly responsible for the fitness loss

displayed by *c14dm⁻* mutants [15].

While sterols are well recognized for their stabilizing effects on the plasma membrane

[16, 17], their functions in intracellular organelles are poorly understood, as the end products of

sterol synthesis are mainly found on the plasma membrane and secretory vesicles in mammalian

cells and yeast [18-20]. In *L. major*, C14DM is primarily found in the ER but also present in the

mitochondria suggesting that this enzyme is needed there [7]. An earlier study reported the

presence of substantial amounts of endogenous and exogenous sterols in the mitochondria of

Trypanosoma cruzi, a related trypanosomatid protozoan [21]. Application of sterol biosynthesis

inhibitors such as ketoconazole and lovastatin altered the ultrastructure of mitochondria in *T.*

cruzi, leading to intense proliferation of the inner-mitochondrial membranes and necrotic cell

death [22]. Similarly, RNAi knockdown of ergosterol pathway genes in *Trypanosoma brucei*

resulted in disorganization of mitochondria structure and depletion of ATP [23]. In *Leishmania*,

the lipid composition of mitochondrial membrane has yet to be determined. Nonetheless,

structural and functional abnormalities were also observed among other defects following

treatment with azole compounds. For example, in *Leishmania amazonensis*, combinatorial

treatment of ketoconazole and terbinafine led to alterations in the distribution and appearance of

cristae and the formation of paracrystalline arrays within the mitochondrial matrix [24]. Another

study reported the collapse of mitochondrial membrane potential ($\Delta\Psi$m) following posaconazole

treatment in *L. amazonensis* [25]. Besides azoles, treatment of *L. amazonensis* and *T. cruzi* with

various azasterols, which inhibit the sterol C-24-methyltransferase, induced a loss of

mitochondrial matrix contents [22, 26, 27]. Although the off-target effects from inhibitors cannot

be excluded, these observations suggest that sterol synthesis play important roles in preserving

the structure and function of mitochondria in trypanosomatids.

Compared to mammalian cells which usually possess multiple mitochondria per cell,

trypanosomatids have only one, relatively large mitochondrion per cell [28]. In addition to

energy production, mitochondria in trypanosomatids have been implicated in fatty acid synthesis

and redox homeostasis during different stages of their life cycle [29, 30]. Mitochondrial

malfunctions, e.g. changes in $\Delta\Psi$m, disruption of mitochondrial membrane integrity, and

increase in reactive oxygen species (ROS) production, are major mediators of cell death under

various stress conditions [31-33].

In this study, we sought to address the following important questions. First, how does

C14DM inactivation affect the integrity and functions of mitochondria in *L. major*? Second,

besides heat sensitivity, do *c14dm⁻* mutants exhibit any other stress response defects? Third, how

do mitochondrial deficiencies influence the fitness of *c14dm⁻* parasites? Our results showed that

genetic or chemical disruption of C14DM severely compromised the mitochondria in *L. major*,

leading to significant ROS accumulation and extreme vulnerability to biologically relevant stress

conditions such as heat and glucose starvation. These findings may guide the development of

new treatment strategies that exploit the fitness cost associated with drug resistance.

RESULTS

***C14dm⁻* promastigotes show increased mitochondrial membrane potential (ΔΨm) and altered mitochondrial morphology.**

To examine if C14DM deletion affects the mitochondria in *L. major*, we labeled promastigotes with MitoTracker CMXRos, a fluorogenic dye that binds to mitochondria in a ΔΨm-dependent manner. As indicated in Fig 1A, *c14dm⁻* mutants showed an intense mitochondrial staining 2-4 times stronger than wild type (WT) and *c14dm⁻*/+C14DM (add-back) parasites. In addition, the mitochondria of *c14dm⁻* appeared to be more swollen with larger volume and surface area based on transmission electron microscopy and quantitative analysis by fluorescence microscopy (Fig 1B and Fig S1A). To examine whether the mitochondrial membrane in *c14dm⁻* was compromised, we analyzed the presence of cytochrome *c* in mitochondrial fraction and cytosolic fraction following extraction with digitonin, with ISCL and HSP83 being used as mitochondrial and cytosolic markers, respectively [34] [35](Fig S1B). Similar to WT and *c14dm⁻*/+C14DM parasites, cytochrome *c* in *c14dm⁻* remained in the mitochondrial enriched fractions when the digitonin extraction was performed at room temperature (Fig S1B). As a control, digitonin treatment at 37 °C disrupted the outer mitochondrial membrane, causing cytochrome *c* release into the cytosol [36] (Fig S1B). Thus, despite of its effects on ΔΨm and mitochondrial size, genetic ablation of C14DM does not affect mitochondrial membrane integrity at least with regard to cytochrome *c* localization under ambient conditions.

Consistent with the MitoTracker staining results, when live parasites were labeled with another ΔΨm-dependent dye tetramethylrhodamine ethyl ester (TMRE), *c14dm⁻* mutants retained 2-4 times more fluorescence than WT and *c14dm⁻*/+C14DM cells as indicated by flow

cytometry and microscopy (Fig 1C-D). The difference between WT and *c14dm⁻* parasites was

more pronounced during the log phase when cells were metabolically active than stationary

phase (Fig 1C). Similar results were observed after staining with rhodamine 123 (Fig S2A). As

expected, pretreatment with the mitochondrial uncoupler CCCp (75 μM for 15 min) dramatically

reduced the incorporation of TMRE and MitoTracker into WT and *c14dm⁻* parasites (Fig 1A and

1C), without significant effect on cell viability (<5% of cells were propidium iodide positive by

flow cytometry). Furthermore, WT promastigotes grown in the presence of itraconazole (ITZ, a

C14DM inhibitor) [7] for 48 hours at sublethal concentrations showed similar ΔΨm as *c14dm⁻*

mutants (Fig 1C). Finally, we also examined if extracellular nutrient level affects ΔΨm by

labelling promastigotes with TMRE in either complete M199 medium, M199 medium without

fetal bovine serum, phosphate-buffered saline (PBS) supplemented with 5.5 mM of glucose (the

glucose concentration in M199 medium), or PBS alone (Fig 2A). In these experiments, *c14dm⁻*

mutants exhibited 2.5-4 times higher ΔΨm than WT/add-back parasites and no significant

differences were detected between nutrient-replete and nutrient limiting conditions (Fig 2A).

Collectively, these observations suggest that C14DM inhibition leads to significantly increased

ΔΨm in *L. major* promastigotes regardless of extracellular nutrient levels.

Inactivation of C14DM leads to ΔΨm-dependent accumulation of ROS in the mitochondria.

In mammalian cells, a high ΔΨm may cause accumulation of ROS in the mitochondria

(mainly in the form of superoxide or $O_2^{\bullet-}$) in part due to reverse electron flow from complex II to

complex I of the electron transport chain (ETC) [37-39]. To assess the ROS level in *c14dm⁻*

mutants, promastigotes were labelled with a mitochondria-specific ROS indicator MitoSox Red

which binds to mitochondrial nucleic acid and fluoresces upon oxidation by superoxide [40]. In

complete M199, *c14dm⁻* mutants showed 3.5 times more MitoSox labeling than WT and add-

back cells (Fig 2B). Unlike ΔΨm, the mitochondrial ROS level in *c14dm⁻* was sensitive to

nutrient availability (Fig 2B): PBS only (the highest) > PBS with glucose > M199 medium

without fetal bovine serum > complete M199 (the lowest). Importantly, pretreatment of *c14dm⁻*

parasites with CCCp significantly reduced MitoSox Red labeling, indicating that the high ΔΨm

is mainly responsible for ROS accumulation (Fig 2A-B and Fig 1C). These findings suggest that

the mitochondria in *c14dm⁻* mutants are under significant oxidative stress during starvation.

Next we examined the mitochondrial ROS levels in log phase and stationary phase

promastigotes after labeling them with MitoSox in PBS alone (Fig 2C). Compared to WT and

add-back cells, *c14dm⁻* mutants contained 10-15 times more mitochondrial ROS by flow

cytometry during log phase and early stationary phase, and the difference became less dramatic

during late stationary phase when cells ceased proliferation (Fig 2C). Microscopic examination

of MitoSox-labelled parasites revealed singular fluorescent spots in every cell representing the

kinetoplast DNA, indicating the mitochondrial origin of ROS (Fig 2D). In these images, MitoSox

signals from *c14dm⁻* were much more prominent than those from WT and *c14dm⁻/+C14DM*

cells (Fig 2D), which was in agreement with flow cytometry analysis (Fig 2B-C). Furthermore,

when cultivated in the presence of ITZ (at 0.1-0.2 µM or near EC25 levels), *L. major* WT

promastigotes and *L. mexicana* axenic amastigotes displayed high levels of MitoSox Red

staining similar to *c14dm⁻* (Fig 2C and 2E), indicating that C14DM inhibition causes ROS built-

up in the mitochondria of *Leishmania* parasites. In comparison to *c14dm⁻*, mitochondrial defects

exhibited by the sterol C-24-methyltransferase mutants (*smt⁻*) were less profound (Fig S2A-

B)[15], suggesting that the accumulation of C-14-methylated sterols is mainly responsible for

these changes.

We then determined the cytoplasmic ROS level in *c14dm⁻* by labeling log phase parasites

with dihydroethidium (DHE), a cytosolic ROS indicator. As illustrated in Fig 2F, the amount of

cytosolic ROS in *c14dm⁻* was about two-fold higher than WT and add-back parasites, which was

not as dramatic as the increase of mitochondrial ROS. As a control, treatment of WT parasites

with hydrogen peroxide (100 μM for 10 min) led to significantly increased staining for DHE but

not MitoSox Red, confirming the specificity of MitoSox Red for mitochondrial superoxide (Fig

S3). Together, these data suggest that genetic or chemical ablation of C14DM leads to significant

accumulation of ROS inside the mitochondria.

Examination of superoxide dismutase (SOD) expression in *c14dm⁻* mutants.

As byproducts of mitochondrial respiration, ROS (mainly the form of $O_2^{\bullet-}$) can be

generated from the partial reduction of oxygen by complexes I, II, and III of the ETC [41]. To

limit the damaging effects of ROS, *Leishmania* cells utilize several antioxidant systems

involving superoxide dismutases (SODs) and tryparedoxin peroxidases [42, 43]. The fact that

c14dm⁻ parasites accumulate high levels of $O_2^{\bullet-}$ suggests that either the ROS removal system is

defective, or the production of $O_2^{\bullet-}$ is elevated due to excessive electron leakage from the ETC in

these mutants. Here we examined whether a dysregulation of SOD, the primary enzyme to

detoxify $O_2^{\bullet-}$, is responsible for the accumulation of mitochondrial ROS in *c14dm⁻*.

Leishmania encode three iron-dependent SODs: the mitochondrial SODA and two glycosomal

SODB (SODB1 and SODB2) [43, 44]. In whole cell lysate, we detected a 14-98% increase in

SOD activity from *c14dm⁻* mutants and the increase was more pronounced during the stationary

phase (Fig 3A). Western blot revealed that *c14dm⁻* mutants had similar levels of SODA

(mitochondrial) and SODB (cytosolic) protein as WT parasites (Fig 3B). The increase in SOD

activity in late stationary phase *c14dm⁻* coincided with the decrease of mitochondrial ROS

(although still higher than WT and add-back parasites, Figs 2C, S2B and 3A), suggesting an

adaptive response from the mutants to upregulate their antioxidant defense machinery.

Collectively, these findings indicate that the ROS is accumulated in *c14dm⁻* mutants despite of

near-normal SOD production.

***C14dm⁻* mutants show reduced cellular respiration and increased glycolytic ATP**

production.

The mitochondrial abnormalities in *c14dm⁻* prompted us to examine if these mutants

could effectively carry on respiration and generate ATP. First, we measured the cellular oxygen

consumption rate using the MitoXpress oxygen probe as described previously [45]. When

incubated a mitochondrial respiration buffer made of Hanks' balanced salt solution (HBSS)

supplemented with sodium pyruvate and 2-deoxy-G-glucose (to block glycolysis) [46], *c14dm⁻*

parasites showed a significant reduction (42-60%) in oxygen consumption compared to WT cells,

and such defect was corrected in the *c14dm⁻*/+C14DM parasites (Fig 4A-B). As expected, the

addition of antimycin A, a respiratory inhibitor that blocks cytochrome *c* reductase or complex

III [47], decreased oxygen consumption in WT parasites (Fig 4A). A similar experiment was

performed when cells were incubated in the Dulbecco's Modified Eagle's Medium (DMEM)

which contained 25 mM of glucose. While all cells exhibited higher oxygen consumption rates in

DMEM than in HBSS, *c14dm⁻* mutants still used less oxygen than WT and *c14dm⁻*/+C14DM

parasites (Fig 4C). Given the considerable ROS accumulation in *c14dm⁻*, the low oxygen

consumption rate may protect these mutants from further oxidative damage.

Next, we determined the ATP production from WT, *c14dm⁻* and *c14dm⁻*/+C14DM

parasites. When log phase promastigotes were incubated in an isotonic buffer (without glucose or

any inhibitor) for 1 hour, a 30-40% increase in ATP content was observed in *c14dm⁻* mutants in

comparison to WT or add-back parasites (Fig 4D, lane 1). Supplementation with glucose led to

increased ATP production in all cells, presumably from both glycolysis and mitochondrial

respiration (Fig 4D, lane 5). Importantly, despite the low oxygen consumption, *c14dm⁻* mutants

produced 30-40% more ATP than WT and add-back parasites from glucose (Fig 4D, lane 5),

suggesting that these cells have increased capacity to carry out glycolysis. Curiously, addition of

sodium pyruvate had only modest effect on ATP production (Fig 4D, lane 9). Although

Leishmania parasites possess putative plasma membrane-bound pyruvate transporters [48], the

efficiency of pyruvate uptake is not known. Another possibility is that parasites can sense the

lack of carbon sources besides pyruvate and regulate ATP biogenesis accordingly. As expected,

addition of 2-deoxy-G-glucose (5.5 mM) significantly reduced ATP production in the absence of

exogenous glucose (Fig 4D, lane 2 and 10). No significant reduction was detected when cells

were incubated in the presence of glucose (5.5 mM), which might be due to inadequate

competition on glucose uptake and utilization (Fig 4D, lane 6). Furthermore, addition of sodium

azide which inhibits cytochrome *c* oxidase [46, 49] dramatically lowered ATP production in all

cells (Fig 4D, lane 3, 7 and 11), suggesting that the glycolysis flux may be coupled to the

efficiency of oxidative phosphorylation respiration in *Leishmania* so that cells can turn down

glycolysis to prevent the buildup of toxic metabolites.

Glucose starvation has deleterious effect on *c14dm⁻* parasites.

Our findings so far suggest that in comparison to WT cells, *c14dm⁻* mutants depend more

on non-respiration pathways such as glycolysis to generate energy. If so, these mutants would be

highly sensitive to glucose restriction. To test this possibility, we exposed parasites to glucose-

limiting conditions and measured cell survival over time. As illustrated in Fig 5A, when

incubated in HBSS, *c14dm⁻* mutants showed substantial cell death after 12 hours. At 20-24 hours,

more than 90% of *c14dm⁻* cells were dead (Fig 5A-B). Similar results were observed when

parasites were incubated in other neutral, isotonic buffers such as PBS (Fig S4). In addition, *L.*

major WT parasites cultivated in the presence of ITZ were highly vulnerable to glucose

starvation like *c14dm⁻* mutants (Fig S5). In these experiments, WT control, *smt⁻*, *smt⁻*/+SMT and

c14dm⁻/+C14DM parasites showed very good survival (<9% death after 24 hours), indicating

this phenotype is caused by the ablation of C14DM (Fig 5B). Importantly, supplementation with

glucose at 5.5 mM (the glucose concentration in M199 medium) completely rescued *c14dm⁻*

mutants, whereas at the same molar concentration, sodium pyruvate (a respiration substrate), 2-

deoxy-G-glucose (a non-metabolizable glucose analog), or aspartate (a gluconeogenic amino

acid) failed to protect the mutants (Fig 5). Further tests revealed that glucose concentrations

above 100 μM were sufficient to rescue *c14dm⁻* mutants (Fig S4 and data not shown).

Recent studies have revealed that bloodstream forms of *Trypanosoma brucei* can use

glycerol to support growth in the absence of glucose [50, 51]. To test if *Leishmania* parasites can

utilize glycerol effectively, we attempted to rescue *c14dm⁻* mutants with glycerol. Remarkably,

when provided at more than 100 μM, glycerol was able to protect *c14dm⁻* mutants under

glucose-free conditions (Fig 5 and Fig S4). Both glucose and glycerol were also able to alleviate

the mitochondrial defects in *c14dm⁻* mutants (Fig S6). Together, our results indicate that

inactivation of C14DM leads to hypersensitivity to glucose starvation, which is likely caused by

energy depletion and excessive ROS accumulation.

C14dm⁻ parasites show increased uptake of glucose and glycerol.

Next, we examined whether the uptake of glucose and glycerol is altered in *c14dm⁻*

parasites using radiolabeled substrates as previously described [52, 53]. If respiration defects

force *c14dm⁻* mutants to rely more on glycolysis to generate energy, these parasites may have

higher demands for glucose or equivalent carbon sources. Trypanosomatid protozoans including

Leishmania species possess a family of glucose transporters [54] and aquaglyceroporins [53] that

can mediate nutrient uptake. As shown in Fig 6, *c14dm⁻* mutants were more efficient than WT

and add-back parasites in the uptake of glucose and glycerol. The incorporation of glucose was

much more robust at room temperature than at 4 °C (Fig 6A), whereas the incorporation of

glycerol was less dependent on temperature (Fig 6B-C). The increased uptake of glucose and

glycerol in *c14dm⁻* may serve as a compensatory mechanism to offset deficiencies in

mitochondrial respiration.

ROS accumulation is partially responsible for the heat sensitivity in *c14dm⁻* mutants.

We have previously shown that *c14dm⁻* parasites are extremely sensitive to heat stress

(e.g. 37 °C treatment), and their increased plasma membrane fluidity (induced by the

accumulation of C-14-methylated sterol intermediates) is a likely cause [7]. To investigate if the

accumulation of ROS is also a contributing factor, WT and *c14dm⁻* parasites were incubated in

PBS at either 27 °C (the regular temperature for promastigotes) or 37 °C. After 3 hours, a

marked reduction in ΔΨm was detected in 30-35% of *c14dm⁻* parasites without substantial cell

death (Fig 7A-B). No significant change in ΔΨm was observed in WT parasites (Fig 7B). To

determine if the loss of ΔΨm was associated with compromised mitochondrial membrane

integrity, we assessed the release of cytochrome *c* from mitochondria into the cytosol (Fig 7C-D).

For WT cells, the majority of cytochrome *c* was found in the mitochondrial enriched membrane

fractions under control (27 °C) and heat stress (37 °C, 3 hours) conditions (Fig 7C-D). In

contrast, ~40% of cytochrome *c* was detected in the cytosolic fraction when *c14dm⁻* parasites

were subjected to heat treatment (Fig 7C-D). Since 3 hours incubation at 37 °C did not induce

substantial cell death (Fig 7A), this finding indicates that the mitochondrial disruption occurred

when the majority of *c14dm⁻* mutants were viable. In addition, we measured the levels of

cytosolic ROS in control and heat stressed cells. Unlike WT and add-back parasites which

showed no significant change, *c14dm⁻* mutants displayed ~3-fold increase in cytosolic ROS

following 37 °C treatment for 3 hours (Fig 7E). Together, these data suggest that heat stress

causes a partial rupture of mitochondrial outer membrane in *c14dm⁻*, leading to the dissemination

of cytochrome *c* and ROS into the cytosol.

To further evaluate the contribution of ROS accumulation to the observed heat response

phenotype, we treated *c14dm⁻* mutants with glutathione (GSH, an antioxidant) before exposing

them to 37 °C. When administered at 5 mM, GSH treatment of *c14dm⁻* reduced the

mitochondrial ROS by ~60% and the heat-induced cell death in 8 hours by ~50% (Fig 7F-G).

The effect was much less prominent when GSH was provided at 1 mM (Fig 7F-G), indicating a

dosage-dependent alleviation of ROS stress. Additionally, we tested the effect of antimycin A, an

oxidative stress inducer (Fig S3A), on the heat sensitivity of WT parasites. At 27 °C, 10 µM of

antimycin A had very mild effect on cell survival (<10% death after 36 hours) but at 37 °C, more

than 60% of WT cells died after 8 hours (Fig 7H). From these studies, we conclude that in

addition to plasma membrane defects, oxidative stress also makes *c14dm⁻* mutants vulnerable to

heat.

***C14dm⁻* mutants exhibit increased sensitivity to pentamidine.**

Finally, to probe whether the mitochondrial defects in *c14dm⁻* can be exploited for

developing better chemotherapy, we tested the susceptibility of these parasites to pentamidine, an

antimicrobial agent with efficacy against trypanosomatids [55]. While the mode of action for

pentamidine is not completely understood, previous reports suggest that this drug needs to be

sequestered into the mitochondria of *Leishmania* in a $\Delta\Psi$m-dependent manner to exert its

activity [55, 56]. Thus, the elevated $\Delta\Psi$m may make *c14dm‾* more vulnerable to pentamidine.

Indeed, in a 48 hours drug response assay [15], *c14dm‾* mutants showed heightened growth

inhibition by pentamidine, as their EC90, EC50 and EC25 (effective concentrations required to

inhibit growth by 90%, 50% and 25% respectively) were 5-15 times lower than those of WT and

add-back parasites (Fig 8A). WT cells grown in the presence of ITZ were also hypersensitive to

pentamidine, suggesting that this drug can work synergistically with azoles (Fig 8A). In addition,

we examined whether short time exposure to pentamidine can cause death or irreversible damage

to *c14dm‾* mutants by treating parasites with 2.5 μM or 25 μM of pentamidine for one hour,

followed by two washes with PBS and then let cells grow in drug-free M199 media [57]. As

shown in Fig 8B, exposure to 2.5 μM of pentamidine for one hour led to significant grow delay

(2-3 days) in *c14dm‾* and these mutants could only grow to 5-9 x 10^6 cells/ml in six days (the

maximal density for *L. major* promastigotes in complete M199 is ~3 x 10^7 cells/ml). In contrast,

for WT and add-back parasites, 2.5 μM of pentamidine had little impact and it took 25 μM of

pentamidine to cause a similar degree of inhibition (Fig 8B-C). Meanwhile, 25 μM of

pentamidine was sufficient to halt the growth of *c14dm‾* nearly completely after the one-hour

exposure (Fig 8C). CCCp treatment was able to rescue the mutants in this wash out experiment

(Fig 8B-C), suggesting that the high $\Delta\Psi$m and/or mitochondrial ROS are responsible the

hypersensitivity of *c14dm‾* to pentamidine.

DISCUSSION

Our previous work has revealed significant plasma membrane defects in *c14dm⁻* mutants including increased membrane fluidity and failure to maintain detergent resistant membrane fractions [7]. While predominantly located in the plasma membrane, sterols are also present in substantial amounts in lipid droplets and intracellular organelle membranes [20, 58]. A more thorough characterization of the *c14dm⁻* mutants would reveal new insight into the roles of sterol metabolism in *Leishmania* and provide new leads to exploit the vulnerability caused by the accumulation of abnormal sterols. In this study, we found a 2-5 folds increase in the ΔΨm of *c14dm⁻* relative to WT and add-back parasites and similar results were observed in ITZ-treated WT parasites (Fig 1 and Fig S2). We also detected a modest swelling of mitochondria in *c14dm⁻* mutants (Fig 1B and Fig S1A). These findings are consistent with the previously reported hyperpolarization of mitochondrial membrane in *Leishmania mexicana* following treatment with triazole compounds [59].

In mammalian cells, elevated ΔΨm may cause electron flow in the reverse direction through the ETC leading to electron leakage and superoxide accumulation in the mitochondrial matrix [37, 39]. By flow cytometry and fluorescence microscopy, we observed significantly higher levels of superoxide in the mitochondria of *c14dm⁻* mutants and ITZ-treated WT parasites in comparison to control cells (Fig 2 and Fig S2-S3). Under the nutrient-replete condition, a 3-5-fold increase was detected in *c14dm⁻* mutants. Under the nutrient-limiting condition, the difference became 10-12-fold (Fig 2B). Meanwhile, the cytosolic ROS level in *c14dm⁻* mutants was only about 2-fold higher than WT and add-back parasites, suggesting that the majority of accumulated ROS was sequestered in the mitochondria under ambient temperature (Fig 2F and Fig S3). In agreement with this notion, the mitochondrial membrane integrity in *c14dm⁻*

appeared to be intact based on the intracellular distribution of cytochrome *c* (Fig S1B). It is not

clear why inactivation of C14DM leads to elevated levels of $\Delta\Psi$m and mitochondrial ROS. The

expression and activity of SODs were not compromised in *c14dm⁻* mutants (Fig 3), suggesting

that ROS production was higher in the absence of C14DM. In mammalian cells and yeast,

mitochondria can synthesize phosphatidylglycerol (PG), cardiolipin (CL), phosphatidic acid, and

a portion of phosphatidylethanolamine (PE, through the decarboxylation of phosphatidylserine)

[19, 60, 61]. Other lipids including phosphatidylcholine, sphingolipids and sterols have to be

imported. Changes in mitochondrial lipidome, especially the abundance of CL and PE, can lead

to ETC defects [62, 63]. It is therefore possible that defects in sterol synthesis at the ER alter the

lipid composition of mitochondrial membrane in *c14dm⁻* mutants. Such membrane modification

may negatively affect the functions of ETC including the terminal F0-F1 complex, leading to the

accumulation of proton motive force ($\Delta\Psi$m), leakage of electron and partial reduction of oxygen.

In comparison to WT and add-back parasites, *c14dm⁻* mutants showed significant

impairment in oxygen consumption yet produced higher than WT-level of ATP in a glucose-free

buffer (Fig 4). Similarly, the cellular ATP content in *c14dm⁻* mutants was 30-40% higher than

WT and add-back parasites when cells were incubated in the presence of glucose (Fig 4). These

results suggest that without C14DM, *Leishmania* parasites downregulate respiration and rely

more on glycolysis to generate energy. Mechanism of this energetics switch remains to be

determined. In many eukaryotes, the heterotrimeric AMP-activated protein kinase (AMPK)

complex is a key regulator of cellular energy metabolism [64-66]. Future investigation will

examine if AMPK activation in response to C14DM ablation is responsible for reduced

respiration and increased glucose consumption via glycolysis.

Consistent with the increased dependency on glucose as an energy source, *c14dm⁻*

mutants were extremely sensitive to glucose starvation (Fig 5). This hypersensitivity may arise

from the lack of energy production after glucose is depleted via glycolysis and the rapid increase

in ROS due to respiration deficiency (Fig 2 and Fig 5). An alternative and not mutually exclusive

explanation is that the accumulation of aberrant sterol intermediates in *c14dm⁻* mutants

compromised the progression of autophagy, a possibility currently under investigation.

Interestingly, supplementation of either glucose or glycerol at concentrations above 100 µM

completely rescued the survival of *c14dm⁻* mutants (Fig 5 and Fig S4). While glycerol could be

converted into glucose, it is unlikely to account for the rescue as the energy required for

gluconeogenesis exceeds that generated from glycolysis. Instead, it is possible that *Leishmania*

parasites metabolize glycerol into glycerol-3-phosphate (by glycerol kinase) and then

dihydroxyacetone phosphate (by the mitochondrial FAD-dependent glycerol-3-phosphate

dehydrogenase) which can be incorporated into the glycolysis pathway, as described in *T. brucei*

[50, 51]. We also observed increased uptake of glucose and glycerol by *c14dm⁻* mutants (Fig 6),

a possible adaptive response to their high demand for carbon sources. Future research will focus

on effects on sterol synthesis on the permeability of plasma membrane to probe the mechanism

of this phenotype.

Next, we examined the involvement of mitochondrial defects in the previously reported

heat sensitivity exhibited by *c14dm⁻* mutants. Despite the high mitochondrial ROS, *c14dm⁻*

parasites were fully viable at ambient temperature in the presence of glucose or glycerol.

However, within 3 hours of exposure at 37 °C, the $\Delta\Psi$m in *c14dm⁻* mutants showed signs of

collapse, along with a significant increase in cytosolic ROS level and the release of cytochrome *c*

to the cytosol (Fig 7). One possibility is that similar to the effect on plasma membrane, the

accumulation of aberrant sterol intermediates compromises the mitochondrial membrane

integrity at high temperature in *c14dm⁻* parasites. Although the presence of a canonical apoptotic

pathway in *Leishmania* is debatable, earlier studies have suggested the involvement of

mitochondrial contents such as cytochrome *c* in causing cell death [32, 67]. In addition,

treatment with antioxidant GSH could partially rescue the heat induced cell death *c14dm⁻*

mutants, and the ROS generating agent antimycin A made WT *L. major* vulnerable to heat (Fig

7). Together, these findings suggest that ROS accumulation is partially responsible for the heat

sensitivity in *c14dm⁻* parasites.

Finally, we also observed heightened sensitivity of *c14dm⁻* to pentamidine, in both the

48-hour growth inhibition assay and the one-hour exposure assay (Fig. 8). Previous studies on

the amphotericin B resistant lines in *L. mexicana* have identified mutations in sterol biosynthesis

and hypersensitivity to pentamidine to be a common feature among these parasites [68, 69]. Our

results here are in line with those previous reports as *c14dm⁻* mutants are highly resistant to

amphotericin B due to their lack of ergostane-based sterols [7]. It is possible that the elevated

ΔΨm and/or mitochondrial ROS cause accumulation of pentamidine in the mitochondria of

c14dm⁻, making them more susceptible to this drug than WT parasites [55, 56].

In summary, we have shown that deletion or inhibition of C14DM leads to significant

alterations in the mitochondria highlighted by increases in ΔΨm and ROS levels, along with

deficiency in respiration. These defects make *c14dm⁻* extremely vulnerable to glucose restriction

and contribute to their hypersensitivity to heat. Thus, in addition to its stabilizing effects on the

plasma membrane, sterol synthesis plays important roles in the regulation of cellular energetics,

redox homeostasis and parasite fitness under stress. Analysis of the dynamics of ER-

mitochondria transport, the mitochondrial lipid composition, and the spatial distributions and

activities of ETC complexes in *c14dm¯* would shed light on the mechanistic aspect of how

endogenous sterols influence mitochondria in *Leishmania*. Furthermore, the hypersensitivity of

c14dm¯ to pentamidine raises the possibility of using azole-pentamidine or amphotericin B-

pentamidine combinations against leishmaniasis. Future studies may also explore the application

of glycolysis inhibitors or ROS inducers in conjugation with sterol biosynthesis inhibitors.

MATERIALS AND METHODS

Materials

Tetramethylrhodamine ethyl ester (TMRE), rhodamine 123, Mitotracker CMXRos, MitoSox Red, and dihydroethidium (DHE) were purchased from Thermo Fisher Scientific. Itraconazole (ITZ) was purchased from LKT Laboratories. Antimycin A was from ENZO Life Sciences. The MitoXpress oxygen consumption assay kit was purchased from Luxcel Biosciences. For ATP analysis, the CellTiter-Glo luminescent assay kit was purchased from Promega. All other chemicals were purchased from VWR International or Sigma Aldrich Inc. unless otherwise specified.

Leishmania culture

L. major LV39 clone 5 WT (Rho/SU/59/P), *c14dm⁻* (*C14DM*-null mutant), *smt⁻*, *c14dm⁻/+C14DM* (episomal add-back), and *smt⁻/+SMT80* promastigotes were cultivated at 27 °C in complete M199 media (with 10% fetal bovine serum and additional supplements, pH 7.4) [70]. *Leishmania mexicana* (MNYC/BZ/62/M379) axenic amastigotes were cultured using an amastigote medium based on the Drosophila Schneider's medium supplemented with 20% fetal bovine serum and 0.0015% hemin (pH 5.5) in vented flasks in a humidified 32 °C/5% CO_2 incubator [71]. Culture density and cell viability were determined by hemacytometer counting and flow cytometry after propidium iodide (PI) staining, respectively as previously described [72]. In this study, log phase promastigotes refer to replicative parasites at 2.0-6.0 x 10^6 cells/ml, and stationary phase promastigotes refer to non-replicative parasites at densities higher than 2.0 x 10^7 cells/ml.

Determination of mitochondrial membrane potential (ΔΨm) and reactive oxygen species (ROS) level

469 ΔΨm was determined by flow cytometry after staining with TMRE or rhodamine 123 as

470 previously described [73]. Briefly, promastigotes were resuspended in complete M199, serum-

471 free M199, phosphate buffered saline (PBS, pH 7.2) containing 5.5 mM of glucose, or PBS alone

472 at 2.0 x 10^6 cells/ml; TMRE was added to a final concentration of 100 nM; after incubation at 27

473 °C for 15 min, the mean fluorescence intensity (MFI) was determined using an Attune® NxT

474 Flow Cytometer. Control samples include WT and *c14dm⁻* parasites pretreated with 75 µM of

475 carbonyl cyanide m-chlorophenyl hydrazone (CCCp) for 15 min and WT parasites grown in the

476 presence of 0.5 µM of ITZ for 48 hours.

477 Mitochondrial superoxide accumulation was determined as described previously with

478 minor modifications [33]. Promastigotes were resuspended in complete M199, serum-free M199,

479 PBS containing 5.5 mM of glucose, or PBS alone at 1.0 x 10^7 cells/ml and labeled with 5 μM of

480 MitoSOX red at 27 °C. After 25 min, MFI was measured by flow cytometry as described above.

481 To examine the level of cytosolic ROS, parasites were labeled with 5 µM of DHE for 30 min at

482 27 °C followed by flow cytometry analysis. Parasites pretreated with 5 µM of antimycin A, 75

483 μM of CCCp, or 100 μM of H_2O_2 for 15 min were used in these assays as controls. Such

484 treatments did not affect parasite viability based on PI-staining (<5% PI positive). *C14dm⁻*

485 mutants grown in the presence of 1 mM or 5 mM of glutathione (GSH) for 48 hours were also

486 included. In all flow cytometry experiments, 20,000 events were acquired for each sample.

487 **Determination of superoxide dismutase (SOD) activity**

488 SOD activity in whole cell extracts was measured as described previously for *L.*

489 *amazonensis* [74]. Briefly, 2 x 10^8 promastigotes were harvested, washed twice with PBS and

490 resuspended in a hypotonic buffer (5 mM of Tris–HCl, 0.1 mM of EDTA, and 1 x complete

491 EDTA-free protease inhibitor cocktail, pH 7.8) at a final concentration of 5 x 10^8 cells/ml. To

prepare lysates, cells were subjected to three freeze-thaw cycles alternating between liquid

nitrogen and a 37 °C water bath. Cell lysis was confirmed by microscopy. Lysates were clarified

by centrifugation (12,000 g for 30 min at 4 °C) and protein concentrations were determined

using a BCA protein assay kit (Thermo Fisher Scientific). SOD activity in whole cell extracts

was measured using the EnzyChrome superoxide dismutase assay kit (BioAssay Systems)

according to the manufacturer's protocol.

Digitonin fractionation and Western blot

Crude mitochondria-enriched fractions were generated by digitonin fractionation as

described previously [75]. Briefly, log phase promastigotes were incubated for 10 min on ice in

20 mM of Tris-HCl (pH 7.5), 0.6 M of sorbitol, 2 mM of EDTA and 0.025% (w/v) of digitonin

at 1 x 10^8 cells/ml at room temperature. As a positive control for mitochondrial membrane

disruption, WT parasites were treated with 0.035% (w/v) of digitonin at 37 °C. After

centrifugation (8000 g, for 30 min at 4 °C), the mitochondria-enriched pellet was separated from

the supernatant (enriched for cytosolic proteins). Fractions of equal cell equivalents were

subjected to SDS-PAGE and immunoblotting. Primary antibodies included rabbit-anti-*T. brucei*

cytochrome *c* (1:500 dilution) [32], rabbit-anti-*L. amazonensis* SODA and SODB (1:500), rabbit-

anti-*L. major* ISCL (1:500) [34], and rabbit-anti-*Leishmania* HSP83 (1:5000) antisera [7]. The

goat-anti-rabbit IgG conjugated with HRP (1:5000) was used as secondary antibody.

Measurement of oxygen consumption and ATP production

Oxygen consumption was determined using the MitoXpress Intracellular Oxygen Assay

kit (Luxcel Biosciences, Cork, Ireland) as described previously [45]. Briefly, log phase parasites

were washed once with PBS and resuspended in DMEM or a mitochondrial respiration buffer

(5.5 mM of sodium pyruvate, 5.5 mM of 2-deoxy-D-glucose in HBSS, pH 7.2) at 2.0 x 10^7 cells/

ml. For each sample, 100 µl of parasite suspension (corresponding to 2.0 x 10^6 cells) was added

in triplicates to a 96 black-well microtiter plate. WT parasites pre-treated with 10 µM of

antimycin A or 75 µM of CCCp were used as controls. 10 µl of MitoXpress (1 µM) was added to

each well. A layer of mineral oil (100 µl/well) was added over the cells to prevent diffusion of

atmospheric oxygen. MitoXpress signals were measured at 90 s intervals for 25-60 min using

excitation/emission wavelengths of 380 nm/650 nm in a BioTek synergy 4 fluorescence

microplate reader. For experiments in DMEM, we could not continue beyond 25 min when the

readings from WT cells reached the saturation point. Wells containing buffer only and buffer +

MitoXpress were included as blanks. For each cell line, the relative OCR was calculated by

subtracting the 0 min reading from the 15 min reading.

To measure ATP production, log phase parasites were washed once with PBS and

resuspended in an assay buffer containing 21 mM of HEPES, 0.7 mM of Na_2HPO4, and 137 mM

of NaCl (pH 7.2) in the presence or absence of glucose (5.5 mM), sodium pyruvate (5.5 mM), 2-

deoxy-D-glucose (5.5 mM) or sodium azide (20 mM) at 4.0 x 10^6 cells/ml. After incubation at

27 °C for one hour, 25 µl of suspension (corresponding to 1.0 x 10^5 cells) were transferred to a

96 black-well plate, mixed with equal volume of the CellTiter-Glo reagent and incubated for 10

min in the dark. Bioluminescence was measured using a DTX880 microplate reader as described

[23]. For each experiment, an ATP standard curve was prepared by diluting pure ATP stock in

the assay buffer.

Microscopy

To stain mitochondria, live parasites were resuspended in DMEM at 1.0×10^6 cells/ml

and labeled with 100 nM of Mitotracker CMXRos or TMRE. After incubation at 27 °C (30 min

for Mitotracker and 15 min for TRME), parasites were washed once with PBS, fixed and

processed for fluorescence microscopy as previously described [15]. MitoSox Red labeling (5

µM for 25 min at 27 °C) of live parasites was performed similarly without fixation using an

Olympus BX51 Upright Fluorescence Microscope. DNA was stained with DAPI (2.5 µg/ml).

Transmission electron microscopy of log and stationary phase promastigotes was performed as

previously described [76].

Starvation and heat stress assays

To measure heat sensitivity, log phase promastigotes were incubated in complete M199

media for 0-24 h at either 27 °C or 37 °C/5% CO_2. Effects of antimycin A (10 µM) and GSH (1

mM or 5 mM) on heat sensitivity were also monitored. For starvation response, log phase

promastigotes were washed once with HBSS and resuspended in HBSS supplemented with

glucose (5.5 mM), 2-deoxy-D-glucose (5.5 mM), aspartate (5.5 mM), sodium pyruvate (5.5 mM),

or glycerol (0.1 mM) at 2.5 x 10^6 cells/ml and incubated for 0-24 h at 27 °C. In these stress

assays, culture densities and percentages of dead cells were determined over time as previously

described [72].

Glucose and glycerol uptake assays

Uptake of glucose and glycerol were monitored as previously described with minor

modifications [53, 77]. For glucose incorporation, log phase promastigotes were washed twice

with PBS and resuspended at 5.0 x 10^7 cells/ml in PBS in the presence of 100 µM of D-[U-^{14}C]

glucose (250 mCi/mmol, PerkinElmer Inc.). The reaction mixture (200 µl each in triplicates) was

immediately loaded onto 100 µl of dibutyl phthalate in 1.5 ml Eppendorf tubes. Incubation was

performed at room temperature and terminated at the indicated times by pelleting cells through

the oil layer through centrifugation (16,000 g, 1 min). Tubes were then flash frozen with liquid

nitrogen and the bottoms (containing cell pellets) were clipped off and submerged in scintillation

fluid containing 1% SDS for 24 h, followed by scintillation counting. To measure glycerol

uptake, log phase promastigotes were washed twice in a transport buffer (33 mM of HEPES, 98

mM of NaCl, 4.6 mM of KCl, 0.55 mM of $CaCl_2$, 0.07 mM of $MgSO_4$, 5.8 mM of Na_2PO4, 0.3

mM of $NaHCO_3$, and 14 mM of glucose, pH 7.3) and resuspended at 1.0×10^8 cells/ml in the

same buffer. Uptake was measured at room temperature by mixing 100 µl of cells with 100 µl of

transport buffer containing 8 µM of [^{14}C] glycerol (0.125 µCi each in triplicates, PerkinElmer

Inc.). Incorporation of radioactivity was determined as described above.

Susceptibility to pentamidine

To determine the effective concentration of pentamidine to inhibit parasite growth by

90%, 50% and 25% (EC90, EC50 and EC25), log phase promastigotes were inoculated in

complete M199 media at 2.0×10^5 cells/ml in 0-25 µM of pentamidine. Culture densities were

measured after 48 hours of incubation in 24-well plates. EC90, EC50 and EC25 were determined

using cells grown in the absence of pentamidine as controls. To assess the leishmanicidal activity

of pentamidine in short time exposure, log phase promastigotes were inoculated in complete

M199 media containing 0, 2.5 µM, or 25 µM of pentamidine. For *c14dm⁻*, half of them were also

treated with 75 µM of CCCp. After one hour, cells were washed twice with PBS and re-

inoculated in complete M199 media at 2.0×10^5 cells/ml. Culture densities were then determined

daily for six days.

Statistical analysis

Unless specified otherwise, assay values in all figures are averaged from three to five

independent repeats and error bars represent standard deviations. Differences were assessed by

the Student's t test (for two groups) or one-way ANOVA (for three or more groups) using the

Sigmaplot 13.0 software (Systat Software Inc, San Jose, CA). *P* values indicating statistical

significance were grouped in all figures (***: $p < 0.001$, **: $p < 0.01$, *: $p < 0.05$).

ACKNOWLEDGMENTS

587   We thank Dr. Norma Andrews (University of Maryland College Park) for providing us

588 the rabbit anti-*L. amazonensis* SODA and SODB antisera, Dr. Wandy Beatty (Washington

589 University in St. Louis) for transmission electron microscopy, and Dr. Catherine Wakeman for

590 usage on the BioTek synergy 4 fluorescence microplate reader.

REFERENCES

1.	Alvar J, Velez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis worldwide and global estimates of its incidence. PLoS One. 2012;7(5):e35671. Epub 2012/06/14. doi: 10.1371/journal.pone.0035671. PubMed PMID: 22693548; PubMed Central PMCID: PMC3365071.

2.	Ghorbani M, Farhoudi R. Leishmaniasis in humans: drug or vaccine therapy? Drug Des Devel Ther. 2018;12:25-40. Epub 2018/01/11. doi: 10.2147/DDDT.S146521. PubMed PMID: 29317800; PubMed Central PMCID: PMC5743117.

3.	Croft SL, Sundar S, Fairlamb AH. Drug resistance in leishmaniasis. Clin Microbiol Rev. 2006;19(1):111-26. PubMed PMID: 16418526.

4.	Goad LJ, Holz GG, Jr., Beach DH. Sterols of *Leishmania* species. Implications for biosynthesis. Mol Biochem Parasitol. 1984;10(2):161-70. Epub 1984/02/01. PubMed PMID: 6700638.

5.	Urbina JA. Lipid biosynthesis pathways as chemotherapeutic targets in kinetoplastid parasites. Parasitology. 1997;114 Suppl:S91-9. PubMed PMID: 9309771.

6.	Urbina JA, Vivas J, Lazardi K, Molina J, Payares G, Piras MM, et al. Antiproliferative effects of delta 24(25) sterol methyl transferase inhibitors on *Trypanosoma (Schizotrypanum) cruzi*: in vitro and in vivo studies. Chemotherapy. 1996;42(4):294-307. PubMed PMID: 8804798.

7.	Xu W, Hsu FF, Baykal E, Huang J, Zhang K. Sterol Biosynthesis Is Required for Heat Resistance but Not Extracellular Survival in *Leishmania*. PLoS Patho. 2014;10(10):e1004427. Epub 2014/10/24. doi: 10.1371/journal.ppat.1004427. PubMed PMID: 25340392.

8.   de Souza W, Rodrigues JC. Sterol Biosynthesis Pathway as Target for Anti-

trypanosomatid Drugs. Interdiscip Perspect Infect Dis. 2009;2009:642502. Epub 2009/08/15. doi:

10.1155/2009/642502. PubMed PMID: 19680554; PubMed Central PMCID: PMC2721973.

9.   de Macedo-Silva ST, de Souza W, Rodrigues JC. Sterol Biosynthesis Pathway as an

Alternative for the Anti-Protozoan Parasite Chemotherapy. Curr Med Chem. 2015;22(18):2186-

98. Epub 2015/03/20. PubMed PMID: 25787966.

10.   Mosimann V, Neumayr A, Paris DH, Blum J. Liposomal amphotericin B treatment of

Old World cutaneous and mucosal leishmaniasis: A literature review. Acta Trop. 2018;182:246-

50. Epub 2018/03/20. doi: 10.1016/j.actatropica.2018.03.016. PubMed PMID: 29550282.

11.   Jefcoate CR, Gaylor JL, Calabrese RL. Ligand interactions with cytochrome P-450. I.

Binding of primary amines. Biochemistry. 1969;8(8):3455-63. Epub 1969/08/01. PubMed PMID:

5809236.

12.   Warrilow AG, Parker JE, Kelly DE, Kelly SL. Azole affinity of sterol 14alpha-

demethylase (CYP51) enzymes from *Candida albicans* and *Homo sapiens*. Antimicrob Agents

Chemother. 2013;57(3):1352-60. Epub 2013/01/01. doi: 10.1128/AAC.02067-12. PubMed

PMID: 23274672; PubMed Central PMCID: PMC3591892.

13.   McCall LI, El Aroussi A, Choi JY, Vieira DF, De Muylder G, Johnston JB, et al.

Targeting Ergosterol biosynthesis in *Leishmania donovani*: essentiality of sterol 14 alpha-

demethylase. PLoS Negl Trop Dis. 2015;9(3):e0003588. Epub 2015/03/15. doi:

10.1371/journal.pntd.0003588. PubMed PMID: 25768284; PubMed Central PMCID:

PMC4359151.

14.   McConville MJ, Thomas-Oates JE, Ferguson MA, Homans SW. Structure of the

lipophosphoglycan from *Leishmania major*. J Biol Chem. 1990;265(32):19611-23.

15.   Mukherjee S, Xu W, Hsu FF, Patel J, Huang J, Zhang K. Sterol methyltransferase is

required for optimal mitochondrial function and virulence in *Leishmania major*. Mol Microbiol.

2019;111(1):65-81. doi: 10.1111/mmi.14139. PubMed PMID: 30260041; PubMed Central

PMCID: PMC6351164.

16.   Dufourc EJ. Sterols and membrane dynamics. J Chem Biol. 2008;1(1-4):63-77. Epub

2009/07/02. doi: 10.1007/s12154-008-0010-6. PubMed PMID: 19568799; PubMed Central

PMCID: PMC2698314.

17.   Barenholz Y. Cholesterol and other membrane active sterols: from membrane evolution

to "rafts". Prog Lipid Res. 2002;41(1):1-5. Epub 2001/11/06. PubMed PMID: 11694266.

18.   Zinser E, Paltauf F, Daum G. Sterol composition of yeast organelle membranes and

subcellular distribution of enzymes involved in sterol metabolism. J Bacteriol.

1993;175(10):2853-8. Epub 1993/05/01. PubMed PMID: 8491706; PubMed Central PMCID:

PMC204601.

19.   van Meer G, de Kroon AI. Lipid map of the mammalian cell. J Cell Sci. 2011;124(Pt

1):5-8. Epub 2010/12/22. doi: 10.1242/jcs.071233. PubMed PMID: 21172818.

20.   Maxfield FR, Menon AK. Intracellular sterol transport and distribution. Curr Opin Cell

Biol. 2006;18(4):379-85. Epub 2006/06/30. doi: 10.1016/j.ceb.2006.06.012. PubMed PMID:

16806879.

21.   Rodrigues CO, Catisti R, Uyemura SA, Vercesi AE, Lira R, Rodriguez C, et al. The

sterol composition of *Trypanosoma cruzi* changes after growth in different culture media and

results in different sensitivity to digitonin-permeabilization. J Eukaryot Microbiol.

2001;48(5):588-94. Epub 2001/10/13. PubMed PMID: 11596924.

22. Kessler RL, Soares MJ, Probst CM, Krieger MA. *Trypanosoma cruzi* response to sterol

biosynthesis inhibitors: morphophysiological alterations leading to cell death. PLoS One.

2013;8(1):e55497. Epub 2013/02/06. doi: 10.1371/journal.pone.0055497. PubMed PMID:

23383204; PubMed Central PMCID: PMC3561218.

23. Perez-Moreno G, Sealey-Cardona M, Rodrigues-Poveda C, Gelb MH, Ruiz-Perez LM,

Castillo-Acosta V, et al. Endogenous sterol biosynthesis is important for mitochondrial function

and cell morphology in procyclic forms of *Trypanosoma brucei*. Int J Parasitol.

2012;42(11):975-89. Epub 2012/09/12. doi: 10.1016/j.ijpara.2012.07.012. PubMed PMID:

22964455.

24. Vannier-Santos MA, Urbina JA, Martiny A, Neves A, de Souza W. Alterations induced

by the antifungal compounds ketoconazole and terbinafine in *Leishmania*. J Eukaryot Microbiol.

1995;42(4):337-46. Epub 1995/07/01. PubMed PMID: 7620457.

25. de Macedo-Silva ST, Urbina JA, de Souza W, Rodrigues JC. In vitro activity of the

antifungal azoles itraconazole and posaconazole against *Leishmania amazonensis*. PLoS One.

2013;8(12):e83247. Epub 2014/01/01. doi: 10.1371/journal.pone.0083247. PubMed PMID:

24376670; PubMed Central PMCID: PMC3871555.

26. Rodrigues JC, Attias M, Rodriguez C, Urbina JA, Souza W. Ultrastructural and

biochemical alterations induced by 22,26-azasterol, a delta(24(25))-sterol methyltransferase

inhibitor, on promastigote and amastigote forms of *Leishmania amazonensis*. Antimicrob Agents

Chemother. 2002;46(2):487-99. PubMed PMID: 11796362.

27. Rodrigues JC, Bernardes CF, Visbal G, Urbina JA, Vercesi AE, de Souza W. Sterol

methenyl transferase inhibitors alter the ultrastructure and function of the *Leishmania*

amazonensis mitochondrion leading to potent growth inhibition. Protist. 2007;158(4):447-56.

PubMed PMID: 17719843.

28.   Orlandi PA, Jr., Turco SJ. Structure of the lipid moiety of the *Leishmania donovani*

lipophosphoglycan. J Biol Chem. 1987;262(21):10384-91.

29.   Stephens JL, Lee SH, Paul KS, Englund PT. Mitochondrial fatty acid synthesis in

Trypanosoma brucei. J Biol Chem. 2007;282(7):4427-36. Epub 2006/12/15. doi:

10.1074/jbc.M609037200. PubMed PMID: 17166831.

30.   Tomas AM, Castro H. Redox metabolism in mitochondria of trypanosomatids. Antioxid

Redox Signal. 2013;19(7):696-707. Epub 2012/10/03. doi: 10.1089/ars.2012.4948. PubMed

PMID: 23025438; PubMed Central PMCID: PMC3739956.

31.   Gros L, Lorente SO, Jimenez CJ, Yardley V, Rattray L, Wharton H, et al. Evaluation of

azasterols as anti-parasitics. J Med Chem. 2006;49(20):6094-103. Epub 2006/09/29. doi:

10.1021/jm060290f. PubMed PMID: 17004723.

32.   Dolai S, Pal S, Yadav RK, Adak S. Endoplasmic reticulum stress-induced apoptosis in

Leishmania through Ca2+-dependent and caspase-independent mechanism. J Biol Chem.

2011;286(15):13638-46. Epub 2011/02/19. doi: 10.1074/jbc.M110.201889. PubMed PMID:

21330370; PubMed Central PMCID: PMC3075708.

33.   Kathuria M, Bhattacharjee A, Sashidhara KV, Singh SP, Mitra K. Induction of

mitochondrial dysfunction and oxidative stress in *Leishmania donovani* by orally active

clerodane diterpene. Antimicrob Agents Chemother. 2014;58(10):5916-28. Epub 2014/07/30. doi:

10.1128/AAC.02459-14. PubMed PMID: 25070112; PubMed Central PMCID: PMC4187897.

34.   Zhang O, Xu, W., Pillai, A., Zhang, K. Developmentally Regulated Sphingolipid

Degradation in *Leishmania major*. PLoS One. 2012;7(1):e31059.

35.   Shapira M, Pinelli E. Heat-shock protein 83 of *Leishmania mexicana amazonensis* is an

abundant cytoplasmic protein with a tandemly repeated genomic arrangement. Eur J Biochem.

1989;185(2):231-6. Epub 1989/11/06. PubMed PMID: 2684665.

36.   Hide M, Ritleng AS, Brizard JP, Monte-Allegre A, Sereno D. *Leishmania infantum*:

tuning digitonin fractionation for comparative proteomic of the mitochondrial protein content.

Parasitol Res. 2008;103(4):989-92. Epub 2008/06/26. doi: 10.1007/s00436-008-1062-9. PubMed

PMID: 18575891.

37.   Drose S, Brandt U. Molecular mechanisms of superoxide production by the

mitochondrial respiratory chain. Adv Exp Med Biol. 2012;748:145-69. Epub 2012/06/26. doi:

10.1007/978-1-4614-3573-0_6. PubMed PMID: 22729857.

38.   Kowaltowski AJ, de Souza-Pinto NC, Castilho RF, Vercesi AE. Mitochondria and

reactive oxygen species. Free Radic Biol Med. 2009;47(4):333-43. Epub 2009/05/12. doi:

10.1016/j.freeradbiomed.2009.05.004. PubMed PMID: 19427899.

39.   Liu Y, Fiskum G, Schubert D. Generation of reactive oxygen species by the

mitochondrial electron transport chain. J Neurochem. 2002;80(5):780-7. Epub 2002/04/12.

PubMed PMID: 11948241.

40.   Moreira W, Leprohon P, Ouellette M. Tolerance to drug-induced cell death favours the

acquisition of multidrug resistance in *Leishmania*. Cell Death Dis. 2011;2:e201. Epub

2011/09/02. doi: 10.1038/cddis.2011.83. PubMed PMID: 21881603; PubMed Central PMCID:

PMC3186901.

41.   Brand MD, Affourtit C, Esteves TC, Green K, Lambert AJ, Miwa S, et al. Mitochondrial

superoxide: production, biological effects, and activation of uncoupling proteins. Free Radic Biol

Med. 2004;37(6):755-67. Epub 2004/08/12. doi: 10.1016/j.freeradbiomed.2004.05.034. PubMed

PMID: 15304252.

42.   Fairlamb AH, Cerami A. Metabolism and functions of trypanothione in the

Kinetoplastida. Annu Rev Microbiol. 1992;46:695-729. Epub 1992/01/01. doi:

10.1146/annurev.mi.46.100192.003403. PubMed PMID: 1444271.

43.   Mittra B, Laranjeira-Silva MF, Miguel DC, Perrone Bezerra de Menezes J, Andrews NW.

The iron-dependent mitochondrial superoxide dismutase SODA promotes *Leishmania* virulence.

J Biol Chem. 2017;292(29):12324-38. Epub 2017/05/28. doi: 10.1074/jbc.M116.772624.

PubMed PMID: 28550086; PubMed Central PMCID: PMC5519379.

44.   Plewes KA, Barr SD, Gedamu L. Iron superoxide dismutases targeted to the glycosomes

of *Leishmania chagasi* are important for survival. Infect Immun. 2003;71(10):5910-20. Epub

2003/09/23. doi: 10.1128/iai.71.10.5910-5920.2003. PubMed PMID: 14500512; PubMed Central

PMCID: PMC201062.

45.   Cardenas D, Carter PM, Nation CS, Pizarro JC, Guidry J, Aiyar A, et al. LACK, a

RACK1 ortholog, facilitates cytochrome *c* oxidase subunit expression to promote *Leishmania*

major fitness. Mol Microbiol. 2015;96(1):95-109. Epub 2015/01/15. doi: 10.1111/mmi.12924.

PubMed PMID: 25582232; PubMed Central PMCID: PMC6055511.

46.   Manzano JI, Carvalho L, Perez-Victoria JM, Castanys S, Gamarro F. Increased glycolytic

ATP synthesis is associated with tafenoquine resistance in *Leishmania major*. Antimicrob

Agents Chemother. 2011;55(3):1045-52. Epub 2011/01/05. doi: 10.1128/AAC.01545-10.

PubMed PMID: 21199921; PubMed Central PMCID: PMC3067095.

47.   Guidarelli A, Brambilla L, Rota C, Tomasi A, Cattabeni F, Cantoni O. The respiratory-

chain poison antimycin A promotes the formation of DNA single-strand breaks and reduces

toxicity in U937 cells exposed to t-butylhydroperoxide. Biochem J. 1996;317 (Pt 2):371-5. Epub

1996/07/15. doi: 10.1042/bj3170371. PubMed PMID: 8713060; PubMed Central PMCID:

PMC1217497.

48.   Sanchez MA. Molecular identification and characterization of an essential pyruvate

transporter from *Trypanosoma brucei*. J Biol Chem. 2013;288(20):14428-37. Epub 2013/04/10.

doi: 10.1074/jbc.M113.473157. PubMed PMID: 23569205; PubMed Central PMCID:

PMC3656298.

49.   Leary SC, Hill BC, Lyons CN, Carlson CG, Michaud D, Kraft CS, et al. Chronic

treatment with azide in situ leads to an irreversible loss of cytochrome *c* oxidase activity via

holoenzyme dissociation. J Biol Chem. 2002;277(13):11321-8. Epub 2002/01/18. doi:

10.1074/jbc.M112303200. PubMed PMID: 11796734.

50.   Pineda E, Thonnus M, Mazet M, Mourier A, Cahoreau E, Kulyk H, et al. Glycerol

supports growth of the *Trypanosoma brucei* bloodstream forms in the absence of glucose:

Analysis of metabolic adaptations on glycerol-rich conditions. PLoS Patho.

2018;14(11):e1007412. Epub 2018/11/02. doi: 10.1371/journal.ppat.1007412. PubMed PMID:

30383867; PubMed Central PMCID: PMC6245841.

51.   Kovarova J, Nagar R, Faria J, Ferguson MAJ, Barrett MP, Horn D. Gluconeogenesis

using glycerol as a substrate in bloodstream-form *Trypanosoma brucei*. PLoS Patho.

2018;14(12):e1007475. Epub 2018/12/28. doi: 10.1371/journal.ppat.1007475. PubMed PMID:

30589893; PubMed Central PMCID: PMC6307712.

52.   Feng X, Tran KD, Sanchez MA, Al Mezewghi H, Landfear SM. Glucose Transporters

and Virulence in *Leishmania mexicana*. mSphere. 2018;3(4). Epub 2018/08/03. doi:

10.1128/mSphere.00349-18. PubMed PMID: 30068561; PubMed Central PMCID:

PMC6070740.

53.   Jeacock L, Baker N, Wiedemar N, Maser P, Horn D. Aquaglyceroporin-null

trypanosomes display glycerol transport defects and respiratory-inhibitor sensitivity. PLoS Patho.

2017;13(3):e1006307. Epub 2017/03/31. doi: 10.1371/journal.ppat.1006307. PubMed PMID:

28358927; PubMed Central PMCID: PMC5388498.

54.   Burchmore RJ, Rodriguez-Contreras D, McBride K, Merkel P, Barrett MP, Modi G, et al.

Genetic characterization of glucose transporter function in *Leishmania mexicana*. Proc Natl Acad

Sci U S A. 2003;100(7):3901-6. PubMed PMID: 12651954.

55.   Basselin M, Denise H, Coombs GH, Barrett MP. Resistance to pentamidine in

Leishmania mexicana involves exclusion of the drug from the mitochondrion. Antimicrob

Agents Chemother. 2002;46(12):3731-8. Epub 2002/11/19. doi: 10.1128/aac.46.12.3731-

3738.2002. PubMed PMID: 12435669; PubMed Central PMCID: PMC132791.

56.   Mukherjee A, Padmanabhan PK, Sahani MH, Barrett MP, Madhubala R. Roles for

mitochondria in pentamidine susceptibility and resistance in *Leishmania donovani*. Mol Biochem

Parasitol. 2006;145(1):1-10. Epub 2005/10/13. doi: 10.1016/j.molbiopara.2005.08.016. PubMed

PMID: 16219371.

57.   Kaminsky R, Mamman M, Chuma F, Zweygarth E. Time-dose-response of *Trypanosoma*

brucei brucei to diminazene aceturate (Berenil) and in vitro simulation of drug-concentration-

time profiles in cattle plasma. Acta Trop. 1993;54(1):19-30. Epub 1993/06/01. doi:

10.1016/0001-706x(93)90065-j. PubMed PMID: 8103625.

58.   Mesmin B, Maxfield FR. Intracellular sterol dynamics. Biochim Biophys Acta.

2009;1791(7):636-45. Epub 2009/03/17. doi: 10.1016/j.bbalip.2009.03.002. PubMed PMID:

19286471; PubMed Central PMCID: PMC2696574.

59.   Figarella K, Marsiccobetre S, Arocha I, Colina W, Hasegawa M, Rodriguez M, et al.

Ergosterone-coupled Triazol molecules trigger mitochondrial dysfunction, oxidative stress, and

acidocalcisomal Ca(2+) release in *Leishmania mexicana* promastigotes. Microb Cell.

2015;3(1):14-28. Epub 2015/12/11. doi: 10.15698/mic2016.01.471. PubMed PMID: 28357313;

PubMed Central PMCID: PMC5354587.

60.   van Meer G, Lisman Q. Sphingolipid transport: rafts and translocators. J Biol Chem.

2002;277(29):25855-8. PubMed PMID: 12011105.

61.   Horvath SE, Daum G. Lipids of mitochondria. Prog Lipid Res. 2013;52(4):590-614.

Epub 2013/09/07. doi: 10.1016/j.plipres.2013.07.002. PubMed PMID: 24007978.

62.   Joshi AS, Thompson MN, Fei N, Huttemann M, Greenberg ML. Cardiolipin and

mitochondrial phosphatidylethanolamine have overlapping functions in mitochondrial fusion in

Saccharomyces cerevisiae. J Biol Chem. 2012;287(21):17589-97. Epub 2012/03/22. doi:

10.1074/jbc.M111.330167. PubMed PMID: 22433850; PubMed Central PMCID: PMC3366806.

63.   Basu Ball W, Neff JK, Gohil VM. The role of nonbilayer phospholipids in mitochondrial

structure and function. FEBS Lett. 2018;592(8):1273-90. Epub 2017/10/27. doi: 10.1002/1873-

3468.12887. PubMed PMID: 29067684; PubMed Central PMCID: PMC5918238.

64.   Hardie DG, Scott JW, Pan DA, Hudson ER. Management of cellular energy by the AMP-

activated protein kinase system. FEBS Lett. 2003;546(1):113-20. Epub 2003/06/28. PubMed

PMID: 12829246.

65.   Carling D. The AMP-activated protein kinase cascade--a unifying system for energy

control. Trends Biochem Sci. 2004;29(1):18-24. Epub 2004/01/20. doi:

10.1016/j.tibs.2003.11.005. PubMed PMID: 14729328.

66.   Chen L, Wang J, Zhang YY, Yan SF, Neumann D, Schlattner U, et al. AMP-activated

protein kinase undergoes nucleotide-dependent conformational changes. Nat Struct Mol Biol.

2012;19(7):716-8. Epub 2012/06/05. doi: 10.1038/nsmb.2319. PubMed PMID: 22659875.

67.   Roy A, Ganguly A, BoseDasgupta S, Das BB, Pal C, Jaisankar P, et al. Mitochondria-

dependent reactive oxygen species-mediated programmed cell death induced by 3,3'-

diindolylmethane through inhibition of F0F1-ATP synthase in unicellular protozoan parasite

Leishmania donovani. Mol Pharmacol. 2008;74(5):1292-307. PubMed PMID: 18703668.

68.   Mwenechanya R, Kovarova J, Dickens NJ, Mudaliar M, Herzyk P, Vincent IM, et al.

Sterol 14alpha-demethylase mutation leads to amphotericin B resistance *in Leishmania mexicana*.

PLoS Negl Trop Dis. 2017;11(6):e0005649. Epub 2017/06/18. doi:

10.1371/journal.pntd.0005649. PubMed PMID: 28622334; PubMed Central PMCID:

PMC5498063.

69.   Pountain AW, Weidt SK, Regnault C, Bates PA, Donachie AM, Dickens NJ, et al.

Genomic instability at the locus of sterol C24-methyltransferase promotes amphotericin B

resistance in *Leishmania parasites*. PLoS Negl Trop Dis. 2019;13(2):e0007052. Epub

2019/02/05. doi: 10.1371/journal.pntd.0007052. PubMed PMID: 30716073; PubMed Central

PMCID: PMC6375703.

70.   Kapler GM, Coburn CM, Beverley SM. Stable transfection of the human parasite

Leishmania major delineates a 30-kilobase region sufficient for extrachromosomal replication

and expression. Mol Cell Biol. 1990;10(3):1084-94. PubMed PMID: 2304458.

71.   Bates PA, Robertson CD, Tetley L, Coombs GH. Axenic cultivation and characterization

of *Leishmania mexicana* amastigote-like forms. Parasitology. 1992;105 (Pt 2):193-202. Epub

1992/10/01. PubMed PMID: 1454417.

72.   Zhang K, Showalter M, Revollo J, Hsu FF, Turk J, Beverley SM. Sphingolipids are

essential for differentiation but not growth in *Leishmania*. EMBO J. 2003;22(22):6016-26.

PMCID: 275442. PubMed PMID: 14609948.

73.   Vergnes B, Gourbal B, Girard I, Sundar S, Drummelsmith J, Ouellette M. A proteomics

screen implicates HSP83 and a small kinetoplastid calpain-related protein in drug resistance in

Leishmania donovani clinical field isolates by modulating drug-induced programmed cell death.

Mol Cell Proteomics. 2007;6(1):88-101. Epub 2006/10/20. doi: 10.1074/mcp.M600319-MCP200.

PubMed PMID: 17050524.

74.   Mittra B, Laranjeira-Silva MF, Perrone Bezerra de Menezes J, Jensen J, Michailowsky V,

Andrews NW. A Trypanosomatid Iron Transporter that Regulates Mitochondrial Function Is

Required for *Leishmania amazonensis* Virulence. PLoS Patho. 2016;12(1):e1005340. Epub

2016/01/08. doi: 10.1371/journal.ppat.1005340. PubMed PMID: 26741360; PubMed Central

PMCID: PMC4704735.

75.   Kaser S, Oeljeklaus S, Tyc J, Vaughan S, Warscheid B, Schneider A. Outer membrane

protein functions as integrator of protein import and DNA inheritance in mitochondria. Proc Natl

Acad Sci U S A. 2016;113(31):E4467-75. Epub 2016/07/21. doi: 10.1073/pnas.1605497113.

PubMed PMID: 27436903; PubMed Central PMCID: PMC4978248.

76.   Zhang K, Hsu FF, Scott DA, Docampo R, Turk J, Beverley SM. *Leishmania* salvage and

remodelling of host sphingolipids in amastigote survival and acidocalcisome biogenesis. Mol

Microbiol. 2005;55(5):1566-78. PubMed PMID: 15720561.

77.   Rodriguez-Contreras D, Landfear SM. Metabolic changes in glucose transporter-deficient

Leishmania mexicana and parasite virulence. J Biol Chem. 2006;281(29):20068-76. PubMed

PMID: 16707495.

 FIGURE CAPTIONS

Fig 1. *C14dm⁻* promastigotes show high mitochondrial membrane potential (ΔΨm). (A) Log

phase promastigotes were labeled with 100 nM of Mitotracker CMXRos followed by DAPI

staining. WT parasites pretreated with CCCp were included as controls. DIC: differential

interference contrast. (**B**) Representative transmission electron micrographs showing

mitochondria in log phase promastigotes. Black arrows indicate areas containing kinetoplast (top

panel) and tubular mitochondria (bottom panel). (**C**) To measure ΔΨm, log phase and stationary

phase (day 1-day 4) promastigotes were labelled with 100 nM of TMRE for 15 min in PBS and

mean fluorescence intensity (MFI) were determined by flow cytometry. Controls include WT

and *c14dm⁻* parasites pretreated with CCCp and WT cells pretreated with 0.2 µM of ITZ. Bar

values represent averages from three experiments and error bars represent standard deviations.

(**D**) Live parasites imaging after TMRE labeling.

Fig 2. Inactivation of C14DM leads to ROS accumulation in the mitochondria. (A, **B** and **F**)

Promastigotes were incubated in complete M199, serum-free M199, PBS contain 5.5 mM of

glucose, or PBS only and labeled with 100 nM of TMRE (**A**, for 15 min), 5 µM of MitoSox Red

(**B**, for 25 min) or 5 µM of DHE (**F**, for 30 min). MFIs were determined by flow cytometry.

Effects of CCCp (75 µM, 15 min) were also monitored in **A** and **B**. (**C**) Log or stationary phase

(day 1-day 4) promastigotes were labeled with 5 µM of MitoSox Red for 25 min in PBS and

MFIs were determined by flow cytometry and normalized to values from log phase WT. WT

cells grown in the presence of ITZ (for 48 hours) were also analyzed. (**D** and **E**) Live cell

imaging of *L. major* promastigotes (**D**) or *L. mexicana* axenic amastigotes (grown in the absence

or presence of 0.1 µM of ITZ for 48 hours, **E**) after labelling with MitoSox Red in PBS. White

arrows indicate kDNA binding of MitoSox. (**E**) Live after labeling with MitoSox Red. In **A-C**

and **F**, averaged values from five experiments are shown and error bars represent standard

deviations.

Fig 3. Increased superoxide dismutase (SOD) activity in *c14dmˉ* parasites. (**A**) Whole cell

lysates from log phase and stationary phase (day 1-day 3) promastigotes were collected as

described in **Materials and Methods**. SOD activity was measured using a SOD assay kit and

results were normalized based on the amount of protein in each sample. Error bars represent

standard deviations from two independent experiments. (**B**) Mitochondria enriched membrane

fractions (Mem.) and cytosolic fractions (Sup.) were isolated from log phase promastigotes as

described. Western blots were performed using antibodies against *L. amazonensis* SODA

(mitochondrial) and SODB (both SODB1 and SODB2, glycosomal).

Fig 4. *C14dmˉ* parasites show compromised cellular respiration. (**A-C**) Log phase

promastigotes were incubated in a mitochondrial respiration buffer (**A**-**B**) or DMEM (**C**) and

oxygen consumption rates (OCR) were determined using the MitoXpress probe as described in

Materials and Methods. Buffer only and buffer with probe (no cells) were included as blanks

(**A**). In **A** and **C**, fluorescence signals (in arbitrary units or A.U.) were measured every 90 s. In **B**,

the relative OCR from the experiments in **A** was calculated by subtracting the 0 min reading

from the 15 min reading. WT parasites treated with antimycin A (AA) were included as controls.

(**D**) Cellular ATP contents in log phase promastigotes were measured after 1 hour incubation in

an assay buffer (21 mM of HEPES, 0.7 mM of Na_2HPO4, and 137 mM of NaCl, pH 7.2) in the

absence or presence of glucose (Glu, 5.5 mM), sodium pyruvate (Pyr, 5.5 mM), 2-deoxy-D-

glucose (2DG, 5.5 mM), or sodium azide (NaN_3, 20 mM). All experiments were repeated three

or four times and error bars represent standard deviations.

Fig 5. *C14dm⁻* parasites are hypersensitive to starvation. Log phase promastigotes were inoculated in HBSS in the absence or presence of 2DG (5.5 mM), Pyr (5.5 mM), aspartate (Asp, 5.5 mM), Glu (5.5 mM), or glycerol (0.1 mM). In **A**, percentages of cell death for *c14dm⁻* mutants were determined at the indicated time points. In **B**, percentages of cell death were determined after 24 hours for WT, *c14dm⁻, c14dm⁻/+C14DM, smt⁻,* and *smt⁻/+SMT* parasites. Error bars represent standard deviations from three independent experiments.

Fig 6. *C14dm⁻* parasites show increased uptake of glucose and glycerol. Incorporation of radiolabeled glucose (**A**) or glycerol (**B** and **C**) were determined as described under **Materials and Methods**. Transport assays were performed at room temperature (**A** and **B** unless specified) or 4 °C (**C**). Error bars represent standard deviations from three independent experiments.

Fig 7. ROS accumulation contributes to heat sensitivity in *c14dm⁻* mutants. (**A**-**E**) Log phase promastigotes were incubated in PBS at 27 °C or 37 °C for 3 hours. In **A**, percentages of cell death were determined by flow cytometry after PI staining. In **B**, ΔΨm was determined after TMRE labeling. Unlabeled WT parasites were included as a negative control (Neg). In **C**-**D**, parasites were lysed and cytosolic fractions (Sup.) were separated from mitochondrial fractions (Mem.) as described in **Materials and Methods**. Distribution of cytochrome *c* and HSP83 (a cytosolic protein marker) were determined by Western blots (**C**) and quantified (**D**). In **E**, parasites were labelled with the cytosolic ROS indicator DHE. Fluorescence intensities were measured by flow cytometry and relative cytosolic ROS levels were determined. (**F**) Log phase promastigotes were incubated at 37 °C in the absence or presence of reduced GSH. After 8 hours, mitochondrial ROS levels were measured as described. (**G**) *C14dm⁻* parasites were incubated at 37 °C in the absence or presence of reduced GSH and percentages of cell death were determined at 0 and 8 hours. (**H**) WT parasites were incubated at 27 °C or 37 °C in the absence or presence

of AA (10 µM) and percentages of dead cells were determined by flow cytometry at the

indicated time points. †: Very few WT+AA cells were detectable after 36 hours at 37 °C. Error

bars represent standard deviations from three independent experiments.

Fig 8. *C14dm⁻* parasites show increased sensitivity to pentamidine (PEN). (**A**) Log phase

promastigotes were inoculated in M199 media containing various concentrations of PEN.

Culture densities were determined after 48 hours and the PEN concentrations required to inhibit

growth by 90%, 50%, and 25% were determined (as EC90, EC50 and EC25 respectively) using

cells grown in the absence of PEN as controls. "WT + ITZ" represent WT parasites grown in the

presence of 0.5 µM of ITZ. (**B**-**C**) Log phase promastigotes were inoculated in M199 media

containing 2.5 µM (**B**) or 25 µM (**C**) of PEN for one hour, followed two washes with PBS. Cells

were then re-inoculated in M199 media and culture densities were determined every 24 hours.

Some *c14dm⁻* mutants were also incubated in the presence of 75 µM of CCCp along with PEN

(for one hour). Parasites originally inoculated in the absence of PEN were included as controls.

Error bars represent standard deviations from three or four independent experiments.

SUPPLEMENTAL FIGURE CAPTIONS

Fig S1. *C14dm⁻* promastigotes show increased mitochondrial surface area but maintain mitochondrial membrane integrity at ambient temperature. (**A**) After Mitotracker staining (Fig 1A), the average mitochondrial surface areas in log phase promastigotes of WT, *c14dm⁻*, and *c14dm⁻/+C14DM* were determined using Image J (~200 cells were analyzed for each parasite line, au: arbitrary unit). (**B**) Log phase promastigotes were lysed with digitonin at 37 °C or room temperature (RT) and mitochondria enriched membrane fractions (Mem.) were separated from cytosolic fractions (Sup.) as described in **Materials and Methods**. Western blots were performed using antibodies against cytochrome *c*, ISCL (a mitochondrial membrane protein), and HSP83 (a cytosolic protein).

Fig S2. *C14dm⁻* and *smt⁻* mutants show increased ΔΨm and mitochondrial ROS. Log phase and stationary phase (day 1-day 4) promastigotes were resuspended in PBS and labeled with 5 µg/ml of rhodamine 123 for 15 min (**A**) or with 5 µM of MitoSox Red for 25 min (**B**) at room temperature. MFIs were determined by flow cytometry. Error bars represent standard deviations from three independent experiments.

Fig S3. *C14dm⁻* mutants accumulate ROS in the mitochondria. Log phase promastigotes were labeled with 5 µM of MitoSox Red for 25 min (**A**) or 5 µM of DHE for 30 min (**B**) and MFIs were determined by flow cytometry. Effects of AA (5 µM) and H_2O_2 (100 µM) on WT parasites were also monitored. Error bars represent standard deviations from three experiments.

Fig S4. 100 µM of glucose or glycerol is sufficient to rescue *c14dm⁻* mutants in PBS. Log phase promastigotes were incubated in PBS in the absence or presence of glucose (100 µM) or glycerol (100 µM) and percentages of dead cells were determined by flow cytometry at the indicated time points. Error bars represent standard deviations from three experiments.

Fig S5. ITZ-treated WT parasites are hypersensitive to glucose depletion. *L. major* WT

promastigotes were cultivated in solvent alone (0.1% w/v of DMSO) or 0.2 µM of ITZ for 48

hours. Cells were then incubated in HBSS and percentages of dead cells were determined by

flow cytometry at the indicated time points. Error bars represent standard deviations from three

experiments.

Fig S6. *C14dm⁻* mutants rescued by glucose or glycerol show less severe mitochondrial

defects. Log phase promastigotes were incubated in PBS in the absence or presence of glucose

(100 µM) or glycerol (100 µM) for 24 hours. $\Delta\Psi$m (**A**) and mitochondrial ROS level (**B**) were

determined by flow cytometry. Control cells represent cells analyzed at the beginning of

incubation. †: no viable *c14dm⁻* mutants were available after 24 hours. Error bars represent

standard deviations from three experiments.

Figure 1. *C14dm⁻* promastigotes show high mitochondrial membrane potential (ΔΨm). (**A**) Log phase promastigotes were labeled with 100 nM of Mitotracker CMXRos followed by DAPI staining. WT parasites pretreated with CCCp were included as controls. DIC: differential interference contrast. (**B**) Representative transmission electron micrographs showing mitochondria in log phase promastigotes. Black arrows indicate areas containing kinetoplast (top panel) and tubular mitochondria (bottom panel). (**C**) To measure ΔΨm, log phase and stationary phase (day 1-day 4) promastigotes were labelled with 100 nM of TMRE for 15 minutes in PBS and mean fluorescence intensity (MFI) were determined by flow cytometry. Controls include WT and *c14dm⁻* parasites pretreated with CCCp and WT cells pretreated with 0.2 μM of ITZ. Bar values represent averages from three experiments and error bars represent standard deviations. **: *p*<0.01, *: *p*<0.05. (**D**) Live parasites imaging after TMRE labeling.

Figure 2. Inactivation of C14DM leads to ROS accumulation in the mitochondria. (**A**, **B** and **F**) Promastigotes were incubated in complete M199, serum-free M199, PBS contain 5.5 mM of glucose, or PBS only and labeled with 100 nM of TMRE (**A**, 15 min), 5 µM of MitoSox Red (**B**, 25 min) or 5 µM of DHE (**F**, 30 min). MFIs were determined by flow cytometry. Effects of CCCp (75 µM, 15 min) were also monitored in **A** and **B**. (**C**) Log or stationary phase (day 1-day 4) promastigotes were labeled with 5 µM of MitoSox Red for 25 min in PBS and MFIs were determined by flow cytometry and normalized to values from log phase WT. WT cells grown in the presence of ITZ (for 48 hr) were also analyzed. (**D** and **E**) Live cell imaging of *L. major* promastigotes (**D**) or *Leishmania mexicana* axenic amastigotes (grown in the absence or presence of 0.1 µM of ITZ for 48 hr, **E**) after labelling with MitoSox Red in PBS. White arrows indicate kDNA binding of MitoSox. (**E**) Live after labeling with MitoSox Red. In **A-C** and **F**, averaged values from five experiments are shown and error bars represent standard deviations. ***: $p<0.001$, **: $p<0.01$, *: $p<0.05$.

Figure 3. Increased superoxide dismutase (SOD) activity in _c14dm⁻_ parasites. (**A**) Whole cell lysates from log phase and stationary phase (day 1 to day 3) promastigotes were collected as described in Materials and Methods. SOD activity was measured using a SOD assay kit (BioAssay Systems) and results were normalized based on the amount of protein in each sample. Error bars represent standard deviations from two independent experiments (*: $p<0.05$). (**B**) Mitochondria enriched membrane fractions (Mem.) and cytosolic fractions (Sup.) were isolated from log phase promastigotes as described. Western blots were performed using antibodies against _L. amazonesis_ SODA (mitochondrial) and SODB (both SODB1 and SODB2, glycosomal).

Figure 4. *C14dm⁻* parasites show compromised cellular respiration. (**A-C**) Log phase promastigotes were incubated in a mitochondrial respiration buffer (**A-B**) or DMEM (**C**) and oxygen consumption rates (OCR) were determined using the MitoXpress probe as described in Materials and Methods. Buffer only and buffer with probe (no cells) were included as blanks (**A**). In **A** and **C**, fluorescence signals (in arbitrary units or A.U.) were measured every 90 s. In **B**, the relative OCR from the experiments in **A** was calculated by subtracting the 0 min reading from the peak reading and then divided by the number of minutes. WT parasites treated with antimycin A (AA) or CCCp were included as controls. (**D**) Cellular ATP contents in log phase promastigotes were measured after 1 hour incubation in an assay buffer (21 mM of HEPES, 0.7 mM of Na_2HPO4, and 137 mM of NaCl, pH 7.2) in the absence or presence of glucose (Glu, 5.5 mM), sodium pyruvate (Pyr, 5.5 mM), 2-deoxyglucose (2DG, 5.5 mM), or sodium azide (NaN_3, 20 mM). All experiments were repeated three times and error bars represent standard deviations (***: $p<0.001$, **: $p<0.01$, *: $p<0.05$).

Figure 5. *C14dm⁻* parasites are hypersensitive to starvation. Log phase promastigotes were inoculated in HBSS in the absence or presence of Glu (5.5 mM), Pyr (5.5 mM), aspartate (Asp, 5.5 mM), 2-deoxyglucose (2DG, 5.5 mM), or glycerol (0.1 mM). In **A**, percentages of cell death for *c14dm⁻* mutants were determined at the indicated time points. In **B**, percentages of cell death were determined after 24 hours for WT, *c14dm⁻, c14dm⁻/+C14DM, smt⁻*, and *smt⁻/+SMT* parasites. Error bars represent standard deviations from three independent experiments (***: *p*<0.001, **: *p*<0.01).

Figure 6. _C14dm⁻_ parasites show increased uptake of glucose and glycerol.
Incorporation of radiolabeled glucose (**A**) or glycerol (**B** and **C**) were determined as described under Materials and Methods. Transport assays were performed at room temperature (**A** and **B** unless specified) or 4 °C (**C**). Error bars represent standard deviations from three independent experiments (***: $p<0.001$, **: $p<0.01$).

Figure 7. ROS accumulation contributes to heat sensitivity in *c14dm¯* mutants. (**A**-**E**) Log phase promastigotes were incubated in PBS at 27 °C or 37 °C for 3 hours. In **A**, percentages of cell death were determined by flow cytometry after PI staining. In **B**, ΔΨm was determined after TMRE labeling. Unlabeled WT parasites were included as negative controls (Neg). In **C**-**D**, parasites were lysed and cytosolic fractions (Sup.) were separated from mitochondrial fractions (Mem.) as described in Materials and Methods. Distribution of cytochrome *c* and HSP83 (a cytosolic protein marker) were determined by Western blots (**C**) and quantified (**D**). In **E**, parasites were labelled with the cytosolic ROS indicator DHE. Fluorescence intensities were measured by flow cytometry and relative cytosolic ROS levels were determined. (**F**) Log phase promastigotes were incubated at 37 °C in the absence or presence of reduced GSH. After 8 hours, mitochondrial ROS levels were measured as described. (**G**) *C14dm¯* parasites were incubated at 37 °C in the absence or presence of reduced GSH and percentages of cell death were determined at 0 and 8 hours. (**H**) WT parasites were incubated at 27 °C or 37 °C in the absence or presence of AA (10 µM) and percentages of dead cells were determined by flow cytometry at the indicated time points. †: Very few WT+AA cells were detectable after 36 hours at 37 °C. Error bars represent standard deviations from three independent experiments (***: $p<0.001$, **: $p<0.01$, *: $p<0.05$).

Figure 8. *C14dm¯* parasites show increased sensitivity to pentamidine. (**A**) Log phase promastigotes were inoculated in M199 media containing various concentrations of pentamidine. Culture densities were determined after 48 hours and the pentamidine concentrations required to inhibit growth by 90%, 50%, and 25% were determined (as EC90, EC50 and EC25 respectively) using cells grown in the absence of pentamidine as controls. "WT + ITZ" represent WT parasites grown in the presence of 0.5 μM of itraconazole. (**B**-**C**) Log phase promastigotes were inoculated in M199 media containing 2.5 μM (**B**) or 25 μM (**C**) of pentamidine for one hour, followed two washes with PBS. Cells were then re-inoculated in M199 media and culture densities were determined every 24 hours. Some *c14dm¯* mutants were also incubated in the presence of 75 μM of CCCp along with pentamidine (for one hour). Parasites originally inoculated in the absence of pentamidine were included as controls. Error bars represent standard deviations from three independent experiments (***: $p < 0.001$, **: $p < 0.01$, *: $p < 0.05$).