Title: Selective REM-Sleep Suppression Increases Next-Day Negative Affect and Amygdala Responses to
Social Exclusion

Authors:

1,2 Robert W Glosemeyer, 3,4 Susanne Diekelmann, 5 Werner Cassel, 5 Karl Kesper, 5 Ulrich Koehler, 6
Stefan Westermann, 7 Armin Steffen, 2 Stefan Borgwardt, 2 Ines Wilhelm, 1,2 Laura Müller-Pinzler, 1,2
Frieder M Paulus, 1,2 Sören Krach, 1,2 David S Stolz

Email:

Robert W Glosemeyer glosemeyerrobert@gmail.com

Susanne Diekelmann susanne.diekelmann@uni-tuebingen.de

Werner Cassel cassel@staff.uni-marburg.de

Ulrich Koehler koehleru@med.uni-marburg.de

Karl Kesper kesperk@med.uni-marburg.de

Stefan Westermann st.westermann@uke.de

Armin Steffen armin.steffen@uksh.de

Stefan Borgwardt stefan.borgwardt@uksh.de

Ines Wilhem Ines.Wilhelm-Groch@uksh.de

Laura Müller-Pinzler mueller-pinzler@snl.uni-luebeck.de

Frieder M Paulus paulus@snl.uni-luebeck.de

Sören Krach* krach@snl.uni-luebeck.de

David S Stolz* david.stolz@uni-luebeck.de

Affiliations:

1 Social Neuroscience Lab, Department of Psychiatry and Psychotherapy, University of Lübeck,
Ratzeburger Allee 160, D-23538 Lübeck, Germany

2 Translational Psychiatry Unit (TPU), Department of Psychiatry and Psychotherapy, University of Lübeck,
Ratzeburger Allee 160, D-23538 Lübeck, Germany

3 Institute of Medical Psychology and Behavioral Neurobiology, University Tübingen, Otfried-Müller-Str.
25, 72076 Tübingen, Germany

4 Department of Psychiatry and Psychotherapy, University Hospital Tübingen, Calwerstr. 14, 72074
Tübingen, Germany

5 Department of Internal Medicine, Division of Pneumology, Intensive Care and Sleep Medicine, Hospital
of the University of Marburg, Germany

6 Department of Psychiatry and Psychotherapy, University Medical Center Hamburg-Eppendorf,
Hamburg, Germany

7 Department of Otorhinolaryngology, University of Lübeck, Germany

Short Title: REM & Affect

Corresponding Authors*:

Sören Krach & David S. Stolz

Department of Psychiatry and Psychotherapy, Social Neuroscience Lab

University of Lübeck, Ratzeburger Allee 160, D-23538 Lübeck, Germany

Phone: +49 45131017529

Pages: 44

Tables: 3

Figures (Including Color Figures): 3

Color Figures: 3

Supplementary Information: 1

Words (including Abstract): 6844

Keywords: REM, selective sleep suppression, social exclusion, general affect, emotion, amygdala

Abstract

Healthy sleep, positive general affect, and the ability to regulate emotional experiences are

fundamental for well-being. In contrast, various mental disorders are associated with altered rapid eye

movement (REM) sleep, negative affect, and diminished emotion regulation abilities. However, the neural

processes mediating the relationship between these different phenomena are still not fully understood.

In the present study of 42 healthy volunteers, we investigated the effects of selective REM sleep

suppression (REMS) on general affect, as well as on feelings of social exclusion, emotion regulation, and

their neural underpinnings. Using functional magnetic resonance imaging we show that REMS increases

amygdala responses to experimental social exclusion, as well as negative affect on the morning following

sleep deprivation. There was no evidence that emotional responses to experimentally induced social

exclusion or their regulation using cognitive reappraisal were impacted by diminished REM sleep. Our

findings indicate that general affect and amygdala activity depend on REM sleep, while specific emotional

experiences possibly rely on additional psychological processes and neural systems that are less readily

influenced by REMS.

Introduction

Sleep is fundamental for general well-being (Howell et al., 2008; Steptoe et al., 2008). Healthy sleep consists of two repeatedly cycling types of sleep: non-rapid eye movement sleep (NREM) including slow-wave sleep (SWS) phases of deep sleep and rapid eye movement sleep (REM). Over the course of the night, the percentage of SWS decreases while REM sleep percentage increases towards the morning (Kryger et al., 2017). In many occasions however, sleep is fragmented or shortened (e.g. due to stress, early school or work starts and late bedtimes) and even minor sleep deprivation can have broad, short-term as well as long-term consequences for health and well-being. Sleep disturbances affect health on the level of immune regulation (Besedovsky et al., 2019; Bryant et al., 2004; Lange et al., 2010) and metabolic markers (Briançon-Marjollet et al., 2015; Koren et al., 2016). Furthermore, sleep loss negatively impacts cognitive processing and emotional reactivity (Durmer & Dinges, 2005; Walker, 2008), as well as social behavior, leading to withdrawal from social interaction and feelings of loneliness (Ben Simon & Walker, 2018).

Interestingly, most mental disorders are associated with sleep peculiarities. Insomnia is comorbid in 85-90% of individuals diagnosed with major depression (Franzen & Buysse, 2017; Geoffroy et al., 2018; Staner, 2010). Alterations in REM sleep architecture (REM sleep latency, density and distribution) are particularly prominent in most mental disorders such as major depression, bipolar disorder and Posttraumatic Stress Disorder (PTSD; Armitage, 2007; Gottesmann & Gottesman, 2007). Given that these disorders are mainly characterized by altered affective experience during daytime, it is interesting to note that inadequate sleep itself has been shown to impact next day emotionality. Sleep loss has been shown to reduce the ability to regulate emotions, leading to exaggerated responses to aversive or even neutral experiences (E. Ben Simon et al., 2015; Zohar et al., 2005a). Recent studies even suggest sleep abnormalities as being causal agents in mood disorders rather than just a symptom (Goldstein & Walker, 2014; Staner, 2010).

Although single sleep stages seem to be differentially involved in mood disorders, the role of these

sleep stages for cognitive and emotional functioning is not well understood. Increasing evidence suggests

that SWS and associated neurophysiological processes support the consolidation of newly encoded

declarative memories (Diekelmann & Born, 2010; Stickgold & Walker, 2013). REM sleep, on the other hand,

is particularly important for emotional processing, including emotional reactivity (Baran et al., 2012;

Wagner, 2002) and the formation of emotional memories (Nishida et al., 2009; Wagner et al., 2001). REM-

sleep dreaming was also found to attenuate residual emotional load from the day before (e.g., Paré et al.,

2002; Walker, 2009). However, only very few studies have examined the effects of selective suppression

of REM sleep during an otherwise normal night of sleep. The existing evidence suggests that the selective

deprivation of REM sleep mainly disturbs the consolidation of emotional memories (Lipinska & Thomas,

2019; Spoormaker et al., 2012; Wiesner et al., 2015), whereas the selective suppression of SWS mainly

impairs emotionally-neutral declarative memory encoding and consolidation (Casey et al., 2016; Van Der

Werf et al., 2009). These experimental findings are in line with clinical observations suggesting that REM

sleep in particular is closely tied with emotional functioning (Goldstein & Walker, 2014).

In order to better understand the processes underlying the interaction of sleep and emotion,

research increasingly addresses the neural mechanisms associated with REM sleep-dependent emotional

functioning. At the neural level, a reduction of emotional reactivity in response to previously learned

emotional events was found to be accompanied by an overnight decrease of amygdala reactivity, a cluster

of nuclei in the temporal lobes and part of the limbic system (van der Helm et al., 2011; Walker, 2009).

The limbic system has been widely associated with the processing of affectively laden stimuli and plays an

important role in the guidance of behavioral responses to such stimuli (Phan et al., 2004; Taylor et al.,

2003). Under conditions of total sleep deprivation, next-day negative affect is accompanied by increased

amygdala reactivity and decreased functional coupling of the medial prefrontal cortex (MPFC) with limbic

structures (Yoo et al., 2007). Since the MPFC is thought to exert inhibitory control on the amygdala

(Davidson, 2002), this finding has been interpreted to reflect a failure of top-down control in the regulation

of appropriate emotional responsivity (Yoo et al., 2007). Simon and colleagues directly examined the effect

of sleep deprivation on the neural correlates of next day emotion reactivity (E. Ben Simon et al., 2015).

Following sleep deprivation, there was no indication for valence-specific processing of affective pictures

in the amygdala, however in the sleep-rested control night, a low amount of REM sleep was associated

with a decline in anterior cingulate cortex (ACC)-amygdala connectivity, possibly reflecting a specific effect

of REM sleep on cognitive control of emotions.

Successful cognitive control of emotions is regarded to be an essential prerequisite of mental

health (Berking & Wupperman, 2012; Gross & Muñoz, 1995). In daily life, emotions are constantly

regulated either implicitly or explicitly by applying specific cognitive strategies like suppression (e.g.

distracting the attention away from unpleasant emotional experiences) or reappraisal (e.g. reinterpreting

an unpleasant emotional situation). Emotion regulation thus refers to the ability to "influence which

emotions we have, when we have them, and how we experience and express these emotions" (Gross,

2008, p.497). Interestingly, correlational evidence indicates that the success of emotion regulation is

associated with sleep quality (Mauss et al., 2013). In this study, participants were asked to engage in

cognitive reappraisal (CRA; Gross, 2002), in this case applying a previously learned cognitive strategy to

"redirect the spontaneous flow of emotions" (Koole, 2009, p. 6), while watching a sadness-inducing film.

The ability to decrease self-reported sadness using CRA compared to baseline was lower in participants

who reported poorer sleep quality during the preceding week (Mauss et al., 2013).

Despite their substantial clinical significance, the neural mechanisms of the effect of REM sleep on

the efficacy of regulation strategies in ameliorating unpleasant affect remain unknown. To fill this gap, it

is important to experimentally induce negative affect that may then be attenuated by applying CRA

strategies. In order to induce negative affect in the present study, we simulated social exclusion in a

laboratory setting using the so-called *Cyberball* (Williams & Jarvis, 2006). *Cyberball* is a virtual ball-tossing

paradigm, where participants are playing with a preset computer program while believing that they are

playing with two other human participants. By manipulating the number of ball-tosses towards the

participant, the degree of social inclusion can be controlled experimentally. Evidence suggests that the

distressing experience of social exclusion might share neural regions and mechanisms with the affective

processing of physical pain (Dewall et al., 2010; Eisenberger & Lieberman, 2004; MacDonald & Leary,

2005). In a neuroimaging study using the *Cyberball* game, participants showed greater activations in

cingulate and prefrontal regions involved in the processing of the affective pain component when excluded

compared to being included (Eisenberger et al., 2003). Although behavioral consequences and neural

activation patterns associated with social exclusion have been studied quite intensively, there is scarce

research on intervening cognitive appraisals and coping mechanisms regarding feelings of social exclusion

(Williams, 2007).

The aim of the present study was to examine how REM sleep suppression impacts the following

day general affect, emotional reactivity, and associated neural mechanisms of emotion regulation during

the acute experience of social exclusion. In a between-subjects design we invited participants to a

combined polysomnography and fMRI study. After a habituation night allowing regular sleep, participants

were randomly allocated to either a REM sleep suppression (REMS) group or one of two control groups: a

non-suppression control group with regular sleep (CTL) and a high-level control group with similar amounts

of awakenings, but where suppression targeted phases of slow wave sleep (SWSS). In the morning after

the suppression night, subjects participated in the *Cyberball* to induce negative emotions during fMRI. All

participants engaged in two sessions of the game. In the first session, participants played the game without

any instructions. In the second session, participants were instructed to actively regulate their emotions by

applying the previously learned CRA.

We hypothesized that selective REMS (vs. SWSS and regular sleep) generally reduces positive and

increases negative affect. Furthermore, we expected that selective REMS (vs. SWSS and regular sleep)

increases emotional reactivity during social exclusion and dampens the effect of CRA on emotional

reactivity. On the level of neural systems we explored whether REMS leads to altered functional activity in

(para-)limbic areas such as the amygdala, ACC, insula, and hippocampus during social exclusion and

whether neural activity of these regions is modulated by targeted REMS during cognitive reappraisal.

Methods

Participants

A total of 45 participants were initially invited to take part in the experiment (28 female, age (years):

M=23.69, SD=2.67), gave written informed consent prior to participation in the study and received

financial compensation for their participation. All participants were recruited at Philipps-University

Marburg, were fluent in German, and had normal or corrected-to-normal vision. None of them were

diagnosed with neurological or psychiatric disorders (present and past), current alcohol or drug abuse, use

of psychiatric medications (present and past), anatomical brain abnormalities (e.g. lesions, strokes etc.),

or sleep disturbances. Due to technical issues with the polysomnographic recording in the experimental

night (n=1) or insufficient quality of the fMRI data (n=2), 3 subjects were not included in the final analyses.

The final sample thus consisted of 42 participants (27 female, age (years): M=23.76, SD=2.74). Participants

were randomly assigned to one of three groups, which differed regarding to the sleep protocol in the

experimental night (i.e. either REMS, SWSS or CTL, details below). Groups did not differ in age (CTL:

M=24.00, SD=3.12; REMS: M=23.06, SD=2.22; SWSS: M=24.60, SD=2.91; $F_{(2,39)}$=1.09, p=.346) or gender

(CTL: 8 female, 7 male; REMS: 12 female, 5 male; SWSS: 7 female, 3 male; $X^2_{(2,42)}$=1.22, p=.543). REM

sleep was successfully suppressed in the REMS group during the experimental night (REMS score defined

as %REM sleep in habituation minus %REM sleep in experimental night; effect of group: $F_{(2,37)}$=13.21,

p<.001, η^2=.42; planned contrast REMS > others: t(38)=5.17, one-sided p<.001). SWSS and CTL were similar

with regard to REMS (t=0.52, two-sided p=.608; see table 1 and figure 2 for details).

Sleep Manipulation Procedure

Participants came to the laboratory for two consecutive nights – one habituation night and a subsequent

experimental night (see figure 1). The purpose of the habituation night was to exclude sleep disorders,

make participants familiar with the polysomnography (PSG) recording procedure and adapt to the

environment in the sleep laboratory. The morning after the experimental night, participants were taken

to the fMRI facility where they completed the experimental task in the scanner (see Experimental task

below). During both nights PSG was recorded to monitor and identify sleep phases (see table 1). A

maximum of three participants was invited to the sleep laboratory on each night. Upon arrival on the

habituation night, participants were informed about the procedure of the study and gave written informed

consent. After the habituation night participants could spend the day as usual but were asked to refrain

from napping, smoking and consuming stimulating foods and drinks (e.g. coffee, tea, energy drinks). On

the subsequent night, participants entered a fully equipped single sleep room for PSG (Embla N7000 PSG,

TNI-Medical, Würzburg, Germany) consisting of electroencephalography (EEG), electrooculogram (EOG),

electromyogram of the mentalis muscle (EMG) and pulse oximetry using EMBLA N7000 (TNI-Medical). The

EEG system had 9 electrodes positioned according to the 10-20 system, adhering to the guidelines of the

American Academy Of Sleep Medicine (AASM; Berry et al., 2015), and placed on the following positions:

F3, F4, C3, Cz, C4, O1, O2, M1 and M2. Limb movements were videotaped with infrared camera. The

experimenter turned off the lights at 10 pm and asked participants to try to fall asleep. In case a

deterioration of the PSG recordings was observed, the experimenter entered the sleep room to improve

the signal by reattaching the respective electrodes. At 6 am, lights were turned on, participants were

woken up and the electrodes were removed.

For those participants in the REMS and SWSS group, during the second night (i.e. the experimental

night), the experimenter started an acoustic beep (80 dB, 500 Hz, 500ms; Nielsen et al., 2010) once the

polysomnographic trajectories indicated that the participant had entered the target sleep phase. After an

awakening, participants were kept awake for 90 seconds. In case the participant did not wake up, the

volume of the acoustic beep was increased, and ultimately the experimenter entered the participant's

room to turn on the lights to make sure the target sleep phase was interrupted. To control for the number

and lengths of awakenings, both suppression groups were disturbed similarly often during the

experimental night. In the REMS group, participants' sleep was disturbed as soon as they entered the REM

sleep stage. Awakenings for the SWSS group were selectively carried out during the non-REM sleep stage

N3, defined according to AASM guidelines (Berry et al., 2015). The control subjects were not awakened at

all. PSG recordings were scored by an experienced sleep technician (C.L.) at the sleep laboratory at the

Department of Otorhinolaryngology at the University of Lübeck according to AASM guidelines (Berry et al.,

2015) and using Somnologica 3.3.1 (Build 1529). The technician was blinded for group assignments and

hypotheses.

General Affect Ratings

The PANAS (Positive and Negative Affect Schedule, Crawford & Henry, 2004) was completed shortly before

going to bed and after waking up on both nights by all participants to assess the effect of sleep

manipulation on general positive and negative affect.

Experimental Task

Upon waking up after the experimental night, participants were accompanied to the functional magnetic

resonance imaging (fMRI) facility. In the MRI, participants were instructed via a screen that they were

about to play a ball-tossing game, i.e. the *Cyberball* task, allegedly with two other persons (see Figure 1; a

second paradigm was presented to the subjects afterwards, which involved IAPS pictures to study

regulation of basic emotions but is not further described in the present manuscript). All three players,

including the participant, were represented by avatars, i.e. green, 3D-animated stick-figures standing in a

triangle on a lawn (see Burgdorf, Rinn, & Stemmler, 2016 for a detailed description of the animation).

Participants were able to control the avatar on the bottom edge of the screen, while the other two avatars

were placed on the left and right of the horizontal midline of the screen. The names of the participant and

of the other two players were displayed next to the respective avatar. For each throw, a short sentence

on the bottom of the screen indicated who threw the ball to whom. When one of the computer avatars

had the ball, they waited a random amount of time between 1000 and 2000 ms before tossing the ball.

When the participant had the ball, they had 2000 ms to decide where to toss the ball. In case they

responded too slowly, the computer randomly selected which of the other avatars would receive the ball.

The duration of each throw was 750 ms and was visualized by a series of 40 frames, showing an avatar

doing a throwing movement and the red ball flying from one avatar to the catching avatar, with the latter

moving its arm to catch the ball. While participants believed that the other avatars were controlled by two

other anonymous persons sitting in adjacent rooms, their behavior in fact followed a predefined script

(see below). Following the instructions, a short sequence of lines of text built up on the screen, making the

subjects believe that the computer was being connected to a gaming server of the university on which the

experiment was run. This included the request of entering an IP-address, as well as lines stating how many

players were logged into the game. Subsequently, participants performed a short training session

consisting of seven trials in which they received and threw the ball three times.

The task then consisted of two sessions, each of which consisted of four experimental blocks.

Before each of the experimental blocks, a line of text was presented on-screen for 1500 ms telling the

participant that a new round would start. In every block, a maximum number of 24 ball tosses were

completed. In two blocks of every session, the participant was included in the game and repeatedly

received the ball throughout the entire block (inclusion blocks, INC), having the possibility to toss the ball

to one of the other avatars eight or nine times by clicking a button using the index finger (left player) or

middle finger (right player). In the other two blocks (exclusion blocks, EXC), after the participant had

received the ball three or four times, the paradigm was programmed so that the two avatars only tossed

the ball between one another, thereby effectively excluding the participant from the game.

After having completed four blocks during which subjects simply participated in the task without

additional instructions (VIEW session), a second session of the task was played. For the second session, a

short on-screen text instructed participants to use cognitive reappraisal to regulate their emotions during

the game (CRA session; see Fig. 1). Participants were specifically asked to reappraise the situation, in case

any negative emotions should arise ("In case negative emotions arise, please try to re-evaluate the

situation"). This was accompanied by the instruction to "partake in the game and try to visualize the

situation as vividly as possible", which was also presented before the first session. Subjects were asked to

press the left button as soon as they were ready, and then a short break varying randomly between 1500

and 2500 ms was presented, instructing participants to wait until the other two players had indicated that

they were ready. After presentation of a fixation cross for one second, the block started.

In each session, two inclusion and two exclusion blocks were presented to the subjects, with the

first block of each session always being an inclusion block. In one session, inclusion and exclusion blocks

were alternating, while in the other session, the initial inclusion block was followed by two consecutive

exclusion blocks and a final inclusion block. The assignment of block order to the VIEW and the CRA

sessions was counterbalanced across subjects. On average, EXC blocks lasted 50.93 seconds (SD=0.62), and

INC blocks lasted 45.87 seconds (SD=2.21).

Between blocks, participants were asked to rate how they felt about the preceding block regarding

the extent to which they felt socially excluded. The low end of the scale was labelled as rejected/despised

and the high end was labelled as accepted/familiar. In addition, subjects rated their sadness, anger, and

shame, but these ratings were intended to distract from the purpose of the study and not analyzed

(sadness was described by: sad, downcast, gloomy; for anger: angry, irritated, furious, mad; for shame:

abashed, embarrassed). Each rating was displayed using a 9-point Likert-type scale ranging from 0 (not at

all) to 8 (very much). Every rating was initialized at 4 (i.e. a neutral rating), and participants used presses

of the right or left response buttons to move the rating to higher or lower values, respectively. The time

for ratings was limited to 4 seconds. After each rating and the start of the next block within a session there

was a short pause, jittered between 4.3 to 5.3 seconds. Emotion ratings for each condition (EXC, INC) and

emotion regulation session (VIEW, CRA) were averaged for each participant and analyzed using repeated

measures analyses of variance (rmANOVA).

Statistical Analysis

Average ratings of feeling excluded for each condition (EXC, INC) and session (VIEW, CRA), as well as the

percentages of the different sleep stages for the habituation and experimental nights were analyzed using

analyses of variance (ANOVA). Significant interactions and main effects were followed up using paired

comparisons. Since we expected that selective REMS would specifically alter affective experience and

associated neural responses (Baran et al., 2012; Wagner, 2002), we furthermore performed an a priori

planned contrast of the REMS group against both control groups, the CTL and SWSS. The alpha-level was

set to .05, and was adjusted using Bonferroni-correction, in case multiple tests were performed.

FMRI Data Acquisition and Preprocessing

For each of the two experimental sessions, 130 functional volumes were recorded at 3T (Siemens Trio,

Erlangen), of which the first three were discarded to allow for equilibration of T1 saturation effects.

Functional volumes consisted of 36 ascending near-axial slices (voxel size=3*3*3 mm, 10% interslice gap,

FOV=192 mm) and were recorded with TR=2200 ms, TE=30 ms, FA=90°. In addition, a high-resolution

anatomical T1 image was recorded consisting of 176 slices (voxel size=1*1*1 mm, FOV=256 mm, TR=1900

ms, TE=2.52 ms, 9° FA).

The MRI data were analyzed using SPM12 in Matlab 2019b. Functional MRI images from each

session of the *Cyberball* paradigm were slice-time corrected to the middle slice and spatially realigned.

Subsequently, spatial normalization to MNI space was performed using unified segmentation (Ashburner

et al., 2013) by estimating the forward deformation fields from the mean functional image and applying

these to the realigned functional images. These spatially normalized images were then resliced to a voxel

size of 2*2*2 mm and smoothed with an 8 mm full-width at half-maximum isotropic Gaussian kernel and

high-pass filtered at 1/256 Hz.

FMRI Data Analysis

The preprocessed functional images were statistically analyzed by a two-level mixed effects GLM

procedure. For each participant, a statistical model was specified, including data from both experimental

sessions (VIEW, CRA). For each session, the INC, EXC and rating phases were modelled as regressors of

interest and the six realignment parameters estimated during spatial realignment were included to

account for variance in the functional data that was due to head motion. The contrast images obtained

from the individual participants were then aggregated in a random effects model on the second level to

test for effects of condition and session, and to test for differences between the three experimental

groups. To disentangle significant interaction effects, we ran a series of post-hoc tests on the parameter

estimates extracted from the peak activation, applying Bonferroni-correction for multiple comparisons.

Regions of Interest (ROI) for FMRI Analysis

In order to focus our analyses on the limbic system as well as the insula, regions commonly associated with

affective processing (Lindquist et al., 2016), we constructed an a priori mask using the Wake Forest

University Pickatlas (v. 2.4; (Lancaster et al., 1997, 2000; Maldjian et al., 2003, 2004). This mask comprised

the anterior cingulate cortex, bilateral insula, bilateral hippocampus, as well as left and right amygdala.

Results

REM Sleep Predicts General Negative Affect

General affect ratings in the Positive and Negative Affect Schedule (PANAS) obtained in the morning

differed significantly between habituation and experimental night (effect of night: PA: F(1,37)=4.55,

p=.040, η_p^2=.11; NA: F(1,37)=9.33, p=.004, η_p^2=.20), while type of suppression had no significant effect (PA:

F(2,37)=0.74, p=.484, η_p^2=.04; NA: F(2,37)=2.21, p=.124, η_p^2=.11). The interaction effect was not significant

for positive affect (PA: F(2,37)=0.34, p=.713, η_p^2=.02) but approached significance for negative affect (NA:

F(2,37)=3.08, p=.058, η_p^2=.14). Post-hoc comparisons demonstrated that across groups positive affect was

lower and negative affect was higher after the experimental night as compared to the habituation night

(PA: t(39)=2.20, two-sided p=.034, d=0.35, 95% CI=[0.03;0.67]; NA: (t(39)=-3.23, two-sided p=.003, Cohen's

d=-0.51, 95% CI=[-0.84;-0.18]), with the increase in negative affect being most pronounced in the REMS

group (t(14)=4.47, p=.001, Cohen's d=1.15, 95% CI=[0.48;1.80]; see figure 2). A planned contrast showed

that this increase in negative affect was stronger in the REMS group as compared to the other groups

(t(38)=2.47, two-sided p=.037, Cohen's d=.81, 95% CI=[0.14;1.47]), whereas no group difference could be

observed for change in positive affect (t(38)=-0.72, two-sided p=.950, Cohen's d=.24, 95% CI=[-0.88;0.41],

all p-values Bonferroni-corrected). Regarding the evening ratings, type of sleep suppression did not have

any significant effects or interactions, neither for PA nor for NA (all ps>.110; see supplementary

information for details). Across groups, increases in morning negative affect from habituation to

experimental night could be predicted by percentage of REM sleep (Pearson's r=.39, one-sided p=.015,

95% CI=[0.08;0.63]; figure 2), even after controlling for changes in SWS, total sleep time (TST) and

wakefulness after sleep onset in a multiple linear regression model (see table 2).

No Effect of Selective REM Sleep Suppression on Emotional Responses to Social Exclusion

By means of the *Cyberball*, we successfully induced the unpleasant feeling of social exclusion (main effect of INC/EXC: $F(1,39)=131.65$, $p<.001$, $\eta_p^2=.771$; see table 3 for descriptive statistics), thereby replicating earlier findings (Eisenberger et al., 2003). Moreover, if participants were asked to engage in emotion regulation using cognitive reappraisal, feelings of being excluded could be toned down significantly (main effect of VIEW/CRA: $F(1,39)=66.99$, $p<.001$, $\eta_p^2=.632$), with this effect being most strongly pronounced during trials of social exclusion (interaction effect INC/EXC x VIEW/CRA: $F(1,39)=44.97$, $p<.001$, $\eta_p^2=.536$).

However, regarding our second hypothesis, type of sleep suppression neither had a significant impact on emotions after social exclusion (INC/EXC * group interaction: $F(2,39)=1.47$, $p=.243$, $\eta_p^2=.070$) nor on the effect of emotion regulation (VIEW/CRA * group interaction: $F(2,39)=.027$, $p=.973$, $\eta_p^2=.001$). Last, we did not find a statistically significant three-way interaction on the self-reported emotions after social exclusion (INC/EXC * VIEW/CRA * group interaction: $F(2,39)=.45$, $p=.639$, $\eta_p^2=.023$). Similarly, no significant main effect or interactions with type of sleep suppression were found when contrasting REMS against the other two groups (all *p*-values >. 41).

REM Sleep Suppression Alters Amygdala Activity During Social Exclusion

Similar to earlier studies, we found that during social exclusion participants' neural activity in the left and right hippocampus (left: x,y,z (mm): -36,-40,-6; *T*=7.15, *k*=89, *p*<.001, FWE-corrected; right: 36,-32,-8; *T*=4.88, *k*=4, *p*=.022, FWE-corrected) as well as the right insula (34,-10,22; *T*=5.07, *k*=13, *p*=.014, FWE-corrected) was significantly increased compared to the inclusion condition (Bolling et al., 2011; Masten et al., 2009). In addition, across groups, neural activity in the right anterior insula was significantly increased during VIEW as compared to CRA blocks (main effect VIEW>CRA: 30,24,-4; *T*=4.58, *k*=1, *p*=.048, FWE-corrected). Testing whether these two main effects interacted or whether they were modulated by type of sleep suppression did not yield any significant effects. This held both when testing for differences

between any of the three experimental groups and when comparing the REMS group against the other

two groups combined.

Last, we tested the three-way interaction of EXC/INC, VIEW/CRA, and type of sleep suppression. A

test of differences between any of the three groups for the two-way interaction contrast

[EXC/VIEW>INC/VIEW] > [EXC/CRA>INC/CRA] was not significant. However, comparing the REMS group to

the other two groups indicated differential neural responses in the right amygdala for the REMS group (26,

-2, -28; F=23.70, k=1, p=.048, FWE-corrected inside the a priori mask; see figure 3). Precisely, the contrast

EXC>INC was positive during VIEW in the REMS group, but did not differ from zero in the other groups

(REMS: t=4.04, two-sided p=.005; CTL: t=-1.07, two-sided p=1.000; SWSS: t=-0.58, two-sided p=1.000; see

figure 3). In addition, EXC>INC was more positive during VIEW than during CRA in the REM group (t=5.27,

two-sided p<.001), but not in the other groups (CTL: t=-2.18, two-sided p=.141; SWSS: t=-1.70, two-sided

p=.369; all p-values Bonferroni-corrected). This pattern of results suggests that the signaling of information

that is relevant to the individual's social well-being and mediated by amygdala activity depends on intact

REM-sleep.

Discussion

Nearly everyone can relate to the devastating effects of a sleepless or interrupted night on one's next day

mood. While the effect of total sleep deprivation on emotional reactivity has been investigated intensively

in the past (Krause et al., 2017; Zohar et al., 2005b) the present study focused on the specific impact of

selective REM sleep suppression on general affect, as well as emotion regulation and its neural correlates

under conditions of social exclusion.

We found that lower amounts of REM sleep across all participants were associated with higher

levels of general negative affect in the next morning, a finding that is in line with previous literature

implicating REM sleep in emotional functioning (for a review, see Vandekerckhove & Cluydts, 2010).

Despite this general effect, however, our findings do not provide evidence for a direct link of REMS with

the subjective emotional response to experimentally induced social exclusion. The ability to regulate one's

negative emotions during social exclusion was also not affected by prior REMS, which was an unexpected

finding. Interestingly though, despite no changes in subjectively reported emotions, neural activity in the

limbic system was altered after REMS when participants experienced social exclusion. Precisely, neural

responses to passively experienced ostracism were specifically increased in right amygdala after REM sleep

was selectively suppressed. The amygdala is strongly associated with emotional processing (LeDoux, 2000;

Phelps, 2006) and has anatomical connections to the anterior insula and the anterior cingulate cortex

(Amaral & Price, 1984; Cerliani et al., 2012). As part of this so-called salience network (Seeley et al., 2007),

the amygdala's assumed function of signaling the relevance of information is central for the domain of

affective experiences (Pessoa & Adolphs, 2010; Phelps, 2006). Previous studies showed that amygdala

responses to viewing negative emotional stimuli increased after total sleep deprivation (Yoo et al., 2007a),

depended on intact REM sleep in particular (van der Helm et al., 2011), and correlated with autonomic

responses to psychosocial stress (Orem et al., 2019). Our study connects these previous findings by

demonstrating that amygdala activity tracks information relevant to the subjects' social well-being in

dependence of REM sleep.

We can only speculate why these alterations in brain function after REMS did not manifest on the

behavioral level in form of increased emotional reactivity to the experience of social exclusion. Previous

research provides several explanations that might be helpful to understand this dissociation. First, the

effects of experimental short-term sleep manipulations might be strongest immediately after awakening

and may be readily washed out thereafter (Tassi & Muzet, 2000), and may thus not have lasted until the

fMRI session in the present study. This may hold in particular for selective suppression of specific sleep

stages rather than total sleep deprivation, which produces stronger and more long-lasting effects (Tassi &

Muzet, 2000). The efficacy of REMS in the present study may have been too small to surface on the

behavioral level later in the morning. However, the effects of REMS nonetheless persisted on the level of

brain systems. The altered brain activity in the amygdala could indicate that even small changes in REM

sleep can influence brain systems implicated in regulating responses to affectively salient stimuli (LeDoux,

2000; Phelps, 2006), while they are too small to penetrate the level of subjective experiences, which are

possibly processed further downstream (LeDoux, 2020; Orem et al., 2019). Yet, what speaks against this

explanation are findings by Wiesner and colleagues (2015) or Morgenthaler and colleagues (2014) who

applied even more rigorous REM sleep deprivation, achieving a mean REM sleep percentage of around

one percent of TST, but nevertheless could not find REM sleep related behavioral effects during emotion

recognition tasks. Similarly, Liu and colleagues compared the effect of a 24h sleep deprivation to regular

sleep on the experience of distress following social exclusion in the *Cyberball* game (Liu et al., 2014). As in

the present study, the authors did not find an effect of sleep deprivation on the subjective experience of

social rejection.

Regardless of the timing and potential washout during the day, the lack of significant findings for

the experience of social exclusion might also relate to the conceptual breadth and the ecological validity

of the affective assessment. While the underlying brain systems show increased reactivity towards

potentially threatening conditions and signal greater homeostatic imbalance, the distinct assessment of

emotional responses to ostracism, such as feelings of exclusion, might touch a different facet of affective

construal. That is, the affective salience of information tracked by amygdala activity (Pessoa & Adolphs,

2010; Phelps, 2006) may be modulated by REMS. However, the construal of emotional experience is

assumed not to rely on the activity in single regions, but on the dynamic interaction of various neural

systems supporting multiple psychological processes of emotional experience apart from affective salience

(Lindquist et al., 2012; Lindquist & Barrett, 2008). Hence, amygdala responses to psychosocial stress do

not necessarily influence the construal of subjective emotion ratings, that may depend on additional

networks involving prefrontal cortical regions, as suggested by previous studies (Motomura et al., 2013;

Orem et al., 2019; Urry et al., 2006). In addition, we did not find evidence that REM sleep deprivation

modulated the ability to apply cognitive reappraisal on the behavioral level. Therefore, REM sleep deprived

subjects may have been able to counteract the impact of increased amygdala responses during social

exclusion. A potential factor diminishing the severity of experimental social exclusion is that despite our

attempts to create a socially immersive context (Krach et al., 2013), laboratory studies per se have limited

relevance beyond the experimental situation itself. Thus, even without explicit reappraisal strategies, this

limited relevance might have entered into the construal of emotional responses during passive social

exclusion, and reduced the influence of altered amygdala responses in the REMS group.

A related explanation for the brain-behavior dissociation is that healthy participants have

resources for compensation. As the neuroimaging findings advocate that REMS impacts limbic circuit

activity during emotional experiences, the question arises whether participants with a priori (sub)clinical

peculiarities in reaction to social exclusion, ostracism or labile sense of belonging would also report altered

subjective experiences. It has been shown that inter-individual dispositions such as differences in rejection

sensitivity (Downey et al., 2004), social anxiety (Zadro et al., 2006), trait self-esteem, depression (Nezlek

et al., 1997), attachment style (Dewall et al., 2012) and situational factors of the exclusion experience

moderate the effect of social exclusion. Last, it is possible that experimental REM-sleep suppression

becomes effective on the emotional level only when applied repeatedly, simulating chronic sleep

disturbances that are associated with psychiatric disorders in a more realistic manner.

Speculatively, a further explanation could relate to the psychological task used to examine

emotional reactivity. Earlier research indicated that sleep deprivation effects might differ depending on

whether the focus was on general state-like morning affect (e.g. as measured using the PANAS), the

processing and responding to affective material (e.g. emotion recognition, pain processing, reactivity to

threatening stimuli) or on the direct induction of emotional states (e.g. inducing the unpleasant experience

of social exclusion). One assumption derived from the present data could be that direct emotion-induction

tasks are so powerful that a single night of sleep deprivation or selective REMS might not suffice to exert

effects on the psychological level. To our knowledge, however, apart from the present study there is no

further work that directly examined the effect of selective REMS on experimentally induced emotional

states. Second, general unspecific affect may function differently than emotional reactions to specific

elicitors (Beedie et al., 2005), potentially moderating the effect of REMS on these different affective

processes. However, at least one study found that general affect was not influenced by REMS, which

contradicts our findings (Wiesner et al., 2015). Last, we are not aware of any study systematically

comparing the extent to which the different psychological aspects of affective experience are susceptible

to selective suppression of sleep stages. Taken together, the specific interaction of selective REMS with

general affect, in contrast to more confined emotional responses, demands more in-depths analyses of

different kinds of experimental designs and dependent variables.

Our findings of increased next morning general negative affect as well as increased limbic activity

after REMS are well in line with findings in patients with mental disorders. A wide range of mental

disorders, including mood and anxiety disorders, are not only accompanied by profound disturbances of

REM sleep (Habukawa et al., 2018; Kobayashi et al., 2007) but also by deficits in generating and controlling

emotions in an appropriate way. Posttraumatic Stress Disorder (PTSD) is one of these mental disorders,

where a link between REM sleep and emotion regulation has been frequently discussed (Colvonen et al.,

2019; Pace-Schott et al., 2015). In these patients, deficits in emotion processing and regulation on the

behavioral level are coincided by alterations in brain networks implicated in emotion regulation such as

the amygdala, hippocampus, insula, and anterior cingulate (Kunimatsu et al., 2019). Apart from that, PTSD

patients as compared to traumatized individuals who did not develop a PTSD, show a profound REM sleep

fragmentation and alterations in REM density accompanied by higher sympathetic drive during this sleep

stage (Habukawa et al., 2018; Kobayashi et al., 2007). Importantly, these REM sleep disturbances early

after trauma predict later PTSD development (Bustamante et al., 2002; Mellman et al., 2007), which let

researchers speculate that REM sleep alterations are not only a secondary symptom but represent one

mechanistic factor contributing to the development and/or the exacerbation of PTSD symptoms (Colvonen

et al., 2019; Pace-Schott et al., 2015). However, empirical evidence for such a mechanistic link is scarce.

Our findings of altered negative affect and changed brain responses to social exclusion after one

night of REMS in healthy subjects resemble patterns typically seen in PTSD patients. Hence, the present

data provide additional evidence for a role of REM sleep in emotion processing and regulation and they

further suggest a mechanistic link between REM sleep disturbances and psychopathology. Future research

should investigate the effect of experimental REM sleep manipulation (i.e. REM sleep increase as well

deprivation/fragmentation) through pharmacological and/or psychological treatments (i.e. cognitive

behavioral therapy for insomnia, CBT-I; see e.g. Ashworth et al., 2015; Manber et al., 2008) on emotion

regulation in healthy subjects and patients suffering from mental disorders. Conducting more research in

this field gains additional importance from the fact that sleep deprivation does not have negative effects

only, but can also have positive effects, as evident in its anti-depressant potentials (Boland et al., 2017).

Limitations

Although in the present study, the use of cognitive reappraisal (CRA) strategies was effective in toning down feelings of social exclusion (Gross, 2007), CRA was always applied in the second session. Therefore, we cannot rule out that habituation to the task during the second session influenced the regulation of emotionality by CRA. However, counterbalancing the view and reappraisal session would have introduced even stronger confounds, considering that participants would have most likely also engaged in CRA in the second session if they had learned about this strategy in the first session. In order to better discriminate between CRA and habituation effects one could apply a three-session-design and randomly instruct participants to use CRA either in the second or third session. In a larger sample than ours, Mauss and colleagues applied such a design to show that poorer sleep quality over the course of the last week was linked to decreased abilities in engaging in CRA (Mauss et al., 2013).

A second limitation may be the presence of social desirability effects. While CRA is regarded as the most efficient emotion regulation technique (Gross, 2001), the experimenters' intention may have been too obvious in the present social exclusion context, which might have increased social desirability effects, rendering this emotion regulation strategy suboptimal. Alternatively, in a different experimental setup, one could implicitly offer subjects an opportunity for diverting their attention away from the unpleasant emotional experience in one session and thereby apply another emotion regulation technique (i.e. suppression) without an explicit instruction and the danger of social desirability effects. Future studies will need to specifically target the effect of different emotion regulation strategies and the impact that selective REMS might exert on such techniques.

Conclusion

In conclusion, the present study examined the effect of selective REMS on general affect as well as the neural correlates of task-induced negative emotionality and the ability to regulate emotional experiences by cognitive reappraisal during experimentally induced social exclusion. While we found that

REMS predicted general negative affect in the next morning, we did not find experimental evidence for

task-induced changes of negative emotionality during the experience of social exclusion. However, limbic

system activity during social exclusion was significantly increased in REM sleep suppressed participants

pointing to a mechanistic link between REM sleep disturbances and affect. Further investigations of the

underlying mechanisms of REM sleep and emotional dysregulation in social situations will pave the way

for a better understanding of the role of sleep disturbance in psychiatric disorders.

Author contributions

Prepared and conceived the study: W.C., K.K., U.K., S.K., S.W.; Collected the data: S.K., F.M.P., S.W;

Analyzed the fMRI and behavioral data: R.G., F.M.P., L.M.P., S.K., D.S.S.; Analyzed the PSG data: A.S.;

Interpreted the data and wrote the manuscript: S.B., S.D., R.G., S.K., F.M.P., L.M.P., D.S.S., I.W.

Acknowledgements

We thank Konrad Whittaker and Tareq Naji for their help in conducting the polysomnography

measurements and Christian Lange for support in scoring the PSG data.

References

Amaral, D. G., & Price, J. L. (1984). Amygdalo-cortical projections in the monkey (Macaca fascicularis).

Journal of Comparative Neurology, *230*(4), 465–496. https://doi.org/10.1002/cne.902300402

Armitage, R. (2007). Sleep and circadian rhythms in mood disorders. *Acta Psychiatrica Scandinavica.*

Supplementum, *433*, 104–115. https://doi.org/10.1111/j.1600-0447.2007.00968.x

Ashburner, J., Barnes, G., Chen, C., Daunizeau, J., Moran, R., Henson, R., Glauche, V., & Phillips, C. (2013).

SPM12 Manual The FIL Methods Group (and honorary members). *Functional Imaging Laboratory*,

475–1. https://doi.org/10.1111/j.1365-294X.2006.02813.x

Ashworth, D. K., Sletten, T. L., Junge, M., Simpson, K., Clarke, D., Cunnington, D., & Rajaratnam, S. M. W.

(2015). A randomized controlled trial of cognitive behavioral therapy for insomnia: An effective

treatment for comorbid insomnia and depression. *Journal of Counseling Psychology*, *62*(2), 115–

123. https://doi.org/10.1037/cou0000059

Baran, B., Pace-Schott, E. F., Ericson, C., & Spencer, R. M. C. (2012). Processing of Emotional Reactivity

and Emotional Memory over Sleep. *The Journal of Neuroscience*, *32*(3), 1035–1042.

Beedie, C. J., Terry, P. C., & Lane, A. M. (2005). Distinctions between emotion and mood. *Cognition and*

Emotion, *19*(6), 847–878. https://doi.org/10.1080/02699930541000057

Ben Simon, E., Oren, N., Sharon, H., Kirschner, A., Goldway, N., Okon-Singer, H., Tauman, R., Deweese,

M. M., Keil, A., & Hendler, T. (2015). Losing Neutrality: The Neural Basis of Impaired Emotional

Control without Sleep. *Journal of Neuroscience*, *35*(38), 13194–13205.

https://doi.org/10.1523/JNEUROSCI.1314-15.2015

Ben Simon, Eti, & Walker, M. P. (2018). Sleep loss causes social withdrawal and loneliness. *Nature*

Communications, *9*(1), 3146. https://doi.org/10.1038/s41467-018-05377-0

Berking, M., & Wupperman, P. (2012). Emotion regulation and mental health: recent findings, current

challenges, and future directions. *Current Opinion in Psychiatry*, *25*(2), 128–134.

https://doi.org/10.1097/YCO.0b013e3283503669

Berry, R. B., Brooks, R., Gamaldo, C. E., Harding, S. M., Lloyd, R. M., & Marcus, C. L. (2015). *The AASM*

Manual for the Scoring of Sleep and Associated Events: Rules, Terminology, and Technical

Specifications (2.2). American Academy of Sleep Medicine.

Besedovsky, L., Lange, T., & Haack, M. (2019). The sleep-immune crosstalk in health and disease.

Physiological Reviews, *99*(3), 1325–1380. https://doi.org/10.1152/physrev.00010.2018

Boland, E. M., Rao, H., Dinges, D. F., Smith, R. V, Goel, N., Detre, J. A., Basner, M., Sheline, Y. I., Thase, M.

E., & Gehrman, P. R. (2017). Meta-Analysis of the Antidepressant Effects of Acute Sleep

Deprivation. *The Journal of Clinical Psychiatry*, *78*(8), e1020—e1034.

https://doi.org/10.4088/jcp.16r11332

Bolling, D. Z., Pitskel, N. B., Deen, B., Crowley, M. J., Mayes, L. C., & Pelphrey, K. A. (2011). Development

of neural systems for processing social exclusion from childhood to adolescence. *Developmental*

Science, *14*(6), 1431–1444. https://doi.org/10.1111/j.1467-7687.2011.01087.x

Briançon-Marjollet, A., Weiszenstein, M., Henri, M., Thomas, A., Godin-Ribuot, D., & Polak, J. (2015). The

impact of sleep disorders on glucose metabolism: endocrine and molecular mechanisms.

Diabetology & Metabolic Syndrome, *7*, 25. https://doi.org/10.1186/s13098-015-0018-3

Bryant, P. A., Trinder, J., & Curtis, N. (2004). Sick and tired: Does sleep have a vital role in the immune

system? *Nature Reviews Immunology*, *4*(6), 457–467. https://doi.org/10.1038/nri1369

Burgdorf, C., Rinn, C., & Stemmler, G. (2016). Effects of personality on the opioidergic modulation of the

emotion warmth-liking: Opioidergic Modulation of Warmth-Liking. *Journal of Comparative*

Neurology, *524*(8), 1712–1726. https://doi.org/10.1002/cne.23847

Bustamante, V., Fins, A. I., Pigeon, W. R., & Nolan, B. (2002). REM Sleep and the Early Development of

Posttraumatic Stress Disorder. *American Journal of Psychiatry*, *159*(October), 1696–1701.

https://doi.org/10.1176/appi.ajp.159.10.1696

Casey, S. J., Solomons, L. C., Steier, J., Kabra, N., Burnside, A., Pengo, M. F., Moxham, J., Goldstein, L. H.,

& Kopelman, M. D. (2016). Slow wave and REM sleep deprivation effects on explicit and implicit

memory during sleep. *Neuropsychology*, *30*(8), 931–945. https://doi.org/10.1037/neu0000314

Cerliani, L., Thomas, R. M., Jbabdi, S., Siero, J. C. W., Nanetti, L., Crippa, A., Gazzola, V., D'Arceuil, H., &

Keysers, C. (2012). Probabilistic tractography recovers a rostrocaudal trajectory of connectivity

variability in the human insular cortex. *Human Brain Mapping*, *33*(9), 2005–2034.

https://doi.org/10.1002/hbm.21338

Colvonen, P. J., Straus, L. D., Acheson, D., & Gehrman, P. (2019). A Review of the Relationship Between

Emotional Learning and Memory, Sleep, and PTSD. *Current Psychiatry Reports*, *21*(1), 2.

https://doi.org/10.1007/s11920-019-0987-2

Crawford, J. R., & Henry, J. D. (2004). The Positive and Negative Affect Schedule (PANAS): Construct

validity, measurement properties and normative data in a large non-clinical sample. *British Journal*

of Clinical Psychology, *43*(3), 245–265. https://doi.org/10.1348/0144665031752934

Davidson, R. J. (2002). Anxiety and affective style: role of prefrontal cortex and amygdala. *Biological*

Psychiatry, *51*(1), 68–80. https://doi.org/10.1016/S0006-3223(01)01328-2

Dewall, C. N., Macdonald, G., Webster, G. D., Masten, C. L., Baumeister, R. F., Powell, C., Combs, D.,

Schurtz, D. R., Stillman, T. F., Tice, D. M., & Eisenberger, N. I. (2010). Acetaminophen reduces social

pain: behavioral and neural evidence. *Psychological Science*, *21*(7), 931–937.

https://doi.org/10.1177/0956797610374741

Dewall, C. N., Masten, C. L., Powell, C., Combs, D., Schurtz, D. R., & Eisenberger, N. I. (2012). Do neural

responses to rejection depend on attachment style? An fMRI study. *Social Cognitive and Affective*

Neuroscience, *7*(2), 184–192. https://doi.org/10.1093/scan/nsq107

Diekelmann, S., & Born, J. (2010). The memory function of sleep. *Nature Reviews Neuroscience*, *11*(2),

114–126. https://doi.org/10.1038/nrn2762

Downey, G., Mougios, V., Ayduk, O., London, B. E., & Shoda, Y. (2004). Rejection Sensitivity and the

Defensive Motivational System: Insights From the Startle Response to Rejection Cues. *Psychological*

Science, *15*(10), 668–673. https://doi.org/10.1111/j.0956-7976.2004.00738.x

Durmer, J. S., & Dinges, D. F. (2005). Neurocognitive consequences of sleep deprivation. In *Seminars in*

Neurology (Vol. 25, Issue 1, pp. 117–129). https://doi.org/10.1055/s-2005-867080

Eisenberger, N. I., & Lieberman, M. D. (2004). Why rejection hurts: A common neural alarm system for

physical and social pain. *Trends in Cognitive Sciences*, *8*(7), 294–300.

https://doi.org/10.1016/j.tics.2004.05.010

Eisenberger, N. I., Lieberman, M. D., & Williams, K. D. (2003). Does rejection hurt? An FMRI study of

social exclusion. *Science (New York, N.Y.)*, *302*(5643), 290–292.

https://doi.org/10.1126/science.1089134

Franzen, P. L., & Buysse, D. J. (2017). Sleep in Psychiatric Disorders. In S. Chokroverty (Ed.), *Sleep*

Disorders Medicine (pp. 977–996). Springer New York.

Geoffroy, P. A., Hoertel, N., Etain, B., Bellivier, F., Delorme, R., Limosin, F., & Peyre, H. (2018). Insomnia

and hypersomnia in major depressive episode: Prevalence, sociodemographic characteristics and

psychiatric comorbidity in a population-based study. *Journal of Affective Disorders*, *226*, 132–141.

https://doi.org/10.1016/j.jad.2017.09.032

Goldstein, A. N., & Walker, M. P. (2014). The Role of Sleep in Emotional Brain Function. *Annual Review of*

Clinical Psychology, *10*(1), 679–708. https://doi.org/10.1146/annurev-clinpsy-032813-153716

Gottesmann, C., & Gottesman, I. (2007). The neurobiological characteristics of rapid eye movement

(REM) sleep are candidate endophenotypes of depression, schizophrenia, mental retardation and

dementia. *Progress in Neurobiology*, *81*(4), 237–250.

https://doi.org/10.1016/j.pneurobio.2007.01.004

Gross, J. J. (2001). Emotion Regulation in Adulthood: Timing Is Everything. *Current Directions in*

Psychological Science, *10*(6), 214–219. https://doi.org/10.1111/1467-8721.00152

Gross, J. J. (2002). Emotion regulation: Affective, cognitive, and social consequences. *Psychophysiology*,

39(3), S0048577201393198. https://doi.org/10.1017/S0048577201393198

Gross, J. J. (2007). *Handbook of emotion regulation*. The Guilford Press.

Gross, J. J. (2008). Emotion Regulation. In M. Lewis, J. M. Haviland-Jones, & L. Feldman Barrett (Eds.),

Handbook of Emotions (3rd ed., pp. 497–512). The Guilford Press.

Gross, J. J., & Muñoz, R. F. (1995). Emotion Regulation and Mental Health. *Clinical Psychology: Science*

and Practice, *2*(2), 151–164. https://doi.org/10.1111/j.1468-2850.1995.tb00036.x

Habukawa, M., Uchimura, N., Maeda, M., Ogi, K., Hiejima, H., & Kakuma, T. (2018). Differences in rapid

eye movement (REM) sleep abnormalities between posttraumatic stress disorder (PTSD) and major

depressive disorder patients: REM interruption correlated with nightmare complaints in PTSD.

Sleep Medicine, *43*, 34–39. https://doi.org/10.1016/j.sleep.2017.10.012

Howell, A. J., Digdon, N. L., Buro, K., & Sheptycki, A. R. (2008). Relations among mindfulness, well-being,

and sleep. *Personality and Individual Differences*, *45*(8), 773–777.

https://doi.org/10.1016/j.paid.2008.08.005

Kobayashi, I., Boarts, J. M., & Delahanty, D. L. (2007). Polysomnographically measured sleep

abnormalities in PTSD: a meta-analytic review. *Psychophysiology*, *44*(4), 660–669.

https://doi.org/10.1111/j.1469-8986.2007.537.x

Koole, S. (2009). The psychology of emotion regulation: An integrative review. *Cognition & Emotion*,

23(1), 4–41. https://doi.org/10.1080/02699930802619031

Koren, D., Dumin, M., & Gozal, D. (2016). Role of sleep quality in the metabolic syndrome. *Diabetes,*

Metabolic Syndrome and Obesity : Targets and Therapy, *9*, 281–310.

https://doi.org/10.2147/DMSO.S95120

Krach, S., Müller-Pinzler, L., Westermann, S., & Paulus, F. M. (2013). Advancing the neuroscience of social

emotions with social immersion. *Behavioral and Brain Sciences*, *36*(04), 427–428.

https://doi.org/10.1017/S0140525X12001951

Krause, A. J., Simon, E. Ben, Mander, B. A., Greer, S. M., Saletin, J. M., Goldstein-Piekarski, A. N., &

Walker, M. P. (2017). The sleep-deprived human brain. *Nature Reviews Neuroscience*, *18*(7), 404–

418. https://doi.org/10.1038/nrn.2017.55

Kryger, M., Roth, T., & Dement, W. C. (2017). *Principles and Practice of Sleep Medicine* (6th ed.). Elsevier.

https://doi.org/10.1016/C2012-0-03543-0

Kunimatsu, A., Yasaka, K., Akai, H., Kunimatsu, N., & Abe, O. (2019). MRI findings in posttraumatic stress

disorder. *Journal of Magnetic Resonance Imaging : JMRI*. https://doi.org/10.1002/jmri.26929

Lancaster, J. L., Summerlin, J. L., Rainey, L., Freitas, C. S., & Fox, P. T. (1997). The Talairach Daemon a

database server for talairach atlas labels. *NeuroImage*, *5*(4 PART II).

Lancaster, J. L., Woldorff, M. G., Parsons, L. M., Liotti, M., Freitas, C. S., Rainey, L., Kochunov, P. V.,

Nickerson, D., Mikiten, S. A., & Fox, P. T. (2000). Automated Talairach Atlas labels for functional

brain mapping. *Human Brain Mapping*, *10*(3), 120–131. https://doi.org/10.1002/1097-

0193(200007)10:3<120::AID-HBM30>3.0.CO;2-8

Lange, T., Dimitrov, S., & Born, J. (2010). Effects of sleep and circadian rhythm on the human immune

system. *Annals of the New York Academy of Sciences*, *1193*(1), 48–59.

https://doi.org/10.1111/j.1749-6632.2009.05300.x

LeDoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience*, *23*, 155–184.

LeDoux, J. E. (2020). How does the non-conscious become conscious? *Current Biology*, *30*, R196–R199.

https://doi.org/10.1016/j.cub.2020.01.033

Lindquist, K. A., & Barrett, L. F. (2008). Constructing Emotion. *Psychological Science*, *19*(9), 898–903.

https://doi.org/10.1111/j.1467-9280.2008.02174.x

Lindquist, K. A., Satpute, A. B., Wager, T. D., Weber, J., & Barrett, L. F. (2016). The Brain Basis of Positive

and Negative Affect: Evidence from a Meta-Analysis of the Human Neuroimaging Literature.

Cerebral Cortex (New York, N.Y. : 1991), *26*(5), 1910–1922. https://doi.org/10.1093/cercor/bhv001

Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F. (2012). The brain basis of

emotion: A meta-analytic review. *Behavioral and Brain Sciences*, *35*(03), 121–143.

https://doi.org/10.1017/S0140525X11000446

Lipinska, G., & Thomas, K. G. F. (2019). The Interaction of REM Fragmentation and Night-Time Arousal

Modulates Sleep-Dependent Emotional Memory Consolidation. *Frontiers in Psychology*, *10*.

https://doi.org/10.3389/fpsyg.2019.01766

Liu, J. C. J., Mulick, D., & Chee, M. W. L. (2014). Odd one out: social ostracism affects self-reported needs

in both sleep-deprived and well-rested persons. *Journal of Sleep Research*, *23*(4), 448–457.

https://doi.org/10.1111/jsr.12141

MacDonald, G., & Leary, M. R. (2005). Why Does Social Exclusion Hurt? The Relationship Between Social

and Physical Pain. *Psychological Bulletin*, *131*(2), 202–223. https://doi.org/10.1037/0033-

2909.131.2.202

Maldjian, J. A., Laurienti, P. J., & Burdette, J. H. (2004). Precentral gyrus discrepancy in electronic versions

of the Talairach atlas. *NeuroImage*, *21*(1), 450–455.

https://doi.org/10.1016/j.neuroimage.2003.09.032

Maldjian, J. A., Laurienti, P. J., Kraft, R. A., & Burdette, J. H. (2003). An automated method for

neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *NeuroImage*.

https://doi.org/10.1016/S1053-8119(03)00169-1

Manber, R., Edinger, J. D., Gress, J. L., San Pedro-Salcedo, M. G., Kuo, T. F., & Kalista, T. (2008). Cognitive

behavioral therapy for insomnia enhances depression outcome in patients with comorbid major

depressive disorder and insomnia. *Sleep*, *31*(4), 489–495. https://doi.org/10.1093/sleep/31.4.489

Masten, C. L., Eisenberger, N. I., Borofsky, L. A., Pfeifer, J. H., McNealy, K., Mazziotta, J. C., & Dapretto, M.

(2009). Neural correlates of social exclusion during adolescence: Understanding the distress of peer

rejection. *Social Cognitive and Affective Neuroscience*, *4*(2), 143–157.

https://doi.org/10.1093/scan/nsp007

Mauss, I. B., Troy, A. S., & LeBourgeois, M. K. (2013). Poorer sleep quality is associated with lower

emotion-regulation ability in a laboratory paradigm. *Cognition & Emotion*, *27*(3), 567–576.

https://doi.org/10.1080/02699931.2012.727783

Mellman, T. A., Pigeon, W. R., Nowell, P. D., & Nolan, B. (2007). Relationships between REM sleep

findings and PTSD symptoms during the early aftermath of trauma. *Journal of Traumatic Stress*,

20(5), 893–901. https://doi.org/10.1002/jts.20246

Morgenthaler, J., Wiesner, C. D., Hinze, K., Abels, L. C., Prehn-Kristensen, A., & Göder, R. (2014). Selective

REM-sleep deprivation does not diminish emotional memory consolidation in young healthy

subjects. *PLoS ONE*, *9*(2), 1–7. https://doi.org/10.1371/journal.pone.0089849

Motomura, Y., Kitamura, S., Oba, K., Terasawa, Y., Enomoto, M., Katayose, Y., Hida, A., Moriguchi, Y.,

Higuchi, S., & Mishima, K. (2013). Sleep Debt Elicits Negative Emotional Reaction through

Diminished Amygdala-Anterior Cingulate Functional Connectivity. *PLoS ONE*, *8*(2).

https://doi.org/10.1371/journal.pone.0056578

Nezlek, J. B., Kowalski, R. M., Leary, M. R., Blevins, T., & Holgate, S. (1997). Personality Moderators of

Reactions to Interpersonal Rejection: Depression and Trait Self-Esteem. *Personality and Social*

Psychology Bulletin, *23*(12), 1235–1244.

Nielsen, T. A., Paquette, T., Solomonova, E., Lara-Carrasco, J., Popova, A., & Levrier, K. (2010). REM sleep

characteristics of nightmare sufferers before and after REM sleep deprivation. *Sleep Medicine*,

11(2), 172–179. https://doi.org/10.1016/j.sleep.2008.12.018

Nishida, M., Pearsall, J., Buckner, R. L., & Walker, M. P. (2009). REM Sleep, Prefrontal Theta, and the

Consolidation of Human Emotional Memory. *Cerebral Cortex*, *19*(May).

https://doi.org/10.1093/cercor/bhn155

Orem, T. R., Wheelock, M. D., Goodman, A. M., Harnett, N. G., Wood, K. H., Gossett, E. W., Granger, D.

A., Mrug, S., & Knight, D. C. (2019). Amygdala and prefrontal cortex activity varies with individual

differences in the emotional response to psychosocial stress. *Behavioral Neuroscience*, *133*(2), 203–

211. https://doi.org/10.1037/bne0000305

Pace-Schott, E. F., Germain, A., & Milad, M. R. (2015). Sleep and REM sleep disturbance in the

pathophysiology of PTSD: the role of extinction memory. *Biology of Mood & Anxiety Disorders*, *5*, 3.

https://doi.org/10.1186/s13587-015-0018-9

Paré, D., Collins, D. R., & Pelletier, J. G. (2002). Amygdala oscillations and the consolidation of emotional

memories. *Trends in Cognitive Sciences*, *6*(7), 306–314.

Pessoa, L., & Adolphs, R. (2010). Emotion processing and the amygdala: From a "low road" to "many

roads" of evaluating biological significance. *Nature Reviews Neuroscience*, *11*(11), 773–782.

https://doi.org/10.1038/nrn2920

Phan, K. L., Wager, T. D., Taylor, S. F., & Liberzon, I. (2004). Functional neuroimaging studies of human

emotions. *CNS Spectrums*, *9*(4), 258–266. https://doi.org/10.1017/S1092852900009196

Phelps, E. A. (2006). Emotion and Cognition: Insights from Studies of the Human Amygdala. *Annual*

Review of Psychology, *57*(1), 27–53. https://doi.org/10.1146/annurev.psych.56.091103.070234

Seeley, W. W., Menon, V., Schatzberg, A. F., Keller, J., Glover, G. H., Kenna, H., Reiss, A. L., & Greicius, M.

D. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control.

The Journal of Neuroscience : The Official Journal of the Society for Neuroscience, *27*(9), 2349–2356.

https://doi.org/10.1523/JNEUROSCI.5587-06.2007

Spoormaker, V. I., Schröter, M. S., Andrade, K. C., Dresler, M., Kiem, S. A., Goya-Maldonado, R., Wetter,

T. C., Holsboer, F., Sämann, P. G., & Czisch, M. (2012). Effects of rapid eye movement sleep

deprivation on fear extinction recall and prediction error signaling. *Human Brain Mapping*, *33*(10),

2362–2376. https://doi.org/10.1002/hbm.21369

Staner, L. (2010). Comorbidity of insomnia and depression. *Sleep Medicine Reviews*, *14*(1), 35–46.

https://doi.org/10.1016/j.smrv.2009.09.003

Steptoe, A., O'Donnell, K., Marmot, M., & Wardle, J. (2008). Positive affect, psychological well-being, and

good sleep. *Journal of Psychosomatic Research*, *64*(4), 409–415.

https://doi.org/10.1016/j.jpsychores.2007.11.008

Stickgold, R., & Walker, M. P. (2013). Sleep-dependent memory triage: Evolving generalization through

selective processing. In *Nature Neuroscience* (Vol. 16, Issue 2, pp. 139–145).

https://doi.org/10.1038/nn.3303

Tassi, P., & Muzet, A. (2000). Sleep inertia. *Sleep Medicine Reviews*, *4*(4), 341–353.

https://doi.org/10.1053/smrv.2000.0098

Taylor, S. F., Phan, K. L., Decker, L. R., & Liberzon, I. (2003). Subjective rating of emotionally salient

stimuli modulates neural activity. *NeuroImage*, *18*(3), 650–659. https://doi.org/10.1016/S1053-

8119(02)00051-4

Urry, H. L., van Reekum, C. M., Johnstone, T., Kalin, N. H., Thurow, M. E., Schaefer, H. S., Jackson, C. A.,

Frye, C. J., Greischar, L. L., Alexander, A. L., & Davidson, R. J. (2006). Amygdala and Ventromedial

Prefrontal Cortex Are Inversely Coupled during Regulation of Negative Affect and Predict the

Diurnal Pattern of Cortisol Secretion among Older Adults. *Journal of Neuroscience*, *26*(16), 4415–

4425. https://doi.org/10.1523/JNEUROSCI.3215-05.2006

van der Helm, E., Yao, J., Dutt, S., Rao, V., Saletin, J. M., & Walker, M. P. (2011). REM Sleep Depotentiates

Amygdala Activity to Previous Emotional Experiences. *Current Biology*, *21*(23), 2029–2032.

https://doi.org/10.1016/j.cub.2011.10.052

Van Der Werf, Y. D., Altena, E., Schoonheim, M. M., Sanz-Arigita, E. J., Vis, J. C., De Rijke, W., & Van

Someren, E. J. W. (2009). Sleep benefits subsequent hippocampal functioning. *Nature*

Neuroscience, *12*(2), 122–123. https://doi.org/10.1038/nn.2253

Vandekerckhove, M., & Cluydts, R. (2010). The emotional brain and sleep: An intimate relationship. *Sleep*

Medicine Reviews, *14*(4), 219–226. https://doi.org/10.1016/j.smrv.2010.01.002

Wagner, U. (2002). Changes in Emotional Responses to Aversive Pictures Across Periods Rich in Slow-

Wave Sleep Versus Rapid Eye Movement Sleep. *Psychosomatic Medicine*, *64*(4), 627–634.

https://doi.org/10.1097/01.PSY.0000021940.35402.51

Wagner, U., Gais, S., & Born, J. (2001). Emotional Memory Formation Is Enhanced across Sleep Intervals

with High Amounts of Rapid Eye Movement Sleep. *Learning & Memory*, *8*(2), 112–119.

https://doi.org/10.1101/lm.36801

Walker, M. P. (2008). Cognitive consequences of sleep and sleep loss. *Sleep Medicine*, *9*(SUPPL. 1).

https://doi.org/10.1016/S1389-9457(08)70014-5

Walker, M. P. (2009). The role of sleep in cognition and emotion. *Annals of the New York Academy of*

Sciences, *1156*, 168–197. https://doi.org/10.1111/j.1749-6632.2009.04416.x

Wiesner, C. D., Pulst, J., Krause, F., Elsner, M., Baving, L., Pedersen, A., Prehn-Kristensen, A., & Göder, R.

(2015). The effect of selective REM-sleep deprivation on the consolidation and affective evaluation

of emotional memories. *Neurobiology of Learning and Memory*, *122*, 131–141.

https://doi.org/10.1016/j.nlm.2015.02.008

Williams, K. D. (2007). Ostracism. *Annual Review of Psychology*, *58*(1), 425–452.

https://doi.org/10.1146/annurev.psych.58.110405.085641

Williams, K. D., & Jarvis, B. (2006). Cyberball: A program for use in research on interpersonal ostracism

and acceptance. *Behavior Research Methods*, *38*(1), 174–180. https://doi.org/10.3758/BF03192765

Yoo, S.-S., Gujar, N., Hu, P., Jolesz, F. A., & Walker, M. P. (2007a). The human emotional brain without

sleep – a prefrontal-amygdala disconnect. *Current Biology*, *17*(20), 1–8.

https://doi.org/10.1016/j.cub.2007.08.007

Yoo, S.-S., Gujar, N., Hu, P., Jolesz, F. A., & Walker, M. P. (2007b). The human emotional brain without

sleep — a prefrontal amygdala disconnect. *Current Biology*, *17*(20), R877–R878.

https://doi.org/10.1016/j.cub.2007.08.007

Zadro, L., Boland, C., & Richardson, R. (2006). How long does it last? The persistence of the effects of

ostracism in the socially anxious. *Journal of Experimental Social Psychology*, *42*(5), 692–697.

https://doi.org/10.1016/j.jesp.2005.10.007

Zohar, D., Tzischinsky, O., Epstein, R., & Lavie, P. (2005a). The effects of sleep loss on medical residents'

emotional reactions to work events: A cognitive-energy model. *Sleep*, *28*(1), 47–54.

https://doi.org/10.1093/sleep/28.1.47

Zohar, D., Tzischinsky, O., Epstein, R., & Lavie, P. (2005b). The Effects of Sleep Loss on Medical Residents'

Emotional Reactions to Work Events: a Cognitive-Energy Model. *Sleep*, *28*(1), 47–54.

https://doi.org/10.1093/sleep/28.1.47

Figures and captions

Figure 1. Summary of the experimental procedure. Three groups of subjects spent two consecutive nights in the sleep laboratory. During both nights, polysomnography was recorded. In the second night, two groups were woken up as soon as they entered REM sleep or SWS (groups REMS and SWSS, respectively), while the control group was not woken up (CTL). After waking up on the second morning, all subjects performed two sessions of the *Cyberball* game while inside the fMRI scanner (see Burgdorf et al., 2016, for a detailed description of the *Cyberball* paradigm). The game included a total of eight blocks, with four inclusion (INC) and four exclusion (EXC) blocks. During inclusion, subjects received the ball continuously throughout the block, while during exclusion, the other two players stopped throwing the ball to the participant soon after the beginning of the block, effectively excluding the participant from the game. Additionally, during the first four blocks (half INC, half EXC), subjects simply performed the task (VIEW), while during the second four blocks (half INC, half EXC) they were instructed to use cognitive reappraisal (CRA) to regulate their emotions.

Figure 2. Selective sleep suppression and general affect. **a** REM sleep was significantly more reduced in the REMS group than in the other two groups, whereas SWS suppression was significantly stronger in the SWSS group than in the other two groups. Suppression scores for REM (SWS) sleep are the difference of percentage REM (SWS) of TST between the habituation night minus the experimental night. **b** Ratings of general positive and negative affect, measured with the Positive And Negative Affect Scale (PANAS; Crawford & Henry, 2004) on the mornings after the habituation night and after the experimental night, separately for the three groups. **c** Changes in negative (right) but not positive affect (left) from the habituation night to the experimental night correlate significantly with REM sleep suppression scores across all groups. Shaded areas represent 95% confidence intervals for simple bivariate regression. Error bars represent standard error of the mean. n.s.=not significant.

Figure 3. Neural responses to ostracism are altered by selective REM sleep suppression. **Left** Neural responses for the contrast [EXC/VIEW>INC/VIEW] > [EXC/CRA>INC/CRA] differed significantly between the REMS group and the other two groups in the right amygdala, (peak MNI-coordinates: x=26, y=-2, z=-28, surviving FWE-correction at $p<.05$ inside the a priori mask). Displayed at $p<.005$, uncorrected, for visualization purposes. **Right** Parameter estimates for the contrast EXC>INC in the VIEW session were positive in the REMS group, but did not deviate from 0 in the other two groups, nor in the CRA session for any of the three groups. In addition, parameter estimates for EXC>INC were significantly more positive in the REMS group during VIEW than during CRA, while sessions did not differ in the other two groups. See main text for details. ***: $p<.001$. Error bars represent standard error of the mean.

 Tables

Table 1. Sleep measures

	CTL				REMS				SWSS			
	Habituation night		Experimental night		Habituation night		Experimental night		Habituation night		Experimental night	
n subjects with valid PSG dataset	15		15		16		17		9		10	
	M	*SD*	*M*	*SD*	*M*	*SD*	*M*	*SD*	*M*	*SD*	*M*	*SD*
TST (min)	369.94	78.06	414.59	26.08	410.86	40.24	399.05	39.02	384.63	54.15	402.96	31.28
N1 (% TST)	1.87	2.76	0.58	0.96	0.60	0.65	1.72	2.54	2.47	2.57	1.18	1.39
N2 (% TST)	61.82	7.27	61.11	8.53	64.83	4.69	68.66	7.62	65.68	8.44	69.33	8.29
SWS (% TST)	17.11	6.21	18.04	7.92	14.18	6.88	16.86	5.62	13.98	7.43	9.83	4.82
REM (% TST)	19.21	6.33	20.27	4.39	20.39	5.86	12.76	6.16	17.42	6.56	19.65	8.87
WASO (min)	67.16	50.46	51.06	29.00	47.01	40.18	71.10	42.66	57.57	38.50	61.63	37.96
WASO/(TST+WASO)	0.16	0.13	0.11	0.06	0.10	0.09	0.15	0.09	0.13	0.10	0.13	0.08
Awakenings	8.13	3.09	9.00	4.21	7.75	3.66	11.47	4.90	11.11	6.64	15.00	6.68
REM latency (min)	142.97	68.32	106.10	51.71	101.19	43.15	154.12	93.11	137.78	86.27	123.35	82.01

Note. In both the REMS group and the SWSS group, PSG data quality in the habituation night from one subject was not suitable for analysis. CTL=Control group with undisturbed sleep; REMS=rapid eye movement sleep suppression group; SWSS=slow wave sleep suppression group; TST=total sleep time (time from sleep onset to morning awakening after the time awake during the night is subtracted); N1=Sleep stage 1, N2=Sleep stage 2, SWS=Sleep stage N3; REM=rapid eye movement sleep; WASO=wakefulness after sleep onset to morning awakening; *M*=Mean; *SD*=Standard Deviation

Table 2. Regression of affective changes on sleep deprivation parameters (dependent variable: change in negative affect (∆ Experimental morning – Habituation morning))

	B	SE	β	t	p	sign.
(Intercept)	0.098	0.060		1.64	.110	n.s.
REM sleep suppression	0.020	0.008	0.406	2.60	.014	*
SWS sleep suppression	-0.000	0.008	-0.002	-0.01	.989	n.s.
TST suppression	0.002	0.002	0.404	1.60	.118	n.s.
WASO increase	0.005	0.002	0.638	2.95	.006	***

Note. n.s.: not significant; *: $p<.05$; ***: $p<.001$

Table 3. Feelings of being excluded, separately for groups and conditions

	EXC/VIEW		INC/VIEW		EXC/CRA		INC/CRA	
	M	*SD*	*M*	*SD*	*M*	*SD*	*M*	*SD*
CTL	6.67	1.92	2.20	1.25	*3.67*	*2.50*	1.50	0.78
REMS	6.38	2.90	2.00	1.21	*2.82*	*2.33*	1.62	0.86
SWSS	5.90	2.49	2.25	0.98	*2.60*	*1.51*	1.75	1.46