Lack of mucosal cholinergic innervation is associated with

increased risk of enterocolitis in Hirschsprung's disease

Simone Keck[1*], Virginie Galati-Fournier[1], Urs Kym[1], Michèle Moesch[1], Jakob

Usemann[2], Ulrike Subotic[1,3], Sasha J. Tharakan[3], Thomas Krebs[4], Eleuthere

Stathopoulos[5], Peter Schmittenbecher[6], Dietmar Cholewa[7], Philipp Romero[8],

Bertram Reingruber[9], Elisabeth Bruder[10], NIG study Group[11,12] and Stefan Holland-

Cunz[1]

*Corresponding author: simone.keck@unibas.ch

[1] Department of Pediatric Surgery, University Children's Hospital Basel (UKBB) and

University of Basel, Basel, Switzerland

[2] Department of Pediatric Pulmonology, University Children's Hospital Basel (UKBB),

Basel and Division of Respiratory Medicine, University Children's Hospital Zurich,

Zurich, Switzerland

[3] Department of Pediatric Surgery, University Children's Hospital Zurich, Zurich,

Switzerland

[4] Department of Pediatric Surgery, Children's Hospital of Eastern Switzerland, St.

Gallen, Switzerland

[5] Department of Pediatric Surgery, University Hospital of Lausanne (CHUV),

Lausanne, Switzerland

[6] Department of Pediatric Surgery, Municipal Hospital, Karlsruhe, Germany

[7] Department of Pediatric Surgery, University Hospital of Bern, Bern, Switzerland

[8] Department of Pediatric Surgery, University Hospital of Heidelberg, Heidelberg,

Germany

[9] Department of Pediatric Surgery, Florence Nightingale Hospital, Düsseldorf,

Germany

[10] Department of Medical Genetics and Pathology, University Hospital Basel, Basel,

Switzerland

[11] NIG Study Group: Isabella Bielicki[1], Sandra Weih[1], Ueli Moehrlen[3], Noëmi Zweifel[3],

Kai-Uwe Kleitsch[4], Carole Gengler[12], Kiarasch Mortazawi[6], Milan Milosevic[7], Vera

Otten[9], Lennart Homrighausen[9]

[12] Department of Pathology, University Hospital of Lausanne (CHUV) and University

of Lausanne, Lausanne, Switzerland

Conflict of interest

38 The authors disclose no conflicts.

Author contributions

S.K.: study design; design, execution, and supervision of the experiments; data

acquisition, analysis, and interpretation; writing and revision of the manuscript;

V.G-F.: execution of the experiments; data acquisition and analysis; contribution to

manuscript drafting; U.K.: execution of the experiments; data acquisition and

analysis; M.M.: execution of the experiments; data acquisition and analysis of

retrospective patient cohort; J.U.: statistical analysis; E.B.: acquisition of patient

Background & Aims

Hirschsprung's disease (HSCR) is a congenital intestinal motility disorder defined by the absence of enteric nervous cells (ganglia). The development of HSCR-associated enterocolitis remains a life-threatening complication. Absence of enteric ganglia implicates extramural innervation of acetylcholine-secreting (cholinergic) nerve fibers. Cholinergic signals have been reported to control excessive inflammation, but the impact on HSCR-associated enterocolitis is unknown.

Methods

We enrolled 44 HSCR patients in a prospective multicenter study and grouped them according to their degree of colonic mucosal cholinergic innervation using immunohistochemistry. The fiber phenotype was correlated with the tissue cytokine profile as well as immune cell frequencies using quantitative reverse-transcribed real-time polymerase chain reaction (qRT-PCR) of whole colonic tissue and fluorescence-activated cell sorting (FACS) analysis of isolated colonic immune cells. Fiber-associated immune cells were identified using confocal immunofluorescence microscopy and characterized by RNA-seq analysis. Microbial dysbiosis was analyzed in colonic patient tissue using 16S rDNA gene sequencing. Finally, the fiber phenotype was correlated with postoperative enterocolitis manifestation.

Results

We provided evidence that extrinsic mucosal innervation correlated with reduced interleukin (IL)-17 cytokine levels and T-helper-17 (Th17) cell frequencies. Bipolar CD14^{high} macrophages colocalized with neurons and expressed significantly less interleukin-23, a Th17-promoting cytokine. HSCR patients lacking mucosal

74 cholinergic nerve fibers showed microbial dysbiosis and had a higher incidence of

75 postoperative enterocolitis.

Conclusion

77 The mucosal fiber phenotype might serve as a new prognostic marker for

78 enterocolitis development in HSCR patients and may offer an approach to

79 personalized patient care and new future therapeutic options.

80 (www.clinicaltrials.gov accessing number NCT03617640)

Keywords

83 Enterocolitis, Neuroimmunology, Cholinergic neurons, Macrophages, Th17 cells,

84 Microbiome

Abbreviations used in this paper

Hirschsprung's disease (HSCR), Fluorescence-activated cell sorting (FACS),

Quantitative reverse-transcribed real-time polymerase chain reaction (qRT-PCR),

Interleukin (IL), T-helper-17 (Th17), Enteric nervous system (ENS), Acetylcholine

(ACh), α7-nicotinic ACh receptor (α7nAChR), Cholinergic-anti-inflammatory-pathway

(CAIP), Vagus nerve (VN), Macrophages (MΦ), Muscle macrophages (MMΦ), tumor

necrosis factor (TNF), Postoperative ileus (POI), Colon descendens (CD), Rectum

(REC), Sigmoid colon (SC), Vasoactive intestinal peptide (VIP), Tyrosine

hydroxylase (TH), Nitric oxide synthase (NOS), Natural killer (NK), NK cell receptors

(NCR), Antigen-presenting cell (APC), Principal component analysis (PCA), Human

leucocyte antigen (HLA), Growth/differentiation factor (GDF), Neuropilin-1 (NRP1),

UNC-5 homology B (UNC5B), Ephrin type B-receptor 2 (EPHB2), Semaphorin

(SEMA), Lamina propria (LP), Fluorescence in situ hybridization (FISH), Operational

taxonomic units (OTUs), Bone morphogenetic protein (BMP), Toll-like receptor (TLR)

Main

Hirschsprung's disease (HSCR) is a multigenetic congenital malformation of the

colon characterized by a distal lack of enteric ganglia cells (aganglionosis) leading to

intestinal obstruction and prestenotic megacolon. In affected infants, the aganglionic

bowel part is surgically removed, and healthy ganglionic tissue is pulled through and

reconnected to the anus. Either pre- or postoperatively, 20% to 50% of patients

suffer from life-threatening HSCR-associated enterocolitis leading to acute

abdominal distension, diarrhea, and fever up to sepsis and organ failure[1-3]. The

underlying pathogenesis is not fully understood but appears to involve changes in

intestinal barrier function[4-6], immune response[7-9], and the microbiome[10-15].

As a consequence of lacking an intrinsic enteric nervous system (ENS), extrinsic

acetylcholine (ACh)-producing (cholinergic) nerve fibers originating from sacral roots

S2-S4 innervate the distal colon. ACh released from these fibers is responsible for

constant muscle contraction in HSCR patients[16].

Intestinal cholinergic signals not only coordinate physiologic bowel function but

additionally prevent excessive inflammation by allowing the intestinal immune

system to tolerate harmless commensals and food antigens[17]. The cholinergic-anti-

inflammatory-pathway (CAIP) was first described by the Tracey group[18]. In a model

of systemic inflammation, vagus nerve (VN) stimulation favored the release of ACh

by memory T cells which in turn bound to α7-nicotinic ACh receptor (α7nAChR)-

positive splenic macrophages (MΦ) to reduce tumor necrosis factor (TNF)-α

production[18,19]. This concept was confirmed in intestinal inflammation models

involving colitis[20-23], food allergy[24], postoperative ileus (POI)[25], and pancreatitis using

VN stimulation or pharmacologic targeting of α7nAChR. In humans, nicotine, which

activates nAChRs, has been associated with clinical improvement of ulcerative

colitis[26]. VN stimulation in Crohn's disease results in amelioration of disease

activity[27]. Contrary to systemic CAIP, intestinal modulation is independent of the

spleen and involves direct activation of local enteric neurons[25,28,29]. Although VN

activation leads to downregulation of inflammatory cytokine production by muscularis

MΦ via α7AChR activation in POI, it remains unclear which receptors and cell types

are involved in colitis amelioration[30,31]. Given the fact that the ENS utilizes more than

30 different neurotransmitters, it is challenging to identify individual immune-

modulating functions. As HSCR patients miss these ENS-specific neurotransmitters,

the study of cholinergic neurons is possible.

Here, we investigated the role of cholinergic mucosal innervation in regulating the

immune cell status, microbial composition, and development of enterocolitis in HSCR

patients.

Materials and Methods

140 Detailed protocols are provided in the Supplementary Materials and Methods

Results

Distinct cholinergic innervation of the colon in HSCR patients

144 In this prospective, multicenter study, we collected fresh tissue of different colonic

145 segments from 44 pediatric HSCR patients undergoing therapeutic pull-through

146 surgery as well as 6 control patients (non-HSCR) undergoing surgery for

147 miscellaneous reasons. The length of aganglionic colon was determined by

148 experienced pathologists of the respective pathology units (see Methods section,

149 Supplementary Table 1 and Supplementary Table 2)[32]. Compared to non-HSCR

150 colonic tissue, the aganglionic colon of HSCR patients lack the myenteric and

151 submucosal plexus but show the characteristic increase of ACh esterase (AChE)

152 activity in the rectosigmoid (Fig. 1a). Increased cholinergic activity is limited to the

153 aganglionic rectosigmoid distal colon and decreases in caudocranial direction[16,33]. As

154 a consequence, no extrinsic mucosal AChE$^+$ nerve fibers can be detected in the

155 proximal aganglionic colon descendens (CD), with only few present in the

156 intermuscular space. Depending on the length of aganglionosis, the proximal CD

157 segment of HSCR patients can be ganglionic or aganglionic (Fig. 1a and

158 Supplementary Table 1). CD segments showing a hypoganglionic phenotype

159 (transition zone) were excluded from the analysis.

160 We performed anti-AChE immunohistochemistry staining in cryosections of

161 longitudinally rolled colonic stripes (Swiss roll), including rectum (REC), sigmoid

162 colon (SC), and colon descendens (CD) from HSCR patients. All patient tissues

163 showed the typical thick AChE$^+$ fibers in the muscle region, but the density of thin

164 mucosal fibers varied between patients (Fig. 1b, c). According to mucosal

165 innervation in the distal rectosigmoid colon, the sections were semi-quantitatively

166 allocated into "fiber-low" (Fig. 1b) and "fiber-high" (Fig. 1c) patient groups in a

167 double-blind investigation (see methods for details). Fiber phenotype was validated

168 using tubulin immunohistochemistry (Supplementary Figure 1). Consequently, for

169 further analysis, aganglionic and ganglionic segments with variable mucosal

170 cholinergic innervation were defined as illustrated by anti-β3tubulin (pan neuronal

171 marker) and anti-AChE (cholinergic marker) immunofluorescent staining (Fig. 2a).

Thus, rectosigmoid fiber-low tissues were those lacking intrinsic nerve cell bodies as

well as extrinsic mucosal cholinergic innervation. Rectosigmoid Fiber-high tissues

were those lacking intrinsic nerve cell bodies but showing extrinsic mucosal

cholinergic innervation (Fig. 2a, arrow). Aganglionic CD sections from HSCR patients

were defined as those lacking intrinsic nerve cell bodies as well as extrinsic mucosal

cholinergic innervation (because of extrinsic nerve fiber restriction to the

rectosigmoid). Hence, aganglionic CD segments were used from both fiber-low and

fiber-high patient tissue. Ganglionic CD sections from HSCR and non-HSCR patients

showed the presence of non-cholinergic intrinsic nerve cell bodies (ganglia) with

mucosal nerve terminals (Fig. 2a, asterisk) and were considered as normally

innervated. Secondary antibody controls showed a high background in the epithelial

crypt region for AChE due to biotin/streptavidin amplification (Fig. 2d). By performing

immunofluorescence studies in fiber-high tissue, we could exclude peptidergic (VIP),

dopaminergic/adrenergic (TH), and nitrinergic (NOS) origins of the mucosal fibers

(Fig. 2b). However, we detected TH^+ nerve bundles in the submucosa and muscle

region. Sporadic fiber bundles were positive for the glia cell marker S100B, and the

majority of mucosal fibers were not associated with glia cells (Fig. 2b). Respective

secondary antibody controls were negative (Fig. 2c). We did not observe any

significant patient bias in terms of anthropometric, clinical, environmental, or

maternal factors between fiber-low and fiber-high tissues, except a higher

percentage of infants delivered by cesarean section in the fiber-high patient group

(Supplementary Table 3).

Absence of cholinergic fibers correlated with inflammatory immune status

A functional ENS is crucial for intestinal immune homeostasis[34]. To investigate if the

absence of neuronal signals impairs the overall inflammatory cytokine status in

HSCR patients, we performed quantitative real-time polymerase chain reaction (q-

RT PCR) analysis of whole colonic tissue from HSCR patients and control patients.

TNF-α did not significantly vary between the different colonic segments from HSCR

patients and control patients (Fig. 3a, Supplementary Table 4). Cytokines involved in

Rorγt$^+$ Th17 cell differentiation, i.e., IL-1β, IL-23, and IL-6[35,36], were significantly

elevated in aganglionic CD segments compared to ganglionic CD segments, but they

were not significantly changed compared to fiber-high segments. IL-17 and IL-22,

two Th17 effector cytokines, were significantly elevated in aganglionic CD sections

compared to ganglionic CD sections from HSCR patients (Fig. 3a, Supplementary

Table 4). Both cytokines were significantly attenuated in cholinergically innervated

fiber-high segments compared to fiber-low or aganglionic CD segments.

Beside innate lymphoid cells 3 (ILC3) and others, Th17 cells are the main producers

of IL-17 and IL-22 and are counter-regulated by forkhead box P3 positive (Foxp3$^+$)

regulatory T cells (Treg)[37-40]. Frequencies of mucosal IL17$^+$ Th17 and Foxp3$^+$ Treg T

cells were assessed by flow cytometry (Fig. 3 b, c, supplementary Fig. 3a, b). Th17

frequencies were significantly increased in fiber-low segments compared to fiber-

high segments or CD segments (Fig. 3b, Supplementary Fig. 2a showing the

percentage of total cells). We observed no difference between aganglionic and

ganglionic CD tissue of HSCR and control patients. Treg frequencies were

significantly elevated in both fiber-low and fiber-high tissues from HSCR patients

(Fig. 3c, Supplementary Fig. 2a showing the percentage of total cells). However, the

ratio of Th17/Treg was significantly higher in fiber-low segments compared to fiber-

high segments (Fig. 3d, Supplementary Fig. 2a showing the percentage of total cells).

Frequencies of natural killer (NK) cell receptors (NCR)$^+$ and NCR$^-$ ILC3s were

comparable between fiber-low and fiber-high tissues from HSCR patients. However,

the NCR⁻ ILC3 population was significantly elevated in fiber-low colonic segments

compared to ganglionic CD of HSCR patients (Supplementary Fig. 2b). Thus, lack of

an ENS appeared to have an impact on inflammatory cytokine production, whereas

lack of cholinergic neurons seemed to affect Th17 cells and their related cytokines.

CX3CR1⁺ MΦ represent the major antigen-presenting cell (APC) type in the colon

and are crucial for CD4 T cell differentiation[41]. These MΦ exhibit a non-inflammatory

M2-like phenotype and are essential for local expansion and maintenance of

Treg[42,14]. In contrast, intestinal CCR2⁺ CX3CR1⁺ blood-monocyte-derived MΦ are

essential for Th17 cell differentiation[43-47]. We performed qRT-PCR analysis in whole

colonic tissue to check for expression of CX3CR1 and CCR2. In fiber-high tissue as

well as in muscle tissue, we found a significantly higher expression of CX3CR1 than

CCR2 while in all other segments, expression levels of the two markers were

comparable (Supplementary Fig. 2c). To check whether these changes resulted from

altered MΦ populations, we investigated viable CD64⁺ MΦ by flow cytometry

(Supplementary Fig. 3d). All MΦ expressed comparable levels of CD11c (Fig. 3f,

upper graph). However, we found a significantly higher expression of CD14, an M2

marker[48], in MΦ isolated from the mucosal lamina propria (LPMΦ) of fiber-high

colonic tissue and in MΦ isolated from muscularis (MMΦ) (Fig. 3f, lower graph).

MMΦ interact with neurons of the myenteric plexus and have been described as

bipolarly shaped[49-51]. Using confocal fluorescence microscopy, we visualized a

bipolar shape of LPMΦ in fiber-high segments similar to that seen with MMΦ (Fig. 3g,

left and right image). In contrast, LPMΦ in fiber-low colonic segments exhibited a

round or stellate phenotype (Fig. 3g, middle image). We found a significantly higher

percentage of bipolar MΦ in fiber-high colonic tissue (Fig. 3g, bar graph). We

mimicked the phenotype of the two colonic macrophage populations by

differentiating blood-derived monocytes from healthy adult donors into M1 and M2

MΦ. After 10 days of differentiation, M2 MΦ showed an elongated bipolar shape

(Supplementary Fig. 2d, e) and increased expression of CD14 and CX3CR1

(Supplementary Fig. 2f, g). In contrast, M1 MΦ exhibited a rounded shape and

expressed moderate levels of CD14 while lacking CX3CR1 expression. To test the

ability of the MΦ to differentiate T cells into certain effector T cells, we co-cultured

M1 and M2 MΦ with autologous naïve CD4 T cells in the presence of anti-CD3 and

performed a Th17 and Treg T cell differentiation assay. After 6 days of co-culture,

the frequency of Th17 and Treg cells were analyzed and the ratio calculated.

Th17/Treg ratio was significantly lower when bipolar CX3CR1$^+$CD14^{high} M2 MΦ were

used (Supplementary Fig. 2h). These observations suggested that fiber-high

segments of HSCR colon harbor a bipolar CD64$^+$CD11c$^+$CD14^{high} MΦ population in

the mucosa which correlates with decreased Th17 cell frequencies. Of note, due to

the low viability of patient-derived colonic macrophages, no functional assays could

be performed, and M2 MΦ did not fully resemble the colonic phenotype.

Bipolar CD14^{high} MΦ colocalized with cholinergic fibers

Both MΦ and T cells express receptors for neurotransmitters, qualifying them for

neuronal interaction[25,49,50,52-54]. Representative fluorescence images (20x and 63x) of

fiber-high and fiber-low distal colonic tissue revealed CD64$^+$ MΦ and CD3$^+$ T cells in

close neighborhood to AChE$^+$ fibers (Fig. 4a). In the muscle layer, MΦ were only

detected between thick cholinergic fibers. To identify the cell type interacting with

mucosal nerve fibers, we performed fast-scanning confocal imaging using a Fiji

software macro to determine the sum of overlapping pixel values (Fig. 3b and

Supplementary Movie 1). Per patient (fiber-low segments, n=4; fiber-high segments,

n=5; muscle, n=2), 100-200 cells were annotated, and the sum of pixel values was

determined. The results showed that LPMΦ in fiber-high segments significantly

colocalized with cholinergic fibers comparable to MMΦ, in contrast to LPMΦ of fiber-

low segments (Fig. 4c). No significant overlap between CD3[+] T cells and cholinergic

fibers was found (Fig. 4d). Furthermore, neuron-associated LPMΦ in fiber-high

tissue from HSCR patients mainly exhibited a bipolar shape (Fig. 4e). These results

suggest that bipolar LPMΦ preferentially interacted with cholinergic fibers in

aganglionic fiber-high tissue from HSCR patients.

Cholinergic fibers suppressed Th17 inducing IL-23 expression in MΦ

We next sought to identify specific genes targeted by neuronal interaction. Therefore,

we performed RNA sequencing in viable FACS sorted mucosal MΦ

(CD45[+]HLA[+]CD64[+]) from fiber-low and fiber-high tissues. As control, we included

MMΦ as well as blood-derived M1 and M2 MΦ differentiated *in vitro*. Principal

component analysis (PCA) showed clear transcriptional separation of blood-derived

M1 and M2 MΦ, whereas colonic LPMΦ and MMΦ accumulated together (Fig. 5a).

Macrophage-specific genes (CD64, CD14, ITGAM, CX3CR1, CCR2, ITGAX, and

CSF1R) were similarly expressed between colonic populations while MMΦ

expressed muscle-specific Lyve1 (Fig. 5b)[55]. When testing for gut-specific M2

markers (AhR, CD200R1, AREG, CD209, TGF-β2, and IL-10), we found elevated

expression in colonic subsets similar to blood-derived M2 MΦ. In contrast to LPMΦ

from fiber-low segments, LPMΦ from fiber-high tissue as well as MMΦ showed

decreased expression of chemokine receptor CCR7 crucial for lymph-node homing,

indicating a tissue-resident population (Fig. 5b). Differential gene expression analysis

yielded 24 significantly changed genes between LPMΦ isolated from fiber-low and

fiber-high colonic tissues (Fig. 5c, d). Strikingly, IL-23a, a subunit of the Th17-

inducing cytokine IL-23, was significantly downregulated in LPMΦ from fiber-high

segments, suggesting a correlation to cholinergic innervation. Additionally, we found

a significant reduction of IL-6 and CCL-20, both of which are involved in Th17

differentiation and recruitment, but no significant change in TNF-α (Fig. 5e).

Using immunofluorescence analysis, we quantitatively analyzed IL-23 protein

expression in CD64$^+$ LPMΦ in cryosections from fiber-low and fiber-high colonic

tissue (Fig. 5f, g). LPMΦ in 4 to 8 regions per patient were analyzed and quantified

using CellProfiler software. Fiber-low colonic tissue showed significantly increased

frequencies of IL-23-positive LPMΦ compared to fiber-high colonic tissue (Fig. 5h).

This result indicated that the lack of mucosal cholinergic fibers correlated with

increased expression of the Th17 inducing cytokine IL-23 in LPMΦ.

Distinct expression of macrophage-neuron crosstalk transcripts

Next, we aimed to identify receptors possibly involved in the neuronal modulation of

LPMΦ. Based on transcriptional analysis, we verified the expression of receptors for

neurotransmitters, neuropeptides, neurotrophic factors, and axon-guidance proteins

(Fig. 6). Considering the crucial role of α7nAChR in mediating the CAIP, we found

elevated levels of the receptor in LPMΦ isolated from fiber-high segments from

HSCR patients (Fig. 6a and f). In contrast, other receptors reported to be involved in

cholinergic immune suppression such as receptors for serotonin[56], catecholamine

(ADRB2)[50,57], or neuropeptides (NMUR1, VIPR1, SSTR2, and CALCRL)[58-65] were

not upregulated in fiber-high LPMΦ compared to fiber-low LPMΦ or were not

expressed (NMUR1) (Fig. 6b, c and f). MΦ secrete different neurotrophic and axon-

guidance proteins as well as respective receptors. Fiber-associated LPMΦ

expressed growth/differentiation factor (GDF) 15 and, to a lesser extent, GDF11

(Fig. 6d and f). Within the family of axon-guidance proteins, we found elevated

expression of the receptors neuropilin-1 (NRP1), UNC-5 homology B (UNC5B), and

ephrin type B-receptor 2 (EPHB2) binding semaphorin 3A (SEMA3A), netrin, and

ephrin-B family members, respectively (Fig. 6e and f). In contrast, LPMΦ isolated

from fiber-low segments showed elevated expression of SEMA3A and SEMA7A

compared to LPMΦ from fiber-high segments (Fig. 6e and f).

Overall, these data suggest a possible involvement of α7nAChR in neuron- MΦ

crosstalk. In addition, fiber-associated MΦ seemed to support neuronal growth and

differentiation via GDF production and axon-guidance proteins. However, data

analysis was limited by low sample numbers and high variance and thus needs

further validation.

Microbial dysbiosis in colonic fiber-low tissues from HSCR patients

Lacking intestinal MΦ or ENS affects the composition of intestinal microbiome[66-69]. In

a chemically induced colitis model in mice, CX3CR1$^+$ MΦ were crucial to maintain

the intestinal barrier function to prevent bacterial translocation[47,70]. To evaluate if the

presence of cholinergic fibers impacts on bacterial translocation, we performed 16S

rDNA FISH analysis of fiber-low and fiber-high cryosections. Bacteria translocated to

lamina propria were detected in both groups, but no significant difference was found

(Supplementary Fig. 4a, b). We then checked the composition of microbial

communities by 16s rDNA sequencing in cryosections of fiber-low, fiber-high, and

ganglionic CD tissue from HSCR patients. Alpha diversity analysis showed an

increase in operational taxonomic units (OTUs) in fiber-low tissue when compared to

ganglionic CD tissue, whereas estimated OTU richness as well as beta diversity

were comparable in all groups (Fig. 7a and Supplementary Fig. 4c). On the phylum

levels, *Actinobacteria* seemed to be less prevalent in aganglionic tissue compared to

ganglionic tissue (Fig. 7b). Within the bacterial family compositions, *Actinobacteria*

were significantly less prevalent in fiber-low segments compared to ganglionic CD,

whereas other families were similarly expressed (Fig. 7c). The main representative

species of the *Actinobacteria* were several *Bifidobacteria* (*B. longum* (OTU1)*; B.*

bifidum (OTU6)*; B. pseudocatenulatum* (OTU5), with *B. longum* showing the highest

frequency in all samples (Supplementary Table 5). Differential bacterial family

analysis between fiber-low and fiber-high colonic tissue revealed a significant

increase in *Staphylococcaceae* in fiber-low colonic segments, whereas in fiber-high

tissue, *Enterobacteriaceae* and *Clostridiales Family_XI* were upregulated (Fig. 7c).

Within the *Staphylococcaceae* family, *S. aureus* (OTU7) was the only detectable

species within the core microbiome (Supplementary Table 5). The *Clostridiales*

Family_XI was represented by 4 genera, i.e., *Finegoldia* (OTU825)*, Peptoniphilus*

(OTU196)*, Anaerococcus* (OTU543) and *Ezakiella* (OTU27, OTU63). The main

representative species of *Enterobacteriaceae* family was *Escherichia coli* (OTU2),

which represented the second frequent bacterial species from the core microbiome

(Supplementary Table 5). These observations suggested a different composition of

colonic translocated bacteria between fiber-low and fiber-high tissue including a

decrease in *Actinobacteria* and prevalence of *Staphylococcaceae* family in fiber-low

segments lacking cholinergic mucosal innervation.

HSCR patients with fiber-low phenotype showed a higher incidence of post-
operative enterocolitis

Next, we evaluated the impact of fiber phenotype on the development of

postoperative enterocolitis. One year after pull-through surgery, 42 HSCR patients

were followed up, and postoperative enterocolitis manifestation was evaluated from

medical records. Enterocolitis was defined to fulfill at least 3 of the following criteria:

diarrhea (>3x/day), vomiting, temperature >38.5°C, antibiotics, hospitalization,

meteorism, and high inflammatory blood parameters (i.e., leucocytes, CRP).

Anthropometric, clinical, environmental, maternal, and dietary factors, collected from

the parent questionnaire, were analyzed for potential enterocolitis risk factors. A total

of 9 (of 42) patients developed enterocolitis one year after surgery (Table 1a).

Development of enterocolitis was significantly associated with cesarean delivery at

birth of the infant (p=0.012). Seven of the 9 patients exhibited a fiber-low and only

two a fiber-high phenotype, showing a higher incidence of fiber-low phenotype

among the children who developed enterocolitis (78% versus 22%, p=0.634). To

validate the observations, we analyzed a retrospective HSCR patient cohort

collected between 2003 and 2018 at the UKBB (Supplementary Table 6). A total of

29 HSCR patients were included. We did not observe any significant patient bias in

terms of anthropometric or clinical factors between fiber-low and fiber-high tissues

(Supplementary Table 7). Enterocolitis development was recorded in 14 of 29

patients but showed no correlation with anthropometric or clinical data (Table 1b).

Strikingly, enterocolitis development correlated significantly with the fiber phenotype

showing a higher incidence in children with the fiber-low phenotype (86%, 12/14

children) compared to the fiber-high phenotype (14%, 2/14). These observations

suggest that HSCR patients with fiber-low phenotype might have a higher risk to

develop enterocolitis and that the fiber phenotype could serve as a prognostic

marker for risk of enterocolitis development.

Discussion

In recent years, multiple studies mainly using different mouse disease models

demonstrated the reciprocal relationship between immune and nervous systems. In

humans however, difficulty to study individual effects of neurotransmitters leads to a

lack of studies providing direct evidence of neuronal-immune crosstalk. Here, we

provided unique evidence of a cholinergic neuron-macrophage crosstalk in the colon

correlated with reduced type 17-mediated immune reactions. By studying colonic

tissue of HSCR patients characterized by the lack of an ENS and extramural

mucosal cholinergic innervation, we deciphered the relationship between neuronal

ACh and the mucosal immune system. Using confocal colocalization studies, we

identified elongated tissue-resident MΦ interacting with cholinergic nerve fibers.

Differently shaped intestinal MΦ populations were described before. Gabanyi et al.

described bipolar-shaped MMΦ with a tissue-protective anti-inflammatory M2

phenotype contrary to stellate-shaped LPMΦ[50]. The elongated profile allows MMΦ to

interact and wrap around neighboring neurons. At steady state, MMΦ regulate ENS

functions and therefore gut motility by providing bone morphogenetic protein (BMP)

2 in response to microbial stimuli[49]. Tamoxifen-induced deletion of CX3CR1[+] tissue-

resident MΦ leads to loss of enteric neurons[71]. Interestingly, MMΦ develop

independently of the ENS[71,72]. The ENS supplies colony-stimulating factor1 (CSF1)

needed for M2 MΦ differentiation and survival[49,73]. Upon *Salmonella typhimurium*

infection, MMΦ interact with sympathetic adrenergic neurons via β2 adrenergic

receptors to induce a tissue- and neuron-protective gene expression program[50]. In a

model of POI which is characterized by surgery-induced inflammatory response of

MMΦ and impaired motility, VN stimulation or pharmacologic triggering of cholinergic

enteric neurons prevented MΦ activation and reduced POI in a α7nAChR-dependent

manner in both mice and humans[56]. However, anti-inflammatory neuro-immune

modulation seems to involve different neuronal receptors in type 2-mediated

intestinal disease models. For example, in a murine model of food allergy, VN

stimulation ameliorated disease severity independently of α7nAChR[74]. Uptake of

food antigens by anti-inflammatory CX3CR1$^+$ MΦ and subsequent induction of food-

specific Tregs effectively prevent allergic reactions[44]. VN stimulation increases

phagocytic activity in CX3CR1$^+$ MΦ, independently of α7nAChR, but phagocytic

activity remains dependent on α4β2nAChR[75]. Using a murine helminth infection

model, the Artis group demonstrated that the cholinergic neuropeptide neuromedin U

activates ILC2 thus increasing the protective immune response and parasitic

clearance[58,59]. In contrast, signals from sympathetic adrenergic neurons lead to

attenuated ILC2 effector function and prevent chronic pathologic type 2 inflammation

via activation of ILC2-specific β2 adrenergic receptors[76]. In our study, IL-23, a

cytokine promoting the maintenance of Th17 cells[77], is significantly increased in

LPMΦ isolated from fiber-low tissue lacking cholinergic innervation compared to

innervated fiber-high tissue. Consistent with this finding, elevated Th17 cells were

detected in fiber-low colonic tissue compared to fiber-high colonic tissue. Th17 cells

are indispensable for the clearance of extracellular bacteria and fungi and therefore

for protective immunity. Dysregulation of cytokines controlling Th17 cells

predisposes to the development of autoimmune diseases and inflammatory bowel

disease[36,78,79]. Genome-wide studies identified genes, including the gene for IL-23

receptor, involved in Th17 differentiation associated with a susceptibility to

inflammatory bowel disease[78,80-82]. IL-23 is produced by dendritic cells and MΦ in

response to extracellular bacteria and fungi recognized by special innate pattern-

recognition receptors, the toll-like receptor (TLR) 2 and nucleotide oligomerization

domain containing protein 2 (NOD2)[83,84]. Under inflammatory conditions, monocyte-

derived CCR2$^+$CX3CR1$^+$ MΦ infiltrate the gut mucosa and prime Th17 cell

responses[43,47]. At steady state, CX3CR1$^+$ MΦ are indispensable for epithelial

integrity, control of bacterial translocation, and development of oral and commensal

tolerance by secreting anti-inflammatory cytokines and maintaining Treg[44,68].

CX3CR1 MΦ further promote intestinal homeostasis by microbiota-dependent

crosstalk with IL-22-secreting ILC3[85].

In our study, we found higher tissue expression of CX3CR1 than CCR2 in fiber-high

colonic segments of HSCR patients and identified two MΦ populations based on

CD14 expression and shape. CD14$^+$ bipolar MΦ were found in close proximity to

nerve fibers in both muscularis and lamina propria, expressing low levels of CCR7,

suggesting a tissue-resident phenotype[86]. CD14^{low} stellate-shaped MΦ were present

in non-innervated lamina propria tissue, expressing CCR7 indicating a migratory

population. Based on their transcriptional profile, LPMΦ isolated from both fiber-low

and fiber-high colonic tissues exhibit a gut-imprinted M2 phenotype (CX3CR1, AREG,

AhR, CSF1R, CD200R, CD209, TGFβ2, and IL-10), but they differed in Th17

cytokines (IL-23 and IL-6) and Th17 chemokines (CCL20). Consequently, cholinergic

signals seemed to be associated with reduced type 17 immune responses. In a

murine model of psoriasis, sensory nociceptive neurons augment dermal dendritic

cell-mediated IL-23 production[64]. The neurons directly sense disease-inducing

Candida albicans and release the neuropeptide calcitonin gene-related peptide

(CGRP) which increases IL-23 production[65]. Surprisingly, another study uncovered

an immunosuppressive effect of nociceptive fibers during *S. aureus* infection

indicating a role for the properties of the underlying pathogen or target immune cell

on nerve fiber-induced immune modulation[87].

Neurotrophic factors and their respective receptors are crucial for ENS development

and function. In fact, the underlying genetic defects in HSCR patients involve

signaling pathways controlling migration and development of the ENS, including the

two major HSCR genes, rearranged during transfection (Ret) and endothelin

receptor type B (EndrB)[88,89]. In a study using Ret-deficient mice, the authors

demonstrated that microbiota-induced glial cell-derived neurotrophic factors induce

IL-22 in Ret-expressing ILC3 therefore maintaining intestinal homeostasis[90]. Ret

expression was not detected in colonic MΦ populations, and we did not detect any

changes in ILC3 frequencies between fiber-low and fiber-high colonic segments.

Here, we used RNA-seq transcriptional analysis to identify factors and receptors

involved in neuro-immune interactions. However, the RNA-seq data were highly

variable within the groups and were not validated in qPCR analyses or on protein

levels. Further studies are needed to fully uncover receptors and mechanisms

involved in neuron-MΦ interactions. Besides the contribution of cholinergic receptors,

we supposed the contribution of neurotrophic factors, axon-guidance proteins, and

their respective receptors in neuron-MΦ interactions. GDF15 has been reported to

suppress inflammatory responses in a murine model of endotoxin-induced sepsis

and prevent maturation of inflammatory dendritic cells in a heart-transplantation

model[91,92]. Using an imiquimod- or IL-23-induced mouse model of psoriasis-like skin

inflammation, GDF11 administration attenuated the severity of skin inflammation via

suppression of the NF-κB signaling pathway[93]. Recently, NRP-1 was identified in

adipose-tissue MΦ protecting from metabolic syndrome by favoring fatty acid uptake

and β-oxidation away from pro-inflammatory M1 polarization and glycolysis[94].

Additionally, in a model of a neurodevelopmental disorder, brown adipose-tissue MΦ

control sympathetic tissue innervation critical for thermogenesis via the plexin-

semaphorin pathway[52].

Inflammation in chemically induced colitis is suppressed by administering

recombinant netrin-1[95] whereas UNC5B deficiency exacerbates chemically induced

colitis[96]. SEMA3A regulates neurite growth cone repulsion suggesting a role of

LPMΦ from fiber-low tissues in the inhibition of mucosal innervation[97,98]. Besides its

crucial role in neurogenesis, SEMA7A supports the generation of monocyte-derived

inflammatory cytokines and differentiation of inflammatory T cells[99,100]. More studies

are needed to evaluate the involvement of the different receptors in our study.

The microbial composition of feces from HSCR patients with enterocolitis is shifted

towards opportunistic pathogens such as *Candida albicans* und *Clostridium*

difficile[1,10,101,102]. In our study, none of the patients was diagnosed with enterocolitis

at the time of the pull-through surgery. However, higher baseline frequencies of Th17

cells and IL-23 might promote *C. difficile* infection, as was recently postulated[103].

Microbial prevalence and composition were altered in aganglionic tissue from fiber-

low and fiber-high phenotypes compared to ganglionic segments. *Actinobacteria*

were significantly reduced in fiber-low segments compared to ganglionic CD tissue.

Compared to fiber-high colonic tissue, fiber-low tissue is associated with higher

levels of *Staphylococcus aureus* previously reported to be associated with an

increased Th17 response[104]. In contrast, *Clostridiales Family_XI* was upregulated

more than 2 log2 fold in fiber-high tissue compared to fiber-low tissue, which was

associated with a decreased risk of *Clostridium difficile* infection and necrotizing

enterocolitis in preterm children[105,106]. In our study, differential bacterial analysis was

limited due to the small patient samples per group and high variability within the

groups. The fact that we analyzed bacterial communities in colonic tissue and rather

than feces further limited the data analysis.

Overall, 9 of 42 patients developed HSCR-associated enterocolitis one year after

pull-through-surgery. Interestingly, development of enterocolitis was significantly

associated with cesarean delivery at birth of the infant in the prospective HSCR

patient cohort. Microbial composition in children delivered by cesarean section is

known to be associated with skin commensals such as *Staphylococcus*, while the

microbiome in children delivered by natural birth mode is characterized by vaginal

bacteria, *e.g., Bifidobacteria* and *Lactobacillus*[107-110]. However, since cesarean

section was not a significant risk factor in the retrospective patient cohort, its role in

enterocolitis development needs to be further examined. Analyzing the correlation of

fiber phenotype with enterocolitis development one year after pull-through surgery

showed a higher incidence for fiber-low (78%) phenotype than fiber-high (22%)

phenotype in patients developing enterocolitis. The result was validated in a

retrospectively analyzed HSCR patient cohort and showed a significant incidence of

86% (fiber-low phenotype) versus 14% (fiber-high phenotype) to develop

enterocolitis. In the retrospective cohort, fiber-low and fiber-high phenotypes were

equally distributed (15 versus 14). Almost 50% of the study population developed

enterocolitis compared to 21% in the prospective cohort.

HSCR represents a rare disease, and low patient numbers limit data analysis.

However, we anticipate validating our results in additional retrospective HSCR

patient cohorts. Interestingly, in HSCR mouse models using EndrB-deficient mice,

no mucosal nerve fibers were detected, and the mice exhibited microbial dysbiosis

leading to enterocolitis[111-113]. The absence of cholinergic fibers could therefore serve

as a predictive marker for enterocolitis and improve personalized patient

management. With the current post-operative therapeutic management, clinicians

can react to inflammatory bowel situations only at the time point of enterocolitis onset.

Using the fiber phenotype as a prognostic marker for the risk of enterocolitis,

clinicians could directly react after pull-through surgery to prevent postoperative

enterocolitis. Possible therapies include high-volume enemas, antibiotic interval

treatment, pre- and probiotic therapy, as well as dietary adaptations. Additionally,

interventional surgical corrections may be considered. For instance, it is still unclear

how the neuronal phenotype of a resected colon segment might predispose for

enterocolitis development. It has been reported that colonic inflammatory Th17 cells

migrate to neighboring mesenteric lymph nodes, triggering enterocolitis development.

Because the mesentery is generally not removed during surgery, this could explain

why patients may develop enterocolitis after removal of affected aganglionic colon

segments. Using the fiber phenotype as a prognostic enterocolitis marker would

enable possible resection of the neighboring mesentery lymph nodes/tissue as a

strategy to prevent enterocolitis. Further, immunotherapies targeting IL-17 or IL-23

rather than TNF-α could open new therapeutic treatment options.

Using a HSCR patient cohort lacking the ENS, we provided evidence that extrinsic

mucosal cholinergic neurons interacted with MΦ and correlated with reduced type-17

immune response most likely mediated by reduced MΦ-dependent IL-23 expression.

HSCR patients lacking mucosal extrinsic cholinergic neurons had a higher risk of

developing postoperative enterocolitis. The mucosal fiber phenotype could serve as

a new prognostic marker for HSCR-associated enterocolitis risk and may offer a

personalized patient care and future therapeutic options.

References

References

1   Demehri, F. R., Halaweish, I. F., Coran, A. G. & Teitelbaum, D. H.

Hirschsprung-associated enterocolitis: pathogenesis, treatment and

prevention. *Pediatr Surg Int* **29**, 873-881, doi:10.1007/s00383-013-3353-1

(2013).

2   Austin, K. M. The pathogenesis of Hirschsprung's disease-associated

enterocolitis. *Semin Pediatr Surg* **21**, 319-327,

doi:10.1053/j.sempedsurg.2012.07.006 (2012).

3   Wang, J. S., Lee, H. C., Huang, F. Y., Chang, P. Y. & Sheu, J. C. Unexpected

mortality in pediatric patients with postoperative Hirschsprung's disease.

Pediatr Surg Int **20**, 525-528, doi:10.1007/s00383-004-1213-8 (2004).

4   Aslam, A., Spicer, R. D. & Corfield, A. P. Turnover of radioactive mucin

precursors in the colon of patients with Hirschsprung's disease correlates with

the development of enterocolitis. *J Pediatr Surg* **33**, 103-105 (1998).

5   Mattar, A. F., Coran, A. G. & Teitelbaum, D. H. MUC-2 mucin production in

Hirschsprung's disease: possible association with enterocolitis development. *J*

Pediatr Surg **38**, 417-421; discussion 417-421, doi:10.1053/jpsu.2003.50071

(2003).

6   Soeda, J., O'Briain, D. S. & Puri, P. Regional reduction in intestinal

neuroendocrine cell populations in enterocolitis complicating Hirschsprung's

disease. *J Pediatr Surg* **28**, 1063-1068 (1993).

7   Imamura, A., Puri, P., O'Briain, D. S. & Reen, D. J. Mucosal immune defence

mechanisms in enterocolitis complicating Hirschsprung's disease. *Gut* **33**,

801-806 (1992).

8   Turnock, R. R., Spitz, L. & Strobel, S. A study of mucosal gut immunity in

infants who develop Hirschsprung's-associated enterocolitis. *J Pediatr Surg*

27, 828-829 (1992).

9   O'Donnell, A. M., Nakamura, H., Tomuschat, C., Marayati, N. F. & Puri, P.

Altered expression of KCNG3 and KCNG4 in Hirschsprung's disease. *Pediatr*

Surg Int **35**, 193-197, doi:10.1007/s00383-018-4394-2 (2019).

10   Hardy, S. P., Bayston, R. & Spitz, L. Prolonged carriage of Clostridium difficile

in Hirschsprung's disease. *Arch Dis Child* **69**, 221-224,

doi:10.1136/adc.69.2.221 (1993).

11   Shen, D. H. *et al.* Detection of intestinal bifidobacteria and lactobacilli in

patients with Hirschsprung's disease associated enterocolitis. *World J Pediatr*

5, 201-205, doi:10.1007/s12519-009-0038-x (2009).

12   Thomas, D. F., Fernie, D. S., Malone, M., Bayston, R. & Spitz, L. Association

between Clostridium difficile and enterocolitis in Hirschsprung's disease.

Lancet **1**, 78-79 (1982).

13   Ward, N. L., Pieretti, A., Dowd, S. E., Cox, S. B. & Goldstein, A. M. Intestinal

aganglionosis is associated with early and sustained disruption of the colonic

microbiome. *Neurogastroenterol Motil* **24**, 874-e400, doi:10.1111/j.1365-

2982.2012.01937.x (2012).

14   Medrano, G. *et al.* B-lymphocyte-intrinsic and -extrinsic defects in secretory

immunoglobulin A production in the neural crest-conditional deletion of

609  endothelin receptor B model of Hirschsprung-associated enterocolitis. *FASEB*

610  *J*, fj201801913R, doi:10.1096/fj.201801913R (2019).

611 15 Cheng, Z. *et al.* Bacterial Microbiome Dynamics in Post Pull-Through

612  Hirschsprung-Associated Enterocolitis (HAEC): An Experimental Study

613  Employing the Endothelin Receptor B-Null Mouse Model. *Front Surg* **5**, 30,

614  doi:10.3389/fsurg.2018.00030 (2018).

615 16 Bruder, E., Terracciano, L. M., Passarge, E. & Meier-Ruge, W. A. [Enzyme

616  histochemistry of classical and ultrashort Hirschsprung's disease]. *Pathologe*

617  **28**, 105-112, doi:10.1007/s00292-007-0901-2 (2007).

618 17 Stakenborg N., D. G. M., Boeckxstaens G.E., Matteoli G. The versatile role of

619  the vagus nerve in the gastrointestinal tract. *EMJ Gastroenterol.*, 106-114

620  (2013).

621 18 Borovikova, L. V. *et al.* Vagus nerve stimulation attenuates the systemic

622  inflammatory response to endotoxin. *Nature* **405**, 458-462,

623  doi:10.1038/35013070 (2000).

624 19 Rosas-Ballina, M. *et al.* Acetylcholine-synthesizing T cells relay neural signals

625  in a vagus nerve circuit. *Science* **334**, 98-101, doi:10.1126/science.1209985

626  (2011).

627 20 Ghia, J. E., Blennerhassett, P., Kumar-Ondiveeran, H., Verdu, E. F. & Collins,

628  S. M. The vagus nerve: a tonic inhibitory influence associated with

629  inflammatory bowel disease in a murine model. *Gastroenterology* **131**, 1122-

630  1130, doi:10.1053/j.gastro.2006.08.016 (2006).

21 Bregeon, J. *et al.* Sacral nerve stimulation enhances early intestinal mucosal

repair following mucosal injury in a pig model. *J Physiol* **594**, 4309-4323,

doi:10.1113/JP271783 (2016).

22 Sun, P. *et al.* Involvement of MAPK/NF-kappaB signaling in the activation of

the cholinergic anti-inflammatory pathway in experimental colitis by chronic

vagus nerve stimulation. *PLoS One* **8**, e69424,

doi:10.1371/journal.pone.0069424 (2013).

23 Gomez-Pinilla, P. J. *et al.* Prucalopride Activates the Intestinal Cholinergic

Anti-Inflammatory Pathway and Prevents Postoperative Ileus.

Gastroenterology **146**, S89-S89 (2014).

24 Di Giovangiulio, M. *et al.* Vagotomy affects the development of oral tolerance

and increases susceptibility to develop colitis independently of the alpha-7

nicotinic receptor. *Mol Med* **22**, 464-476, doi:10.2119/molmed.2016.00062

(2016).

25 Matteoli, G. *et al.* A distinct vagal anti-inflammatory pathway modulates

intestinal muscularis resident macrophages independent of the spleen. *Gut* **63**,

938-948, doi:10.1136/gutjnl-2013-304676 (2014).

26 Bastida, G. & Beltran, B. Ulcerative colitis in smokers, non-smokers and ex-

smokers. *World J Gastroenterol* **17**, 2740-2747, doi:10.3748/wjg.v17.i22.2740

(2011).

27 Bonaz, B. *et al.* Chronic vagus nerve stimulation in Crohn's disease: a 6-

month follow-up pilot study. *Neurogastroenterol Motil* **28**, 948-953,

doi:10.1111/nmo.12792 (2016).

28   Cailotto, C. *et al.* Neuro-anatomical evidence indicating indirect modulation of

macrophages by vagal efferents in the intestine but not in the spleen. *PLoS*

One **9**, e87785, doi:10.1371/journal.pone.0087785 (2014).

29   Gautron, L. *et al.* Neuronal and nonneuronal cholinergic structures in the

mouse gastrointestinal tract and spleen. *J Comp Neurol* **521**, 3741-3767,

doi:10.1002/cne.23376 (2013).

30   Goverse, G., Stakenborg, M. & Matteoli, G. The intestinal cholinergic anti-

inflammatory pathway. *J Physiol* **594**, 5771-5780, doi:10.1113/JP271537

(2016).

31   Kimura, H. *et al.* Neural anti-inflammatory action mediated by two types of

acetylcholine receptors in the small intestine. *Sci Rep-Uk* **9**, doi:ARTN 5887

10.1038/s41598-019-41698-w (2019).

32   Szylberg, L. & Marszalek, A. Diagnosis of Hirschsprung's disease with

particular emphasis on histopathology. A systematic review of current

literature. *Prz Gastroenterol* **9**, 264-269, doi:10.5114/pg.2014.46160 (2014).

33   Wetherill, C. & Sutcliffe, J. Hirschsprung disease and anorectal malformation.

Early Hum Dev **90**, 927-932, doi:10.1016/j.earlhumdev.2014.09.016 (2014).

34   Yoo, B. B. & Mazmanian, S. K. The Enteric Network: Interactions between the

Immune and Nervous Systems of the Gut. *Immunity* **46**, 910-926,

doi:10.1016/j.immuni.2017.05.011 (2017).

35   Revu, S. *et al.* IL-23 and IL-1beta Drive Human Th17 Cell Differentiation and

Metabolic Reprogramming in Absence of CD28 Costimulation. *Cell Rep* **22**,

2642-2653, doi:10.1016/j.celrep.2018.02.044 (2018).

36   Burkett, P. R., Meyer zu Horste, G. & Kuchroo, V. K. Pouring fuel on the fire:

Th17 cells, the environment, and autoimmunity. *J Clin Invest* **125**, 2211-2219,

doi:10.1172/JCI78085 (2015).

37   Awasthi, A. *et al.* Cutting edge: IL-23 receptor gfp reporter mice reveal distinct

populations of IL-17-producing cells. *J Immunol* **182**, 5904-5908,

doi:10.4049/jimmunol.0900732 (2009).

38   Zhou, L. *et al.* IL-6 programs T(H)-17 cell differentiation by promoting

sequential engagement of the IL-21 and IL-23 pathways. *Nat Immunol* **8**, 967-

974, doi:10.1038/ni1488 (2007).

39   Korn, T., Bettelli, E., Oukka, M. & Kuchroo, V. K. IL-17 and Th17 Cells. *Annu

Rev Immunol* **27**, 485-517, doi:10.1146/annurev.immunol.021908.132710

(2009).

40   Tosiek, M. J., Fiette, L., El Daker, S., Eberl, G. & Freitas, A. A. IL-15-

dependent balance between Foxp3 and RORgammat expression impacts

inflammatory bowel disease. *Nat Commun* **7**, 10888,

doi:10.1038/ncomms10888 (2016).

41   Lee, S. H., Starkey, P. M. & Gordon, S. Quantitative analysis of total

macrophage content in adult mouse tissues. Immunochemical studies with

monoclonal antibody F4/80. *J Exp Med* **161**, 475-489 (1985).

42   Kuhl, A. A., Erben, U., Kredel, L. I. & Siegmund, B. Diversity of Intestinal

Macrophages in Inflammatory Bowel Diseases. *Front Immunol* **6**, 613,

doi:10.3389/fimmu.2015.00613 (2015).

43　Panea, C. *et al.* Intestinal Monocyte-Derived Macrophages Control

Commensal-Specific Th17 Responses. *Cell Rep* **12**, 1314-1324,

doi:10.1016/j.celrep.2015.07.040 (2015).

44　Hadis, U. *et al.* Intestinal tolerance requires gut homing and expansion of

FoxP3+ regulatory T cells in the lamina propria. *Immunity* **34**, 237-246,

doi:10.1016/j.immuni.2011.01.016 (2011).

45　Scott, C. L. *et al.* CCR2(+)CD103(-) intestinal dendritic cells develop from DC-

committed precursors and induce interleukin-17 production by T cells.

Mucosal Immunol **8**, 327-339, doi:10.1038/mi.2014.70 (2015).

46　Regoli, M., Bertelli, E., Gulisano, M. & Nicoletti, C. The Multifaceted

Personality of Intestinal CX3CR1(+) Macrophages. *Trends Immunol* **38**, 879-

887, doi:10.1016/j.it.2017.07.009 (2017).

47　Medina-Contreras, O. *et al.* CX3CR1 regulates intestinal macrophage

homeostasis, bacterial translocation, and colitogenic Th17 responses in mice.

J Clin Invest **121**, 4787-4795, doi:10.1172/JCI59150 (2011).

48　Dionne, S., Duchatelier, C. F. & Seidman, E. G. The influence of vitamin D on

M1 and M2 macrophages in patients with Crohn's disease. *Innate Immun-*

London **23**, 557-565, doi:10.1177/1753425917721965 (2017).

49　Muller, P. A. *et al.* Crosstalk between Muscularis Macrophages and Enteric

Neurons Regulates Gastrointestinal Motility. *Cell* **158**, 1210,

doi:10.1016/j.cell.2014.08.002 (2014).

50　Gabanyi, I. *et al.* Neuro-immune Interactions Drive Tissue Programming in

Intestinal Macrophages. *Cell* **164**, 378-391, doi:10.1016/j.cell.2015.12.023

(2016).

51   De Schepper, S., Stakenborg, N., Matteoli, G., Verheijden, S. & Boeckxstaens,

G. E. Muscularis macrophages: Key players in intestinal homeostasis and

disease. *Cell Immunol* **330**, 142-150, doi:10.1016/j.cellimm.2017.12.009

(2018).

52   Wolf, Y. *et al.* Brown-adipose-tissue macrophages control tissue innervation

and homeostatic energy expenditure. *Nat Immunol* **18**, 665-674,

doi:10.1038/ni.3746 (2017).

53   Galitovskiy, V. *et al.* Cytokine-induced alterations of alpha7 nicotinic receptor

in colonic CD4 T cells mediate dichotomous response to nicotine in murine

models of Th1/Th17- versus Th2-mediated colitis. *J Immunol* **187**, 2677-2687,

doi:10.4049/jimmunol.1002711 (2011).

54   Levite, M. Neurotransmitters activate T-cells and elicit crucial functions via

neurotransmitter receptors. *Curr Opin Pharmacol* **8**, 460-471,

doi:10.1016/j.coph.2008.05.001 (2008).

55   Lim, H. Y. *et al.* Hyaluronan Receptor LYVE-1-Expressing Macrophages

Maintain Arterial Tone through Hyaluronan-Mediated Regulation of Smooth

Muscle Cell Collagen. *Immunity* **49**, 1191, doi:10.1016/j.immuni.2018.12.009

(2018).

56   Stakenborg, N. *et al.* Preoperative administration of the 5-HT4 receptor

agonist prucalopride reduces intestinal inflammation and shortens

postoperative ileus via cholinergic enteric neurons. *Gut*, doi:10.1136/gutjnl-

2018-317263 (2018).

57   Noh, H. *et al.* Beta 2-adrenergic receptor agonists are novel regulators of

macrophage activation in diabetic renal and cardiovascular complications.

Kidney Int **92**, 101-113, doi:10.1016/j.kint.2017.02.013 (2017).

58   Cardoso, V. *et al.* Neuronal regulation of type 2 innate lymphoid cells via

neuromedin U. *Nature* **549**, 277-281, doi:10.1038/nature23469 (2017).

59   Klose, C. S. N. *et al.* The neuropeptide neuromedin U stimulates innate

lymphoid cells and type 2 inflammation. *Nature* **549**, 282-286,

doi:10.1038/nature23676 (2017).

60   Delgado, M. & Ganea, D. Vasoactive intestinal peptide: a neuropeptide with

pleiotropic immune functions. *Amino Acids* **45**, 25-39, doi:10.1007/s00726-

011-1184-8 (2013).

61   Chiu, I. M. *et al.* Bacteria activate sensory neurons that modulate pain and

inflammation. *Nature* **501**, 52-57, doi:10.1038/nature12479 (2013).

62   Baliu-Pique, M., Jusek, G. & Holzmann, B. Neuroimmunological

communication via CGRP promotes the development of a regulatory

phenotype in TLR4-stimulated macrophages. *Eur J Immunol* **44**, 3708-3716,

doi:10.1002/eji.201444553 (2014).

63   Assas, B. M., Pennock, J. I. & Miyan, J. A. Calcitonin gene-related peptide is

a key neurotransmitter in the neuro-immune axis. *Front Neurosci* **8**, 23,

doi:10.3389/fnins.2014.00023 (2014).

64   Riol-Blanco, L. *et al.* Nociceptive sensory neurons drive interleukin-23-

mediated psoriasiform skin inflammation. *Nature* **510**, 157-161,

doi:10.1038/nature13199 (2014).

65   Kashem, S. W. *et al.* Nociceptive Sensory Fibers Drive Interleukin-23

769   Production from CD301b+ Dermal Dendritic Cells and Drive Protective

770   Cutaneous Immunity. *Immunity* **43**, 515-526,

771   doi:10.1016/j.immuni.2015.08.016 (2015).

772   66   Earley, A. M., Graves, C. L. & Shiau, C. E. Critical Role for a Subset of

773   Intestinal Macrophages in Shaping Gut Microbiota in Adult Zebrafish. *Cell Rep*

774   **25**, 424-436, doi:10.1016/j.celrep.2018.09.025 (2018).

775   67   Rolig, A. S. *et al.* The enteric nervous system promotes intestinal health by

776   constraining microbiota composition. *PLoS Biol* **15**, e2000689,

777   doi:10.1371/journal.pbio.2000689 (2017).

778   68   Schneider, K. M. *et al.* Cx3cr1 Is a Gatekeeper for Intestinal Barrier Integrity:

779   Limiting Steatohepatitis by Promoting Intestinal Homeostasis. *Journal of*

780   *Hepatology* **62**, S225-S225, doi:Doi 10.1016/S0168-8278(15)30084-2 (2015).

781   69   Johnson, C. D. *et al.* Deletion of choline acetyltransferase in enteric neurons

782   results in postnatal intestinal dysmotility and dysbiosis. *FASEB J* **32**, 4744-

783   4752, doi:10.1096/fj.201701474RR (2018).

784   70   Niess, J. H. *et al.* CX3CR1-mediated dendritic cell access to the intestinal

785   lumen and bacterial clearance. *Science* **307**, 254-258,

786   doi:10.1126/science.1102901 (2005).

787   71   De Schepper, S. *et al.* Self-Maintaining Gut Macrophages Are Essential for

788   Intestinal Homeostasis. *Cell* **175**, 400-415 e413,

789   doi:10.1016/j.cell.2018.07.048 (2018).

72   Avetisyan, M. *et al.* Muscularis macrophage development in the absence of

an enteric nervous system. *P Natl Acad Sci USA*, 4696-4701,

doi:10.1073/pnas.1802490115 (2018).

73   Bogunovic, M. *et al.* Origin of the Lamina Propria Dendritic Cell Network.

Immunity **31**, 513-525, doi:10.1016/j.immuni.2009.08.010 (2009).

74   Bosmans, G. *et al.* Vagus nerve stimulation dampens intestinal inflammation

in a murine model of experimental food allergy. *Allergy*, doi:10.1111/all.13790

(2019).

75   van der Zanden, E. P. *et al.* Vagus nerve activity augments intestinal

macrophage phagocytosis via nicotinic acetylcholine receptor alpha4beta2.

Gastroenterology **137**, 1029-1039, 1039 e1021-1024,

doi:10.1053/j.gastro.2009.04.057 (2009).

76   Klose, C. S. & Artis, D. Neuronal regulation of innate lymphoid cells. *Curr*

Opin Immunol **56**, 94-99, doi:10.1016/j.coi.2018.11.002 (2019).

77   Stritesky, G. L., Yeh, N. & Kaplan, M. H. IL-23 promotes maintenance but not

commitment to the Th17 lineage. *J Immunol* **181**, 5948-5955,

doi:10.4049/jimmunol.181.9.5948 (2008).

78   Barrett, J. C. *et al.* Genome-wide association defines more than 30 distinct

susceptibility loci for Crohn's disease. *Nature Genetics* **40**, 955-962,

doi:10.1038/ng.175 (2008).

79   Galvez, J. Role of Th17 Cells in the Pathogenesis of Human IBD. *ISRN*

Inflamm **2014**, 928461, doi:10.1155/2014/928461 (2014).

80   Franke, A. *et al.* Replication of signals from recent studies of Crohn's disease

identifies previously unknown disease loci for ulcerative colitis. *Nature*

Genetics **40**, 713-715, doi:10.1038/ng.148 (2008).

81   Duerr, R. H. *et al.* A genome-wide association study identifies IL23R as an

inflammatory bowel disease gene. *Science* **314**, 1461-1463,

doi:10.1126/science.1135245 (2006).

82   Gazouli, M. *et al.* NOD2/CARD15, ATG16L1 and IL23R gene polymorphisms

and childhood-onset of Crohn's disease. *World J Gastroentero* **16**, 1753-1758,

doi:10.3748/wjg.v16.i14.1753 (2010).

83   Lyakh, L., Trinchieri, G., Provezza, L., Carra, G. & Gerosa, F. Regulation of

interleukin-12/interleukin-23 production and the T-helper 17 response in

humans. *Immunol Rev* **226**, 112-131, doi:DOI 10.1111/j.1600-

065X.2008.00700.x (2008).

84   Gerosa, F. *et al.* Differential regulation of interleukin 12 and interleukin 23

production in human dendritic cells. *Journal of Experimental Medicine* **205**,

1447-1461, doi:10.1084/jem.20071450 (2008).

85   Mortha, A. *et al.* Microbiota-Dependent Crosstalk Between Macrophages and

ILC3 Promotes Intestinal Homeostasis. *Science* **343**, 1477-+, doi:ARTN

1249288

10.1126/science.1249288 (2014).

86   Schulz, O. *et al.* Intestinal CD103(+), but not CX3CR1(+), antigen sampling

cells migrate in lymph and serve classical dendritic cell functions. *Journal of*

Experimental Medicine **206**, 3101-3114, doi:10.1084/jem.20091925 (2009).

87   Chiu, I. M. *et al.* Bacteria activate sensory neurons that modulate pain and

inflammation. *Nature* **501**, 52-+, doi:10.1038/nature12479 (2013).

88   Heuckeroth, R. O. Hirschsprung disease - integrating basic science and

clinical medicine to improve outcomes. *Nat Rev Gastroenterol Hepatol* **15**,

152-167, doi:10.1038/nrgastro.2017.149 (2018).

89   Tilghman, J. M. *et al.* Molecular Genetic Anatomy and Risk Profile of

Hirschsprung's Disease. *N Engl J Med* **380**, 1421-1432,

doi:10.1056/NEJMoa1706594 (2019).

90   Ibiza, S. *et al.* Glial-cell-derived neuroregulators control type 3 innate

lymphoid cells and gut defence. *Nature* **535**, 440-+, doi:10.1038/nature18644

(2016).

91   Rozakis-Adcock, M. *et al.* Association of the Shc and Grb2/Sem5 SH2-

containing proteins is implicated in activation of the Ras pathway by tyrosine

kinases. *Nature* **360**, 689-692, doi:10.1038/360689a0 (1992).

92   Zhang, Y. *et al.* GDF15 Regulates Malat-1 Circular RNA and Inactivates

NFkappaB Signaling Leading to Immune Tolerogenic DCs for Preventing

Alloimmune Rejection in Heart Transplantation. *Front Immunol* **9**, 2407,

doi:10.3389/fimmu.2018.02407 (2018).

93   Wang, W. *et al.* GDF11 Antagonizes Psoriasis-like Skin Inflammation via

Suppression of NF-kappaB Signaling Pathway. *Inflammation* **42**, 319-330,

doi:10.1007/s10753-018-0895-3 (2019).

94   Wilson, A. M. *et al.* Neuropilin-1 expression in adipose tissue macrophages

protects against obesity and metabolic syndrome. *Sci Immunol* **3**,

doi:10.1126/sciimmunol.aan4626 (2018).

95   Aherne, C. M. *et al.* Neuronal guidance molecule netrin-1 attenuates

inflammatory cell trafficking during acute experimental colitis. *Gut* **61**, 695-705,

doi:10.1136/gutjnl-2011-300012 (2012).

96   Ranganathan, P., Jayakumar, C., Li, D. Y. & Ramesh, G. UNC5B receptor

deletion exacerbates DSS-induced colitis in mice by increasing epithelial cell

apoptosis. *J Cell Mol Med* **18**, 1290-1299, doi:10.1111/jcmm.12280 (2014).

97   Hu, S. & Zhu, L. Semaphorins and Their Receptors: From Axonal Guidance to

Atherosclerosis. *Front Physiol* **9**, 1236, doi:10.3389/fphys.2018.01236 (2018).

98   Ji, J. D., Park-Min, K. H. & Ivashkiv, L. B. Expression and function of

semaphorin 3A and its receptors in human monocyte-derived macrophages.

Hum Immunol **70**, 211-217, doi:10.1016/j.humimm.2009.01.026 (2009).

99   Suzuki, K. *et al.* Semaphorin 7A initiates T-cell-mediated inflammatory

responses through alpha1beta1 integrin. *Nature* **446**, 680-684,

doi:10.1038/nature05652 (2007).

100   Xie, J. & Wang, H. Semaphorin 7A as a potential immune regulator and

promising therapeutic target in rheumatoid arthritis. *Arthritis Res Ther* **19**, 10,

doi:10.1186/s13075-016-1217-5 (2017).

101   Frykman, P. K. *et al.* Characterization of Bacterial and Fungal Microbiome in

Children with Hirschsprung Disease with and without a History of Enterocolitis:

A Multicenter Study. *Plos One* **10**, doi:ARTN e0124172

10.1371/journal.pone.0124172 (2015).

102   Thomas, D. F. M., Malone, M., Fernie, D. S., Bayston, R. & Spitz, L.

Association between Clostridium-Difficil and Enterocolitis in Hirschsprungs-

Disease. *Lancet* **1**, 78-79 (1982).

103   Saleh, M. M. *et al.* Colitis-Induced Th17 Cells Increase the Risk for Severe

884   Subsequent Clostridium difficile Infection. *Cell Host & Microbe* **25**, 756-+,

885   doi:10.1016/j.chom.2019.03.003 (2019).

886   104   Schirmer, M. *et al.* Linking the Human Gut Microbiome to Inflammatory

887   Cytokine Production Capacity. *Cell* **167**, 1125-1136 e1128,

888   doi:10.1016/j.cell.2016.10.020 (2016).

889   105   Vincent, C. *et al.* Bloom and bust: intestinal microbiota dynamics in response

890   to hospital exposures and Clostridium difficile colonization or infection.

891   *Microbiome* **4**, 12, doi:10.1186/s40168-016-0156-3 (2016).

892   106   Vincent, C. *et al.* Reductions in intestinal Clostridiales precede the

893   development of nosocomial Clostridium difficile infection. *Microbiome* **1**, 18,

894   doi:10.1186/2049-2618-1-18 (2013).

895   107   Stokholm, J. *et al.* Cesarean section changes neonatal gut colonization. *J*

896   *Allergy Clin Immunol* **138**, 881-889 e882, doi:10.1016/j.jaci.2016.01.028

897   (2016).

898   108   Shin, H. *et al.* The first microbial environment of infants born by C-section: the

899   operating room microbes. *Microbiome* **3**, 59, doi:10.1186/s40168-015-0126-1

900   (2015).

901   109   Shi, Y. C. *et al.* Initial meconium microbiome in Chinese neonates delivered

902   naturally or by cesarean section. *Sci Rep* **8**, 3255, doi:10.1038/s41598-018-

903   21657-7 (2018).

904   110   Mueller, N. T. *et al.* Delivery Mode and the Transition of Pioneering Gut-

905   Microbiota Structure, Composition and Predicted Metabolic Function. *Genes*

906   *(Basel)* **8**, doi:10.3390/genes8120364 (2017).

111 Erickson, C. S. *et al.* Sacral neural crest-derived cells enter the aganglionic

colon of Ednrb-/- mice along extrinsic nerve fibers. *J Comp Neurol* **520**, 620-

632, doi:10.1002/cne.22755 (2012).

112 Pierre, J. F. *et al.* Intestinal dysbiosis and bacterial enteroinvasion in a murine

model of Hirschsprung's disease. *J Pediatr Surg* **49**, 1242-1251,

doi:10.1016/j.jpedsurg.2014.01.060 (2014).

113 Druckenbrod, N. R., Powers, P. A., Bartley, C. R., Walker, J. W. & Epstein, M.

L. Targeting of endothelin receptor-B to the neural crest. *Genesis* **46**, 396-400,

doi:10.1002/dvg.20415 (2008).

114 Georgeson, K. E. & Robertson, D. J. Laparoscopic-assisted approaches for

the definitive surgery for Hirschsprung's disease. *Semin Pediatr Surg* **13**, 256-

262 (2004).

115 Chen, Y. *et al.* Transanal endorectal pull-through versus transabdominal

approach for Hirschsprung's disease: a systematic review and meta-analysis.

J Pediatr Surg **48**, 642-651, doi:10.1016/j.jpedsurg.2012.12.036 (2013).

116 Romero, P. *et al.* Outcome of transanal endorectal vs. transabdominal pull-

through in patients with Hirschsprung's disease. *Langenbecks Arch Surg* **396**,

1027-1033, doi:10.1007/s00423-011-0804-9 (2011).

117 Struijs, M. C., Diamond, I. R., de Silva, N. & Wales, P. W. Establishing norms

for intestinal length in children. *J Pediatr Surg* **44**, 933-938,

doi:10.1016/j.jpedsurg.2009.01.031 (2009).

Acknowledgment

We thank all the patients and their parents as well as volunteer blood donors for their

contribution. We also thank T. Lopes, E. Traunecker, and L. Raeli at the Flow

Cytometry Facility of the Department of Biomedicine (Basel) for cell sorting,

A. Loynton-Ferrand, K. Schleicher, and O. Biehlmaier at the Imaging Core Facility of

the Biocenter (Basel) for assistance in confocal microscopy, D. Calabrese at the

Histology Facility of the Department of Biomedicine (Basel) for histological support,

P. Weigold at Microsynth AG (Balgach) for microbiome analysis, and S. Rogers at

MediWrite (Basel) for her excellent manuscript revision.

Funding

This work was supported by the Stay on Track (awarded to S.K.) and the Research

Fund Junior Researcher (awarded to S.K.; DMS2343) from the University of Basel,

Switzerland.

Figure legends

Fig. 1 Distribution of neuronal innervation and fiber scoring of HSCR patients

a, Schematic representation of neuronal innervation in non-HSCR (control group) and HSCR patients. Images (5 μm colonic cryosections) show immunohistochemical tubulin staining of ganglionic CD region (non-HSCR) or AChE staining of aganglionic rectum region (HSCR). CD regions of HSCR patients were ganglionic or aganglionic dependent on the length of aganglionosis (see Supplementary Table 1). Rec (rectum); SC (sigmoid colon); CD (colon descendens). **b-c**, Longitudinal colonic strips were rolled up from the distal to proximal ends and cryopreserved as a 'Swiss roll'. Cholinergic nerve fibers were visualized in 5 μm cryosections by AChE immunohistochemical staining using brightfield microscopy. The presence and density of AChE$^+$ fibers in rectum (1 REC) and sigmoid colon (2 SC) were evaluated by four individuals. **b**, Representative image of a fiber-low colonic sections. **c**, Representative image of a fiber-high colonic section. Close-up view shows epithelial region from the different colonic segments.

Fig. 2 Mucosal cholinergic innervation in distal aganglionic colon of fiber high HSCR patient

a, Differently innervated colonic segments from aganglionic and ganglionic regions of HSCR patients and control patients used for subsequent laboratory analysis. Immunofluorescence (5 μm cryosections) of epithelial/mucosal region using tubulin (Alexa647, green), a general neuronal marker, and AChE (Cy3, red), a cholinergic marker. DAPI (blue) shows cell nuclei. Images were taken by fluorescence microscopy (20x, scale bar 50 μm) and processed by Fiji software. Encircled areas mark the lumen. Arrows mark cholinergic fibers; asterisk mark non-cholinergic fibers.

b, Immunofluorescence (5 μm Swiss roll cryosections) of aganglionic rectum region

from a fiber-high patient using tubulin (Alexa647, green), a general neuronal marker,

combined with VIP (vasoactive intestinal peptide; A555, red), labeling peptidergic

neurons; TH (tyrosine hydroxylase; A555, red), labeling dopaminergic and

adrenergic neurons; NOS (nitric oxide synthase, A555, red), labeling nitrinergic

neurons and S100B (S100 calcium-binding protein B, A555, red), labeling glia cells.

DAPI (blue) shows cell nuclei. Images were taken by fluorescence microscopy (20x,

scale bar 50 μm) and processed by Fiji software. Close-up views (white boxes) show

mucosal/epithelial region. **c-d**, Immunofluorescence (5 μm Swiss roll cryosections) of

aganglionic rectum region from a fiber-high patient using secondary antibody

controls.

Fig. 3 Cytokine profile and immune cell populations in colonic tissue

a, Quantitative RT-PCR analysis of total colonic tissue isolated from different

aganglionic (AG) and ganglionic (G) colonic segments of HSCR and control patients.

Fiber-low (n=31); fiber-high (n=13); AG CD (n=12); G CD (n=11); control CD (n=4).

IL-17 expression was not detectable in 1 AG CD sample and in 1 G CD sample. IL-

22 expression was not detectable in 2 G CD sample. **b-d**, Flow cytometric analysis

of mononuclear cells isolated from different aganglionic and ganglionic colonic

segments of HSCR and control patients. **b**, Frequencies of IL-17$^+$ Th17 T cells were

determined in viable CD3$^+$CD4$^+$ lymphocytes restimulated with PMA/Ionomycin

(Supplementary Fig. 3a). **c**, Treg cells were defined as CD127$^-$Foxp3$^+$ cells in

unstimulated viable CD3$^+$CD4$^+$ lymphocytes using Foxp3/transcription factor staining

buffer set (Supplementary Fig. 3b). Fiber-low (n=30); fiber-high (n=13); AG CD

(n=10); G CD (n=10); control CD (n=5) **d,** Th17/Treg ratio is shown. **e**, Frequencies

of CD11c$^+$ and CD14$^+$ MΦ were evaluated in viable HLA$^+$CD64$^+$ cells by flow

cytometry (Supplementary Fig. 2d). Fiber-low (n=26); Fiber-high (n=13); AG CD

(n=10); muscle (n=10); G CD (n=10); control CD (n=5). **f,** CD64$^+$ MΦ were assessed

by immunofluorescence in 5 µm colonic cryosections from a muscle region (left), a

mucosal region from a fiber-low patient, and a mucosal region from fiber-high patient

(right). Scale bar 30 µm (muscle) and 10 µm (fiber-low; fiber-high). Images from 3

fiber-high and 3 fiber-low patients were analyzed for MΦ shape. A total of 100-200

CD64$^+$ cells per patient were counted and the percentage of bipolar cells calculated

(bar graph). Scatter plots show means ± SEM. Significance was determined using

unpaired nonparametric Mann-Whitney test (* p≤ 0.05; ** p≤ 0.01; *** p≤ 0.001).

Source data are provided as a Source Data file.

Fig. 4 MΦ colocalize with cholinergic nerve fibers

a, Immunofluorescence of epithelial region in cryosections (5 µm, Swiss roll of

rectum and sigmoid colon) of representative fiber-high and fiber-low patients. Anti-

AChE (red); anti-CD64 or anti-CD3 (green); DAPI (blue). Image from muscle region

(right) shows thick fiber bundles (AChE) together with CD64$^+$ cells. Images were

obtained by fluorescence (20x, scale bar 30 µm) and confocal (63x, scale bar 10 µm)

microscopy and processed by Fiji software. **b**, Confocal immunofluorescence image

of an intercrypt region from a fiber-high patient. Overlapping pixels (yellow, arrows)

were visualized on a Z-projection image. See Supplementary movie.

c-d, Quantification of the pixel intensity colocalization of CD64$^+$ MΦ **(c)** or CD3$^+$

T cells **(d)** with AChE$^+$ fibers was performed by a Fiji software macro. Confocal

images from distal colon regions of fiber-low (n=4 for CD64; n=3 for CD3) and fiber-

high (n=5 for CD64; n=3 for CD3) patients as well as from muscle regions of HSCR

patients (n=2) were taken. The following negative staining controls were used:

CD64-A488 **(c)** or CD3-A488 **(d)** with biotin + streptavidin-Cy3 and IgG1-A488 +

biotin/streptavidin-Cy3 (n=2). Per patient (dot), 200-300 cells were analyzed. Scatter

plots show mean raw integrated density (RawIntDen) ± SEM. **e**, CD64$^+$ MΦ from 3

fiber-high patients were investigated with respect to bipolar (n=170) or round/stellate

(n=152) morphological phenotype. Raw integrated density of colocalized pixels

between CD64 and AChE are shown. Significance was determined using unpaired

nonparametric Mann-Whitney test (* p≤ 0.05; **** p≤ 0.0001). Source data are

provided as a Source Data file.

Fig. 5 Fiber-associated MΦ express diminished IL-23 levels

RNA sequencing was performed on sorted viable colonic MΦ (CD45$^+$HLA$^+$CD64$^+$,

Supplementary Fig.3d and Methods) and blood-derived M1 and M2 MΦ.

a, PCA plot shows similarity between the different populations based on their first

two principal components. The top 500 genes, selected by highest row variance,

were used. PC, principal component. Fiber-low (green, n=4); fiber-high (blue, n=5);

muscle (red, n=3); M1 (black, n=3); M2 (grey, n=3). **b**, Normalized sequence reads

of selected macrophage-related genes. Scatter plots show means ± SEM.

Significance was determined using unpaired nonparametric Mann-Whitney test or

unpaired t-test (for M1/M2 MΦ) (* p≤ 0.05; ** p≤ 0.01; *** p≤ 0.001) **c**, Global

transcriptional changes between macrophages isolated from fiber-low colonic tissue

and fiber-high colonic tissue visualized by a volcano plot. Genes with an adjusted

p-value < 0.05 and log2 fold change greater than 1 (red dots) and less than -1 (blue

dots) are indicated. **d**, Heat map of differentially expressed genes (from b) between

MΦ in fiber-low colonic tissue and fiber-high colonic tissue. Each column represents

one patient. **e**, Normalized sequence reads of selected genes. Scatter plots show

means ± SEM. Significance was determined using unpaired nonparametric Mann-

Whitney test or unpaired t-test (for M1/M2 MΦ) (* p≤ 0.05; ** p≤ 0.01; *** p≤ 0.001).

f-g, IL-23$^+$CD64$^+$ MΦ were assessed by immunofluorescence in 5 µm colonic

cryosections. A representative image of mucosal/epithelial region from fiber-low

tissue (**f**) and fiber-high tissue (**g)** is shown. Scale bar 30 µm. CD64 (A647, green);

IL23p19-PE (red) **h**, Quantitative analysis of IL-23 expression in macrophages

evaluated in images from 4 fiber-high and 4 fiber-low patients. A total of 900-4700

CD64$^+$ cells per patient were analyzed for IL-23 expression and the percentage of

IL23$^+$CD64$^+$ cells (% of total macrophages) calculated. Scatter plots show means ±

SEM. Significance was determined using unpaired nonparametric Mann-Whitney test

(* p≤ 0.05). Source data are provided as a Source Data file.

Fig. 6 Receptors involved in macrophage-neuron crosstalk

a-e, Heat maps of selected genes possibly involved in macrophage-neuron crosstalk.

Normalized reads are shown. RNA sequencing was performed on sorted viable

colonic MΦ (CD45$^+$HLA$^+$CD64$^+$) from mucosa of fiber-low (n=4) and fiber-high (n=5)

colonic tissue as well as muscle tissue (n=3). Heat maps show mean values of each

group. **a**, Expression of nicotinic (CHRNA) and muscarinic (CHRM) ACh receptors. **b**,

Expression of serotonergic (HTR) and catecholaminergic (ADRB) receptors.

c, Expression of receptors for neuromedin U (NMUR1), vasoactive-intestinal-peptide

(VIPR1), somatostatin (SSTR2), calcitonin-gene-related peptide (CALCRL/RAMP1).

d, Expression of neurotrophins (bone-morphogenic proteins [BMP] and

growth/differentiation factors [GDF]) and neurotrophin receptors (NTRK2 binds brain-

derived-neurotrophic factor (BDNF) and neurotrophin (NT)-4; GFRA2 and RET bind

GDNF and neurturin. **e**, Expression of axon-guidance proteins and respective

receptors. Semaphorins (SEMA) bind to plexins or neuropilin-1 (NRP1). UNC-5

represent the receptor for netrin. Receptor proteins ROBO1 and 3 bind slit proteins.

Neogenin (Neo) receptor binds to repulsive guidance protein (RGM). Ephrins bind to

Eph receptors EPHA and EPHB. **f**, Normalized sequence reads of selected genes.

Scatter plots show means ± SEM. Significance was determined using unpaired

nonparametric Mann-Whitney test (*p≤ 0.05). Source data are provided as a Source

Data file.

Fig. 7 Colonic microbial dysbiosis in fiber-low colonic tissue from HSCR
patients

a-d, 16S rRNA analysis of colonic tissue from aganglionic (fiber-low n=14; fiber-high

n=11) and ganglionic (CD gang. n=4) segments of age-matched HSCR patients.

a, Alpha diversity analysis show effects on observed operational taxonomic units

(OTUs) and estimated OTU richness by Shannon and Simpson indices. **b**, Pie charts

show microbial composition in each group on phylum level. **c**, Scatter plots show

normalized reads of bacterial classes dedicated to respective phyla. Scatter blots

show mean values ± SEM. Significance was determined using unpaired

nonparametric Mann-Whitney test (* p≤ 0.05). **d**, Differential bacterial family analysis

of fiber-low versus fiber-high groups. Asterisk mark significantly changed families

with an adjusted p value< 0.05 (***p= 4.01×10^{-15}; **p= 2.08×10^{-4}; *p= 0.022).

Families are assigned to respective phyla by color code. Source data are provided

as a Source Data file.

Tab. 1 Risk factors for post-operative enterocolitis

a, Prospective HSCR patient cohort (n=42). Values are median (range) or number

(percentage); (n): number of data available; length of aganglionosis: rectum (1),

sigmoid colon (2), colon descendens (3), colon transversum (4), colon ascendens (5).

Differences between enterocolitis development (yes/no) were tested with the

Wilcoxon rank-sum test and χ2-test, as appropriate. No follow-up for patients no.

0023 and 0032 available. **b**, Retrospective HSCR patient cohort (n=29). Values are

mean (SD), median (range), or number (percentage); n: number; SD: standard

deviation. Differences between enterocolitis development (yes/no) were tested with

the Wilcoxon rank-sum test and χ2-test, as appropriate. Source data are provided as

a Source Data file.

1097

Fig. 1 Distribution of neuronal innervation and fiber scoring of HSCR patients

a

Non-HSCR

HSCR

Fiber low   **Fiber high**

proximal

CD

SC

Rec

distal

Ganglionic

Aganglionic or ganglionic

Aganglionic

IHC: anti-tubulin

IHC: anti-AchE

IHC: anti-AchE

mucosa sub-mucosa muscle

mucosa sub-mucosa muscle

mucosa sub-mucosa muscle

Myenteric plexus   ● Submucous plexus with mucosal terminals   cholinergic extrinsic nerve fibers

Rec: rectum; SC: sigmoid colon; CD: colon descendens

b

IHC: anti-AchE

proximal

distal (anus)

3 2 1

1Rec

2 SC

3 CD

HSCR Fiber low

c

3 2 1

1Rec

2 SC

3 CD

HSCR Fiber high

Fig. 2 Mucosal cholinergic innervation in distal aganglionic colon of fiber high HSCR patient

a

HSCR patients | control patient

Fiber Low | Fiber High | CD AG | CD G | CD control

Tubulin

AchE

merged

Aganglionic | Ganglionic

b

Tubulin | VIP | Merged

Tubulin | TH | Merged

Tubulin | NOS | Merged

Tubulin | S100B | Merged

c

d

guinea pig-A555 + mouse IgG2a-A647

mouse IgG2a-A555 + mouse IgG2b-A647

mouse IgG1-A555 + mouse IgG2a-A647

mouse IgG2b-biotin/ SA-cy3 mouse IgG2a-A647

Fig. 3 Cytokine profile and immune cell populations in colonic tissue

Fig. 4 Macrophages colocalize with cholinergic nerve fibers

Fig. 5 Fiber-associated macrophages express diminished IL-23 levels

Fig. 6 Receptors involved in macrophage-neuron crosstalk

Fig. 7 Colonic microbial dysbiosis in fiber low colonic tissue from HD patients

Table 1: Risk factors for post-operative enterocolitis development

a

Prospective HSCR patient cohort	Enterocolitis		p-value
Anthropometric data and potential risk factors, (n)	Yes n=9	No n=33	
Age at operation, months (42)	3 (1–127)	5 (2–70)	0.211
Sex, male (42)	7 (78)	27 (81)	0.784
Preterm birth (38)	2 (22)	1 (3)	0.068
Operation procedure trans-anal (42)	8 (89)	29 (88)	0.934
Length of aganglionosis (42)	1 (1-5)	2 (1-5)	0.061
Pets (37)	1 (11)	11 (39)	0.116
Patient living in rural area (40)	3 (33)	17 (55)	0.256
Antibiotic treatment (38)	2 (22)	3 (10)	0.357
Siblings (39)	8 (89)	22 (73)	0.331
Breastfed at operation (39)	9 (100)	27 (90)	0.323
Maternal age at delivery, years (35)	27 (23-39)	29 (21-43)	0.788
Parental smoking (38)	4 (44)	5 (17)	0.094
Caesarean Section (38)	6 (75)	8 (27)	0.012
Maternal Allergy (38)	1 (11)	8 (31)	0.236
Fiber low (30)	7 (78)	23 (70)	0.634
Fiber high(12)	2 (22)	10 (30)	0.634

b

	Enterocolitis		p-value
Retrospective HSCR patient cohort	Yes (n=14)	No (n=15)	
Age at operation, months	4 (1–48)	9 (1–171)	0.229
Sex, male	11 (79)	14 (93)	0.249
Preterm birth	1 (7)	1 (7)	0.960
Antibiotic treatment	10 (7)	11 (73)	0.909
Caesarean Section	7 (50)	5 (33)	0.124
Fiber low	12 (86)	3 (20)	<0.001
Fiber high	2 (14)	12 (80)	<0.001

Supplementary Table 1: Overview of prospective HSCR patient cohort

Study Number	Sex	Age at Surgery (months)	OP technique	Site of operation	Diagosis	Lenght of AG	Fiber score	CD aganglionic	CD ganglionic	Postoperative clinical symptoms: Obstipation (1); Incontinence (2); Enterocolitis (3); Pain (4)	Postoperative clinical treatment: Laxative (1); Clyster (2); Botox (3); Antibiotics (4); Colonic irrigation (5); Pre-/probiotics (6)
HSCR patients											
0002	F	6.6	abdominal Swenson	01	HSCR	TCA	low	x		1,2,3,4	1,2,4,5,6
0004	M	6.7	transanal Soave	02	HSCR	S-HSCR	low			no	no
0005	M	127.5	transanal Soave	10	HSCR	S-HSCR	low			2,3,4	1,4,5
0006	F	4.3	transanal Soave	10	HSCR	S-HSCR	low			1	1
0010	M	3.9	transanal Swenson	07	HSCR	S-HSCR	high			no	no
0012	M	4.9	transanal Swenson	09	HSCR	L-HSCR	high	x		1	1,2
0013	M	6.9	transanal Swenson	07	HSCR	S-HSCR	low			no	no
0014	M	3.9	transanal Soave	10	HSCR	S-HSCR	low		x	3	4
0015	F	2.8	transanal Soave	01	HSCR	S-HSCR	low		x	no	no
0016	M	3.2	transanal Soave	10	HSCR	S-HSCR	high			no	4
0017	M	5.9	abdominal Soave	09	HSCR	TCA	low	x		no	no
0018	M	3.2	transanal Soave	10	HSCR	S-HSCR	high		x	no	4
0020	M	11.8	transanal Soave	07	HSCR	L-HSCR	low	x		no	no
0021	M	23.5	transanal Soave	04	HSCR	S-HSCR	low	x		1	2
0022	M	7.2	abdominal Soave	09	HSCR	TCA	low	x		no	4
0023	F	14.2	transanal Soave	04	HSCR	S-HSCR	low		x	no	no
0025	M	3	transanal Soave	10	HSCR	S-HSCR	high	x	x	no	4
0029	M	10.1	transanal Soave	09	HSCR	L-HSCR	high	x	x	2,4	no
0032	M	35.9	transanal Soave	04	HSCR	S-HSCR	high		x	no	no
0034	M	2.9	abdominal Soave	06	HSCR	TCA	low	x		no	no
0038	M	19.2	abdominal Soave	10	HSCR	TCA	low	x		4	no
0039	M	12.5	transanal Soave	09	HSCR	S-HSCR	low			no	no
0040	M	3.8	transanal Soave	02	HSCR	S-HSCR	low			no	no
0041	M	5.9	transanal Soave	03	HSCR	L-HSCR	low	x		1	1
0042	M	2.2	transanal Soave	06	HSCR	S-HSCR	low			1,3	1,3,4,5
0044	M	13.6	transanal Soave	03	HSCR	S-HSCR	low		x	no	no
0047	M	44.5	transanal Soave	02	HSCR	S-HSCR	high			2	no
0048	M	3.8	transanal Soave	07	HSCR	S-HSCR	low			1	3,5
0049	M	70.5	transanal Soave	10	HSCR	S-HSCR	low		x	no	4
0050	M	5.7	transanal Soave	10	HSCR	L-HSCR	low	x		no	no
0051	M	2.5	transanal Soave	08	HSCR	S-HSCR	low			3	no
0053	M	1	transanal Soave	06	HSCR	S-HSCR	low			1,3	1,2,3,4,5
0054	F	6.1	transanal Soave	10	HSCR	S-HSCR	high			no	no
0056	M	4.2	transanal Soave	08	HSCR	S-HSCR	high	x		1	1
0057	F	7.7	transanal Soave	08	HSCR	S-HSCR	high			1,3	1,4,5,6
0058	M	5.7	transanal Soave	08	HSCR	S-HSCR	high	x		3	4,5
0059	M	5	transanal Soave	10	HSCR	S-HSCR	low			no	no
0060	M	5	abdominal Soave	10	HSCR	TCA	high			4	5
0061	M	2.8	transanal Soave	02	HSCR	S-HSCR	low	x		1	1
0063	M	3.8	transanal Soave	08	HSCR	S-HSCR	low		x	3,4	1,4,5
0064	F	64	abdominal Soave	01	HSCR	TCA	low	x		no	4
0065	F	34.5	transanal Soave	09	HSCR	S-HSCR	low			no	no
0066	F	5.3	transanal Soave	01	HSCR	L-HSCR	low	x		no	6
0068	M	3.9	transanal Soave	09	HSCR	S-HSCR	high		x	no	no
Non-HSCR patients											
0024	M	0.4	transabdominal	01	Anal atresia/Ostomy			x	x		
0030	M	7	transabdominal	06	Anal atresia/Ostomy			x	x		
0037	M	5	transabdominal	01	Anal atresia/Ostomy			x	x		
0062	M	5.9	transabdominal	01	Anal atresia/Ostomy			x	x		
0067	F	6.8	transabdominal	01	Colonic perforation/Ostomy			x	x		
0045	M	11.9	transabdominal	10	misdiagnosed HSCR			x	x		

Site of operation: Basel (CH) 01; Bern (CH) 02; Düsseldorf (DE) 03; Freiburg (DE) 04; Heidelberg (DE) 06; Karlsruhe (DE) 07; Lausanne (CH) 08; St. Gallen (CH) 09; Zürich (CH) 10. Length of aganglionose: TCA (Total colonic aganglionose); L-HSCR (long-segmented HSCR); S-HSCR (short segmented HSCR). Female (F); Male (M).

Supplementary Table 2: Enzyme histochemistry performed at the different pathology units of recruiting clinics used for HSCR diagnosis

Pathology unit	Detection of ganglia	Detection of extrinsic nerve fibers
Basel, Freiburg, Karlsruhe	LDH, NOS, SDH	AChE
Zürich	LDH	AChE
Bern	LDH, NOS	AChE
St. Gallen	SDH	AChE
Lausanne	LDH	AChE
Heidelberg	Calretinin	AChE
Düsseldorf	NADPH, LDH	AChE

Lactic dehydrogenase (LDH), succinic dehydrogenase (SDH), nitroxide synthase (NOS) Nicotinamide adenine dinucleotide (NADPH), acetylcholinesterase (AChE)

Supplementary Table 3: Characteristics prospective HSCR patient cohort

Anthropometric data and potential risk factors (n)	All (n=44)	Fiber-low (n=31)	Fiber–high (n=13)	p-value
Age at operation, months (44)	5 (1–127)	5 (1–127)	4 (3–44)	0.678
Sex, male (44)	35 (79)	23 (74)	12 (92)	0.174
Preterm birth (40)	3 (8)	1 (4)	2 (18)	0.114
Pets (41)	13 (33)	9 (33)	4 (33)	1.000
Patient living in rural area (42)	21 (50)	15 (50)	6 (50)	1.000
Antibiotic treatment (40)	5 (13)	4 (15)	1(8)	0.602
Siblings (41)	32 (78)	21 (72)	11 (92)	0.175
Breastfed at operation (41)	38 (93)	27 (90)	11 (100)	0.276
Maternal age at delivery, years (37)	29 (21–43)	30 (23–43)	28 (21–35)	0.111
Parental smoking (40)	9 (23)	5 (18)	4 (33)	0.283
Caesarean Section (40)	14 (35)	7 (25)	7 (58)	0.043
Maternal Allergy (40)	11 (28)	7 (25)	4 (33)	0.685

Values are median (range) or number (percentage); (n): number of data available; Differences between the fiber high/low group of children were tested with the Wilcoxon rank-sum test and χ2-test, as appropriate. Source data are provided as a Source Data file.

Supplementary Figure 1:
Fiber scoring of distal colon of HSCR patients using tubulin and AChE immunohistochemistry

Tubulin and AChE immunohistochemistry of 5µm cryosections of distal aganglionic colon of two fiber high (**a**) and two fiber low (**b**) HSCR patients.

Supplementary Figure 2:
Immune cell populations in colonic tissue from HSCR and control patients and *in vitro* T-cell conversion assay

a, Flow cytometric analysis of mononuclear cells isolated from different aganglionic and ganglionic colonic segments of HSCR and control patients. Frequencies of total cells is shown. IL-17⁺ Th17 T cells were gated in viable CD3⁺CD4⁺ lymphocytes restimulated with PMA/Ionomycin. Treg cells were defined as CD127⁻Foxp3⁺ cells in unstimulated viable CD3⁺CD4⁺ lymphocytes. Fiber-low (n=30); fiber-high (n=13); AG CD (n=10); G CD (n=10); control CD (n=5). CD14⁺ MΦ were gated in viable HLA⁺CD64⁺ cells. Fiber-low (n=26); Fiber-high (n=13); AG CD (n=10); muscle (n=10); G CD (n=10); control CD (n=5). Exact p values indicated in Supplementary Table 4 **b** Frequencies of NCR⁺ (Nkp44⁺) and NCR⁻ (Nkp44⁻) ILC3 (CD117⁺) subsets. Fiber-low (n=17); fiber-high (n=9); AG CD (n=6); G CD (n=6); control CD (n=3). Supplementary Fig. 3c shows flow cytometric gating strategy. Scatter plots show means ± SEM. Significance was determined using unpaired nonparametric Mann-Whitney test (* p≤ 0.05; ** p≤ 0.01; *** p≤ 0.001 and **** p≤ 0.0001); Exact p values indicated in Supplementary Table 4. **c,** Quantitative RT-PCR analysis of total colonic tissue or isolated muscle tissue from different aganglionic and ganglionic colonic segments of HSCR and control patients (CX3CR1;circles and CCR2;squares). Significance was determined using Wilcoxon matched-pairs signed rank test. Fiber-low (n=31); Fiber-high (n=13); AG CD (n=12); G CD (n=11); control CD (n=5). **d-e,** Representative brightfield microscopic images (10x) of M2 **(d)** and M1**(e)** after 10 days of differentiation. Scale bar 50 μm. **f-g**, Flow cytometric analysis of blood-derived M1 and M2 macrophages from adult healthy donors. Histograms show CD14 (**f**) and CX3CR1 (**g**) expression in M1 (green) and M2 (blue) macrophages as well as in unstained controls (filled histograms). Representative analysis out of 3 independent experiments is shown. **h,** Th17 and Treg conversion assay using blood-derived M1 and M2 macrophages. Frequencies of Th17 and Treg cells were determined and Th17/Treg ratio calculated. One out of 3 independent experiments is shown. Significance was determined using unpaired t-test. Source data are provided as a Source Data file.

Supplementary Table 4: Statistical analysis of gene and protein expression

	Fiber low vs. Fiber high	Fiber low vs. CD AG	Fiber low vs. CD G	Fiber low vs. CD control	Fiber high vs. CD AG	Fiber high vs. CD G	Fiber high vs. CD control	CD AG vs. CD G	CD AG vs. CD control	CD G vs. CD control	Muscle vs. CD AG	Muscle vs. CD G	Muscle vs. CD control
FACS Th17	0.0338	0.0002	0.0073	0.0978	0.3434	0.7844	>0.9999	0.4813	0.6787	>0.9999			
FACS Treg	0.6665	0.0158	0.0308	0.1413	0.0257	0.0358	0.1433	0.1433	0.8969	0.953			
FACS ILC3 NCR+	0.4762	0.4356	0.7273	0.8509	0.9003	0.7446	0.6	0.8718	0.6667	0.8333			
FACS ILC3 NCR-	0.2339	0.9956	0.0487	0.7763	0.6698	0.7292	0.4818	0.1638	0.8833	0.1833			
FACS Ratio Th17/Treg	0.0099	0.0817	0.2177	0.157	0.5629	0.2316	0.7028	0.5288	0.953	0.8591	0.1383	0.2711	0.2428
FACS CD11c	>0.9999	0.909	0.3889	0.8962	0.7038	0.1263	0.6877	0.4688	0.7632	0.8358			
FACS CD14	0.0176	0.05196	0.614	0.2171	0.017	0.1514	0.7589	0.3238	0.2544	0.308	0.0279	0.0131	0.6064
qPCR IL-17	0.0129	0.0222	0.0275	0.2372	< 0.0001	0.775	0.4773	0.0011	0.0103	0.7333			
qPCR IL-22	0.0151	0.2887	0.0453	0.4117	0.0114	0.9738	0.8697	0.0336	0.1934	0.8224			
qPCR IL-6	0.1968	0.3354	0.0363	0.4366	0.0788	0.4037	0.078	0.0188	> 0.9999	0.078			
qPCR TNF-α	0.5322	0.6971	0.0759	0.5692	0.4968	0.0863	> 0.9999	0.1584	0.7066	0.122			
qPCR IL-1β	0.7296	0.4609	0.0837	0.6752	0.401	0.4302	0.8882	0.0447	0.2868	0.4462			
qPCR IL-23	0.6935	0.1832	0.2552	0.5235	0.1185	0.7184	0.7227	0.0398	0.0381	> 0.9999			
Luminex IL-17	0.123	0.469	0.522	0.506	0.123	0.564	0.095	0.355	>.999	0.428			
Luminex IL-22	0.291	0.907	0.298	0.426	0.621	0.924	0.095	0.523	0.521	0.153			
Luminex IL-6	0.151	0.65	0.276	0.802	0.087	0.894	0.443	0.101	>.999	0.279			
Luminex TNF-α	0.744	0.082	0.071	0.909	0.06	0.22	0.849	0.006	0.166	0.317			
Luminex IL-1β	0.265	0.117	0.049	>.999	0.04	0.699	0.443	0.014	0.383	0.195			
ELISA IL-23	0.439	0.701	0.374	0.963	0.315	0.101	0.485	0.888	0.909	0.4			
FACS Th17 (% of total)	0.0009	0.3938	0.0765	0.1194	0.0255	0.1264	0.2546	0.2393	0.4396	0.8838			
FACS Treg (% of total)	0.0337	0.7365	0.0278	0.4709	0.0441	0.9153	0.2355	0.013	0.2439	0.2398			
FACS Ratio Th17/Treg (% of total)	0.0313	0.0974	0.954	0.5448	0.8938	0.0638	>0.9999	0.3357	0.7551	0.5237			
FACS CD14 (% of total cells)	0.5076	0.8214	0.925	0.6578	0.8918	0.7378	0.3359	0.955	0.371	0.4222	0.0789	0.0372	0.2398

Statistical significance between sample groups was determined using unpaired nonparametric two-tailed Mann-Whitney test, and differences were considered statistically significant with * p≤ 0.05; ** p≤ 0.01; *** p≤ 0.001 and **** p≤ 0.0001. Grey boxes mark statistically significant values.

Supplementary Figure 3: Gating strategy for flow cytometric analysis and cell sorting

a-e, Representative gating strategies for the FACS analysis of lamina propria mononuclear cells from HSCR and control patients. a, T helper 17 (TH17) cells were evaluated in viable CD3$^+$CD4$^+$ lymphocytes restimulated with PMA (phorbol 12-myristate 13-acetate)/Ionomycin in the presence of GolgiStop^{TM} and Brefeldin A. Unstimulated Lymphocytes (Golgi-stop/Brefeldin A only) served as negative control. **b**, CD127$^-$Foxp3$^+$ regulatory T cells (Tregs) were gated in viable CD3$^+$CD4$^+$ lymphocytes. **c**, Subsets of ILC3 were detected in viable lin$^-$CD127$^+$CD161$^+$CRTH2$^-$ lymphocytes. ILC3 NCR$^+$ were defined as c-kit$^+$NKp46$^+$ and ILC3 NCR$^-$ as c-kit$^+$NKp46$^-$. **d**, Macrophage gating strategy for FACS analysis as well as for cell sorting. HLA$^+$CD64$^+$ macrophages were detected in viable CD45$^+$ leucocytes.

Supplementary Movie 1: Co-localization of macrophages and cholinergic nerve fibers

Colocalization of mucosal macrophages with cholinergic nerve fibers at the epithelial region of an inter-crypt, related to Figure 4b. Cryosection (5 μm) from a rectum part of a fiber-high patient was stained using human anti-CD64 (green), anti-AChE (red), and nuclear stain (DAPI). 3D reconstruction movie from a fast scanning confocal image using Imaris software is shown.

Supplementary Figure 4: Microbial dysbiosis between fiber low and fiber high colonic tissue

a-b, 16s rRNA FISH analysis was performed on 5 µm cryosections of distal colon (rectum and sigmoid colon) of fiber-low (n=27) and fiber-high (n=8) HSCR patients using an EUB338-specific probe. **a**, Representative images of a fiber-low and fiber-high colonic tissue probed with Cy3-labelled sense and anti-sense (AS; negative control) EUB338. **b**, Frequencies of translocated bacteria (16sRNA$^+$/DAPI$^+$) were determined using CellProfiler software and are shown as percentage of total DAPI$^+$ cells. **c**, 16S rRNA analysis of colonic tissue from aganglionic (fiber-low n=14; fiber-high n=11) and ganglionic (CD gang. n=4) segments of age-matched HSCR patients. Beta diversity 2D plot show patterns of inter-sample relations using detrended correspondence analysis (DCA). Source data are provided as a Source Data file.

Supplementary Table 5: Core microbiome of 16S rDNA sequencing of colonic tissue

OTU	Samples	Size	Freq	DomOTU	DomSize	DomId	Min	LoQ	Med	HiQ	Max	Taxonomy
OTU1	36	748468	0.339	-	-	-	11	637	13721	36291	84291	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Bifidobacteriales,f:Bifidobacteriaceae,g:Bifidobacterium,s:Bifidobacterium_longum_F18
OTU4	36	187671	0.0851	-	-	-	5	477	2619	7769	26234	k:Bacteria,p:Proteobacteria,c:Gammaproteobacteria,o:Enterobacteriales,f:Enterobacteriaceae,g:Escherichia-Shigella,s:Escherichia_coli_KOEGE_62_[175a]
OTU2	36	48058	0.0218	-	-	-	5	33	219	847	21388	k:Bacteria,p:Proteobacteria,c:Gammaproteobacteria
OTU10	36	48045	0.0218	-	-	-	1	150	423	2473	8624	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Enterococcaceae,g:Enterococcus,s:Enterococcus_faecalis_EnGen0246
OTU35	36	38523	0.0175	-	-	-	22	77	132	297	23544	k:Bacteria,p:Firmicutes,c:Bacilli,o:Bacillales,f:Staphylococcaceae,s:Staphylococcus_aureus_M1256
OTU21	36	29874	0.0135	-	-	-	18	94	178	530	9954	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Corynebacteriales,f:Corynebacteriaceae
OTU42	36	21712	0.00984	-	-	-	28	116	316	668	2844	k:Bacteria,p:Proteobacteria,c:Alphaproteobacteria,o:Rhizobiales,f:Rhizobiales_Incertae_Sedis,g:Phreatobacter
OTU53	36	2804	0.00127	-	-	-	15	15	36	70	465	k:Bacteria,p:Proteobacteria,c:Betaproteobacteria,o:Burkholderiales,f:Comamonadaceae,g:Delftia
OTU43	36	2639	0.0012	-	-	-	3	12	29	68	501	k:Bacteria,p:Firmicutes,c:Bacilli,f:Bacillaceae,g:Anaerobacillus
OTU17	35	39122	0.0177	-	-	-	24	106	264	668	7842	k:Bacteria,p:Proteobacteria,c:Alphaproteobacteria,o:Rhizobiales,f:Phyllobacteriaceae,g:Phyllobacterium
OTU6	34	83714	0.0379	-	-	-	1	12	349	3129	26184	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Bifidobacteriales,f:Bifidobacteriaceae,g:Bifidobacterium,s:Bifidobacterium_bifidum_ATCC_29521_=_JCM_1255_=_DSM_20456
OTU19	34	7827	0.00355	-	-	-	1	8	36	132	1980	k:Bacteria,p:Firmicutes,c:Bacilli,o:Bacillales,f:Bacillaceae,s:Bacillus_subtilis_subsp._subtilis
OTU29	34	5688	0.00258	-	-	-	1	13	47	108	1491	k:Bacteria,p:Firmicutes,c:Bacilli,o:Bacillales,f:Listeriaceae,g:Listeria,s:Listeria_monocytogenes
OTU26	33	11261	0.0051	-	-	-	1	11	48	257	2658	k:Bacteria,p:Proteobacteria,c:Gammaproteobacteria,o:Pseudomonadales,f:Pseudomonadaceae,g:Pseudomonas,s:Pseudomonas_aeruginosa_PAC52
OTU35	33	5295	0.0024	-	-	-	1	8	41	129	1018	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Lactobacillaceae,g:Lactobacillus,s:Lactobacillus_fermentum
OTU5	33	5228	0.00237	-	-	-	2	18	56	136	1566	k:Bacteria,p:Proteobacteria,c:Gammaproteobacteria,o:Xanthomonadales,f:Xanthomonadales_Incertae_Sedis,g:Acidibacter
OTU12	32	20721	0.00939	-	-	-	1	1	5	291	5809	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU50	31	40841	0.0185	-	-	-	1	3	21	100	20809	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Bifidobacteriales,f:Bifidobacteriaceae,g:Bifidobacterium,s:Bifidobacterium_pseudocatenulatum_DSM_20438_=_JCM_1200_=_LMG_10505
OTU831	31	7624	0.00346	-	-	-	1	1	7	63	2538	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Bifidobacteriales,f:Bifidobacteriaceae,g:Bifidobacterium
OTU8	30	21177	0.0096	-	-	-	3	3	34	303	13401	k:Bacteria,p:Firmicutes,c:Negativicutes,o:Selenomonadales,f:Veillonellaceae,g:Veillonella
OTU7	30	23263	0.0105	-	-	-	1	2	21	83	20960	k:Bacteria,p:Proteobacteria,c:Gammaproteobacteria,o:Enterobacteriales,f:Enterobacteriaceae
OTU812	28	12449	0.00564	-	-	-	1	7	29	68	5886	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Family_XI,g:Finegoldia
OTU75	27	60864	0.0276	-	-	-	1	2	36	1975	26467	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU73	27	4646	0.00211	-	-	-	2	10	36	112	2356	k:Bacteria,p:Proteobacteria,c:Alphaproteobacteria,o:Sphingomonadales,f:Sphingomonadaceae,g:Sphingomonas
OTU361	26	9680	0.00439	-	-	-	10	20	70	170	2600	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Enterococcaceae,g:Enterococcus
OTU69	26	3322	0.00151	-	-	-	5	5	15	56	2100	k:Bacteria,p:Proteobacteria,c:Betaproteobacteria,o:Burkholderiales,f:Comamonadaceae,g:Acidovorax
OTU44	25	12070	0.00547	-	-	-	1	3	23	496	4016	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Bifidobacteriales,f:Bifidobacteriaceae,g:Bifidobacterium
OTU32	25	9946	0.00451	-	-	-	2	2	10	98	4090	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Enterococcaceae,g:Enterococcus,s:Enterococcus_pallens_ATCC_BAA-351
OTU99	24	93012	0.0422	-	-	-	2	6	88	4570	36070	k:Bacteria,p:Actinobacteria,c:Coriobacteriia,o:Coriobacteriales,f:Coriobacteriaceae,g:Collinsella,s:Collinsella_aerofaciens
OTU196	24	6264	0.00284	-	-	-	1	6	12	172	2096	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Family_XI,g:Peptoniphilus
OTU541	22	40265	0.0183	-	-	-	1	2	9	318	16032	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Family_XI,g:Anaerococcus
OTU22	22	21390	0.0097	-	-	-	1	8	157	507	13925	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Actinomycetales,f:Actinomycetaceae,g:Varibaculum
OTU114	21	40558	0.0184	-	-	-	1	7	55	1183	27312	k:Bacteria,p:Proteobacteria,c:Betaproteobacteria,o:Burkholderiales,f:Alcaligenaceae,g:Sutterella
OTU14	21	19704	0.00893	-	-	-	1	3	6	30	11191	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU24	21	5238	0.00237	-	-	-	1	6	22	320	1720	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Clostridiaceae_1,g:Clostridium_sensu_stricto_1,s:Clostridium_paraputrificum
OTU41	20	13038	0.00591	-	-	-	1	4	28	1340	3766	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Flavonifractor
OTU20	20	15971	0.00724	-	-	-	1	2	7	202	9414	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU23	19	8516	0.00386	-	-	-	1	1	3	78	5426	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU82	19	47772	0.0217	-	-	-	4	10	28	100	1362	k:Bacteria,p:Proteobacteria,c:Betaproteobacteria,o:Rhodocyclales,f:Rhodocyclaceae,g:Methyloversatilis,s:Methyloversatilis_universalis_EHg5
OTU16	19	2996	0.00136	-	-	-	2	2	6	26	23950	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Actinomycetales,f:Actinomycetaceae,g:Actinomyces,s:Actinomyces_turicensis_ACS-279-V-Col4
OTU55	17	78452	0.0356	-	-	-	2	8	24	31	77484	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Lactobacillaceae,g:Lactobacillus,s:Lactobacillus_casei_5b
OTU11	17	15444	0.007	-	-	-	2	6	88	164	10006	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Actinomycetales,f:Actinomycetaceae,g:Actinomyces,s:Actinomyces_europaeus_ACS-120-V-Col10b
OTU34	16	13394	0.00607	-	-	-	1	2	5	1078	13099	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Actinomycetales,f:Actinomycetaceae,g:Actinomyces
OTU38	16	4860	0.0022	-	-	-	1	2	8	34	4592	k:Bacteria,p:Cyanobacteria,o:Chloroplast
OTU30	16	17280	0.00783	-	-	-	2	2	18	334	7174	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Rikenellaceae,g:Alistipes,s:Alistipes_sp._AL-1
OTU31	16	5940	0.00269	-	-	-	2	2	25	159	3221	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Lachnoclostridium
OTU56	16	2690	0.00122	-	-	-	2	2	46	150	1418	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Blautia
OTU869	15	18179	0.00824	-	-	-	4	5	45	1432	11357	k:Bacteria,p:Firmicutes,c:Negativicutes,o:Selenomonadales,f:Veillonellaceae,g:Veillonella,s:Veillonella_sp._FFA-2014
OTU25	15	47392	0.0215	-	-	-	4	4	12	384	38992	k:Bacteria,p:Actinobacteria,c:Coriobacteriia,o:Coriobacteriales,f:Coriobacteriaceae,g:Atopobium,s:Atopobium_minutum_10063974
OTU36	15	13810	0.00626	-	-	-	2	4	42	118	5966	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Porphyromonadaceae,g:Parabacteroides
OTU51	13	4162	0.00189	-	-	-	2	5	48	308	2476	k:Bacteria,p:Proteobacteria,c:Deltaproteobacteria,o:Desulfovibrionales,f:Desulfovibrionaceae,g:Bilophila,s:Bilophila_wadsworthia_3_1_6
OTU47	13	3704	0.00168	-	-	-	2	2	3	162	3021	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Blautia
OTU49	13	3246	0.00147	-	-	-	1	2	12	94	2146	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Fusicatenibacter
OTU54	13	2864	0.0013	-	-	-	1	11	53	372	1164	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Clostridiaceae_1,g:Clostridium_sensu_stricto_1,s:Clostridium_perfringens_C_str._JGS1495
OTU37	13	2692	0.00122	-	-	-	2	2	95	135	952	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Bacteroidaceae,g:Bacteroides
OTU28	12	31270	0.0142	-	-	-	3	4	16	59	30279	k:Bacteria,p:Verrucomicrobia,c:Verrucomicrobiae,o:Verrucomicrobiales,f:Verrucomicrobiaceae,g:Akkermansia
OTU68	11	5809	0.00263	-	-	-	1	1	2	897	3587	k:Bacteria,p:Firmicutes,c:Bacilli,o:Lactobacillales,f:Lactobacillaceae,g:Lactobacillus,s:Lactobacillus_rhamnosus_DSM_20021_=_JCM_1136_=_NBRC_3425
OTU925	10	15240	0.00691	-	-	-	1	5	637	1952	9648	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae
OTU40	10	6673	0.00302	-	-	-	1	5	5	207	3953	k:Bacteria,p:Actinobacteria,c:Actinobacteria,o:Actinomycetales,f:Actinomycetaceae,g:Actinobaculum,s:Actinobaculum_massiliae_ACS-171-V-Col2
OTU62	9	3685	0.00167	-	-	-	1	1	215	761	1500	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Faecalibacterium
OTU33	9	8666	0.00393	-	-	-	2	2	10	95	5933	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Anaerotruncus,s:Clostridiales_bacterium_VE202-13
OTU52	9	3327	0.00151	-	-	-	1	1	2	3	3079	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Roseburia
OTU45	8	2509	0.00114	-	-	-	1	1	1	37	1398	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Faecalibacterium
OTU27	8	7846	0.00356	-	-	-	2	38	124	1712	5452	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Blautia
OTU57	8	5962	0.0027	-	-	-	1	4	70	208	5591	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:Lachnoclostridium
OTU70	8	3781	0.00171	-	-	-	1	1	1	445	2946	k:Bacteria,p:Proteobacteria,c:Betaproteobacteria,o:Neisseriales,f:Neisseriaceae,g:Eikenella
OTU101	8	3273	0.00148	-	-	-	3	3	132	1346	1581	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Prevotellaceae,g:Prevotella
OTU77	7	14998	0.0068	-	-	-	3	3	17	481	14392	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Subdoligranulum
OTU46	7	6330	0.00287	-	-	-	2	4	6	88	6174	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Faecalibacterium
OTU63	7	3574	0.00162	-	-	-	1	3	35	275	3217	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Ruminococcaceae,g:Ruminococcaceae_UCG-005
OTU100	7	2778	0.00126	-	-	-	2	2	18	958	1250	k:Bacteria,p:Bacteroidetes,c:Bacteroidia,o:Bacteroidales,f:Porphyromonadaceae,g:Parabacteroides
OTU87	6	3612	0.00164	-	-	-	1	30	46	1616	1884	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:[Eubacterium]_hallii_group
OTU	6	2824	0.00128	-	-	-	1	1	74	172	2574	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Family_XI,g:Ezakiella
OTU	5	4055	0.00184	-	-	-	1	30	300	387	3364	k:Bacteria,p:Actinobacteria,c:Coriobacteriia,o:Coriobacteriales,f:Coriobacteriaceae,g:Slackia
OTU	5	2420	0.0011	-	-	-	8	38	136	156	2082	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Coriobacteriaceae,g:Slackia
OTU	4	2856	0.00129	-	-	-	2	32	950	1872	1872	k:Bacteria,p:Firmicutes,c:Clostridia,o:Clostridiales,f:Lachnospiraceae,g:[Eubacterium]_hallii_group

Supplementary Table 6: Overview of retrospective HSCR patient cohort

Study Number	Sex	Age at Surgery (months)	OP technique: Del la Torre (1); Soave transnal (2); Soave transabdominal (3); Swenson transanal (4); Swenson transabdominal (5); Duhamel (6); Rehbein (7)	Lenght of AG	Fiber score	Postoperative clinical symptoms: Constipation (1); Incontinence (2); Enterocolitis (3); Pain (4); Impaired growth (5); Stenose/Ileus (6)	Postoperative clinical treatment: Laxative (1); Clyster (2); Botox (3); Antibiotics (4); Colonic irrigation (5); Pre-/probiotics (6)
1	F	22	2	L-HSCR	low	3, 6	4,5,6
2	F	7	6	TCA	low	1,3,5,6	2,4
3	M	2	1	S-HSCR	high	1	3,5,6
4	M	171	1	TCA	low	no	no
5	M	9.5	1	TCA	high	1	6
6	M	2	1	L-HSCR	low	3, 6	3,4,5,6
7	M	3.5	1	S-HSCR	high	1,2	2
8	M	1	1	S-HSCR	high	no	no
9	M	10.7	1	S-HSCR	high	1,6	5
10	F	25	2	S-HSCR	high	no	no
11	M	48	1	S-HSCR	high	1,3,6	1,2,3,4,5
12	M	52	2	S-HSCR	high	2	1,5,6
13	M	1	1	L-HSCR	low	1,2,3	2,4,5
14	M	28.5	1	S-HSCR	high	2	6
15	M	4	7	TCA	low	1,3,4,5,6	4,5,6
16	F	2.8	1	TCA	low	3	4,5
17	M	3.7	1	S-HSCR	low	3	no
18	M	8	6	S-HSCR	low	1,3	1,5
19	M	5	1	L-HSCR	high	2	no
20	M	7	1	S-HSCR	high	3	4
21	M	9	1	TCA	low	3,5	3,4,5
22	M	60	2	S-HSCR	high	no	no
23	M	67	1	L-HSCR	high	2,6	6
25	M	1.5	1	S-HSCR	low	3	1,2,3,4,5
27	M	3.5	1	S-HSCR	low	no	no
30	M	3	1	L-HSCR	low	1,3	3,4,5,6
31	M	4.3	1	S-HSCR	low	3,6	5
32	M	1.5	1	S-HSCR	low	no	no
33	M	2	1	L-HSCR	high	5	no

Length of aganglionose: TCA (Total colonic aganglionose); L-HSCR (long-segmented HSCR); S-HSCR (short segmented HSCR). Female (F); Male (M). Source data are provided as a Source Data file.

Supplementary Table 7: Characteristics retrospective HSCR patient cohort

Characteristics of retrospective HSCR patient cohort

Anthropometric data and potential risk factors	All (n=29)	Fiber-low (n=15)	Fiber–high (n=14)	p-value
Age at operation, months	5 (1–171)	3 (1–127)	9.5 (1–67)	0.063
Sex, male	25 (86)	12 (80)	13 (93)	0.316
Preterm birth	2 (7)	1 (7)	1 (7)	0.960
Antibiotic treatment	21 (72)	12 (80)	9 (64)	0.550
Caesarean Section	12 (41)	6 (40)	6 (43)	0.876

Values are mean (SD), median (range) or number (percentage); n: number; SD: standard deviation; Differences between the fiber high/low group of children were tested with the Wilcoxon rank-sum test and $\chi2$-test, as appropriate.

Supplementary Table 8: Antibodies for FACS and Histology

Antibodies for Flow cytometry

Name	Clone	Fluorocrome	Supplier
CD3	SK7	PerCP	BioLegend
HLA-DR	L243	FITC	eBioscience
CD45	HI30	APC	BD Biosciences
CD14	HCD14	PerCP	BioLegend
CD64	10.1	PE-Cy7	BioLegend
CD11c	3.9	Alexa700	eBioscience
CD4	RPA-T4	FITC	BD Biosciences
CD127	A019D5	PE	BioLegend
CD25	BC96	Alexa 647	BioLegend
NKp44	p44-8	BB515	BD Biosciences
CD11c	3.9	PerCP/Cy5.5	BioLegend
CD19	HIB19	PerCP	BioLegend
CD56	HCD56	PerCP	BioLegend
FcεRI	AER-37	PerCP	Novus Biologicals
TCR α/β	IP26	PerCP/Cy5.5	BioLegend
TCR γ/δ	B1	PerCP/Cy5.5	BioLegend
CD117 (c-kit)	104D2	APC	BioLegend
CD294 (CRTH2)	BM16	PE-Cy7	BioLegend
CD161	HP-3G10	Alexa 700	BioLegend
IL-17A	eBio64DEC17	APC	eBioscience
IFN-γ	B27	Alexa 700	BioLegend
Rorγt	Q21-559	Alexa 488	BD Biosciences
Foxp3	PCH101	PE-Cy7	eBioscience
CD16/CD32 (Fc block)		purified	BD Biosciences

Primary Histology antibodies

Name	clone	fluorocrome	host/subtype	supplier
CD64	10.1	purified	mouse IgG1	BioLegend
CD3	UCHT1	Alexa 488	mouse IgG1	BioLegend
Acetylcholinesterase (AchE)	HR2	purified	mouse IgG2b	Abcam
Beta III tubulin	2G10	purified	mouse IgG2a	Abcam
Beta III tubulin	GT11710	purified	mouse IgG2b	Genetex
Tyrosine Hydroxylase (TH)	1A5	purified	mouse IgG2a	Invitrogen
nNOS	16/nNOS/NOS type	purified	mouse IgG2a	BD Bioscience
S100b	11C12E12	purified	mouse IgG1	Biolegend
VIP	polyclonal	purified	guinea pig	Peninsula Laboratories
IL-23p19	23dcdp	PE	mouse IgG2b	eBioscience

Secondary Histology antibodies

Name	clone	fluorocrome	host/subtype	supplier
anti-mouse IgG1	polyclonal	Alexa 488	goat	Invitrogen
anti-mouse IgG1	polyclonal	Alexa 555	goat	Invitrogen
anti-mouse IgG1	polyclonal	Alexa 647	goat	Invitrogen
anti-mouse IgG2a	polyclonal	Alexa 647	goat	Invitrogen
anti-mouse IgG2a	polyclonal	Alexa 555	goat	Invitrogen
anti-mouse IgG2b	polyclonal	Biotin	goat	BD Biosciences
anti-mouse IgG2b	polyclonal	Alexa 647	goat	Invitrogen
anti-guinea pig	polyclonal	Alexa 555	goat	Invitrogen
Streptavidin		Cy3		BioLegend

Supplementary Table 9: Primers

List of Primers

Gene	Forward Sequence 5'-3'	Reverse Sequence 5'-3'
IL-17A	CTA CCA CCG ATC CAC CTC AC	GCG TTG ATG CAG CCC AAG TG
IL-6	TAC ATC CTC GAC GGC ATC TC	GCC ATC TTT GGA AGG TTC AG
IL-22	ACG GAG TCA GTA TGA GTG AG	CAC CAC CTC CTG CAT ATA AG
TNF-α	GCT GCA CTT TGG AGT GAT CG	GGG CTA CAG GCT TGT CAC TC
IL-23	GTG GGA CAC ATG GAT CTA AG	AGA CCC TGG TGG ATC CTT TG
IL-1β	ACC TCC AGG GAC AGG ATA TG	CGC AGG ACA GGT ACA GAT TC
CX3CR1	CAT CGT GGT CTT TGG GAC TG	TTG GTG AGG GCA AAC ACT AC
CCR2	ATA CCA GGA CTG CCT GAG AC	CGC TCT CGT TGG TAT TTC TG
β2M	CAG CGT ACT CCA AAG ATT CA	GAA TGC TCC ACT TTT TCA AT

Supplementary Figure 5: Fiber innervations grades for Fiber scoring

a

grade 1:
absent innervation

grade 2:
low innervation

grade 3:
intermediate innervation

grade 4:
high innervation

b

a, AChE immunohistochemistry of 5µm cryosections of distal aganglionic colon showing 4 different mucosal innervation grads. Grades 1 and 2 were grouped into fiber-low and 3 and 4 were grouped into fiber-high HSCR patients. **b**, Using brightfield microscopy and CellSens Dimension Software the innervation grade was quantitatively confirmed in 3 representative patients per group (for details see methods). Scatter dot plots show means ± SEM. Source data are provided as a Source Data file.