Collective motion as a distinct behavioral state of the individual

D. Knebel[1,2], C. Sha-ked[1], N. Agmon[2], G. Ariel[3]* and A. Ayali[1,4]*

[1] School of Zoology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv, Israel.

[2] Department of Computer Science, Bar-Ilan University, Ramat-Gan, Israel.

[3] Department of Mathematics, Bar Ilan University, Ramat-Gan, Israel.

[4] Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel.

* Corresponding author

Abstract

Collective motion is an important biological phenomenon, ubiquitous among

different organisms. Locusts are a quintessential example of collective motion: while

displaying a minimal level of cooperation between individuals, swarms of millions are highly

robust and persistent. Using desert locusts in a series of carefully controlled laboratory

experiments, we show that locust coordinated marching induces a distinct behavioral mode

in the individual that we term a "collective-motion-state". Importantly, this state is not

induced by aggregation or crowding per-se, but only following collective motion. It is

manifested in specific walking behavior kinematics and has a lasting effect. Simulations

confirm that the collective-motion-state provides an individual-based mechanism, which

increases the robustness and stability of swarms in the presence of fluctuations. Overall, our

findings suggest that collective-motion is more than an emergent property of the group, but

a specific behavioral mode that is rooted in endogenous biological mechanisms of the

individual.

Introduction

The ability to form groups that move collectively is a key behavioral feature of many

species [1,2], that increases the survival of both individuals and groups [3,4]. Collectively moving

organisms, however, differ in the levels of peer-to-peer interactions, ranging from minimal

cooperation to complex social behaviors [5,6]. Analysis of both experiments and theoretical

modeling of collective motion suggests a highly stochastic (or chaotic) movement pattern,

which makes the trajectory of individuals difficult to predict in practice [7–9]. Furthermore,

endogenous variability among individuals, heterogenic environments, and variability in the

interactions between the individual and its direct environment, are all sources of

randomness that may affect the coordinated behavior of the collective. Accordingly, it is not

clear how synchronized collective motion constitutes such a robust phenomenon,

maintaining its form across various group sizes and densities, and under heterogeneous and

unpredictable environmental conditions.

One of the most interesting, albeit disastrous, examples of collective motion is that

of the marching of locusts. These insects swarm in groups of millions, migrating in mass

across large distances, devastating crops and agriculture [10–12]. In the context of social

interactions, locust swarming is characterized by a minimal level of cooperation between

individuals: collectivity is mostly manifested in alignment among neighboring individuals and

in maintaining the overall movement in the same general direction. Nonetheless, the locust

swarming phenomenon is extremely robust, with huge swarms demonstrating moderate to

high collectivity on huge scales (up to 6-7 orders of magnitudes), in terms of both the

number of animals and their spatio-temporal distribution [13,14] (see further references in [7]).

Thus, locusts exhibit a considerable disparity between little local cooperation and large-

scale collectivity.

What is the key to this ability of locust swarms to maintain their integrity? Here, we

show by a series of carefully controlled behavioral experiments that collective movement

induces an internal switch in individual gregarious locusts, activating a behavioral mode we

refer to as a "collective-motion-state". In this state, the kinematic behavior of individuals

notably differs from that during the non-collective state. It is important to emphasize that

by "collective" and the "non-collective" states we are not referring to the well-known

solitarious-gregarious phase transition in locusts [10,11].

How does the collective-motion-state affect the formation and robustness of the

swarm? Interestingly, the switch into this state seems to occur rapidly, and in response to

coordinated walking. In particular, our experiments indicate that aggregation alone is not

sufficient. Switching out of the collective-motion-state occurs over a longer time scale –

significantly longer than the typical time scale of fluctuations in the swarm dynamics. Hence,

stochastic fluctuations are "smoothed-out", leading to highly robust dynamics of the swarm

collective behavior.

This ability to switch between socio-behavioral states appears to be beneficial for

the swarm integrity. Using a simplified computer model, we simulated the swarming

properties of locust-like agents with different kinematic parameters, representing the

different behavioral states. The results demonstrate that the ability to switch between the

distinct behavioral states, as a consequence of both instantaneous and past stimuli, leads to

swarms with tighter groups, compared for example to fixed or continuously changing

dynamics, and without affecting the within-group alignment. Hence, we conclude that the

collective-motion-state provides an individual-based mechanism that increases the stability

of swarms in the presence of fluctuations, preventing the swarm from collapsing.

Results

The main objective of this report was to examine the behavior of individual animals

upon joining or leaving a group of conspecifics. Here we studied gregarious locusts, reared

in dense populations, one developmental stage before becoming adults and developing

functional wings (i.e. fifth larval instar). The experiments comprised three consecutive

stages, representing different conditions (Fig. 1 and Vid. S1): (1) Isolation stage: a single

animal was taken from its highly dense rearing cage, tagged with a barcode, and introduced

alone into a ring-shaped arena (outer and inner diameters: 60 cm and 30 cm, respectively);

(2) Grouping stage: nine other individually tagged animals were added to the arena; and (3)

Re-isolation stage: the nine added animals were removed from the arena, leaving the

original animal alone. The duration of each stage was one hour, during which the animals'

trajectories were fully reconstructed using a barcode tracking system. The middle 40

minutes of each stage were analyzed, as detailed in the Methods section. A range of

kinematic statistics was collected in order to classify and compare the locusts' behavior in

the different stages.

Swarm formation – validation of collective motion

In order to verify that our grouping conditions were indeed inducing collective

motion (swarming), we calculated the synchronization in movement of the grouped animals

(the order parameter, see Methods for definition). The median order parameter in the

grouping stage was found to be significantly higher than that obtained for computationally

randomized groups (as presented in our previous report, Knebel et al., 2019); medians:

0.348 and 0.239 respectively; Wilcoxon rank-sum test: $p < 0.01$). Consequently, we

conclude that the groups in our experimental setup indeed demonstrate swarming and

collective motion.

Figure 1. A schematic flow of the experimental procedure. Locusts were reared in high density conditions. The

Experiments comprised the following consecutive stages: (1) isolation for one hour in the arena; (2) grouping

for one hour; and (3) re-isolation for one hour.

Kinematic differences among the isolation, grouping, and re-isolation stages

Locusts walk in an intermittent motion pattern [16,17], i.e., movement occurs in

sequences of alternating walking bouts and pauses. To characterize individual locust

kinematics, we measured four parameters: (1) the fraction of time an animal spends

walking; (2) the average speed while walking; (3) the average walking bout duration; and (4)

the average pause duration. Comparing these values across the three experimental stages,

we found several statistically significant differences (Fig. 2). In the following, *p*-values

correspond to a Friedman test followed by a multiple comparison test using the Bonferroni

method.

We found that when comparing between isolation and grouping, the fraction of time

spent walking and the pause durations differed significantly ($p<0.01$ and $p<0.001$,

respectively), showing a larger fraction of time walking and shorter pauses while grouping.

These findings are in accordance with previous reports [15], and are consistent with the

known propensity of locusts to walk more and rest for shorter times while in a swarm [15–18].

However, our experiments also revealed a new effect of swarming. Comparing the isolation

and re-isolation stages, we found that all the studied parameters differed significantly.

Specifically, the fraction of time walking, speed, and walking bout duration were all higher in

the re-isolation stage, while the pause duration decreased ($p<0.001$, $p<0.001$, $p<0.05$ and

$p<0.05$, respectively). Furthermore, comparing the grouping and re-isolation conditions

revealed that the walking bout duration increased significantly following re-isolation

($p<0.01$). The data for all other comparison combinations were found not to differ

significantly. The increase in activity following re-isolation is surprising, and suggests that

the marching behavior of locusts is not dictated by instantaneous or immediate interactions

among individuals per-se. Rather, our findings indicate that the interactions with other

marching locusts induce a switch to a new internal behavioral state, which outlasts the

presence of the swarm. In accordance with our results, we term this internal state the

"collective-motion-state".

In order to verify that the observed behavioral changes indeed represent a transient

state, rather than a permanent behavioral modulation, a simple control experiment was

performed. Following the re-isolation stage, the locusts were returned to their rearing cage

(in high crowding conditions without collective motion), and tested the next day again alone

in the arena (isolation condition). We found no significant behavioral difference between

this latter isolation and the first isolation stage of the previous day (n = 6). Hence, the

collective-motion state is transient. A second series of control experiments was performed

in order to exclude potential time effects on the locusts' behavior due to the duration of the

experiments. To this end, locusts were tested in isolation for three consecutive hours. The

kinematic analysis procedure described above was then performed separately on three 40

min segments of the 3 h tests and no significant differences were found (n = 6).

Figure 2. Kinematic changes throughout the three experimental conditions. (a) the fraction of walking, (b) the averaged walking speed, (c) the average duration of walking bouts, and (d) the average duration of pauses, of the traced animals in the isolation, grouping, and re-isolation stages. *$p<0.05$, **$p<0.01$, ***$p<0.001$.

Consistency in individual behavioral tendencies

Despite the major differences in behavioral kinematics among the three experimental stages, we found that individual behavioral tendencies persisted. The fraction of walking, speed, and pause durations all showed high within-individual correlation across the three stages. The walking bout durations, however, were significantly correlated only between the isolation and grouped stages (see Table 1 for numerical details).

The increased values of this latter parameter upon re-isolation may reflect a mechanism that is activated when an animal finds itself away from the group, and overrides its prior behavioral tendencies. To test this hypothesis, and further explore the effects of the walking bout duration on the ability (and efficiency) of locusts to swarm, the effect of the different measured kinematic parameters was further studied using a simplified computer model.

Modeling and simulations

The experiments described above demonstrated that individual locusts introduced into a collectively moving swarm undergo a switch into a distinct internal socio-behavioral state. However, whether this change confers a benefit on the swarm formation and maintenance, and if so, of what kind, remained unanswered. To explore this aspect of the collective-motion-state, we developed a model that simulates locust swarms, in which individual kinematics and population size could be manipulated. We focused ,in particular, on the walking bout duration, as this specific parameter seems instrumental in differentiating the locust behavior after re-isolation (Table 1).

To simulate swarms, we used a simplified agent-based model in a square domain with periodic boundaries (size: 10 x 10). Agents were designed as rectangles with a circular

receptive field around their center. The agents' kinematics were programed to resemble

that of locusts, i.e. to move in an intermittent motion (pause-and-go) pattern: at every step

of the simulation, each agent made an individual decision whether to walk or stop, based on

its current state (walking or stopping) with predefined probabilities (p_W and p_S, respectively).

While moving, the speed was constant. The individual direction of movement was allowed

to change only when an agent changes its state from stopping to walking [16]. An agent's new

direction was a weighted sum of its own direction (inertia), the direction of other agents in

its visual field, with a short memory (see [19] for the importance of memory in pause-and-go

simulations), and noise. See Methods for details and Table S1 for parameter values.

The spatial and temporal scales of the model were set as follows: 1 cm was

considered as a distance of 0.1 in the simulated arena, and 1 sec corresponded to 1

simulation step. Thus, we fixed the model dimensions to correspond with real locusts' size

and movement parameters. The size of agents (0.1 x 0.4) maintains the proportions of 5th

larval instar locusts. Additionally, the visual range of 1 radius, ten times larger than the

agent's rectangle width, represents the proportional visual field of locusts (range just under

10 cm [16]). To adjust the temporal scale, an average stopping duration in the model was set

to be about 3 steps (by setting p_S to be 0.67; see Fig. S1), parallel to the median of 3 second

pause duration found in the grouping condition in the empirical experiment (Fig. 2d). The p_W

values were set to range between 0.6 and 0.92, making the walking bout durations

comparable with the empirical data (compare Fig. 2c and Fig. S1). Finally, the speed was set

to 0.25, corresponding to the typical swarming speed shown in Fig 2b.

In order to study the effect of individual bout duration on the swarm, we

concentrate on the effect of swarm size, N, and on p_W, describing the propensity of animals

to keep walking. We evaluated the effect of these parameters on four statistics that

characterize collective motion in the swarm:

1. The order parameter (the size of the average direction vector).

2. Spread (the average distance between all pairs).

3. The number of neighbors (within the field of view).

4. Regrouping time (the average number of steps it takes an agent that has lost all

other agents in its receptive field to re-obtain at least one neighbor).

Statistics were averaged over all agents and simulation steps. In order to allow comparison

with our experimental findings, we focus on relatively small swarms (up to 40 agents). Fig. 3

shows the median value over 40 independent repetitions for each parameter set. We found

that the order parameter increases both with p_W and with the number of agents, N (Fig. 3a).

The spread shows a more complex dependence on p_W, depending on N (Fig. 3b). For small

swarms, large p_W leads to a smaller spread, but the trend is reversed for large swarms,

showing a smaller spread with small p_W. The number of neighbors (NN) is decreases with p_W

but increases with N (Fig. 3c), and the slope is much larger for large swarms. For small

swarms, the NN is small and practically independent of p_W. The regrouping time varies

significantly with swarm size (increasing by a factor of up to 9) and is mostly independent of

p_W for large groups (Fig. 3d). However, small groups (up to 20) reduce their time to regroup

with higher p_W. This result makes sense assuming that isolated individuals perform a

random search until they join a new cluster. Because agents only change direction when

starting to move, the length of walking segments between rotations increases with p_W.

Figure 3. The influence of different p_W values on the collectivity parameters of simulated swarms. (a) The order

parameters, (b) spread , (c) the average number of agents within each agent's visual field , and (d) the average

number of steps to regroup, for different p_W values (rows), and different group sizes (columns). Each colored

box represents the median of 40 simulations.

Fig. S2-S6 show a detailed sensitivity analysis, demonstrating the differences

obtained when changing individual parameters. This analysis shows that, despite some

quantitative differences, our qualitative observation that the propensity of animals to walk

(modelled by the parameter p_W) has a different effect on small and large groups, holds in a

wide range of parameters. However, some parameters do show a notable effect on the

collective dynamics: increasing the noise level substantially (Fig. S3) may completely destroy

collectivity, leading to a small order parameter. This observation is related to the well-

known phase transition in Vicsek-like models [20]. In addition, several physical parameters,

such as the visual range (Fig. S4), speed (Fig. S5), and the size of agents (Fig. S6) have a

partial influence on some of the statistics. It is therefore important that such parameters are

assigned realistic values, as measured from experiments.

Overall, these findings indicate that large p_W values, namely, longer walking bouts,

increase the chances that low- density groups will stay together (smaller spreads, shorter

time to regroup). However, for larger swarms, smaller p_W – shorter walking bouts – is

favorable (more neighbors, smaller spread and a marginal cost in the order parameter and

regrouping time). Therefore, if aggregation is essential to individuals, our results suggest

that it might be beneficial for an agent to change its p_W as a function of the group size.

Sociality-dependent p_W

Several previous theoretical models of collective motion have found that adaptable

interactions, i.e., interactions that can change according to the local environment, can

improve the cohesiveness and performance of swarms [21–23]. Accordingly, in light of our

experimental results, we sought to study the effect of the individual's ability to change its

walking properties. To this end, we introduced an experience-dependent p_W to the

simulated agents (Fig. 4). The p_W could change during a simulation as a result of the agent's

receptive field of view in the last simulation steps. For simplicity, we assume linear

dependence (Ariel et al., 2014b). Biologically, such a dependence may be due to the way

locusts perceive their surrounding: many locusts walking in the same direction around one

locust would affect it more than only a few. To be precise, denote by $\bar{n}_i(t)$ the average

number of neighbors of a focal animal i in the previous k (memory length) steps before time

t, the probability that animal i keeps walking, $p_W(i,t)$ was given by

$$p_w(i,t) = \begin{cases} 0.92 - \alpha\bar{n}_i(t), & 0.92 - \alpha\bar{n}_i(t) \geq 0.6 \\ 0.6, & 0.92 - \alpha\bar{n}_i(t) < 0.6 \end{cases}$$

where α is a constant determining the slope of the linear effect ($\alpha = 0.1$ in Fig. 4). As can be

seen, p_W was ranged between 0.6 and 0.92, in order to avoid phlegmatic behavior or

unnatural hyperactive of the agent (respectively). See Fig. S7 for the obtained p_W values and

walking bout durations in those simulations.

Figure 4. The influence of adaptive p_W values with different memory lengths on the collectivity parameters of

simulated swarms. (a) The order parameters, (b) spread , (c) the average number of steps to regroup, and (d)

the average number of agents within each agent's visual field, for different memory length (rows) and group

sizes (columns). Each colored box represents the median of 40 simulations.

This conditional individual behavior was found to minimize the spread and

regrouping time, while maximizing the agent's number of neighbors for almost all swarm

sizes (Compare Fig. 3 and 4). This conditional behavior, however, did not increase the order

parameter above that in the fixed p_W case (Fig. 4a).

We tested different memory lengths that affected the adaptive p_W. The average p_W

did not change dramatically (Fig. S7b) but did result in higher variance across the simulation

steps – the shorter the memory was, the higher the step-to-step p_W fluctuations were (Fig.

S7c). This behavior is expected, as the sensitivity to the current situation of the agent

decreased with memory length. Fig. 4 shows that while the order and spread favors short

memory, the NN and steps to regroup were largely indifferent to it.

Overall, our simulations show that increasing the walking bout duration when an

agent finds itself alone has a beneficial impact on its chances of meeting other agents and

maintaining high group density. In regard to the empirical results, our simulations suggest

an instrumental role for the increased bout duration in regrouping when an animal is

isolated after swarming.

Discussion

Our findings reported here suggest that, in locusts, the sensory-motor act of

collective motion is accompanied by an internal state of the individual locust – a collective-

motion-state that is manifested in specific behavioral kinematics. This state is induced by

the experience of synchronous, collective marching. In turn, it has an important role in

maintaining the integrity and consistency of the swarm. Below we discuss several key

aspects and implications of this finding.

In a recent study [15], we have introduced a comparison between the walking behavior

kinematics of individual gregarious (crowded-reared) locusts in different social (density)

contexts. Our reported findings are reconfirmed and further elucidated here by the results

of the initial isolation and grouping stages in our experiments. The novel idea posited here is

that these differences represent and are induced by not only the spatially and temporally

immediate social environment, and the instantaneous local interactions among locusts, but

also by the effects of an internal state induced by the general experience of collective

motion. A fundamental aspect of the concept of the collective-motion-state arises from our

findings related to its persistent effect in time: Upon re-isolation, the individual locust

adopted behavioral kinematics that critically differed from that in the first experimental

stage (initial isolation). We also showed that, as expected, the collective-motion-state is

transient. If the locust does not experience collective motion for some time, and is then

isolated once more, it loses the unique walking-related kinematics it previously adopted in

response to the collective motion, i.e. the internal collective-motion-state.

The individual locusts in our experiments retained the variability observed among

them (see also Knebel et al., 2019), while demonstrating a second layer of variability or

plasticity upon experiencing collective motion, when entering the collective-motion-state.

Considerable research has been devoted in recent years to understanding the effect of

variability among individuals on the group's collective behavior, both experimentally—

ranging from bacteria to primates [24–33]—and theoretically [34–40]; (see [41–43] for recent reviews).

The interactions between variability in specific aspects of the individuals' behavior and

group-level processes were found to be complex and, moreover, bidirectional (e.g. [15]).

Variability among individual animals was found to have important consequences for the

collective behavior of the group (e.g. [44,45]).

However, beyond the variability among the individuals composing a group,

variability is also expected in the behavior of the individual animal over time, as it

experiences changes in environmental and social conditions. The swarm (or flock, shoal,

herd, etc.) is a heterogeneous entity, moving in a heterogeneous environment. The

individual is bound on occasion to find itself in different locations within the swarm (leading

edge, at the outskirts, trailing), and it may also find itself separated from the group by

natural obstacles (vegetation, rocks and boulders). It is essential for the robustness and

consistency of the swarm that throughout these changing conditions, the behavior of the

individual will adapt accordingly, such as to be appropriate for the changing context. For

example, if temporarily separated from the core of the swarm, a locust's walking kinematics

should change to support rapid reunion with the group, as reported in both our

experimental and simulation findings (e.g. increased fraction of walking and duration of

walking bouts). If previously naïve to collective motion, that individual's kinematics would

however be disadvantageous, or even hinder the formation of a swarm.

In [17], the authors suggest that behavioral variability can be explained by the existence of

two internal states. Studying single locusts in isolation for eight consecutive hours, they

have observed changes in behavioral kinematics that were suggested to result from

"internal state behavioral modulation". The observed variations, however, were merely

attributed to changes in "starvation/satiation state", i.e. as the locust becomes starved, it

changes its walking behavior, searching more vigorously for food. Moreover, they conclude

that animals continually switch between the two states on a scale of minutes. The

collective-motion-state reported here is, of course, a very different type of internal

behavioral state, which is strongly involved with the locust past and current social

environment. It may be viewed as a form, or a manifestation of a social carry-over effect [46],

where a social environment experienced by a focal individual affects aspects of its

locomotion behavior at a later, non-social context. As noted, however, the change in

behavioral state described here is induced by collective marching, i.e., a particular mode of

social interaction, rather than by aggregation or being around other conspecifics per-se.

Moreover, as our simulations show, the enhanced marching displayed in the re-isolation

stage is advantageous for maintaining collective swarming - it is still much related to the

social context rather than carried-over to a non-social one.

It should be stressed once again that the current study focused on gregarious,

crowded-reared locusts only. The described behavioral states should not be confused,

therefore, with the well-known and much researched locust density-dependent phase

polyphenism [10,11]. Collective motion is limited to the gregarious, swarming and migrating

phase. Accordingly, all our experimental animals were taken from our gregarious breeding

colony, maintained in crowded conditions for many consecutive generations. In their

breeding cages, mostly due to the physical constraints and abundance of food, despite

experiencing high density the locusts very rarely, if at all, demonstrate collective motion.

Thus, they adopted the collective-motion-state only upon experiencing and taking part in

collective marching within the experimental arena.

The reported collective-motion-state is also in accord with the overall daily

behavioral changes of marching locust swarms. The swarm will spend the night (as well as

times of low temperature or other unfavorable climatic conditions) roosting among the

vegetation. Upon suitable conditions, after a period of feeding, the locusts will initiate

marching – highly synchronized, collective motion. Frequently, when temperature becomes

too high around noon, or when dusk arrives, the swarm will again switch to feeding and

roosting. These daily patterns call for corresponding changes in the internal behavioral

states of the individual locusts and mostly a dedicated collective-motion-state.

In the current work we are cautious in talking about *behavioral* states. Although this

is beyond the scope of this study, it is clear that these behavioral states represent

physiological states. With some confidence, we can speculate about the nature or the

physiological mechanisms involved in the demonstrated behavioral states. Behavioral

plasticity in locust behavior has been attributed to various second messengers or

neuromodulators, or to the balance among them. Most notable are the biogenic amines

(e.g. Serotonin, a prominent bio-amine, was recently reported to inhibit walking behavior in

Drosophila [47]). Hence, it may well be that the (spatial and temporal) immediate social

environment affects biogenic amine levels, and these in turn modulate the walking-related

behavioral kinematics manifested in the different behavioral states.

Another candidate that may be involved in the collective-motion-state is the locust

adipokinetic hormone (AKH). AKH is a metabolic neuropeptide principally known for its

mobilization of energy substrates, notably lipid and trehalose, during energy-requiring

activities such as flight and locomotion, but also during stress (e.g. [48]). It is well accepted

that the metabolic state affects the level of general activity of an organism, and AKHs are

reported to stimulate locomotor activity, either directly by way of their activity within the

central nervous system (e.g. [49]), or via octopamine – a biogenic amine with ample

behavioral effects [50,51].

Furthermore, as noted, we have demonstrated here an extended effect of the

experience of collective motion. Hence, learning and memory-related mechanisms would

also seem to be involved. Again, previous work may suggest some candidate molecules and

pathways, including cGMP-dependent protein kinase (PKG), and Protein Kinase A [52–54].

Last, as noted, solitarious phase locusts lack the capacity to demonstrate collective

motion, and thus also the collective-motion-state. Accordingly, they differ from gregarious

locusts in all the above physiological pathways (bioamines: e.g. [10,55,56]; AKH: [57,58]; PKG: [53];

PKA: [54]. An in-depth investigation of the development of gregarious-like states in solitary

locusts should prove to be very enlightening.

A central question is whether a collective (herd, flock, or swarm) is merely a sum of

its parts, or a new entity. Most related studies have perceived collectivity as a self-emergent

phenomenon, suggesting that new dynamics and behavior are the result of intricate, multi-

body, typically non-linear interactions (e.g. [20,59]). One hidden assumption underlying this

perception, is that individuals remains inherently unchanged when isolated or in a crowd.

Even studies of heterogeneous swarms, in which conspecifics may differ from each other,

still assume consistency in the properties of the individual over time. This is essentially a

physical point of view, in the sense that agents/individuals possess certain properties that

determine their behavior across a range of situations. Thus, the collective motion is an

emergent property that builds up in particular contexts, such as a sufficiently high local

density of animals. This point of view allows, among others, extrapolation from experiments

with one, two, or a few animals to large swarms (e.g. [30]).

Our findings reported here suggest a fundamentally different point of view. We

perceive the sensory-motor act of collective motion as accompanied by an internal state – a

collective-motion-state that is manifested in specific behavioral kinematics. This state is

induced by the experience of synchronous, collective motion. Most importantly, it is *not*

induced by spatial aggregation alone. Collectivity, therefore, is not just self-emerging.

Rather, the collective-motion-state has an important role in maintaining the integrity and

consistency of the swarm. The robustness of the swarm is also a major challenge and

requirement in swarming robotics, making the current novel insights applicable and even

important also to this emerging field.

In the case of locusts, our far from complete understanding to date of this

phenomenon is also proving crucial for human well-being and even survival, as evident from

the current devastating locust situation in large parts of Africa and Asia [60]. Much scientific

attention has been dedicated to the perception, decision-making, and individual kinematics

of locusts in a swarm. These efforts have led to various models that attempt to explain the

collective behavior on the basis of local interactions among the individual locusts (see [7] for

review). The current study is, to the best of our knowledge, the first to provide a more in-

depth focus, by including the internal state of the individual locust as an important factor in

dictating its behavior, and in turn affecting the maintenance and the properties of the

swarm.

Methods

Animals

The empirical experiments were conducted on gregarious desert locusts,

Schistocerca gregaria (Forsskål), which are bred at Tel Aviv University, School of Zoology.

The animals are kept at high density: over 100 animals in cages of 60 l, under a 12 h:12 h

light/dark regime, a temperature of 30℃, and 30-65 % humidity. Animals are fed daily with

wheat seedlings and dry oats. All experimental locusts were the offspring of many

generations of gregarious locusts reared in these conditions.

Experimental setup

The ring-shaped arena comprised blue plastic walls (60 cm diameter and 55 cm high)

with an inner central cone dome (30 cm diameter). The bottom of the structure was

covered with a thin layer of Fluon (Whitford Plastics Ltd., Runcorn, UK), in order to prevent

locusts from climbing. A white paper covered the floor of the arena and was changed before

each experiment. Light sources around the arena lit its floor evenly. A video camera (Sony

FDR-AXP35: 4K Ultra HD) was positioned above the arena and filmed the locusts throughout

the experiment.

Monitoring the locust movements

Prior to the experiments, each of the experimented locusts was tagged with a

unique barcode glued to the dorsal side of its prothorax using a drop of Epoxy glue. Using

the video footage at 25/3 frames per second, the trajectory of each animal was

reconstructed offline using BugTag software (Robiotec Ltd., Israel). To this we added a

custom-designed multiple-target tracking and a trajectory-smoothing method (for details:

Ariel et al., 2014b). briefly, segments in which tags were not identified (less than 5 cm or 25

s) by the system positions were interpolated. We thus obtained the positions of the tags'

center of mass relative to the arena center, at a final resolution of 2-3 pixels (ca. 0.5 mm),

for about 99% of the video frames. In order to reach 100% recognition, we completed the

analysis manually for the remaining frames. In three out of the 26 experiment the tracking

system failed to recognise 1-2 animals in the grouping stage. Therefore, the order

parameter was calculated based on the recognized animals only. These miss-identifications,

which only slightly affect the order parameter calculated in the grouping stage, have no

influence on any of the kinematic statistics in any of the stages (that only describe the focal

animal).

Experimental procedure

In the morning of each experiment day, 10 locusts were taken from their cage and

individually tagged. Thereafter, one locust was randomly chosen and placed in the arena

alone for one hour. The other 9 locusts were then added for an additional hour; after which

they were removed. The remaining locust was then filmed for another hour.

Analyzed parameters

Following the recognition process, we analyzed the middle 40 min of each of the 3 hours

of the experiment using MATLAB (MathWorks, Natick, MA, USA). The following parameters

were calculated:

1. Order parameter – the absolute value of the average direction of movement of the

walking animals (clockwise denoted as -1 and anticlockwise as +1), averaged for all

the frames.

2. Fraction of walking – the number of frames an animal walked divided by the total

number of analyzed frames, averaged over all experimental animals.

3. Detection of walking bouts – a walking period is defined as walking at a speed above

0.25 cm/sec for more than one third of a second.

4. Speed – averaged for all experimental animals and analyzed frames when the animal

walked and the speed differed from zero.

5. Walking bout duration – the time an animal moved continuously between one pause

to the next, averaged for all walking bouts and experimental animals.

6. Pause duration – the time an animal did not walk between one walking bout and the

next, averaged for all pauses and experimental animals.

Statistical analysis

All statistical tests were conducted with MATLAB. To compare the order parameters,

Wilcoxon rank-sum test was used. To compare among the three stages, Friedman Test was

used. Since all comparisons were significant according to the latter test, Bonferroni multiple

comparison post-hoc test accompanied each Friedman Test, in order to reveal the

significantly different groups. The p values noted are of the Bonferroni multiple comparison

post-hoc test. Correlations were conducted using Spearman's Rank Correlation test.

Modeling

We employed a two dimensional model comprising N agents moving at a fixed

speed v in a square domain of linear size L and periodic boundaries. Position and heading

(direction) of each rectangle center are denoted by $x_i(t)$ and $\hat{v}_i(t)$, respectively, where

$x_i(t) \in [0, L]^2$ and $\|\hat{v}_i(t)\| = 1$. In addition, $w_i(t)$ is a Boolean variable that denotes

whether at time t, agent i is moving ($w_i(t) = 1$) or pausing ($w_i(t) = 0$). In order to take into

account the physical size of animals, we assume that each agent is a rectangle with sides

$a_1 \times a_2$.

In each simulation step, the position of moving agents changes by amount v in the direction

$\hat{v}_i(t)$,

$x_i(t + 1) = x_i(t) + vw_i(t)\hat{v}_i(t) \pmod{L}.$

The new velocity of agent i depends on three terms: i. Inertia (its own direction); ii. the

direction of other agents in its receptive field (see below), possibly with a memory; and iii. a

random vector with unit norm $\xi_i(t)$. The receptive field view is a circle around $x_i(t)$. Unlike

previous models, we assume that an agent sees all other agents whose rectangle is within a

distance r from $x_i(t)$, and not merely the center of the rectangle. Denoting the four corners

of the rectangle describing agent j as indexes $c_i^1(t), \dots, c_i^4(t)$, we define the set of

observable neighbors of agent i at time t as

$I_i(t) \coloneqq \{j \colon \left\| x_i(t) - c_j^k(t) \right\| \leq r \text{ for some } k = 1 \dots 4\},$

the average direction in the receptive field at time t, is $\hat{u}_i(t) = u_i(t)/\|u_i(t)\|$, where

502
$$u_i(t) = \left(\frac{1}{n_i(t)} \sum_{j \in I_i(t)} \sin(\hat{v}_j), \frac{1}{n_i(t)} \sum_{j \in I_i(t)} \cos(\hat{v}_j) \right).$$

Here, $n_i(t)$ is the number of elements in $I_i(t)$.

Overall, the new heading $\hat{v}_i(t+1)$ is the weighted average of four terms: inertia,

direction of moving intersecting rectangles at time t, the average heading of moving

intersecting rectangles at time $[t - 5 \dots t - 1]$, and independent random vector with unit

norm $\xi_i(t)$. Let the set of the weights be $[q_1 \dots q_4]$. The updated weighted averaged

direction, is, therefore, $\hat{v}_i(t) = v_i(t)/\|v_i(t)\|$, where

511
$$v_i(t+1) = q_1 \hat{v}_i(t) + q_2 \hat{u}_i(t) + q_3 \frac{1}{5} \sum_{s=1}^{5} \hat{u}(t-s) + q_4 \xi_i(t)$$

In Fig. 3 and 4, the weights were set to $Q = [0.2, 0.3, 0.2, 0.3]$.

Switching between walking and pausing is determined as follows. Assuming

exponential distributions of moving and pausing durations, the probability to proceed

moving is given by a single parameter, p_w, describing the probability per-step of an agent to

continue walking, $p_w = P(w_i(t+1) = 1|w_i(t) = 1)$. Similarly, $p_s = P(w_i(t+1) =$

$0|w_i(t) = 0)$ is the probability per-step that a stopping agent will continue stopping.

In the simulations which introduce adaptive p_w, the latter was updated each time "walking"

started, and ranged between 0.4 and 0.8, as a linear function of the average number of

neighbors , $\bar{n}_i(t)$ in the last k. steps.

The following parameters were calculated over 200-1000 stimulation steps. Brackets

denote averaging over all analyzed frames.

1. Order: $\theta(t) = \frac{1}{N}\langle\|\sum_{i=1}^{N}\hat{v}_i\|\rangle$.

2. Spread: $S(t) = \frac{1}{N(N-1)}\langle\sum_{i=1}^{N}\sum_{j\neq i}|x_i(t) - x_j(t)|\rangle$

3. Number of neighbors: $n_i(t) = \frac{1}{N}\langle\sum_{i=1}^{N}I_i(t)\rangle$

4. Steps to regroup: the number of steps that passed from the first time an agent is

527   alone $(n_i(t) = 0)$ until it has a first neighbor $(n_i(t+s) > 0)$.

Acknowledgements

530   This research has been supported by the Israel Science Foundation (research grant 2306/18)

Author contributions

D.K., G.A., and A.A. designed the study. D.K. and C.S. performed the experiments.

D.K. and C.S. analyzed the data. D.K. N.A. and G.A constructed the model. D.K. wrote the

code, performed simulations and analyzed the data. D.K., G.A., and A.A. wrote the

manuscript. All authors reviewed and approved the paper.

Competing interests

The authors declare that they have no competing interests.

Data availability

The data is available upon request.

References

1. Sumpter, D. J. T. The principles of collective animal behaviour. *Philos. Trans. R. Soc. B Biol. Sci.* **361**, 5–22 (2006).

2. Ward, A. & Webster, M. *Sociality: The Behaviour of Group-Living Animals*. (Springer International Publishing, 2016). doi:10.1007/978-3-319-28585-6

3. Be'er, A. & Ariel, G. A statistical physics view of swarming bacteria. *Mov. Ecol.* **7**, 1–17

(2019).

4.   Yang, W. C. & Schmickl, T. Collective Motion as an Ultimate Effect in Crowded Selfish

Herds. *Sci. Rep.* **9**, 1–11 (2019).

5.   Cavagna, A. *et al.* Scale-free correlations in starling flocks. *Proc. Natl. Acad. Sci. U. S.*

A. **107**, 11865–11870 (2010).

6.   Attanasi, A. *et al.* Collective Behaviour without Collective Order in Wild Swarms of

Midges. *PLoS Comput. Biol.* **10**, 1–10 (2014).

7.   Ariel, G. & Ayali, A. Locust Collective Motion and Its Modeling. *PLoS Comput. Biol.* **11**,

1–25 (2015).

8.   Algar, S. D., Stemler, T. & Small, M. From Flocs to Flocks. in *A Mathematical Modeling*

Approach from Nonlinear Dynamics to Complex Systems (ed. E., M.) 157–175

(Springer, 2019). doi:10.1007/978-3-319-78512-7_9

9.   Escaff, D., Toral, R., Van Den Broeck, C. & Lindenberg, K. A continuous-time persistent

random walk model for flocking. *Chaos* **28**, (2018).

10.   Cullen, D. A. *et al. From Molecules to Management: Mechanisms and Consequences*

of Locust Phase Polyphenism. Advances in Insect Physiology **53**, (Elsevier Ltd., 2017).

11.   Ayali, A. The puzzle of locust density-dependent phase polyphenism. *Curr. Opin.*

Insect Sci. **35**, 41–47 (2019).

12.   Zhang, L., Lecoq, M., Latchininsky, A. & Hunter, D. Locust and Grasshopper

Management. *Annu. Rev. Entomol.* **64**, 15–34 (2019).

13.   Ellis, P. E. & Ashall, C. Field Studies on diurnal Behaviour, Movement and Aggregation

in the Desert Locust (Schistocerca gregaria Forskål). *Anti-Locust Bull.* **25**, 1–103

(1957).

14.   Uvarov, B. *Grasshoppers and locusts: A handbook of general acridology*. (Centre for

574   Overseas Pest Research, 1977).

575   15.   Knebel, D., Ayali, A., Guershon, M. & Ariel, G. Intra- versus intergroup variance in

576   collective behavior. *Sci. Adv.* **5**, eaav0695 (2019).

577   16.   Ariel, G., Ophir, Y., Levi, S., Ben-Jacob, E. & Ayali, A. Individual pause-and-go motion is

578   instrumental to the formation and maintenance of swarms of marching locust

579   nymphs. *PLoS One* **9**, e101636 (2014).

580   17.   Bazazi, S., Bartumeus, F., Hale, J. J. & Couzin, I. D. Intermittent motion in desert

581   locusts: behavioural complexity in simple environments. *PLoS Comput. Biol.* **8**,

582   e1002498 (2012).

583   18.   Bazazi, S. *et al.* Collective Motion and Cannibalism in Locust Migratory Bands. *Curr.*

584   *Biol.* **18**, 735–739 (2008).

585   19.   Rimer, O. & Ariel, G. Kinetic order-disorder transitions in a pause-and-go swarming

586   model with memory. *J. Theor. Biol.* **419**, 90–99 (2017).

587   20.   Vicsek, T. & Zafeiris, A. Collective motion. *Phys. Rep.* **517**, 71–140 (2012).

588   21.   Shklarsh, A., Ariel, G., Schneidman, E. & Ben-Jacob, E. Smart swarms of bacteria-

589   inspired agents with performance adaptable interactions. *PLoS Comput. Biol.* **7**,

590   (2011).

591   22.   Shaukat, M. & Chitre, M. Adaptive behaviors in multi-agent source localization using

592   passive sensing. *Adapt. Behav.* **24**, 446–463 (2016).

593   23.   Gorbonos, D. & Gov, N. S. Stable swarming using adaptive long-range interactions.

594   *Phys. Rev. E* **95**, 1–10 (2017).

24.   Benisty, S., Ben-Jacob, E., Ariel, G. & Be'er, A. Antibiotic-induced anomalous statistics

of collective bacterial swarming. *Phys. Rev. Lett.* **114**, 1–5 (2015).

25.   Fürtbauer, I. & Fry, A. Social conformity in solitary crabs, Carcinus maenas, is driven

by individual differences in behavioural plasticity. *Anim. Behav.* **135**, 131–137 (2018).

26.   Planas-Sitjà, I., Deneubourg, J.-L., Gibon, C. & Sempo, G. Group personality during

collective decision-making: a multi-level approach. *Proceedings. Biol. Sci.* **282**,

20142515 (2015).

27.   Brown, C. & Irving, E. Individual personality traits influence group exploration in a

feral guppy population. *Behav. Ecol.* **25**, 95–101 (2014).

28.   Dyer, J. R. G., Croft, D. P., Morrell, L. J. & Krause, J. Shoal composition determines

foraging success in the guppy. *Behav. Ecol.* **20**, 165–171 (2009).

29.   Farine, D. R., Strandburg-Peshkin, A., Couzin, I. D., Berger-Wolf, T. Y. & Crofoot, M. C.

Individual variation in local interaction rules can explain emergent patterns of spatial

organisation in wild baboons. *Proc. R. Soc. B Biol. Sci.* **284**, 20162243 (2017).

30.   Calovi, D. S. *et al.* Collective response to perturbations in a data-driven fish school

model. *J. R. Soc. Interface* **12**, (2015).

31.   Herbert-Read, J. E. *et al.* The role of individuality in collective group movement. *Proc.*

R. Soc. B Biol. Sci. **280**, (2013).

32.   Jolles, J. W., Laskowski, K. L., Boogert, N. J. & Manica, A. Repeatable group differences

in the collective behaviour of stickleback shoals across ecological contexts. *Proc. R.*

Soc. B Biol. Sci. **285**, 13–16 (2018).

33.   Crall, J. D. *et al.* Social context modulates idiosyncrasy of behaviour in the gregarious

cockroach Blaberus discoidalis. *Anim. Behav.* **111**, 297–305 (2016).

34.   Aplin, L. M., Farine, D. R., Mann, R. P. & Sheldon, B. C. Individual-level personality

619   influences social foraging and collective behaviour in wild birds. *Proceedings. Biol. Sci.*

620   **281**, 20141016 (2014).

621   35.   Ariel, G., Rimer, O. & Ben-Jacob, E. Order–Disorder Phase Transition in

622   Heterogeneous Populations of Self-propelled Particles. *J. Stat. Phys.* **158**, 579–588

623   (2014).

624   36.   Copenhagen, K., Quint, D. A. & Gopinathan, A. Self-organized sorting limits behavioral

625   variability in swarms. *Sci. Rep.* **6**, 1–11 (2016).

626   37.   Guisandez, L., Baglietto, G. & Rozenfeld, A. Heterogeneity promotes first to second

627   order phase transition on flocking systems. 1–11 (2017).

628   38.   Jolles, J. W., Boogert, N. J., Sridhar, V. H., Couzin, I. D. & Manica, A. Consistent

629   Individual Differences Drive Collective Behavior and Group Functioning of Schooling

630   Fish. *Curr. Biol.* (2017). doi:10.1016/j.cub.2017.08.004

631   39.   Menzel, A. M. Collective motion of binary self-propelled particle mixtures. *Phys. Rev.*

632   *E - Stat. Nonlinear, Soft Matter Phys.* **85**, 31–33 (2012).

633   40.   Mishra, S., Tunstrøm, K., Couzin, I. D. & Huepe, C. Collective dynamics of self-

634   propelled particles with variable speed. *Phys. Rev. E* **86**, 011901 (2012).

635   41.   Mar Delgado, M. del *et al.* The importance of individual variation in the dynamics of

636   animal collective movements. *Philos. Trans. R. Soc. B Biol. Sci.* **373**, (2018).

637   42.   Modlmeier, A. P., Keiser, C. N., Wright, C. M., Lichtenstein, J. L. L. & Pruitt, J. N.

638   Integrating animal personality into insect population and community ecology. *Curr.*

639   *Opin. Insect Sci.* **9**, 77–85 (2015).

640   43.   Webster, M. M. & Ward, A. J. W. Personality and social context. *Biol. Rev.* **86**, 759–

641   773 (2011).

44.   Szorkovszky, A. *et al.* Assortative interactions revealed by sorting of animal groups.

Anim. Behav. **142**, 165–179 (2018).

45.   O'shea-Wheller, T. A., Masuda, N., Sendova-Franks, A. B. & Franks, N. R. Variability in

individual assessment behaviour and its implications for collective decision-making.

Proc. R. Soc. B Biol. Sci. **284**, 1–7 (2017).

46.   Niemelä, P. T. & Santostefano, F. Social carry-over effects on non-social behavioral

variation: Mechanisms and consequences. *Front. Ecol. Evol.* **3**, 1–12 (2015).

47.   Howard, C. E. *et al.* Serotonergic Modulation of Walking in Drosophila. *Curr. Biol.* **29**,

4218-4230.e8 (2019).

48.   Perić-Mataruga, V., Nenadović, V. & Ivanović, J. Neurohormones in insect stress: A

review. *Arch. Biol. Sci.* **58**, 1–12 (2006).

49.   Wicher, D. Metabolic Regulation and Behavior: How Hunger Produces Arousal - An

Insect Study. *Endocrine, Metab. Immune Disord. Targets* **7**, 304–310 (2007).

50.   Verlinden, H. *et al.* The role of octopamine in locusts and other arthropods. *J. Insect*

Physiol. **56**, 854–867 (2010).

51.   Yang, Z. *et al.* Octopamine mediates starvation-induced hyperactivity in adult

Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 5219–5224 (2015).

52.   Geva, N., Guershon, M., Orlova, M. & Ayali, A. Memoirs of a locust: Density-

dependent behavioral change as a model for learning and memory. *Neurobiol. Learn.*

Mem. **93**, 175–182 (2010).

53.   Lucas, C. *et al.* The locust foraging gene. *Arch. Insect Biochem. Physiol.* **74**, 52–66

(2010).

54.   Ott, S. R. *et al.* Critical role for protein kinase A in the acquisition of gregarious

665  behavior in the desert locust. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 381–387 (2012).

666 55. Alessi, A. M., O'Connor, V., Aonuma, H. & Newland, P. L. Dopaminergic modulation of

667  phase reversal in desert locusts. *Front. Behav. Neurosci.* **8**, 1–15 (2014).

668 56. Ma, Z. *et al.* Octopamine and tyramine respectively regulate attractive and repulsive

669  behavior in locust phase changes. *Sci. Rep.* **5**, 1–11 (2015).

670 57. Ayali, A. & Pener, M. P. Density-dependent phase polymorphism affects response to

671  adipokinetic hormone in Locusta. *Comp. Biochem. Physiol. -- Part A Physiol.* **101**, 549–

672  552 (1992).

673 58. Pener, M. P., Ayali, A. & Golenser, E. Adipokinetic hormone and flight fuel related

674  characteristics of density-dependent locust phase polymorphism: A review. *Comp.*

675  *Biochem. Physiol. - B Biochem. Mol. Biol.* **117**, 513–524 (1997).

676 59. Cucker, F. & Smale, S. Emergent Behavior in Flocks. *IEEE Trans. Automat. Contr.* **52**,

677  852–862 (2007).

678 60. FAO. Desert Locust situation update - 21 April 2020. *Food and Agriculture*

679  *Organization of the United Nations* (2020).

680  doi:http://www.fao.org/ag/locusts/en/info/info/index.html

683   **Tables**

	p-Value	*rho*-Value
Fraction of Walking		
Isolation - Grouping	0.000	0.706
Isolation - Re-isolation	0.001	0.679
Grouping - Re-isolation	0.002	0.638
Speed		
Isolation - Grouping	0.001	0.679
Isolation - Re-isolation	0.010	0.561
Grouping - Re-isolation	0.002	0.636
Walking bout duration		
Isolation - Grouping	*0.002*	0.628
Isolation - Re-isolation	*0.148*	0.391
Grouping - Re-isolation	*0.354*	0.314
Puase duration		
Isolation - Grouping	0.007	0.580
Isolation - Re-isolation	0.003	0.612
Grouping - Re-isolation	0.000	0.708

685   **Table 1. Correlation values between kinematic parameters throughout the three experimental conditions.** The

686   fraction of walking, the averaged walking speed, the average duration of walking bouts, and the average

687   duration of pauses were tested for correlation across the three experimental conditions: (1) isolation, (2)

688   grouping, and (3) re-isolation. The underlined values mark significant correlations.