Dopaminergic Neurons Regulates Aging and Longevity in Flies

Xiaolin Tian[1]*

[1]Neuroscience Center of Excellence, Department of Cell Biology and Anatomy, Louisiana State University Health Sciences Center, New Orleans, LA 70112, USA

*Corresponding author

LSUHSC-NO, Neuroscience Center of Excellence

2020 Gravier St. STE.D

New Orleans, LA 70112

Phone: 504-599-0912

Email address: xtian@lsuhsc.edu

Abstract

Dopaminergic neurons (DANs) modulate essential brain functions involving memory formation, reward processing, and decision making. Here, I demonstrate a novel and important function of the DANs in the fly brain in regulating aging and longevity. I show that two distinct groups of DANs, when overexpressing the putative scaffolding protein Mask, can each significantly extend lifespan. Dopamine transmission is required to elicit beneficial effects when Mask is overexpressed in these neurons, but elevating dopamine production or supplementation of L-dopa is not sufficient to yield similar effects on longevity. The lifespan extension induced by Mask-overexpressing in the long-lived flies is accompanied by sustained adult locomotor and fecundity. And other physiological functions in the adult such as food intake, energy storage, and insulin production in the brain are not consistently altered by overexpressing Mask in either group of DANs. Our recent work demonstrated a novel cellular function of Mask in promoting microtubule (MT) dynamics. Overexpressing Unc-104 or moderately reducing p150[Glued] level, interventions that can induce MTs dynamics, in the same groups of DANs also extend lifespan in flies. All together, these data unraveled an undocumented DANs-dependent mechanism that regulate systemic aging and longevity in flies, alterations of the cellular properties in subsets of DANs can have a profound impact on aging.

Dopaminergic neurons, the central component of a key modulatory system in the brain, are greatly affected by age. Many essential brain functions involving memory formation, motor control, reward processing, arousal, metabolism, and decision making are modulated by the dopaminergic system[1,2], but it is unclear whether it also regulates the aging process in animals. In the course of studying a putative scaffolding protein Mask, I uncovered a novel role of dopaminergic neurons in regulating longevity and aging in fruit flies. I found that overexpressing Mask in dopaminergic neurons leads to a ~40% -increase in lifespan in flies (median lifespan in male flies increased from ~76 days to ~104 days) (Fig. S1). This effect seems to be specific to the dopaminergic neurons, as overexpressing Mask is neither the entire body nor in the whole nervous system (neurons or glial cells) showed significant effects on the lifespan of flies (Fig. S2). Dopaminergic neurons in the adult fly brain cluster into eight groups[3]. I next asked whether all or only subsets of dopaminergic neurons are responsible for their lifespan-control function. I express Mask in distinct groups of dopaminergic neurons in the fly brains using three well-characterized Gal4 drivers[4-6], and I found that expressing Mask in the PAM dopaminergic neurons by the 0273-Gal4 driver does not alter the lifespan (Fig. 1A). However, expressing Mask driven by either TH-C'- or TH-D'-Gal4 significantly extends the lifespan in flies — the median lifespan in male flies increased from 64 days to 96 days, or from 63 days to 86 days, respectively (Fig. 1 B&C).

Although the dopaminergic system provides essential modulation on various behaviors and physiological functions, flies devoid of dopamine in their brains[7] and worms lacking the rate-limiting enzymes for dopamine synthesis live a normal lifespan[8]. Consistently, blocking neuronal activity in the TH-C' or TH-D' dopaminergic neurons by the DREADD-Di[9] does not affect the lifespan in flies (Fig. S3). These results together suggest that, although dopamine systems may

not be required for a normal lifespan, overexpressing Mask in these dopaminergic neurons possibly induces a gain-of-function effect in the dopaminergic neurons which consequentially confers a beneficial outcome on aging and longevity. Before I can fully understand the underlying molecular mechanism for this phenomenon, I first set out to confirm the potentials of these dopaminergic neurons to impact lifespan in flies. Using established interventions that affect neuronal activities, I examined the effects on longevity by fully activating the TH-C' or TH-D' dopaminergic neurons in the adult flies. Next, I found that continuous activation of the TH-C' or TH-D' dopaminergic neurons using the thermosensitive cation channel TrpA1 shortens the lifespan in the male flies, while the lifespan in the female flies was not affected (Fig. S4). After, I employed the inducible activation system, DREADD-D$_s$[9], in the TH-C' or TH-D' dopaminergic neurons to quantitatively modulate their neuronal activity instead of fully activating these neurons. I found that the 1μM CNO treatment (only induces very modest activation of the DREADD-D$_s$ receptors) induces prominent and consistent extension of lifespan in flies that express the engineered Gαs receptor in their TH-C' or TH-D' neurons (Fig. 2), suggesting that moderate modulation of these neurons is sufficient to impact longevity.

Next, I asked whether dopamine transmission is required for lifespan extension induced by Mask-overexpression in the dopaminergic neurons. I reasoned that if dopamine transmission is needed, reducing dopamine signaling in these neurons would suppress the ability of Mask-overexpression to extend lifespan. Indeed, genetically heterozygosity that reduces gene dosages of either of the four genes that are essential for dopamine transmission: two enzymes responsible for dopamine synthesis - Tyrosine Hydroxylase (TH) and Dopa Decarboxylase (Ddc); and two transporters for dopamine - dopamine transporter (DAT) that controls dopamine reuptake, and vesicular monoamine transporter (VMAT) that mediates synaptic vesicular loading of dopamine,

can each blunt the extend of lifespan extension when Mask is overexpressed in either TH-C' or

TH-D' neurons in the heterozygous backgrounds of all four genes compared to wild type

background (Fig. S5). These results further support the notion that the dopamine system plays an

essential role in Mask-mediated lifespan extension. I then tested whether upregulation of

dopamine signaling is sufficient to induce lifespan extension in flies. It appeared that feeding

flies with dopamine precursor L-DOPA[10], dopamine D1 receptor agonist SKF-82958[11],

dopamine D2 receptor agonist Quinpirole[12], or D1 and D2 agonist together all fail to extend the

lifespan in flies (Fig. S6). The lifespan of female flies fed with L-DOPA is even moderately

shortened (Fig. S6). Since the adverse effects may be attributable to global elevation of

dopaminergic signaling which has been shown to cause neural toxicity in mammals[13], I next used

genetic approaches to selectively increase dopamine production in the TH-C' or TH-D' neurons

via overexpressing Tyrosine Hydroxylase (TH), or Dopa Decarboxylase (Ddc), two enzymes that

are responsible for dopamine synthesis, in the TH-C' or TH-D' dopaminergic neurons. I found

that overexpressing these two enzymes did not affect longevity (Fig. S7). Similar to L-DOPA

feeding, these treatment lead to moderate reduction in the lifespan in female flies. These results

suggest that although neurons require dopamine transmission to communicate and to induce

lifespan extension, increasing dopamine production alone in these dopaminergic neurons is not

sufficient to elicit similar effects on aging and longevity induced by Mask overexpression or

moderate DREADD-Ds activation in the same neurons.

To fully understand how overexpressing Mask in TH-C' or TH-D' neurons impact aging and

longevity at the systematic level, I next examined age-related phenotypes including feeding,

energy storage/metabolic, adult locomotor and brain insulin productions in Mask-expressing and

control flies. Dopaminergic neurons regulate feeding and food-foraging in flies[14-17], raising the

possibility that the long lived flies that overexpressing Mask in the TH-C' or TH-D' neurons may consume less food and therefore undertake voluntary caloric restriction, a potential contributing factor to their prolonged lifespan. However, the Mask-overexpressing long-lived flies consume comparable amount of food compared to the control flies at young and mid age (10 and 30 day old) in the Capillary Feeder assay (CAFE assay)[18] (Fig. S8). These flies also show moderately increased levels of whole body glucose and Triglyceride (TAG) levels at mid and late ages (30 and 50 days old) (Fig. S9). I next assessed adult locomotor activity in flies using the rapid iterative negative geotaxis (RING) assay[19,20], and I found that expressing Mask in TH-C' dopaminergic neurons significantly enhances the locomotor performance during the entire lifespan (Fig. S10). To determine whether there is a correlation between the prolonged lifespan and potential changes in insulin production in the brain, I next measured transcription levels of Dilp2, 3, 5 and 6 in the adult brains of Mask-overexpression and control flies. I did not detect any consistent reduction in transcription levels of the Dilp genes between gender or age (Fig. S11- 14). All together, the results of the aging analysis suggests that the Mask-overexpressing flies live healthy and active lives throughout the entire lifespan.

Delayed reproductive maturation or reduced fecundity correlate closely with lifespan extension[21]. To test whether there is a tradeoff between fertility and lifespan in Mask-overexpressing flies, I examined the fecundity of male and female flies at different ages. I did not detect any delay or reduction in reproduction in either male or female flies. Rather, overexpressing Mask in the TH-C' neurons increased fertility in female flies beginning right after eclosion, and the TH-D' overexpressing flies showed a significantly higher level of fecundity at older ages (Fig. 3). While previous studies reveal a trade-off between lifespan and fertility, the Mask-overexpressing flies, as well as a few other previously reported mutant flies[22-]

[25] show a trait of extended lifespan at no cost of reproduction. Given the lack of consensus theory to explain the relationship between aging and reproduction, the phenomenon that increased fecundity can be a concomitant of lifespan extension calls for an alternative theory that can provide a unifying explanation for the intricate relationship between aging and reproduction.

Since overexpressing Mask in the dopaminergic neurons very likely induces a gain-of-function effects that leads to lifespan extension, I next investigated the primary cellular effector in the Mask-overexpressing dopaminergic neurons that mediate the outcome on aging and longevity. We recently demonstrated that Mask promotes microtubule (MT) dynamics in fly larval motor neurons and body wall muscles[26]. This function of Mask prompted me to postulate that altered MT dynamics in the Mask-expressing dopaminergic neurons is the key effector that induces changes in neuronal behaviors leading to lifespan extensions in flies. If this is true, reagents that manipulate MT dynamics in the same manner would also induce lifespan extension when applied to the same subsets of dopaminergic neurons. I chose two interventions that have been previously shown to impact MT dynamics: overexpressing the Kinesin heavy chain Unc-104[27] or knocking down a component of the Dynein/Dynactin complex, p150[Glued28], has each been shown to reduce MT length and stability and enhance MT-based transportations. When we applied these manipulations to the TH-C' or TH-D' dopaminergic neurons, both treatments extended lifespan in flies (Fig. 4). These results strongly argue that increasing MT dynamics and reducing MT stability in the TH-C' or TH-D' dopaminergic neurons is sufficient to induce lifespan extension in flies.

I next asked whether other cellular functions of Mask also contribute to its ability to induce this dopaminergic-dependent lifespan extension. Mask is a large protein (4001 amino acids) containing several conserved functional domains (Fig. S15), and the structures of Mask enables

its versatile functions in the cells: such as its ability to promote microtubule (MT) dynamics[26], autophagy[29], HIPPO pathway[30,31] or vesiculation[29,32]. Mask is a KH-domain-containing protein. KH domains bind RNA or single-stranded DNA and regulate transcription, translation, mRNA stability and alternative splicing[33-35]. Mutating the GXXG loop to GDDG in the KH minimal motif greatly reduces the ability of the KH domain to bind RNAs[36]. The GXXG loop of Mask resides in amino acids 3053-3056 as GRGG, which is fully conserved between fly and human. To assess whether the KH domain is required for normal functions of Mask, I generated a UAS-Mask transgene that carries a GRGG to GDDG mutation in its KH domain (named Mask-KH-Mutant) (depicted in Fig. S15A). Our previous work demonstrated that overexpressing Mask promotes autophagic degradation[29], and an intact KH domain is required for Mask to enhance autophagic degradation because overexpression of Mask-KH-Mutant fail to induce this effect (Fig. S16). However, the KH-Mutant transgene can rescue the MT defects in larval muscle[26], and more importantly, expression of Mask-KH-Mutant in the TH-C' dopaminergic neurons extended lifespan to a similar level as expression of the wild type Mask protein. (Fig. S15 B&C). The fact that KH domain is dispensable for Mask to induce lifespan extension indicates that autophagy may not be a primary cellular effector in Mask-mediated lifespan-extension. Also, overexpressing ATG-1, a potent inducer of autophagy, in all neurons in the adult flies can extend lifespan[37], however, overexpressing Mask in all neurons in the adult flies failed to induce similar lifespan extension (Fig. S17). These data together suggest that autophagy is unlikely a primary cellular effector in Mask-mediated lifespan-extension.

The Ankyrin repeats in ANKHD1, the mammalian homolog of Mask, regulates membrane vesiculation[32], suggesting a possible role of its fly counterpart Mask to promote vesicle formation and recycling at the nerve terminals. Indeed, overexpressing Mask in the fly larval neuromuscular junction (NMJ) increases the size of Rab5-positive vesicles pool at the presynaptic terminal (Fig. S18) and promotes vesicular recycling (measured by FM-143 dye

loading assay, Fig. S19). Expressing wild type or a constitutively active form of Rab5 can both enhance synaptic vesicle fusion efficacy and firing strength at the presynaptic terminals[38]. But, when expressed in the TH-C' or TH-D' dopaminergic neurons, the active form of Rab5 protein can mildly extend lifespan in the male flies (Fig. S20). It appears that although Mask can enhance endocytosis and vesicle recycling at the nerve terminal, promoting these events alone is only able to induce at much less extend as Mask the cellular output by dopaminergic neurons that translates to lifespan extension. Although there is still no answer for whether the function of Mask to promote MT dynamics and to enhance endocytosis at the nerve terminals are consequentially related. It is possible that altered vesicle trafficking is one of the downstream effectors that mediate Mask-depend lifespan extension.

In this study I report a novel role of dopaminergic neurons in regulating aging and longevity, and I show that enhancing the MT dynamics in subsets of dopaminergic neurons profoundly prolonged lifespan in flies. Interestingly, the lifespan extension is accompanied by the sustained fecundity in the long-lived flies, a phenotype that contradicts the disposable soma theory for aging which states an inversely related energy allocation between reproduction and somatic maintenance. To reconcile such a paradox, I propose an explanation that could potentially provide a consensus theory to aging and reproduction. This theory states a reproduction-center mechanism for aging, and that there is no competition for resource between somatic maintenance and reproduction, but rather, somatic tissues adapt and cop with the animal's need for reproduction. In detail, reproduction is centered by other physiological functions, and, in a "live fast and die young" scenario, the fulfilled reproduction in early life lifts the need for somatic maintenance and the animal die early. In the nutrition-sensing scenario, when the animals sense a lack a nutrition in the environment that poses threatens for the survival of the progenies, they halt

the reproductive activities, and adjust the somatic system accordingly to survive and prolong their lifespan to reproduce later. In social insects, reproductive females undergo somatic transformation to live much longer to meet the prolonged reproduction demand. This theory roots in the notion that natural selection centers on fitness on reproduction in a way that the animals possess the ability to adapt to different environment changes and challenges to ensure successful reproduction and the survival of the species. This reproduction-centered aging theory, posit that the organism would set an internal status, or a goal for their reproduction capacity; and to fulfill this goal the body not only engage mechanisms to monitor the progress of the fulfillment of it but also possess a system that can efficiently support somatic maintenance to suit the reproduction need. The brain, with the dopaminergic systems being an essential part for information evaluation and decision making, is the best candidate to centrally organize the information on internal status as well as the environmental cues, and to relay out to the peripheral the decision of adjusting the soma to match the reproduction status. The identification of the dopaminergic system as a regulator for aging regulation provides an entry point to further explore the molecular and cellular mechanisms underpin this new reproduction-centered aging theory.

Method and Material

Drosophila strains and genetics

Flies were maintained at 25°C on standard food containing agar (0.75% w/v), cornmeal (5% w/v), molasses (4.15% v/v), and yeast (1.575% w/v) at 25 °C in a 12:12 light:dark cycle. The following strains were used in this study: UAS-Mask[39], TH-C'-Gal4[6], TH-D'-Gal4[6] (from Mark Wu), 0273-Gal4 (from Thomas R. Clandinin), UAS-DREADD-D$_s$[9] (from Charles Nichols), Ddc-

Gal4 (Bloomington Stock #7010); UAS-unc104-GFP (Bloomington Stock #24787); UAS-p150 Glued RNAi (Bloomington Stock #24761). Control flies used in the study are w^{1118} from BestGene Inc. (Chino Hills, CA, USA) in which the UAS-Mask line was generated[39] and Iso[31] (from Mark Wu) in which TH-C'-Gal4 and TH-D'-Gal4 flies were generated[6].

Fly Longevity Assay

The procedures are adapted from previously described protocol[40]. 15-20 male or female flies were separated into one vial, and a total of 4-6 vials of flies were used for each genotype for longevity measurement. The flies are maintained at 25°C on standard food or food containing drugs and are transferred into fresh vials every 3 days.

DREADD Activation

10mM Clozapine N-oxide (CNO) (Tocris 4936) stock solution was prepared in DMSO. Food containing control (DMSO only) or 1uM CNO was prepared freshly every three days before the flies were transferred to new vials during the longevity recording experiments.

Fly Fecundity Assay

Newly eclosed individual virgin male or female flies (day1) were crossed with 2-3 w^{1118} virgin female or male flies respectively and kept together for 3 days. The total progeny produced within the 3-day intervals were counted after they eclose as adult flies. Sexually exposed and active male or female flies reared together with w^{1118} female or male, were aged till day 10, 30 and 50. Each female fly was then separated into a single vial and allow to lay eggs for 3 days. The total progeny produced within the 3-day intervals were counted after they eclose. The aged male flies were individually mated with three 5-7 day old w^{1118} virgin females for 3 days, and fecundity was measured as the percentage of males that can successfully produce progenies.

Statistical analysis

Data analysis for longevity: The Kaplan-Meier estimator was used to analyze the data and the survivorship curves were generated in Origin. The log-rank test was performed with Evan's A/B tools.

Each sample will be compared with other samples in the group (more than two) using ANOVA, or with the other sample in a group of two using a t-test. The graphs were generated in Origin (Origin Lab, Northampton, MA).

Acknowledgments

I greatly appreciate the help of Chunlai Wu for technical supports and critical comments on the manuscript. I am especially grateful to Dr. Mark Wu for providing many helpful reagents. I also thank Dr. Thomas R. Clandinin for fly stocks, as well as the Bloomington Stock Center for other fly stocks and Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242, for antibodies. I also thank Liz McGehee for editorial assistance. This work is supported by a Start-up fund from LSUHSC to X.T.

References

1. Schultz, W. Multiple dopamine functions at different time courses. *Annu Rev Neurosci* **30**, 259-288 (2007). PubMed PMID: 17600522
2. Arias-Carrión, O., Stamelou, M., Murillo-Rodríguez, E., Menéndez-González, M. & Pöppel, E. Dopaminergic reward system: a short integrative review. *Int Arch Med* **3**, 24 (2010). PubMed PMID: 20925949
3. Mao, Z. & Davis, R. L. Eight different types of dopaminergic neurons innervate the Drosophila mushroom body neuropil: anatomical and physiological heterogeneity. *Front Neural Circuits* **3**, 5 (2009). PubMed PMID: 19597562
4. Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., Gohl, D., Silies, M., Certel, S. & Waddell, S. Layered reward signalling through octopamine and dopamine in Drosophila. *Nature* **492**, 433-437 (2012). PubMed PMID: 23103875
5. Galili, D. S., Dylla, K. V., Ludke, A., Friedrich, A. B., Yamagata, N., Wong, J. Y., Ho, C. H., Szyszka, P. & Tanimoto, H. Converging circuits mediate temperature and shock aversive olfactory conditioning in Drosophila. *Curr Biol* **24**, 1712-1722 (2014). PubMed PMID: 25042591
6. Liu, Q., Liu, S., Kodama, L., Driscoll, M. R. & Wu, M. N. Two dopaminergic neurons signal to the dorsal fan-shaped body to promote wakefulness in Drosophila. *Curr Biol* **22**, 2114-2123 (2012). PubMed PMID: 23022067
7. Riemensperger, T., Isabel, G., Coulom, H., Neuser, K., Seugnet, L., Kume, K., Iche-Torres, M., Cassar, M., Strauss, R., Preat, T., Hirsh, J. & Birman, S. Behavioral consequences of dopamine deficiency in the Drosophila central nervous system. *Proc Natl Acad Sci U S A* **108**, 834-839 (2011). PubMed PMID: 21187381
8. Murakami, H. & Murakami, S. Serotonin receptors antagonistically modulate Caenorhabditis elegans longevity. *Aging Cell* **6**, 483-488 (2007). PubMed PMID: 17559503
9. Becnel, J., Johnson, O., Majeed, Z. R., Tran, V., Yu, B., Roth, B. L., Cooper, R. L., Kerut, E. K. & Nichols, C. D. DREADDs in Drosophila: a pharmacogenetic approach for controlling behavior, neuronal signaling, and physiology in the fly. *Cell Rep* **4**, 1049-1059 (2013). PubMed PMID: 24012754
10. Kayser, M. S., Yue, Z. & Sehgal, A. A critical period of sleep for development of courtship circuitry and behavior in Drosophila. *Science* **344**, 269-274 (2014). PubMed PMID: 24744368
11. Walters, M. R., Dutertre, M. & Smith, C. L. SKF-82958 is a subtype-selective estrogen receptor-alpha (ERalpha) agonist that induces functional interactions between ERalpha and AP-1. *J Biol Chem* **277**, 1669-1679 (2002). PubMed PMID: 11700319
12. Wiemerslage, L., Schultz, B. J., Ganguly, A. & Lee, D. Selective degeneration of dopaminergic neurons by MPP(+) and its rescue by D2 autoreceptors in Drosophila primary culture. *J Neurochem* **126**, 529-540 (2013). PubMed PMID: 23452092
13. Yamamoto, Branden J. Stansley & Bryan K. L-Dopa and Brain Serotonin System Dysfunction. *Toxics* **3**, 75-88 (2015). PubMed PMID:
14. Liu, Q., Tabuchi, M., Liu, S., Kodama, L., Horiuchi, W., Daniels, J., Chiu, L., Baldoni, D. & Wu, M. N. Branch-specific plasticity of a bifunctional dopamine circuit encodes protein hunger. *Science* **356**, 534-539 (2017). PubMed PMID: 28473588
15. Tsao, C. H., Chen, C. C., Lin, C. H., Yang, H. Y. & Lin, S. Drosophila mushroom bodies integrate hunger and satiety signals to control innate food-seeking behavior. *Elife* **7**(2018). PubMed PMID: 29547121
16. Landayan, D. & Wolf, F. W. Shared neurocircuitry underlying feeding and drugs of abuse in Drosophila. *Biomed J* **38**, 496-509 (2015). PubMed PMID: 27013449

17. Marella, S., Mann, K. & Scott, K. Dopaminergic modulation of sucrose acceptance behavior in Drosophila. *Neuron* **73**, 941-950 (2012). PubMed PMID: 22405204

18. Diegelmann, S., Jansen, A., Jois, S., Kastenholz, K., Velo Escarcena, L., Strudthoff, N. & Scholz, H. The CApillary FEeder Assay Measures Food Intake in Drosophila melanogaster. *J Vis Exp* (2017). PubMed PMID: 28362419

19. Nichols, C. D., Becnel, J. & Pandey, U. B. Methods to assay Drosophila behavior. *J Vis Exp* (2012). PubMed PMID: 22433384

20. Gargano, J. W., Martin, I., Bhandari, P. & Grotewiel, M. S. Rapid iterative negative geotaxis (RING): a new method for assessing age-related locomotor decline in Drosophila. *Exp Gerontol* **40**, 386-395 (2005). PubMed PMID: 15919590

21. Yuan, R., Meng, Q., Nautiyal, J., Flurkey, K., Tsaih, S. W., Krier, R., Parker, M. G., Harrison, D. E. & Paigen, B. Genetic coregulation of age of female sexual maturation and lifespan through circulating IGF1 among inbred mouse strains. *Proc Natl Acad Sci U S A* **109**, 8224-8229 (2012). PubMed PMID: 22566614

22. Simon, A. F., Shih, C., Mack, A. & Benzer, S. Steroid control of longevity in Drosophila melanogaster. *Science* **299**, 1407-1410 (2003). PubMed PMID: 12610309

23. Hwangbo, D. S., Gershman, B., Tu, M. P., Palmer, M. & Tatar, M. Drosophila dFOXO controls lifespan and regulates insulin signalling in brain and fat body. *Nature* **429**, 562-566 (2004). PubMed PMID: 15175753

24. Marden, J. H., Rogina, B., Montooth, K. L. & Helfand, S. L. Conditional tradeoffs between aging and organismal performance of Indy long-lived mutant flies. *Proc Natl Acad Sci U S A* **100**, 3369-3373 (2003). PubMed PMID: 12626742

25. Grandison, R. C., Piper, M. D. & Partridge, L. Amino-acid imbalance explains extension of lifespan by dietary restriction in Drosophila. *Nature* **462**, 1061-1064 (2009). PubMed PMID: 19956092

26. Zhu, Mingwei, Martinez, Daniel, Guidry, Jessie J., Majeste, Niles, Mao, Hui, Yanofsky, Sarah, Tian, Xiaolin & Wu, Chunlai. Mask, the Drosophila Ankyrin Repeat and KH domain-containing protein, regulates microtubule dynamics. *bioRxiv*, 2020.2004.2022.056051 (2020). PubMed PMID:

27. Randall, T. S., Yip, Y. Y., Wallock-Richards, D. J., Pfisterer, K., Sanger, A., Ficek, W., Steiner, R. A., Beavil, A. J., Parsons, M. & Dodding, M. P. A small-molecule activator of kinesin-1 drives remodeling of the microtubule network. *Proc Natl Acad Sci U S A* **114**, 13738-13743 (2017). PubMed PMID: 29229862

28. Lazarus, J. E., Moughamian, A. J., Tokito, M. K. & Holzbaur, E. L. Dynactin subunit p150(Glued) is a neuron-specific anti-catastrophe factor. *PLoS Biol* **11**, e1001611 (2013). PubMed PMID: 23874158

29. Zhu, M., Zhang, S., Tian, X. & Wu, C. Mask mitigates MAPT- and FUS-induced degeneration by enhancing autophagy through lysosomal acidification. *Autophagy* **13**, 1924-1938 (2017). PubMed PMID: 28806139

30. Sansores-Garcia, L., Atkins, M., Moya, I. M., Shahmoradgoli, M., Tao, C., Mills, G. B. & Halder, G. Mask is required for the activity of the Hippo pathway effector Yki/YAP. *Curr Biol* **23**, 229-235 (2013). PubMed PMID: 23333314

31. Sidor, C. M., Brain, R. & Thompson, B. J. Mask proteins are cofactors of Yorkie/YAP in the Hippo pathway. *Curr Biol* **23**, 223-228 (2013). PubMed PMID: 23333315

32. Kitamata, M., Hanawa-Suetsugu, K., Maruyama, K. & Suetsugu, S. Membrane-Deformation Ability of ANKHD1 Is Involved in the Early Endosome Enlargement. *iScience* **17**, 101-118 (2019). PubMed PMID: 31255983

33. Brooks, A. N., Duff, M. O., May, G., Yang, L., Bolisetty, M., Landolin, J., Wan, K., Sandler, J., Booth, B. W., Celniker, S. E., Graveley, B. R. & Brenner, S. E. Regulation of alternative splicing in

Drosophila by 56 RNA binding proteins. *Genome Res* **25**, 1771-1780 (2015). PubMed PMID: 26294686

34. Garcia-Mayoral, M. F., Hollingworth, D., Masino, L., Diaz-Moreno, I., Kelly, G., Gherzi, R., Chou, C. F., Chen, C. Y. & Ramos, A. The structure of the C-terminal KH domains of KSRP reveals a noncanonical motif important for mRNA degradation. *Structure* **15**, 485-498 (2007). PubMed PMID: 17437720

35. Grishin, N. V. KH domain: one motif, two folds. *Nucleic Acids Res* **29**, 638-643 (2001). PubMed PMID: 11160884

36. Hollingworth, D., Candel, A. M., Nicastro, G., Martin, S. R., Briata, P., Gherzi, R. & Ramos, A. KH domains with impaired nucleic acid binding as a tool for functional analysis. *Nucleic Acids Res* **40**, 6873-6886 (2012). PubMed PMID: 22547390

37. Ulgherait, M., Rana, A., Rera, M., Graniel, J. & Walker, D. W. AMPK modulates tissue and organismal aging in a non-cell-autonomous manner. *Cell Rep* **8**, 1767-1780 (2014). PubMed PMID: 25199830

38. Wucherpfennig, T., Wilsch-Brauninger, M. & Gonzalez-Gaitan, M. Role of Drosophila Rab5 during endosomal trafficking at the synapse and evoked neurotransmitter release. *J Cell Biol* **161**, 609-624 (2003). PubMed PMID: 12743108

39. Zhu, M., Li, X., Tian, X. & Wu, C. Mask loss-of-function rescues mitochondrial impairment and muscle degeneration of Drosophila pink1 and parkin mutants. *Hum Mol Genet* **24**, 3272-3285 (2015). PubMed PMID: 25743185

40. Linford, N. J., Bilgir, C., Ro, J. & Pletcher, S. D. Measurement of lifespan in Drosophila melanogaster. *J Vis Exp* (2013). PubMed PMID: 23328955

Figure Legend

Figure 1. Overexpressing Mask in TH-C' or TH-D' DANs Extends Lifespan.

Survivorship curves of control flies and flies overexpressing Mask under the control of A) 0273-Gal4, B) TH-C'-Gal4, and C) TH-D'-Gal4 drivers. D) Schematic of expression patterns of 0273-, TH-C'- and TH-D'-Gal4 drivers.

Figure 2. Moderately activating the ectopically expressed DREADD system in the TH-C' or TH-D' DANs Extends Lifespan in flies.

Survivorship curves of male and female adult flies expressing the DREADD Gαs receptor and raised on food containing DMSO only (No Drug) or 1uM Clozapine N-oxide (CNO). A, B) moderately activating the DREADD-Ds receptor in the TH-C' neurons leads to extension of lifespan in flies. The median lifespan increased from ~87 days to ~98 days (~12.6% increase) in the male flies and from ~95 days to ~108 days (~12.6% increase) in the female flies. C, D) moderate activation of the DREADD-Ds receptor expressed in the TH-D' neurons leads to significant extension of lifespan in flies. The median lifespan increased from ~73 days to ~95 days (~30% increase) in the male flies and from ~67 days to ~107 days (~50% increase) in the female flies.

Figure 3. Mask-overexpressing Flies show sustained Fecundity at old ages. A) Numbers of progenies produced by female flies in a 3 day interval. The female flies that overexpress Mask in the TH-C' or TH-D' DA neurons produce significantly more progenies at 30- and 50-day old ages. B) Reproduction in male flies is measured as the percentage of male flies that are fertile.

At age 50-day, majority of the control male flies lose the ability to produce progeny, while over 80% of male flies that overexpress Mask in the TH-C' and 60% of those overexpress Mask in the TH-D' DANs are still reproductively active.

Figure 4. Overexpressing Unc-104 or Knocking down p150^{Glued} in the TH-C' or TH-D' DANs extend lifespan in flies.

Survivorship curves of female flies expressing UAS-GFP-Unc104 or a weak RNAi transgene for p150^{Glued} in the TH-C' or TH-D' DANs. A) Overexpressing the Unc-104-GFP transgene in the TH-C' or TH-D' neurons leads to extension of lifespan in flies. The median lifespan in the flies overexpressing Unc-104-GFP increased from ~67 days in control flies) to ~92 days (~37% increase). In the flies overexpressing Unc-104-GFP in the TH-D' neurons, the median lifespan increased ~40% (median lifespan ~94 days). B) Moderately knocking down p150^{Glued} in the TH-C' neurons leads to moderate extension of lifespan in flies, median lifespan increased from ~86 days to ~98 days (~13.95% increase). Same treatment in the TH-D' neurons leads to an increase of the median lifespan to ~92 days (~6.97% increase).

Figure 1

Figure 2

Figure 3

Figure 4

Female