Chemical Systems Biology Reveals Mechanisms of Dissociated Glucocorticoid Receptor Signaling

Nelson E. Bruno[1], Jerome C. Nwachukwu[1], Sathish Srinivasan[1], Charles C. Nettles[1], Tina Izard[1], Zhuang Jin[2*], Siddaraju V. Boregowda [2], Donald G. Phinney[2], Olivier Elemento[3], Xu Liu[4], Eric A. Ortlund[4], René Houtman[5], Diana A. Stavreva[6], Gordon L. Hager[6], Theodore M. Kamenecka[2], Douglas J. Kojetin[1,2], Kendall W. Nettles[1].

1. Department of Integrative Structural and Computational Biology, The Scripps Research Institute, Jupiter, FL 33458 USA

2. Department of Molecular Medicine, The Scripps Research Institute, Jupiter, FL 33458 USA

3. Caryl and Israel Englander Institute for Precision Medicine, Institute for Computational Biomedicine, Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY 10065, USA

4. Department of Biochemistry, Emory University School of Medicine, Atlanta, GA 30322, USA

5. Department of Research and Development, PamGene, 's-Hertogenbosch, 5211 HH, the Netherlands.

6. Laboratory of Receptor Biology and Gene Expression, Building 41, B602, 41 Library Dr., National Cancer Institute, NIH, Bethesda, MD 20892-5055, USA

* Current affiliation: Corbus Pharmaceuticals, Inc. 500 River Ridge Dr. Norwood, MA 02062.

Correspondence: Kendall Nettles, knettles@scripps.edu

Abstract

Glucocorticoids display remarkable anti-inflammatory activity, but their use is limited by on-target

adverse effects including insulin resistance and skeletal muscle atrophy. We used a chemical

systems biology approach Ligand Class Analysis (LCA) to examine ligands designed to modulate

glucocorticoid receptor activity through distinct structural mechanisms. These ligands displayed

diverse activity profiles, providing the variance required to identify target genes and coregulator

interactions that were highly predictive of their effects on myocyte glucose disposal and protein

balance. Their anti-inflammatory effects were linked to glucose disposal but not muscle atrophy.

This approach also predicted dissociated activity *in vivo*, identifying compounds that were muscle

sparing or anabolic for protein balance and mitochondrial potential. LCA defines the mechanistic

links between the ligand-receptor interface and the ligand-driven physiological outcomes, a

general approach that can be applied to any ligand-regulated allosteric signaling system.

Introduction

Glucocorticoids (GCs) are among the most prescribed medicines due to their remarkable anti-inflammatory effects. GCs also cause insulin resistance, reduce glucose uptake in skeletal muscle—the organ responsible for ~80% of glucose disposal—and facilitate muscle wasting by reducing protein synthesis and increasing protein degradation[1,2]. Efforts to develop improved GCs have been hampered by poor understanding of the structural and molecular mechanisms through which glucocorticoid receptor (GR) ligands drive diverse phenotypic outcomes, such as effects on inflammation, glucose disposal, or skeletal muscle atrophy. This phenomenon is called selective modulation or dissociated signaling, and is a common feature of allosteric signaling through GR and other nuclear receptor superfamily transcription factors[3], G protein-coupled receptors, and other small molecule drug targets.

GCs are secreted by the adrenal gland in a circadian fashion and in acute response to nutrient deprivation or other stressors. In skeletal muscle, GCs stimulate catabolic signaling that opposes the insulin/PI3K/AKT signaling pathway, inhibits glucose uptake, and releases amino acids by inhibiting protein synthesis and stimulating protein degradation[2]. Upon GC binding, GR dissociates from inhibitory heat shock protein complexes and translocated to the nucleus, oligomerizes upon binding to DNA, and recruits an ensemble of coregulators, which include chromatin-remodeling and histone-modifying enzymes that regulate gene expression. Several GR coregulators and target genes have been identified in skeletal muscle[4-7], but factors that drive ligand-selective modulation are unclear.

We developed a chemical systems biology approach called Ligand Class Analysis (LCA) to identify molecular mechanisms that drive ligand-selective signaling[8-10]. With LCA, we characterize a series of related compounds that display a full range of variance in phenotypes such

as insulin-mediated glucose uptake or protein synthesis and use that variance to identify molecular

predictors of the phenotype. This enables us to identify ligand-dependent gene expression and

protein-interaction profiles that predict molecular phenotypes induced by different classes of

ligands that perturb receptor structure differently[8-10]. We then overexpress or knockdown

predictive genes or interacting proteins to determine which factors are required for ligand-

dependent phenotypes.

In this work, we evaluated the physiological effects of GCs during overnight nutrient

deprivation when they coordinate acute metabolic adaptations, including effects on glucose

disposal and protein balance. We tested a series of GCs modified at different positions on the

steroidal scaffold[11], providing distinct strategies to perturb GR structure and dynamics. Using a

machine learning approach with LCA, we found that the effects of GCs on protein balance in

skeletal muscle can be dissociated from their effects on glucose disposal and the inflammatory

response, allowing us to discover the underlying coregulators and genes driving these activities,

and discover orally available dissociated glucocorticoids with muscle-sparing activity profiles in

vivo.

Results

Development of a physiologically relevant skeletal muscle profiling platform for effects of

GCs on glucose disposal and protein balance during nutrient deprivation

We designed a series of GCs to test the role of three different structural perturbations found in

compound PF-0251802 (PF802) (**1**, **Figure 1a**)[12], a dissociated GR ligand, the pro-drug of which

reached clinical trials [13]. PF802 has a benzyl group to drive antagonism by disrupting the agonist-

bound GR conformation (**Figure 1b**), similar to the carbon-11 (C11) substitution in the antagonist

RU-486 (**Figure 1c**), and a trifluoromethyl group attached at the equivalent position of C17 on a

steroid to increase affinity (**Figure 1c-d**). PF802 also contains a methylpyridinyl acetamide group

attached at the equivalent position to C3 in the steroid scaffold. We predict that this group enters

the solvent channel underneath the coregulator binding site, as first seen with deacylcortivazol

(**Figure 1e**), to alter the shape of AF2, and the ensemble of coregulators that bind this surface, thus

driving dissociated effects as we previously showed for the estrogen receptor[9,14].

To understand the how these different substitutions control GR activity, we generated a set

of 22 related compounds based on a steroidal core, which enabled us to understand where

substitutions are directed within the ligand-binding site (**6-27**, **Figure 1f-g**). These included a

number of substitutions at C11 to drive agonism/antagonism (**13-20 Figure 1g**), different

substitutions at C17 to optimize affinity (**21-24**, **Figure 1g**), and the methylpyridinyl acetamide at

C3 to perturb the AF2 surface (**6-12**, **Figure 1g**), where the position of methyl and nitrogen in the

pyridine ring were altered. We have published the synthesis and preliminary structure-activity

relationships for these compounds[11].

We developed a profiling platform to examine how ligand-regulated GR nuclear

localization, structure, and target gene expression in skeletal muscle control glucose disposal and

protein balance (**Figure 1h, Methods**), and whether these effects could be dissociated from anti-

inflammatory effects such as the suppression of IL-6 secretion by A549 lung cells. We assayed

ligand-induced nuclear translocation of a GFP-GR fusion protein, and probed for ligand-selective

changes in surface structure by characterizing the GR-interaction profiles of 154 peptides derived

from nuclear receptor coregulators and other interacting proteins using the MARCoNI FRET

assay[15]. We also completed nascent RNA expression profiling of C2C12 myotubes, one hour after

vehicle or Dexamethasone (Dex) treatment and selected 28 target genes as potential regulators of

glucose disposal and protein balance (the rationale for compound selection is in **Supplementary**

Information Methods). We then compared ligand-dependent regulation of these target genes in

myotubes using the nanoString nCounter for direct multiplexed transcript counting without PCR

amplification, enabling highly quantitative and reproducible assays[16]. These datasets comprise the

independent variables that we used in a machine learning approach described below, to identify

ligand-selective regulators of glucose disposal and protein balance in skeletal muscle, the

dependent variables.

To assay physiologically relevant effects of GCs on skeletal muscle, we treated the

myotubes in the context of nutrient deprivation and a brief insulin challenge. One of the barriers

to understanding GC action in skeletal muscle is that effects on protein balance appear to require

50–100 μM Dex *in vitro* [17], despite a ~5 nM Kd for GR. The physiological effects of GCs on

protein balance and Glut4 translocation vary *in vivo*, depending upon the stressor. For example,

GCs are secreted in response to fasting or exercise, but these stressors have very different effects

on skeletal muscle. GCs rapidly induce skeletal muscle atrophy and insulin resistance during

fasting, while exercise has opposite effects[18,19]. We found that serum starving the myotubes

overnight and administering a brief insulin challenge was sufficient to generate assays with robust

signal to noise, suggesting that effects of GR are indeed through inhibiting the insulin/PI3K/AKT

signaling pathway[20]. These optimizations enabled highly robust and reproducible assays for

glucose disposal and protein balance (see **Methods**), with maximal effects of Dex in the 1-10 nM

range (**Supplementary Figure 1a–g**).

We developed a screening assay for phosphorylation of AKT at Thr308 (pAKT), reflecting

its joint role in coordinating both protein synthesis and glucose disposal (**Supplementary Figure**

1a–b). For glucose disposal, we measured the rate limiting step, transfer of the glucose receptor,

Glut4, to the cell membrane (**Supplementary Figure 1c**). For protein balance, we developed two

assays: protein degradation was assayed by release of tritiated phenylalanine (degradation)

(**Supplementary Figure 1d**); and insulin-stimulated protein synthesis (synthesis) was assayed

using the SUnSET assay (**Supplementary Figure 1e**), which measures the incorporation of

puromycin into nascent polypeptide chains. We assessed the relationship between protein balance

and mitochondrial potential (ψm) measured by high content imaging after staining with a

MitoTracker dye, where accumulation of the dye is dependent upon membrane potential.

(**Supplementary Figure 1f–h**). We also compared ligand-dependent effects on IL-1β-induced

secretion of IL-6 by A549 lung cells, a standard assay for the inflammatory response. Ligand

activity in these assays represent the dependent variables to identify predictive coregulator

interactions and target genes, and below, individual compounds with interesting pharmacological

properties.

After profiling the compounds, we asked which dependent variables show similar ligand

response patterns, and identified two clusters, connected by pAKT. In the first cluster, effects of

the ligands on protein synthesis were similar to effects on protein degradation, pAKT, and ψm,

suggesting transcriptional control via a common set of coregulators and target genes (**Figure 1i,**

Supplementary Figure 1i, Supplementary Dataset shows the correlation matrix). In the second

cluster, effects of the ligands on pAKT showed patterns similar to effects on IL-6 and Glut4

(**Figure 1i**), while effects on nuclear translocation of GR also correlated with their effects on IL-6

and Glut4. However, effects on pAKT and GR nuclear translocation were dissociated

(**Supplementary Figure 1j**). This result indicates that some ligand-dependent outcomes of GR

signaling (e.g. Glut4, IL-6) involve a graded response to the amount of nuclear GR, while others

require a threshold amount of nuclear GR. Nevertheless, these results suggest that the ligands can

use different transcriptional signaling pathways—coregulators and target genes—that coordinate

some processes but allow others to be dissociated.

Machine learning defines the best sets of predictors for GC effects in skeletal muscle

Using Random Forest, a classification and regression tree algorithm [21], we performed predictive

modeling of each GR-mediated phenotype or dependent variable (**Figure 1h**), based on all other

features including the independent variables (i.e. GR-peptide interactions and -target genes). To

set a statistical threshold, we used *Boruta*, a Random Forest-based wrapper algorithm, to identify

a set of predictors for each GR-mediated phenotype, by iteratively comparing the importance of

each feature in the dataset, to the highest importance observed in a "shadow" dataset obtained by

shuffling every variable [22]. In each of 500 iterative decision rounds, a feature is selected as a

predictor if it is more important than the most important shadow feature. In contrast, a feature is

rejected if it has not been selected after multiple consecutive decision rounds. We subsequently

identified a minimal set of predictors, by applying a forward selection-based strategy. Predictors

of each GR-mediated phenotype previously identified via *Boruta*, were tested one-by-one, starting

with the most important, for their ability to improve the predictive capacity of the model. Predictors

that improved the model were retained, while those that worsened the model were removed, until

the set of predictors was expended. With this approach we were able to predict between 46.8%

(degradation) and 84.3% (Glut4) of the variance in the dependent variables (**Figure 2a,**

Supplementary Dataset).

This analysis revealed which target genes and interacting peptides were selective or

common predictors of the effects of GR ligands on skeletal muscle (**Figure 2a–b, Supplementary**

Figure 2a–b). The selective prediction of Glut4 but not ψm by the *Bcl2l1* gene was especially

striking (**Figure 2b**, **Supplementary Figure 2b**). *Bcl2l1* encodes Bcl-xL, a mitochondrial anti-

apoptotic BCL-2 family protein that inhibits glucose metabolism and mitochondrial-mediated

secretion of insulin by pancreatic β-cells [23], suggesting that it may also contribute to glucose

disposal in skeletal muscle.

Linear regression showed that a SMRT peptide interaction with GR significantly predicted

synthesis, but not Glut4 or the best Glut4-predictive target genes (**Figure 2c**). There was a striking

connection between the coregulator peptides that predicted Glut4, showing very low r^2 with the

protein balance assays, but very high predictive power for the specific genes that best predicted

Glut4 activity (**Figure 2c**). While the MarCoNI peptide set is best viewed as a probe for ligand-

regulated effects on surface structure, all of the peptides in **Figure 2c** represent bona fide nuclear

receptor interaction motifs. Thus, we can exploit the variance in ligand activity to identify the

common and dissociated transcriptional networks underlying GR-mediated phenotypes in skeletal

muscle.

The variance in the ligand activity profiles also allowed LCA to identify signal in what

would typically be considered noise. Both *Bcl2l1* and *Socs2* showed less than two-fold induction,

the standard threshold for noise in gene expression data. Much of the gene expression data

displayed an inflection point (**Supplementary Figure 2c**). When we removed the data below the

inflection point, *Fkbp5, Blc2l1*, *Tsc22d3* still predicted Glut4 translocation (**Supplementary**

Figure 2d), suggesting that insulin-mediated glucose disposal can be fine-tuned by very small

changes in target gene expression. We call these super-resolution analyses, similar to what we

have seen with super-resolution X-ray crystallography, where comparing many structures bound

to a series of compounds identified ligand-induced perturbations within the noise of an individual

crystal structure that drive biological outcomes[8]. Thus, subtle changes in gene expression that

would typically be considered noise and disregarded, are sufficient to identify transcriptional

networks underlying GR-mediated phenotypes through LCA.

We also found that different substitutions on the steroidal scaffold, designed to perturb

different GR differently, modulated distinct transcriptional networks. A comparison of ligand-

dependent effects on Glut4 revealed a dramatic increase in predictive power by the C11-substituted

compounds compared to the whole compound set, while the C3-substituted compounds exhibited

a widespread reduction in r^2 (**Figure 2d–e, Supplementary Figure 2e,** compare to **Figure 2b**).

This was evident despite their similar variances in the skeletal muscle profiling assays

(**Supplementary Figure 2f**), supporting the idea that C3- and C11-substituted compounds utilize

different structural mechanisms and transcriptional networks to modulate GR-mediated

phenotypes, which we explore below with molecular dynamics simulations.

Gene perturbation studies define underlying causal relationships for GR-mediated skeletal

muscle phenotypes

We performed gene perturbation studies to determine which correlations are due to causal

relationships. The Dex-induced gene, *Fkbp,5* is a good candidate to coordinate metabolic effects

of GCs at an organismal level, as its protein product facilitates dephosphorylation of pAKT by the

phosphatase, PHLPP1[24]. In addition, Fkbp5 knockout mice show greater skeletal muscle insulin

sensitivity and reduced adiposity on a high fat diet[25]. We electroporated *GFP* and either *Fkbp5* or

Foxo1 (an atrophy inducing control gene) expression plasmids into contralateral tibialis anterior

(TA) muscles. The *Fkbp5*- or *Foxo1*-transduced muscles were approximately 10% smaller than

control (**Figure 3a**). Electroporation with *Foxo1* or *Fkbp5* inhibited insulin-induced *de novo*

protein synthesis (**Figure 3b**) and pAKT *in vivo* (**Figure 3c**), demonstrating that *Fkbp5* regulates

protein balance in skeletal muscle.

Knockdown of NCoR or SMRT (gene names *Ncor1* and *Ncor2*, respectively) enhanced

Dex-dependent inhibition of protein synthesis (**Figure 3d, Supplementary Figure 3**), consistent

with the peptide binding data showing that GR interaction with NCoR and SMRT corepressor

peptides were stimulated by anabolic compounds (**Figure 2b**). In contrast, knockdown of NCOA6

and PELP1 fully reversed the inhibitory effect of Dex, while knockdown of SRC2 (gene name

Ncoa2), but not SRC1 (gene name *Ncoa1*), partially reversed this effect (**Figure 3d**). In contrast,

pAKT was regulated by SRC1, as well as PELP1 (**Figure 3e**). These data support a model where

GR recruits specific coregulators to distinct subsets of target genes that can have overlapping or

different effects on skeletal muscle metabolism and protein balance.

Discovery of dissociated GCs with improved skeletomuscular activity profiles

Using our systems chemical biology approach, we developed and characterized compounds that—

as a set—showed dissociated effects on glucose disposal and protein balance (**Figure 1**) and

identified some unique genes and coregulator interactions associated with these effects (**Figures**

2–3). We now wanted to ask whether any of the individual compounds showed dissociated effects

(see **Supplementary Figure 4** for compound SAR summary or **Supplementary Dataset** for all

data) that might have therapeutic potential and identified two promising glucocorticoids with

dissociated effects on inflammation and protein balance (**Figure 4a–d**). Three compounds in the

C11 series of compounds inhibited IL-6 with efficacies comparable to Dex (**13-15**,

Supplementary Figure 4a). Among these, SR11466 (**15**) is a partial GR agonist (EC50 = 0.1 nM)

that showed no effects on protein balance, and a slight improvement (i.e. less inhibition) compared

to Dex in the Glut4 assay (**Figure 4a–d**, **Supplementary Figure 4b**). We also identified SR16024

(**18**) as a full GR antagonist (IC50 = 1.4 nM) that stimulated protein synthesis but did not suppress

protein degradation or increase Glut4 translocation (**Figure 4b-d**, **Supplementary Figure 4b**).

SR11466 and SR16024 increased mitochondrial potential, with the latter doubling mitochondrial

potential (**Figure 4e–f**).

As a further test for dissociation, the compounds were profiled in primary human

osteoblasts for effects on mineralization during 4 weeks of differentiation from mesenchymal stem

cells. SR11466 showed a slight improvement in the inhibition of mineralization compared to Dex

and PF802, while SR16024 showed a better profile (**Figure 4g**). We also noticed SR15421 (**9**), a

lower affinity compound with EC50 of 240 nM (**Supplementary Figure 7a**), that was anti-

inflammatory with less impact on mineralization (**Figure 4g-h**). SR15421 is part of the C3-

substituted series, where modifying the position of the methyl and the nitrogen on the

methylpyridine had profound effects on inhibition of IL-6 secretion (**Figure 4h**). Dose response

curves reveal a moderately, but significantly improved effect of SR11466 on mineralization

compared to Dex (2-way ANOVA, drug p = 0.0018, drug x time p = 1.4×10^{-4}), while SR15421

and SR16024 inhibited mineralization only at the 10 μM dose (**Figure 4h**). The mineralization

data showed very low correlations with the other assays, except for protein synthesis (Pearson r =

0.54, p = 4×10^{-3}), supporting the joint regulation of bone and muscle mass. These data suggest

that it is possible to find GCs with improved bone- and muscle-sparing properties compared to

Dex using our in vitro profiling platform.

We opted to test the two high affinity ligands further in mechanism of action and animal

studies, based on their favorable pharmacokinetics (**Supplementary Figure 4c**), and on-target

mechanism of action (**Figure 5a–b**, **Supplementary Figure 5**). In vivo, SR11466 strongly

suppressed LPS-induced TNFα levels in the blood, while the full antagonist SR16024 was not

inhibitory (**Figure 5c**). We also assessed loss of lean mass following a larger dose of LPS, which

was significantly worsened by Dex. Both SR11466 and SR16024 were significantly better than

Dex, while SR16024 was protective against loss of lean mass compared to vehicle (**Figure 5d**).

The differential effects of SR11466 on inflammation and proteostasis demonstrate that it is a bona

fide dissociated glucocorticoid, while SR16024 displayed stronger muscle sparing activities.

A lactate tolerance test was then administered to probe for differences in liver

gluconeogenesis following an overnight fast. Here again, Dex caused significantly greater loss of

lean mass after the fast; both SR11466 and SR16024 were significantly better than Dex (**Figure**

5e); SR16024 attenuated the loss of overall body weight (**Figure 5f**). The lactate tolerance test

showed that both Dex and SR11466 increased the rate of glucose production, while SR16024 did

not (**Figure 5g**), suggesting that ligand-specific GC effects on glucose uptake in skeletal muscle

and gluconeogenesis in the liver may utilize common signaling mechanisms, such as ligand-

selective coregulator recruitment, to integrate organismal stress responses.

Molecular dynamics simulations revealed that suppression of IL-6 is associated with specific

structural perturbations.

To understand how substitutions to the C3 position regulate allosteric signaling in GR, we

performed all atom molecular dynamics (MD) simulations comparing **7** and **9**, which differ only

in the placement of the methyl on the pyridine (**Figure 1g**) but showed very different effects on

suppression of IL-6 (**Figure 4h**). Molecular docking demonstrated that the compounds bound

similarly to Dex, with the pyridine extending off the A-ring into a solvent channel, in between

helix 3 (h3) and h5 (**Supplementary Methods**, **Supplementary Figure 6a**). In three independent

1 µs simulations of ligand-bound and ligand-free (apo) GR LBD, we observed no significant

structural changes in the conformation of helix 12 (h12). However, ligand anti-inflammatory

activity (Dex > **9** >**7** > apo) was associated with a multi-Å decrease in the distance between the C-

terminus of h11 and the N-terminus of h3, and a decrease in the heterogeneity of this h3/h11

interface (**Figure 6a–c**). We also assessed patterns of correlated motion and found that the **7**

showed less correlation between residues h3 and h11/h12 residues compared to dex, with 9 again

showing intermediate effects (**Supplementary Figure 6b**).

To further probe for effects of the C3 substitutions on correlated motion between h12 and

the solvent channel, we used dynamical network analysis[26] of ligand-free and the ligand-bound

complexes by connecting pairs of nodes with their edges if they have satisfied a distance

requirement (<4.5 Å) for at least 75% of the simulation time. Edge distance is inversely

proportional to the pairwise correlations between two nodes; thus, a short path length indicates a

strong correlation/communication. A pair of distal nodes are connected by the optimal (shortest)

path and the suboptimal (longer) paths with the length defined as the sum of edge distances along

the path. We analyzed the top 1000 suboptimal paths between residue E755 and R614 as two nodes

to study the allosteric communication between h12 and the C-terminus of h5, where the

modifications on the steroidal A ring extend to the solvent channel. With apo GR, this

communication was through h11 and along h5, while with dex it wrapped around the ligand

binding pocket from h12 to h3 and then through the β-sheet (**Figure 6d**), enabled by the dex-

induced closing of the distance between h3 and h11 (**Figure 6b**). With Dex, this was associated

with much longer path lengths (**Figure 1e**), representing weaker allosteric communication overall.

With **9**, there were less of these extended path lengths, while **7** more closely resembled the apo

GR (**Figure 6d-e**). An analysis of residues contacting the C3-substituted pyridine showed that

differential positioning of the R611 side chain on h5, which can form h-bonds with the pyridine or

A-ring ketone of Dex, can drive the orientation of the steroid core to control the dynamics and

surface structure and associated receptor activity (**Figure 6f–g, Supplementary Figure 6c–e**).

Discussion

In this work, we used a structure-based design strategy and LCA to reveal how GCs can use

different coregulators and transcriptional networks for the physiological control of glucose

disposal and protein balance during acute nutrient deprivation. By examining closely related

compounds with LCA, we determined that protein balance and mitochondrial potential were

intercorrelated but dissociated from Glut4 translocation and suppression of IL-6. Using a

physiologically relevant skeletal muscle profiling platform optimized to study insulin signaling

during nutrient deprivation, we were able to identify SR11466 as a partial GR agonist that was not

catabolic, demonstrating in vivo dissociated anti-inflammatory activity. In fact, SR11466 was

slightly anabolic for protein balance, while maintaining strong anti-inflammatory activity in vivo.

Such compounds may be useful treatments for cachexia, muscular dystrophies, back pain or

osteoarthritis. We also identified SR16024 as a highly anabolic GC that inhibits both fasting- and

LPS-induced weight loss, while stimulating mitochondrial potential and protein synthesis in

response to insulin. Further studies are needed to evaluate the long term effects of SR16024, but

GR antagonists have been tested clinically for depression[27], and may have efficacy in muscular

dystrophies and enzalutamide-resistant prostate cancers that have switched from androgen to

glucocorticoid dependency[28]. While preliminary, the lack of SR15421 effects on mineralization

suggests that it may also be possible to dissociate the anti-inflammatory and osteoporotic effects

of GCs.

The unbiased nature of LCA enabled the identification of a number of unexpected signaling

and biophysical properties of GR. Insulin receptor signaling pathways control both anabolic and

catabolic signaling, and these can be dissociated by transcriptional control of pathway-selective

regulators, often with very subtle changes in gene expression. Covariance among assays reflects

common underlying signaling mechanisms driven by common receptor conformations,

coregulators, and target genes. The correlation between effects on bone and skeletal muscle was

not surprising, but the connection between anti-inflammatory and metabolic effects of GCs in

different cell types was unexpected and may point to an organismal level of evolutionary

connection between underlying signaling pathways. GC coordination of the inflammatory

response often coincides with immune differentiation[29,30], which requires metabolic adaptations

[31,32]. This suggests that GCs may use a common set of coregulators to coordinate anti-

inflammatory effects with modulation of metabolism. LCA provides a mechanism to identify

common signaling mechanisms among different cell types and processes.

GC-regulated transcription is controlled via underlying biophysical properties of GR, such

as oligomerization, nuclear translocation, and on/off rates of interactions with an ensemble of

response elements, coregulators, and collaborating transcription factors. We have shown here that

LCA predicts that ligand-selective surface conformers—identified from a structural peptide

interaction assay—can predict differential effects of the ligands, which we validated with

knockdown of selected coregulators. We and others have observed a similar dichotomy where

Ncoa2/SRC2/GRIP1 is required for NF-κB-induced cytokine production, but then switches to a

repressor upon recruitment of GR to the promoter[33,34], representative of a more general phenomena

of receptors using many coregulators to regulate a single gene, but also some shared and some

differential coregulator usage across genes[14,35,36]. The selective association of GR nuclear

translocation with glucose disposal and not protein balance was also surprising. It suggests that

GCs modulate insulin dependent Glut4 translocation via a graded transcriptional program that is

highly sensitive to the amount of nuclear GR, and control protein balance through transcriptional

responses that require a threshold amount of nuclear GR. This is consistent with other data

suggesting that only 10% of GR may be DNA-bound at any given time [37]. Similarly, we previously

found that the mammalian two-hybrid assay was a much better predictor of proliferation by

estrogen receptor ligands than a chromatin immunoprecipitation assay [9], suggesting that complex

biology can be simulated by a set of simple biophysical models and their importance identified

with LCA.

370
371 **References**
372

1.   Watson, M.L. et al. A cell-autonomous role for the glucocorticoid receptor in skeletal
muscle atrophy induced by systemic glucocorticoid exposure. Am J Physiol Endocrinol Metab
302, E1210-20 (2012).
2.   Bodine, S.C. & Furlow, J.D. Glucocorticoids and Skeletal Muscle. Adv Exp Med Biol 872,
145-76 (2015).
3.   Nettles, K.W. & Greene, G.L. Ligand control of coregulator recruitment to nuclear
receptors. Annu Rev Physiol 67, 309-33 (2005).
4.   Kuo, T. et al. Genome-wide analysis of glucocorticoid receptor-binding sites in myotubes
identifies gene networks modulating insulin signaling. Proc Natl Acad Sci U S A 109, 11160-5
(2012).
5.   Chen, S.L., Dowhan, D.H., Hosking, B.M. & Muscat, G.E. The steroid receptor
coactivator, GRIP-1, is necessary for MEF-2C-dependent gene expression and skeletal muscle
differentiation. Genes Dev 14, 1209-28 (2000).
6.   Tobimatsu, K. et al. Overexpression of the transcriptional coregulator Cited2 protects
against glucocorticoid-induced atrophy of C2C12 myotubes. Biochem Biophys Res Commun 378,
399-403 (2009).
7.   Amat, R., Solanes, G., Giralt, M. & Villarroya, F. SIRT1 is involved in glucocorticoid-
mediated control of uncoupling protein-3 gene transcription. J Biol Chem 282, 34066-76 (2007).
8.   Nwachukwu, J.C. et al. Systems Structural Biology Analysis of Ligand Effects on ERalpha
Predicts Cellular Response to Environmental Estrogens and Anti-hormone Therapies. Cell Chem
Biol 24, 35-45 (2017).
9.   Nwachukwu, J.C. et al. Predictive features of ligand-specific signaling through the estrogen
receptor. Mol Syst Biol 12, 864 (2016).
10.   Srinivasan, S. et al. Ligand-binding dynamics rewire cellular signaling via estrogen
receptor-alpha. Nat Chem Biol 9, 326-32 (2013).

11.   Jin, Z. et al. Synthesis of novel steroidal agonists, partial agonists, and antagonists for the
glucocorticoid receptor. Bioorg Med Chem Lett 27, 347-353 (2017).
12.   Hu, X. et al. The antagonists but not partial agonists of glucocorticoid receptor ligands
show substantial side effect dissociation. Endocrinology 152, 3123-34 (2011).
13.   Stock, T., Fleishaker, D., Wang, X., Mukherjee, A. & Mebus, C. Improved disease activity
with fosdagrocorat (PF-04171327), a partial agonist of the glucocorticoid receptor, in patients with
rheumatoid arthritis: a Phase 2 randomized study. Int J Rheum Dis (2017).
14.   Nwachukwu, J.C. et al. Resveratrol modulates the inflammatory response via an estrogen
receptor-signal integration network. Elife 3, e02057 (2014).
15.   Aarts, J.M. et al. Robust array-based coregulator binding assay predicting ERalpha-agonist
potency and generating binding profiles reflecting ligand structure. Chem Res Toxicol 26, 336-46
(2013).
16.   Geiss, G.K. et al. Direct multiplexed measurement of gene expression with color-coded
probe pairs. Nat Biotechnol 26, 317-25 (2008).
17.   Stitt, T.N. et al. The IGF-1/PI3K/Akt pathway prevents expression of muscle atrophy-
induced ubiquitin ligases by inhibiting FOXO transcription factors. Mol Cell 14, 395-403 (2004).
18.   Goodyear, L.J. & Kahn, B.B. Exercise, glucose transport, and insulin sensitivity. Annu Rev
Med 49, 235-61 (1998).
19.   Magomedova, L. & Cummins, C.L. Glucocorticoids and Metabolic Control. Handb Exp
Pharmacol 233, 73-93 (2016).
20.   Kuo, T., Harris, C.A. & Wang, J.C. Metabolic functions of glucocorticoid receptor in
skeletal muscle. Mol Cell Endocrinol 380, 79-88 (2013).
21.   Breiman. Random Forests. Machine Learning 45, 5-32 (2001).
22.   Kursa, M.B. Robustness of Random Forest-based gene selection methods. BMC
Bioinformatics 15, 8 (2014).
23.   Luciani, D.S. et al. Bcl-2 and Bcl-xL suppress glucose signaling in pancreatic beta-cells.
Diabetes 62, 170-82 (2013).
24.   Pei, H. et al. FKBP51 affects cancer cell response to chemotherapy by negatively
regulating Akt. Cancer Cell 16, 259-66 (2009).
25.   Balsevich, G. et al. Stress-responsive FKBP51 regulates AKT2-AS160 signaling and
metabolic function. Nat Commun 8, 1725 (2017).
26.   Bowerman, S. & Wereszczynski, J. Detecting Allosteric Networks Using Molecular
Dynamics Simulation. Methods Enzymol 578, 429-47 (2016).
27.   Block, T. et al. Mifepristone Plasma Level and Glucocorticoid Receptor Antagonism
Associated With Response in Patients With Psychotic Depression. J Clin Psychopharmacol 37,
505-511 (2017).
28.   Arora, V.K. et al. Glucocorticoid receptor confers resistance to antiandrogens by bypassing
androgen receptor blockade. Cell 155, 1309-22 (2013).
29.   Yu, B. et al. Epigenetic landscapes reveal transcription factors that regulate CD8(+) T cell
differentiation. Nat Immunol 18, 573-582 (2017).
30.   Coppo, M., Chinenov, Y., Sacta, M.A. & Rogatsky, I. The transcriptional coregulator
GRIP1 controls macrophage polarization and metabolic homeostasis. Nat Commun 7, 12254
(2016).
31.   Geltink, R.I.K., Kyle, R.L. & Pearce, E.L. Unraveling the Complex Interplay Between T
Cell Metabolism and Function. Annu Rev Immunol 36, 461-488 (2018).

32.   Van den Bossche, J., O'Neill, L.A. & Menon, D. Macrophage Immunometabolism: Where Are We (Going)? Trends Immunol 38, 395-406 (2017).

33.   Hudson, W.H. et al. Cryptic glucocorticoid receptor-binding sites pervade genomic NF-kappaB response elements. Nat Commun 9, 1337 (2018).

34.   Chinenov, Y. et al. Role of transcriptional coregulator GRIP1 in the anti-inflammatory actions of glucocorticoids. Proc Natl Acad Sci U S A 109, 11776-81 (2012).

35.   Stallcup, M.R. & Poulard, C. Gene-Specific Actions of Transcriptional Coregulators Facilitate Physiological Plasticity: Evidence for a Physiological Coregulator Code. Trends Biochem Sci 45, 497-510 (2020).

36.   Stashi, E., York, B. & O'Malley, B.W. Steroid receptor coactivators: servants and masters for control of systems metabolism. Trends Endocrinol Metab 25, 337-47 (2014).

37.   Voss, T.C. et al. Dynamic exchange at regulatory elements during chromatin remodeling underlies assisted loading mechanism. Cell 146, 544-54 (2011).

38.   Heikkinen, S., Argmann, C.A., Champy, M.F. & Auwerx, J. Evaluation of glucose homeostasis. Curr Protoc Mol Biol Chapter 29, Unit 29B 3 (2007).

39.   Bruno, N.E. et al. Creb coactivators direct anabolic responses and enhance performance of skeletal muscle. EMBO J 33, 1027-43 (2014).

40.   Hong, D.H. & Forsberg, N.E. Effects of dexamethasone on protein degradation and protease gene expression in rat L8 myotube cultures. Mol Cell Endocrinol 108, 199-209 (1995).

41.   Walker, D., Htun, H. & Hager, G.L. Using inducible vectors to study intracellular trafficking of GFP-tagged steroid/nuclear receptors in living cells. Methods 19, 386-93 (1999).

42.   Stavreva, D.A. et al. Prevalent glucocorticoid and androgen activity in US water sources. Sci Rep 2, 937 (2012).

Acknowledgements. N.E.B. was supported by the BallenIsles Men's Golf Association. K.W.N.

is supported by G.S. Humane Corp. W.M. Keck Foundation Grant to EAO.

Author contributions. Conceptualization, N.E.B. and K.W.N.; Methodology, N.E.B., K.W.N.,

S.V.B, J.C.N., G.L.H., M.C., R.H.; Investigation, N.E.B., S.S., S.V.B., J.C.N., C.C.N., D.A.S.,

R.H., D.J.K., X.L., O.E., Z.J., T.M.K.; Resources, Z.J., T.M.K.; Writing – Original Draft, K.W.N.,

N.E.B., J.C.N.; Writing–revision, K.W.N., J.C.N., X.L., E.A.O., D.J.K, E.O.; Supervision,

K.W.N., N.E.B., G.L.H., T.M.K., E.A.O., T.I.

Methods

Quantification and Statistical analyses

All statistics were done using GraphPad Prism unless otherwise stated. Random Forest/Boruta

were done with R. For Pearson correlation, the p-value (two-tail) tested the null hypothesis that

the true population correlation coefficient for that pair of variables is zero. Two-tailed unpaired

Student's t-test was done to compare only two groups. One-way or two-way ANOVA was used

for comparisons of more than two groups with correction for multiple hypothesis testing. All

values were expressed as the mean ± S.E.M. The compound injections were done by one person

and the determination of weight/body composition done by another person blinded to the treatment

groups. For the high content imaging assay of myotube diameter and the in cell western blots,

images were visually inspected in a blinded fashion for the presence of intense foci or speckles

and those images were not included in the analyses. For the IL-6 dose curves, data were identified

through the linear regression function in GraphPad Prism

Machine learning

A decision tree-based classification algorithm, *random forest*, can be used to divide the data into

subsets many times to create a forest of many classifications, and determine which classification

best explains the data [21]. This produces a rank order of features (i.e. the assays) that predict the

dependent variable but does not provide a statistical cut-off that indicates which features are

important. The Boruta algorithm resolves this issue by creating shuffled copies of the features and

evaluates the degree to which random forest performs better on the real data compared to the

shuffled data [22].

After scaling the full dataset to Z scores, the *Boruta* package was used to identify predictors

of GR-regulated phenotypes. Each Boruta prediction was performed with three different random

seeds, the results of which were then compiled. Variables that were confirmed in 2 or more runs

were selected as predictors. The minimal or best predictive model was obtained using the

randomForest package. To this end, predictors were added to the model one-by-one, in order of

importance. Each model was assessed by growing 500 decision trees and testing one predictor at

each tree branch. If a predictor improves the predictive power of model, it is retained in the model.

If a predictor fails to improve the model it is discarded, before the next predictor is tested. This

process is repeated until the last predictor is tested. The best model therefore consists of only the

minimal   set   of   predictors.   Sample   *R*   scripts   are   available   at:

https://github.com/jnwachuk/Ligand_Class_Analysis_Reveals_Dissociated_Signaling_Networks

_in_Skeletal_Muscle.

Animals

All animal procedures were approved by the Scripps Research Institute IACUC. Male C57BL/6J

mice, 8–10 wk old, were acquired from the Jackson Laboratories (Bar Harbor, ME) and used for

all conditions. Mice were housed under a 12-h light/dark cycle with *ad libitum* access to food and

water unless otherwise stated. Before all surgical procedures, mice were anesthetized with gaseous

isoflurane (Baxter, USP, USA). Before tissue extraction, the mice were euthanized with CO_2 and

cervical dislocation.

In vivo LPS assay

Mice were treated with 1.5 µg LPS (Sigma L2880) for 1 hr and then blood drawn for determination

of TNFα (ELISA, R&D Systems, MTA00B). Mice were then given an injection of 30 µg LPS for

23 hrs. Body composition was determined using time domain nuclear magnetic resonance (TD-

NMR) in a Bruker minispec.

Lactate tolerance test

Mice were administered GCs at 10 mg/kg IP and then fasted overnight. The next morning

lactate (1.5 g/kg) was injected intraperitoneally and blood glucose assayed at the indicated times

with an Accu-check glucose monitor as described[38].

Electric-pulse-mediated gene transfer and in vivo SUnSET assay

Plasmid DNA (50 μg) was transfected into mouse tibialis anterior (TA) muscles by electroporation

as described previously [39]. 1 hr before electroporation mice were anesthetized, and a small incision

was made through the skin covering the TA muscle and then injected with 30 Al of 0.5 U/Al

hyaluronidase and then injected with plasmid DNA. *Fkbp5* and *Foxo1* expression plasmids were

from the MGC Expression Ready cDNA Library Plasmid DNA was grown in DH5 *Escherichia*

coli, purified with an EndoFree plasmid kit (Qiagen, Valencia, CA, USA), and resuspended in 71

mM sterile PBS. After the injections, electric pulses were administered by using non-invasive

platinum-coated tweezertrodes. Eight 100-ms square-wave electric pulses at a frequency of 1 Hz

and at 200-ms intervals were delivered with an ECM 830 electroporation unit (BTX-Harvard

Apparatus, Holliston, MA, USA) with a field strength of 40 V/cm. After the electroporation

procedure, the incision was closed with Vetbond surgical glue. At 7 d after transfection, muscles

were subjected to nonradioactive *in vivo* measurements of protein synthesis. mice were

anesthetized as described above and then given an intraperitoneal injection of 0.040 mol/g

puromycin dissolved in 100 μl of PBS. At exactly 30 min after injection, tissues were extracted

and frozen in liquid N_2 for WB analysis.

Western blot analysis

Whole cell extracts were prepared in HEPES whole cell lysis buffer (20 mM HEPES, pH 7.4, 1%

TX-100,1x phosphoSTOP complete inhibition cocktail tablets [Roche], and 1X Complete EDTA-

Free protease inhibitor cocktail [Roche]). Frozen muscle tissue was homogenized in lysis buffer

T-PER® Tissue Protein Extraction reagent (Thermo Scientific) by a polytron homogenizer and

sonication. The protein concentration was determined using the protein assay reagent (Bio-Rad).

Sixty to eighty μg of protein was separated by SDS-PAGE and transferred to nitrocellulose

membranes for immunoblotting.

Cell Culture and screening assays

C2C12 myoblasts, L6 myoblasts, and A549 cells (ATCC) were cultured in high glucose DMEM

supplemented with glutamine and 10% FBS. For steroid-free culture conditions, cells were grown

in charcoal:dextran-stripped FBS (Gemini Bio-products, cat no. 100-119).

Optimization of culture conditions for GR ligand assays

Our goal was to develop assays that replicate the in vivo effects of GCs at low nM doses, not the

50-100 μM doses that have been reported as necessary to induce atrophy in vitro. We noticed that

Dex strongly inhibits myoblast proliferation, and promotes fusion into myotubes, and thus we

might expect the test compounds to have differential effects on differentiation, with agonists

producing a more fully developed and mature network of myotubes, which are then subject to GC

mediated atrophy of the myotubes. To avoid such effects on myogenesis, we switched myoblasts

to differentiation media, and then on day 4, added 10 μM cytarabine, a chemotherapy agent that

kills dividing cells, and then adding GCs on day 6 or 7 and completing assays on day 8.

We also hypothesized that GCs would induce atrophy more readily in the normal

physiological context, nutrient deprivation. We optimized the timing of nutrient deprivation and

administration of GCs to obtain the best signal to noise ratio of Dex vs vehicle. We were initially

unable to observe much effect on pAKT, because it was elevated basally. This was optimized by

adding an hr of serum free treatment. Adding an insulin challenge increased the signal to noise in

all the assays and focused the experimental design on GC regulation of the insulin signaling

pathways effects on protein balance and glucose disposal. Variation was further reduced by use of

a BioMek 100-nl pin-tool dispensing robot and using a series of partial media changes to prevent

cells from detaching. To reduce edge effects and media evaporation over the 8 days of the assays

we added a gas permeable membrane to the 384-well plates for all assays except the protein

degradation assay. All assays were performed in C2C12 cells (ATCC) except for the Glut4

translocation assay, which was performed in rat L6 myocytes. This is because the C2C12 have

upregulated Glut1, which is constitutively active, and overwhelms any insulin-induced effects on

Glut4 translocation. Assays were optimized until the they could be repeated with a Pearson

correlation > 0.8 between biological replicates from separate passages of cells.

Optimized protocol

On the day of the experiments cell were seeded at 70% confluency and 48 hrs latter washed with

PBS and switch to differentiation media consisting of phenol-free DMEM supplemented with

glutamine and charcoal stripped serum (Sigma 18l289). Partial media changes were performed

every 48 hrs to prevent lifting of the tubes from the wells. After six days in differentiation media,

the cells were serum starved, treated with ligands and used as described on each particular assay.

348-well plates were used unless otherwise noted, and a gas permeable sealing membrane

(Breathe-Easy membrane) was applied immediately after seeding and replaced after every partial

media change.

Glut4 translocation assay

Low passage, L6 rat myoblast (ATCC), were cultured in high glucose DMEM supplemented with

glutamine and 10% serum (proliferation media). On day 1 of differentiation, 2,500 cells were

seeded on 384-well plates in 50 µl of proliferation media that was replaced every other day. On

day 5 cells were washed and replaced with differentiation media, which was then replaced daily.

After 3 days of differentiation the cells were ligand treated and serum starved overnight. Next day

we replace the medium with 100 µl DMEM ± 1 µM insulin (Sigma I0516) without serum or

glucose and incubate for 1 hour and then fixed with 3.7% formaldehyde for 20 minutes. Non-

permeabilized cell were then washed in TBS and blocked for 2 hours (Rockland MB-070) then

incubated in primary anti-Glut-4 antibody in blocking buffer for 48 hours at 4 °C. Then the plates

were washed 5 times for 5 minutes with TBS-T and then incubated with secondary antibody for 1

hr wash and analyzed using a LI-COR Odyssey imaging system.

SUnSET assay for insulin-mediated protein synthesis

C2C12 myoblasts were cultured and differentiated in 384-well plates as described above. After

differentiation, the different ligands were added with a Biomek pin tool dispenser in serum-free

DMEM overnight, next treated with 100 nM insulin. After 60 min, 1 µM puromycin was added to

all wells, and the cells were incubated for an additional 30 min. The in cell western (LI-COR) was

used to quantitate uptake of puromycin. Cells were fixed with 3.7% formaldehyde for 20 minutes.

Non-permeabilized cell were then washed in TBS and blocked for 2 hours (Rockland MB-070)

then incubated in primary anti-puromycin antibody in blocking buffer for 48 hours at 4 °C. Then

the plates were washed 5 times for 5 minutes with TBS-T and then incubated with secondary

antibody for 1 hr wash and analyzed using a LI-COR Odyssey imaging system.

.

Protein degradation

Rates of protein degradation were determined by measuring the release of TCA-soluble

radioactivity from proteins prelabeled with [3H]-phenylalanine. In brief C2C12 cells were seed in

24 well plates and after completing differentiation, myotubes were labeled with 1.0 µCi/ml of L-

[3,5-3H]- phenylalanine for 48 h dedifferentiation media. Cells were then treated for 24 h with

ligands in DMEM containing 2 mM unlabeled phenylalanine. After treatment, the culture medium

was transferred into a microcentrifuge tube containing 100 µl of bovine serum albumin (10 mg/ml)

and TCA was added to a final concentration of 10% (w/v). Samples were incubated at 4° C for 1

hr followed by centrifugation for 5 min. The supernatant was used for determination of TCA-

soluble radioactivity. The protein precipitates were dissolved with a tissue solubilizer

(Solvable™). Cell monolayers were washed with ice-cold PBS and solubilized with 0.5 M NaOH

containing 0.1% Triton X-100. Radioactivity in the cell monolayer and TCA soluble and insoluble

fractions were measured using a Packard TRI-CARB 1600 TR liquid scintillation analyzer. Protein

degradation was expressed as the percentage protein degraded over the 24 hr period and was

calculated as 100 times the TCA-soluble radioactivity in the medium divided by the TCA-soluble

plus the TCA-insoluble radioactivity in the medium plus the cell layer (i.e., myotube) radioactivity

[40].

High-content imaging and analysis

Primary or C2C12 myotubes or mycocytes in black 96- or 384-well tissue culture plates with clear

bases (Greiner Bio-One, North America, Inc.) were stained for 15 min with 200 nM MitoTracker®

Orange CM-H_2TMRos dye (Invitrogen™ by ThermoFisher Scientific) for imaging of

mitochondrial potential. The cells were rinsed with steroid-free DMEM to remove un-incorporated

dye, and then incubated in the differentiation media for 45 min to reach peak fluorescence. Cells

were fixed in 4% formaldehyde for 20 min, stained with 300 nM DAPI for 5 min, permeabilized

in PBS containing 0.1% Triton X-100 for 20 min, blocked for 1 h with 1x TBS containing 0.1%

Tween-20 (TBS-T) and 2.5% normal goat serum, and incubated at 4 °C overnight with an

AlexaFluor® 488-conjugated anti-skeletal muscle myosin (F59) antibody (Santa Cruz

Biotechnology, Inc. cat no. sc-32732 AF488). The next day, the myotubes were washed 4 times

with TBS-T to remove unbound antibodies and rinsed twice with PBS. The stained myotubes were

imaged at 10–20X magnification on the IN Cell Analyzer 6000 platform (GE Healthcare). For

each treatment condition, a stack of 24–27 images containing an average of 50–100 myotubes per

image were analyzed using the IN Cell Developer Toolbox image analysis software, with a

customized segmentation protocol for myotubes. The average (mean) diameter and mitochondrial

potential (i.e. MitoTracker staining density x area) of myotubes in these images were then

calculated.

Nuclear translocation

The cell line used for these experiments (3617) expresses green fluorescent protein (GFP)-tagged

GR (GFP-GR) from a chromosomal locus under control of the tetracycline-repressible promoter

[41]. Cytoplasm-to-nuclear translocation of the GFP-GR in response to different compounds was

determined as previously described [42]. Cells were plated on a 384-well plate in DMEM medium

containing 10% charcoal stripped serum (Hyclone, Logan, UT) without tetracycline (to allow the

expression of the GFP-GR) at density of 2,500 cells per well. Cells were treated with compounds

or a diluent as control for 30 min, fixed with paraformaldehyde, the nuclei stained with 4′,6-

diamindino-2-phenylindole (DAPI) and Perkin Elmer Opera Image Screening System was used

for fully automated collection of images. An image analysis pipeline was customized using the

Columbus software (PerkinElmer) to automatically segment the nucleus using the DAPI channel

and then construct a ring region (cytoplasm) around the nucleus mask for each cell in the digital

micrographs. Translocation was calculated as a ratio of the mean GFP-GR intensity in nucleus and

cytoplasm, and each value was further normalized to the value for the control (i.e., DMSO) sample

on the same plate.

664   ***Osteogenic Differentiation of Human Mesenchymal Stem Cells***

Human MSCs at passage 2 were maintained in growth media consisted of alpha-MEM (Life

Technologies, 32561--037) supplemented with 17% FBS (Sigma-Aldrich, 12303C), 2mM L-

glutamine (Life Technologies, 25030-081), and 100 units/mL penicillin/streptomycin (Life

Technologies, 15140-122). Effect of glucocorticoid receptor modulators on osteogenic

differentiation of MSCs was quantified using Alizarin Red S (Sigma-Aldrich, A5533) staining.

Briefly, human MSCs were seeded in the middle 8 wells of 48-well plates at plating density of

10,000 cells per well. PBS (Life Technologies, 10010-031) was added to surrounding wells to

reduce evaporation of media in the middle wells during prolonged incubation. 24h later, growth

media was replaced by osteogenic induction media (OIM) consisted of low-glucose DMEM (Life

Technologies, 10567-014) supplemented with 10% FBS (Sigma-Aldrich, 12303C), 50 µg/ml L-

ascorbic-2-phosphate (Sigma-Aldrich, A8960), 10mM β-glycerolphosphate (Sigma-Aldrich,

G9891), and glucocorticoid modulators. OIM containing test compounds was replaced every

week. At the end of 2 weeks, monolayers were washed with PBS (Life Technologies, 10010-031),

incubated in Richard-Allan Scientific™ Neutral Buffered Formalin (10%) (Thermo Scientific,

5725) for 2 h at room temperature, washed with deionized water and stained with Alizarin Red S

for 30 min at room temperature. Monolayers were then rinsed 3 × with deionized water until clear,

stain was then extracted with 10% (w/v) cetylpyridinium chloride (Sigma-Aldrich, C0732) in

10 mM sodium phosphate (Sigma, S5011 and S5136), pH 7.0 for 15 min at room temperature and

the amount of extracted dye quantified spectroscopically at 562 nm. Spectroscopic analysis

performed using a synergy™ NEO HTS Multi-Mode Microplate Reader (BioTek Instruments,

Inc.).

IL-6 secretion

IL-1β (R&D systems) was dissolved in phosphate-buffered saline plus 0.1% bovine serum albumin

(Sigma). A549 cells were plated in DMEM containing charcoal stripped serum in 384 well plates.

The next day, cells were treated for 24 with GCs at 1 μM and then treated with 1 ng/ml IL-1β for

24 hrs. supernatants were collected and IL-6 assayed with AlphaLISA (PerkinElmer) using the

manufacturer's instructions.

Micro array Assay for Real-time Coregulator-Nuclear receptor Interaction (MARCoNI)

This method has been previously described [15]. Assay mixes with 1 nM GST-tagged GR (aa 521-

777, Invitrogen #PV4689), 25 nM Alexa-488 conjugated GST antibody (Fisher Scientific,

#10368552), 5 mM DTT, in Coregulator buffer F (Invitrogen, #PV4547) in the presence of solvent

only (2% DMSO) or 1 μM compound as indicated, were prepared on ice. Experiments were

conducted on a PamStation96 (PamGene) using 2 cycles per minute at room temperature. In short,

PamChip arrays (PamGene #88101) were blocked for 20 cycles with 1% Bovine serum albumin

(Calbiochem #126609) in Tris-buffered saline (TBS), incubated for 40 minutes with 25 μl assay

mix using 3 arrays per mix and finally rinsed twice with TBS. Binding on array (tif) images were

quantified with BioNavigator software (PamGene). A modulation index, i.e. the log-transformed

ratio of binding in presence over absence (solvent) of compound, was calculated for each

interaction. Significance of the modulation was assessed using Student's t-Test and FDR post hoc

correction.

Antibodies

Anti-puromycin antibody: 3RH11 (Kerafast, Inc. cat no. EQ0001), or clone 4G11

(MilliporeSigma, cat no. MABE342). Anti-AMPKα Antibody (Cell Signaling Technology, Inc.

#2532). Phospho-AMPKα (Thr172) (40H9) (Cell Signaling Technology, Inc. #2535), Anti-AKT1

(2H10) antibody (Cell Signaling Technology, Inc. #2967), Anti-pAKT-T308 (C31E5E) antibody

(Cell Signaling Technology, Inc. #2965)

Lentiviral vectors and transduction

Lentiviruses were produced from sets of four mouse piLenti-siRNA-GFP lentiviral vectors

(Applied Biological Materials, Inc. Richmond, BC, Canada) targeting *Pelp1* (cat no.

i029598), *Ncoa2* (cat no. i033105), *Ncoa1* (cat no. i038883), *Ncoa6* (cat no. i042804), *Ncor1* (cat

no. i044954), or *Ncor2* (cat no. i042611), or the Control piLenti-siRNA-GFP lentiviral vector (cat

no. LV015-G). HEK-293T cells in 3x 10 cm dishes per construct, were co-transfected with 10 µg

lentiviral vector, 7.5 µg psPAX2, and 2.5 µg pVSV-G per dish using ProFection® kit (Promega)

according to manufacturer's protocol. At ~15 h post-transfection, the media was replaced with 2.5

ml fresh media. Conditioned media containing lentiviral particles (LVM) were collected every 12

h for another 48 h, pooled, and stored at 4 °C. LVM were passed through a 0.45-micron filter to

remove cell debris, and then centrifuged at 4,000 RPM overnight to pellet the viruses. The pellets

were resuspended in 50 µl sterile PBS and added with 8 µg/ml polybrene to day-1 C2C12

myoblasts.

Figure Legends

Figure 1. Structure-based design approach and glucocorticoid profiling platform.

a) Chemical structure of the dissociated glucocorticoid, PF802.

b) Crystal structure of the dexamethasone (Dex) bound GR ligand-binding domain (LBD). Helix

12 is colored red and the SRC2 coregulator peptide binding in the AF2 binding surface is colored

coral. Carbons-3, -11, and -17 are indicated in the chemical structure (1M2Z.pdb).

c) The bulky dimethylanaline group attached at C11 in RU-486 displaces h12 from the agonist

position to disrupt the AF2 surface and generate antagonism (1NHZ.pdb).

d) Substitutions at C3, as seen with the furoate group in momethasone furoate or the propyne in

RU-486 target a small internal pocket to increase affinity (4P6W.pdb).

e) Substitutions at C3 of the steroidal A-ring enter the solvent channel underneath the AF2 surface,

potentially changing the shape of the surface and the ensemble of interacting coregulators

(3BQD.pdb).

f) Model of SR11466 (**15**) bound to the GR LBD

g) Glucocorticoids used in this study.

h) Compound profiling and computational strategy. Effects on the independent variables were

used in a machine learning approach, Random Forest, to identify predictors of skeletal muscle

phenotypes, the dependent variables, described in **Figure 2** and below.

i) Relationships among GR-mediated phenotypes. Lines indicate significant Pearson correlation

between variables using Bonferroni $p_{Adj} < 0.0071$.

Figure 2. Machine learning reveals best predictors of selective modulation and common

signaling

a) Compounds **1-3**, **6-27** were profiled for regulation of specific target genes in skeletal muscle and GR interaction with specific peptides. Predictors of GR-mediated phenotypes were identified using the random forest classifier[21,22] and statistical thresholds established with the Boruta wrapper algorithm for feature selection[22]. The numbers following the peptide names represent the amino acid beginning of the peptide from the full-length protein.

b) Predictive models for protein synthesis and Glut4 translocation include common and unique predictors (coregulator peptide interactions and target genes), with specific examples shown and significance determined by linear regression. Each point represents the activity of a distinct ligand in the cognate assay.

c) Linear regression was used to compare the predictive power of the indicated peptide interactions for the indicated skeletal muscle phenotypes and target genes.

d) Linear regression was performed for ligand-dependent effects on Glut4 translocation. The predictive power, r^2 of target genes and peptide interactions observed with all compounds (**1-3** and **6-27**) were rank ordered, and then compared to the r^2 observed within compound series modified at either C11 (**13–20**) or C3 (**6-12**) of the steroid scaffold.

e) Linear regression on the effects of the C11 substituted compounds on *Bcl2l1* expression or Glut4 translocation. Each point represents the effects of a separate compound.

See also **Methods**, **Supplementary Dataset**, and **Supplementary Figure S2.**

Figure 3. Functional validation of the predictive target gene, Fkbp5, and GR coregulators

a–c) Empty vector (GFP) control, and either Fkbp5 or Foxo1 expression plasmids were electroporated into contralateral Tibias Anterior (TA) muscles of mice. After 7 days, the mice were fasted overnight and treated with puromycin and insulin for 1 hr the TA muscles collected for

weighting and western blots. N= 3–4.

a) Weight of transduced TA muscles.

b–c) Western blot and quantitation of **b**) puromycin (measuring incorporation into growing

polypetide chains to assay insulin-mediated protein synthesis) or **c**) pAKT in transduced TA

muscles.

d) Effects of Dex on insulin-induced protein synthesis in C2C12 myotubes transduced with

lentivirus expressing the indicated shRNAs. N = 6 wells x 4 shRNAs targeting each gene.

e) Effects of Dex on insulin-induced pAKT in C2C12 myotubes transduced with lentivirus

expressing the indicated shRNAs. N = 6 wells x 4 shRNAs targeting each gene.

Data are mean + S.E.M. See also **Supplementary Figure 3.**

Figure 4. In vitro characterization of two dissociated GR ligands with musculoskeletal

sparing activities.

a) IL-1β-induced IL-6 production by A549 cells treated with the indicated compounds was

compared by AlphaLISA. N = 3.

b–d) Myotubes were assayed as described **Supplementary Figure 1** and **Methods**. N = 4-6

e-f) Mitochondrial potential of C2C12 myoblasts was compared by high content imaging. f)

Representative myoblasts imaged during the assay.

g) The effects of all tested compounds on IL-1β-induced secretion of IL-6 by A549 cells (y axis)

and primary human osteoblast mineralization (x axis). Right, Dose curves of indicated compounds

in the IL-6 assay.

h) Effects of the indicated C3-substituted compounds on IL-6 secretion. N = 3.

Data are mean + SEM. See also **Supplementary Figure 4**

Figure 5. In vivo compound profiling and validation of on target mechanism of action

a) Dex inhibits SR16024-dependent protein synthesis in C2C12 myotubes and right-shifts the dose curve, demonstrating the SR1604 activities are on-target through GR. N = 4

b) RU-486 relieves SR11466-suppressed IL-6 secretion. IL-1β-induced IL-6 production by A549 cells treated with the indicated compounds was compared by AlphaLISA. N = 3

c) Plasma TNFα levels of mice treated with the indicated GR ligands (10 mg/kg Dex or SR16024, or 50 mg/kg SR11466) overnight before a 1-hr LPS challenge of 1.5 mg per mouse. N = 10 mice per group (5 x 2 experiments).

d) Changes in lean mass of mice in panel J were determined by whole-body NMR after an additional 18-hr LPS treatment of 30 mg per mouse. N = 10 mice per group (5 x 2 experiments).

e-f) Changes in the lean mass and body weights of male C57BL/6 mice treated with (10 mg/kg Dex or SR16024, or 50 mg/kg SR11466) and fasted overnight. N = 10 per group (5 x 2 experiments).

g) Dex and SR11466 increased the rate of glucose production, unlike SR16024. Mice were subjected to a lactate-tolerance test (LTT) after overnight fast. N = 5 mice per group.

Data are mean+SEM. See also **Supplementary Figure 5 and Methods.**

Figure 6. Molecular dynamics simulations reveal that C3 substitutions in the steroid scaffold alter allosteric communication between ligand and activity.

a) Ribbon diagram of the GR-LBD illustrates the C-terminus of helix 11 (h11) and N-terminus of h3 where we observed ligand-induced conformational effects in three 1 μs simulations.

b) Distance distribution plots of the Cα distance between Met560 (h3) and Thr739 (h11).

c) Backbone Cα, C', N, and O RMSD distribution plots of the h3 and h11 regions colored magenta

in (**a**).

d) Dynamical network analysis of suboptimal pathways for correlated motion between E755 and

R614. Red arrows indicate pathways found with Dex-bound GR.

e) Histogram showing the suboptimal pathlengths with the indicated ligands.

f) Structures extracted from the Dex-bound simulations showing the R611 in (left) and out (right)

side chain conformations.

g) Distance distribution plots of Q570 (Cδ) and R611 (Cζ) side chain atom distances as a proxy to

determine the relative populations inward R611 conformations that can interact with ligand.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6