Habitat patchiness drives spatial structure in morphological trait variation and co-

variation in spotted salamanders (*Ambystoma maculatum*)

Elizabeth T. Green[1,2], Anthony I. Dell[1,3], John A. Crawford[1], Elizabeth G. Biro[3,4], David R.

Daversa[1*]

[1] National Great Rivers Research and Education Center (NGRREC), East Alton, IL

62024

[2] Middlebury College, Middlebury, VT 05753

[3] Department of Biology, Washington University in St. Louis, St. Louis, MO 63130

[4] Tyson Research Center, Washington University in St. Louis, St. Louis, MO 63130

*Corresponding Author: Dave Daversa, ddaversa@gmail.com, ORCID: 0000-0002-

8984-8897

Author contributions

DRD and ETG designed the study and executed field sampling and trait measurements,

with supervisorial support from AID and JAC. ETG completed the morphometrics and

principle components analyses. DRD performed the rest of the statistical analyses. All

authors contributed to the writing and revision of the manuscript.

Abstract

The influence of intraspecific trait variation in species interactions makes trait-based

approaches critical to understanding eco-evolutionary processes. Given that species

occupy habitats that are patchily distributed in space, advancement in trait-based

ecology hinges on understanding how trait variation is spatially structured across the

landscape. We sampled larval spotted salamanders (*Ambystoma maculatum*) across

spatially discrete ponds to quantify spatial structure in morphology. Spatial structure

explained 7-35% of total observed variation in the length and shape of salamander

larvae, depending on the body segment measured (i.e., head, body, tail). Salamander

tail morphology was more variable and exhibited more spatial structure than head or

body morphology. Salamander mass was also highly variable, and was strongly

correlated with total length. Analysis of allometry revealed that the slopes of mass-length

relationships were similar across space, but the intercepts differed spatially. Preliminary

evidence hints that newly constructed ponds are drivers of spatial differences in

allometric intercepts. Pond construction may therefore bolster diversity in trait co-

variation, and in so doing instil more adaptive potential of salamanders under current

and future environmental change.

Key Words: Intraspecific trait variation, eco-morphology, allometry, phenotypic

plasticity, allometry, amphibian, ecology, morphometrics, conservation

Introduction

Introduction

Individual morphological traits underpin many eco-evolutionary processes. The

range of morphology exhibited by individuals of a species, a form of intraspecific trait

variation (Bolnick et al., 2011), shapes the niche breadth of populations, which in turn

affects their resiliency to environmental disturbances and biological invasions (Tack,

Horns & Laine, 2014). Models of population and community dynamics that incorporate

intraspecific trait variation have therefore become central to ecology (Bolnick et al., 2011;

Moran, Hartig & Bell, 2016). A key outstanding challenge for this 'trait-based' paradigm

is to understand trait variation in a spatially explicit context (Violle et al., 2012; Moran,

Hartig & Bell, 2016).

Species occur in landscapes comprised of spatially discrete habitat patches.

Patchy habitat structure creates networks of spatially isolated groups of individuals,

potentially structuring how traits of those individuals are distributed in space. For

example, environmental or genetic variability across habitat patches may drive group-

level differences in trait expression, especially when movement between patches is

limited. Group-level trait differences produce spatially structured trait variation across

the landscape. Theory predicts that spatial structure in trait variation fundamentally alters

population and community dynamics (Moran, Hartig & Bell, 2016; Banitz, 2019). By

partitioning trait variation among multiple habitat patches, spatial structure potentially

allows for a broader range of adaptive responses to environmental disturbances (Moran,

Hartig & Bell, 2016) and antagonistic interactions (Tack, Horns & Laine, 2014).

Alternatively, spatial structure in trait variation may heighten extinction risk by

geographically isolating trait diversity into specific habitat patches. Thus, distinguishing

how traits are distributed across space, both within and among habitat patches, should

aid with predicting community dynamics under environmental change.

Characterizing spatial structure of trait variation in natural populations is not

straightforward. Within-patch sampling must be comprehensive enough to capture the

range of traits expressed by individuals, and the number of habitat patches sampled

must be extensive enough to capture the spatial structure. Additionally, trait

measurements should be rapidly collected. Otherwise, temporal trait variation arising

from seasonal life histories and ontogeny may confound spatial effects. The studies that

have overcome these logistical challenges have focused primarily on plants (Des

Roches et al., 2018). Empirical data on the spatial structure of trait variation in animals

remains highly limited, particularly for vertebrates.

Even fewer studies have assessed spatial structure in trait co-variation

(Evangelista et al., 2019). Co-variation describes relationships between multiple

individual traits. Measures of trait co-variation are useful for describing species growth

patterns (Hirst, Glazier & Atkinson, 2014), their adaptive constraints (Voje et al., 2014),

and complex morphological characteristics such as body shape (Laughlin & Messier,

2015). Practically, they provide a tool for filling data gaps in ecological models (Madin et

al., 2016). Allometric scaling of mass with body length is a basic form of trait co-variation

that is widely used for this purpose (Madin et al., 2016). Morphometrics integrates co-

variation between body length, depth, and sometimes width, to characterize body

shapes of individuals. Body shape, being a strong proxy of performance and fitness, is

arguably a better predictor of species adaptive responses to environmental change than

linear body measurements (Laughlin & Messier, 2015). Understanding the eco-

evolutionary processes in patchy landscapes therefore requires consideration of spatial

structure in trait co-variation as well.

In this study, we assessed the spatial structure in trait variation and co-variation

in a patchy metapopulation of spotted salamanders (*Ambystoma maculatum*). We

sought to quantify the extent to which habitat patchiness drove spatial structure in

salamander mass and length, as well as allometric and morphometric relationships

among multiple traits. We sampled larval spotted salamanders among a network of

spatially discrete wetlands (i.e. ponds) and measured body length and mass of 519

individuals. We used these data to quantify individual variation in mass and length within

and among ponds. We then examined the spatial structure in variation and co-variation

of those traits by: a) quantifying the contribution of pond-level differences in mass and

length to total observed variation in these traits; b) testing whether mass-length allometry

differed among ponds; and c) quantifying the contribution of pond-level differences in

salamander shape to total observed variation in shape.

Methods

Study species

Spotted salamanders are broadly distributed throughout the Northeastern and

Midwestern United States. These salamanders are semi-terrestrial pond breeders,

annually migrating from terrestrial hibernacula to reproduce in fishless ponds, which in

our study areas occurs between March–April. After hatching from eggs, aquatic larvae

develop and metamorphose in 6–10 weeks. Larvae feed on invertebrates and anuran

tadpoles, and are themselves prey for adult salamanders and larger aquatic

invertebrates such as odonate larvae and beetles (Urban, 2010).

Spotted salamander populations provide useful systems to study morphological

variation in a spatial context because: i) they occupy and move among spatially discrete

ponds that comprise a functional metapopulation with patchy habitat structure (Patrick,

Calhoun & Hunter, 2008); and ii) larval stages exhibit substantial morphological plasticity

in response to heterogeneity in biotic and abiotic conditions (Scott, 1990; Urban, 2010),

which enables a range of trait expressions across individuals and habitat patches.

Field sampling and husbandry

We sampled multiple ponds in east-central Missouri, distributed across three

reserves – Tyson Research Center (800 ha), Forest 44 State Park (400 ha) and Shaw

nature reserve (700 ha) (Fig. 1). We focused on six ponds in which pilot surveys

confirmed salamander larvae were present. Three ponds (Mincke Pond, Arthur Christ

Pond, Beth's Pond) were constructed in 2008 for research purposes and had similar

sizes and dimensions (Burgett 2015). As part of a separate experiment, Rotenone was

applied to Beth's pond in 2008. While this initially reduced microbial biodiversity (Woods

et al., 2016) the micro-organismal community structure had returned to pre-treatment

conditions well before sampling for this study (Woods et al., 2016). The other three focal

ponds (Salamander Pond, Forest 44 Pond, Shaw Pond) were older and variable in size

(Table S1). Salamander Pond was created in 1965. Forest 44 pond and Shaw pond

were excavated between 1990 and 1996 (data extracted from Google Earth Historical

Imagery). All ponds contained multiple predators of larval spotted salamanders:

including dragonfly (*Anax* sp.) and damselfly (*Lestes* sp.) nymphs, diving beetles

(*Dytiscidae* sp.), hyrdrophilid beetles (*Tropisternus* sp), and adult newts (*Notophthalmus*

viridescens) (Tables S1 and S4; Biro unpublished). All ponds were located within

forested habitats typical of temperate deciduous ecosystems found in the Midwestern

U.S. A major highway bisected the study area, separating Forest 44 and Shaw ponds

from the other four ponds.

We intensively sampled one pond per week (30 June–08 August, 2016) by dip-

netting around the perimeter. We dipnetted until 10 minutes passed without a capture to

maximize coverage of morphological variation within ponds. We kept all larvae that did

not show overt signs of injury or illness (e.g. damaged tails or legs, tumorous growth)

and immediately transported them to the National Great Rivers Research and Education

Center (NGRREC) – less than 1 hr drive - where they were housed for seven days

before being returned to their original ponds. At NGRREC, we housed larvae individually

in circular plastic arenas (28 cm diameter) filled with 500 mL dechlorinated water

(approximately 2.5 cm depth). Larvae were maintained at 18 °C with a 14:10 h light:dark

cycle, consistent with ambient conditions at the surveyed ponds. Salamanders were fed

a single gray tree frog (*Hyla chrysoscelis-versicolor*) tadpole on the fifth day of the

housing period. Observing that not all salamanders ate the tree frogs fed to them the

prior day, we tested whether feeding influenced measures of mass for a random subset

of 237 individuals for which feeding data were available (see Supplementary Material for

detailed methods).

Trait measurements

On the sixth day after capture we measured the length and mass of salamanders,

distinguishing between head length, body length, and tail length. We photographed

lateral and dorsal images of larvae placed into clear tanks that minimized movement (Fig.

S1). We blot-dried individuals on paper towels before weighing. We measured the length

of salamander heads, bodies, and tails from images using ImageJ (Fig. S1) (Rasband,

1997). We classified larval development stage from our images according to Harrison,

(1969).

To obtain measurements of the shape of larvae we digitized landmarks on lateral

images using the software *tpsDig2* (Rohlf, 2006). Following Van Buskirk & Schmidt,

(2000) we tagged twenty landmarks that outlined larval shape (Fig. S1). Landmarks 1–3

described the shape of the head of the larvae, landmarks 4–11 described body shape,

and landmarks 9–20 outlined tail shape. Landmarks were rotated, scaled by size, and

aligned to a coordinated system using the Procrustes least-squares superimposition

available in the *geomorph* package for R statistical software (Adams and Otárola-Castillo

2013). We conducted four principal component analyses to explore the scaled two-

dimensional shape variation, again distinguishing head shape, body shape, and tail

shape. The first principal component (PC) score accounted for most of the variation in

head shape (37 %), body shape (33%), and tail shape (33%). We therefore used PC1 as

the shape metric in our analyses.

Data Analysis

179   We calculated the coefficient of variation – the ratio of the standard deviation to

the mean – as a standardized measure of individual morphological variation within

ponds. We assessed spatial structure of variation in salamander length and mass with

generalized linear mixed models (GLMM), using the *lme4* package in R (R Core Team,

2013). Specifically we calculated the intra-class correlation coefficient (ICC) from

GLMMs that included 'pond' as a random intercept term. The ICC, also called the

variance partitioning coefficient, is the proportion of total variation in response variables

that is attributable to group-level differences. In our analysis, the ICC indicated the

proportion of total observed variation in salamander length and mass that came from

pond-level differences in those traits. Trait variation would become increasingly spatially

structured as pond-level differences explained proportionally more of total trait variation,

that is, with higher ICC values. We ran separate GLMMs for mass and the length

measurements – head length, body length, tail length, and total length. All models used

a Gaussian error structure, as all morphological data were normally distributed (Fig. 3).

We omitted developmental stage from the models because all larvae were in the final

Harrison stages (45-46).

195   We then tested for spatial structure in mass-length allometry, a form of trait co-

variation. Specifically, we ran GLMMs to test whether the slopes and intercepts of mass-

length regressions differed across the six focal ponds, again distinguishing head length,

body length, and tail length. GLMMs included mass as the response, length as a fixed

effect, pond as a random intercept term, and length as a random slope term. We log-

transformed both length and mass and used a Gaussian error structure for the

normalized data in all models. To enable convergence of these more complex models,

we multiplied (log-transformed) mass by a factor of 10 to standardize the units with

length measurements. To test for differences in regression intercepts and slopes across

focal ponds, we used likelihood ratio tests comparing the fit of models that included both

terms with models omitting the random intercept or random slope term. We also

calculated the marginal and conditional R^2 of the models using the *MUMIn* package in R.

Marginal R^2 is a measure of the amount of variation in mass that was explained by the

fixed effect of length, whereas conditional R^2 considers variation explained by both fixed

and random effect terms (Johnson, 2014).

To further assess spatial structure in co-variation of morphological traits, we

determined the extent to which pond-level variation in salamander head, body, tail, and

overall (all segments combined) shape contributed to total observed variation in these

multidimensional morphological traits. Again, we calculated the ICC from GLMMs

including 'pond' as a random intercept term. PC1 scores were used as response

variables. The shape data were also normally distributed (Fig. 3), and so we used a

Gaussian error structure for all models.

Results

We obtained measurements of 519 spotted salamander larvae (Forest 44: N =

101, Shaw: N = 88, Salamander pond: N = 65, Arthur Christ: N = 116, Beth's: N = 30,

Mincke: N = 119). Salamander mass was not influenced by their feeding the previous

day in the subset of 227 individuals tested ($F_{(1-82)}$ = 3.43, p = 0.068). Within all ponds,

salamander mass varied more among individuals than any of the length measurements

(Fig. 2). Morphological variation was consistently lower in Beth's pond (Fig. 2), though

this pattern was likely due to the lower sample size. Salamanders in Mincke pond

generally varied more morphologically than salamanders from other ponds (Fig. 2).

Otherwise, there was no indication that specific ponds harbored most of the

morphological variation (Figs. 2 and 3).

Differences in average trait values across ponds accounted for 7–35% of the

total observed variation in salamander mass and length, depending on the specific body

section measured (Table 1). Specifically, pond-level differences accounted for

proportionally more of the observed variation in salamander mass (35%), total length

(35%), and tail length (27%) than in head (11%) and body length (17%) (Table 1).

Salamander mass co-varied most strongly with total length, which makes sense

because we took single measures of mass that incorporated all body segments. Mass

also co-varied strongly with body and tail lengths, but was associated less with head

lengths, likely because heads are the smallest body segment of salamanders (Fig. S1).

Accounting for pond-level differences in the intercepts of mass-length relationships

improved explanations of variance in mass (Table 2), resulting in detectable differences

in intercepts of mass-length relationships across ponds in terms of their intercepts

(including pond as a random intercept term improved model fit; mass-head length: X^2_1 =

111.84, p < 0.001; mass-body length: X^2_1 = 141.09, p < 0.001; mass-tail length: X^2_2 =

39.089, p < 0.001; total length: X^2_1 = 81.136, p < 0.001; Fig. 4). For a given length,

individuals from Salamander pond tended to be heavier than individuals from other

ponds (Fig. 4) whereas individuals form Beth's pond were generally lighter in mass per

unit length (Fig. 4). Pond-level differences in the slopes of mass-length relationships

played a lesser role in explaining variation in salamander mass (Table 2). Only the

slopes for mass-head length and mass-tail length relationships differed across ponds

(mass-head length: X^2_2 = 8.68, p = 0.013; mass-tail length: X^2_2 = 11.25, p = 0.004; Fig.

4); the slopes of mass-body length and mass-total length relationships were consistent

across sampled ponds (mass-body length: X^2_2 4.24, p = 0.120; mass-total length: X^2_2 =

5.05, p = 0.080; Fig. 3). The slope exponents were always < 3 (Table S3), indicating

that larger salamander larvae generally had more elongate heads, bodies and tails than

smaller larvae.

Similar to length measures, the shape of salamander tails exhibited more spatial

structure than did the heads or bodies. Pond-level trait differences contributed 25% of

the total observed variation in tail shape, compared with 11%, 9%, and 7% of head, body,

and overall shape, respectively (Table 1). Further, there was little evidence that PC

scores were clustered by pond (Fig. 5), indicating a weak signal of spatial structure in

salamander shapes.

Discussion

There was a stronger signal of spatial structure in larval salamander tails

compared with their heads or bodies. Spatial structure in trait co-variation was also

evident in salamander tails. Scaling of mass with salamander tail lengths differed across

ponds both in terms of the intercepts and the slopes of the relationships, and pond-level

differences contributed more to total observed variation in tail shapes than for head or

body shapes. The spatial structure of pond habitats therefore seems to act particularly

strongly on salamander tails, perhaps because tails play an important role in swimming

performance. Being meso-predators, swimming performance for salamanders is critical

to both capturing prey and evading predators (Van Buskirk & Schmidt, 2000; Urban,

2010). Tails may therefore be more closely linked to fitness, hence under stronger

selection, than heads and bodies, at least in habitats where predation is a significant

threat. Multiple predators of larval salamanders were found in our focal ponds, and so

predation risk is likely to be a strong selective force in the salamander metapopulation

studied here. In a preliminary analysis of the data, we did not detect an influence of

predator density on tail length of salamanders (see Supplementary Material for detailed

methods and results), but this analysis was based on our limited sample of ponds and

warrants further investigation. Regardless of the factors driving salamander tail variation,

our findings suggest that salamander responses to habitat alterations, biological

invasions, and other pond-level disturbances may manifest as changes in tail

morphology as opposed to changes in head and body morphology. If this prediction

holds, the more common body size measure for amphibians, snout-to-vent length, which

only takes head and body lengths into account, would be insufficient for predicting the

eco-evolutionary responses of this species to landscape-scale environmental changes in

aquatic habitats.

Even with respect to salamander tails, much of total morphological variation that

we observed was unstructured (i.e. random). The substantial within-pond variation in

salamander morphology suggests that local factors, such as microhabitat heterogeneity,

also influence salamander morphology and mediate the extent to which habitat

patchiness spatially structures morphological variation. Given that many ponds were

spaced within the documented dispersal ranges of salamanders (Zamudio & Wieczorek,

2006; Patrick, Calhoun & Hunter, 2008), movement between ponds could also have

mediated spatial structure in morphological variation by sustaining mixing of genotypes

and phenotypes. At the metapopulation scale, local and spatial factors likely interact to

shape varying degrees of spatial structure in morphological variation like that we

observed in salamanders. This points out that spatial structure in morphological

variation occurs along a continuum rather than in a binary, structured versus

unstructured, fashion. The impact of trait variation on population and community

dynamics likely depends the degree to which traits are spatially structured. This

argument is supported by recent modelling (Banitz, 2019) and calls for empirical testing

of how population and community dynamics change as trait variation becomes

increasingly spatially structured by habitat patchiness.

Spatial structure in the allometric relationship between salamander mass and

total length arose specifically from pond-level differences in intercepts; slopes of the

relationship were highly consistent. Spatial pond structure therefore appears to drive

spatial differences in salamander mass relative to total body length, but does not appear

to alter how mass scales with total body length. This spatially robust scaling of mass and

total body length may explain why the two traits separately exhibited nearly identical

degrees of spatial structure. More broadly, this pattern of allometric scaling, in which

intercepts but not slopes of trait relationships differ, is consistent with allometric

relationships documented across many other taxa (Voje et al., 2014), suggesting a

general constraint to the plasticity and evolution of the slopes of trait relationships.

The inclusion of morphological diversity data in biodiversity conservation stems

from the idea that different populations of the same species are not equal in terms of

eco-evolutionary history. As such, exploring various approaches to the conservation of

morphological diversity is important to developing strategies for reducing biodiversity

losses under global change (Des Roches et al., 2018). The mix of within- and between-

pond morphological variation in salamanders provides promise that pond construction

can utilize local and spatial processes to bolster morphological diversity. Capitalizing on

the presence of new constructed ponds in our study area, we made a preliminary

comparison of salamander morphology and allometry between 'new' and 'old ponds'

(see Supplementary Material for detailed methods and results). This analysis did not

detect differences in mass or length of salamanders between 'new' and 'old' ponds, but

pond age did influence mass-length relationships and body shapes (Supplementary

Material Table S4). Habitat restoration through pond construction may then bolster

diversity in trait co-variation, and in so doing may instill more adaptive potential under

environmental change (Laughlin & Messier, 2015). Although there are a number of

studies for various taxa that quantify functional connectivity between habitat patches

(and local populations) using genetic techniques, we encourage additional studies on

morphological parameters and patterns to better understand the mechanisms that

promote long-term population persistence in fragmented landscapes.

Acknowledgements

We thank members of the Tyson Research Station for their support of our field sampling,

John Grady for assistance with data analyses and visualization, Julie Messier for

assistance with the statistical analyses, and S. Trombulak for assistance with data

analyses and comments that significantly improved this manuscript. This project was

conducted in accordance with University of Illinois IACUC #16203, and funded in part by

the NGRREC intern program. None of the authors experienced conflict of interest that

could have influenced the objectivity of this study.

Conflicts of interest

We declare not conflicts of interest

References

Banitz T. 2019. Spatially structured intraspecific trait variation can foster biodiversity in

disturbed, heterogeneous environments. *Oikos* 128:1478–1491. DOI:

10.1111/oik.05787.

Bolnick DI, Amarasekare P, Araújo MS, Bürger R, Levine JM, Novak M, Rudolf VHW,

Schreiber SJ, Urban MC, Vasseur DA. 2011. Why intraspecific trait variation matters

in community ecology. *Trends in Ecology & Evolution* 26:183–192. DOI:

10.1016/j.tree.2011.01.009.

Des Roches S, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, Schweitzer JA,

Palkovacs EP. 2018. The ecological importance of intraspecific variation. *Nature*

Ecology & Evolution 2:57–64. DOI: 10.1038/s41559-017-0402-5.

Evangelista C, Olden JD, Lecerf A, Cucherousset J. 2019. Scale-dependent patterns of

intraspecific trait variations in two globally invasive species. *Oecologia* 189:1083–

1094. DOI: 10.1007/s00442-019-04374-4.

Harrison RG. 1969. Harrison stages and description of the normal development of the

spotted salamander, *Ambystoma punctatum* (Linn.). *Organization and Development*

of the Embryo 1:44–66.

Hirst AG, Glazier DS, Atkinson D. 2014. Body shape shifting during growth permits tests

that distinguish between competing geometric theories of metabolic scaling. *Ecology*

Letters 17:1274–1281. DOI: 10.1111/ele.12334.

Johnson PCD. 2014. Extension of Nakagawa & Schielzeth's R^2_{GLMM} to random slopes

models. *Methods in Ecology and Evolution* 5:944–946. DOI: 10.1111/2041-

210X.12225.

Laughlin DC, Messier J. 2015. Fitness of multidimensional phenotypes in dynamic

adaptive landscapes. *Trends in Ecology & Evolution* 30:487–496. DOI:

10.1016/j.tree.2015.06.003.

Madin JS, Hoogenboom MO, Connolly SR, Darling ES, Falster DS, Huang D, Keith SA,

Mizerek T, Pandolfi JM, Putnam HM, Baird AH. 2016. A trait-based approach to

ddvance coral reef science. *Trends in Ecology & Evolution* 31:419–428. DOI:

10.1016/j.tree.2016.02.012.

Moran EV, Hartig F, Bell DM. 2016. Intraspecific trait variation across scales:

implications for understanding global change responses. *Global Change Biology*

22:137–150. DOI: 10.1111/gcb.13000.

Patrick DA, Calhoun AJK, Hunter ML. 2008. The importance of understanding spatial

population structure when evaluating the effects of silviculture on spotted

salamanders *(Ambystoma maculatum)*. *Biological Conservation* 141:807–814. DOI:

10.1016/j.biocon.2007.12.026.

R Core Team. 2013. *R: A language and environment for statistical computing*. Vienna,

Austria: R Foundation for Statistical Computing.

Rasband WS. 1997. ImageJ. US National Institutes of Health, Bethesda, MD, USA.

Rohlf FJ. 2006. tpsDig, version 2.10. *http://life. bio. sunysb. edu/morph/index. html*.

Scott DE. 1990. Effects of Larval Density in Ambystoma Opacum: An Experiment Large-

Scale Field Enclosures. *Ecology* 71:296. DOI: 10.2307/1940269.

Tack AJM, Horns F, Laine A-L. 2014. The impact of spatial scale and habitat

configuration on patterns of trait variation and local adaptation in a wild plant parasite.

Evolution 68:176–189. DOI: 10.1111/evo.12239.

Urban MC. 2010. Microgeographic adaptations of spotted salamander morphological

defenses in response to a predaceous salamander and beetle. *Oikos* 119:646–658.

DOI: 10.1111/j.1600-0706.2009.17970.x.

Van Buskirk J, Schmidt BR. 2000. Predator-induced phenotypic plasticity in larval newts:

trade-offs, selection, and variation in nature. *Ecology* 81:3009–3028.

Violle C, Enquist BJ, McGill BJ, Jiang L, Albert CH, Hulshof C, Jung V, Messier J. 2012.

The return of the variance: intraspecific variability in community ecology. *Trends in*

Ecology & Evolution 27:244–252. DOI: 10.1016/j.tree.2011.11.014.

Voje KL, Hansen TF, Egset CK, Bolstad GH, Pélabon C. 2014. Allometric constraints

and the evolution of allometry. *Evolution* 68:866–885. DOI: 10.1111/evo.12312.

Woods LM, Biro EG, Yang M, Smith KG. 2016. Does regional diversity recover after

disturbance? A field experiment in constructed ponds. *PeerJ* 4:e2455. DOI:

10.7717/peerj.2455.

Zamudio KR, Wieczorek AM. 2006. Fine-scale spatial genetic structure and dispersal

among spotted salamander *(Ambystoma maculatum)* breeding populations. *Molecular*

Ecology 16:257–274. DOI: 10.1111/j.1365-294X.2006.03139.x.

Tables and Figures

body segment	pond-level variance	SD	residual variance	SD	ICC (%)	95% CI - lower	95% CI - upper
mass							
	0.013	0.112	0.023	0.152	35.3	8.62	58.71
length							
head	0.15	0.39	1.27	1.13	10.8	0.73	26.11
body	0.92	0.96	4.36	2.09	17.4	8.72	53.62
tail	4.27	2.07	11.29	3.36	27.5	4.08	52.07
combined	18.23	4.27	32.91	5.74	35.6	6.25	57.80
shape							
	pond-level variance	SD	residual variance	SD	ICC	95% CI - lower	95% CI - upper
head	0.0004	0.019	0.003	0.053	11.0	0.7	26.5
body	0.0002	0.016	0.002	0.049	9.4	0.3	23.6
tail	0.001	0.026	0.002	0.049	22.8	1.2	35.0
combined	0.000	0.009	0.001	0.033	7.60	0.2	17.7

Table 1. Proportion of mass, length, and shape variation attributable to pond-level differences. Displayed are the variance components of generalized linear mixed models used for our analyses of spatial structure in salamander mass, length, and shape. The intra-class correlation coefficient (ICC), or the variance partitioning component, is the proportion of pond-level variation explaining total observed trait variation (pond-level variance + residual variance). Higher ICC values denote higher degree of spatial structure in salamander trait variation. CI = confidence intervals of the ICC derived from bootstrapping over 500 resampling events.

mass-length co-variation								
body segment	intercept variance	SD	slope variance	SD	residual variance	SD	marginal R^2	conditional R^2
head	0.11	0.33	0.09	0.29	0.02	0.13	0.26	0.49
body	0.04	0.20	0.02	0.13	0.01	0.09	0.65	0.77
tail	0.08	0.28	0.04	0.19	0.01	0.08	0.73	0.79
combined	0.07	0.26	0.00	0.07	0.00	0.07	0.81	0.87

Table 2. Variation in intercepts and slopes of mass-length relationships.

Displayed are the variance components of generalized linear mixed models used for our

analyses of spatial structure in the scaling of salamander mass with different length

measurements. Models included length (log-transformed) as a fixed effect and a random

slope term, and pond as a random intercept term. Marginal R^2 denotes the amount of

variation in mass explained by the length measurement alone, whereas conditional R^2

considers variation explained by the random intercept and slope terms.

437
438 **Figure 1. Map of study area.** The six focal ponds were located in Eastern Missouri

439 (U.S.A.) across three conservation areas. Mincke Pond, Arthur Christ Pond, and Beth's

440 Pond were located in Tyson Research Center. Shaw pond was located in the Shaw

441 Nature Reserve. Forest 44 pond was located in Forest 44 Conservation Area. All ponds

442 occurred in Oak-Hickory forests typical of the region.

443
Fig. 2. Within-pond variation in salamander morphology. The coefficient of variation,

or the extent of variance in relation to mean trait values, is shown for **(a)** mass and

length measures and **(b)** the six ponds where we sampled salamanders, ordered from

left to right in the chronological order in which they were sampled. Colors distinguish

pond of capture in (a) and the focal trait in (b). Note that the same data are reported in (a)

and (b).

Fig. 3. Salamander mass and length variation within and between ponds. Panels on

left **(a,c,e,g,i)** are frequency distributions of mass and length measurements across all

focal ponds. Violin plots on the right-side panels **(b,d,f,h,j)** distinguish individual-level

and pond-level morphological variation to show how the traits were spatially structured.

Box plots within the violin plots denote the mean, standard error, and 95% confidence

intervals of trait measures. Ponds in the right-side panels are ordered from left to right in

the chonological order in which they were sampled.

Fig. 4. Spatial structure in mass-length allometry in salamanders. Regression lines

of salamander mass with (a) head length, (b) body length, (c) tail length, and (d) are

shown for the six focal ponds, distinguished by line colors. Shaded areas show the 95%

confidence intervals of the regression lines. Mass and length are plotted on a $\log_{10}$ scale

in all cases.

Fig. 5. Salamander shape variation within and among ponds. The shape values are

based on sets of landmarks at different points along the lateral surface of salamander

bodies (a). The overall (b), head (c), body (d), and tail (e) shape of salamanders

collected from the six focal ponds. The shape values are based on sets of landmarks at

different points along the lateral surface of salamander bodies. PC1 and PC2 values

increase with elongation of shape and increasing length:height ratio.

485