Citrulline supplementation improves spatial memory in a murine model for Alzheimer's disease

Katia Martínez-González[a,b], Leonor Serrano-Cuevas[c], Eduardo Almeida-Gutiérrez[d],

Salvador Flores-Chavez[e], Juan Manuel Mejía-Aranguré[f], Paola Garcia-delaTorre[a]*

[a] Unidad de Investigación Médica en Enfermedades Neurológicas, Hospital de Especialidades, Centro Médico Nacional Siglo XXI, Instituto Mexicano del Seguro Social, México

[b] Posgrado en Ciencias Biológicas, Unidad de Posgrado, Edificio A, 1° Piso, Circuito de Posgrados, Ciudad Universitaria, Coyoacán, C.P. 04510, Distrito Federal, México, Universidad Nacional Autónoma de México.

[c] Coordinación de Unidades Médicas, División de Evaluación y Rendición de Cuentas de los Procesos de Atención Médica en Unidades Médicas de Alta Especialidad, Instituto Mexicano del Seguro Social, México

[d] Head of Medical Education and Research, Hospital de Cardiología, Centro Médico Nacional Siglo XXI, Instituto Mexicano del Seguro Social, México

[e] Unidad de Investigación Médica en Nutrición, Hospital de Pediatría, Centro Médico Nacional Siglo XXI, Instituto Mexicano del Seguro Social, México

[f] Coordinación de Investigación Científica, Instituto Mexicano del Seguro Social, México

*Corresponding author

Cuauhtémoc 330, Col. Doctores 06720, Ciudad de México, México

pgarciatorre@gmail.com

ABSTRACT

INTRODUCTION: Alzheimer's disease (AD) correlates with the dysfunction of metabolic pathways that translates into neurological symptoms. An arginine deficiency, a precursor of nitric oxide (NO), has been reported for AD patients. We aimed to evaluate the effect of citrulline oral supplementation on cognitive decline in an AD murine model.

METHODS: Three-month citrulline or water supplementation was blindly given to male and female wild type and 3xTg-AD mice trained and tested in the Morris Water Maze (MWM). Cerebrospinal fluid and brain tissue were collected. Ultra Performance Liquid Chromatography was used for arginine determinations while the Griess method was used for NO.

RESULTS: Eight-month-old male 3xTg-AD mice supplemented with citrulline performed significantly better in the MWM task; arginine levels increased in cerebrospinal fluid although no changes were seen in brain tissue and only a tendency of increase of NO was observed.

DISCUSSION: Citrulline oral administration is a viable treatment for memory improvement in the early stages of AD pointing to NO as a viable, efficient target for memory dysfunction in AD.

Keywords: nitric oxide, arginine, dementia, water maze, 3xTg-AD, hippocampus, nutritional supplementation

1. BACKGROUND

Alzheimer's disease (AD) is the most common cause of dementia and consists of progressive neurodegeneration characterized by initial memory impairment and cognitive decline [1]. Most studies are focused on amyloid and tangle pathologies, which are specific to AD, and although its precise etiology is still unclear, cardiovascular risk factors such as hypertension, hypercholesterolemia, diabetes mellitus, aging, and sedentary lifestyle are associated with a higher incidence of AD [2]. At this time, more and more evidence suggests that chronic brain hypoperfusion and cerebrovascular pathology are the earliest clinical markers of AD [3].

In an attempt to shed novel insight into its etiology, one alternative is to view Alzheimer's disease as a metabolic disease. Some studies support the hypothesis that AD correlates with the dysfunction of metabolic pathways that are translated into neurological symptoms [4]. In this regard, changes in bioenergetics are part of the normal aging process and even when these anomalies are body-wide, they affect the brain most substantially because of its exceptionally high-energy requirements [5]. In AD, changes in the glycolysis pathway, a decrease in oxidative metabolism, diminished mitochondrial potential, defective or reduced mitochondria, and IGF-1/insulin resistance have been reported [6].

In general, almost all amino acids and ammonia levels are significantly lower in AD patients when compared with controls, arginine being one of the most relevant since a reduction in this amino acid can affect the citrulline/nitric oxide (NO) pathway [7, 8]. Arginine and citrulline concentrations are slightly lower in CSF from

possible AD patients [9,10], but arginine has also been found to be lower in diagnosed AD patients [11]. However, there are some studies in which arginine levels in AD did not show any changes [12] or have been found to be elevated compared to mild cognitive impairment patients [13].

NO has been shown to be biologically active in the central nervous system and is involved in learning processes and cognitive function [14] as it helps to regulate synaptic plasticity [15] and contributes to long-term potentiation and long-term depression [16]. Deficiency of NO has been implicated in neurodegeneration by promoting endothelial dysfunction, accelerating formation and accumulation of amyloid peptides, reducing synaptic plasticity, activating microglia, and evoking neuroinflammation [15,17].

Citrulline is a precursor of L-Arginine (L-Arg) which is oxidized to NO and Citrulline by the action of endothelial NO synthase (NOS). It has been reported that exogenous administration of L-Arg restores NO bioavailability [18] and may offer a therapeutic strategy for controlling NO metabolism disorders and improving cardiovascular function [19].

On the matter, citrulline supplementation for 3 months reduced age-related hippocampus raft changes, resulting in tightly close raft patterns to those in young animals. All citrulline treated rats had low levels of amyloid protein precursor (APP) and low levels of C99-APP-Cter, the first-step fragment of amyloidogenic APP [20].

Our aim in this study was to evaluate the effect of citrulline oral supplementation on cognitive decline in the progression of Alzheimer's disease in a murine model and a possible way in which citrulline could increase nitric oxide levels.

2. METHODS

2.1 Animals

All procedures were performed in accordance with the current rulings in Mexican law (NOM-062-ZOO-1999) and with the approval of the local Science and Ethics Committees (R-2012-785-049).

Homozygous male and female 3xTg-AD (3xTg-AD; B6; 129-Psen1 <tm1Mpm> Tg (APPSwe, tauP301L) 1Lfa) mice were used as an AD murine model, and B6129SF2/J WT male and female mice were used as controls since this is the genetic background originally described for the transgenic mice used. Mice were bred in our lab from parents purchased from The Jackson Laboratory, and housed in the *Bioterio de la Coordinación de Investigación en Salud*. Mice were housed in plastic cages, wood chip bedding, and metal wire covers in groups of 2-4 same-sex littermates. They were housed in a colony room maintained at 12:12 light/dark cycle at 20-22ºC and provided with water and food *ad libitum*. All animals were genotyped using polymerase chain reaction of the tail tissue samples by Hot Shot Tail DNA preparation.

2.2 Citrulline supplementation

Mice from each strain and age (5 or 9 months old) were randomized into two groups, one control and one experimental. Mice from the experimental group received 1g/kg of malate citrulline dissolved in water fed orally with an esophageal cannula stainless steel, semi-curved 30° with Roma tip, and a length of 80 mm, once daily for 12 weeks. The control group was fed with the same volume of water

according to the mouse's weight.

2.3 Memory task

Following citrulline or placebo supplementation, learning and memory skills were tested using the Morris Water Maze (MWM) modified from Bromley-Brits et al. [21]. The MWM consists of a circular tank (150 cm in diameter, 50 cm in height) filled with opaque water. A 14cm x 14 cm platform submerged 1.5 to 2 cm below the water surface was placed inside the tank. Throughout the experiment, the temperature of the pool was 22 ± 1 ºC and the location of the platform was maintained constant.

The task consisted of four trials per day for 4 consecutive days. Trials were assigned with pseudorandom starting points (E, SE, NW, N) each day. A total of 16 trials were performed facing the wall at one of the four starting positions and allowed to locate the submerged platform. Mice were allowed to swim freely until they found the platform and the time required to find the platform (escape latency) was recorded. Mice were allowed to stay on the platform for 10 more seconds prior to their removal to the rest cage. If a mouse was unable to locate the platform within 60 s, it was gently guided to the platform and allowed to stay on it for 10 s and its escape latency was recorded as 60 s.

Twenty four hours after the last trial, a 60 s probe trial was performed without the escape platform. Mice were released into the tank individually and allowed to swim for 60 s. Latency, time spent in the target quadrat, and crosses were recorded.

2.4 Cerebrospinal fluid (CSF) collection

Two days after probe-day, 8-month-old animals were anesthetized with pentobarbital and NaCl 0.9% (50 mg/kg) administered intraperitoneally. The neck was shaved, and CSF samples were taken from the cisterna magna by puncture, the volume obtained was approximately 4ul per animal; a pull of four mice's CSF was used to measure arginine levels. The whole procedure took 10 min per mouse (including anesthesia).

2.5 Tissue collection

Mice were decapitated by a guillotine, the brain was carefully removed and tissue from the hippocampus was dissected, frozen in dry ice, and stored at -80ºC.

2.6 Arginine determination

Ultra Performance Liquid Chromatography (UPLC) analyses were performed by a modified method based on Ivanova [22] and Jong [23] updated for Waters UPLC®. Briefly, homogenates from the hippocampus in saline solution (100 ul) were diluted in 100 μl PBS buffer (NaH_2PO_4 10mM, NaCl 140mM, pH 8.0), centrifuged to precipitate and separate debris, and supernatants were saved and frozen to -70 °C for further analysis. A stock solution of derivatization reagent was prepared by dissolving 10 mg of o-phthalaldehyde in 0.2 ml methanol followed by the addition of 1.8 ml of potassium-borate buffer (pH 9.5) and 10 μl 3-mercaptopropionic acid. The working solution was prepared by two-fold dilution of the stock solution with a borate buffer.

Samples were extracted by solid-phase extraction (SPE) on 1-ml Waters Oasis MCX SPE® columns. The SPE columns were loaded with a mixture of 25 μl of

PBS-extracted tissue, 175 µl water, 100 µl of internal standard (10 µmol/L monomethyl arginine dissolved in HCl 10 mM), 100 µl HCl 10 mM and 600 µl PBS buffer. The dried residue was dissolved on 50 µl working solution of derivatization reagent. After completion of the reaction (2 to 3 minutes), the reaction was stopped by the addition of 50 µl of 0.2 M KH_2PO_4.

Chromatographic separations were performed on Waters UPLC® Accquity system using an Accquity C-18 BEH UPLC column 1.7 µm particle size protected with a guard pre-column filter. Fluorescence was monitored at λ excitation 340 nm and λ emission 455 nm. Arginine was clearly separated with baseline resolution between signals; retention times were 2.5 to 2.8 min (max. peak signal).

2.7 Nitric oxide determination

The method used here, developed by Miranda et al., consists of the vanadium (III)-dependent reduction of nitrate to nitrite, and the spectrophotometric detection of nitrite by means of the Griess reaction in a single step. A rapid, simple spectrophotometer method for simultaneous detection of nitrate and nitrite [24].

Tissue homogenate was deproteinized by filtration in Microcon YM-50 and centrifugated at 12000 rpm during 40 minutes at 20º C, 50ul were obtained and 100ul of deionized H2O, 150 ul of 0.8% VCL3 in 1N HCl, 150 ul of 2% sulfanilic acid, and 0.1% NED were consecutively added. Samples were incubated at 37º C for 60 minutes and centrifuged at 9000 rpm for 10 minutes. The supernatant was measured at 545 nm in a monochromator-based optic Spectrophotometer EPOCH (controlled with the Gen5 Software interface). A standard curve was done by serial

dilutions of NaNO3 100uM.

2.8 Statistical analysis

An ANOVA for repeated measures was used to analyze Morris Water Maze training and a Fisher *post hoc* was then used to detect differences between groups. A Kruskal Wallis test was used to analyze latency time in probe trial test, a *post hoc* of Dunn's was then used to detect differences between strains and a Mann Whitney was used to compare groups (treatment, sex, age). A Kruskal Wallis test was used to analyze arginine levels in tissue with all the groups, and a *post hoc* if Dunn's analysis was then used to detect differences between groups. Finally, a Spearmen's r test was used to evaluate the correlation between arginine and ON levels, and arginine and memory performance in both strains.

3. RESULTS

3.1 Memory improvement with citrulline supplementation

After a three-month period of citrulline or water administration, animals were trained and tested in the Morris Water Maze task. During training, all groups in the 8 month-old group, reduced their escape latency (time to reach the platform) as shown by an ANOVA with repeated measures with an effect on trials ($F_{(15,35)}$ = 10.93, $p < 0.001$; $F_{(15,27)}$ = 14.30, $p < 0.001$) but no effect of strain ($F_{(1,525)}$ = 1.60, $p > 0.05$; $F_{(1,405)}$ = 0.85, $p > 0.05$) or treatment ($F_{(1,525)}$ = 0.92, $p > 0.05$; $F_{(1,405)}$ = 0.80, $p > 0.05$) for male and female mice respectively (Fig.1a and 1b).

Figure 1. Morris Water Maze training. Latency (time in seconds) is presented for transgenic and wild type strains; dotted lines represent water treatment and solid lines citrulline supplementation. Letter a) represents 8-month-old males and b) 8-month-old females. *p < 0.05. TG: transgenic, WT: wild type

A Mann Whitney test showed differences in latency time on probe day (24 h after the last trial) between female (Mdn= 6.61) and male transgenic mice (Mdn= 33.54), (U = 0, p < 0.05). Since differences between sex were found, every subsequent analysis was done separately.

On probe day, a Kruskal Wallis test showed significant differences between male groups (H(3) = 14.45, p < 0.05), and a Dunn's *post hoc* test showed differences between transgenic water-treated mice (higher latency time) and wild type water-treated mice (Z = 3.74; p < 0.05) showing that transgenic mice had cognitive impairment at the time of the test. A Mann Whitney U test (Table 1) showed that transgenic mice that received citrulline supplementation had a shorter escape latency than water-treated mice and similar to water-treated wild type mice, suggesting an improvement in the memory test due to citrulline supplementation. No significant differences were found between treatments for wild type male mice

On the other hand, a Kruskal Wallis test for female mice strains showed no statistically significant difference of escape latency (H(3)= 3.28, p > 0.05) as can been seen in Figure 2b, suggesting that female transgenic mice do not show memory deficit in this task compared to its control group. Moreover, citrulline supplementation had no effect on wild type or transgenic female mice (Table 1).

Figure 2. Morris Water Maze Probe for 8-month-old mice. Latency (seconds) is presented for wild type (grey) and transgenic (black) mice. Letter a) represents males and b) females. *p < 0.05. TG: transgenic, WT: wild type

We then tested citrulline supplementation in nine-month-old mice; after a three-month period of citrulline or water administration, animals were trained and tested in the MWM task. During training, all groups in the 12 month-old group, reduced their escape latency (time to reach the platform) as shown by an ANOVA with repeated measures with an effect on trials (F(11, 14)= 4.40 , p < .001; F(11, 14)= 4.04 , p < 0.001); an effect of strain in males (F(1,154)= 10.49, p < 0.05) but not in

females (F(1,154)= 1.40, p > 0.05); and no effect of treatment for male and female mice respectively (F(1,154)= 0.11, p > 0.05 ; F(1,154)= 0.44, p > 0.05).

As can been seen in Figure 3a significant differences in latency for probe day between 3xTgAD and wild type water-treated males were found. A Kruskal Wallis test (H(3)= 8.45, p < 0.05) and a Dunn's *post hoc* test showed differences between transgenic water-treated mice (higher latency time) and wild type water-treated mice (Z = 2.57; p < 0.05) showing that transgenic mice had cognitive impairment at the time of the test. However, a Mann Whitney U test showed no effect between treatments on either strain as shown in Table 1.

Finally, a Kruskal Wallis test for female mice strains showed no statistical differences on escape latency (H(3) = 2.73, p > 0.05) as can been seen in figure 3b, suggesting again that transgenic female mice do not show memory deficit in this task compared to its control group. Since no changes were found on the 12-month-old groups, no further experiments were performed for this age group.

Figure 3. Morris Water Maze Probe for 12-month-old mice. Latency (seconds) is presented for wild type (grey) and transgenic (black) mice. Letter a) represents males and b) females. *p < 0.05. TG: transgenic, WT: wild type

Table 1. Mann Whitney U test for water maze probe day divided by treatment, strain, and age group.

		Treatment				U	P
		Citrulline		**Water**			
		Mean	SD	Mean	SD		
Male 8-month-old	Wid type	14.11	9.06	7.36	3.26	11	0.18
	3xTg-AD	30.8	11.68	16.04	5.53	10	**0.01**
Female 8-month-old	Wid type	8.45	4.61	10.53	4.23	16	0.31
	3xTg-AD	7.3	4.22	7.72	5.03	14.5	0.66
Male 12-month-old	Wid type	5.75	3.09	6.25	1.7	6	0.65
	3xTg-AD	20.5	11.21	12.87	7.35	5	0.48
Female 12-month-old	Wid type	10.1	7.85	15.01	8.97	5	0.48
	3xTg-AD	6.31	4.72	9.16	6.43	5	0.48
		Strain				U	P
		Wild type		**3xTg-AD**			
		Mean	SD	Mean	SD		
Male 8-month-old	Water	7.36	3.26	30.8	11.68	1	**0.002**
	Citrulline	14.11	9.06	16.04	5.53	29.5	0.618
Female 8-month-old	Water	8.45	4.61	7.3	4.22	22	0.805
	Citrulline	10.53	4.23	7.72	5.03	11	0.343
Male 12-month-old	Water	5.75	3.09	20.5	11.21	0	**0.029**
	Citrulline	6.25	1.7	12.87	7.35	3.5	0.229
Female 12-month-old	Water	10.1	7.85	6.31	4.72	7	0.886
	Citrulline	15.02	8.97	9.16	6.43	5	0.486

3.2 Arginine levels in CSF and brain tissue

Arginine levels were determined in both cerebrospinal fluid (CSF) and brain tissue (hippocampus) to determine whether the changes seen in the behavioral performance of mice could be related to changes in arginine levels due to oral citrulline supplementation.

Arginine levels in CSF of citrulline treated transgenic (180%) and citrulline treated wild type (134%) male mice showed an increase compared to water-treated mice, as did levels in citrulline treated transgenic (180%) and citrulline treated wild type (126%) female mice. However, the collected CSF was only sufficient for one measurement by HPLC of a pull of mice; hence, no statistical analyses were performed to these data.

Arginine levels were also determined in the hippocampus of both strains with citrulline or water treatments as can been seen in Figure 4. A Kruskal Wallis test for male hippocampus showed statistically significant differences between groups (H(3)= 8.96; p < 0.05) and a Mann Whitney test indicated that the levels of arginine in transgenic water-treated mice was higher (Mdn= 1413) than water-treated wild type mice (Mdn= 427) (U= 3, p < 0.01), and an increase in arginine levels in citrulline-treated transgenic mice (Mdn= 2280) compare with water-treated wild type mice (Mdn= 427) (U= 5 p < 0.05) and a tendency to increase in citrulline-treated transgenic mice (Mdn= 2280) compare with citrulline-treated wild type mice (Mdn= 694) (U=19 p = 0.68); no significant differences in arginine levels in the hippocampus (H(3) = 5,35; p > 0.05) between treatments or strains for female mice were found.

Figure 4. Arginine levels in hippocampus tissue are presented for wild type (circles) and transgenic (triangles) strains. Letter a) represents males and b) females. *p < 0.05. TG: transgenic, WT: wild type

3.3 Nitric oxide

It has been reported that the administration of citrulline can increase arginine

levels, and one of the pathways that arginine could follow is the synthesis of NO, in

order to know if the citrulline that we supplement was raising NO levels in brain

tissue we decided to evaluate it using the Griess method which measures the nitrate and nitrite metabolites.

Nitrite levels were determined in the hippocampus of 3xTgAD and wild type male mice, a Kruskal Wallis showed differences between groups (H (3) = 13.03, $p < 0.01$), and a Dunn's *post hoc* showed significant differences between transgenic citrulline treated mice and water and citrulline treated wild type mice ($Z = 3.05$; $p < 0.05$; $Z = 2.78$; $p < 0.05$).

Something similar happened with nitrite levels of females; a Kruskal Wallis showed differences between groups (H(3) = 12.52; $p > 0.05$) and a Dunn's *post hoc* revealed differences between citrulline-treated transgenic mice and wild type water and citrulline treated mice ($Z = 2.87$; $p < 0.05$; $Z = 2.59$ $p < 0,05$).

Finally, a Spearman r test showed no correlation between arginine levels and latency in seconds ($r = -0.50$; $p = 0.21$) and no correlation between arginine levels and NO levels ($r = -0.60$; $p = 0.35$) for citrulline-treated male 3xTgAD mice.

No correlation was found for either arginine and NO ($r = 0.60$; $p = 0.41$) or arginine and latency in seconds ($r = 0.50$; $p = 0.45$) when analyzing female citrulline treated 3xTgAD mice.

Figure 5. Nitic oxide levels in hippocampus tissue are presented for wild type (circles) and transgenic (triangles) strains. Letter a) represents males and b) females. *$p < 0.05$. NO: nitric oxide, TG: transgenic, WT: wild type

4. DISCUSSION

Three months of citrulline supplementation to male AD transgenic mice improved

memory performance in a spatial task when the administration started at 5 months

of age. We found that 8-month-old male transgenic mice show cognitive decline

compared to the wild type group and that this deterioration can be prevented if the

animals have had citrulline supplementation. No changes in the acquisition were found for either group, hence, the improvement we see here is specific to long term memory.

However, when the same treatment was given to 9-month-old male 3xTg-AD mice, no significant improvement in the memory task was observed. As Alzheimer's disease progresses, cognitive impairment increases, and with that, the possibility of reversing the symptoms diminishes. For this model, cognitive impairment is observed first at 4 months of age [25], we began citrulline supplementation at 5 months of age in order to test an effect in an early stage of Alzheimer's disease in this model. We found that citrulline supplementation improves memory impairment as long as it is administered at an early stage of the disease in this murine model.

By contrast, transgenic female mice of either 8 or 12 months of age showed no memory deficit compared to their wild type counterparts and hence, no improvement could be recorded. Noteworthy, there is a significant effect of sex in the performance of the Morris Water Maze for the 3xTg-AD model. We found differences between male and female mice without taking into account citrulline supplementation, female mice have a better performance independent of the strain for this specific memory task. Female mice of both strains had lower escape latencies on probe day, but no differences with males during acquisition.

On the matter, other studies using 3xTg-AD mice have reported inconsistencies between sexes in their results depending on the memory tasks used [26], which constitutes one of the limitations of working with this murine model. Stover and

coworkers evaluated performance in different behavioral tests of cognitive function such as spontaneous alternation in the Y-Maze, novel object recognition, the Barnes maze, and cued and contextual fear memory tasks. They showed that in general, females have better cognitive performance than males at 6.5 months of age [26].

A possible explanation for differences between sexes is related to immunocompetence; it has been reported that male 3xTgAD mice have a worse immune dysfunction and a higher mortality rate than female 3xTgAD mice [27]. Accordingly, it has been shown that female mice, rats, and monkeys show stronger antibody [28] and cell-mediated immune responses than males, were estrogens have been suggested to be responsible [29,30]. Inflammation has been strongly implicated in the pathogenesis of AD; production of pro-inflammatory cytokines can perpetuate the cycle of neuroinflammatory processes including amyloidosis, neuronal death, cortical thinning, reduced brain volume, cerebrovascular disease-related events, and neurodegeneration [31]. The hypothesis of a direct contribution of the inflammatory response to amyloid plaque progression and, thus, to the neurodegeneration associated with AD [27] could explain the influence of inflammation on cognition.

Neither female nor male wild type mice at 8 or 12 months of age, showed memory improvement when administered with citrulline. Conversely, other preliminary experiments were C57BL6J 24 month-old mice were citrulline-fed for 3 months found memory improvement in a Y-maze task [20]. Nonetheless, our oldest mice were still half the age of the ones used here and do not show memory impairment.

One issue that arises when attempting to compare the results of behavioral tests of 3xTgAD mice across studies is the variety of control strains that have been used, each of which may have distinct behavioral phenotypes. The original background strain by LaFerla (C7BL/6;129X1/SvJ;129S1/Sv, B6129SF2) has been shown to have distinct behavioral phenotypes compared to the second generation offspring of mice created by a cross between C57BL/6J females and 129S1/SvlmJ males B6129SF2, which approximates the original strain and is recommended by the supplier of the 3xTgAD mice [26] and the one used here. However, since we are testing the effect of treatment on memory impairment, our best control is the 3xTgAD treated with water.

Since we saw changes in memory performance after citrulline administration in transgenic male mice, we evaluated the biological implications citrulline could have had. Citrulline metabolism in mammals can be classified into three pathways; the first metabolic pathway is arginine biosynthesis; the second pathway is the nitric oxide (NO) cycle [32], a final substrate that is involved in learning processes and cognitive function [14]; and the third pathway is the urea cycle, taking place in the liver [32].

Based on the above, we evaluated changes in arginine and nitric oxide levels after citrulline administration. We found that arginine increases in CSF due to citrulline supplementation in both strains and both sexes. However, we were only able to analyze one pull of 4 mice samples leaving us with no statistical analysis and constituting one of the limitations of this study. We also found that arginine in the

hippocampus shows only a tendency to increase in male transgenic citrulline-treated mice and observed no changes in female mice.

On the matter, intracellular arginine can be derived from 3 possible sources: uptake of extracellular arginine, de novo synthesis, or arginine released by degradation of cellular proteins. Multiple arginine pools can exist within cells, depending on how rapidly arginine exchanges between compartments [33] and their exchangeability with the extracellular space via exchange transporters as well as their accessibility for NO synthase (NOS) [34]. It has been shown that neuronal NOS (nNOS) in A673 neuroepithelioma and TGW-nu-I neuroblastoma cells can fast and efficiently be nourished by extracellular arginine, under all experimental conditions NO synthesis fully recovered within 4 min upon addition of extracellular arginine [34]. In human neuronal cells, arginine transported into cells was the preferred substrate for nNOS [34].

This is perhaps why we did not find significant changes in arginine levels in hippocampus tissue in citrulline-treated 3xTgAD mice, arginine pools could remain intact because the arginine required for different pathways is taken from the extracellular space. On the matter, Tachikawa and coworkers showed that in male Wistar rats (21 days postnatal) L-arginine is transported into the brain as an intact form and metabolizes immediately to either NO or polyamines [35].

On that matter, it has been shown that geriatric patients with cerebrovascular disease treated with oral L-arginine at a dose of 1.6 g per day for 3 months improve their cognitive function and reduce their levels of lipid peroxidation [36].

Furthermore, it has recently been reported that intracerebroventricular administration of L-arginine for 28 days improves spatial memory acquisition of 6.5 months-old female 3xTgAD [37].

When measuring NO as a product of arginine, we found no statistical significance for 3xTgAD citrulline-treated and water-treated mice (p = 0.19); however, an increase of NO can be seen. The regulatory roles of NO are complex, in part because NO concentrations can vary greatly in magnitude; physiological concentrations can vary from low nanomolar up to low micromolar. Lower concentrations of NO may not be sufficient to activate all signaling pathways but instead will stimulate pathways triggered by highly reactive substrates for NO [38]. A canonical signaling role for NO modifications is the activation of soluble guanylate cyclase (sCG) which generates cyclic guanosine monophosphate (cGMP). The NO-mediated cGMP/Protein Kinase G pathway can modify other important signaling pathways, such as the MAP kinases, leading to enhanced ERK-mediated transcription and have a variety of effects on synaptic plasticity both pre and postsynaptically [39,40]. NO-cGMP-PKG signaling may promote activation of protein kinase signaling cascades critical for long-term synaptic plasticity and memory formation [41]. Hence, no statistically significant changes can still have a biological impact, in this case, a small increase of NO could ultimately have an effect on memory.

Regarding transgenic and wild type mice, we found that NO levels in both male and female transgenic water-treated mice were higher than water-treated wild type mice. These results are in concordance with that reported by Lourenc□o and

coworkers; they observed that in young male 3xTgAD mice (3 and 6 months) the NO peak amplitude tended to be lower compared to age-matched controls. During age progression (12 months of age) the NO peak amplitude increased in 3xTgAD mice while it remained unchanged in control mice, and in old aged mice (18 months) the NO peak amplitude was significantly higher as compared with the age-matched controls [42]. On the other hand, Dias and coworkers reported that in an early stage of the AD pathology, NMDA evoked NO production is significantly augmented in 3xTgAD mice [43].

Finally, we found a significant increase in NO of female transgenic mice compared to their male counterparts. In this regard, it is known that sex hormones influence endothelial function through their effects on agonists and contribute to key functional differences between males and females related to endothelial function and cardiovascular disease risk [44]. It is known that estrogen binding to activated eNOS promotes the production of NO [45] in the endothelial caveola [46] which could affect cognition.

Accordingly, it has been shown that treatment with 17 beta-estradiol to ovariectomized female 3xTgAD mice effectively prevented the acceleration of AD-like neuropathology and behavioral impairment caused by hormone depletion [47]. The same treatment significantly reduced both beta-amyloid accumulation and working memory deficits [48]. This, in part, could explain the lack of memory impairment seen for this model, in this particular task. However, we did not measure estradiol levels since it was not the main objective of the project, but rather a side observation which in concordance with our findings on cognition,

could explain how female 3xTgAD mice do not show memory impairment at the same age as male mice.

At this time, our findings suggest that the memory improvement observed in the 8-month-old 3xTgAD male mice supplemented with citrulline could be due to an increase in NO production from arginine. It is known that NO produced by neuronal NOS (nNOS) functions as an important modulator of neuronal function acting on the release of neurotransmitters [49]. NO plays an important role in synaptogenesis, long term potentiation, and long-term depression [50]. Furthermore, the NO deficiency has been implicated in neurodegeneration by promoting endothelial dysfunction, accelerating formation, and accumulation of amyloid peptides, reducing synaptic plasticity, activating microglia, and evoking neuroinflammation [17].

Citrulline oral administration, a non-toxic supplement that has been previously used for other ailments [51–53], is a viable treatment for memory improvement in the early stages of Alzheimer's Disease. These results point to NO as a viable, efficient target for memory dysfunction in AD. Further research would help define if the results observed here apply to human subjects as well.

5. ACKNOWLEDGMENTS

This paper constitutes part of the Ph.D. in Biological Science graduate requisites of Katia Leticia Martínez-González who thanks Posgrado en Ciencias Biológicas, Biología Experimental, and acknowledges the scholarships provided by CONACyT (294248) and IMSS (99096757). This research was possible thanks to the support of B. Eng. Sergio Becerril Zavala from Pronat.

Funding

This work was supported by the Fondo de Investigación en Salud, Instituto Mexicano del Seguro Social with project number FIS/IMSS/PROT/G17-2/1754.

Disclosure statement

No competing interests to declare.

6. REFERENCES

[1] DeTure MA, Dickson DW. The neuropathological diagnosis of Alzheimer's disease. Molecular Neurodegeneration 2019;14. https://doi.org/10.1186/s13024-019-0333-5.

[2] Austin SA, Santhanam AV, Katusic ZS. Endothelial nitric oxide modulates expression and processing of amyloid precursor protein. Circ Res 2010;107:1498–502.

[3] Bourasset F, Ouellet M, Tremblay C, Julien C, Do TM, Oddo S, et al. Reduction of the cerebrovascular volume in a transgenic mouse model of

Alzheimer's disease. Neuropharmacology 2009;56:808–13.

https://doi.org/10.1016/j.neuropharm.2009.01.006.

[4] Cai H, Cong W-N, Ji S, Rothman S, Maudsley S, Martin B. Metabolic

Dysfunction in Alzheimers Disease and Related Neurodegenerative Disorders.

Current Alzheimer Research 2012;9:5–17.

https://doi.org/10.2174/156720512799015064.

[5] Sonntag K-C, Ryu W-I, Amirault KM, Healy RA, Siegel AJ, McPhie DL, et al.

Late-onset Alzheimer's disease is associated with inherent changes in

bioenergetics profiles. Scientific Reports 2017;7.

https://doi.org/10.1038/s41598-017-14420-x.

[6] Ferrer I. Altered mitochondria, energy metabolism, voltage-dependent anion

channel, and lipid rafts converge to exhaust neurons in Alzheimer's disease.

Journal of Bioenergetics and Biomembranes 2009;41:425–31.

https://doi.org/10.1007/s10863-009-9243-5.

[7] Hurtado MO, Kohler I, de Lange EC. Next-generation biomarker discovery in

Alzheimer's disease using metabolomics - from animal to human studies.

Bioanalysis 2018;10:1525–46.

[8] Gueli MC, Taibi G. Alzheimer's disease: amino acid levels and brain metabolic

status. Neurological Sciences 2013;34:1575–9.

https://doi.org/10.1007/s10072-013-1289-9.

[9] Arlt S, Schwedhelm E, Kölsch H, Jahn H, Linnebank M, Smulders Y, et al.

Dimethylarginines, homocysteine metabolism, and cerebrospinal fluid markers

for Alzheimer's disease. J Alzheimers Dis 2012;31:751–8.

[10] Fonteh AN, Harrington RJ, Tsai A, Liao P, Harrington MG. Free amino acid

and dipeptide changes in the body fluids from Alzheimer's disease subjects. Amino Acids 2007;32:213–24. https://doi.org/10.1007/s00726-006-0409-8.

[11] Ibáñez C, Simó C, Martín-Álvarez PJ, Kivipelto M, Winblad B, Cedazo-Mínguez A, et al. Toward a Predictive Model of Alzheimer's Disease Progression Using Capillary Electrophoresis–Mass Spectrometry Metabolomics. Analytical Chemistry 2012;84:8532–40. https://doi.org/10.1021/ac301243k.

[12] Mulder C, Wahlund L-O, Blomberg M, de Jong S, van Kamp GJ, Scheltens P, et al. Alzheimer's disease is not associated with altered concentrations of the nitric oxide synthase inhibitor asymmetric dimethylarginine in cerebrospinal fluid. J Neural Transm 2002;109:1203–8.

[13] Kaiser E, Schoenknecht P, Kassner S, Hildebrandt W, Kinscherf R, Schroeder J. Cerebrospinal Fluid Concentrations of Functionally Important Amino Acids and Metabolic Compounds in Patients with Mild Cognitive Impairment and Alzheimer's Disease. Neurodegenerative Diseases 2010. https://doi.org/10.1159/000287953.

[14] Hawkins RD. NO Honey, I Don't Remember. Neuron 1996;16:465–7. https://doi.org/10.1016/s0896-6273(00)80064-1.

[15] Dubey H, Gulati K, Ray A. Amelioration by nitric oxide (NO) mimetics on neurobehavioral and biochemical changes in experimental model of Alzheimer's disease in rats. Neurotoxicology 2018;66:58–65.

[16] Liu C, Liang MC, Soong TW. Nitric Oxide, Iron and Neurodegeneration. Front Neurosci 2019;13:114.

[17] Katusic ZS, Austin SA. Endothelial nitric oxide: protector of a healthy mind. Eur

Heart J 2014;35:888–94.

[18] Lorin J, Zeller M, Guilland J-C, Cottin Y, Vergely C, Rochette L. Arginine and nitric oxide synthase: regulatory mechanisms and cardiovascular aspects. Mol Nutr Food Res 2014;58:101–16.

[19] Citrulline: From metabolism to therapeutic use. Nutrition 2013;29:479–84.

[20] Marquet-de Rougé P, Clamagirand C, Facchinetti P, Rose C, Sargueil F, Guihenneuc-Jouyaux C, et al. Citrulline diet supplementation improves specific age-related raft changes in wild-type rodent hippocampus. Age 2013;35:1589–606.

[21] Bromley-Brits K, Deng Y, Song W. Morris Water Maze Test for Learning and Memory Deficits in Alzheimer's Disease Model Mice. Journal of Visualized Experiments 2011. https://doi.org/10.3791/2920.

[22] Ivanova M, Artusi C, Boffa GM, Zaninotto M, Plebani M. HPLC determination of plasma dimethylarginines: method validation and preliminary clinical application. Clin Chim Acta 2010;411:1632–6.

[23] de Jong S, Teerlink T. Analysis of asymmetric dimethylarginine in plasma by HPLC using a monolithic column. Anal Biochem 2006;353:287–9.

[24] Espey MG, Miranda KM, Thomas DD, Wink DA. Distinction between nitrosating mechanisms within human cells and aqueous solution. J Biol Chem 2001;276:30085–91.

[25] Billings LM, Oddo S, Green KN, McGaugh JL, LaFerla FM. Intraneuronal Aβ Causes the Onset of Early Alzheimer's Disease-Related Cognitive Deficits in Transgenic Mice. Neuron 2005;45:675–88. https://doi.org/10.1016/j.neuron.2005.01.040.

[26] Stover KR, Campbell MA, Van Winssen CM, Brown RE. Early detection of cognitive deficits in the 3xTg-AD mouse model of Alzheimer's disease. Behav Brain Res 2015;289:29–38.

[27] Giménez-Llort L, Arranz L, Maté I, De la Fuente M. Gender-Specific Neuroimmunoendocrine Aging in a Triple-Transgenic 3×Tg-AD Mouse Model for Alzheimer's Disease and Its Relation with Longevity. Neuroimmunomodulation 2008;15:331–43. https://doi.org/10.1159/000156475.

[28] Weinstein Y, Ran S, Segal S. Sex-associated differences in the regulation of immune responses controlled by the MHC of the mouse. J Immunol 1984;132:656–61.

[29] Fuente MDL, De La Fuente M, Baeza I, Guayerbas N, Puerto M, Castillo C, et al. Changes with ageing in several leukocyte functions of male and female rats. Biogerontology 2004;5:389–400. https://doi.org/10.1007/s10522-004-3201-8.

[30] Keller ET, Zhang J, Yao Z, Qi Y. The impact of chronic estrogen deprivation on immunologic parameters in the ovariectomized rhesus monkey (Macaca mulatta) model of menopause. J Reprod Immunol 2001;50:41–55.

[31] McGrattan AM, McGuinness B, McKinley MC, Kee F, Passmore P, Woodside JV, et al. Diet and Inflammation in Cognitive Ageing and Alzheimer's Disease. Curr Nutr Rep 2019;8:53–65.

[32] Curis E, Nicolis I, Moinard C, Osowska S, Zerrouk N, Bénazeth S, et al. Almost all about citrulline in mammals. Amino Acids 2005;29:177–205. https://doi.org/10.1007/s00726-005-0235-4.

[33] Morris SM. Arginine Metabolism Revisited. The Journal of Nutrition

2016;146:2579S – 2586S. https://doi.org/10.3945/jn.115.226621.

[34] Simon A, Karbach S, Habermeier A, Closs EI. Decoding the substrate supply to human neuronal nitric oxide synthase. PLoS One 2013;8:e67707.

[35] Tachikawa M, Hirose S, Akanuma S-I, Matsuyama R, Hosoya K-I. Developmental changes of l-arginine transport at the blood-brain barrier in rats. Microvasc Res 2018;117:16–21.

[36] Ohtsuka Y, Nakaya J. Effect of oral administration of L-arginine on senile dementia. Am J Med 2000;108:439.

[37] Fonar G, Polis B, Meirson T, Maltsev A, Elliott E, Samson AO. Intracerebroventricular Administration of L-arginine Improves Spatial Memory Acquisition in Triple Transgenic Mice Via Reduction of Oxidative Stress and Apoptosis. Transl Neurosci 2018;9:43–53.

[38] Thomas DD, Heinecke JL, Ridnour LA, Cheng RY, Kesarwala AH, Switzer CH, et al. Signaling and stress: The redox landscape in NOS2 biology. Free Radic Biol Med 2015;87:204–25.

[39] Batchelor AM, Bartus K, Reynell C, Constantinou S, Halvey EJ, Held KF, et al. Exquisite sensitivity to subsecond, picomolar nitric oxide transients conferred on cells by guanylyl cyclase-coupled receptors. Proceedings of the National Academy of Sciences 2010;107:22060–5. https://doi.org/10.1073/pnas.1013147107.

[40] Ota KT, Monsey MS, Wu MS, Young GJ, Schafe GE. Synaptic plasticity and NO-cGMP-PKG signaling coordinately regulate ERK-driven gene expression in the lateral amygdala and in the auditory thalamus following Pavlovian fear conditioning. Learn Mem 2010;17:221–35.

[41] Chien W-L, Liang K-C, Teng C-M, Kuo S-C, Lee F-Y, Fu W-M. Enhancement of long-term potentiation by a potent nitric oxide-guanylyl cyclase activator, 3-(5-hydroxymethyl-2-furyl)-1-benzyl-indazole. Mol Pharmacol 2003;63:1322–8.

[42] Lourenço CF, Ledo A, Barbosa RM, Laranjinha J. Neurovascular uncoupling in the triple transgenic model of Alzheimer's disease: Impaired cerebral blood flow response to neuronal-derived nitric oxide signaling. Exp Neurol 2017;291:36–43.

[43] Dias C, Lourenço CF, Ferreiro E, Barbosa RM, Laranjinha J, Ledo A. Age-dependent changes in the glutamate-nitric oxide pathway in the hippocampus of the triple transgenic model of Alzheimer's disease: implications for neurometabolic regulation. Neurobiol Aging 2016;46:84–95.

[44] Stanhewicz AE, Wenner MM, Stachenfeld NS. Sex differences in endothelial function important to vascular health and overall cardiovascular disease risk across the lifespan. American Journal of Physiology-Heart and Circulatory Physiology 2018;315:H1569–88. https://doi.org/10.1152/ajpheart.00396.2018.

[45] Kim KH, Bender JR. Rapid, estrogen receptor-mediated signaling: why is the endothelium so special? Sci STKE 2005;2005:e28.

[46] Mineo C, Shaul PW. Regulation of eNOS in Caveolae. Advances in Experimental Medicine and Biology 2012:51–62. https://doi.org/10.1007/978-1-4614-1222-9_4.

[47] Carroll JC, Rosario ER, Chang L, Stanczyk FZ, Oddo S, LaFerla FM, et al. Progesterone and Estrogen Regulate Alzheimer-Like Neuropathology in Female 3xTg-AD Mice. Journal of Neuroscience 2007;27:13357–65. https://doi.org/10.1523/jneurosci.2718-07.2007.

[48] Carroll JC, Pike CJ. Selective estrogen receptor modulators differentially regulate Alzheimer-like changes in female 3xTg-AD mice. Endocrinology 2008;149:2607–11.

[49] Christopherson KS, Bredt DS. Nitric oxide in excitable tissues: physiological roles and disease. J Clin Invest 1997;100:2424–9.

[50] Prast H, Philippu A. Nitric oxide as modulator of neuronal function. Progress in Neurobiology 2001;64:51–68. https://doi.org/10.1016/s0301-0082(00)00044-7.

[51] Schwedhelm E, Maas R, Freese R, Jung D, Lukacs Z, Jambrecina A, et al. Pharmacokinetic and pharmacodynamic properties of oral L-citrulline and L-arginine: impact on nitric oxide metabolism. Br J Clin Pharmacol 2008;65:51–9.

[52] Kim I-Y, Schutzler SE, Schrader A, Spencer HJ, Azhar G, Deutz NEP, et al. Acute ingestion of citrulline stimulates nitric oxide synthesis but does not increase blood flow in healthy young and older adults with heart failure. Am J Physiol Endocrinol Metab 2015;309:E915–24.

[53] Vincent MA, Clerk LH, Lindner JR, Klibanov AL, Clark MG, Rattigan S, et al. Microvascular recruitment is an early insulin effect that regulates skeletal muscle glucose uptake in vivo. Diabetes 2004;53:1418–23.