# Negative density-dependent parasitism in a group-living carnivore

Gregory F Albery*[1], Chris Newman[2], Julius Bright Ross[2], Shweta Bansal[1], Christina Buesching[2]

1. Department of Biology, Georgetown University, Washington, DC, USA
2. Wildlife Conservation Research Unit, The Recanati-Kaplan Centre, Department of Zoology, University of Oxford, Oxford, U.K.

*Corresponding author: gfalbery@gmail.com

# Abstract

Animals living at high population densities commonly experience greater exposure to disease, leading to increased parasite burdens. However, social animals can benefit immunologically and hygienically from cooperation, and individuals may alter their socio-spatial behaviour in response to infection, both of which could counteract density-related increases in exposure. Consequently, the costs and benefits of sociality for disease are often uncertain. Here, we use a long-term study of a wild European badger population (*Meles meles*) to investigate how within-population variation in host density determines infection with multiple parasites. Four out of five parasite taxa exhibited consistent spatial hotspots of infection, which peaked among badgers living in areas of low local population density. Combined movement, survival, spatial, and social network analyses revealed that parasite avoidance was the likely cause of this negative density dependence, with possible roles for localised mortality, encounter-dilution effects, and micronutrient-enhanced immunity. These findings demonstrate that animals can organise their societies in space to minimise parasite infection, with important implications for badger behavioural ecology and for the control of badger-associated diseases.

# Introduction

A wild animal's infectious disease burden is determined by its exposure and susceptibility to

infective pathogens. Typically, higher population density results in increased contact rates

and thus greater exposure [1,2]; however, sociality carries immunological and hygienic

benefits that can counteract this exposure-exacerbating effect [3,4]. For instance, cooperative

foraging improves nutrition, alleviating infection costs [5]; mutual grooming removes

ectoparasites [6–8]; and "social immune responses" maintain collective group health [9,10],

e.g. through extirpating sick individuals [11]. Additionally, the environmental distribution of

resources can influence spatial behaviour [12–14] and determine local population densities

[15], while altering susceptibility [16] and transmission efficiency [17]. Sociality and disease

may thus be confounded through shared causal origins rather than being mechanistically

linked [17]. Finally, individuals can minimise exposure by avoiding environmental cues

[18,19] or infected conspecifics [8,20], creating a population-level "landscape of disgust"

analogous to predatory "landscapes of fear" [21,22]. These processes could produce negative

density effects, sometimes depending on parasite transmission mode [1,17,23], but their role

in defining observed density-infection relationships is poorly understood.

Uncertainty concerning density-infection relationships largely arises because studies of social

effects are often carried out across discrete groups [1,24], between populations [25,26], or

between species (e.g. [1,2,24,27]). Continuous density measures are rarely linked to parasite

burden within one contiguous host population, and studies that do so generally use purely

social metrics rather than spatial density gradients (e.g. [28–30]). Continuous spatial density

measures are advantageous because: density is a continuous, spatially explicit variable [31–

33]; for the many parasites that are spread through the environment, spatial density measures

better represent transmission than direct (social network) metrics [17,34–36]; and between-

population and cross-species comparisons are fraught with confounding factors such as

shared environmental causality or compensatory evolutionary changes [1,3], which within-

population studies can more easily avoid or overcome. Furthermore, accounting for spatial

effects may help anticipate or test the caveats of cooperation, environmental causality, and

behavioural avoidance outlined above.

European badgers (*Meles meles*) are nocturnal, ground-dwelling mustelids with a pan-

European distribution, eating a varied diet composed largely of earthworms [37]. Badgers are

of particular interest for cattle health in the United Kingdom because of their role in

maintaining and spreading bovine tuberculosis [37,38]. Although intended as a control

measure, intense culling perturbs badger social systems, causing stressed survivors to

disperse, thereby spreading bovine tuberculosis to neighbouring farms [39–41].

Understanding badgers' socio-spatial behaviours and their epidemiological implications is

therefore an important research priority (e.g. [38,42–44]). The demography, behaviour, and

parasitism of badgers in Wytham Woods, Oxfordshire have been studied continuously since

1987 [37,45]. Dens (termed "setts") are situated dependent on soil composition and landscape

topography [46]. These badgers reside in cohesive mixed-sex social groups, with around 35%

consistent philopatry; however, these groups are also fluid: 16.4% of individuals are trapped

at a different social group to their previous capture history, and 19.8% away from their natal

group [47]. Furthermore, from genetic pedigree, 48% of cubs have extra-group paternity [48].

Badgers host several arthropod parasites, including badger-specific fleas (*Paraceras melis*)

and lice (*Trichodectes melis*) and generalist ticks (*Ixodes* sp.) [49]. They also carry two

gastrointestinal protozoans: *Eimeria melis* and *Isospora melis*. *E. melis* predominantly infects

young individuals, causing substantial mortality [50,51]. Although studies have examined

social grooming effects on ectoparasite burden [6,52] and roles of denning behaviour in

parasite transmission [49], the within-population spatial-social parasite epidemiology has yet

to be investigated.

Here, we investigate parasite burdens in the Wytham badger population and their associations

with spatial and social behaviour. We establish parasite distributions using spatial

autocorrelation models. We quantify social drivers using both spatial density gradients and

social network metrics, postulating that greater badger densities would drive higher parasite

burdens through increased exposure. Finally, we examine survival effects, investigating

whether parasite-linked mortality might alter spatial patterns of badger population density.

We consider a range of potential social/spatial covariates of parasite burden, including

density-related exposure changes, benefits of cohabitation, condition/susceptibility effects,

encounter-dilution effects, and parasite avoidance behaviours.

# 84    Methods

## 85    Data collection

Badgers were sampled as part of a long-term study in Wytham Woods, Oxfordshire, UK

(51.778°N, 1.313°W), established in 1987 and recording one of the highest local badger

densities ever reported [37,45]. Badgers were trapped overnight at their setts in steel-mesh

cages baited with peanuts, collected the next morning, transported to a handling facility and

sedated. Individuals were identifiable by tattoo, applied on first trapping (typically as cubs).

Measures were taken of head-body length (mm) and weight (to the nearest 0.1kg). We

calculated a standardised index of body condition dividing log(weight) by log(body length).

The population was trapped seasonally (winter: Jan/Feb/March; spring: April/May/June;

summer: July/Aug/Sept; autumn: Oct/Nov/Dec) in 4 quadrants, for 3 days per quadrant. Our

dataset included 9016 captures of 1369 badgers spanning 29 years (1990-2018).

The population currently comprises 23 social groups (mean group size=11; range 2-29

individuals), each using more than one sett, with sett/social group affiliation established using

baitmarking [46,53]. Badgers were assigned to groups using established residency rules [48].

We computed social networks based on co-trapping using a "gambit of the group" approach ,

where individuals trapped in the same sett in the same year were assumed to have interacted

[54].

Fleas were counted during a stereotyped 20-second inspection of the badger's full body, and

ticks (*Ixodes* sp.) and lice (*Trichodectes melis*) were counted within  a 4x4cm square of

preferentially-parasitised skin near the groin (per Cox *et al.* 1999). Faecal sampling for two

protozoan endoparasites, *Eimeria melis* and *Isospora melis,* was carried out through 1993-

1997 and 2009-2017 (N=1287 counts). Endoparasite  were counted using salt flotation and

microscopy [51,55]. Each count was duplicated, and data were reduced to a binary

infected/uninfected status (rather than counts) because of their highly overdispersed

distribution.

# Statistical Analysis

## Covariates

Statistical analysis and data manipulation used R version 3.6.0 [56]. All data and code are

available at https://github.com/gfalbery/Badgeworks. Models were constructed using the

`inla` package [57,58]. Response variables included counts of fleas and lice (negative

binomial distribution) and prevalence of ticks, *Eimeria*, and *Isospora*. Explanatory covariates

included: Year (continuous); Month (categorical); Age Category (categorical: cub, yearling,

and adult); Sex (male and female); and Body Condition (continuous). Continuous covariates

were standardised (mean=0; standard deviation=1). Individual Identity and Year were fitted

throughout as categorical random effects.

## 1. Spatial autocorrelation models

To identify generalised spatial patterns we first fitted models accounting for continuous spatial autocorrelation across the population [33,57,58]. We fitted a base model using only Year, Month, Age Category, and Sex, with individual and annual random effects. We then added an SPDE random effect controlling for spatial autocorrelation in the response variable in space, based on individuals' point locations. We then compared these models using Deviance Information Criterion (DIC) to establish whether accounting for spatial autocorrelation improved model fit; models within 2ΔDIC were judged competitive.

## 2. Density models

We then fitted models that replaced the SPDE effects with individual measures of local population density. Models included a range of social/spatial density measures, calculated using various methods. Spatial density metrics represent numbers of individuals in a given space, and therefore tread the line between social and spatial behavioural traits [17]. We computed these by creating space use distribution kernels with the `adehabitathr` package in R [59]. We rasterised the usage distribution, assigning each individual a local density based on the raster value for their map location (Figure 1A-B). Measures included: 1) **Lifetime density**: the density of individuals' centroids across the study period. 2) **Trapping density**: the density of trapping events, incorporating multiple captures of the same individual per year. 3) **Annual density**: the density of individuals' centroids in the sampling year, calculated from annual density kernels. Only one spatial density measure was allowed in a given model, as the measures correlate (R>0.5) and co-fitting several measures produced spurious, unexpected correlations in the opposite direction expected from data exploration. We also used two social metrics based on co-trapping patterns. **Degree centrality** was the number of unique badgers with whom each individual was trapped in the same sett per year. **Group size** was the total number of individuals trapped in a given social group per year. Following spatial autocorrelation model procedures, we conducted a model selection procedure for behavioural metrics, using 2ΔDIC as the cutoff, and only including the best-fitting metric. We conducted these models for: the overall dataset (N=9016); adults only (age 2+, fitting age in years as a fixed covariate; N=6159); and juveniles only (N=1639). For these models, we display only the best-fitting metric, and for the overall dataset if it exhibited a significant trend (and a subset-only model otherwise).

## 3. Multi-response models

To investigate whether parasites covaried with badger density at the within- or between-individual level, we constructed multi-response models in MCMCglmm with an unstructured covariance matrix [60,61]. We fitted parasites and local density as response variables, estimating their covariance when accounting for other fixed effects and decomposing this covariance at the within- and between-individual level. A negative within-individual (residual) correlation would imply that higher burden/prevalence individuals (compared to their baseline) moved to lower density areas during their lives, supporting either social ostracism or sickness behaviours. In contrast, negative between-individual covariance would imply that individuals inhabiting lower densities generally had inherently greater parasitism. We only constructed these models for parasite-host age category combinations that demonstrated density effects in the INLA models.

## 4. Survival models

To investigate parasites' mortality effects, we fitted models with survival as a binary response (1=animal seen in any subsequent year; 0 otherwise). These models included the same covariates as the parasite models, plus parasite prevalence/count and badger density measures. Parasite measures were either log(x+1)-transformed integers (fleas, lice) or binary (ticks, *Eimeria*, and *Isospora*). We conducted another model selection procedure, adding parasite metrics successively if they improved model fit. This was carried out for adults and juveniles separately.

# Results

INLA autocorrelation models revealed considerable spatial structuring of parasite burdens. Models for 4/5 parasites were substantially improved by incorporating spatial effects (ΔDIC>8; Table 1). Only *Isospora*'s model was not improved by spatial autocorrelation (Table 1).

Density models provided substantial support for negative associations between badger density and parasite infection. Including at least one density measure as a covariate improved the model for at least one age category for 4/5 parasites; all had significant negative slopes for density effects (Figure 2; Figure SI1-4; Table SI1-3). Slopes were relatively shallow, but extremely significant and robust (Figure 2). Flea counts were best described by Lifetime Density for the overall model (ΔDIC=-16.31; P<0.0001; Figure 2A), while lice burdens decreased with Trapping Density in the juveniles-only model (ΔDIC=-8.77; P=0.0017; Figure

2B). Tick prevalence was lower in areas of higher Lifetime Density ($\Delta$DIC=-2.47; P=0.0038;

Figure 2C), and *Eimeria* prevalence was lower with greater Trapping Density ($\Delta$DIC=-5.83;

P=0.0053; Figure 2D). Only one positive social effect was detected: adults with greater

degree had slightly more lice ($\Delta$DIC=-2.39; P=0.027; Figure SI2). All DIC changes

associated with the model addition procedure are presented in Table SI1-3.

MCMCglmm multi-response models revealed similar trends to INLA univariate models

(Figure SI1-4), and allowed us to decompose density-parasite correlations into within- and

between-individual changes (Figure 3B). Correlations were greater and much more

significant between- than within-individuals in all cases (Figure 3B). This demonstrates that

movement between high- and low-density areas was unlikely to produce changes in parasite

burden over an individual's lifetime, but rather that individuals' home ranges exhibited

inherently different local density-parasite relationships. Only louse infection predicted

survival probability; the effect was relatively small, minimally significant, and specific to

juveniles (P=0.038; Figure 3A).

Our models also revealed various other effects (Figures SI1-4). Briefly, in the overall models,

juveniles had fewer fleas, more lice, and greater *Eimeria* prevalence than did yearlings and

adults, and lower *Isospora* prevalence than adults (Figure SI1). Males had more lice than did

females, with substantial monthly variation in all parasites (Figure SI1). Additionally, body

condition was negatively associated with fleas, lice, and *Eimeria* infection in all age/sex

classes (Figure SI1). The adults-only and juveniles-only models were very similar to the

overall models (Figures SI1-3); notably, in adults, lice burden increased with age in years,

whereas *Eimeria* and *Isospora* prevalence decreased (Figure SI2).

# Discussion

Using a combination of spatial and social network analysis, we uncovered negative

associations between local population density and multiple parasites in this wild group-living

carnivore. We found strong but contrasting spatial structuring of fleas, lice, ticks, and

*Eimeria*, and, contrary to expectations, all four were most prevalent or abundant in areas of

lowest badger density. Co-trapping network and grouping metrics were not predictive of

parasitism, implying that "direct" social behaviours, such as mutual allo-grooming, were

unlikely to explain the negative density effects. Additionally, badger density effects

manifested independent of survival and body condition effects, implying that spatial

structuring of the host population did not originate from: 1) localised host mortality, 2)

reduced susceptibility arising from co-habitation and implied cooperation benefits [3,5], or 3)

greater local resource availability influencing susceptibility [16,62,63]. Additionally, multi-

response models revealed low within-individual covariance between density and parasites,

providing little support for heavily-parasitised individuals becoming ostracised during the

course of their lives [11]. Taking spatial structuring together with setts' propensity to harbour

parasites [49,64], the most parsimonious interpretation is that badgers avoid infection

behaviourally, preferring to inhabit areas poorly disposed to parasite transmission (Table 2).

These individual-level avoidance behaviours amplify at the population level to produce

patterns of badger density inversely related to parasite distributions in the environment, as

expected under a "landscape of disgust" [19,21,22]. As well as providing rare evidence of

non-consumptive effects of parasites in a wild carnivore, these results imply that animals can

arrange their society in space according to parasite transmission.

Our study demonstrates observationally that parasites can determine a society's structure in

the environment, and that this phenomenon may counteract the more conventional prediction

that host density exacerbates parasitism through increased exposure (e.g. [1,2,65]).

Consequently, studies aiming to quantify social covariates of disease should explicitly  seek

to investigate individual movement and avoidance behaviours, socio-spatial structuring, and

feedbacks between sociality and space use [17]. Previous studies on other mammals have

revealed avoidance of infected conspecifics [8] and faeces [18,66], but it remained unclear

whether animals avoid spatial hotspots of transmission themselves, and whether these could

have population-level consequences [21,22]. Badgers respond to social scent cues and may

use these to detect and avoid infested or infectious individuals [67]; furthermore, they move

between setts regularly to avoid accumulating parasites [64,68], abandoning highly infested

setts and chambers [49];  we posit that this behaviour has emergent population-level

consequences. Wytham badgers preferentially establish their setts on northwest-facing slopes

in areas with sandy soils [46], and variation in internal sett temperature and humidity are

associated with reproductive success [69,70]. Our data suggest additional selection for sett

traits and sites resistant to parasite infestation and transmission, which produce an emergent

trend for fewer setts, with fewer occupants, in more highly parasitised areas. Quantifying

parasites in the environment or inside setts and correlating them with badger behaviour could

provide support for this hypothesis. Environmental (sett chamber-based) data on parasite

presence would be needed to determine whether badgers avoid infection *actively* (pre-

infection), using environmental cues that predict parasitism, or *reflexively* (post-infection), by

moving away from areas of high parasitism due to irritation or sickness.

Four parasites exhibited negative density effects, which could impose tradeoffs on avoidance

behaviour. The parasites' distributions were highly contrasting, likely driven by different

environmental factors: for example, sett microclimates favouring flea survival will facilitate

their transmission [49,64]; similarly, *Eimeria* is transmitted faecal-orally, and oocysts are

likely acquired from warm, moist sett chambers, which may explain the gradual decline

moving away from the Thames river in the Northwest toward drier parts of the woods (Figure

1F). Ultimately, in combination with other socio-ecological factors (e.g., finding suitable

mates), badgers may be incapable of completely avoiding all parasites via denning decisions,

which could mediate co-infection and promote diverse parasite communities, giving rise to

local hotspots across the population. This may lead to parasite avoidance being traded off

against foraging, reproductive success and survival [22,71]. If badgers exhibit between-

individual variation in movement and foraging specialisation [72], avoidance of multiple co-

infecting parasites could maintain between-individual immune heterogeneity [55]. Notably,

only cub density was negatively associated with lice. This observation could be linked to the

detectable mortality effects in cubs, driving local cub mortality and/or motivating

reproducing females to avoid spatial hotspots of lice.

All of the parasites we examined employ some degree of indirect transmission, likely

yielding different relationships with host density than more directly transmitted parasites

[1,17]. Ultimately, only avoiding conspecifics can help avoid direct parasite transmission

(e.g. [8]), which may not be achievable through purely spatial structuring. Future studies

could examine sexually transmitted infections or aerosol-transmitted viruses to investigate

whether individuals living in areas of lower density gain any benefit in terms of direct

parasite transmission. This may be of particular importance for bovine tuberculosis (bTB),

which has a complex, nonlinear relationship with badger sociality [73–75]. Notably,

previous bTB studies have generally used social group size as a metric of sociality; given that

bTB has an environmental transmission component, particularly between badgers and cattle

[76], spatial density metrics, such as those employed here, could be revealing.

We note two potential sources of negative density dependence untested by our modelling

approach: the encounter-dilution effect [77,78] and micronutrient impacts on immunity [79].

For the former, where parasites exhibit a constant attack rate in space, greater host densities

actually drive a lower per-individual burden of disease because the same number of parasites

is divided among more available hosts [77,78]. Because this requires that the spatial

distributions of parasites are not tightly linked to the host distribution (e.g. through

reproducing on the host), it is unlikely to apply for any of the badger-specific parasites

studied here. However, non-host-specific ticks (*Ixodes* sp.) could transmit from other species

to badgers in a given area, producing an encounter-dilution effect; therefore, we are unable to

rule out this mechanism (Table 2). Regarding nutritional status, we used body condition

index as a coarse predictor of the ability to mount a healthy immune response. However,

micronutrients are essential to effective immune function [79] and would not be detected

from body condition indices; therefore, it is plausible that badgers might congregate in areas

of high micronutrient availability. If this behavioural tendency functions to combat pathogens

specifically, it amounts to a "landscape of disgust" acting through reduced susceptibility

rather than reduced exposure. This possibility could be tested in this or similar systems by

comparing the spatial distributions of high-resolution individual-level immune measures with

host density distributions.

Our findings have important implications for the socio-spatial dynamics of this system and its

resilience to pathogens and ecological change. If badger populations are organised optimally

to occupy areas of least parasite transmission, even small disturbances (e.g. setts lost to

forestry) could disrupt its socio-spatial structure and force individuals into unfavourable,

more highly parasitised areas, exacerbating their disease burden. Therefore, disruptive

anthropogenic activities such as culling-linked perturbation could have unseen consequences

for badger disease beyond larger-scale movement impacts (e.g. [39,40]). These findings

further inform our understanding of the drivers of badger spatial behaviour, offering insights

that may be invaluable for their conservation and disease control [38,46].

# Acknowledgements

304    We would like to thank two anonymous reviewers, and Dan Becker, Amy Sweeny, and

305    Quinn Webber for comments on the manuscript. We also warmly acknowledge the many

306    students and co-workers who helped to collect this substantial data set. GFA and SB were

307    supported by NSF grant number 1414296.

309     Figure 1: Spatial distributions of badger population density and parasites in Wytham Woods,
310    Oxfordshire, between 1989 and 2018. A: badger population density distributed across Wytham
311    Woods, calculated based on a space use kernel for individuals' annual centroids. B: Individual
312   badgers trapped at setts (represented by points) were assigned a local density value based on their
313   location on the rasterised space use kernel. Darker blue colours in A and B correspond to greater
314  population density. C-F: the spatial distribution of the four spatially distributed parasites, estimated
315 using the INLA SPDE effect. Darker colours correspond to increased parasitism. The density values
316  in B were fitted as covariates in linear models to explain individual parasite burdens, revealing a
317 negative correlation between density and parasitism (see Fig. 2). All axes are in kilometres, with the
318 0,0 point at the bottom left of the study area. We examined a fifth parasite, *Isospora melis*, but it was
319        not significantly spatially distributed (see Table 1).

Figure 2: Negative associations between population density and parasite infection. Badgers living at higher densities had fewer fleas (A), fewer lice (B), lower tick prevalence (C), and lower *Eimeria melis* prevalence (D). B represents juveniles only. Points represent individual samples, with colours randomised along a colour palette for plotting clarity. Opaque black lines are derived from linear model fits, taking the mean of the posterior distribution. Transparent grey lines represent 100 fits drawn randomly from the posterior estimate distributions of each model, to demonstrate error in the slope and intercepts. Text at the top of the Figures communicates slope estimates, with lower and upper 95% credibility intervals in brackets, and P values. NB: in panel C, the y axis scale and the location of the positive points have been altered for plotting clarity and to better visualise the slope, due to low tick prevalence.

Figure 3: Associations of parasitism with survival and density-parasitism covariance partitioning. A)
Louse burden was negatively associated with individual survival probability in juveniles. Opaque
black line=linear model fit. Transparent grey lines=100 fits drawn randomly from the posterior
estimate distribution, to demonstrate error in the slope and intercept. Text at the top of panel A
communicates slope estimate, with lower and upper credibility interval in brackets and P value. B)
Estimates for within- and between-individual covariance between parasitism and density, taken from
MCMCglmm multivariate models. Points=posterior mean effect size estimates; error bars=95%
credibility intervals.

|  | **Fleas** | **Lice** | **Ticks** | **Eimeria** | **Isospora** |
|---|---|---|---|---|---|
| **Base** | 153.03 | 36.57 | 49.25 | 8.28 | 0 |
| **SPDE** | 0 | 0 | 0 | 0 | 2.25 |

Table 1: DIC changes associated with spatial autocorrelation terms for the five examined parasites.
Lower numbers denote better models; best-fitting models ΔDIC=0. All models except *Isospora* were
improved by addition of spatial fields ("SPDE").

| Potential mechanism | Conclusion | Reason for conclusion |
|---|---|---|
| Co-habitation/ cooperation benefits | No | No direct social effects or competition with body condition effects (Model Set 2; Figure SI1-3) |
| Allo-grooming | No | Spatial rather than direct social effects; grooming not possible for endoparasites (Model Set 2; Figure SI1-3; Table SI1-3) |
| Nutrition-associated immune benefits | No | No competition with body condition effects (Model Set 2; Figure SI1-3) |
| Social ostracism/self-isolation | No | Low within-individual covariance of density and parasitism (Model Set 3; Figure 3B) |
| Local host die-offs | ~No | Possible for lice, in juveniles, but no other mortality effects were evident (Model Set 4; Figure 3A) |
| Encounter-Dilution | ~No | Possible for generalist ticks, but not for the other (badger-specific, non-mobile) parasites [77,78] |
| Avoidance | Yes | All other possibilities eliminated; consistent with individual-level behavioural responses [64] |

353 Table 2: Reasoning surrounding our hypothesis testing. We rejected most hypotheses for our observed
354 negative density effects other than parasite avoidance.

# References

1.  Cote IM, Poulin R. 1995 Parasitism and group-size in social animals: a meta-analysis.
*Behav. Ecol.* **6**, 159–165. (doi:10.1093/beheco/6.2.159)

2.  Altizer S *et al.* 2003 Social Organization and Parasite Risk in Mammals: Integrating
Theory and Empirical Studies. *Annu. Rev. Ecol. Evol. Syst.* **34**, 517–547.
(doi:10.1146/annurev.ecolsys.34.030102.151725)

3.  Ezenwa VO, Ghai RR, McKay AF, Williams AE. 2016 Group living and pathogen
infection revisited. *Curr. Opin. Behav. Sci.* **12**, 66–72.
(doi:10.1016/j.cobeha.2016.09.006)

4.  Kappeler PM, Cremer S, Nunn CL. 2015 Sociality and health: impacts of sociality on
disease susceptibility and transmission in animal and human societies. *Philos. Trans.*

*R. Soc. B Biol. Sci.* **370**, 20140116–20140116. (doi:10.1098/rstb.2014.0116)

5. Almberg ES, Cross PC, Dobson AP, Smith DW, Metz MC, Stahler DR, Hudson PJ.
2015 Social living mitigates the costs of a chronic illness in a cooperative carnivore.
*Ecol. Lett.* **18**, 660–667. (doi:10.1111/ele.12444)

6. Stewart PD, Macdonald DW. 2003 Badgers and Badger Fleas: Strategies and Counter-
Strategies. *Ethology* **109**, 751–764. (doi:10.1046/j.1439-0310.2003.00910.x)

7. Madden JR, Clutton-Brock TH. 2009 Manipulating grooming by decreasing
ectoparasite load causes unpredicted changes in antagonism. *Proc. R. Soc. B Biol. Sci.*
**276**, 1263–1268. (doi:10.1098/rspb.2008.1661)

8. Poirotte C, Massol F, Herbert A, Willaume E, Bomo PM, Kappeler PM, Charpentier
MJE. 2017 Mandrills use olfaction to socially avoid parasitized conspecifics. *Sci. Adv.*
**3**. (doi:10.1126/sciadv.1601721)

9. Meunier J. 2015 Social immunity and the evolution of group living in insects. *Philos.*
*Trans. R. Soc. B Biol. Sci.* **370**, 19–21. (doi:10.1098/rstb.2014.0102)

10. Cremer S, Armitage SAO, Schmid-Hempel P. 2007 Social Immunity. *Curr. Biol.* **17**,
693–702. (doi:10.1016/j.cub.2007.06.008)

11. Baracchi D, Fadda A, Turillazzi S. 2012 Evidence for antiseptic behaviour towards
sick adult bees in honey bee colonies. *J. Insect Physiol.* **58**, 1589–1596.
(doi:10.1016/j.jinsphys.2012.09.014)

12. Bachorec E *et al.* 2020 Spatial networks differ when food supply changes꞉ Foraging
strategy of Egyptian fruit bats. , 1–15. (doi:10.1371/journal.pone.0229110)

13. Methion S, Díaz López B. 2019 Individual foraging variation drives social
organization in bottlenose dolphins. *Behav. Ecol.* , 1–10. (doi:10.1093/beheco/arz160)

14. Webber QMR, Laforge MP, Bonar M, Robitaille AL, Hart C, Zabihi-Seissan S, Wal E
Vander. 2020 The ecology of individual differences empirically applied to space-use
and movement tactics. The American Naturalist. *Am. Nat.* , 1–4. (doi:10.1086/708721)

15. Webber QMR, Vander Wal E. 2018 An evolutionary framework outlining the
integration of individual social and spatial ecology. *J. Anim. Ecol.* **87**, 113–127.
(doi:10.1111/1365-2656.12773)

16. Becker DJ, Albery GF, Kessler MK, Lunn TJ, Falvo CA, Czirják GÁ, Martin LB,

Plowright RK. 2019 Macroimmunology: the drivers and consequences of spatial

patterns in wildlife immune defense. *J. Anim. Ecol.* **89**, 972–995. (doi:10.1111/1365-

2656.13166)

17.   Albery GF, Kirkpatrick L, Firth JA, Bansal S. 2020 Unifying spatial and social

network analysis in disease ecology. *J. Anim. Ecol.* , 1–17. (doi:10.1111/1365-

2656.13356)

18.   Weinstein SB, Moura CW, Mendez JF, Lafferty KD. 2017 Fear of feces? Tradeoffs

between disease risk and foraging drive animal activity around raccoon latrines. *Oikos*

**00**, 1–8. (doi:10.1111/oik.04866)

19.   Curtis VA. 2014 Infection-avoidance behaviour in humans and other animals. *Trends*

*Immunol.* **35**, 457–464. (doi:10.1016/j.it.2014.08.006)

20.   Stockmaier S, Bolnick DI, Page RA, Carter GG. 2018 An immune challenge reduces

social grooming in vampire bats. *Anim. Behav.* **140**, 141–149.

(doi:10.1016/j.anbehav.2018.04.021)

21.   Weinstein SB, Buck JC, Young HS. 2018 A landscape of disgust. *Science (80-. ).* **359**,

1213–1215.

22.   Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary Consequences

of Parasite Avoidance. *Trends Ecol. Evol.* **xx**, 1–14. (doi:10.1016/j.tree.2018.05.001)

23.   Hopkins SR, Fleming-Davies AE, Belden LK, Wojdak JM, Fleming☐Davies AE,

Belden LK, Wojdak JM. 2020 Systematic review of modeling assumptions and

empirical evidence: does parasite transmission increase nonlinearly with host density?

*Methods Ecol. Evol.* **11**, 0–2. (doi:10.1111/2041-210x.13361)

24.   Ezenwa VO. 2004 Host social behavior and parasitic infection: A multifactorial

approach. *Behav. Ecol.* **15**, 446–454. (doi:10.1093/beheco/arh028)

25.   Downs CJ, Stewart KM, Dick BL. 2015 Investment in constitutive immune function

by north American elk experimentally maintained at two different population densities.

*PLoS One* **10**, 1–17. (doi:10.1371/journal.pone.0125586)

26.   Brown CR, Brown MB. 1986 Ectoparastism as a cost of coloniality in cliff swallows (

Hirundo pyrrhonota). *Ecology* **67**, 1206–1218. (doi:10.2307/1938676)

27.   Rifkin JL, Nunn CL, Garamszegi LZ. 2012 Do Animals Living in Larger Groups

Experience Greater Parasitism? A Meta-Analysis. *Am. Nat.* **180**, 70–82.

(doi:10.1086/666081)

28.  MacIntosh AJJ, Jacobs A, Garcia C, Shimizu K, Mouri K, Huffman MA, Hernandez

AD. 2012 Monkeys in the Middle: Parasite Transmission through the Social Network

of a Wild Primate. *PLoS One* **7**, 15–21. (doi:10.1371/journal.pone.0051144)

29.  VanderWaal KL, Atwill ER, Isbell LA, McCowan B. 2014 Linking social and

pathogen transmission networks using microbial genetics in giraffe (Giraffa

camelopardalis). *J. Anim. Ecol.* **83**, 406–414. (doi:10.1111/1365-2656.12137)

30.  Drewe JA. 2010 Who infects whom? Social networks and tuberculosis transmission in

wild meerkats. *Proc. Biol. Sci.* **277**, 633–42. (doi:10.1098/rspb.2009.1775)

31.  Pullan RL, Sturrock HJW, Soares Magalhaes RJ, Clements ACA, Brooker SJ. 2012

Spatial parasite ecology and epidemiology: a review of methods and applications.

*Parasitology* **139**, 1870–1887. (doi:10.1017/S0031182012000698)

32.  Kirby RS, Delmelle E, Eberth JM. 2017 Advances in spatial epidemiology and

geographic information systems. *Ann. Epidemiol.* **27**, 1–9.

(doi:10.1016/j.annepidem.2016.12.001)

33.  Albery GF, Becker DJ, Kenyon F, Nussey DH, Pemberton JM. 2019 The fine-scale

landscape of immunity and parasitism in a wild ungulate population. *Integr. Comp.*

*Biol.* **icz016**, 1–11. (doi:10.1093/icb/icz016)

34.  Manlove K, Aiello C, Sah P, Cummins B, Hudson PJ, Cross PC. 2018 The ecology of

movement and behaviour: a saturated tripartite network for describing animal contacts.

*Proceedings. Biol. Sci.* **285**, 20180670. (doi:10.1098/rspb.2018.0670)

35.  Richardson TO, Gorochowski TE. 2015 Beyond contact-based transmission networks:

The role of spatial coincidence. *J. R. Soc. Interface* **12**. (doi:10.1098/rsif.2015.0705)

36.  Leu ST, Sah P, Krzyszczyk E, Jacoby A, Mann J. 2020 Sex, synchrony, and skin

contact: integrating multiple behaviors to assess pathogen transmission risk. *Behav.*

*Ecol.* , 1–10. (doi:10.1093/beheco/araa002)

37.  Macdonald DW, Newman C, Buesching CD. 2015 Badgers in the rural landscape—

conservation paragon or farmland pariah? Lessons from the Wytham Badger Project.

In *Wildlife Conservation on Farmland Volume 2*, pp. 65–95.

(doi:10.1093/acprof:oso/9780198745501.003.0004)

38.    McDonald JL, Robertson A, Silk MJ. 2018 Wildlife disease ecology from the
459            individual to the population: Insights from a long-term study of a naturally infected
460            European badger population. *J. Anim. Ecol.* **87**, 101–112. (doi:10.1111/1365-
461            2656.12743)

462    39.    Ham C, Donnelly CA, Astley KL, Jackson SYB, Woodroffe R. 2019 Effect of culling
463            on individual badger Meles meles behaviour: Potential implications for bovine
464            tuberculosis transmission. *J. Appl. Ecol.* **56**, 2390–2399. (doi:10.1111/1365-
465            2664.13512)

466    40.    Tuyttens F a M, Delahay RJ, MacDonald DW, Cheeseman CL, Long B, Donnelly C a.
467            2000 Spatial perturabtion caused by a badger (Meles meles) culling operation:
468            implication for the function of territoriality and the control of bovine tuberculosis
469            (Mycobacterium bovis. *J. Appl. Ecol.* **69**, 815–828.

470    41.    Donnelly CA *et al.* 2007 Impacts of widespread badger culling on cattle tuberculosis:
471            concluding analyses from a large-scale field trial. *Int. J. Infect. Dis.* **11**, 300–8.
472            (doi:10.1016/j.ijid.2007.04.001)

473    42.    Rozins C, Silk MJ, Croft DP, Delahay RJ, Hodgson DJ, McDonald RA, Weber N,
474            Boots M. 2018 Social structure contains epidemics and regulates individual roles in
475            disease transmission in a group-living mammal. *Ecol. Evol.* , 1–12.
476            (doi:10.1002/ece3.4664)

477    43.    Silk MJ, Weber NL, Steward LC, Hodgson DJ, Boots M, Croft DP, Delahay RJ,
478            McDonald RA. 2018 Contact networks structured by sex underpin sex-specific
479            epidemiology of infection. *Ecol. Lett.* **21**, 309–318. (doi:10.1111/ele.12898)

480    44.    Weber N, Carter SP, Dall SRX, Delahay RJ, McDonald JL, Bearhop S, McDonald RA.
481            2013 Badger social networks correlate with tuberculosis infection. *Curr. Biol.* **23**,
482            R915–R916. (doi:10.1016/j.cub.2013.09.011)

483    45.    Macdonald DW, Newman C. 2002 Population dynamics of badgers (Meles meles) in
484            Oxfordshire, U.K.: Numbers, density and cohort life histories, and a possible role of
485            climate change in population growth. *J. Zool.* **256**, 121–138.
486            (doi:10.1017/S0952836902000158)

487    46.    Macdonald DW, Newman C, Dean J, Buesching CD, Johnson PJ. 2004 The
488            distribution of Eurasian badger, *Meles meles*, setts in a high-density area: Field

observations contradict the sett dispersion hypothesis. *Oikos* **106**, 295–307.
(doi:10.1111/j.0030-1299.2004.12879.x)

47.  Macdonald DW, Newman C, Buesching CD, Johnson PJ. 2008 Male-biased
Movement in a High-density Population of the Eurasian Badger (Meles meles). *J.*
*Mammal.* **89**, 1077–1086. (doi:10.1644/07-mamm-a-185.1)

48.  Annavi G, Newman C, Dugdale HL, Buesching CD, Sin YW, Burke T, Macdonald
DW. 2014 Neighbouring-group composition and within-group relatedness drive extra-
group paternity rate in the European badger (Meles meles). *J. Evol. Biol.* , 2191–2203.
(doi:10.1111/jeb.12473)

49.  Cox R, Stewart PD, Macdonald DW. 1999 The ectoparasites of the european badger,
Meles meles, and the behavior of the host-specific flea, Paraceras melis. *J. Insect*
*Behav.* **12**, 245–265.

50.  Anwar MA, Newman C, Macdonald DW, Woolhouse MEJ, Kelly DW. 2000
Coccidiosis in the European badger (Meles meles) from England, an epidemiological
study. *Parasitology* **120**, 255–260. (doi:10.1017/S0031182099005491)

51.  Newman C, Macdonald DW, Anwar MA. 2001 Coccidiosis in the European badger,
Meles meles in Wytham Woods: Infection and consequences for growth and survival.
*Parasitology* **123**, 133–142. (doi:10.1017/S0031182001008265)

52.  Johnson DDP, Stopka P, Macdonald DW. 2004 Ideal flea constraints on group living:
Unwanted public goods and the emergence of cooperation. *Behav. Ecol.* **15**, 181–186.
(doi:10.1093/beheco/arg093)

53.  Buesching CD, Newman C, Service K, Macdonald DW, Riordan P. 2016 Latrine
marking patterns of badgers ( Meles meles ) with respect to population density and
range size. **7**, 1–11.

54.  Franks DW, Ruxton GD, James R. 2010 Sampling animal association networks with
the gambit of the group. *Behav. Ecol. Sociobiol.* **64**, 493–503. (doi:10.1007/s00265-
009-0865-8)

55.  Sin YW, Annavi G, Dugdale HL, Newman C, Burke T, MacDonald DW. 2014
Pathogen burden, co-infection and major histocompatibility complex variability in the
European badger (Meles meles). *Mol. Ecol.* **23**, 5072–5088. (doi:10.1111/mec.12917)

56.  R Core Team. 2020 R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

57.    Lindgren F, Rue H. 2015 Bayesian Spatial Modelling with R-INLA. *J. Stat. Softw.* **63**,
1–25. (doi:10.18637/jss.v063.i19)

58.    Lindgren F, Rue H, Lindstrom J. 2011 An explicit link between Gaussian fields and
Gaussian Markov random fields: the stochastic partial differential equation approach.
*J. R. Stat. Soc. B* **73**, 423–498.

59.    Calenge C. 2011 Home range estimation in R: the adehabitatHR package. See
https://cran.r-project.org/web/packages/adehabitatHR/index.html (accessed on 10
March 2020).

60.    Hadfield JD. 2010 MCMC methods for multi-response generalized linear mixed
models: the MCMCglmm R package. *J. Stat. Softw.* **33**, 1–22. (doi:10.1002/ana.22635)

61.    Brommer JE, Covarrubias-Pazaran G. 2019 Multivariate mixed models in ecology and
evolutionary biology: inference and implementation in R. *EcoEvoRxiv Prepr.*
(doi:10.1017/CBO9781107415324.004)

62.    Strandin T, Babayan SA, Forbes KM. 2018 Reviewing the effects of food provisioning
on wildlife immunity. *Philos. Trans. R. Soc. B Biol. Sci.* **373**.
(doi:10.1098/rstb.2017.0088)

63.    Sweeny AR, Clerc M, Pontifes PA, Venkatesan S, Babayan SA, Pedersen AB. 2019
Supplemented nutrition increases immunity and drug efficacy in a natural host-
helminth system. , 1–43.

64.    Butler JM, Roper TJ. 1996 Ectoparasites and sett use in European badgers. *Anim.*
*Behav.* **52**, 621–629. (doi:10.1006/anbe.1996.0203)

65.    Ezenwa VO. 2016 Helminth-microparasite co-infection in wildlife: lessons from
ruminants, rodents, and rabbits. *Parasite Immunol.* **38**, 527–534.
(doi:10.1111/pim.12348)

66.    Poirotte C, Kappeler PM. 2019 Hygienic personalities in wild grey mouse lemurs vary
adaptively with sex. *Proc. R. Soc. B Biol. Sci.* **286**, 20190863.
(doi:10.1098/rspb.2019.0863)

67.    Newman C, Buesching CD. 2019 Detecting the Smell of Disease and Injury: Scoping
Evolutionary and Ecological Implications. In *Chemical Signals in Vertebrates 14*, pp.

238–250. Cham: Springer International Publishing. (doi:10.1007/978-3-030-17616-
7_17)

68.  Brøseth H, Bevanger K, Knutsen B. 1997          Function of multiple badger
Meles meles          setts: distribution and utilisation          . *Wildlife Biol.* **3**,
89–96. (doi:10.2981/wlb.1997.011)

69.  Tsunoda M, Newman C, Buesching CD, Macdonald DW, Kaneko Y. 2018 Badger
setts provide thermal refugia, buffering changeable surface weather conditions. *J.*
*Therm. Biol.* **74**, 226–233. (doi:10.1016/j.jtherbio.2018.04.005)

70.  Kaneko Y, Newman C, Buesching CD, MacDonald DW. 2010 Variations in badger
(meles meles) sett microclimate: Differential cub survival between main and
subsidiary setts, with implications for artificial sett construction. *Int. J. Ecol.* **2010**.
(doi:10.1155/2010/859586)

71.  Hutchings MR, Judge J, Gordon IJ, Athanasiadou S, Kyriazakis I. 2006 Use of trade-
off theory to advance understanding of herbivore-parasite interactions. *Mamm. Rev.*
**36**, 1–16. (doi:10.1111/j.1365-2907.2006.00080.x)

72.  Robertson A, McDonald RA, Delahay RJ, Kelly SD, Bearhop S. 2014 Individual
foraging specialisation in a social mammal: the European badger (Meles meles).
*Oecologia* **176**, 409–421. (doi:10.1007/s00442-014-3019-2)

73.  Benton CH, Delahay RJ, Robertson A, McDonald RA, Wilson AJ, Burke TA,
Hodgson D. 2016 Blood thicker than water: Kinship, disease prevalence and group
size drive divergent patterns of infection risk in a social mammal. *Proc. R. Soc. B Biol.*
*Sci.* **283**. (doi:10.1098/rspb.2016.0798)

74.  Delahay RJ, Walker N, Smith GS, Wilkinson D, Clifton-Hadley RS, Cheeseman CL,
Tomlinson AJ, Chambers MA. 2013 Long-term temporal trends and estimated
transmission rates for Mycobacterium bovis infection in an undisturbed high-density
badger (Meles meles) population. *Epidemiol. Infect.* **141**, 1445–1456.
(doi:10.1017/S0950268813000721)

75.  Woodroffe R *et al.* 2009 Social group size affects Mycobacterium bovis infection in
European badgers (Meles meles). *J. Anim. Ecol.* **78**, 818–827. (doi:10.1111/j.1365-
2656.2009.01545.x)

76.  Woodroffe R, Donnelly CA, Ham C, Jackson SYB, Moyes K. 2016 Badgers prefer

cattle pasture but avoid cattle□: implications for bovine tuberculosis control. *Ecol.*
*Lett.* **19**, 1201–1208. (doi:10.1111/ele.12654)

77.   Buck JC *et al.* 2017 Host density increases parasite recruitment but decreases host risk
in a snail — trematode system Published by□: Wiley on behalf of the Ecological
Society of America Stable URL□: http://www.jstor.com/stable/26601143 Host density
increases parasite recruit. *Ecology* **98**, 2029–2038.

78.   Mooring MS, Hart BL. 1992 Animal Grouping for Protection from Parasites: Selfish
Herd and Encounter-Dilution Effects. *Behaviour* **123**, 173–193.
(doi:10.2307/4535069)

79.   Calder PC, Jackson AA. 2000 Undernutrition, infection and immune function. *Nutr.*
*Res. Rev.* **13**, 3–29. (doi:10.1079/095442200108728981)

## Supplementary figures and tables: Negative density-dependent parasitism in a group-living carnivore

Figure SI1: Model effect outputs for the full dataset INLA density models. Points represent modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with zero. Only the behavioural traits that improved the model fit (ΔDIC<-2) and were kept in the final model are shown.

Figure SI2: Model effect outputs for the adults-only dataset INLA density models. Points represent modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with zero. Only the behavioural traits that improved the model fit ($\Delta$DIC<-2) and were kept in the final model are shown.

Figure SI3: Model effect outputs for the juveniles-only dataset INLA density models. Points represent

modes of the posterior distribution of the effect estimates; error bars represent the 95% credibility

intervals. Asterisks denote significant results, whose 95% credibility intervals did not overlap with

zero. Only the behavioural traits that improved the model fit ($\Delta$DIC<-2) and were kept in the final

model are shown.

Figure SI4: Model effect outputs for the multivariate MCMCglmm models closely represent the INLA

univariate models. Points represent modes of the posterior distribution of the effect estimates; error

bars represent the 95% credibility intervals. Asterisks denote significant results, whose 95%

credibility intervals did not overlap with zero. Only the behavioural traits that improved the model fit

($\Delta$DIC<-2) and were kept in the final model are shown.

| PARASITE | ROUND | VARIABLE | DELTA | KEPT |
|----------|-------|----------|-------|------|
| **FLEAS** | 1 | Degree | 2.08 | 0 |
| | 1 | GroupSize | 2.055 | 0 |
| | 1 | **LifetimeDensity** | -16.001 | 1 |
| | 1 | TrappingDensity | 1.115 | 0 |
| | 1 | AnnualDensity | -6.889 | 0 |
| **LICE** | 1 | Degree | 1.796 | 0 |
| | 1 | GroupSize | 1.3 | 0 |
| | 1 | LifetimeDensity | 1.694 | 0 |
| | 1 | TrappingDensity | 0.166 | 0 |
| | 1 | AnnualDensity | 1.117 | 0 |
| **TICKS** | 1 | Degree | 0.773 | 0 |
| | 1 | GroupSize | 3.644 | 0 |
| | 1 | **LifetimeDensity** | -2.192 | 1 |
| | 1 | TrappingDensity | 1.007 | 0 |
| | 1 | AnnualDensity | -0.266 | 0 |
| **EIMERIA** | 1 | Degree | 1.134 | 0 |
| | 1 | GroupSize | 1.79 | 0 |
| | 1 | LifetimeDensity | 1.084 | 0 |
| | 1 | **TrappingDensity** | -6.035 | 1 |
| | 1 | AnnualDensity | 1.851 | 0 |
| **ISOSPORA** | 1 | Degree | 1.897 | 0 |
| | 1 | GroupSize | 1.861 | 0 |

| | 1 | LifetimeDensity | 1.873 | 0 |
|---|---|---|---|---|
| | 1 | TrappingDensity | 1.507 | 0 |
| | 1 | AnnualDensity | 1.717 | 0 |

625 Table SI1: DIC Changes (ΔDIC) associated with density measure model addition for the full

626 models. Negative ΔDIC values correspnd to increased model fit. In each round, the variable

627 that increased model the most was retained, and then the process was repeated, until no

628 variables improved the model by decreasing ΔDIC by more than -2. Variables that were

629 retained in each round are highlighted by asterisks.

| PARASITE | ROUND | VARIABLE | DELTA | KEPT |
|----------|-------|----------|-------|------|
| FLEAS | 1 | Degree | 1.835 | 0 |
|  | 1 | GroupSize | 1.251 | 0 |
|  | 1 | **LifetimeDensity** | -8.444 | 1 |
|  | 1 | TrappingDensity | 1.509 | 0 |
|  | 1 | AnnualDensity | -4.549 | 0 |
| LICE | 1 | **Degree** | -2.388 | 1 |
|  | 1 | GroupSize | 1.747 | 0 |
|  | 1 | LifetimeDensity | 1.332 | 0 |
|  | 1 | TrappingDensity | 1.943 | 0 |
|  | 1 | AnnualDensity | 0.944 | 0 |
| TICKS | 1 | Degree | 1.93 | 0 |
|  | 1 | GroupSize | -12.573 | 0 |
|  | 1 | LifetimeDensity | -5.864 | 0 |
|  | 1 | TrappingDensity | 1.2 | 0 |
|  | 1 | **AnnualDensity** | -14.067 | 1 |
| EIMERIA | 1 | Degree | 14.117 | 0 |
|  | 1 | GroupSize | 1.964 | 0 |
|  | 1 | LifetimeDensity | 1.817 | 0 |
|  | 1 | TrappingDensity | -0.283 | 0 |
|  | 1 | AnnualDensity | 1.237 | 0 |
| ISOSPORA | 1 | Degree | 1.569 | 0 |
|  | 1 | GroupSize | 0.882 | 0 |

| | 1 | LifetimeDensity | 1.9 | 0 |
|---|---|---|---|---|
| | 1 | TrappingDensity | 1.65 | 0 |
| | 1 | AnnualDensity | 1.8 | 0 |

633 Table SI2: DIC Changes (ΔDIC) associated with density measure model addition for the
634 adults-only models. Negative ΔDIC values correspond to increased model fit. In each round,
635 the variable that increased model the most was retained, and then the process was repeated,
636 until no variables improved the model by decreasing ΔDIC by more than -2. Variables that
637 were retained in each round are highlighted by asterisks.

| PARASITE | ROUND | VARIABLE | DELTA | KEPT |
|----------|-------|----------|-------|------|
| **FLEAS** | 1 | Degree | 0.985 | 0 |
| | 1 | GroupSize | -2.362 | 0 |
| | 1 | **LifetimeDensity** | -15.209 | 1 |
| | 1 | TrappingDensity | 1.337 | 0 |
| | 1 | AnnualDensity | -10.016 | 0 |
| | | | | |
| **LICE** | 1 | Degree | -2.862 | 0 |
| | 1 | GroupSize | 1.747 | 0 |
| | 1 | LifetimeDensity | -3.907 | 0 |
| | 1 | **TrappingDensity** | -8.087 | 1 |
| | 1 | AnnualDensity | -2.572 | 0 |
| | | | | |
| **TICKS** | 1 | Degree | 1.602 | 0 |
| | 1 | GroupSize | 0.462 | 0 |
| | 1 | LifetimeDensity | -0.244 | 0 |
| | 1 | TrappingDensity | 0.918 | 0 |
| | 1 | AnnualDensity | -0.459 | 0 |
| | | | | |
| **EIMERIA** | 1 | Degree | 1.42 | 0 |
| | 1 | GroupSize | 1.067 | 0 |
| | 1 | LifetimeDensity | -0.127 | 0 |
| | 1 | **TrappingDensity** | -7.465 | 1 |
| | 1 | AnnualDensity | -0.945 | 0 |
| | | | | |
| **ISOSPORA** | 1 | Degree | 1.73 | 0 |
| | 1 | GroupSize | 1.096 | 0 |

| | | | |
|---|---|---|---|
| 1 | LifetimeDensity | 1.775 | 0 |
| 1 | TrappingDensity | 1.567 | 0 |
| 1 | AnnualDensity | 1.709 | 0 |

641 Table SI3: DIC Changes ($\Delta$DIC) associated with density measure model addition for the
642 juvenile-only models. Negative $\Delta$DIC values correspond to increased model fit. In each
643 round, the variable that increased model the most was retained, and then the process was
644 repeated, until no variables improved the model by decreasing $\Delta$DIC by more than -2.
645 Variables that were retained in each round are highlighted by asterisks.

647