Title: Diagnostic Evidence GAuge of Single cells (DEGAS): A flexible deep-transfer learning

framework for prioritizing cells in relation to disease

Authors: Travis S. Johnson[1,2,3], Christina Y. Yu[1,2], Zhi Huang[4], Siwen Xu[5], Tongxin Wang[6],

Chuanpeng Dong[5], Wei Shao[1], Mohammad Abu Zaid[1], Xiaoqing Huang[3], Yijie Wang[6],

Christopher Bartlett[7], Yan Zhang[2,8], Brian A. Walker[9], Yunlong Liu[5,10], Kun Huang[1,3,10,11*], Jie

Zhang[10*]

Affiliations

[1]Department of Medicine, Indiana University School of Medicine

[2]Department of Biomedical Informatics, College of Medicine, The Ohio State University

[3]Department of Biostatistics and Health Data Science, Indiana University School of

Medicine

[4]School of Electrical and Computer Engineering, Purdue University

[5]Center for Computational Biology and Bioinformatics, Indiana University School of

Medicine

[6]Department of Computer Science, Indiana University

[7]Nationwide Children's Hospital

[8]The Ohio State University Comprehensive Cancer Center (OSUCCC - James)

[9]Division of Hematology Oncology, Melvin and Bren Simon Comprehensive Cancer

Center, Indiana University

[10]Department of Medical and Molecular Genetics, Indiana University School of Medicine

[11]Regenstrief Institute

* To whom correspondence should be addressed (jizhan@iu.edu or kunhuang@iu.edu)

Abstract

We propose *DEGAS* (Diagnostic Evidence GAuge of Single cells), a novel deep transfer

learning framework, to transfer disease information from patients to cells. We call such

transferrable information "impressions," which allow individual cells to be associated with

disease attributes like diagnosis, prognosis, and response to therapy. Using simulated data and

ten diverse single cell and patient bulk tissue transcriptomic datasets from Glioblastoma

Multiforme (GBM), Alzheimer's Disease (AD), and Multiple Myeloma (MM), we demonstrate the

feasibility, flexibility, and broad applications of the *DEGAS* framework. *DEGAS* analysis on

newly generated myeloma single cell transcriptomics led to the identification of $PHF19^{high}$

myeloma cells associated with progression.

Keywords

Prognostic models, Survival, Cox proportional hazards, Single cell RNA sequencing, scRNA-

seq, Machine Learning, Deep learning, Transfer learning, Multiple Myeloma, Alzheimer's

Disease

Background

The emergence of single cell RNA sequencing (scRNA-seq) in 2009 has revolutionized the

medical research community with single cell level resolution, providing a much deeper

understanding of transcriptomic heterogeneity in tissues and diseases. Now that scRNA-seq is a

standard part of the biomedical research toolbox, increasing numbers of scRNA-seq studies have

been published [1, 2], and databases have quickly accumulated with scRNA-seq data, such as

Hemberg lab [3], scRNASeqDB [4], SCPortalen [5], Allen Institute Cell Types Database, and the

NCBI Gene Expression Omnibus (GEO) [5]. Many methods have been developed to analyze

scRNA-seq data, the most notable being *Seurat*, which includes ways to cluster and normalize

cell expression as well as perform integrative analysis with other data types (*e.g.,* CITE-seq and

ATAC-seq) [6]. These methods are important for understanding many prognostic and diagnostic

disease attributes in scRNA-seq data. Here we use "disease attributes" as a broad term inclusive

of many types of information and labeling such as diagnostic information, disease subtypes,

disease status, prognostic information like survival, and responses to therapy. For *Seurat* and

similar methods, while cell types/clusters can be identified and associated with disease attributes

[7-10], individual cells are unable to be associated in the same manner. This may result in failing

to identify subsets of cells associated with disease attributes, especially if the disease-associated

cells cluster together with non-disease-associated cells.

Currently, disease associated cell types can be identified by transferring molecular heterogeneity

information from cells to patients using single cell expression deconvolution [11-13]. However,

this approach is limited as it focuses on the changes in relative abundance of subtypes of cells

instead of transcription changes of these cells. The resolution of the cell subtyping is constrained

by the clustering experiment. Therefore, novel machine learning methods that can transfer

information from patients to cells and identify latent links between them are sorely needed to

leverage the relative strengths of single cell and patient level data. For example, in cancer studies,

bulk transcriptomic data is ideal for studying inter-tumor heterogeneity and scRNA-seq is ideal for

studying intra-tumor heterogeneity. However, such integration faces numerous challenges since

different data modalities and different data sources can have different characteristics in terms of

quantity, quality, distribution and resolution [1]. For instance, it is common to find studies with a

large number of patient samples for bulk tissue RNA sequencing (RNA-seq), whereas studies

with scRNA-seq data usually contain a small number of patient samples. Most scRNA-seq

experiments generate a large number of cells per sample, making the scaling of such data to

multiple tissue samples computationally difficult [1]. On the other hand, a large patient sample

size is often required for statistical studies such as prediction of disease attributes [14]. If

traditional methods were used, the resulting scRNA-seq data could end up with cell numbers on

the scale of millions making such studies more difficult.

To address these challenges, previous studies have directly established associations of diseases with cell types derived from scRNA-seq without using deconvolution. These methods mainly utilize unsupervised methods and focused primarily on the number of differentially expressed genes (DEGs) in a given cell type corresponding to DEGs related to some disease attribute [15, 16]. For example, Gawel *et al.* used enrichment of the cell cluster specific DEGs and multicellular disease models (MCDMs) to visualize the cell types for prioritization [7]. *Muscat* identified DEGs between treatment groups in scRNA-seq samples which were used to identify cell types related to sample treatment [17]. Alternatively, k nearest neighbor (*kNN*) graphs have been used to identify cell types that undergo transcriptional changes related to biological perturbations [18]. The cell type prioritization tool *Augur* did not primarily rely on DEGs, but still focused the biological resolution to the cell type level [19]. They trained classifiers on each cell type with respect to the disease state of the tissue from which those cells were sampled. The accuracy of the classifier in each cell type was used to prioritize its relation to the disease state of interest [19]. These methods rely on either prior knowledge to calculate enrichment of DEGs or require scRNA-seq data from both disease and normal samples. Furthermore, all of these existing methods are reliant on accurately defining the cell types within a scRNA-seq experiment. In summary, these methods assign disease associations to the previously defined cell types and not to the individual cells.

To address such challenges as prioritizing individual cells in relation to disease with considerations on sample size and computational cost, we established the combined deep learning and transfer learning framework called *DEGAS* (Diagnostic Evidence GAuge of Single cells) to integrate scRNA-seq and bulk tissue transcriptomic data with the goal to transfer clinical information from patients to cells. The ability of *DEGAS* to assign patient-level disease attributes to single cells, among other functions, provides a flexible and useful tool to prioritize cells, cell types, patients, and patient subtypes in relation to disease attributes. In this paper, we focus on

104 the most relevant use case of associating disease attributes from patients to individual cells since

105 there is no current state-of-the art technique to perform this task.

107 We use transcriptomic data as an example where bulk expression is referred to as patients and

108 scRNA-seq is referred to as cells. The rationale behind the *DEGAS* framework is that scRNA-seq

109 data and patient-level transcriptomic data (*e.g.,* RNA-seq with clinical information) share the same

110 feature space (*i.e.,* common set of genes). In addition, a natural connection exists between the

111 two data types that can be leveraged to further identify the associations between patients and

112 cells. Viewing this association as a graph (**Fig. 1**), we can connect the disease attributes in

113 patients to individual cells, via a latent representation of the common feature space (selected

114 genes). This latent representation fitting two datasets can be learned using a transfer learning

115 technique called domain adaptation [20-23]. Domain adaptation applies linear or non-linear

116 transformations on the features for both datasets so that their distributions are similar after the

117 transformations. Our biological intuition is thus: the expression patterns of genes in cells and

118 tissues should carry a portion of the same biological patterns such as molecular pathways,

119 signaling cascades, and/or metabolic processes, making the information learned from this portion

120 of gene expression patterns transferable between patients and cells. Our hypothesis is that the

121 latent representation learned from these shared gene expression patterns will be simultaneously

122 predictive of patient disease attributes and cellular subtypes. Similar hypotheses are already

123 adopted to transfer information between different single cell experiments [6, 24-28] and to transfer

124 information from bulk transcriptomic cell type atlases to single cell experiments [29].

126 In our *DEGAS* framework, we incorporate patient-level disease attributes information with cell

127 type information from disparate datasets to perform cell prioritization on scRNA-seq data. These

128 disease associations in cells can be attributed to disease-related biological perturbations

129 identified in the patients. This novel deep transfer learning approach simultaneously trains a

model on single cell data and patient data along with their labels and learns a representation in

which the cells and patients occupy the same latent space. Multitask learning, also known as

parallel transfer learning, is precisely designed to achieve these two goals. Used extensively in

computer vision, multitask learning learns a low dimensional representation of the input data to

optimally address multiple tasks. Examples of such application in medical science include

predicting benign versus malignant tumor samples and subclassification in breast cancer

histology images [30, 31]. In this paper, we further extend this line of research to include datasets

with patient disease attributes that can be trained simultaneously so that the disease attributes

can be transferred (or cross-mapped) between single cells and patients. Specifically, our

framework enables knowledge learned from patients using deep learning models to be transferred

to single cells and vice versa. The major advantages of our transfer learning framework are that

the single-cell gene expression data and clinical bulk gene expression data can come from

different patient cohorts of the same disease without matched data while the disease associations

can still be directly assigned to individual cells. This flexibility not only presents an ingenious way

to integrate molecular omics data analysis in different levels, but also virtually merges them into

the same cohort, which makes studying a broad variety of heterogeneous diseases possible.

Various types of workflows can integrate the *DEGAS* framework, which can be tailored to user

preference and data availability. These workflows consist of preprocessing, formatting data,

training *DEGAS* models using the *DEGAS* framework, predicting disease associations in cells

using the *DEGAS* framework, and downstream analysis (**Fig. 1A**). The *DEGAS* framework in its

simplest form can be broken into three tasks during model training: 1) correctly labeling cells with

a cellular subtype using multitask learning; 2) correctly assigning clinical labels to patients using

multitask learning; and 3) generating a latent space in which patients and cells are comparable

using domain adaptation (**Fig. 1B**). To perform *DEGAS* analysis, first we select representative

gene features that are predictive of cell type, predictive of patient disease attributes, and present

at measurable levels in both scRNA-seq and bulk transcriptomic data. Secondly, we apply deep

learning models to learn the latent representation of the single-cell and patient-level transcriptomic

data, with the goal to simultaneously minimize cell type classification error, patient disease

attribute prediction error, and the differences between cells and patients in their latent

representation. Finally, the patient-level disease attributes such as survival and clinical subtypes

is predicted in the single cells using the patient label output layer and cell types are predicted in

patients using the cell type output layer (**Fig. 1C**). We call these transferrable label probabilities

"impressions" since information from gene expression of disparate data types and studies can be

extracted and the characteristics from one data type can be mapped to another. These

impressions of disease attributes in single cells can be wide ranging characteristics of the patient

samples but must be categorical or time to event. The most interesting of them that can be used

in *DEGAS* are disease status, disease subtype, survival, and response to therapy. Disease status,

subtype, and survival were used in our current experiments but there would also be much utility

in identifying cells associated with poor response to treatment as the data become available.

Furthermore, we emphasize the ability to make predictions of patient disease attributes in

individual cells since there is a lack of such method to perform this task to the best of our

knowledge. *DEGAS* is developed as a generalizable model generating deep transfer learning

framework that can be applied to any disease data as long as the data contain clinical information

for a cohort of patients or a separate clustering analysis result on sets of cells from single cell

level omic experiments of the same disease. Since there is not an inherent limitation to the use

of transcriptomic data, *DEGAS* can be potentially expanded to accommodate other modalities of

data with proper normalization steps.

To demonstrate the feasibility and effectiveness of the *DEGAS* framework, we first tested it on

simulated data and glioblastoma (GBM) transcriptomic data, which contain ground-truth labels of

cell types on single cell gene expression data and clinical labels for patient bulk tissue gene

expression data. Then we applied *DEGAS* to multiple Alzheimer's disease (AD) gene expression

datasets from Mount Sinai/JJ Peters VA Medical Center, Allen Institute, Grubmann *et al.* [32], and

Mathys et al. [15] in which certain cell type changes (microglia and neuron) are largely known [33-

39]. Finally, as an exploratory tool, we applied *DEGAS* to study multiple myeloma (MM)

transcriptomic data, where the disease associated subtypes of cells are largely unknown.

MM is a late stage of myeloma that stems from the proliferation of aberrant clonal plasma cells

in the bone marrow that secrete monoclonal immunoglobulins and is the second most common

blood cancer in the United States [40]. Patient level transcriptomic data for MM has been widely

available for some time and has been used to identify subtypes of MM with different prognoses

[41]. However, only recently has scRNA-seq become available for MM [9, 42, 43] and few

studies have identified the most high-risk subtypes of cells [9]. Here we combined our newly

generated late-stage myeloma scRNA-seq data from four local samples and bulk tissue data

from the Multiple Myeloma Research Foundation CoMMpass study, and then applied *DEGAS* to

infer clinical impressions for myeloma cell subtypes and successfully identified a $PHF19^{high}$

myeloma cell subgroup associated with a high-risk of progression.

Methods

Experimental design and datasets

For a *DEGAS* cell prioritization experiment, one scRNA-seq dataset, one bulk expression dataset,

and patient sample labels (matched with the bulk data samples) are required as input. After

feature selection and scaling (see ***Feature selection and scaling***) of the raw input expression

data, there should be two expression matrices with rows corresponding to samples/cells and

matching columns corresponding to genes. The bulk patient sample labels should be one-hot

encoded in a matrix with rows corresponding to each sample and the columns corresponding to

each class of label. For survival sample labels the first column should be time and the second

column should be the even indicator (1 event and 0 censored). If cell labels are also available,

they should also be one-hot encoded with each row corresponding to a cell and each column

corresponding to a class of label. The *DEGAS* models can be trained and predicted on these

formatted data (**Fig. 1A**).

In this study we analyzed simulated data and data from three different diseases, GBM, AD, and

MM, to test the *DEGAS* framework and apply it for novel discoveries. The simulation, GBM, and

AD experiments were primarily used as validation datasets since the ground truth is known. The

simulated data were generated so that cell types are directly related to disease status in patients.

For GBM data, we used scRNA-seq data for five tumors from Patel *et al*. [44] and microarray data

for the GBM TCGA cohort [45] (**Table 1**). For AD data, we used human scRNA-seq from Allen

Institute for Brain Science (AIBS) Cell Types Database (https://celltypes.brain-map.org/) and AD

patient RNA-seq from the Mount Sinai/JJ Peters VA Medical Center Brain Bank (MSBB) study

[46] (**Table 1**).

Table 1. Summary of the clinical features in each patient cohorts used in training. * Final

age category is >90 years.

Glioblastoma Multiforme TCGA	
Feature	**Details**
Sex	74 Male, 37 Female
Age (years)	Range: 14-83, Mean: 56, Median: 58
Clinical GBM subtype	34 Classical, 33 Mesenchymal, 9 Neural, 35 Proneural
Alzheimer's Disease MSBB	
Feature	**Details**
Sex	90 Male, 131 Female

Age (years)	Range: 61-90+, Mean* > 82, Median = 84
AD diagnosis	135 AD, 86 Control
Multiple Myeloma MMRF	
Feature	**Details**
Sex	387 Male, 260 Female
Age (years)	Range: 27-93, Mean: 64, Median: 64
Relapse-free survival time (days)	Range: 13-1753, Mean: 665.4, Median: 629 200 patients progressed

We further expanded our inquiry into MM, which served as a discovery study. Since the plasma

cell subtypes are less understood in relation to MM clinical outcomes, we aimed to identify

subtypes of plasma cells associated with worse prognosis. We first utilized 647 $CD138^+$-

enriched bone marrow patient samples from the Multiple Myeloma Research Foundation

CoMMpass study (MMRF). These data were generated as part of the Multiple Myeloma

Research Foundation Personalized Medicine Initiatives (https://research.themmrf.org). The

dataset consisted of tumor tissue RNA-seq data and corresponding clinical information including

progression free survival (PFS) time and survival status. PFS was defined as the time taken for

a patient to relapse, progress, or die after treatment of the initial tumor. The demographic

information of the MMRF patients are shown in **Table 1**. The first scRNA-seq data used in this

study were generated by us using samples consisting of $CD138^+$ plasma cells purified from

bone marrow from four myeloma patients including two MM patients.

There were six total samples collected from myeloma patients. Of these, four samples passed

initial quality control checks. Sample 1 and 6 were dropped due to sample degradation and data

quality issues. This in turn left with four usable samples, *i.e.,* samples 2, 3, 4, and 5 for our

study. The low number of patients was a good test case considering most scRNA-seq

experiments frequently have few patients. The single cells were sequenced using 10x

Genomics and Illumina NovaSeq6000 sequencer. *CellRanger 2.1.0*

(http://support.10xgenomics.com/) was utilized to process the raw sequence data. Briefly,

CellRanger used *bcl2fastq* (https://support.illumina.com/) to demultiplex raw base sequence

calls generated from the sequencer into sample-specific FASTQ files. The FASTQ files were

then aligned to the human reference genome GRCh38 with RNA-seq aligner *STAR*. The aligned

reads were traced back to individual cells and the gene expression level of individual genes

were quantified based on the number of UMIs (unique molecular indices) detected in each cell.

The filtered gene-cell barcode matrices generated by *CellRanger* were used for further analysis.

A second publicly available myeloma scRNA-seq dataset was used for validation, which

consisted of NHIP (normal control), MGUS (monoclonal gammopathy of undetermined

significance), SMM (smoldering multiple myeloma), and MM [42]. A second bulk tissue dataset

was used for validating the proportional hazards modeling. This dataset consisted of bulk

expression profiling by microarray of CD138+ plasma cells with overall survival (OS) information

for 559 MM patients [41]. The detailed information of the four datasets is shown in **Table 2**.

Table 2. Overview of all datasets used in the analysis. *The simulated patients were

generated from the splatter simulated cells by combining known proportions of cell types.

"None" is used to denote the lack of labels for the cells/samples in a given dataset. [†]Cells were

down-sampled from the total number of cells because some cell types were over-represented.

Study	Dataset	Sample size	Data type	Attribute

Simulation	Simulated cells*	5000 cells	scRNA-seq	Cell type
	Simulated patients*	600 patients	RNA-seq	Disease status
Glioblastoma	Patel *et al*., 2014	532 cells (5 patients)	scRNA-seq (SMART-seq)	None
	TCGA GBM	111 patients	Microarray	GBM subtype
Alzheimer's disease	AIBS	47,396 cells (11 patients)	scRNA-seq (SMART-seq)	Brain cell types
	Grubman *et al*., 2019	13,214 cells (12 patients)	snRNA-seq (10x Genomics)	AD and normal brain cell types
	Mathys et al., 2019	5288 cells† (48 patients)	snRNA-seq (10x Genomics)	AD and normal brain cell types
	MSBB	682 samples (221 patients)	RNA-seq	AD diagnosis
Multiple myeloma	MMRF	647 patients	RNA-seq	PFS
	IUSM	22,968 cells (4 patients)	scRNA-seq (10x Genomics)	Subtype cluster (Subtype 1-5)
	Ledergor *et al*., 2019	13,440 cells (35 patients)	scRNA-seq (MARS-seq)	Malignancy (NHIP, MGUS, SMM, MM)
	Zhan *et al*., 2006	559 patients	Microarray	OS

Transfer learning using DEGAS

Several types of labels including Cox proportional hazards, patient classification, and cell type

classification, along with maximum mean discrepancy (MMD), a technique used to match

distributions across different sets of data [22], were combined to create the multitask transfer

learning framework *DEGAS*.

The first step was to find a set of gene expression features that were both informative of cell type

and of patient disease attribute (*e.g.,* recurrence). The intersection of high variance genes found

in the scRNA-seq and bulk expression data of patient samples are used for further analysis. The

definition of this gene set is up to the user but *Seurat-CCA*, *LASSO* selection, and even statistical

tests such as t-test and f-test can be used to define the gene set. Since these features are the

same between patients and single cells, the patients and cells share the same input layer. This

makes it possible to predict proportional hazard and cell type regardless of the input sample type

(patient or single cell data).

All experiments in this manuscript use a five-bootstrap aggregated three-layer DenseNet-based

implementation of *DEGAS*, but the simplest form of the *DEGAS* framework is a single layer

network. In our description of the overall architecture below (shown in **Fig. 1B,C**), we used a

single layer network for the purpose of simplicity. The following **Eq. 1** can nevertheless be

extrapolated to multiple layers and architectures, some of which we have already included in our

open-source software package. First, a hidden layer was used to transform the genes into a lower

dimension using a sigmoid activation function (**Eq. 1**). Where X represents an input expression

matrix, θ_{Hidden} represents the hidden layer weights, and b_{Hidden} represents the hidden layer bias.

$$f_{Hidden}(X) = sigmoid(X^{\mathrm{T}}\theta_{Hidden} + b_{Hidden}) \qquad \textbf{Eq. 1}$$

Next, output layers were added for both the patient output and for the single cell output. For the

single cells, there could be classification output or no output. No output means there are no known

labels for the single cells to match. Similarly, patients could have Cox proportional hazard output,

classification output, or no output (implying no known labels for patients).

The Cox proportional hazards estimates consisted of a linear transformation to a single output

followed by a sigmoid activation function (**Eq. 2**):

$$f_{Cox}(X) = sigmoid(f_{Hidden}(X)^{\mathrm{T}}\theta_{Cox} + b_{Cox}), \qquad \textbf{Eq. 2}$$

where the variable X represents an input expression matrix, θ_{Cox} represents the Cox proportional

hazard layer weights [47], and b_{Cox} represents the Cox proportional hazard layer bias. The

classification output consisted of a transformation to the same number of outputs as the number

of labels, *i.e.*, patient subtypes, cellular subtypes, using a softmax activation function (**Eq. 3**).

$$f_{Class}(X) = softmax\left(f_{Hidden}(X)^{\mathrm{T}}\theta_{Class} + b_{Class}\right), \qquad \textbf{Eq. 3}$$

θ_{Class} represents the classification layer weights and b_{Class} represents the classification layer bias.

To train the *DEGAS* model, we need to compute three types of loss functions for the Cox

proportional hazards output, classification output, and MMD [22] respectively. The Cox

proportional hazards loss [47] was calculated only for the patient expression data (X_{Pat}) using the

followup period (C), and event status (t) (**Eq. 4**). Similarly, the patient classification loss was only

calculated for the patient data (X_{Pat}) using the patient labels (Y_{Pat}). Alternatively, the cellular

classification loss was only calculated for the single cell expression data (X_{Cell}) and true subtype

label (Y_{Cell}) (**Eq. 5**). The MMD loss was calculated between the patient expression data (X_{Pat}) and

the single cell expression data (X_{Cell}) (**Eq. 6**), which is the key for mapping the distributions of the

data representations between the single-cell and patient bulk tissue data.

$$Loss_{Cox} = \sum_{C(i)=1}\left(f_{Cox}(X_{Pat})_i - \sum_{t_j \geq t_i}(\exp\left(f_{Cox}(X_{Pat})_j\right))\right) \qquad \textbf{Eq. 4}$$

$$Loss_{Class} = \frac{1}{n}\sum_{i=1}^{n}\left(\sum(Y_{type,i} - f_{Class}\left(X_{type}\right)_i)\right) \; where \; type \; \epsilon \; \{Pat, Cell\} \qquad \textbf{Eq. 5}$$

$$Loss_{MMD} = MMD(X_{Cell}, X_{Pat}) \qquad \textbf{Eq. 6}$$

317 Besides the three losses, we also add a L_2-regularization loss term to constrain for the complexity

318 of the model. The overall loss function was the weighted sum of the four types of loss using the

319 hyper-parameters λ_0 (single cell loss function), λ_1 (patient loss function), λ_2 (MMD loss), and λ_3

320 (regularization loss), so that the importance of each loss term and regularization term could be

321 adjusted (**Eq. 7**):

$$Loss_{ClassCox} = \lambda_0 Loss_{Class} + \lambda_1 Loss_{Cox} + \lambda_2 Loss_{MMD} + \lambda_3\|\theta\|_2^2. \qquad \textbf{Eq. 7}$$

324 To address more diverse scenarios, we can also adapt **Eq. 7** for two classification outputs (**Eq.**

325 **8**), a single classification output without patient disease attribute (**Eq. 9**), a single classification

326 output without cell type label (**Eq. 10**), or a single Cox output without cell type label (**Eq. 11**):

$$Loss_{ClassClass} = \lambda_0 Loss_{Class} + \lambda_1 Loss_{Class} + \lambda_2 Loss_{MMD} + \lambda_3 \|\theta\|_2^2, \qquad \textbf{Eq. 8}$$

$$Loss_{ClassBlank} = \lambda_0 Loss_{Class} + \lambda_2 Loss_{MMD} + \lambda_3 \|\theta\|_2^2, \qquad \textbf{Eq. 9}$$

$$Loss_{BlankClass} = \lambda_1 Loss_{Class} + \lambda_2 Loss_{MMD} + \lambda_3 \|\theta\|_2^2, \qquad \textbf{Eq. 10}$$

$$Loss_{BlankCox} = \lambda_1 Loss_{Cox} + \lambda_2 Loss_{MMD} + \lambda_3 \|\theta\|_2^2. \qquad \textbf{Eq. 11}$$

In summary, a common hidden layer was used to merge the single cells and patient data. Next,

an output layer was added to predict the proportional hazards or classes of the patient samples

[47]. The loss function for the proportional hazards prediction or patient classification was back-

propagated across both layers for each patient. The single cells also had an output layer

consisting of a softmax output to predict the cellular subtype of each cell. Error was back-

propagated across both layers from the label output for each cell. Finally, a model was learned

that can model both the single cells and the patients. To perform this task, we utilized the MMD

method [22] to reduce the differences between patients and cells in a low dimensional

representation. Both single cell and patient bulk tissue data were combined into a single group

such that the MMD loss was minimized between patient bulk tissue data and single cell data from

multiple patients. Because there are many different combinations of these outputs, *i.e.*, single cell

output followed by patient output, we implemented ClassCox, ClassClass, ClassBlank,

BlankClass, and BlankCox models based on equations (7)-(11) in the current version but intend

to provide more options in the future.

To keep the analyses consistent, we used the same network architecture and hyperparameters

throughout all of the experiments. Specifically, we used a three-layer DenseNet architecture

bootstrap aggregated five times such that **Eq. 1** would consist of a DenseNet instead of a single

layer feedforward network and five such models were trained. The same set of hyper-parameters

were used in all of the experiments in this study, except for the robustness to hyper-parameters

experiment, where they were intentionally altered to test the influences on the output results.

These are considered the default hyper-parameters in the *DEGAS* package but can be changed.

They are: training steps 2000, single cell batch size 200, patient batch size 50, hidden layer nodes

50, drop-out retention rate 50%, single cell loss weight (λ_0) 2, patient loss weight (λ_1) 3, MMD

loss weight (λ_2) 3, and L_2-regularization weight (λ_3) 3.

Feature selection and scaling

There are already multiple feature selection techniques available in a wide range of general

statistical packages and scRNA-seq packages. For this reason, *DEGAS* does not focus mainly

on feature selection, data cleaning, scRNA-seq clustering, but rather on transferring clinical traits

from patient to cells for the purpose of prioritizing those cells. For these reasons, a wide range of

feature selection techniques can be used before the *DEGAS* framework is applied.

Data from scRNA-seq experiments are generally very sparse. As a result, there are few genes

with viable expression for any given cell. Due to this, it is necessary to perform feature selection

to remove genes that are lowly expressed or have very low variance. When we select for high

variance and expressed genes in the bulk expression data, more genes are filtered out. After the

intersection of these two gene sets of expressed and high variance genes, we are left with less

than 1000 genes. It is worth noting that such number of gene features is comparable to Seurat

analysis, when usually hundreds to a couple of thousand highly variable genes are selected. The

feature selection steps were tailored to each dataset because the data sparsity and variance vary

greatly from one another, thus the tailored selection insured that enough genes with high enough

variability were available to train on. The feature selection steps are described individually in each

of the simulated, GBM, AD, and MM experiment sections.

For each experiment, the final feature scaling steps were consistent. The gene expression was

converted to sample-wise z-scores because it allows the genes to be more comparable between

samples and has been performed in multiple other studies [27, 48-50]. As the input to our deep

learning models, we scaled these z-scores to a range of [0,1]. This form of z-score scaling and

[0,1] scaling is commonly used in machine learning and deep learning to help model training [51-

53]. We follow this same convention for our deep learning models.

Disease association scores

The final *DEGAS* output is either the output of a sigmoid or a softmax activation. For these

reasons, it can be useful to convert the [0,1] label output to an association score which can be

interpreted like a correlation coefficient. For these reasons, the output probability matrix from

DEGAS can be converted to a [-1,1] value using the *toCorrCoeff* function in the *DEGAS* package.

This function transforms the [0,1] output value matrix (P) with k labels to [-1,1] using **Eq. 12**.

$$disease\ association = 2\left(\frac{P-\frac{1}{k}}{2-\frac{2}{k}}+\frac{1}{2}\right)-1 \qquad\qquad \textbf{Eq. 12}$$

Validating DEGAS using Simulated single cell data

First we generated 5,000 single cells in four cell types where the cell type 4 had two subtypes

(cell type 4 disease and cell type 4 normal). Each of these five groups described above contains

1,000 cells. We split randomly these cells into 2 parts with 2,000 cells used for patient bulk tissue

data generation and 3,000 cells to use directly as single cell data. The 2,000 single cells used to

generate 600 patients across three different experiments (designated as simulation 1, 2, and 3)

where in simulation 1 the cell type 1 is associated with disease, in simulation 2 only the cell type

4 disease is associated with disease, and in simulation 3 the entire cell type 4 is associated with

disease. Each patient bulk tissue data was generated by randomly combining 400 single cells

using the proportions in **Table 3**.

Table 3. Patient cellular makeup for simulation experiments. The abbreviations are:

Simulation (sim), Normal (N), and Disease (D). The high-risk cell types are in bold.

	Cell type 1	Cell type 2	Cell type 3	Cell type 4N	Cell type 4D
Patients sim1D	**50.0%**	16.6%	16.6%	16.6%	00.0%
Patients sim1N	25.0%	25.0%	25.0%	25.0%	00.0%
Patients sim2D	25.0%	25.0%	25.0%	00.0%	**25.0%**
Patients sim2N	25.0%	25.0%	25.0%	25.0%	00.0%
Patients sim3D	16.6%	16.6%	16.6%	**30.0%**	**20.0%**
Patients sim3N	25.0%	25.0%	25.0%	25.0%	00.0%

We then performed 10-fold cross validation by training the *DEGAS* ClassClass models using cell

type and disease attribute. A total of 1000 gene features were used during training. We evaluated

the model capacity for mapping patient labels on patients and cell type labels on single cells using

PR-AUC and ROC-AUC. We then recapitulated the known cell type associations in each

simulation by overlaying disease association onto the simulated cells. As a comparison, we also

deconvoluted the patients using the 4 cell types using least squares. Deconvolution should be

able to correctly identify the cells of interest in simulation 1 and simulation 3. In contrast, cell type

prioritization using *Augur* [19] should be able to correctly identify the disease associated cell types

in simulation 2. In the simulation 1 *Augur* experiment, cell type 1, cell type 2, cell type 3, and cell

type 4 normal were randomly assigned to the disease or normal groups. In the simulation 2 *Augur*

experiment, cell type 1, cell type 2, and cell type 3 were randomly assigned to disease or normal

groups. The cell type 4 disease cells were all assigned to the disease group and the cell type 4

normal cells were all assigned to the normal group. In the simulation 3 *Augur* experiment, cell

type 1, cell type 2, and cell type 3 were randomly assigned to disease and normal groups. The

cell type 4 cells assigned to the disease group consisted of 60% cell type 4 normal and 40% cell

type 4 disease and cell type 4 cells assigned to the normal group consisted of 100% cell type 4

normal. These cell type proportions match those in the simulation 3 patients used by *DEGAS*.

The *Augur* output for each cell type is an ROC-AUC score that reflects how much a cell type

changes transcriptionally between disease and normal samples. To make the comparison fair

between our two methods, we use the output of our algorithm scaled from [0,1] where 0.5 implies

no association, 0 implies a negative association, and 1 implies a positive association. ROC-AUC

is on the same scale. In this way we compare the strength of signal between *Augur* and our

method to identify that cell type 4 has cell-intrinsic changes related to disease.

Validating DEGAS using GBM data

The scRNA-seq data from the Patel *et al*. study [44] were downloaded from NCBI Gene

Expression Omnibus (GSE57872). The single cell expression values were previously normalized

to TPM containing 5,948 genes with $mean(\log_2(TMP))>4.5$ retained in the data table. The top 20%

variance genes were retained for training. These values were converted to z-scores then

standardized to a range of [0,1] for each sample. The TCGA GBM microarray expression data

was downloaded from *Firebrowse* (http://firebrowse.org/). Microarray data were used since it

contains more patient samples for training with GBM subtype information than RNA-seq data.

Likewise, the top 20% variance genes were retained for training and these expression values

were converted to z-scores then standardized to a range of [0,1] for each sample. The GBM

subtype labels for the TCGA patients were downloaded from Verhaak *et al*. [54]. The intersection

of genes between single cells and patients (199 genes) were used for the final model training.

Since subtype labels were only available for the GBM patient samples, we trained a BlankClass

DEGAS model (**Eq. 10**). This model minimizes the MMD loss between single cells and patients

while minimizing the classification loss only in GBM patients. We split the dataset into 10 groups

and performed 10-fold cross-validation by leaving out a single patient group during training. After

cross-validation, we converted the [0,1] *DEGAS* output to an association [-1,1] using the *DEGAS*

toCorrCoeff function. These association scores were overlaid on the GBM single cells and now

referred to as GBM subtype association scores because GBM subtype from patients is overlaid

on single cells. We plotted these association scores stratified by GBM subtype for each tumor

individually. We then compared the proportions of these cell types to the previously defined GBM

types from the original publication were marked with red boxes. We also visualized the GBM

subtypes association in single cells by calculating a low dimensional representation using *tSNE*

and overlaying the *kNN* smoothed GBM subtype associations. To make the scatter plots of cells

and patients more informative, *kNN* smoothing was used by averaging each point's GBM subtype

association value with its five nearest neighbors in *tSNE*. The model performance was shown with

the PR-AUC and ROC-AUC for each of the GBM subtype labels in the TCGA patients from cross-

validation.

In a second analysis on the GBM scRNA-seq and bulk expression data, using the same input

features, we overlaid risk derived from the overall survival in the TCGA GBM cohort onto the

individual cells from the Patel et al. study [44]. GBM has an extremely low 5-year survival rate

resulting only three patients being censored. We introduced more censoring in the data by

generating a uniformly distributed random vector of censoring times in the range 1 to 1063 days,

where 1063 days is the 90th percentile of survival times. If the censor time was lower than the

survival time, the patient was censored at that time instead of having an event at their true survival

time. We then trained 10 BlankCox *DEGAS* models based on the patient survival input during 10-

fold cross validation. The output from these *DEGAS* models were *kNN* smoothed based on the

tSNE coordinates using the *DEGAS knnSmooth* function and converted to death associations

using the *DEGAS toCorrCoeff* function. To highlight the differences in death association of cells,

these associations were centered to 0 using the *DEGAS centerFunc* function. We evaluated the

accuracy of the labels in patients using a rank-sum test based on the cox output in the GBM

patients.

Validating DEGAS and exploration using AD data

For AD datasets, we were primarily interested in identifying known relationships between cell

types and AD diagnosis. For these reasons, we downloaded all of the adult Human scRNA-seq

data from the AIBS. Only inhibitory neurons, excitatory neurons, oligodendrocytes, astrocytes,

microglia, and oligodendrocyte progenitor cells (OPCs) were retained in the analysis due to the

extremely low sample sizes for the remaining cell types. The inhibitory and excitatory neuron

groups were merged into a single neuron group. These data were then $\log_2$ transformed,

converted to sample-wise z-scores, and then standardized to [0,1] by each sample. In the primary

analysis, only the top 50 up-regulated DEGs for each cell type (calculated by *Seurat*) were

retained in the single cell data (see **RESULTS**). In a distinct secondary analysis, features were

selected with >25% non-zero samples and top 20% variance genes (see **Supplementary**

Materials). The labels for the single cells consisted of the major cell types listed above. The AD

brain data was downloaded from Mount Sinai/JJ Peters VA Medical Center Brain Bank

(https://www.synapse.org/#!Synapse:syn3157743). Each of the RNA-seq samples were either

from an AD patient's brain sample or a normal control brain sample. The binary disease attribute

of AD case or normal were used as the label for the model. Like in the previous experiment, the

RNA-seq values were $\log_2$ transformed, converted to sample-wise z-scores, and standardized to

[0,1] for each sample. The top 50% variance genes were retained for training to keep the feature

set larger. The intersection of the patient genes and single cell genes (Primary analysis: 169

genes, Secondary analysis: 456 genes) were using to train the final models. Using the cell type

classification for each AIBS single cell and the AD/normal classification for each MSBB patient

we were able to train a *DEGAS* ClassClass model (**Eq. 8**). The performance was evaluated using

10-fold cross-validation by leaving out each group during training once. As in the GBM

experiments, we converted the *DEGAS* output to an association using the *DEGAS toCorrCoeff*

function for each single cell so that each single cell now had an AD association. Correlation

analysis was performed on AD association scores for different cells with each cell type by taking

the median score and calculating the p-value by treating it as a correlation. In addition, single cells

were plotted overlaid with *kNN* smoothed AD association. Furthermore, to evaluate *DEGAS*

performance, PR-AUC and ROC-AUC were computed for the single cells during cross-validation

for each cell type in the single cell data. Similarly, AD diagnosis PR-AUC and ROC-AUC were

computed from the MSBB patient RNA-seq. For both the primary and secondary AIBS analysis,

DEGs were identified for the high AD association astrocytes and microglia based on the median

AD association then compared to their respective disease associated astrocyte (DAA) [55]

(**Supplementary File 1**), human Alzheimer's microglia (HAM) gene markers [56]

(**Supplementary File 2**), or disease associated microglia (DAM) gene markers [57]

(**Supplementary File 3**). A detailed description of these gene lists can be found in the

Supplementary Materials DAA, HAM, and DAM markers section.

To further highlight the cellular associations to AD, we also performed experiments using a

scRNA-seq dataset from Grubman *et al.* [32]. Since this dataset was sparser, genes were used

with >25% non-zero samples then the top 50% variance genes were selected from these. For the

MSBB data, the same initial feature selection was used (top 50% variance). The same

normalization and standardization procedure as the AIBS scRNA-seq and MSBB were used

again. The intersecting genes between Grubman *et al.* scRNA-seq constituted the final feature

set (61 genes). 10-fold cross validation was performed using a ClassClass model and the AD

associations were overlaid onto the Grubman *et al.* scRNA-seq in the same fashion as the

previous experiment. In addition, a targeted analysis on only the microglia cells was performed.

A single BlankClass model was trained using the same 61 features on the entire Grubman *et al.*

microglia scRNA-seq and MSBB RNA-seq. For both analyses, the AD associations were overlaid

onto the cells, AD associations were compared between cells from AD and normal patient

samples, and DEGs were identified for the high AD association astrocytes and microglia based

on the median AD association then compared to their respective DAA [55] (**Supplementary File**

1), HAM gene markers [56] (**Supplementary File 2**), or DAM gene markers [57] (**Supplementary**

File 3). For the targeted analysis on only microglia, correlation tests were performed between AD

associations and HAM gene markers [56] (**Supplementary File 2**). Also, DEGs were identified

for the high AD association microglia based on the median AD association then compared to the

HAM gene markers [56] (**Supplementary File 2**) and DAM gene markers [57] (**Supplementary**

File 3).

Lastly, *DEGAS* analysis was performed on the Mathys *et al.* scRNA-seq dataset [15]. In this

analysis, the same gene set as the AIBS Primary analysis, *i.e.,* all overlapping genes (157 genes)

were used as input features. 1000 cells or all cells if total number was less than 1000 were

sampled from each cell type since some cell types were over-represented. The same

normalization and standardization procedure was used as the previous analyses. 10-fold cross

validation was performed using these cells from Mathys et al. and the MSBB patient RNA-seq

data using cell type and patient AD status as outcomes respectively. These outcomes represent

a ClassClass DEGAS model. From the cross-validation results, the ROC-AUCs and PR-AUCs for

each cell type label and the patient AD status were calculated. AD associations were calculated

in the same fashion as all previous analyses. The Disease associations were then compared with

AD status of the scRNA-seq donors and across the cell types. DEGs were identified for the high

AD association astrocytes and microglia based on the median AD association then compared to

their respective DAA [55] (**Supplementary File 1**), HAM gene markers [56] (**Supplementary File**

2), or DAM gene markers [57] (**Supplementary File 3**).

Preprocessing of MM scRNA-seq

The scRNA-seq data were first combined into a dataset using *Seurat-CCA* [28]. This initial dataset

integration allowed conserved subtypes of cells to be identified across datasets. All four patient

dataset counts were loaded into a *Seurat* object. *Seurat* normalized, scaled, removed poor quality

cells, and identified high variance genes. Using the union of high variance genes, multi-canonical

correlation analysis was run across all four datasets, the subspaces were aligned across patients,

the aligned single cells were plotted with *tSNE* [58], and clusters of cells were identified. The raw

expression values for the high variance genes identified by *Seurat* were $\log_2$ transformed,

converted to z-scores, and then scaled to [0,1].

Furthermore, each IUSM scRNA-seq patient was individually clustered using *Seurat* to check the

replicability of the clusters and were plotted with *UMAP* [59]. We used Rand, Fowlkes and

Mallows's index (FM), and Jaccard index (JI) to measure the cluster consistency between single

patient clustering experiments and the merged all-patient clustering results. The four single

patient clustering results, one for each IUSM scRNA-seq patient, were used as input into

BERMUDA [25] to visualize and evaluate the original Seurat clustering.

Preprocessing of MMRF patient data

MMRF patients with bulk tissue RNA-seq and clinical data were used in MM analysis. We used

PFS as the disease attribute of interest. TPM values for the MMRF patient gene expression data

and the PFS data were used as the input for *DEGAS*, these values were $\log_2$ transformed,

converted to z-scores, and scaled to [0,1]. The union of the features (502 genes) identified by

Seurat in the single cell data and the features selected in the MMRF patient data were used as

the final feature set. The features retained in the MMRF data were identified by fitting an elastic-

net Cox model [60] to the TPM values based on the PFS.

Evaluate DEGAS performance on MM datasets

PR-AUC and AUC were calculated for each of the output labels for the single cells and for patient

labels if a classification output was used for the patient data. Cox proportional hazard output was

used on patients, a log-rank test was calculated for each patient so that the hazard ratio and p-

value could be evaluated based on patient stratification by median proportional hazard.

Additionally, the same models were used to predict risk in the GSE2658 dataset which had

information on OS. The output for each GSE2658 sample averaged across all 10 *DEGAS* models

and stratified by median risk to show the robustness of the cox output across datasets.

Identification of CD138+ cell types associated with MM prognosis

The single cells from MM patients can be assigned proportional hazards based on the MMRF Cox

output of the model. Each single cell in the validation set was assigned progression association

by feeding those samples through the Cox output layer. In this way, we can infer the association

with progression risk of specific cell types as well as the cell type enrichment contained in each

MMRF sample. Since the Cox output is a proportional hazard, we centered the outputs to zero

for each step of cross validation to produce a PFS association using the *DEGAS centerFunc*. We

plotted these relationships and conducted Student's t-tests on the subtype vs. PFS association in

IUSM single cells, PFS association vs. MM malignancy from Ledergor *et al*., and subtype 2

enrichment vs. MM malignancy from Ledergor *et al* [42].

Analysis of differential gene expression in prognostic cell types

T-tests were calculated cell subtype 1 vs all cell subtypes and cell subtype 2 vs. all cell subtypes

using the batch corrected gene expression values from *Seurat*. These values were stored in

(**Supplementary File 4** and **Supplementary File 5**) respectively. For the marker set of *PHF19*,

HELLS, *EZH2*, *TYMS*, *ZWINT*, and *MKI67* we performed t-tests for each patient individually.

Evaluation of DEGAS robustness to hyper-parameters in GBM

DEGAS 25

Using the GBM dataset, we evaluated the robustness of DEGAS model outputs to hyper-

parameters by repeating 10-fold cross-validation 100 times with randomly generated hyper-

parameters following a uniform distribution. The range of hyper-parameters used in training

consisted of training steps 1,000-3,000, single cell batch size 100-300, patient batch size 20-100,

hidden features 10-100, drop-out retention rate 0.1-0.9, Cell loss weight (λ_0) held at constant 2,

Patient loss weight (λ_1) 0.2-5, MMD loss weight (λ_2) 0.2-5, L_2-regularization weight (λ_3) 0.2-5.

Using these outputs, we performed two tests. One was to evaluate the loss in performance based

on changing the hyper-parameters where performance was measured with ROC-AUC among the

TCGA GBM patients labeled by patient GBM subtype (Mesenchymal, Classical, Proneural,

Neural). In this test, we calculated the spearman correlation and plotted the scatter plot between

the AUC of each of the four GBM subtype labels and the hyper-parameters used.

Next, we evaluated whether or not the correct GBM subtype labels (Mesenchymal, Classical,

Proneural, Neural) could be recapitulated in the GBM scRNA-seq tumors that had known GBM

subtypes (MGH26: Proneural, MGH28: Mesenchymal, MGH29: Mesenchymal, MGH30:

Classical). To do this for each tumor (MGH26, MGH28, MGH29, MGH30), the rank of the

correct label was calculated by calculating the mean of each GBM subtype association across

all of the cells in that tumor. This resulted in each of the 100 random hyper-parameters having a

rank for each GBM subtype for each of the GBM scRNA-seq tumors (4 highest ranked, 1 lowest

ranked). Ideally all GBM scRNA-seq tumors would have a rank of 4 indicating the correct GBM

subtype was ranked the highest regardless of hyper-parameters. Similarly, we also calculated

the spearman correlation and plotted the scatter plot between correct label rank and the hyper-

parameters used.

Evaluation of domain adaptation for DEGAS disease association transfer

We evaluate the necessity for domain adaptation to transfer disease associations to single cells

using 30 total experiments. These experiments evaluated disease associations in cells by

training with MMD loss vs. those without MMD loss for a variety of biases added between the

cells and patients. It is important to highlight the fact that without bias between different

datasets, in this case cells and patients, there is no need for domain adaptation. Practically in

real transcriptomic data, there will always be bias between datasets. For these reasons we

added bias for these 30 experiments. These experiments were conducted for every combination

of MMD loss (with and without MMD), simulation (three simulations), and cellular subtype (five

total subtypes since cell type 4 has two subtypes) totaling 30 combinations. The experiments

were conducted as follows. In each experiment, the counts of 300 cells from a given subtype

were aggregated together and multiplied by 1000 constituting a large systematic bias

associated with a single subtype. This bias vector was added to all of the patients in the given

simulation, both disease and normal. A single three-layer DenseNet *DEGAS* model with five-fold

bootstrap aggregation was trained on all the cells and all the patients then the disease

associations were predicted in the cells. We evaluated error by subtracting the expected

disease association from the predicted disease associations, *e.g.,* cell type 1 in simulation 1

should be 1. We then compared the error rates between the *DEGAS* models with and without

MMD using a t-test.

Evaluation of regularization in DEGAS performance

Regularization is an important method in machine learning to prevent model overfitting. Here we

utilized three such techniques to prevent overfitting, namely, L_2-regularization, dropout, and

bootstrap aggregation. Since all of these techniques may work better or worse in different

scenarios, we perform a simple experiment where all of these regularization techniques are

removed and compared with the regularized results. We performed experiments using each of

the simulated datasets. To evaluate the robustness of our models we performed 10-fold cross

validation in each simulation. The simulated cells were split into 10 groups and the simulated

patients were split into 10 groups. For each fold of cross validation, our default *DEGAS* three-

layer DenseNet model with L_2-regularization, dropout, and bootstrap aggregated 5 times was

trained then a three-layer DenseNet *DEGAS* model was trained on the same data without L_2-

regularization, dropout, and bootstrap aggregation. Both models were then used to predict the

patient disease attributes in the holdout group of patients, the cell types in the hold out group of

cells, and the patient disease attributes in the cells. We compare the performances using ROC-

AUC and PR-AUC for patient disease status in patients and cell type in cells. Furthermore, we

evaluate the label transfer of patient labels to cells by calculating the error based on the

expected cell type association for each cell. We compare between the regularized and

unregularized error in cells with a t-test.

Results

DEGAS clinical impression framework

In this study, we applied *DEGAS* to integrate and analyze scRNA-seq, bulk gene expression, and

clinical data (**Fig. 1**) from simulated data as well as three different diseases: GBM, AD, and MM.

The simulated, GBM, and AD datasets primarily served as validation to demonstrate the feasibility

and universality of the *DEGAS* transfer learning approach since the ground truth of the simulated

data was known, the correct GBM subtypes were known, and neuron loss with microglia gain in

AD brains were also known. We then further expand our study to MM data, which serves as the

discovery dataset, since the myeloma cell subtypes and high-risk factors related to MM are not

as well understood at the single-cell level. In the MM study, we applied *DEGAS* on patient data

from the Multiple Myeloma Research Foundation CoMMpass study (MMRF) and scRNA-seq data

that we generated from myeloma patients. Our aim was to identify the cell subtypes using the

impressions of progression risk on the single cells. We then applied the results to two separate

MM validation datasets, one of which contained plasma cells from normal bone marrow (NHIP),

two MM precursor conditions - monoclonal gammopathy of undetermined significance (MGUS)

and smoldering multiple myeloma (SMM), and MM. We tested if *DEGAS* assignment of

progression risk to cell subtypes were higher for more malignant conditions. An additional external

validation dataset of patient level expression data with OS was used to evaluate whether the

patient stratification learned by *DEGAS* was robust enough to be generalized to an external

survival dataset.

DEGAS correctly identifies high-risk cell types and subtypes in simulated data

To evaluate *DEGAS* in a controlled context, 5,000 single cells were generated with *Splatter [61]*

(**Fig. 2A**) where 2,000 of the cells were held-out to generate simulated patients. Using this group

of held-out cells, 600 simulated patients were generated by aggregating sets of 400 simulated

cells (**Fig. 2B-D**). We conducted three simulation experiments, denoted Simulation 1, Simulation

2, and Simulation 3, where the single cells were aggregated in known proportions for each patient

so that we could generate a "disease" patient group with different cellular composition than the

"normal" patient group (**see Methods**). To highlight the utility of *DEGAS*, the experiments were:

Simulation 1: cell type 1 is enriched in disease patients (**Fig. 2B**); Simulation 2: one subtype of

cell type 4, *i.e.,* cell type 4 disease, is enriched in disease patients (**Fig. 2C**); and Simulation 3:

both subtypes of cell type 4 are enriched in disease patients (**Fig. 2D**).

Please note that the optimal number of clusters for the simulated single cells would be

determined to be four based on a standard scRNA-seq workflow (*i.e., tSNE* followed by *K-*

Medoids where optimal cluster number is selected based on average silhouette width) (**Fig. 2E**).

This would cluster the cells into the four cell types while ignoring the two subtypes in the cell

type 4 (**Fig. 2F**). As a result, deconvolution algorithms will not be able to detect the subtype

level risk associations. Fortunately, cell type prioritization algorithms like *Augur* can detect these

changes within cell types due to disease. However, for situations that do not have a new cell

type or missing cell type in the disease (simulation 1), *Augur* cannot detect the association

between cell type 1 and disease since there is no disease associated cell type change (**Fig.**

2G). *Augur* can detect the disease-associated cell type 4 in simulation 2 (**Fig. 2H**). In simulation

3 where there is a mix of disease and normal subtypes for the cell type 4 in the disease group,

Augur again has difficulty in identifying the cell type 4 disease association (**Fig. 2I**). In contrast

to *Augur*, deconvolution can easily identify the correct cell type for Simulation 1 (**Fig. 2J**) and

Simulation 3 (**Fig. 2L**) but not for simulation 2 (**Fig. 2K**). In comparison, *DEGAS* not only

identified the correct cell type and subtypes in each experiment, it also correctly detected all of

the simulated disease associations (**Fig. 2G-L**). Additionally, *DEGAS* had high precision-recall

area under the curve (PR-AUC) predicting disease status of simulated patients (0.96-0.98)

(**Table S1**) and almost perfectly predicted the cell type of simulated cells (~1.0) during cross-

validation (**Table S2**). Since *DEGAS* directly assigns disease risk to cells, many of the problems

with cell type level analyses can be avoided and the correct groups of cells can be identified by

overlaying impressions of disease risk.

DEGAS correctly mapped single cells to corresponding GBM subtypes

We first demonstrate *DEGAS* in a straightforward case to show the performance of our framework

using real data from GBM. We use single-cell data from Patel *et al*. [44], in which researchers

assigned four major GBM tumor subtypes (Proneural, Mesenchymal, Classical, and Neural) to

the scRNA-seq data obtained from five GBM tumors. Of the five tumor samples, four had been

labeled in the original publication with a single subtype based on the major proportion of cells

assigned to each GBM subtype. For GBM bulk tumor tissue expression data, we obtained

microarray data for 111 GBM patients from The Cancer Genome Atlas (TCGA), for which the

same labels of GBM subtypes were also provided. The OS was also available in a subset of 109

patients. As the simplest form of validation, we used these two datasets as input for the *DEGAS*

model to test if it could re-identify the same GBM subtypes for both single cells and the TCGA

GBM cohort simultaneously. Then we overlaid OS-derived death associations onto the cells to

visualize their association with OS. The resulting *DEGAS* models also proved to be accurate with

high PR-AUCs (0.79-0.97) when predicting each of the GBM subtypes in the TCGA patients

during 10-fold cross validation (**Table S3**). The OS BlankCox *DEGAS* models were able to stratify

the patients into high and low risk groups based on median patient risk (log-rank p-value < 0.05).

DEGAS correctly re-identified the same labels for all four tumors by overlaying GBM subtypes

associations on each single cell, as indicated by the groups of cell subtypes with the highest

association score determined by the median value (indicated with a red box) (**Fig. 3A-D**). For the

fifth tumor sample, MGH31, it was labeled as a combination of multiple GBM subtypes in the

original study, so we did not use it in our evaluation although *DEGAS* identified mesenchymal as

its most associated GBM subtype (**Fig. 3E**). Additionally, these relationships can be visualized by

plotting the single cells and overlaying the GBM subtype association or OS-derived death

association. It is clear that MGH28 and MGH29 have a high association with the mesenchymal

GBM subtype (**Fig. S1A**) and contain populations of cells with high death associations (**Fig. 3F**).

DEGAS identifies increased microglia, reduced neuron populations, DAAs and DAMs

Aside from GBM, AD also has well documented characteristics that can be used as a test bed for

DEGAS. Specifically, there is a well-documented reduction in neurons [36-38], increase in

microglia [33-35, 39], and more recently, AD subtypes of astrocytes [55] and microglia [56, 57].

AD brain scRNA-seq data was obtained from the AIBS and bulk AD RNA-seq were retrieved from

MSBB [46]. During 10-fold cross-validation, *DEGAS* models for both primary and secondary AIBS

analyses achieved high AD diagnosis status PR-AUC (0.82 and 0.76) in MSBB patients **(Table**

S4) and high cell type prediction PR-AUCs (>0.99) for AIBS single cells (**Table S5**).

From the AIBS primary analysis *DEGAS* results, we confirmed that at the single cell level, the

AD associations were negative in neurons as previously described [62], which is shown by the

dark shade of neurons compared to other cell types (**Fig. 3G**, **Table 4**). In contrast, we

observed positive AD associations in microglia cells (**Fig. 3G**, **Table 4**). A strength of the

DEGAS framework is that it can detect intra-cell type differences in disease risk. Within the

astrocyte cell type, we identified an astrocyte subtype that had a positive association with AD

(**Fig. 3G**) that corresponded to the *Astro L1 FGFR3 FOS* subtype from the AIBS brain cell atlas

(*i.e., FOS* is a DAA marker) [63] (**Fig. 3H**), had up-regulated DAA marker *GFAP* (**Fig. 3I**) [55],

and was enriched for DAA markers (OR = 30.93, Fisher's exact p-value < $2.2 \cdot 10^{-16}$, **Table S6**).

Furthermore, the high AD association microglia were enriched for DAM markers (OR = 17.07,

Fisher's exact p-value = $2.11 \cdot 10^{-10}$, **Table S7**). In the secondary AIBS analysis using high

variance genes, we again identified the strong negative AD association for neurons (**Fig. S2A**),

positive AD association in microglia (**Fig. S2A**), high AD association astrocytes enriched for

DAA markers (OR = 5.65, Fisher's exact p-value < $1.66 \cdot 10^{-8}$, **Fig. S2B,C, Table S8**), and high

AD association microglia enriched for DAM markers (OR = 14.34, Fisher's exact p-value <

$4.01 \cdot 10^{-11}$, **Table S9**). When we performed *DEGAS* analysis on a separate dataset from

Grubman et al. [32] with single cells from both AD and normal brains, we found that the major

cell types from AD brains were significantly more associated with AD than their counterparts in

normal brains as judged by median value (**Fig. 3J**).

Table 4 Comparison of AD association scores in single cells between cell types as

visualized in Fig. 3G. The *DEGAS* models were trained using neuron, oligodendrocyte,

astrocyte, OPC, and microglia cell types. The single cells were split into groups based on their

cell type and the mean AD associations of each cell type was evaluated as a correlation. The

neuron and microglia groups are bolded to highlight their much higher mean AD association. P-

values are calculated by treating the association score as a pearson correlation coefficient.

Cell type	Cell-type mean association	Number of cells	p-value

Neuron	**-0.35**	**1329**	**<2.2•10^{-16}**
Oligodendrocyte	0.05	1795	3.42•10^{-2}
Astrocyte	0.03	809	3.94•10^{-1}
OPC	-0.12	738	1.09•10^{-3}
Microglia	**0.22**	**741**	**1.42•10^{-9}**

In the Grubman *et al.* scRNA-seq data, the astrocytes in AD brains were highly positively

associated with AD (AD association = 0.22, pearson correlation p-value = 7.89•10^{-7}) whereas

the astrocytes in normal brains were negatively associated with AD (AD association = -0.06,

pearson correlation p-value = 1.06•10^{-2}, **Fig. 3J**). Astrocytes from AD brains also expressed

GFAP at greater levels than astrocytes from normal brains (t-test p-value < 2.20•10^{-16}) and high

AD association astrocytes were significantly enriched for DAA markers (OR = 21.90, Fishers

exact p-value = 2.21•10^{-12}, **Table S10**). Furthermore, the high AD association microglia were

moderately enriched for DAM markers (OR = 4.15, Fishers exact p-value = 4.11•10^{-2}, **Table**

S11). This provides evidence for DAA and DAM cells in the Grubman *et al.* dataset.

DAM and HAM marker enriched high AD association cells were independently identified in the

targeted analysis of the Grubman *et al.* microglia cells (**Fig. 3K**). AD associations were higher in

cells derived from AD patient samples than Normal patient samples (**Fig. 3L**, t-test p-value =

6.66•10^{-12}), HAM up-regulated markers were more likely to be significantly positively correlated

to AD association than HAM down-regulated markers (**Fig. 3M**, t-test p-value = 2.63•10^{-3}). The

HAM marker APOE [56, 57] was positively correlated with AD association (**Table S12,**

PCC=0.18, p-value = 1.15•10^{-4}). High AD association microglia were significantly enriched for

HAM markers (OR = 21.47, Fishers exact p-value = 6.33•10^{-4}, **Table S13**) and DAM markers

(OR = 11.52, Fishers exact p-value = 1.50•10^{-11}, **Table S14**). It is important to note that there

was no overlap between the input feature set used to train the DEGAS model and HAM marker

DEGAS

genes that were identified, which shows DEGAS is a useful tool to identify disease associated

cells within a single cell type even without prior knowledge of marker genes.

After applying *DEGAS* to the Mathys *et al.* scRNA-seq dataset, the DEGAS models achieved

high AUCs for patient AD status (0.77), patient AD status PR-AUC (0.81), cell types (>0.98), as

well as cell type PR-AUCs (0.82-0.99) during cross validation (**Table S15-16**). The positive AD

association of microglia and negative AD association of neurons were recapitulated (**Fig. S3A,**

Table S17). Within the astrocyte cluster, there existed a subset of astrocytes with higher AD

association (**Fig. S3B**). High AD association astrocytes were significantly enriched for DAA

markers (OR = 14.75, Fishers exact p-value = $3.16 \cdot 10^{-15}$, **Table S18**). A closer comparison of

the scRNA-seq revealed that the top 10% AD association astrocytes, had 2.5 times higher

GFAP expression than the other astrocytes (t-test p-value = $6.36 \cdot 10^{-8}$, **Fig. S3C**). In fact, like

the Grubman *et al.* analysis, the AD association scores were higher in cells coming from AD

patients than normal patients for every cell type in the Mathys *et al.* analysis (**Table S17**).

Notably, we see increased AD association in AD derived astrocytes and microglia likely

representing DAAs and HAMs respectively (**Table S17**). Furthermore, high AD association

microglia were highly enriched for DAM markers (OR = 19.35, Fishers exact p-value < $2.2 \cdot 10^{-16}$,

Table S19) and high AD association in astrocytes correlated well with neuritic plaque count, a

marker for disease severity in AD patients (PCC = 0.22, p-value = $6.36 \cdot 10^{-12}$, **Table S17**).

Again, the Mathys *et al.* analysis provides another example to demonstrate that DEGAS

recapitulates the findings from the AIBS and Grubman *et al.* analyses and shows that DEGAS

models can capture cell type level as well as intra-cell type differences in disease association.

Identification of plasma cell subtypes in CD138+ scRNA-seq of MM

In the MM study, unlike the previous two datasets, there were no predefined cell type labels, but

DEGAS was still capable of analyzing such data and give clinical perspective to the clusters of

cells in the MM scRNA-seq data. In order to cluster cells into groups, we first used Seurat [28], a

commonly used scRNA-seq data analysis tool, to merge and cluster all the CD138+ bone marrow

cells from four patients (two SMM and two MM) whose samples were collected at the IUSM. Using

Seurat, five major clusters of cells were identified (**Fig. 4A**). Cluster 1 consisted of the majority of

the cells in each sample and was most likely the main clone in each of the patients. Cluster 2 was

present in many of the patients and is described in detail after the *DEGAS* analysis. Cluster 3 and

5 were only present in patient 2 representing possible subclones in patient 2. Cluster 4 was shared

between multiple patients. These five clusters were used as the subtype labels in the *DEGAS*

framework. We verified these cell clusters by clustering cells from each patient individually with

Seurat and another scRNA-seq normalization tool *BERMUDA* (Batch Effect ReMoval Using Deep

Autoencoders) [25] for all four patients. We found that the individual clustering results closely

mirrored the *Seurat-CCA* clusters (**Fig. S4A-D**, **Table S20**) and that the subtype 2 was consistent

across all MM patients using *BERMUDA* (**Fig. S4E**). For bulk tissue data from MMRF, the clinical

outcomes of PFS for 647 patients were used as the patient-level input to *DEGAS* and overlaid

onto the CD138+ single cells from the four IUSM patients (**Fig. 4B**).

DEGAS patient stratification and cell type classification on MM

A *DEGAS* model was trained on IUSM patient scRNA-seq data with subtype labels defined above

and MMRF patients with bulk tissue data and PFS information. The performance metrics were

calculated via 10-fold cross-validation. It is worth noting that for PR-AUC, random no skill

classifiers will achieve a performance equal to the percentage of the class of interest and in the

case of uncommon classes like subtype 4, the random classifier performance will be close to zero

(0.02). When predicting cellular subtype label in single cells, *DEGAS* was able to achieve a PR-

AUC between 0.44-0.98 for all of the five CD138+ cellular subtypes identified in the above scRNA-

seq data while the PR-AUC for subtype 2 reached 0.91 (**Table S21**). The receiver operating curve

AUCs (ROC-AUCs) were between 0.90-0.98 for these five subtypes (**Table S21**). Due to class

DEGAS   35

imbalance some of the subtypes did not perform as well as others based on PR-AUC but all of

the PR-AUCs were substantially greater than a purely random model. Aside from correctly

classifying the single cells, *DEGAS* was able to stratify the MMRF patients into high and low risk

groups based on median progression risk (log-rank p-value = $4.72 \cdot 10^{-10}$, **Fig. 4C**). We then

applied the trained model on an external patient transcriptomic dataset from Zhan *et al*. [41] for

validation. We demonstrated that the Cox proportional hazards portion for patient OS time of the

DEGAS model was robust across datasets, and the impressions extracted from the *DEGAS*

framework were capable of stratifying patients into low- and high-risk groups in the validation

dataset (log-rank p-value = $1.12 \cdot 10^{-3}$, **Fig. 4D**).

DEGAS identifies CD138+ cellular subtypes with high progression association

The MM scRNA-seq data provided an example of an exploratory analysis with *DEGAS* which

can be used to generate hypotheses for future studies. The *DEGAS* model for the MM study

transfers clinical impressions to single cells (*i.e.,* single cells were directly assigned a

progression association score), as well as transfers cellular/molecular impressions to patients

(*i.e.,* patients are assigned subtype enrichment score). We found that the subtype 2 cells were

the most associated with prognosis (**Fig. 4B**) based on the *DEGAS* results. Specifically, the

subtype 2 cells were associated with a shorter time to progression (**Fig. 4E**, t-test p-value <

$2.2 \cdot 10^{-16}$). On an external validation scRNA-seq dataset from Ledergor *et al.* [42], the

progression association increased from NHIP (no disease) to SMM (**Fig. 4F**, t-test p-value =

$1.50 \cdot 10^{-2}$) and MM (**Fig. 4F**, t-test p-value = $1.70 \cdot 10^{-2}$), which is consistent with the order of

precursor conditions for MM (NHIP $\rightarrow$ MGUS $\rightarrow$ SMM $\rightarrow$ MM). In addition, the enrichment of the

subtype 2 cells increased from NHIP to near-MM stage SMM (**Fig. 4G**, t-test p-value = $3.10 \cdot 10^{-2}$

) and MM (**Fig. 4G**, t-test p-value = $3.40 \cdot 10^{-2}$).

MM prognostic subtypes have distinct gene signatures

DEGAS   36

Differential gene expression analysis was performed between subtype 2 and all other subtypes

(**Supplementary File 5**), and we found that subtype 2 had significantly up-regulated *PHF19*

expression in all four of the patients (**Fig. 4H**). *PHF19* is a known marker for malignant disease in

MM [64]. Besides *PHF19*, its associated markers such as *HELLS*, *EZH2*, *TYMS*, *ZWINT*, and

MKI67 were also significantly up-regulated in subtype 2. These results suggested the possible

existence of a more malignant CD138+/*PHF19^{high}* subpopulation of plasma cells represented by

the subtype 2 cluster. It is important to notice that the gene feature set that was used as input into

DEGAS only contained the *HELLS* gene, which further highlights the ability of *DEGAS* to predict

high-risk cellular subtypes that can be further studied.

DEGAS is robust to hyper-parameter choice

To assess the robustness of *DEGAS*, we also analyzed how the hyper-parameter choices

influence its results using a set of 100 randomly generated hyper-parameters with 10-fold cross-

validation on each set of those 100 sets of hyper-parameters on the GBM datasets. The hyper-

parameters that we evaluated include: the number of training steps, batch size for single cells,

batch size for patients, number of hidden layer nodes, drop-out retention rate (the percentage of

nodes randomly retained at the hidden layer), patient loss weight, MMD loss weight, and L_2-

regularization weight. The detailed information about the range of hyper-parameters that were

randomly sampled can be found in subsection titled *Evaluation of DEGAS robustness to hyper-*

parameters in the **Methods** section while the default parameters used for all previous experiments

can be found in the subsection titled *Transfer learning using DEGAS*. We discovered that among

the eight hyper-parameters, the majority of them did not significantly affect the ROC-AUC for

predicting GBM subtypes in TCGA GBM patients with the exception of three hyperparameters –

namely the drop-out retention rate, number of hidden layer nodes, and L_2-regularization weight

with spearman correlation p-value < 0.1 (**Fig. S5, Table S22**). Similarly, the majority of hyper-

parameters did not significantly affect the correct assignment of subtype to GBM scRNA-seq

tumor, except for a few exceptions in training steps, patient loss weight, and MMD loss weight

with spearman correlation p-value < 0.1 (**Fig. S6**, **Table S22**). We therefore suggest users to keep

default settings for at least patient loss weight, MMD loss weight, and L_2-regularization weight.

The percentage of GBM subtype labels ranking in the top two predicted labels improves from 74%

to 82% if the default parameters or greater values are used for patient loss weight, MMD loss

weight, and L_2-regularization weight (**Fig. S7**).

Domain adaptation improves DEGAS disease association transfer

Without any bias, MMD and no MMD performances were not different from one another. After

bias was added, MMD did improve the ability of *DEGAS* to transfer disease associations onto

cells (**Fig. S8**). MMD is important for our algorithm because the bias added to the patients

represents the types of systematic bias that are present between bulk and single cell

transcriptomic data. In the example of simulation 2 with cell type 2 bias added, it is clear that all

of the patients tended to cluster adjacent to the cell type 2 cluster (**Fig. S8A**). We defined high-

risk cells in this example as cells with a disease association >0.2 on a [-1,1] scale. Once the

DEGAS model had been trained and the disease associations overlaid onto the cells, the

DEGAS model trained without MMD predicted many cells in cell types other than cell type 1 as

being high-risk (**Fig. S8B**). In contrast, the *DEGAS* model trained with MMD only identified cell

type 1 cells opposed to other cell types as high-risk (**Fig. S8C**). Over all 30 experiments, we

found that the disease association error was lower in the *DEGAS* models with MMD than in the

DEGAS models without MMD (t-test p-value < $2.2 \cdot 10^{-16}$, **Fig. S8D**). When the cells were

ordered by their error, there was no experiment where the *DEGAS* model without MMD

consistently outperformed the *DEGAS* model with MMD (Kolmogorov-Smirnov p-value < $2.2 \cdot 10^{-16}$,

Fig. S8E).

Regularization improves the robustness of DEGAS models

Regularization is an important part of the *DEGAS* model which prevents the data from being

overfit. Without regularization, *DEGAS* models perform worse during cross-validation (**Table.**

S24-25, **Fig. S9**). There is no case where an unregularized model performed better than a

regularized model in predicting patient labels during cross validation. Specifically, in simulation

3, the unregularized models performed 5% worse in PR-AUC when predicting patient labels in

patients (**Table. S24**). Similarly, the unregularized *DEGAS* models performed 9% worse in PR-

AUC when predicting cell type labels in cells (t-test p-value = $4.34 \cdot 10^{-3}$, **Table. S25**). The

regularization also improved the transfer of disease associations to the cells in 2/3 simulations

(**Fig. S9**).

Discussion

In this work, we developed the transfer learning framework *DEGAS* to integrate scRNA-seq and

patient-level transcriptomic data in order to infer the transferrable "impressions" between patient

characteristics in single cells and cellular characteristics in patients. Using transfer learning, we

trained a model with both scRNA-seq and patient bulk tissue gene expression data, then reduced

the differences between the distributions of the representations for the two data types in the final

hidden layer of our model via domain adaptation. This process allows information about patient

disease attributes as well as cell types to be transferred between the two data types. We focus

on the transfer of patient disease attributes to cells because there are far fewer available methods

addressing this task than deconvolution. We tested and validated the *DEGAS* framework on

datasets from one simulation and two diseases: GBM, which contained ground truth tumor

subtype labels, and AD, which contained ground truth cell type-disease associations.

These experiments on validation datasets demonstrate the necessity for *DEGAS* especially as it

relates to the current methods that rely on accurate clustering, cell type annotation, or case-

control scRNA-seq. For datasets that contain case and control scRNA-seq data, tools like *Augur*

are very effective to prioritize cell types. When no patient level transcriptomic data is available but

case-control scRNA-seq is available, tools like *Augur* should be used since *DEGAS* requires

patient level transcriptomic data. If patient level transcriptomic data and single cell transcriptomic

data are available and there is a necessity to overlay disease associations onto individual cells,

then only *DEGAS* can be used. Furthermore, if the scRNA-seq dataset does not contain case and

control samples then *DEGAS* needs to be used instead of *Augur* since *Augur* requires case and

control samples. The *DEGAS* framework in this sense can be used in a wide variety of study

designs as long as there is scRNA-seq and patient transcriptomic data.

Another challenging issue in scRNA-seq analysis is that it is difficult to determine the best

clustering options. In our simulation examples, we can determine that the correct number of

clusters based on average silhouette width would be four clusters. However, if the number of

clusters was increased in the clustering algorithm there would be a stronger correlation between

some clusters and disease. Therein lies the challenge – should the clustering results be optimized

to reflect the relative transcriptomic signals or should they be optimized to create the greatest

correlations with disease state? Furthermore, the different resolutions of clusters may capture

different correlations with disease. For these reasons, assigning disease associations directly to

cells alleviates some of these problems with cluster resolution decisions. Assigning disease

associations directly to cells not only solves the cluster resolution problem but also allows

simultaneous identification of cell-intrinsic and cell proportional changes.

The *DEGAS* algorithm can identify both cell-intrinsic changes and cell proportional changes as

demonstrated in the simulation examples and the AD study. In simulation 1, the disease is

associated with proportional changes in cell type 1. In simulation 2, the disease is associated

with a cell-intrinsic change of cell type 4. In simulation 3, there are both cell-intrinsic changes

and cell proportional changes in cell type 4. In the AD experiments, two of the single-cell

datasets did include data from both AD and normal brains [15, 32]. The cells that came from the

AD patients tended to have a higher association with AD, which indicates the detection of cell-

intrinsic changes. The importance of cell prioritization at the individual cell level is highlighted in

simulation and AD examples. Simulation 2 shows an example where cell level associations are

necessary due to clustering results that do not capture the disease associations. Specifically,

there are cases where cells will cluster together but have dissimilar associations to disease. If

the cells of cell type 4 are not evaluated individually, the association of the cell type 4 disease

subtype with disease could be lost. In the AD example, the astrocyte cell type is overall not

associated with AD. However, a subset of astrocytes expressing markers for DAAs were found

to have a positive disease association while still clustering with the astrocytes that were not

associated with disease. Similarly, microglia cells are broadly positively associated with AD but

the highest AD association microglia were enriched for DAM markers. When a targeted analysis

was performed on only microglia from AD and normal brains, highest AD association microglia

were enriched for both HAM and DAM markers. These examples show how DEGAS can identify

disease associated cells that cluster within a larger cell type.

In short, the *DEGAS* analysis on AD data further validated our model by correctly identifying the

decreased neuron and increased microglia proportions in AD patients. Aside from these known

characteristics of AD pathology, we also identified a *GFAP+* astrocyte subtype taken from normal

human brain tissue that is associated with AD and is supported from AD mouse models [55]. We

further validated this by finding that *GFAP* expression in Astrocytes was significantly increased in

Astrocytes taken from AD patients and concluded that there may be an expansion of this Astrocyte

subtype in AD. This is also a convincing example of the utility of *DEGAS* as it assigned disease

association at the single cell level, allowing us to identify intra-cell type differences in disease risk

that constitute disease-associated cells.

1009

For the GBM single cell patient cohort, each GBM tumor, from which scRNA-seq data was

generated, had a GBM subtype label [44]. The *DEGAS* results showed that the majority of cells

in each tumor were labeled with the same GBM subtype as previously defined in Patel *et al.* [44].

Specifically, *DEGAS* correctly mapped Proneural, Mesenchymal, Classical, and Neural GBM

subtypes to single cells in four GBM tumor samples. This experiment also shows the broad

applicability of the model since the single cells had no labels and the patient samples had

multiclass labels. *DEGAS* is highly flexible and allows for different categories of output labels to

be combined, which may include but are not limited to classification labels, Cox proportional

hazard, and even no labels. This allows for a wide variety of applications to adopt the *DEGAS*

framework so that impressions are not limited to only one type of disease attribute.

1020

To explore disease with less understood cellular subtypes, we applied *DEGAS* to multiple MM

datasets. The models were able to assign PFS metrics to individual cells and subtype populations

of CD138+ cells identified by cell type clustering methods Seurat [28] and *BERMUDA* [25]. Among

the identified subtypes of cells, subtype 2 was the most consistent between patients visualized

by *BERMUDA* (**Fig. S4E**). Furthermore, we found that the subtype 2 cell population appeared to

have a gradient of cells moving away from the main subtype 1 group, possibly associated with a

certain degree of differentiation (**Fig. S4A-D**). We did experience a lower PR-AUC for subtype 4

than the other subtypes used during model training. However, this subtype was extremely

uncommon in the samples and as a result the random PR-AUC would be close to zero making

the PR-AUC of 0.44 well above random. Considering that subtype 4 was not found to be highly

associated with progression, the lower PR-AUC did not greatly affect our interpretation of the

data, which mainly focused on subtype 1 and subtype 2. We believe that *DEGAS* could be

improved for highly imbalanced data.

1034

Upon further examination, we found evidence that the subtype 2 cells may represent a population

of malignant plasma cells expressing high levels of *PHF19*. *PHF19* is known to play a role in

hematopoietic stem cell state and differentiation [65-67] and is a marker for aggressive disease

in MM [64]. Furthermore, knock down of *PHF19* has been shown to shift myeloma cells into a less

proliferative state [64]. The subtype 2 cells express *SDC1* (also known as *CD138)* and showed

significantly increased *PHF19* expression in comparison to the other subtypes. Since all of the

IUSM MM cells in our study had already been FACS sorted for CD138+, it is possible we have

identified a subpopulation of CD138+/*PHF19^{high}* cells in MM tumors. This could prove a useful

finding since currently the association between *PHF19* and tumor aggressiveness is at the patient

level whereas our results imply that only a fraction of malignant plasma cells in a MM tumor

actually overexpress *PHF19.*

1046

This subtype could be targeted using precision immunotherapies that are not restricted to a single

patient since the CD138+/*PHF19^{high}* cells (*i.e.,* subtype 2) were found to be present in multiple

(3/4) patients. Of the three patients with detectable levels of subtype 2 in the CD138+ fraction,

two patients (patient 2 and patient 4) had relapsed MM at time of biopsy and the other patient

(patient 5) was SMM at biopsy and later progressed to MM. The other patient (patient 3) had little

to no detectable subtype 2 cells in the CD138+ fraction and was SMM at time of biopsy and has

not progressed to MM. These signs again seem to indicate a common cellular phenotype

associated with progression in MM.

1055

1056   Based on the validated results in a variety of disease data analyses, we find that *DEGAS* has

1057   broad applications in virtually all diseases with available patient-level and single cell level omic

1058   data. The tensorflow [68] machine learning code is integrated with a simple R package interface

1059   (https://github.com/tsteelejohnson91/DEGAS) which will facilitate researchers to manipulate

1060   scRNA-seq and bulk expression data on their own.

1061

Conclusion

DEGAS is a powerful transfer learning tool for integrating different levels of omic data and

identifying the latent molecular relationships between populations of cells and disease

attributes, which we refer to as impressions. We validated the *DEGAS* framework on simulated

data, GBM and AD by showing *DEGAS* models were capable of accurately predicting patient

characteristics at single-cell level. We then leveraged this transfer learning approach on MM

data and identified a CD138+/*PHF19^{high}* subtype population in MM that was significantly

associated with disease progression. This subtype contains unique RNA profiles and gene

correlations that could be both leveraged as a prognostic biomarker and possibly targeted

directly to reduce the risk of progression. We believe that *DEGAS* can be a powerful solution to

overcome the challenge of integrating patient single-cell data with bulk tissue data so that

researchers can identify populations of cells associated with an disease attribute of interest.

Furthermore, *DEGAS* can accommodate flexible data types. This makes it a highly general

framework that can be applied in multiple diseases and data types to identify cellular

populations that are associated with prognosis or treatment response, or to identify specific

patient groups with certain cell subtypes for personalized treatment.

1078

Acknowledgements

We thank the Center for Computational Biology and Bioinformatics for the computational

resources and work space to complete the research. We also thank the MMRF for the data

generated as part of the Multiple Myeloma Research Foundation Personalized Medicine

Initiatives (https://research.themmrf.org and www.themmrf.org), the Allen Institute for Brain

Science for the data generated as part of their cell types database, and Mount Sinai/JJ Peters

VA Medical Center for the data generated as a part of their brain bank.

1086

Funding

National Institutes of Health NLM-NRSA Fellowship F31LM013056 to TSJ and The Ohio State

University (Columbus, OH) and departmental start-up funding from the Indiana University

School of Medicine (Indianapolis, IN) to KH and TSJ.

Author contributions

TSJ, CYY, JZ, and KH conceived and designed the project. TSJ, CYY, SX performed the

analyses. TSJ and ZH designed the software package. TSJ, CYY, XH, SX, CD, MA, YW, CB,

YZ, YL, JZ, BW, and KH interpreted the results. ZH, TW, WS, YW, and CB provided technical

guidance. TSJ, CYY, JZ, and KH wrote the manuscript. JZ and KH supervised the project.

Competing interests

The authors declare that this research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Data and materials availability

The *DEGAS* R package is freely available on GitHub

(https://github.com/tsteelejohnson91/DEGAS). A minimum reproducible example of the IUSM

myeloma scRNA-seq data is also deposited on Github. Our complete myeloma scRNA-seq has

been deposited on GEO (GSE161722). All other data are publicly available.

Figure legends

Fig. 1 A workflow diagram of the *DEGAS* framework. **A**) The workflow for a typical experiment

with *DEGAS*. Note that *DEGAS* is not meant to replace the abundant packages available to load,

preprocess, select features, cluster, and visualize scRNA-seq data. It is rather meant to augment

these packages to assign disease associations to cells. **B**) The scRNA-seq and patient

DEGAS   45

expression data are preprocessed into expression matrices. Next, Bootstrap aggregated

DenseNet *DEGAS* models are trained using both single cell and patient disease attributes using

a multitask learning neural network that learns latent representation reducing the differences

between patients and single cells at the final hidden layer using maximum mean discrepancy

(MMD). **C**) The output layer of this model can be used to simultaneously infer disease attribute

impressions in single cells and cellular composition impressions in patients.

1119

Fig. 2 Simulation study and baseline comparisons of *DEGAS* framework. A) 5,000 simulated

cells from *Splatter* with 4 cell types where one of the cell types has two subtypes. Cell type 4 is

composed of two subtypes that are specific to either disease or normal patients. 2,000 of these

cells were used to generate the 600 simulated patients in **B-D** and 3,000 were used as the cell

input to our *DEGAS* models. **E)** Optimal cluster number (4 clusters) based on average silhouette

width for the 3,000 cells not used to generate patients. **F)** The same 3,000 cells used as the

cellular input colored by their cluster. **G)** *DEGAS* comparison to *Augur* in simulation 1. **H)** *DEGAS*

comparison with *Augur* in simulation 2. **I)** *DEGAS* comparison with *Augur* in simulation 3. **J-L)**

DEGAS-calculated disease association from each simulation overlaid onto 3,000 cells. The violin

plot in the bottom left corner is deconvolution cell type proportion for cell type 1 in simulation 1

patients (**J**), cell type 4 proportion in simulation 2 patients (**K**), and cell type 4 proportion in

simulation 3 patients (**L**).

1132

Fig. 3 *DEGAS* validation in GBM and AD. *DEGAS* output of the distribution of GBM subtypes

in single cells from five GBM tumors. Four of the five tumors had known GBM subtype information

from Patel *et al*. (MGH26: Proneural, MGH28: Mesenchymal, MGH29: Mesenchymal, and

MGH30: Classical, indicated by red boxes) which were recapitulated by *DEGAS*. The subtype

information for the tumors, MGH26, MGH28, MGH29, and MGH30 were derived from Patel *et al.*

where MGH31 did not have a clearly defined GBM subtype. The association of cells assigned to

each subtype were plotted for each tumor; **A)** MGH26, **B)** MGH28, **C)** MGH29, **D)** MGH30 and **E)**

MGH31. Median values are marked by a diamond in each of the violin plots. **F)** The death

association centered around 0 is overlaid on all of the single cells from the five tumors (indicated

by color). **G)** *DEGAS* output of AD association for each single cell. The AD association score is

indicated by the color and is overlaid onto AIBS single cells. This plot shows the negative AD

association in neuron cells and positive AD association in Microglia. **H-I)** There also appeared to

be a subpopulation of astrocytes with positive AD association. The astrocytes were plotted

separately and colored by AIBS Astrocyte subtypes **(H)** and *GFAP* expression, a disease-

associated astrocyte marker **(I)**. **J)** Comparison of *DEGAS*-derived AD associations for single

cells from AD and Normal control samples from Grubman *et al.* **K-M)** Targeted analysis of

microglia from Grubman *et al.* including the AD associations overlaid onto microglia **(K)**, AD

association comparing AD status of patient sample from which the cells were sampled **(L)**, and

PCC between AD association with HAM marker genes comparing up- and down-regulated HAM

marker genes **(M)**. Significance values: n.s. (not significant), • (0.1), * (0.05), ** (0.01), *** (0.001).

1153

Fig. 4 Association between subtypes and progression risk in MM. IUSM CD138+ scRNA-

seq subtype clusters generated from *Seurat* colored by **A)** cluster, *i.e.,* subtype and **B)**

progression association. **C)** Kaplan-Meier curves of PFS from cross-validation for the MMRF

patients stratified by median proportional hazard. **D)** Kaplan-Meier curves of OS from Zhan *et al.*

external dataset stratified by median proportional hazard. **E)** Progression association for IUSM

CD138+ subtypes **F)** Progression association for NHIP, MGUS, SMM, and MM in the external

dataset Ledergor *et al*. **G)** Subtype 2 enrichment for NHIP, MGUS, SMM, and MM in the

external dataset Ledergor *et al*. NHIP: normal hip bone marrow, MGUS: monoclonal

gammopathy of undetermined significance, SMM: smoldering multiple myeloma, MM: multiple

myeloma. Significance values: • (0.1), * (0.05), ** (0.01), *** (0.001). All plots were generated

using the default parameters for the *DEGAS* package described in the section of **Methods**:

Transfer learning using DEGAS.

1166

References

1. Lahnemann, D., et al., *Eleven grand challenges in single-cell data science.* Genome Biol,
2020. **21**(1): p. 31.
2. Ma, A., et al., *Integrative Methods and Practical Challenges for Single-Cell Multi-omics.*
Trends Biotechnol, 2020. **38**(9): p. 1007-1022.
3. Kiselev, V.Y., A. Yiu, and M. Hemberg, *scmap: projection of single-cell RNA-seq data*
across data sets. Nat Methods, 2018. **15**(5): p. 359-362.
4. Cao, Y., et al., *scRNASeqDB: A Database for RNA-Seq Based Gene Expression Profiles in*
Human Single Cells. Genes (Basel), 2017. **8**(12).
5. Abugessaisa, I., et al., *SCPortalen: human and mouse single-cell centric database.* Nucleic
Acids Res, 2018. **46**(D1): p. D781-D787.
6. Stuart, T., et al., *Comprehensive integration of single-cell data.* Cell, 2019. **177**(7): p.
1888-1902. e21.
7. Gawel, D.R., et al., *A validated single-cell-based strategy to identify diagnostic and*
therapeutic targets in complex diseases. Genome Med, 2019. **11**(1): p. 47.
8. Chen, S., et al., *Single-cell analysis reveals transcriptomic remodellings in distinct cell*
types that contribute to human prostate cancer progression. Nature Cell Biology, 2021.
23(1): p. 87-98.
9. Jang, J.S., et al., *Molecular signatures of multiple myeloma progression through single*
cell RNA-Seq. Blood cancer journal, 2019. **9**(1): p. 1-10.
10. Maynard, A., et al., *Therapy-induced evolution of human lung cancer revealed by single-*
cell RNA sequencing. Cell, 2020. **182**(5): p. 1232-1251. e22.
11. Cobos, F.A., et al., *Benchmarking of cell type deconvolution pipelines for transcriptomics*
data. Nature communications, 2020. **11**(1): p. 1-14.
12. Johnson, T.S., et al., *Combinatorial analyses reveal cellular composition changes have*
different impacts on transcriptomic changes of cell type specific genes in Alzheimer's
Disease. Sci Rep, 2021. **11**(1): p. 353.
13. Johnson, T.S., et al., *Spatial cell type composition in normal and Alzheimers human*
brains is revealed using integrated mouse and human single cell RNA sequencing. Sci
Rep, 2020. **10**(1): p. 18014.
14. Jung, S.-H. and S.-C. Chow, *On sample size calculation for comparing survival curves*
under general hypothesis testing. Journal of biopharmaceutical statistics, 2012. **22**(3): p.
485-495.
15. Mathys, H., et al., *Single-cell transcriptomic analysis of Alzheimer's disease.* Nature,
2019. **570**(7761): p. 332-337.
16. Rossi, M.A., et al., *Obesity remodels activity and transcriptional state of a lateral*
hypothalamic brake on feeding. Science, 2019. **364**(6447): p. 1271-1274.

17. Crowell, H.L., et al., *muscat detects subpopulation-specific state transitions from multi-*
sample multi-condition single-cell transcriptomics data. Nat Commun, 2020. **11**(1): p.
6077.
18. Burkhardt, D.B., et al., *Quantifying the effect of experimental perturbations at single-cell*
resolution. Nat Biotechnol, 2021. **39**(5): p. 619-629.
19. Skinnider, M.A., et al., *Cell type prioritization in single-cell data.* bioRxiv, 2019: p.
2019.12.20.884916.
20. Kouw, W.M. and M. Loog, *A Review of Domain Adaptation without Target Labels.* IEEE
Transactions on Pattern Analysis and Machine Intelligence, 2021. **43**(3): p. 766-785.
21. Hardoon, D.R., S. Szedmak, and J. Shawe-Taylor, *Canonical correlation analysis: An*
overview with application to learning methods. Neural computation, 2004. **16**(12): p.
2639-2664.
22. Gretton, A., et al., *A kernel two-sample test.* Journal of Machine Learning Research,
2012. **13**(Mar): p. 723-773.
23. Andrew, G., et al. *Deep canonical correlation analysis.* in *International conference on*
machine learning. 2013. PMLR.
24. Zhang, F., Y. Wu, and W. Tian, *A novel approach to remove the batch effect of single-cell*
data. Cell discovery, 2019. **5**(1): p. 1-4.
25. Wang, T., et al., *BERMUDA: a novel deep transfer learning method for single-cell RNA*
sequencing batch correction reveals hidden high-resolution cellular subtypes. Genome
Biol, 2019. **20**(1): p. 165.
26. Tran, H.T.N., et al., *A benchmark of batch-effect correction methods for single-cell RNA*
sequencing data. Genome biology, 2020. **21**(1): p. 1-32.
27. Johnson, T.S., et al., *LAmbDA: label ambiguous domain adaptation dataset integration*
reduces batch effects and improves subtype detection. Bioinformatics, 2019. **35**(22): p.
4696-4706.
28. Butler, A., et al., *Integrating single-cell transcriptomic data across different conditions,*
technologies, and species. Nat Biotechnol, 2018. **36**(5): p. 411-420.
29. Aran, D., et al., *Reference-based analysis of lung single-cell sequencing reveals a*
transitional profibrotic macrophage. Nature immunology, 2019. **20**(2): p. 163-172.
30. Araujo, T., et al., *Classification of breast cancer histology images using Convolutional*
Neural Networks. PLoS One, 2017. **12**(6): p. e0177544.
31. Bardou, D., K. Zhang, and S.M. Ahmad, *Classification of Breast Cancer Based on*
Histology Images Using Convolutional Neural Networks. IEEE Access, 2018. **6**: p. 24680-
24693.
32. Grubman, A., et al., *A single-cell atlas of entorhinal cortex from individuals with*
Alzheimer's disease reveals cell-type-specific gene expression regulation. Nature
neuroscience, 2019. **22**(12): p. 2087-2097.
33. Holtman, I.R., et al., *Induction of a common microglia gene expression signature by*
aging and neurodegenerative conditions: a co-expression meta-analysis. Acta
Neuropathol Commun, 2015. **3**: p. 31.
34. Hemonnot, A.L., et al., *Microglia in Alzheimer Disease: Well-Known Targets and New*
Opportunities. Front Aging Neurosci, 2019. **11**: p. 233.

35.   Glass, C.K., et al., *Mechanisms underlying inflammation in neurodegeneration.* Cell,
1248   2010. **140**(6): p. 918-34.
1249   36.   Donev, R., et al., *Neuronal death in Alzheimer's disease and therapeutic opportunities.* J
1250   Cell Mol Med, 2009. **13**(11-12): p. 4329-48.
1251   37.   DeKosky, S.T. and S.W. Scheff, *Synapse loss in frontal cortex biopsies in Alzheimer's*
1252   *disease: correlation with cognitive severity.* Ann Neurol, 1990. **27**(5): p. 457-64.
1253   38.   de Wilde, M.C., et al., *Meta-analysis of synaptic pathology in Alzheimer's disease reveals*
1254   *selective molecular vesicular machinery vulnerability.* Alzheimers Dement, 2016. **12**(6):
1255   p. 633-44.
1256   39.   Akiyama, H., *Inflammatory response in Alzheimer's disease.* Tohoku J Exp Med, 1994.
1257   **174**(3): p. 295-303.
1258   40.   Institute, N.C., *Cancer Statistics*, N.C. Institute, Editor. 2019: Cancer.gov.
1259   41.   Zhan, F., et al., *The molecular classification of multiple myeloma.* Blood, 2006. **108**(6): p.
1260   2020-8.
1261   42.   Ledergor, G., et al., *Single cell dissection of plasma cell heterogeneity in symptomatic*
1262   *and asymptomatic myeloma.* Nat Med, 2018. **24**(12): p. 1867-1876.
1263   43.   Cohen, Y.C., et al., *Identification of resistance pathways and therapeutic targets in*
1264   *relapsed multiple myeloma patients through single-cell sequencing.* Nature medicine,
1265   2021: p. 1-13.
1266   44.   Patel, A.P., et al., *Single-cell RNA-seq highlights intratumoral heterogeneity in primary*
1267   *glioblastoma.* Science, 2014. **344**(6190): p. 1396-401.
1268   45.   Cancer Genome Atlas Research, N., *Comprehensive genomic characterization defines*
1269   *human glioblastoma genes and core pathways.* Nature, 2008. **455**(7216): p. 1061-8.
1270   46.   Wang, M., et al., *The Mount Sinai cohort of large-scale genomic, transcriptomic and*
1271   *proteomic data in Alzheimer's disease.* Sci Data, 2018. **5**: p. 180185.
1272   47.   Ching, T., X. Zhu, and L.X. Garmire, *Cox-nnet: An artificial neural network method for*
1273   *prognosis prediction of high-throughput omics data.* PLoS Comput Biol, 2018. **14**(4): p.
1274   e1006076.
1275   48.   Couturier, C.P., et al., *Single-cell RNA-seq reveals that glioblastoma recapitulates a*
1276   *normal neurodevelopmental hierarchy.* Nat Commun, 2020. **11**(1): p. 3406.
1277   49.   Guo, M., et al., *SINCERA: A Pipeline for Single-Cell RNA-Seq Profiling Analysis.* PLoS
1278   Comput Biol, 2015. **11**(11): p. e1004575.
1279   50.   Iacono, G., R. Massoni-Badosa, and H. Heyn, *Single-cell transcriptomics unveils gene*
1280   *regulatory network plasticity.* Genome Biol, 2019. **20**(1): p. 110.
1281   51.   Grus, J., *Data science from scratch: first principles with python.* 2019: O'Reilly Media.
1282   52.   Ioffe, S. and C. Szegedy. *Batch normalization: Accelerating deep network training by*
1283   *reducing internal covariate shift*. in *International conference on machine learning*. 2015.
1284   PMLR.
1285   53.   Juszczak, P., D. Tax, and R.P. Duin. *Feature scaling in support vector data description*. in
1286   *Proc. asci*. 2002. Citeseer.
1287   54.   Verhaak, R.G., et al., *Integrated genomic analysis identifies clinically relevant subtypes of*
1288   *glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1.* Cancer
1289   Cell, 2010. **17**(1): p. 98-110.

1290   55.   Habib, N., et al., *Disease-associated astrocytes in Alzheimer's disease and aging.* Nature
1291   Neuroscience, 2020. **23**(6): p. 701-706.
1292   56.   Srinivasan, K., et al., *Alzheimer's Patient Microglia Exhibit Enhanced Aging and Unique*
1293   *Transcriptional Activation.* Cell Rep, 2020. **31**(13): p. 107843.
1294   57.   Keren-Shaul, H., et al., *A Unique Microglia Type Associated with Restricting Development*
1295   *of Alzheimer's Disease.* Cell, 2017. **169**(7): p. 1276-1290 e17.
1296   58.   Maaten, L.v.d. and G. Hinton, *Visualizing data using t-SNE.* Journal of machine learning
1297   research, 2008. **9**(Nov): p. 2579-2605.
1298   59.   Becht, E., et al., *Dimensionality reduction for visualizing single-cell data using UMAP.* Nat
1299   Biotechnol, 2018.
1300   60.   Friedman, J., T. Hastie, and R. Tibshirani, *Regularization Paths for Generalized Linear*
1301   *Models via Coordinate Descent.* J Stat Softw, 2010. **33**(1): p. 1-22.
1302   61.   Zappia, L., B. Phipson, and A. Oshlack, *Splatter: simulation of single-cell RNA sequencing*
1303   *data.* Genome Biol, 2017. **18**(1): p. 174.
1304   62.   Fu, H., et al., *Tau Pathology Induces Excitatory Neuron Loss, Grid Cell Dysfunction, and*
1305   *Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease.* Neuron, 2017. **93**(3):
1306   p. 533-541 e5.
1307   63.   Xu, J., et al., *Multimodal single-cell/nucleus RNA sequencing data analysis uncovers*
1308   *molecular networks between disease-associated microglia and astrocytes with*
1309   *implications for drug repurposing in Alzheimer's disease.* Genome research, 2021: p. gr.
1310   272484.120.
1311   64.   Mason, M.J., et al., *Multiple Myeloma DREAM Challenge reveals epigenetic regulator*
1312   *PHF19 as marker of aggressive disease.* Leukemia, 2020. **34**(7): p. 1866-1874.
1313   65.   Bagger, F.O., S. Kinalis, and N. Rapin, *BloodSpot: a database of healthy and malignant*
1314   *haematopoiesis updated with purified and single cell mRNA sequencing profiles.* Nucleic
1315   acids research, 2019. **47**(D1): p. D881-D885.
1316   66.   Lara-Astiaso, D., et al., *Chromatin state dynamics during blood formation.* science, 2014.
1317   **345**(6199): p. 943-949.
1318   67.   Vizán, P., et al., *The Polycomb-associated factor PHF19 controls hematopoietic stem cell*
1319   *state and differentiation.* Science advances, 2020. **6**(32): p. eabb2745.
1320   68.   Abadi, M., et al. *Tensorflow: A system for large-scale machine learning.* in *12th {USENIX}*
1321   *Symposium on Operating Systems Design and Implementation ({OSDI} 16)*. 2016.
1322