1 **Impact of transposable elements on methylation and gene expression across**
2 **natural accessions of *Brachypodium distachyon.***
3

4 Michele Wyler[1], Christoph Stritt[1], Jean-Claude Walser[2], Célia Baroux[1] and Anne C.
5 Roulin[1*]
6
[1]Institute for Plant and Microbial Biology, University of Zurich, Zollikerstrasse 107,
Zurich 8008, Switzerland

[2] Genetic Diversity Centre, ETH Zürich, Universitätstrasse 16, Zurich, Switzerland

*Corresponding author: Anne C. Roulin; Institute for Plant and Microbial Biology,
University of Zurich, Zollikerstrasse 107, Zurich 8008, Switzerland; +41 (0)44 63
48398; anne.roulin@botinst.uzh.ch

Abstract
15

Transposable elements (TEs) constitute a large fraction of plant genomes and are

mostly present in a transcriptionally silent state through repressive epigenetic

modifications such as DNA methylation. TE silencing is believed to influence the

regulation of adjacent genes, possibly as DNA methylation spreads away from the

TE. Whether this is a general principle or a context-dependent phenomenon is still

under debate, pressing for studying the relationship between TEs, DNA methylation

and nearby gene expression in additional plant species. Here we used the grass

Brachypodium distachyon as a model and produced DNA methylation and

transcriptome profiles for eleven natural accessions. In contrast to what is observed in

Arabidopsis thaliana, we found that TEs have a limited impact on methylation

spreading and that only few TE families are associated to a low expression of their

adjacent genes. Interestingly, we found that a subset of TE insertion polymorphisms is

associated with differential gene expression across accessions. Thus, although not

having a global impact on gene expression, distinct TE insertions may contribute to

specific gene expression patterns in *B. distachyon*

Key words: Transposable elements, DNA methylation, gene expression,
Brachypodium distachyon

Significance statement: Transposable elements (TEs) are a major component of

plant genomes and a source of genetic and epigenetic innovations underlying

adaptation to changing environmental conditions. Yet molecular evidence linking TE

silencing and nearby gene expression are lacking for many plant species. We show

that in the model grass Brachypodium DNA methylation spreads over very short
distances around TEs, with an influence on gene expression for a small subset of TE
families.
40
41

Introduction

Since their discovery by Barbara McClintock in the late 1940's (McClintock 1947), transposable elements (TEs) have been shown to populate a large fraction of eukaryotic genomes and to be an important source of genetic and phenotypic diversity. This is explained by the observation that TEs have the ability to modulate nearby gene expression and can remodel genomes when activated by stress (Grandbastien 1998; Grandbastien 2005; Thieme et al. 2017), thereby providing raw diversity necessary for adaptation to changing environments (Bonchev and Parisod 2013; Rey et al. 2016). Remarkable examples of single-trait modulation by a TE insertion providing a selective advantage have been reported in both plants and animals (Guio et al. 2014; Van't Hof et al. 2016; Horváth et al. 2017).

While TEs can provide selective advantages, their mutagenic properties may at the same time represent a threat for the integrity of the host genome (McClintock 1950; Shang et al. 2017; Thind et al. 2018). In response, genomes have evolved several layers of defence to contain TE activity. In plants, transcriptional repression of TEs is mediated by the RNA-directed DNA methylation (RdDM) pathway (for review Sigman and Slotkin 2016; Zhang et al. 2018). Initiated by the plant-specific RNA polymerase IV (*Pol IV*), TE transcription leads to the formation of 24 nt siRNA targeting DNA methylation at the TE source locus. Acting as a double-edged sword, DNA (and histone) methylation may spread to neighbouring regions and possibly contribute the alteration of adjacent gene expression (Hollister et al. 2010; Wang et al. 2013).

Recent studies reported on the impact of TEs on DNA methylation and expression of nearby genes in different plant species (Eichten et al. 2012; Makarevitch et al. 2015; Eichten et al. 2016; Choi and Purugganan 2017; Wang et al. 2018). Yet, most of our knowledge on the regulatory mechanisms largely rely on the eudicot model *Arabidopsis thaliana* (for review Sigman and Slotkin 2016; Zhang et al. 2018). In *A. thaliana*, the presence of TEs tends to be associated with a reduced expression of adjacent genes (Wang et al. 2013), but whether this also occurs in other species remains to be demonstrated (Choi and Purugganan 2017). In this context, we contributed an analysis of DNA methylation and gene expression in the model grass *Brachypodium distachyon*.

Established as an important model for functional genomics, the small genome

of *B. distachyon* (272 Mb; 30% of TEs) is fully sequenced and assembled into five

chromosomes (The International Brachypodium Consortium 2010). Comparative

genomics and cytogenetic analyses (Hasterok et al. 2019) showed that it underwent

multiple and independent nested chromosome fusions which resulted in a higher gene

density at the former distal regions of the inserted chromosomes. Furthermore, TEs

are much more pervasive than in *A. thaliana,* and although their dynamics is well

characterized in *B. distachyon* (Stritt et al. 2018; Stritt et al. 2019), little is know

about their functional impact. We therefore asked to what extent TEs influence DNA

methylation and gene expression in *B. distachyon,* and how these patterns differ from

what is observed in *A. thaliana.*

Results and Discussion

TEs largely shape genome-wide DNA methylation patterns in Brachypodium

We produced bisulfite- and RNA-sequencing data for eleven *B. distachyon* accessions

that belong to three genetic clusters: Western (ABR2, Koz3, Bd30-1 and Sig2; Spain),

Eastern (the reference accession Bd21, Bd21-3, Bd3-1, BdTR12C and Koz3; Iraq and

Turkey) and accessions from a third cluster (BdTR7a, Bd1-1; Turkey) showing a

delay in flowering time (Gordon et al. 2017; Stritt et al. 2018). We also used

publically available bisulfite- and RNA-sequencing data for the reference accession

Col-0 of *A. thaliana* (Table S1).

We first estimated methylation levels (expressed as the percentage of

methylated cytosines), gene and TE density in 100 kb windows for the reference

genomes of *A. thaliana* (Col-0) and *B. distachyon* (Bd21). The circular visualisation

of these three genomic features reveals two distinct architectures: genes and TEs are

clearly compartmentalised in *A. thaliana* (Figure 1A) while largely interspersed in

B. distachyon (Figure 1B). Yet we found that DNA methylation is positively

correlated with TE density and negatively correlated with gene density in both species

(Figure S1). We then estimated the methylation level of genes and TEs, aggregating

the latter at the superfamily level. TE body methylation is significantly higher than

gene body methylation in the two species, although variability is observed among TE

superfamilies (Figure 1C and D, Table S2-S4 for P-values). Genome-wide DNA

methylation patterns are thus largely shaped by TEs in *B. distachyon,* like in other

flowering plants species (Zhang et al. 2006; Regulski et al. 2013; Takuno and Gaut
2013; Seymour et al. 2014). We also found that most TE superfamilies and genes
show a higher level of body methylation in *B. distachyon* than in *A. thaliana* (Figure
1C and D, Table S5 for P-values), confirming a low genome-wide level of body
methylation in *A. thaliana* compared to other species (Seymour et al. 2014, Eichten et
al. 2016).

We finally assessed how far methylation spreads around TEs. To do so, we
estimated methylation levels in 100 bp windows spanning 2.5 kb around each TE
insertion and averaged the values per TE superfamily. We also categorized TE
insertions depending on their location on the chromosome (arm vs. centromere) and
visualised the results by producing heatmaps. We observe that DNA methylation
around TEs spreads over longer distances in gene-poor centromeric regions than in
distal regions of the chromosome arm in *A. thaliana* (Figure 2). In *B. distachyon*,
spreading mostly reaches a few 100 bp, except around centromeric *Gypsy* elements
where methylation tends to spread beyond 2 kb (Figure 2). Thus, in contrast to our
expectations, high levels of TE body methylation in *B. distachyon* are generally not
accompanied with a long spreading of methylation, neither in the centromeres nor on
the chromosome arms. This is surprising given that extended methylation spreading
around TE is associated to gene-poor pericentromeric regions in rice (Choi and
Purugganan, 2017), a species closely related to *B. distachyon*.

Impact of TEs on nearby gene expression

In *A. thaliana*, genes flanked by a TE within 2 kb show reduced expression compared
to genes devoid of TEs in their vicinity (Wang et al. 2013). It was proposed that this
may result from a combined effect of DNA methylation spreading and DNA structure
variation (Wang et al. 2013). For both the reference accessions Col-0 and Bd21 we
estimated the average level of expression for genes having a TE (regardless of the
type of TE) in their sequence or within windows ranging from near vicinity (0-200
bp) to up to 2 kb (Figure S2). While we reproduced the pattern described above with
publically available data in *A. thaliana*, our analysis shows that TEs do not have such
an effect on nearby gene expression in *B. distachyon*. To rule out potential technical
artefacts, we extended the analysis to the ten additional *B. distachyon* accessions we

sampled but did not observe any overall effect of TEs on nearby gene expression
either (Figure S2).

The effect of TEs on methylation and gene expression is known to be family-
dependent (Eichten et al. 2012; Choi and Purugganan 2017), which might explain
why no effect is observed when all TEs are pooled. We therefore repeated this
analysis separately for the ten most active TE families (Stritt et al. 2018) in all eleven
B. distachyon accessions to explore potential genetic background effects. We also
produced heatmaps as described above to assess a potential link between methylation
spreading around these ten TE families and nearby gene expression. Methylation
spreading was in a second step used to statistically cluster the ten families.
Eventually, as young TEs might be more efficiently targeted by silencing
mechanisms, we extracted the age of each copy from Stritt et al. (2019). For four of
the ten families, this analysis allowed to uncover clear effects on gene expression
following two distinct patterns (Figure 3A): a short-distance effect (< 400 bp; all P-
values <0.001) conveyed by two *Gypsy* families (RLG_Bdis0039 and RLG_Bdis004)
and one *Copia* (RLC_Bdis022) family; and a long-distance effect (1 kb; P-values
<0.001) for the *Gypsy* element RLG_Bdis180. However, methylation spreading is
unlikely to be involved as our hierarchical clustering shows that spreading varies from
long (RLG_Bdis004, RLG_Bdis180) to intermediate (RLG_Bdis0039) and short
(RLG_Bdis022) around these four families (Figure 3B). In the same manner, there is
no clear association neither with the age of the family (Figure 3B) nor with the
genetic background as all accessions show comparable gene expression patterns
(Figure 3A). It is nonetheless worth noting that RLG_Bdis004 and RLC_Bdis022 are
two of the most recently active families in the genome of *B. distachyon* (Stritt et al.
2019), suggesting a possible scenario where *de novo* silencing of these two elements
may influence nearby-gene regulation.

We eventually investigated the impact of TE insertion polymorphisms (TIPs)
on gene expression. Overall, we identified 1,833 high-confidence TIPs, i.e. insertions
present in at least one of the ten natural accessions but absent from the reference
genome Bd21 (Table S6). Due to the relative small number of TIPs located nearby
genes, we lacked power to investigate their impact on gene expression at the family
level. To circumvent this difficulty, we measured the log-fold change in gene
expression for loci harbouring a TIP in their vicinity (<1 kb) compared to the same

loci without a TIP in a different accession. We found that the majority of the TIPs

have no effect on adjacent gene expression (Figure S3), consistent with previous

findings that TE polymorphisms have little impact on methylation spreading in

B. distachyon (Eichten et al. 2016). These results contrast again with what is found in

A. thaliana, where polymorphic TEs are associated with differential expression of

nearby-genes among natural accessions (Stuart et al. 2016). A likely explanation for

this difference may be the different types of TIPs in the two species, notably the large

number of non-functional solo-LTR copies in *B. distachyon* (Stritt et al. 2019).

Despite the lack of global effect, a subset of 32 TIPs show a significant effect on

nearby gene expression at high confidence (Table S7), providing candidates for

further functional validation.

Conclusions

In contrast to the situation in *A. thaliana*, an eudicot of similar genome size and

complexity, TEs tend to have a limited impact on methylation spreading and only

punctual effects on gene expression in the model grass *B. distachyon*. Our findings

yet align to what is described in rice where TEs also have little impact on gene

expression. Methylation spreading has been suggested to be under purifying selection

and such a mechanism may largely constrain the impact of TEs on adjacent genes.

Compared to *A. thaliana*, TEs and genes are largely interspersed in *B. distachyon*. We

thus suggest that the differences we observed between the two species regarding the

impact of TEs are largely inherent to the contrasting structure of their genomes.

Materials and Methods

Plant material

We used eleven accession of *B. distachyon* originating from Iraq (Bd21, Bd21-3, Bd3-1), Turkey (BdTR12c, BdTR7a, Koz3, Bd1-1) and Spain (ABR2, Sig2, Luc1, Bd30-1). Seeds were sown on soil-sand mixture in 50 ml pots. For bisulfite and for RNA sequencing, three biological replicates per accession were grown. The second and third leaves of each replicate were sampled after 24 days of 16/8 hours of day/night conditions.

Whole genome bisulfite sequencing (WGBS) and RNA-sequencing analysis

For *B. distachyon*, WGBS libraries were prepared as described in the supplementary material Method S1. For *A. thaliana,* we used publicly available data (Table S1 for references). Raw data were processed as described in Method S1 and mapped onto the reference genomes Bd21 for *B. distachyon* (Phytozome 12, https://phytozome.jgi.doe.gov) and Col-0 for *A. thaliana* (TAIR10, www.arabidopsis.org). Methylation levels were estimated following the method described in (Schultz et al. 2012; Method S1) and by averaging the percentage of methylated cytosines over the three replicates per genotype within each focal unit (e.g. 100 kb/100 bp genomic windows, TE and gene body). We drew Circos plots (Connors et al. 2009) to visualize the architecture of the two reference genomes. We eventually used GLM models with logit transformation to characterize the relationship between gene or TE density and methylation levels. P-values were obtained with the R package multcomp (Hothorn et al. 2008).

For each of the eleven *B. distachyon* accessions, mRNA was extracted using the RNAeasy Plant Mini Kit from Qiagen. Libraries were prepared by the D-BSSE Basel and paired-end sequencing was performed on a Illumina HiSeq 2500. For *A. thaliana,* we again used publicly available data (Table S1). Reads were mapped to the reference genomes with Salmon quant v.0.12.0 (Patro et al. 2017). The abundance of gene expression was quantified averaging transcripts per million (TPM) per accession for each single genotype over the three replicates.

 Methylation spreading

 We estimated methylation spreading by calculating methylation levels in 100 bp
 windows spanning 2.5 kb upstream and downstream of each TE insertion. TE
 insertions were categorized with regard to their position in
 centromeric/pericentromeric regions or on chromosome arms and aggregated at the
 family- (for the ten most active TE families) or superfamily levels. Heatmaps
 depicting methylation levels around TEs were drawn in R using ggplot2 (Wickham
 2009). We also used the levels of methylation spreading around TEs to calculate
 Euclidean distances between the ten most active TE families and to perform a
 hierarchical clustering using hclust in R (R development core team 2013).

 Impact of TEs on gene expression

 We assessed the potential link between the presence of a TE in the vicinity of a gene
 and expression by assigning genes to different categories depending on whether they
 harbor a TE in their sequence or within windows ranging from near vicinity (0-200
 bp) to up to 2 kb. Average gene TPM and standard errors were calculated per distance
 class with all TEs pooled together as well as for the ten most active families
 separately and plotted using the R package plotrix (Lemon 2006). For a subset of
 candidate families, we also compared the expression level (log1p transformed) of
 genes displaying a TE in their vicinity (0-400 bp for RLG_Bdis0039, RLG_Bdis004
 and RLC_Bdis022; 0-1 kb for RLG_Bdis180) to the expression level of genes
 displaying a TE further away (between 400 bp-2 kb and 1 kb-2 kb respectively) with
 a Wilcoxon test.

 Transposable element insertion polymorphism analysis

 Detection of transposable element insertion polymorphisms (TIP) was performed with
 detettore (github.com/cstritt/detettore) using sequencing data from (Gordon et al.
 2017). The expression of genes harboring a TIP in their vicinity (<1 kb) in at least one
 of the accessions was compared to the corresponding genes in the other accessions.
 For each single gene, difference in expression between groups was tested using a t-
 test with a confidence interval of 0.95. Change in gene expression was considered
 significant when the P-value was <0.05 and the log2 change >1.

Acknowledgment

This work was supported by the PSC-syngenta fellowship, the University Research

Priority Program Evolution in Action of the University of Zurich and the Swiss

National Science Foundation (PZ00P3_154724). The authors would like to thank the

D-BSSE for the sequencing effort, the Genetic Diversity Center at ETH Zurich for

providing access to their computing resources and Michael Thieme for his comments

on the manuscript.

Data availability

All the data produced for the analysis will be made publically available upon
acceptance of the manuscript. Scripts used in this study are available at
https://github.com/mwylerCH.
272

Literature

Bonchev G, Parisod C. 2013. Transposable elements and microevolutionary changes in natural populations. Mol. Ecol. Resour. 13:765–775.

Choi JY, Purugganan MD. 2017. Evolutionary Epigenomics of Retrotransposon-Mediated Methylation Spreading in Rice. Mol. Evol. Biol. 35:365–382.

Connors J, Krzywinski M, Schein J, Gascoyne R, Horsman D, Jones SJ, Marra MA. 2009. Circos : An information aesthetic for comparative genomics. Genome Res. 19:1639–1645.

Eichten S, Stuart T, Srivastava A, Borevitz J. 2016. DNA Methylation profiles of diverse Brachypodium distachyon aligns with underlying genetic diversity. Genome Res. 26:1520–1531.

Eichten SR, Ellis NA, Makarevitch I, Yeh C, Gent JI, Guo L, Mcginnis KM, Zhang X, Schnable PS, Vaughn MW, et al. 2012. Spreading of Heterochromatin Is Limited to Specific Families of Maize Retrotransposons. PLoS Genet 8(12): e1003127.

Gordon SP, Contreras-moreira B, Woods DP, Marais DL Des, Burgess D, Shu S, Stritt C, Roulin AC, Schackwitz W, Tyler L, et al. 2017. Extensive gene content variation in the Brachypodium distachyon pan-genome correlates with population structure. Nat. Commun. 8:2184.

Grandbastien M-A. 1998. Activation of plant retrotransposons under stress conditions. Trends Plant Sci. 3:181–187.

Grandbastien M. 2005. Stress activation and genomic impact of Tnt1 retrotransposons in Solanaceae. Cytogenet. Genome Res. 110:229–241.

Guio L, Barron MG, Gonzalez J. 2014. The transposable element Bari-Jheh mediates oxidative stress response in Drosophila. Mol Ecol 23:2020–2030.

Hasterok R, Wang K, Jenkins G. 2019. Progressive refinement of the karyotyping of Brachypodium genomes. New Phytol. epub ahead.

Hollister JD, Smith LM, Guo Y, Ott F, Weigel D, Gaut BS. 2010. Transposable elements and small RNAs contribute to gene expression divergence between Arabidopsis thaliana and Arabidopsis lyrata. Proc Natl Acad Sci U S A 108: 2322–2327.

Hothorn T, Bretz F, Westfall P. 2008. Simultaneous Inference in General Parametric Models. Biometrical Journal, 50(3), 346–363.

Horváth V, Merenciano M, González J. 2017. Revisiting the Relationship between Transposable Elements and the Eukaryotic Stress Response. Trends in Genetics 33:832–841.

Liaw and M. Wiener. 2002. Classification and Regression by randomForest. R News 2:18-22.

Lemon J. 2006. Plotrix: a package in the red light district of R. R-News 6:8-12.

Makarevitch I, Waters AJ, West PT, Stitzer M, Hirsch CN, Ross-Ibarra J, Springer NM. 2015. Transposable elements contribute to activation of maize genes in response to abiotic stress. PLoS Genet 11:e1004915.

Mcclintock B. 1947. Cytogenetic Studies of Maize and Neurospora. Carnegie Inst

Wash Yrbk 46:146–152.

McClintock B. 1950. The origin and behavior of mutable loci in maize. Proc Natl Acad Sci U S A. 36:344–355.

Negi P, Rai AN, Suprasanna P. 2016. Moving through the Stressed Genome: Emerging Regulatory Roles for Transposons in Plant Stress Response. Front Plant Sci 7:1448.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-aware quantification of transcript expression. Nature methods 14(4)_417-418.

R development core team. 2013. R: A language and environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. Available from: http://www.R-project.org.

Regulski M, Lu Z, Kendall J, Donoghue MTA, Reinders J, Llaca V, Deschamps S, Smith A, Levy D, McCombie WR, et al. 2013. The maize methylome influences mRNA splice sites and reveals widespread paramutation-like switches guided by small RNA. Genome Res. 23:1651–1662.

Rey O, Danchin E, Mirouze M, Loot C, Blanchet S. 2016. Adaptation to Global Change : A Transposable Adaptation to Global Change : A Transposable Element – Epigenetics Perspective. Trends Ecol. Evol. 31(7):514-526

Schultz MD, Schmitz RJ, Ecker JR. ' Leveling ' the playing field for analyses of single-base resolution DNA methylomes. Trends Genet. 28:583–585.

Seymour DK, Koenig D, Becker C, Weigel D. 2014. Evolution of DNA Methylation Patterns in the Brassicaceae is Driven by Differences in Genome Organization. PLoS Genet. 10: e1004785.

Shang Y, Yang F, Schulman AH, Zhu J, Jia Y, Wang J. 2017. Gene Deletion in Barley Mediated by LTR-retrotransposon BARE. Sci. Rep. 7: 43766.

Sigman MJ, Slotkin RK. 2016. The First Rule of Plant Transposable Element Silencing: Location, Location, Location. Plant Cell 28:304–313.

Stritt C, Gordon SP, Wicker T, Vogel JP, Roulin AC. 2018. Recent Activity in Expanding Populations and Purifying Selection Have Shaped Transposable Element Landscapes. Genome Biol. Evol.10:304–318.

Stritt C, Wyler M, Gimmi EL, Pippel M, Roulin AC. 2019. Diversity , dynamics and effects of long terminal repeat retrotransposons in the model grass Brachypodium distachyon. New Phytol. Epub ahead of print doi: 10.1111/nph.16308

Stuart T, Eichten S, Cahn J, Borevitz JO, Lister R. 2016. Population scale mapping of transposable element diversity uncovers novel genetic diversity linked to gene regulation and epigenomic variation. Elife 5:e20777.

Takuno S, Gaut BS. 2013. Gene body methylation is conserved between plant orthologs and is of evolutionary consequence. Proc. Natl. Acad. Sci. U. S. A. 110:1797–1802.

The international brachypodium Consortium. 2010. Genome sequencing and analysis of the model grass Brachypodium distachyon. Nature 463:763–768.

Thieme M, Lanciano S, Balzergue S, Daccord N, Mirouze M, Bucher E. 2017.
Inhibition of RNA polymerase II allows controlled mobilisation of
retrotransposons for plant breeding. Genome Biol. 18:134:1–10.

Thind AK, Wicker T, Müller T, Ackermann PM, Steuernagel B, Wulff BBH,
Spannagl M, Twardziok SO, Felder M, Lux T, et al. 2018. Chromosome-scale
comparative sequence analysis unravels molecular mechanisms of genome
dynamics between two wheat cultivars. Genome Biol. 19:104

Van't Hof AE, Campagne P, Rigden DJ, Yung CJ, Lingley J, Quail MA, Hall N,
Darby AC, Saccheri IJ. 2016. The industrial melanism mutation in British
peppered moths is a transposable element. Nature 534:102–105.

Wang X, Weigel D, Smith LM. 2013. Transposon variants and their effects on gene
expression in Arabidopsis. PLoS Genet. 9:e1003255.

Wang Y, Liang W, Tang T. 2018. Constant conflict between Gypsy LTR
retrotransposons and CHH methylation within a stress-adapted mangrove
genome. New Phytol. 220: 922-935.

Wickham. H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag
New York. ISBN 978-3-319-24277-4.

Zhang X, Yazaki J, Sundaresan A, Cokus S, Chan SW, Chen H, Henderson IR, Shinn
P, Pellegrini M, Jacobsen SE, et al. 2006. Genome-wide high-resolution
mapping and functional analysis of DNA methylation in arabidopsis. Cell
126:1189–1201.

Zhang H, Lang Z, Zhu J. 2018. Dynamics and function of DNA methylation in plants.
Nat. Rev. Mol. Cell Biol. 19:489–506.

Figures and Tables

Figure 1: Genome organization and methylation distribution

Circos plot displaying methylation levels as well as gene and TE density in 100 kb windows in A) *A. thaliana* and B) *B. distachyon*. Centromeric and pericentromeric regions are highlighted in grey. Panel C and D display CpG methylation levels (expressed as % of methylated cytosines) for the most common TE superfamilies and for genes in *A. thaliana* and *B. distachyon* respectively.

Figure 2: Methylation spreading around transposable elements

Heatmaps representing methylation (CpG) spreading around TEs located in the pericentromere/centromere regions or on the arm of the chromosomes for *A. thaliana* and *B. distachyon*.

Figure 3: Impact of transposable elements on gene expression

A) Average gene expression as a function of the distance to the nearest TE belonging to the four LTR-retrotransposons RLG_BdisC039, RLG_BdisC004, RLG_BdisC180 and RLC_BdisC022 in the natural accessions of *B. distachyon*. Each coloured line represents a different accession. The dashed line represents the genome-wide level of expression. Boxplots display the expression of genes harbouring a TE in their vicinity for two distance classes only, all accessions pooled together. Stars indicate Wilcoxon test levels of significance: ***<0.001 B) Heatmap representing methylation (CpG) spreading around the 10 most active LTR retrotransposon families. Families were clustered based on the level of methylation spreading (long, intermediate and short) around each insertion. Names on the right of the heatmap describe the TE lineage to which each family belongs. Boxplots display the age of each families estimated by (Stritt et al. 2019).

Supplementary Materials

Figure S1: Relationship between gene or transposable element density and
methylation levels.

Figure S2: Impact of TEs on gene expression. Average gene expression as a
function of the distance to the nearest TE in *A. thaliana* and in the natural accessions
of *B. distachyon,* all TEs pooled together.

Figure S3: Transposable element insertion polymorphisms. Volcano plot
displaying expression changes in genes harbouring a TIP in their vicinity. Significant
changes are displayed in red.

Table S1: WGB- and RNA-sequencing output information

Table S2: Within-species glm outputs (CpG context)

Table S3: Within-species glm outputs (CHG context)

Table S4: Within-species glm outputs (CHH context)

Table S5: glm outputs - comparison *A. thaliana* vs. *B. distachyon*

Table S6: Information about the 1,833 TIPs identified across the natural
accession of *B. distachyon.*

Table S7: Information about the 32 high-confidence TIPs

Method S1: Extended Materials and Methods

A) *A. thaliana*

B) *B. distachyon*

A. thaliana

Distance from TE insertion (in kb)

B. distachyon

Distance from TE insertion (in kb)

Methylation (in %)

Methylation (in %)

A)

B)