Targeted control of toxin production by a mobile genetic element in

Streptococcus pneumoniae.

Emily Stevens[1], Daniel J. Morse[1], Dora Bonini[1], Seána Duggan[1], Tarcisio Brignoli[1], Mario

Recker[2], John A. Lees[3], Nicholas J. Croucher[3], Stephen Bentley[4], Daniel J. Wilson[5], Sarah

G. Earle[5], Robert Dixon[5], Angela Nobbs[6], Howard Jenkinson[6], Tim van Opijnen[7], Derek

Thibault[7], Oliver J. Wilkinson[8], Mark S. Dillingham[8], and Ruth C. Massey[1]*.

1: School of Cellular and Molecular Medicine, University of Bristol, Bristol, BS8 1TD, UK.

2: College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter,

EX4 4QF, UK.

3: MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease

Epidemiology, St. Mary's Campus, Imperial College London, London, W2 1PG, UK.

4: The Wellcome Trust Sanger Institute, Wellcome Trust Genome Campus, Hinxton,

Cambridge CB10 1SA, UK.

5: Big Data Institute, Nuffield Department of Population Health, University of Oxford, Oxford

OX3 7LF, UK.

6: Bristol Dental School, University of Bristol, Bristol, BS1 2LY, UK.

7: Biology Department, Boston College, Chestnut Hill, MA, USA.

8: DNA-Protein Interactions Unit, School of Biochemistry, University of Bristol, Bristol, BS8

1TD, UK.

* Corresponding author (email: ruth.massey@bristol.ac.uk)

Introductory Paragraph

Streptococcus pneumoniae is a major human pathogen that can cause severe invasive diseases such as pneumonia, septicaemia and meningitis[1-3]. Young children are at a particularly high risk, with an estimated 800,000 deaths worldwide in those under five attributable to invasive pneumococcal disease each year[1-3]. The cytolytic toxin pneumolysin (Ply) is a primary virulence factor for this bacterium, however, despite its importance to both the colonisation and pathogenic capabilities of this pathogen, the regulation of its expression is not well defined[4-7]. Using a genome-wide association approach we identified over a hundred potential affectors of Ply activity, including the Integrative and Conjugative Element (ICE) ICE*Sp*23FST81[8]. This regulatory effect is mediated through the activity of a novel modular protein, ZomB, which has an N-terminal UvrD-like helicase domain followed by two Cas4-like nuclease domains. The ZomB protein has potent ATP-dependent nuclease activity and binds specifically to the DNA containing a BOX repeat region that forms part of the *ply* operon. We hypothesise that with over 100 BOX regions across the pneumococcal genome, the acquisition of the *zomB* gene on ICE*Sp*23FST81 has the potential to re-wire the transcriptional landscape of this major human pathogen.

Main Text

Ply is a cytolytic pore-forming toxin that binds to cholesterol in the membranes of host cells, where it inserts into the lipid bilayer forming a transmembrane pore that ultimately causes lysis of the host cell[4-7]. Ply has also been shown to interfere with several aspects of the host immune system[9,10], been implicated in the initial colonization of the nasopharynx[11], and shown to promote transmission of the bacteria to new hosts[12]. Given its importance, we sought to define the genetic basis of Ply activity by analysing a collection of 165 isolates belonging to the *S. pneumoniae* PMEN1 lineage. We constructed a *ply* mutant in the pneumococcal isolate ATCC 700669[8] by replacing the *ply* gene with an erythromycin resistance cassette. Using this wild type and mutant strain as positive and negative controls we quantified the Ply activity (lysis of sheep red blood cells (RBCs)) of 165 clinical isolates in

triplicate, demonstrating that it varied significantly across this collection of closely related

isolates (fig. 1a).

As the genomes of each of these isolates has been sequenced[13], we applied three

complementary genome-wide association (GWAS) approaches to identify loci associated

with Ply activity. In addition to a linear regression approach using the SNP (single nucleotide

polymorphism) data, we also applied two methods that make use of kmer data: *BugWAS*[14]

and *pyseer*[15]. The results from each of these methods are presented in supplementary

tables 1-3. A Manhattan plot in Fig. 1b shows the significantly associated genetic loci

determined through the linear regression method, highlighting those loci where the GWAS

methods agree. There were two notable observations from these analyses. The first was that

five loci were associated with Ply activity across all three methods, with the *pbpX* gene which

encodes the Penicillin Binding Protein 2x being the most significantly associated locus. This

was followed in order of significance by the intergenic region between a gene with the locus

tag SPN23F05820 and *bgaA,* a gene with the locus tag SPN23F00840, the intergenic region

between a gene with the locus tag SPN23F19120 and *msmG,* and the intergenic region

between a gene with the locus tag SPN23F14800 and *greA*. What is striking about these

four latter loci is that they each contain a BOX repeat region, which are a family of short

repetitive sequences of DNA unique to the pneumococcal species[16]. Interestingly, a further

67 BOX containing regions were associated with Ply activity across the three GWAS

analyses. The second notable observation was that 48 individual genes or intergenic regions

on the Integrative and Conjugative Element (ICE) ICE*Sp*23FST81 were associated with Ply

activity.

Fig. 1: Genome wide association study (GWAS) to identify novel effectors of pneumolysin (Ply) production by *S. pneumoniae*. (**a**) Pneumolysin activity of 165 *S. pneumoniae* of the PMEN1 lineage, as measured by cell lysis. A wild type (WT) and isogenic pneumolysin mutant (Δ*ply*) have been included as controls. (**b**) Manhattan plot of the GWAS results using the linear regression method. The blue dotted line indicates the threshold for significance. SNPs in loci identified by two or all three GWAS methods are indicated in blue and orange respectively. The black arrows indicate the regions of interest; 1: SPN23F00840, 2: *pbpX,* 3: intergenic between SPN23F05820 and *bgaA*, 4: ICE*Sp*23FST81, 5: intergenic between SPN23F14800 and *greA,* and 6: intergenic between SPN23F19120 and *msmG.*

Fitness trade-offs between antibiotic resistance and virulence in bacteria are well

established[17-20], and for both clinical isolates and isogenic mutants the acquisition of

penicillin resistance has been shown to decrease the virulence of *S. pneumoniae* in murine

models of infection[21,22]. Given the association of the *pbpX* gene and Ply activity by all three

GWAS methods, we hypothesised that a similar trade-off may be occurring here, such that

the polymorphisms in the *pbpX* gene may be increasing the levels of resistance to penicillin,

which may as a consequence reduce the levels of Ply being produced, or vice versa.

Although bacterial GWAS results are typically validated through mutation of the associated

locus, the contribution of the protein encoded by *pbpX* to the biosynthesis of the

peptidoglycan layers in the bacterial cell wall is such that it is essential and cannot be

inactivated. Instead, we tested our hypothesis by examining the levels of resistance to

penicillin (minimum inhibitory concentrations, MICs) for the isolates, but found no significant

correlation between the MICs and Ply activity (Pearson product-moment correlation r^2 =

0.076, P = 0.36). An alternative hypothesis is that the polymorphisms in the *pbpX* gene

affects the stem peptide composition of peptidoglycan, as this has been shown to inhibit the

release of Ply from the bacterial cells[23]. Further work to test this hypothesis is underway.

The high number of associated loci on ICE*Sp*23FST81 is particularly intriguing. This mobile

genetic element (MGE), which can be found both integrated and in plasmid form (illustrated

in fig. 2a), is believed to be critical to the success of this lineage of *S. pneumoniae* due to the

antibiotic resistance capabilities it brings to the bacteria[8]. Given its ability to move

horizontally as a single contiguous unit between bacteria, it is likely that of the associated

loci only one is an affector of Ply activity, whilst the others are associated through their

physical linkage to this. A survey of the putative activity of all 48 associated loci revealed that

the majority of these are genes typical to such elements, involved in antibiotic resistance and

the mechanics of its movement. An interesting exception is a gene with the locus tag

SPN23F12470. This locus has been annotated as encoding a UvrD-like helicase, a family of

proteins typically associated with core house-keeping activities for bacteria. Further *in silico*

analysis of the encoded protein, that we have named ZomB, suggests that it is a multi-

domain protein with a putative DNA binding helicase domain, followed by two Cas4-like

nuclease domains which are predicted to harbour 4Fe-4S clusters[24] (fig. 2b). With several

known examples of Cas-like proteins regulating bacterial virulence[25-27], we hypothesised that

it is this gene on ICE*Sp*23FST81 that is the affector of Ply activity.

To establish the role of ZomB in Ply activity we replaced the *zomB* gene with an

erythromycin resistance cassette in *S. pneumoniae* strain ATCC 700669. The RBC lytic

activity of the *zomB* mutant was significantly impaired relative to the wild type strain (fig. 2c).

Using anti-Ply antibodies in a Western blot, we further demonstrated that this reduction in

lytic activity was due to a decrease in the amount of Ply protein being produced by the

bacteria (fig. 2d). We also quantified the relative transcription of the *ply* gene by qRT-PCR

and found that the reduced abundance of Ply was likely due to a significant decrease in *ply*

transcription (fig. 2e). Multiple unsuccessful attempts to construct a strain containing the

zomB gene on a plasmid to complement its activity in the mutant led us to conclude that high

levels of expression of this protein are toxic to the bacterial cells.

2a

2b

ZomB protein (approx. 150kDa)

2c
2d
2e

133
Fig. 2: Inactivation of the *zomB* gene on the ICE*Sp*23FST81 affects pneumolysin
production. **(a)** Cartoon illustration of ICE*Sp*23FST81 in both its linear chromosomally
integrated form and circularised plasmid form. Some genes of interest have been included:
SPN23F12410, SPN23F13170 and *rplL* because these flank the element in its linear form
and come into close association when in the plasmid form. Genes encoding the antibiotic
resistance gene are also indicated. **(b)** Schematic of the ZomB protein with its helicase and
two Cas4-like nuclease domains indicated. **(c)** the lytic activity of the *zomB* mutant is
comparable to that of the *ply* mutant as determined by a sheep RBC lysis assay. **(d)** The
presence of Ply in the extracellular medium is reduced in the *zomB* mutant, determined
using anti-Ply antibodies in a Western blot on concentrated bacterial supernatant. **(e)** The
effect the inactivation of *zomB* has on Ply production is mediated at a transcriptional level, as
determined by qRT-PCR. This data was made relative to the expression of the

housekeeping gene *recA* in each sample, and normalised to the level of expression on the
ply gene in the wild type strain. In the box plots each dot represents an individual replicate
data point.
149

The *ply* gene is transcribed as part of an operon with four other genes that encode a

transcriptional regulator (SPN23F19460) and proteins implicated in the movement of Ply

from the cytosol to the bacterial cell wall (SPN23F19460-19500) (fig. 3a) [28]. It also contains a

BOX repeat region immediately downstream of the *ply* gene. Due to their internal repeating

sequences, BOX regions can form stable secondary structures, and their presence has been

associated with effects on the transcription of neighbouring genes[29]. The molecular details of

how their presence affects gene transcription has not yet been determined but is likely due

to these secondary structures where they can either enhance or interfere with transcription

processes depending on their relative positioning[30]. As ZomB appears to be a positive

effector of *ply* transcription, and given its likely DNA binding capability, we hypothesised that

it may be interacting with this BOX regions to elicit its effect on the expression of *ply*.

To characterise the biochemical activities of the ZomB protein, we expressed and purified

recombinant ZomB with an N-terminal 6x histidine tag (fig. 3b). Using a coupled-assay, we

found that ZomB hydrolyses ATP with Michaelis-Menten kinetics and displays a turnover

number of approximately 20 s^{-1} and a K_m value of 90 μM ATP (fig. 3c). These experiments

were performed in the presence of saturating quantities of ssDNA, which was shown to

strongly stimulate ATP hydrolysis with an apparent dissociation constant of approximately 1

μM (ntds) (fig. 3d). This behaviour is typical of the UvrD-like DNA helicases of which ZomB

is a member[31]. To examine the putative nuclease activity, ZomB protein (50nM) was

incubated with duplex DNA with and without ATP and $Mg^{2+.}$ Degradation of the DNA by

ZomB was monitored by gel electrophoresis, which showed that the protein possesses a

potent ATP-dependent nuclease activity (fig. 3e).

We next amplified the region of DNA containing the BOX element from within *ply* operon, as

well as an equivalently-sized region of control DNA within the *ply* coding region, and

performed electrophoretic mobility shift assays (EMSAs) with increasing concentrations of

ZomB protein. As visualised in fig. 3f, the ZomB protein caused a shift in the BOX containing

DNA even at the lowest concentration of protein used, demonstrating its high affinity for this

substrate. There was also a shift in mobility of the control, *ply*-encoding DNA, but this

occurred at higher concentrations of ZomB and resulted in the formation of less distinctly

shifted DNA bands, which is indicative of non-specific binding. The binding activity of ZomB

was not specific to this BOX containing region as we found that it also bound with high

affinity to a second BOX containing region of DNA situated approximately 3kb upstream of

the *ply* gene (supp. fig. 1). These results are consistent with ZomB displaying binding

specificity for BOX elements. Although we have still much to learn about the *in vivo*

biochemical functions of ZomB, we believe it likely that its nuclease activity may unwind the

secondary structures that result from the BOX repeats, thereby relieving the effects these

have on the transcription of neighbouring genes.

191
Fig. 3: The ZomB protein specifically binds to the BOX region within the Ply encoding
operon. (**a**) cartoon of the *ply* encoding operon with the ply gene, the SPN23F- locus tags of
the neighbouring genes, and promoter and transcript length (black arrow) indicated. (**b**) An
SDS-PAGE gel showing the purified his tagged ZomB protein. (**c**) Steady-state ATPase
activity of ZomB (50 nM) was measured at saturating ssDNA concentration to determine the

Michaelis-Menten parameters. (**d**) Steady-state ATPase activity of ZomB is strongly
stimulated by ssDNA with an apparent dissociation constant of approximately 1 µM ntds. (**e**)
Nuclease assays were performed with linear DNA demonstrating that it was degraded by the
ZomB protein in the presence of both ATP and divalent cations. (**f**) EMSAs demonstrating
the specificity of binding of a BOX element by the ZomB protein. An equivalent length of
DNA from with the *ply* coding region was used as a control. The concentrations of ZomB
used were 2.5, 5, 10, 20, 40 and 80nM.
204
205
Through the application of a functional genomics approach to a large collection of

sequenced *S. pneumoniae* isolates, we have identified over a hundred novel putative

affectors of Ply activity (supp. tables 1-3). Of these associated loci we have determined the

molecular detail of the interaction between the ZomB protein encoded on ICE*Sp*23FST81

and the BOX repeat region within the *ply* operon. This however does not explain why so

many other BOX containing regions were associated with Ply activity. One potential

explanation we are exploring is whether the sequence variability within the associated BOX

regions affects the binding affinity of ZomB to them, which would affect the availability of

ZomB for binding to the BOX region within the *ply* operon. It will also be interesting to see

with future work how ZomB contributes to the lifecycle of the ICE*Sp*23FST81, as this MGE

does not contain any BOX-like regions, suggesting that if ZomB regulates the expression of

any genes on the ICE, it uses a different sequence motif to achieve this.

Through their carriage of cargo genes, MGEs confer significant advantages to their bacterial

hosts that enhance the success of both. ICE*Sp*23FST81 confers resistance to both

chloramphenicol and tetracycline, which provides a clear benefit upon exposure to

antibiotics. However, for many bacterial species antibiotic resistance often incurs a fitness

cost in the absence of the antibiotic, which can be offset, for example, by reducing the

energetically costly production of toxins[17,18]. Here, in stark contrast, we observe the

simultaneous acquisition of increased resistance with increased toxin production. Future

work will explore how these combined costs are compensated from the perspective of both

the host cell and MGE, to result in such a globally successful pathogen.

Acknowledgements

ES was a BBSRC SWBio DTP funded PhD student. R.C.M is a Wellcome Trust funded

Investigator. D.J.W. is a Sir Henry Dale Fellow, jointly funded by the Wellcome Trust and the

Royal Society (Grant 101237/Z/13/B) and is supported by a Big Data Institute Robertson

Fellowship. We would also like to thank Prof. Tim Mitchell for helpful discussions during the

preparation of this manuscript.

Materials and Methods

Bacterial strains and growth conditions.

Clinical isolates used in this study (listed in supp. table 4) had been previously sequenced[13]

and all belonged to the PMEN1 clone of which *S. pneumoniae* ATCC 700669 is the

reference strain. Strains were grown for 16-24 hours in 5% CO_2 at 37°C, on either brain-

heart infusion (BHI) blood agar plates containing 5% defibrinated horse blood, or in BHI

broth without blood.

Construction of deletion mutants in *S. pneumoniae*.

Genes were deleted in *S. pneumoniae* using linear PCR products as described previously[32].

In brief, a stitch PCR approach using the primers listed in table 1 were used to generate a

single PCR product consisting of 1kb to either side if the gene to be deleted, with a gene

encoding resistance to erythromycin (*ermAM* - amplified from plasmid pVA838) in the centre.

To transform the bacteria with this PCR product, the wild type bacteria were grown overnight

in 5ml BHI broth, and 2ml of this culture was used to inoculate a pre-warmed tube of 20ml

BHI broth. This was incubated for a further 2 hours, and 0.5ml was added to 9.5ml pre-

warmed BHI broth and incubated for 30 minutes. 1ml of this sub-culture was transferred to

sterile tubes, 10μl of 10μg/ml competence-stimulating peptide-2 (CSP-2) was added each,

and these were incubated for 10-15 minutes. To 200μl aliquots of this either the PCR

product, or sterile water was added and the mixture incubated for a further 2 hours. This was

then added to molten BHI agar (10ml) containing 5% defibrinated horse blood, allowed to set

in a petri dish and incubated for 2 hours. Plates were then overlaid with a further 10ml

molten agar containing 5% defibrinated horse blood with erythromycin (2µg/ml) to selected

for successfully transformed cells. Plates were incubated for up to five days at 37°C, or until

colonies appeared within the agar.

Amplification site	Primer Sequence
ply (LHS) forward	5' CCCTTGCTCTGGTTAAAAAAAGAAGC 3'
ply (LHS) reverse	5' ATATTTTTGTTCATATTTGCCATCTTCTACC 3'
ery-ply forward	5' GGTAGAAGATGGCAAATATGAACAAAAATATAAAA 3'
ery-ply reverse	5' CTACCTGAGGTTATTTCCTCCCGTT 3'
ply (RHS) forward	5' GAGGAAATAACCTCAGGTAGAAGATAAG 3'
ply (RHS) reverse	5' GATCACCTTTTTTAGCTGCTACATAG 3'
zomB (LHS) forward	5' TGCCCACTATTTTTATCTAGTTGCTTACC 3'
zomB (LHS) reverse	5' ATTTTTGTTCATTGTTGTCATCGTTTTACCTC 3'
ery-zomB forward	5' CGATGACAACAATGAACAAAAATATAA 3'
ery-zomB reverse	5' CTTTTCCGGATTCTTATTTCCTCCC 3'
zomB (RHS) forward	5' GGAGGAAATAAGAATCCGGAAAAG 3'
zomB (RHS) reverse	5' AATTAATTCCTGAATACAAGTTAACAAAATAG 3'

Quantification of Ply activity

Strains were grown overnight in BHI broth, then sub-cultured at a 1:1 ratio into fresh broth

and grown for one hour until OD_{600} of 0.4-0.7 was reached. These cultures were then serially

diluted 4-fold in a 96 well plate containing BSA assay buffer (0.05g bovine serum

albumin and 77mg dithiothreitol dissolved in 50ml sterile phosphate buffered saline. To each

well 50µl of triple washed sheep red blood cells were added and the plates were incubated

for one hour in 5% CO_2 at 37°C. The plates were then centrifuged at 2000rpm for 10

minutes at room temperature to separate the intact cells from the soluble lysed material. The

supernatants from each well were transferred to a fresh 96-well plate and absorbance values

read at 596nm were obtained using a FLUOstar Omega microplate reader.

GWAS

The initial Genome-wide associations between single nucleotide polymorphisms (SNP) and

bacterial toxicity were determined by means of linear regression. We performed a singular

value decomposition of the SNP data and used the first four principal components, which

together explained around 50% of the variance, in the linear regression model to adjust for

population structure. Subsequent GWAS analyses of the data by pyseer and BugWas were

performed as previously described[14,15].

Quantification of Ply production

The bacteria were grown overnight in 15ml BHI broth and harvested by centrifugation.

Proteins were extracted from the cell pellets of by resuspending it in 1.5ml of whole cell

lysate buffer (0.01% SDS, 0.1% sodium deoxycholate and 0.015M sodium citrate) followed

by an incubation for 30 minutes at 37°C. The individual proteins were separated on an SDS-

PAGE gels and Western blotting of this conducted using anti-pneumolysin antibody followed

by goat anti-mouse IgG HRP was used as the secondary antibody. The HRP signal was

detected using the Immun-Blot Opti-4CN Colourimetric kit from BioRad.

Quantification of transcription of the *ply* gene.

Bacteria were grown overnight in 5ml BHI broth and RNA was extracted using a Zymo

Quick-RNA Fungal/Bacterial Miniprep kit following the manufacturer's protocol. Turbo DNase

digest kit was used to remove any contaminating DNA. The quality and quantity of the RNA

was determined using a Nanodrop and the RNA was reverse transcribed to cDNA using a

qScript cDNA synthesis kit, as per the manufacturer's protocol. RNA samples were

standardised such that 100ng was added to each cDNA synthesis reaction. cDNA was

subsequently quantified using a Nanodrop and where necessary was standardised again to

ensure the same quantity in each sample. qPCR was performed using a Mic qPCR cycler

(Bio Molecular Systems) and reactions were set up using a SensiFAST SYBR

No-ROX master mix. Three technical repeats were conducted for each cDNA

Sample and the data anlaysed using the $2^{(Ct\ ply\ -\ Ct\ recA)}$ method. The primer pairs were used

where *recA* was used as the control housekeeping gene:

ply FW: GGCTGATTTCGCTGAACAAG

ply RV: CCTTGAGTTGTTCCATGCTG

recA FW: ATCGGAGATAGCCATGTTGG

recA RV: ATAGAGGCGCCAAGTTTACG

Expression and purification of ZomB.

The *zomB* gene was cloned into the expression plasmid pET15b, and expressed in E. coli

strain BL21(DE3) with a 6x histidine tag at its N-terminus. The *E. coli* cells were grown at

37°C in LB broth containing 100µg/ml ampicillin to an OD(600) of around 0.5, whereupon

cells weretemperature acclimatised to 18°C and gene expression was induced for 12 hours

via the addition of 1mM IPTG. Cells were harvested by centrifugation for 20minutes at

6000rpm and resuspended in buffer containing 50mM Tris pH 7.5,150mM NaCl, 10%

sucrose and 0.1mM PMSF. Cells were lysed by sonication in buffer containing 1mM TCEP,

500mM NaCl, 20mM imidazole and a proteaseinhibitor cocktail, and cell debris was removed

by centrifugation for 30 minutes at21,500rpm. Cell lysate was loaded onto a 5ml HisTrap

Nickel binding columnwhich had been equilibrated in HisB buffer (20mM Tris pH7.5, 500mM

NaCl,1mM TCEP, 5% glycerol) plus 20mM imidazole. Proteins possessing His-tagswere

eluted with a gradient from 20mM to 500mM imidazole over 32 minutes with a flow rate of

3ml/min. Protein elution was monitored by measuring absorbanceat 280nm. Fractions

containing the most protein were pooled and applied to a HiTrap Heparin affinity column

equilibrated with HepQ/B buffer (20mM TrispH7.5, 1mM TCEP) plus 100mM NaCl. Heparin-

binding proteins were eluted witha gradient from 100mM to 1M NaCl over 30 minutes,

monitored by absorbance at280nm. 2ml fractions containing the highest protein

concentrations were pooledand applied to a MonoQ anion exchange chromatography

column equilibratedwith HepQ/B buffer plus 100mM NaCl. Proteins were eluted with a

gradient from100mM to 1M NaCl over 30 minutes, with protein absorbance monitored at

280nm, and 0.3ml fractions were collected. Fractions containing the highest concentrations

of eluted protein were pooled and stored in HepQ/B buffer plus 330nM NaCl, corresponding

to the salt concentration at which the protein was eluted from the column. Nanodrop A280

and the ZomB protein's predicted extinction coefficient (118390mol/L/cm) were used to

determine that ZomB had been purified to a concentration of 7.0µM.

Biochemical characterisation of ZomB activity.

ATPase activity was measured by coupling the hydrolysis of ATP to the oxidation of NADH

which gives a change in absorbance at 340 nm. Reactions were performed in a buffer

containing 20 mM Tris-Cl pH 8.0, 50 mM NaCl, 2 mM DTT, 1 mM $MgCl_2$, 50 U/mL lactate

dehydrogenase, 50 U/mL pyruvate dehydrogenase, 1 mM PEP and 100 µg/mL NADH.

Rates of ATP hydrolysis were measured over 1 min at 25℃ and the ssDNA substrate used

was Poly(dT). For calculation of K_{DNA} (defined as the concentration of DNA at which ATP

hydrolysis is half-maximal), the ATP concentration was fixed at 2 mM. The Michaelis-Menten

plot was performed at saturating DNA concentration which is defined as 10x the K_{DNA} value.

The concentration of ZomB was 50 nM in these assays unless indicated otherwise.

Electrophoretic mobility shift assays.

The intergenic region between *ply* and the neighbouring gene SPN23F19460, containing

BOX repeat regions was amplified (using the following primers; BoxF:

GAGAGGAGAATGCTTGCGAC and BoxR:

TAGGAATCTCCTTTTTTCACATTTTAATCTTTC). A region of the *ply* gene of equivalent

size was also amplified (using plyF: ATGGCAAATAAAGCAGTAAATGACTTTATAC and

plyR: GCCCCCTAAAATAACCGCCTTC). The purified ZomB protein was added in a range

of concentrations (5µM, 2.5µM, 1.25µM, 0.625µM, 0.3125µM and 0.15625µM) to 10nM

of the PCR products in a buffer containing 20mM Tris (pH 8), 200mM sodium chloride, 1mM

Tris(2-carboxyethyl) phosphine (TCEP) and 10% glycerol. Samples were then run on a 1.5%

agarose gel in 1X TAE buffer for 110 minutes at 90V. Following this the gel was stained in

TAE containing 1X SYBR Safe DNA gel stain for 30 minutes, and bands were visualised

using a Typhoon FLA 9500.

362 REFERENCES

1. Deng, X. *et al.* Whole-genome sequencing reveals the origin and rapid evolution of an
emerging outbreak strain of *Streptococcus pneumoniae* 12F. *Clin. Infect. Dis.* **62**, 1126-
1132. (2016).
2. Ioachimescu, O.C., Ioachimescu, A.G. & Iannini, P.B. Severity scoring in community
acquired pneumonia caused by *Streptococcus pneumoniae*: a 5-year experience. *Int. J.*
Antimicrob. Agents **24**, 485-490 (2004).
3. O'Brien, K.L. *et al.* Burden of disease caused by *Streptococcus pneumoniae* in children
younger than 5 years: global estimates. *Lancet* **374,** 893-902 (2009).
4. Canvin, J.R. *et al.* The role of pneumolysin and autolysin in the pathology of pneumonia
and septicaemia in mice infected with a type 2 pneumococcus. *J. Infect. Dis.* **172**, 119-
123 (1995).
5. Hirst, R.A. *et al.* The role of pneumolysin in pneumococcal pneumonia and meningitis.
Clin. Exp. Immunol. **138**, 195-201 (2004).
6. Hirst, R.A. *et al. Streptococcus pneumoniae* deficient in pneumolysin or autolysin has
reduced virulence in meningitis. *J. of Infect. Dis.* **197**, 744-751 (2008).
7. Mitchell, A.M. & Mitchell, T.J. *Streptococcus pneumoniae*: virulence factors and
variation. *Clin.l Microbiol. Infect. Dis.* **16**, 411-418 (2010).
8. Croucher, N.J. *et al.* Role of Conjugative Elements in the Evolution of the Multidrug-
Resistant Pandemic Clone *Streptococcus pneumoniae* Spain23F ST81. *J Bacteriol.*
191, 1480-9 (2009).
9. Mitchell, T.J. *et al.* Complement activation and antibody binding by pneumolysin via a
region of the toxin homologous to a human acute-phase protein. *Mol. Microbiol.* **5**, 1883-
8 (1991).
10. Paton, J.C., Rowan-Kelly, B. & Ferrante, A. Activation of human complement by the
pneumococcal toxin pneumolysin. *Infect. Immun.* **43**, 1085-7 (1984).
11. Hotomi, M. *et al.* Pneumolysin plays a key role at the initial step of establishing
pneumococcal nasal colonization. *Folia Microbiologica*. **61**, 375-83 (2016).
12. Zafar, M.A., Wang, Y., Hamaguchi, S. & Weiser, J.N. Host-to-Host Transmission of
Streptococcus pneumoniae Is Driven by Its Inflammatory Toxin, Pneumolysin. *Cell Host*
Microbe. **21**, 73-83 (2017).
13. Croucher, N.J. *et al.* Rapid pneumococcal evolution in response to clinical interventions.
Science. **331**, 430-4 (2011).
14. Earle, S.G. *et al.* Identifying lineage effects when controlling for population structure
improves power in bacterial association studies. *Nat. Microbiol.* **1**, 16041 (2016).

15. Lees, J.A. *et al.* pyseer: a comprehensive tool for microbial pangenome-wide
association studies. *Bioinformatics.* **34**, 4310-2 (2018).
16. Martin, B. *et al.* A highly conserved repeated DNA element located in the chromosome
of Streptococcus pneumoniae. *Nucleic Acids Res.* **20**, 3479–83 (1992).
17. Rudkin, J.K. *et al.* Methicillin resistance reduces the virulence of healthcare-associated
methicillin-resistant *Staphylococcus aureus* by interfering with the *agr* quorum sensing
system. *J Infect Dis* **205**, 798–806 (2012).
18. Collins, J. *et al.* Offsetting virulence and antibiotic resistance costs by MRSA. *ISME J.* **4**,
577–84.
19. Hraiech, S. *et al.* Impaired virulence and fitness of a colistin-resistant clinical isolate of
Acinetobacter baumannii in a rat model of pneumonia. *Antimicrob. Agents Chemother.*
57, 5120–1 (2013).
20. Linares, J.F., *et al.* Overexpression of the multidrug efflux pumps MexCD-OprJ and
MexEF-OprN is associated with a reduction of type III secretion in *Pseudomonas*
aeruginosa. J. Bacteriol. **187**, 1384–91 (2005).
21. Azoulay-Dupuis, E. *et al.* Relationship between Capsular Type, Penicillin Susceptibility,
and Virulence of Human *Streptococcus pneumoniae* Isolates in Mice. *Antimicrob.*
Agents Chemother. **44**, 1575–7 (2000).
22. Rieux, V., Carbon, C. & Azoulay-Dupuis, E. Complex relationship Between Acquisition
of Beta-Lactam Resistance and Loss of Virulence in *Streptococcus Pneumoniae. J*
Infect Dis. **184**, 66-72 (2001).
23. Greene, N.G., Narciso, A.R., Filipe, S.R. & Camilli, A. Peptidoglycan Branched Stem
Peptides Contribute to Streptococcus pneumoniae Virulence by Inhibiting Pneumolysin
Release. *PLoS Pathog.* **11**, e1004996 (2015).
24. Zhang, J., Kasciukovic, T. & White, M.G. The CRISPR Associated Protein Cas4 Is a 5′
to 3′ DNA Exonuclease with an Iron-Sulfur Cluster. *PLoS One.* **7**, e47232 (2012).
25. Sampson, T.R. *et al.* A CRISPR/Cas system mediates bacterial innate immune evasion
and virulence. *Nature.* **497**, 254-7 (2013).
26. Heidrich, N. *et al.* The CRISPR/Cas system in *Neisseria meningitidis* affects bacterial
adhesion to human nasopharyngeal epithelial cells. *RNA Biol.* **16**, 390-396 (2019).
27. Ma, K. *et al.* cas9 Enhances Bacterial Virulence by Repressing the regR Transcriptional
Regulator in *Streptococcus agalactiae. Infect Immun.* **86**, e00552-17 (2018).
28. Kimaro Mlacha, S.Z., *et al.* Phenotypic, genomic, and transcriptional characterization of
Streptococcus pneumoniae interacting with human pharyngeal cells. *BMC Genomics.*
14, 383 (2013).
29. Croucher, N.J., Vernikos, G.S., Parkhill, J. & Bentley, S.D. Identification, variation and
transcription of pneumococcal repeat sequences. *BMC Genomics.* **12**, 120 (2011).
30. Knutsen, E. *et al.* BOX Elements Modulate Gene Expression in *Streptococcus*
Pneumoniae: Impact on the Fine-Tuning of Competence Development. *J Bacteriol.* **188**,
8307-12 (2006).
31. Gilhooly, N.S., Gwynn, E.J. & Dillingham, M.S. Superfamily 1 Helicases. *Front Biosci.* **5**,
206-16 (2013).
32. Chen, L., Ge, X. & Xu, P. Identifying essential *Streptococcus sanguinis* genes using
genome-wide deletion mutation. *Methods Mol. Biol.* **1279**, 15-23 (2015).

Supp 1a

ZomB protein concentrations

Supp 1b

ZomB concentration (nM)

449
450
451
Supplementary Figure 1: ZomB binds to both BOX regions that flank the *ply* gene. EMSA
was performed as described previously, but lower concentrations of both ZomB 2.5-80nM
and DNA (50nM), and with the DNA radiolabelled gamma-32P-ATP to facilitate visualisation.
A third 882bp PCR product was amplified for this experiment containing the BOX repeat
regions upstream of ply, using the following primer sequences:
FW: CAGTAAGATACAGACTGCAAGGC
RV: TGAAAAAGAAACCATTCAGGGGTATTCGG
50nM of each of the three PCR product was added to a reaction containing 2µl
phosphorous-32 (gamma-32P-ATP), 2µl T4 polynucleotide kinase (PNK) and 5µl PNK
buffer,made up to 50µl with water. This reaction mix was incubated at 37°C for 1 hour,
then at 65°C for 20 minutes to heat inactivate the PNK enzyme. Radiolabelled
DNA was purified using Microspin S-400 columns as per the manufacturers
protocol. 2nM DNA was used in an EMSA exactly as described above, using the
protein concentrations from 2.5-80nM, with the exception that the gel was not
stained with SYBR safe, but squashed for three hours before being dried using a
vacuum-heated gel dryer (BioRad). Bands were visualised as before using a
Typhoon FLA 9500.