Title:

Maternal cortisol is associated with neonatal amygdala microstructure and connectivity in a sexually

dimorphic manner

Authors names:

David Q Stoye[a], Manuel Blesa[a], Gemma Sullivan[a], Paola Galdi[a], Gillian J Lamb[a], Gill S Black[a],

Alan J Quigley[b], Michael J Thrippleton[c], Mark E Bastin[c], Rebecca M Reynolds[d,*], James P

Boardman[a,c,*]

Author affiliations:

[a] MRC Centre for Reproductive Health, University of Edinburgh, UK

[b] Department of Radiology, Royal Hospital for Sick Children, Edinburgh, UK

[c] Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK

[d] Centre for Cardiovascular Science, University of Edinburgh, UK

*Equal contributions

Corresponding author:

Professor James P Boardman

MRC Centre for Reproductive Health

W1.26 Queen's Medical Research Institute,

47 Little France Crescent,

Edinburgh, EH16 4TJ, UK.

E: james.boardman@ed.ac.uk

T: +44 131 242 2567

Abstract

The mechanisms linking maternal stress in pregnancy with infant neurodevelopment in a sexually dimorphic manner are poorly understood. We tested the hypothesis that maternal hypothalamic-pituitary-adrenal axis activity, measured by hair cortisol concentration, is associated with microstructure, structural connectivity and volume of the infant amygdala. In 78 human mother-infant dyads, maternal hair was sampled postnatally, and infants underwent magnetic resonance imaging at term-equivalent age. Higher hair cortisol concentration was associated with higher left amygdala fractional anisotropy ($\beta=0.677$, $p=0.010$), lower left amygdala orientation dispersion index ($\beta=-0.597$, $p=0.034$), and higher fractional anisotropy in connections between the right amygdala and putamen ($\beta=0.475$, $p=0.007$) in girls compared to boys. Maternal cortisol during pregnancy is related to newborn amygdala architecture and connectivity in a sexually dimorphic manner. Given the fundamental role of the amygdala in the emergence of emotion regulation, these findings offer new insights into mechanisms linking maternal stress with adverse neuropsychiatric outcomes of children.

Impact Statement

Prenatal stress is transmitted to infant development through cortisol, which imparts sex-specific effects on the development and connectivity of the amygdalae.

Main Text

Introduction

Prenatal exposure to maternal stress is estimated to affect 10-35% of children worldwide, which is a major concern because early life stress is linked to impaired cognitive development, negative affectivity, autism spectrum disorder (ASD), and psychiatric diagnoses including attention deficit hyperactivity disorder (ADHD), addiction, depression and schizophrenia[1]. Neural correlates of prenatally stressed children include alternations in brain structural and functional connectivity, especially in networks involving the amygdala and prefrontal cortex[2].

Adaptation of the maternal hypothalamic-pituitary-adrenal (HPA) axis is a key mechanism by which maternal stress modulates offspring neurodevelopment[3], and there is evidence that this mechanism operates in a sexually dimorphic manner[4]. For, example, higher waking maternal salivary cortisol in pregnancy is associated with increased internalizing behaviours in female infants and reduced internalizing behaviours in males[5, 6]. Higher maternal salivary cortisol in pregnancy is also associated with stronger amygdala functional connectivity with networks involved in sensory processing and integration in newborn girls, with weaker connectivity to these brain regions in boys[7]; and in childhood, with larger amygdalae[8] and reduced segregation of structural networks in girls but not boys[9]. The amygdala is further implicated as a neural target of prenatal stress exposure by observations from studies that have characterised maternal stress by symptomatology of depression and / or anxiety, which report alterations in amygdala volume[10], microstructure[11], and functional and structural connectivity among offspring[12]

Candidacy of the amygdala as an important neural target of prenatal stress exposure comes from the following observations in pre-clinical and clinical studies. First, the amygdala develops early in embryonic life[13] and contains a high concentration of glucocorticoid receptors[14]; second, increased maternal glucocorticoids modulate amygdala development and anxiety-like behaviours in experimental models[15, 16]; third, lesion studies in non-human primates support its critical role in early development of emotion regulation[17]; fourth, newborn amygdala functional connectivity is consistently linked with internalizing behaviours in children up to the age of two years[7, 18]; fifth, early disruption to cell composition of the amygdala is reported in a model of early life stress[19], and in children with autism[20]; and sixth, in pre-clinical models, stress and glucocorticoid exposure induce dendritic arborization, amygdala hypertrophy and induce anxiety-like behaviours[21, 22].

Neonatal magnetic resonance imaging (MRI) serves as an intermediate phenotype for investigating the impact of early life exposures on brain and health because it is distal to the aetiological process,

in this case prenatal stress, and is also more proximal to cognitive, behavioural and disease
outcomes. Structural and diffusion MRI (dMRI) have been used to characterise brain structural
maturation and emerging network connectivity during the perinatal period, and to investigate
pathways to atypical development(23, 24). It is a suitable tool to investigate the impact of prenatal
stress exposure on the amygdala because age-specific templates enable accurate parcellation of
the amygdala and associated structures(25); and diffusion tensor imaging and neurite orientation
and dispersion density imaging (NODDI) support inference about tissue microstructure and network
connectivity, modelled by fractional anisotropy (FA), mean diffusivity (MD), orientation dispersion
index (ODI) and neurite density index (NDI)(26).
91
Hair cortisol concentration (HCC) measured in 3cm hair samples collected from close to the scalp
reflects basal HPA axis activity over the 3 months prior to sampling, and in contrast to single
measures from saliva or blood, it is not influenced by short-term activation of the HPA axis in
response to acute stressors(27). Studies in pregnant women have shown HCC to be an efficient
method of retrospective assessment of long-term cortisol secretion, and thus long-term HPA axis
activity(28, 29).
98
Previous studies have reported sex-specific differences between maternal stress and amygdala
functional connectivity and behavioural outcomes among children(5-9), but study designs leave
uncertainty about the mechanism linking maternal stress with amygdala development, the potential
confounding role of events and environmental exposures during childhood, and the impact of stress
on structural connectivity. Resolving these uncertainties is necessary for developing strategies
designed to improve socio-emotional development of children born to women who are stressed
during pregnancy. Based on studies of the imaging, biochemical and clinical phenotype of prenatal
stress exposure, we hypothesised that higher levels of maternal HPA activity in the final months of
pregnancy ascertained from maternal HCC would impact amygdala development and structural
connectivity of offspring infants in a sexually-dimorphic manner, and that these effects would be
apparent around the time of birth.
110
111 **Results**
112
113 ***Participant characteristics***
114
The parents of 102 infants consented to take part. Of these, 2 preterm infants died before term
equivalent age, 12 did not complete the MRI protocol or images were not amenable processing
due to movement artefact; 1 had an incidental structural anomaly detected at MRI; and 9 withdrew
before MRI scan. This left data from 78 mother-infant dyads for analysis, the maternal and infant

characters for whom are shown in Table 1. Maternal hair was sampled at mean 3.5±2.5 days

after delivery, and the median HCC concentration was 5.6pg/mg (0.5-107.1). Maternal HCC was

not associated with gestational age (GA) at birth (r=0.200, p=0.094). HCC did not differ between

mothers of male and female infants (p=0.997). MRI was carried out at term-equivalent age: median

41.9 weeks' GA (range 38.6-45.9).

Table 1. Maternal and neonatal characteristics

Maternal characteristics, n = 71	
Age (years)	33.1 ± 5.2
BMI (kg/m^2)	25.2 ± 4.2
Primiparous (%) Multiparous (%)	41 (58%) 30 (42%)
SIMD 2016 quintile n (%)[†] 1 2 3 4 5	 4 (6%) 14 (20%) 10 (14%) 14 (20%) 29 (41%)
Tobacco smoked during pregnancy, n (%) No tobacco smoked during pregnancy, n (%)	5 (7%) 66 (93%)
Gestational diabetes (%) Preeclampsia (%) Receiving pharmacological treatment for depression	1 (1%) 4 (6%) 3 (4%)
Infant characteristics, n = 78	
Birthweight (grams)	2895 (454-4248)
Birth weight z-score[‡]	0.2 ± 1.1
Birth gestation (weeks)	38.4 (24.0-42.0)
Male n (%) Female	44 (56%) 34 (44%)
European ancestry n (%) Other	68 (87%) 10 (13%)
Singleton n (%) Twin	63 (81%) 15 (19%)

Normally distributed data is presented as mean ± SD. Non-normally distributed data is presented

as median (range). [†]Scottish Index of Multiple Deprivation (SIMD) 2016 quintile. First quintile

indicates most deprived, and fifth quintile the least deprived. [‡]Calculated according to

INTERGROWTH-21[st] standards

Amygdala microstructure

In univariate analysis, there were moderately strong correlations between both FA and MD, and
GA at birth and age at scan (r=0.41-0.64), and weak correlations with birth weight z-score and
Scottish 2016 quintile (r=0.24-0.30). There were no significant correlations between FA and MD in
amygdalae and ethnicity or infant sex, or maternal parity, age or BMI. There were moderate to
strong correlations between NDI in the amygdalae and GA at birth and age at scan (r=0.43-0.74),
and a weak correlation with Scottish Index of Multiple Deprivation (SIMD) 2016 quintile (r=0.23-
0.26). Weak to moderate correlations were observed between ODI in amygdalae with GA at birth
and age at scan (r=0.28-0.42), (Supplementary Table 1).

In multiple linear regression models, there was a significant interaction effect between maternal
HCC and infant sex in left amygdala FA (p=0.010) and ODI (p=0.034), with higher maternal HCC
being associated with higher left amygdala FA and lower ODI in girls compared to boys (Table 2).
When we stratified by sex, there were associations between maternal HCC and infant amygdala
microstructure in boys, but not girls. Table 3 shows that in boys, higher maternal HCC was
associated with lower left amygdala FA (β=-0.339), lower right amygdala FA (β=-0.287) and NDI
(β=-0.215), and higher right amygdala MD (β=0.264) and ODI (β=0.309), after FDR correction.

Table 2. Associations of maternal hair cortisol concentration (HCC) and its interaction with
infant sex on amygdala microstructure

Side tested	Image metric	Model 1	Model 2			Model 3		
		R^2	R^2	HCC β	HCC p-value	R^2	'HCC x sex' β	'HCC x sex' p-value
Left	FA	.267	.269	-.048	.858	.359	.677	.010
	MD	.405	.405	.018	.858	.413	-.191	.358
	ODI	.105	.111	.082	.858	.181	-.597	.034
	NDI	.530	.534	-.064	.858	.561	.375	.052
Right	FA	.289	.308	-.148	.269	.342	.415	.083
	MD	.492	.523	.189	.143	.530	-.183	.326
	ODI	.214	.228	.128	.269	.283	-.527	.083
	NDI	.554	.562	-.094	.269	.583	.330	.083

Model 1: Age at MRI, gestational age at birth, birth weight z-score, Scottish Index of Multiple

Deprivation 2016 quintile, infant sex. Model 2: Model 1 + (maternal HCC). Model 3: Model 2 +

(maternal HCC x infant sex interaction). p-values are FDR adjusted.

Abbreviations: FA, fractional anisotropy; MD, mean diffusivity; ODI, orientation dispersion index;

NDI, neurite density index; HCC, hair cortisol concentration, R^2, coefficient of determination; β,

standardised beta coefficient; p-value, FDR adjusted probability value

Table 3. Associations of maternal hair cortisol concentration (HCC) with amygdala microstructural parameters assessed separately in boys and girls

Side tested	Image metric	Boys Model 1 R^2	Boys Model 2 R^2	Boys Model 2 HCC β	Boys Model 2 HCC p-value	Girls Model 1 R^2	Girls Model 2 R^2	Girls Model 2 HCC β	Girls Model 2 HCC p-value
Left	FA	.433	.537	-.339	.023	.157	.239	.340	.372
	MD	.489	.497	.090	.462	.445	.449	-.072	.667
	ODI	.124	.215	.317	.085	.132	.159	-.194	.648
	NDI	.622	.654	-.189	.089	.522	.530	.109	.648
Right	FA	.368	.443	-.287	.047	.301	.306	.091	.736
	MD	.508	.571	.264	.047	.497	.506	.111	.736
	ODI	.149	.236	.309	.047	.362	.378	-.149	.736
	NDI	.581	.623	-.215	.047	.571	.573	.050	.736

Model 1: Gestational age at MRI, gestational age at birth, birth weight z-score, Scottish Index of Multiple Deprivation 2016 quintile. Model 2: Model 1 + (maternal HCC). p-values are FDR adjusted. Abbreviations: FA, fractional anisotropy; MD, mean diffusivity; ODI, orientation dispersion index; NDI, neurite density index; HCC, hair cortisol concentration; R^2, coefficient of determination; β, standardised beta coefficient; p-value, FDR adjusted probability value

Structural connectivity of the amygdala

For both hemispheres, the networks with the top 20% number of streamlines were connected to eight structures: thalamus, putamen, insula, superior temporal gyrus, inferior temporal gyrus, middle temporal gyrus, caudate and lateral orbitofrontal cortex, Figure 1. Quantification of streamline counts is given in Supplementary Table 2 and illustrated in Figure 2. Maternal HCC was not associated with streamline counts of the left and right amygdala with these regions.

***Figure 1. Segmentations of the amygdalae and connected regions defined by the top 20% streamline counts.** Figure 1a shows the lateral view of the sagittal plane and 1b the medial view. The same eight regions had the highest streamline counts to the amygdalae bilaterally.*

Figure 2. Chord diagram of the streamline counts between the amygdalae and unilateral regions of interest (ROIs). *The number of streamlines between ROIs are demonstrated by the corresponding arcs thickness. ROIs connected by the top 20% of streamlines are shown in green.*

In fully adjusted analyses, the interaction between maternal HCC and infant sex was significant for mean FA of connections between the right amygdala and putamen. Higher maternal HCC was associated with higher FA for amygdala-putamen connectivity in girls compared with boys (p=0.007). The interaction was also seen for connections to left thalamus, putamen and insula, but the interaction term did not remain after correction for multiple tests (Supplementary Table 3). In sex-stratified analysis, girls had higher FA values in association with high maternal HCC in connections between left amygdala with thalamus, putamen and inferior temporal gyrus, and the

193 right amygdala with putamen and inferior temporal gyrus, but these were not significant after
194 correction for multiple tests (Supplementary Table 4).
195
196 *Amygdala volume*
197
Mean volumes of the left and right amygdala were 877±111 mm^3 and 823±91 mm^3, respectively.
In univariate analysis, there were weak associations (r=0.24-0.3) between amygdala volume and
GA at birth and birth weight z-score, but not with age at scan, SIMD 2016 quintile, sex, ethnicity, or
maternal BMI, parity or age (Supplementary Table 1).   Maternal HCC was not associated with
infant right or left amygdala volume in regression models adjusted for potential covariates, and
interaction terms between maternal HCC and infant sex were not significant (Supplementary Table
5).
205
206 *Sensitivity and sub-group analyses*
207
208 There were seven twin sets in the whole sample. When we repeated analyses including only
209 singletons and the first born of twin pairs, significant associations between maternal HCC, sex and
210 image feature remained, with little change to the value of regression coefficients (Supplementary
211 Table 6).
212
In subgroup analysis of preterm and term infants, the direction and magnitude of interaction effects
for both groups were similar to those of the whole sample. Specifically, when tested in term and
preterm infants, respectively, higher maternal hair cortisol concentration is associated with higher
left amygdala fractional anisotropy (β=0.735 and 0.640), lower left amygdala orientation dispersion
index (β=-0.710 and -0.614), and higher fractional anisotropy in connections between the right
amygdala and putamen (β=0.733 and 0.426) in girls compared to boys (see Supplementary Table
6).
220
221 **Discussion**
222
We report a mechanism that could explain the impact of maternal stress on infant brain
development. We found that maternal HCC, a stable marker of chronic maternal HPA axis activity
in pregnancy, is associated with microstructure and structural connectivity of the newborn
amygdala, a region of functional importance for early social development and emotion regulation.
Specifically, HCC interacts with infant sex to modify amygdala FA, ODI and NDI, which supports
the inference that maternal chronic HPA activity has an impact on dendritic structure, axonal
configuration, and the packing density of neurites, in a sexually dimorphic manner (30-33).

The findings are consistent with recent reports from the GUSTO (Growing Up in Singapore Towards
Health Outcomes) cohort that describe associations between prenatal depression and alterations
in offspring amygdala development(10, 11). That study highlighted the role of maternal mental
health on newborn brain development, and focussed attention on the amygdala. Here, we provide
mechanistic insights into the relationship between maternal stress and amygdala development with
the use of maternal HCC to characterise chronic HPA activity, and the NODDI model for enhanced
inference about tissue microstructure. We chose to measure NODDI parameters for assessing
microstructure because ODI and NDI in grey matter appear to be functionally tractable. For
example, diffusion markers of dendritic density and arborization in grey matter predict differences
in intelligence(34), reduced ODI in grey matter is reported in psychosis and in neurodegenerative
disease, and reduced grey matter NDI is reported in Parkinson's disease, Alzheimer's disease,
autism spectrum disorder, and temporal lobe epilepsy (for review see (33)).

Maternal HCC was also related to structural connectivity of the amygdala in a sex-discordant
manner. Higher maternal HCC was associated with higher FA in girls than boys in tracts between
right amygdala and putamen. These observations were not explained by differences in streamline
counts in relation to maternal HCC. Furthermore, in sex-stratified analysis, there were consistent
trends for girls born to women with higher HCC to have higher mean FA between the left amygdala
and left thalamus, putamen and inferior temporal gyrus, and between right amygdala and right
putamen and right inferior temporal gyrus, although these did not survive FDR correction. During
the neonatal period, higher FA in white matter tracts is typically taken to imply microstructural
maturation, through increased axon diameter, density or myelination. Therefore, increased mean
FA demonstrated in connections between the amygdala and putamen, in girls exposed to higher
cortisol, could be interpreted as increased maturation of these connections.

The mechanisms underpinning differences in the relationship between maternal cortisol and
neurodevelopment of male and female infants are unknown but could occur due to sex differences
in the placental metabolism of glucocorticoids(35), regulation of glucocorticoid receptors(36) and
secretion and actions of corticotropin-releasing hormone (CRH)(37).

Strengths of this study are the use of biophysical tissue modelling (NODDI) to enable inference
about neurite density and organisation in the amygdala; and use of a data driven approach to
investigate amygdala structural connectivity. A second strength is use of maternal HCC to
operationalise stress because it is a quantitative stable marker of cortisol secretion that represents
HPA activity over 3 months; as such HCC is unlikely to reflect transient stresses that can occur in
pregnancy, and it overcomes the problems of diurnal variation that occur with plasma and saliva

measurements. To our knowledge this is the first study to investigate a physiological measure of
chronic maternal HPA activity with quantitative biomarkers of brain development, and to include
infants born very preterm. This was important because preterm birth is associated with both
exposure to maternal HPA axis dysregulation(38), and an increased risk of inattention and affective
disorders(39). The relationships we describe appear to apply across the whole GA range because
GA at birth was included in all regression models that were used to investigate association between
maternal HCC and image metrics, and in sub-group analyses the magnitude and direction of 'HCC
x sex' interaction effects were maintained between term and preterm groups. The study has some
limitations: first it was not powered to detect both sex and birth gestation interactions, but this should
be considered in future study design. Second, follow-up studies that include measures of socio-
emotional development are needed to understand functional consequences of these findings.
Finally, the newborn amygdalae are relatively small anatomical regions so could be susceptible to
partial volume effects influencing microstructural characteristics. To mitigate this risk, we used an
age-specific atlas for segmentation, and excluded voxels with a uiso <0.5.
281
In conclusion, dMRI and HCC were used to investigate mechanisms underlying the transmission
of prenatal stressors on infant development. Maternal HCC in pregnancy is associated with
newborn amygdala microstructure and structural connectivity, in a sex-dimorphic manner. These
findings reveal that the amygdala, a structure of known importance for child development, is
susceptible to variations in the prenatal stress environment, and that cortisol imparts sex specific
effects on human fetal neurodevelopment.
288
289 **Materials and Methods**
290
291 *Participants*
292
The 'Stress Response Systems in Mothers and Infants' cohort recruited mother-infant dyads from
the Royal Infirmary, Edinburgh, between March 2018 and August 2019. It prospectively tests
associations of perinatal glucocorticoid exposure with brain development, and early life exposures
including preterm birth with infant HPA axis regulation. This study recruited mother-infant dyads
with birth at ≤32 completed weeks of gestation, and dyads with birth ≥37 weeks' gestation.
Exclusion criteria were congenital fetal abnormality, chromosomal abnormality or regular maternal
corticosteroid use. All women gave written informed consent. Ethical approval was granted by
South East Scotland 01 Regional Ethics Committee (18/SS/0006).
301
302 *Maternal hair cortisol concentrations (HCC)*
303

Maternal hair was sampled within 10 days of delivery. Hair was cut close to the scalp, at the

posterior vertex, and stored in aluminium foil at -20°C. The proximal 3cm of hair were analysed by

liquid chromatography-tandem mass spectrometry (LC-MS/MS), at Dresden Lab Service GmbH

(Dresden, Germany), using an established protocol(40). Adult hair commonly grows at

1cm/month(41) and thus hair segments represented maternal HPA axis activity over the last three

months of pregnancy.

Demographic and clinical information

Participant demographic information was collected through maternal questionnaire and review of

medical records. Collected maternal information included: age at delivery (years), parity

(primiparous/multiparous), clinical diagnosis of gestational diabetes, pre-eclampsia,

pharmacological treatment for depression during pregnancy, antenatal corticosteroid exposure for

threatened preterm birth; body mass index (BMI) calculated at antenatal booking; smoking status

defined as having smoked any tobacco in pregnancy; SIMD 2016 quintile rank, a score generated

by the Scottish government which measures localities' deprivation according to local income,

employment, health, education, geographic access to services, crime and housing. Infant

demographics included whether participants were a singleton or twin, ethnicity, GA at birth (weeks),

and birth weight z-score calculated according to intergrowth standards(42).

Magnetic Resonance Imaging

Image Acquisition

Infants underwent MRI at term-equivalent age, at the Edinburgh Imaging Facility, RIE. Infants were

fed, wrapped and allowed to sleep naturally in the scanner. Flexible earplugs and neonatal earmuffs

(MiniMuffs, Natus) were used for acoustic protection. Scans were supervised by a doctor, or nurse

trained in neonatal resuscitation.

A Siemens MAGNETOM Prisma 3 T MRI clinical scanner (Siemens Healthcare Erlangen,

Germany) and 16-channel phased-array paediatric head and neck coil were used for

acquisition(43). In brief, we acquired 3D T1-weighted MPRAGE (T1w) (acquired voxel size = 1mm

isotropic) with TI 1100 ms, TE 4.69 ms and TR 1970 ms; 3D T2-weighted SPACE (T2w) (voxel size

= 1mm isotropic) with TE 409 ms and TR 3200 ms; and axial dMRI. dMRI was acquired in two

separate acquisitions to reduce the time needed to re-acquire any data lost to motion artefact: the

first acquisition consisted of 8 baseline volumes (b = 0 s/mm^2 [b0]) and 64 volumes with b = 750

s/mm^2, the second consisted of 8 b0, 3 volumes with b = 200 s/mm^2, 6 volumes with b = 500 s/mm^2

and 64 volumes with b = 2500 s/mm^2 ; an optimal angular coverage for the sampling scheme was

applied(44). In addition, an acquisition of 3 b0 volumes with an inverse phase encoding direction

was performed. All dMRI images were acquired using single-shot spin-echo echo planar imaging

(EPI) with 2-fold simultaneous multislice and 2-fold in-plane parallel imaging acceleration and 2

mm isotropic voxels; except where stated above, all three diffusion acquisitions had the same

parameters (TR/TE 3400/78.0 ms).

Conventional images were reported by an experienced paediatric radiologist (A.J.Q.) using a

structured system(45). Images with focal parenchymal injury (defined as posthaemorrhagic

ventricular dilatation, porencephalic cyst or cystic periventricular leukomalacia) were not included

in the final sample.

Image Pre-processing

Diffusion MRI processing was performed as follows: for each subject the two dMRI acquisitions

were first concatenated and then denoised using a Marchenko-Pastur-PCA-based algorithm(46);

the eddy current, head movement and EPI geometric distortions were corrected using outlier

replacement and slice-to-volume registration(47-50); bias field inhomogeneity correction was

performed by calculating the bias field of the mean b0 volume and applying the correction to all the

volumes(51).

The T2w images were processed using the minimal processing pipeline of the developing human

connectome project (dHCP) to obtain the bias field corrected T2w, the brain masks and the different

tissue probability maps(52). The mean b0 EPI volume of each subject was co-registered to their

structural T2w volume using boundary-based registration(53).

Tissue segmentation and parcellation

The ten manually labelled subjects of the M-CRIB atlas(25) were registered to the bias field

corrected T2w using rigid, affine and symmetric normalization (SyN)(54). Next, the registered labels

of the ten atlases were merged using joint label fusion(55), resulting in a parcellation containing 84

regions of interest (ROIs).

Microstructure and volumetric assessments

Volumes were calculated from ROIs derived in the structural images. ROIs were propagated to the

diffusion native space using the previously computed transformation.

To calculate the tensor derived metric, only the first shell was used. NODDI metrics were calculated

using the recommended values for neonatal grey-matter of the parallel intrinsic diffusivity (1.25

$\mu m^2 \cdot ms^{-1}$)(56). The obtained metrics are: neurite density index (NDI), isotropic volume fraction

(uiso) and orientation dispersion index (ODI). The mean FA, MD, ODI and NDI were calculated for

the left and right amygdalae M-CRIB ROIs, after exclusion of voxels with a uiso <0.5. Voxels with

a uiso <0.5 were excluded, in order to minimise partial volume effects(57).

Network construction and analysis

Tractography was performed using constrained spherical deconvolution(CSD) and anatomically-

constrained tractography(58, 59) The required 5-tissue type file, was generated by combining the

tissue probability maps obtained from the dHCP pipeline with the subcortical structures derived

from the parcellation process. Multi-tissue response function was calculated, with a FA threshold

of 0.1. The average response functions were calculated. Then, the multi-tissue fiber orientation

distribution (FOD) was calculated(60), and global intensity normalization on the FODs images was

performed. Finally, the tractogram was created, generating 10 million streamlines, with a minimum

length of 20 mm and a maximum of 200 mm and a cut-off of 0.05 (default), using backtrack and a

dynamic seeding(61). To be able to quantitatively assess connectivity, spherical-deconvolution

informed filtering of tractograms two (SIFT2) was applied to the resulting tractograms(61). The

connectivity matrix was constructed using a robust approach, a 2-mm radial search at the end of

the streamline was performed to allow the tracts to reach the GM parcellation(62). The final

connectivity matrices were multiplied by the μ coefficient obtained during the SIFT2 process.

These connectomes gave a quantification of the SIFT2 weights (referred to as the streamline

counts), and the mean FA of connections, between both the left and right amygdala to 41 unilateral

regions of interest defined through M-CRIB parcellation. In order to focus analysis on to amygdala's

most structurally connected areas, these 82 ROIs were thresholded according to the number of

streamlines connecting them to the left or right amygdala, with the top 20% (N=16) of connections

taken forward for further analysis testing relationships with maternal HCC.

Statistical analysis

Analyses were performed using IBM SPSS Statistics Version 25 Armonk, NY: IBM Corp.

Continuous data are summarised as mean ± SD if they had a normal distribution, and median

(range) if skewed. Maternal HCC was positively skewed, and log-10 transformed for analysis. The

relationship between maternal HCC with infant characteristics was tested using independent t-test

and Pearson's correlation for categorical and continuous variables, respectively. Associations
between maternal HCC with i) left and right amygdala microstructure (FA, MD, NDI, ODI), ii)
structural connectivity (number of streamlines and mean FA of connections), iii) amygdalae
volumes were tested using multiple linear regression. In all models, image feature was the
dependent variable and maternal HCC was an independent variable. Covariates included infant
sex and clinical or demographic factors that were correlated with either left or right amygdala
microstructure or volume using Pearson's correlation. Associations with the following were tested:
GA at birth, age at scan, birth weight z-score, SIMD2016 quintile, infant ethnicity, infant sex, and
maternal parity, BMI and age. Antenatal corticosteroid treatment for threatened preterm birth was
not included as a covariate because it was given to n=36 (100%) women in the preterm group, was
highly correlated with GA at birth (r=0.958, p<0.001), so its inclusion as a covariate would have
introduced multicollinearity in regression analysis. For descriptive purposes correlations of infant
and maternal factors considered as potential covariates are described as weak if r< 0.3, moderate
if r=0.3-0.7, and strong if r>0.7.

Sex differences in the relationship between maternal HCC and newborn imaging features were
assessed by adding an interaction term between maternal HCC and infant sex in the whole group
regression model. If a significant interaction was present, sex stratified analysis was conducted
independently in boys and girls. Benjamini and Hochberg false discovery rate (FDR) correction was
used to adjust p-values for multiple testing. FDR corrections were conducted separately for
assessments of left amygdala microstructure (n=4), right amygdala microstructure (n=4), left
amygdala connectivity (n=8) and right amygdala connectivity (n=8).

One sensitivity analysis was carried out to assess whether associations between maternal HCC
and image features might be enhanced by inclusion of twins. We repeated analysis of features with
a significant 'HCC x sex' interaction in the whole sample, using only singleton pregnancies and the
first-born infant of twin pairs. One sub-group analysis of preterm (GA at birth ≤ 32 weeks) and term
infants (GA at birth ≥ 37 weeks) was carried out because the relationship between maternal HCC
and infant brain development may be gestation specific.

Materials and Data Availability

All data generated or analysed during this study are included in the manuscript and supporting files.

Acknowledgments

The work was funded by Theirworld (www.theirworld.org) and was undertaken in the MRC Centre for Reproductive Health, which is funded by MRC Centre Grant (MRC G1002033). RMR acknowledges the support of the British Heart Foundation (RE/18/5/34216). Participants were scanned in the University of Edinburgh Imaging Research MRI Facility at the Royal Infirmary of Edinburgh which was established with funding from The Wellcome Trust, Dunhill Medical Trust, Edinburgh and Lothians Research Foundation, Theirworld, The Muir Maxwell Trust and many other sources; we thank the University's imaging research staff for providing the infant scanning.

Competing interests

The authors have no competing interests to declare.

References

1. B. R. H. Van den Bergh *et al.*, Prenatal developmental origins of behavior and mental health: The influence of maternal stress in pregnancy. *Neuroscience and biobehavioral reviews* 10.1016/j.neubiorev.2017.07.003 (2017).

2. D. Scheinost *et al.*, Does prenatal stress alter the developing connectome? *Pediatr Res* **81**, 214-226 (2017).

3. V. G. Moisiadis, S. G. Matthews, Glucocorticoids and fetal programming part 1: Outcomes. *Nat Rev Endocrinol* **10**, 391-402 (2014).

4. S. Sutherland, S. M. Brunwasser, Sex Differences in Vulnerability to Prenatal Stress: a Review of the Recent Literature. *Curr Psychiatry Rep* **20**, 102 (2018).

5. E. C. Braithwaite, S. E. Murphy, P. G. Ramchandani, J. Hill, Associations between biological markers of prenatal stress and infant negative emotionality are specific to sex. *Psychoneuroendocrinology* **86**, 1-7 (2017).

6. E. C. Braithwaite *et al.*, Maternal prenatal cortisol predicts infant negative emotionality in a sex-dependent manner. *Physiology & behavior* **175**, 31-36 (2017).

7. A. M. Graham *et al.*, Maternal Cortisol Concentrations During Pregnancy and Sex-Specific Associations With Neonatal Amygdala Connectivity and Emerging Internalizing Behaviors. *Biol Psychiatry* **85**, 172-181 (2019).

8. C. Buss *et al.*, Maternal cortisol over the course of pregnancy and subsequent child amygdala and hippocampus volumes and affective problems. *Proc Natl Acad Sci U S A* **109**, E1312-1319 (2012).

9. D. J. Kim *et al.*, Prenatal Maternal Cortisol Has Sex-Specific Associations with Child Brain Network Properties. *Cereb Cortex* 10.1093/cercor/bhw303 (2016).

10.   D. J. Wen *et al.*, Influences of prenatal and postnatal maternal depression on amygdala
volume and microstructure in young children. *Transl psychiatry* **7**, e1103-e1103 (2017).

11.   A. Rifkin-Graboi *et al.*, Prenatal maternal depression associates with microstructure of
right amygdala in neonates at birth. *Biological psychiatry* **74**, 837-844 (2013).

12.   J. Posner *et al.*, Alterations in amygdala-prefrontal circuits in infants exposed to prenatal
maternal depression. *Transl Psychiatry* **6**, e935 (2016).

13.   T. Humphrey, The development of the human amygdala during early embryonic life. *J
Comp Neurol* **132**, 135-165 (1968).

14.   Q. Wang *et al.*, Distribution of the glucocorticoid receptor in the human amygdala;
changes in mood disorder patients. *Brain Struct Funct* **219**, 1615-1626 (2014).

15.   L. A. Welberg, J. R. Seckl, M. C. Holmes, Inhibition of 11beta-hydroxysteroid
dehydrogenase, the foeto-placental barrier to maternal glucocorticoids, permanently
programs amygdala GR mRNA expression and anxiety-like behaviour in the offspring.
The European journal of neuroscience **12**, 1047-1054 (2000).

16.   A. Barbazanges, P. V. Piazza, M. Le Moal, S. Maccari, Maternal glucocorticoid secretion
mediates long-term effects of prenatal stress. *J Neurosci* **16**, 3943-3949 (1996).

17.   C. M. Schumann, M. D. Bauman, D. G. Amaral, Abnormal structure or function of the
amygdala is a common component of neurodevelopmental disorders. *Neuropsychologia*
49, 745-759 (2011).

18.   C. E. Rogers *et al.*, Neonatal Amygdala Functional Connectivity at Rest in Healthy and
Preterm Infants and Early Internalizing Symptoms. *J Am Acad Child Adolesc Psychiatry*
56, 157-166 (2017).

19.   M. Kraszpulski, P. A. Dickerson, A. K. Salm, Prenatal stress affects the developmental
trajectory of the rat amygdala. *Stress* **9**, 85-95 (2006).

20.   T. A. Avino *et al.*, Neuron numbers increase in the human amygdala from birth to
adulthood, but not in autism. *Proc Natl Acad Sci U S A* **115**, 3710-3715 (2018).

21.   A. Vyas, S. Bernal, S. Chattarji, Effects of chronic stress on dendritic arborization in the
central and extended amygdala. *Brain Res* **965**, 290-294 (2003).

22.   R. Mitra, R. M. Sapolsky, Acute corticosterone treatment is sufficient to induce anxiety
and amygdaloid dendritic hypertrophy. *Proc Natl Acad Sci U S A* **105**, 5573-5578 (2008).

23.   D. Batalle, A. D. Edwards, J. O'Muircheartaigh, Annual Research Review: Not just a
small adult brain: understanding later neurodevelopment through imaging the neonatal
brain. *J Child Psychol Psychiatry* **59**, 350-371 (2018).

24.   J. P. Boardman, S. J. Counsell, Invited Review: Factors associated with atypical brain
development in preterm infants: insights from magnetic resonance imaging. *Neuropathol
Appl Neurobiol* 10.1111/nan.12589 (2019).

25.   B. Alexander *et al.*, A new neonatal cortical and subcortical brain atlas: the Melbourne
Children's Regional Infant Brain (M-CRIB) atlas. *Neuroimage* **147**, 841-851 (2017).

26.   P. Galdi *et al.*, Neonatal morphometric similarity mapping for predicting brain age and
characterizing neuroanatomic variation associated with preterm birth. *Neuroimage Clin*
25, 102195 (2020).

27.   S. M. Staufenbiel, B. W. Penninx, A. T. Spijker, B. M. Elzinga, E. F. van Rossum, Hair
cortisol, stress exposure, and mental health in humans: a systematic review.
Psychoneuroendocrinology **38**, 1220-1235 (2013).

28.   C. Kirschbaum, A. Tietze, N. Skoluda, L. Dettenborn, Hair as a retrospective calendar of
cortisol production-Increased cortisol incorporation into hair in the third trimester of
pregnancy. *Psychoneuroendocrinology* **34**, 32-37 (2009).

29.   K. L. D'Anna-Hernandez, R. G. Ross, C. L. Natvig, M. L. Laudenslager, Hair cortisol
levels as a retrospective marker of hypothalamic-pituitary axis activity throughout
pregnancy: comparison to salivary cortisol. *Physiology & behavior* **104**, 348-353 (2011).

30.   S. N. Jespersen, L. A. Leigland, A. Cornea, C. D. Kroenke, Determination of Axonal and
Dendritic Orientation Distributions Within the Developing Cerebral Cortex by Diffusion
Tensor Imaging. *IEEE Transactions on Medical Imaging* **31**, 16-32 (2012).

31.   F. Grussu *et al.*, Neurite dispersion: a new marker of multiple sclerosis spinal cord
pathology? *Annals of clinical and translational neurology* **4**, 663-679 (2017).

32.   K. Sato *et al.*, Understanding microstructure of the brain by comparison of neurite
orientation dispersion and density imaging (NODDI) with transparent mouse brain. *Acta*
radiologica open **6**, 2058460117703816 (2017).

33.   A. Nazeri, C. Schifani, J. A. E. Anderson, S. H. Ameis, A. N. Voineskos, In Vivo Imaging
of Gray Matter Microstructure in Major Psychiatric Disorders: Opportunities for Clinical
Translation. *Biol Psychiatry Cogn Neurosci Neuroimaging* 10.1016/j.bpsc.2020.03.003
(2020).

34.   E. Genç *et al.*, Diffusion markers of dendritic density and arborization in gray matter
predict differences in intelligence. *Nature Communications* **9**, 1905 (2018).

35.   C. S. Rosenfeld, Sex-Specific Placental Responses in Fetal Development. *Endocrinology*
156, 3422-3434 (2015).

36.   C. H. Bourke, C. S. Harrell, G. N. Neigh, Stress-induced sex differences: adaptations
mediated by the glucocorticoid receptor. *Horm Behav* **62** (2012).

37.   D. A. Bangasser, K. R. Wiersielis, Sex differences in stress responses: a critical role for
corticotropin-releasing factor. *Hormones (Athens)* **17**, 5-13 (2018).

38.   L. Duthie, R. M. Reynolds, Changes in the maternal hypothalamic-pituitary-adrenal axis
in pregnancy and postpartum: influences on maternal and fetal outcomes.
Neuroendocrinology **98**, 106-115 (2013).

39.   S. Johnson, Cognitive and behavioural outcomes following very preterm birth. *Seminars*
in fetal & neonatal medicine **12**, 363-373 (2007).

40.   W. Gao *et al.*, Quantitative analysis of steroid hormones in human hair using a column-
switching LC-APCI-MS/MS assay. *J Chromatogr B Analyt Technol Biomed Life Sci* **928**,
1-8 (2013).

41.   R. Wennig, Potential problems with the interpretation of hair analysis results. *Forensic Sci*
Int **107**, 5-12 (2000).

42.   J. Villar *et al.*, International standards for newborn weight, length, and head
circumference by gestational age and sex: the Newborn Cross-Sectional Study of the
INTERGROWTH-21st Project. *The Lancet* **384**, 857-868 (2014).

43.   J. P. Boardman *et al.*, Impact of preterm birth on brain development and long-term
outcome: protocol for a cohort study in Scotland. *BMJ Open* **10**, e035854 (2020).

44.   E. Caruyer, C. Lenglet, G. Sapiro, R. Deriche, Design of multishell sampling schemes
with uniform coverage in diffusion MRI. *Magn Reson Med* **69**, 1534-1540 (2013).

45.   L. J. Woodward, P. J. Anderson, N. C. Austin, K. Howard, T. E. Inder, Neonatal MRI to
predict neurodevelopmental outcomes in preterm infants. *N Engl J Med* **355**, 685-694
(2006).

46.   J. Veraart *et al.*, Denoising of diffusion MRI using random matrix theory. *Neuroimage*
142, 394-406 (2016).

47.   S. M. Smith *et al.*, Advances in functional and structural MR image analysis and
implementation as FSL. *Neuroimage* **23 Suppl 1**, S208-219 (2004).

48.   J. L. R. Andersson, S. Skare, J. Ashburner, How to correct susceptibility distortions in
spin-echo echo-planar images: application to diffusion tensor imaging. *Neuroimage* **20**,
870-888 (2003).

49.   J. L. R. Andersson, S. N. Sotiropoulos, An integrated approach to correction for off-
resonance effects and subject movement in diffusion MR imaging. *Neuroimage* **125**,
1063-1078 (2016).

50.   J. L. R. Andersson *et al.*, Towards a comprehensive framework for movement and
distortion correction of diffusion MR images: Within volume movement. *Neuroimage* **152**,
450-466 (2017).

51.   N. J. Tustison *et al.*, N4ITK: improved N3 bias correction. *IEEE Trans Med Imaging* **29**,
1310-1320 (2010).

52.   A. Makropoulos *et al.*, The developing human connectome project: A minimal processing
pipeline for neonatal cortical surface reconstruction. *Neuroimage* **173**, 88-112 (2018).

53.   D. N. Greve, B. Fischl, Accurate and robust brain image alignment using boundary-based
registration. *Neuroimage* **48**, 63-72 (2009).

54.   B. B. Avants, C. L. Epstein, M. Grossman, J. C. Gee, Symmetric diffeomorphic image
598   registration with cross-correlation: evaluating automated labeling of elderly and
599   neurodegenerative brain. *Med Image Anal* **12**, 26-41 (2008).

600   55.   H. Wang *et al.*, Multi-Atlas Segmentation with Joint Label Fusion. *IEEE Trans Pattern*
601   *Anal Mach Intell* **35**, 611-623 (2013).

602   56.   J. M. Guerrero *et al.*, Optimizing the intrinsic parallel diffusivity in NODDI: An extensive
603   empirical evaluation. *PLoS One* **14**, e0217118 (2019).

604   57.   D. Batalle *et al.*, Different patterns of cortical maturation before and after 38 weeks
605   gestational age demonstrated by diffusion MRI in vivo. *NeuroImage* **185**, 764-775 (2019).

606   58.   J. D. Tournier *et al.*, MRtrix3: A fast, flexible and open software framework for medical
607   image processing and visualisation. *Neuroimage* **202**, 116137 (2019).

608   59.   R. E. Smith, J. D. Tournier, F. Calamante, A. Connelly, Anatomically-constrained
609   tractography: improved diffusion MRI streamlines tractography through effective use of
610   anatomical information. *Neuroimage* **62**, 1924-1938 (2012).

611   60.   B. Jeurissen, J. D. Tournier, T. Dhollander, A. Connelly, J. Sijbers, Multi-tissue
612   constrained spherical deconvolution for improved analysis of multi-shell diffusion MRI
613   data. *Neuroimage* **103**, 411-426 (2014).

614   61.   R. E. Smith, J. D. Tournier, F. Calamante, A. Connelly, SIFT2: Enabling dense
615   quantitative assessment of brain white matter connectivity using streamlines
616   tractography. *Neuroimage* **119**, 338-351 (2015).

617   62.   R. E. Smith, J. D. Tournier, F. Calamante, A. Connelly, The effects of SIFT on the
618   reproducibility and biological accuracy of the structural connectome. *Neuroimage* **104**,
619   253-265 (2015).