Ventral hippocampus-lateral septum circuitry mediates foraging-related spatial memory

Elizabeth A. Davis[1*], Clarissa M. Liu[1,2*], Léa Décarie-Spain[1], Logan Tierno Lauer[1], Isabella H. Gianatiempo[1], Andrea N. Suarez[1], Alyssa M. Cortella[1], Joel D. Hahn[3], and Scott E. Kanoski[1,2]

*These two authors contributed equally

[1]Human and Evolutionary Biology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089; [2]Neuroscience Graduate Program, University of Southern California, 3641Watt Way, Los Angeles, CA, USA. 90089; [3]Neurobiology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089

Corresponding author:
Scott E. Kanoski, PhD
University of Southern California
3616 Trousdale Parkway, AHF-252
Los Angeles, CA 90089-0372
Phone: 213-821-5762
Email: kanoski@usc.edu

ABSTRACT

Remembering the location of a food or water source is essential for survival. Here we identify a hippocampal-septal neural circuit that is necessary for spatial memory based specifically on appetitive reinforcement. Both reversible and chronic disconnection of ventral hippocampus CA1 subregion (CA1v) projections to the lateral-septum (LS) using pathway-specific dual viral approaches impaired spatial memory retention for the location of either food or water reinforcement. However, disconnection of this pathway did not affect performance in an aversive escape-motivated spatial memory task that used the same apparatus and visuospatial cues. The selectivity of this pathway in mediating foraging-related memory was further supported by results showing that CA1v-LS disconnection did not affect anxiety-like behavior, locomotor activity, or social and olfactory-based appetitive learning. To examine whether CA1v-LS mediation of foraging-related spatial memory involves collateral projections of CA1v neurons, we utilized virus-based neural pathway tracing analyses to identify the medial prefrontal cortex (mPFC) as a collateral target of LS-projecting CA1v neurons. However, functional disconnection of CA1v and mPFC did not affect appetitive spatial memory, thus further supporting the selectivity of CA1v-LS signaling for this behavior. The lateral hypothalamic area, which is associated with food-motivated behavior and more recently with spatial memory, was identified as a second-order target of CA1v-LS signaling using a multisynaptic anterograde tracing approach. Collective results reveal that CA1v to LS communication plays a critical role in remembering an environmental location based on appetitive (food or water) but not aversive (escape) reinforcement, thus identifying a novel neural pathway regulating foraging-related memory processes.

INTRODUCTION

A survival advantage common to early humans and lower-order mammals is to accurately remember the location of a food or water source in the environment, and to then efficiently navigate back to a shelter or other safe location. The neurobiological substrates that regulate visuospatial navigation are therefore critical to effectively implement food-directed or water-directed foraging behavior. In rodents a number of maze tasks have been developed that have been instrumental in identifying brain structures within the telencephalon that are essential for visuospatial mapping and egocentric orientation in the environment, including the hippocampus and the medial entorhinal cortex, respectively (1, 2). However, despite that reliably locating food and water sources in the environment is a key selection pressure driving the evolution of visuospatial navigation, the overwhelming majority of rodent model research on the neural substrates of spatial learning and memory have utilized procedures such as the Morris water maze and the Barnes maze that each involve escaping aversive stimulus conditions (3, 4). Furthermore, while single unit recordings have been used to identify specific populations of neurons that subserve distinct navigational functions (e.g., hippocampal "place cells", medial entorhinal "grid cells") (5, 6), the bulk of this research has recorded neural activity under neutral conditions void of either appetitive or aversive reinforcement. Very little research has been dedicated to identify brain regions and neural circuits that may specifically promote spatial memory based on the location of appetitive reinforcers (e.g., food, water), as well as the extent to which the nature of the reinforcement is a critical factor in deciphering the brain's control of visuospatial navigation.

The majority of rodent model research investigating brain regions that mediate visuospatial navigational memory has focused on the anterior and "dorsal" subregion of the hippocampus (septal pole; HPCd). However, the posterior and "ventral" hippocampus subregion (temporal pole; HPCv), while classically associated with stress- and affect-associated memory processes (7), also plays a role in visuospatial learning and memory (8-10). For example, under some testing conditions selective HPCv lesions impair spatial memory performance in the Morris water maze (10, 11). Moreover, place cells that are responsive to changes in the visuospatial environment are present within both the HPCd and HPCv pyramidal layers, with a linear increase in the spatial scale of representation from the dorsal to the ventral pole (9). Despite a

common role for the HPCd and HPCv in mediating spatial memory, there is also evidence for a
functional distinction between the subregions (7, 12, 13). For instance, lesions of the HPCv but
not the HPCd alter stress responses (14) and anxiety-like behavior (15), whereas HPCd but not
HPCv lesions impair spatial memory in an incidental (nonreinforced) procedure (16). These two
HPC subregions also have disparate neuroanatomical connectivity supporting a framework for a
functional diversity in which the HPCd preferentially processes cortical-derived sensory
information and the HPCv preferentially processes metabolic and limbic-derived affective
information (7, 13). The functional distinction between these two subregions is further supported
by the generation of the hippocampus gene expression atlas, which provides a comprehensive
integration of gene expression and connectivity across the HPC septo-temporal axis (17). Given
that both the HPCv and the HPCd participate in spatial memory but have distinct
neuroanatomical connectivity and contributions when it comes to regulating other cognitive and
mnemonic processes, it is feasible that the HPCv and HPCd support different forms of spatial
memory depending on the type of reinforcement and/or the context associated with the behavior.
Recent findings identify the HPCv as a critical brain substrate in regulating feeding
behavior and food-directed memory processes. Reversible inactivation of HPCv neurons after a
meal increases the size of and reduces the latency to consume a subsequent meal (18, 19). In
addition, receptors for feeding-related hormones are more abundantly expressed in the HPCv
compared to the HPCd (e.g., ghrelin receptors (20, 21), glucagon-like peptide-1 [GLP-1]
receptors (22)), and these HPCv endocrine and neuropeptide receptor systems alter food intake
and feeding-related memory (23-28). Olfactory information, which is intimately connected with
feeding behavior, is also preferentially processed within the HPCv compared with the HPCd.
The CA1v, specifically, is bidirectionally connected to brain regions that process olfactory
information (7, 29, 30), and CA1v neurons respond more robustly to olfactory contextual cues
compared with CA1d (8). Given these HPCv appetite-relevant neuroanatomical connections,
endocrine and neuropeptide receptor expression profiles, and functional evidence linking this
subregion with food-motivated behaviors, we hypothesize that HPCv mediation of visuospatial
memory is preferentially biased to food and water-reinforced foraging behavior.
HPCv pyramidal neurons have extensive projection targets throughout the brain (31), yet
the functional relevance of these pathways is poorly understood. The lateral septum, in
particular, is a robust target of HPCv glutamatergic pyramidal neurons (32, 33). Given that this

pathway has been shown to affect feeding behavior (34) and that neuroplasticity changes occur

in the LS after learning a spatial memory task (35, 36), we hypothesize that that HPCv to LS

projections participate in regulating foraging-related spatial memory for food and water location.

To investigate this hypothesis, we developed two novel appetitive reinforcement-based spatial

foraging behavioral tasks that allow for direct comparison with an aversive reinforcement-based

task of similar difficulty that uses the same apparatus and spatial cues. Performance in these

tasks was assessed following pathway-specific dual viral-based reversible (chemogenetic

inhibition) or chronic (targeted apoptosis) disconnection of the CA1v to LS pathway. To further

expand neural network knowledge on CA1v to LS signaling, we used conditional viral-based

neuroanatomical tracing strategies to identify both first-order collateral and second-order targets

of LS-projecting CA1v neurons. Collective results from the present study identify novel neural

circuitry of specific relevance to foraging behavior.

RESULTS

The ventral hippocampus (HPCv) is required for remembering the spatial location of food

To examine the importance of the HPCv in memory for the spatial location of food, animals received bilateral *N*-Methyl-D-aspartate excitotoxic lesions of the HPCv (n=12) or bilateral sham injections (control n=12) (histological analyses for the neuron-specific NeuN antigen in Fig. 1A) and were tested in a novel appetitive visuospatial memory task developed in our lab that utilizes a Barnes Maze apparatus. Results revealed no significant differences in errors (incorrect hole investigations) (Fig. 1E) or latency to locate the food source hole (Fig. 1F) during training. However, memory probe results show that animals with HPCv lesions decreased the ratio of correct + adjacent / total holes explored in the first minute compared with controls (Figure 1G; $p < 0.05$, cohen's d=-1.0174; analysis for the combined 2-min period did not yield a statistical group difference), indicating impaired memory for the spatial location of food reinforcement.

Post-surgical analyses of food intake (Supp. Fig. 1A) and body weight (Supp. Fig. 1B) found no significant group differences in these measures over time. These collective results indicate the HPCv is required for memory retention but not learning of the spatial location of food in the environment.

Ventral hippocampus CA1 (CA1v) projections to the lateral septum (LS) mediate appetitive spatial memory for food and water location, but not aversive spatial memory for escape location

To identify a specific neural pathway mediating the results described above, we investigated the functional relevance of the CA1v to LS signaling in learning and recalling the spatial location of either food or water reinforcement using conditional dual viral approaches to either reversibly (via cre-dependent pathway-specific inhibitory chemogenetics; diagram of approach in Fig. 2A) or permanently (via cre-dependent pathway-specific caspase-induced lesions; diagram of approach in Fig. 2B) disconnect CA1v to LS communication. Results from the appetitive food seeking spatial memory task reveal no significant group differences during training in either errors before locating the correct hole (Fig. 2E) or latency to locate the food source hole (Fig. 2F). However, memory probe results demonstrate that both acute (pathway-specific DREADDs) and chronic (pathway-specific caspase) CA1v to LS disconnection decreased the ratio of correct

+ adjacent / total holes explored in the entire two minutes of the probe compared with controls

(Fig. 2G; $p<0.05$, cohen's d=-1.142; analysis for the 1st minute period alone did not yield a

statistical group difference, $p=0.059$), indicating impaired memory for the spatial location of

food reinforcement.

Results from an analogous appetitive water seeking spatial memory task in a separate cohort of

water-restricted rats also reveal no significant group differences during training in either errors

before locating the correct hole (Fig. 2H) or latency to locate the food source hole (Fig. 2I).

However, a significant impairment was observed following either acute or chronic CA1v to LS

disconnection of the ratio of correct + adjacent / total holes explored in 2-min memory probe

compared with controls (Fig 2J; $p<0.05$, cohen's d=-1.18; analysis for the 1st minute period alone

did not yield a statistical group difference), indicating impaired memory for the spatial location

of water reinforcement.

Histological analyses confirmed successful viral transfection of LS-projecting CA1v

neurons with DREADDs (Fig. 2C-D), as well as successful lesioning of LS-projecting CA1v

neurons for the caspace approach (Supplemental Fig. 2H-I. The reversible DREADDs-based

disconnection approach was further confirmed by c-Fos analyses in the LS, where the number of

c-Fos immunoreactive neurons in the LS in response to the food reinforcement-based spatial

memory probe test was reduced following CNO treatment relative to vehicle/control treatment

(Supp. Fig. 2E-G). Post-surgical analyses of food intake (Supp. Fig 2A) and body weight (Supp.

Fig 2B) found no group differences in these measures over time (in the absence of CNO

treatment). These data demonstrate that the CA1v-LS disconnection approaches were successful,

that CA1v to LS communication is critical for remembering the spatial location of food and

water, and that these effects are unlikely to be based on differences in energy status or food

motivation.

To evaluate whether CA1v to LS signaling is specifically involved in appetitive

reinforcement-based spatial memory (food or water) vs. spatial memory in general, in a separate

cohort of animals we tested the effect of reversible and chronic CA1v to LS disconnection in a

spatial learning task based on aversive reinforcement rather than food. Importantly, this task uses

the same apparatus and visuospatial cues as the spatial location food and water memory tasks

described above, but the animals are motivated to locate the tunnel to escape mildly aversive

stimuli (bright lights, loud noise) with no food or water reinforcement in the tunnel (see Fig. 1B).

Training results revealed no significant group differences in errors before correct hole (Fig. 2K)

nor latency to locate the escape hole (Fig. 2L). During the memory probe test, there were no

group differences in the ratio of correct + adjacent / total holes investigated during the first

minute nor the entire two minutes (Fig. 2M) of the memory probe. Post-surgical analyses of food

intake (Supp. Fig. 2C) and body weight (Supp. Fig. 2D) found no group differences in these

measures over time. These collective findings suggest that CA1v to LS signaling specifically

mediates spatial memory in a reinforcement-dependent manner, playing a role in appetitive (food

or water reinforcement) but not aversive (escape-based) spatial memory.

Reversible, but not chronic disconnection of CA1v to LS signaling increases chow intake

under free-feeding conditions.

179   While the results above suggest that the impairments in food or water-based spatial

memory were not based on long-term changes in food intake or body weight, these experiments

were conducted under conditions of either chronic food or water restriction. To examine the role

of CA1v to LS signaling on feeding behavior under free-feeding conditions, we examined meal

pattern feeding analyses in the caspase vs. the control group (chronic CA1v-LS disconnection),

and in the DREADDs group following CNO or vehicle (within-subjects design; reversible

CA1v-LS disconnection). Results averaged over 5d revealed no significant effect of chronic

CA1v to LS disconnection (caspase group) on 24h meal frequency, meal size, meal duration, or

24h food intake in comparison to controls (Supp. Figs. 3A-D). However, CNO-induced acute

reversible silencing of LS-projecting CA1v neurons (DREADDs group) significantly increased

24h chow intake (Supp. Fig. 3H; $p<0.05$, cohen's d=1.0) relative to vehicle treatment,

accompanied by a trend towards increased meal frequency (Supp. Fig. 3E; $p=0.11$) and no effect

on meal size or meal duration (Supp. Figs. 3F&G). These results are unlikely based on

nonselective effects of the CNO, as we've recently shown that ICV CNO injections do not affect

food intake in rats when tested under similar conditions (37). Collectively these findings show

that reversible CA1v to LS disconnection increases food intake under free-feeding conditions,

whereas food intake is not affected by chronic CA1v to LS ablation.

CA1v to LS disconnection does not affect nonspatial HPC-dependent appetitive memory,

anxiety-like behavior, or levels of locomotor activity

The HPCv plays an important role in the social transmission of food preference

procedure (STFP) (28, 38, 39), which tests socially-mediated food-related memory based on

previous exposure to socially-communicated olfactory cues (diagram in Fig. 3A). However,

results revealed that neither chronic nor acute disconnection of CA1v to LS signaling affected

the STFP preference ratio between groups (Fig. 3B), with all three groups performing

significantly above chance (based on within-group one sample t-tests, ps<0.05).

HPCv to LS circuitry has been shown to play a role in mediating anxiety and stress

behavior (40-42). Given that altered anxiety-like behavior and/or locomotion could produce

behavioral changes that may lead to an apparent spatial memory deficit in any of the spatial

learning tasks used, we examined whether HPCv to LS disconnection yields differences in

anxiety-like behavior (zero maze task; diagram of apparatus in Fig. 3C) and/or locomotion (open

field test). Zero maze results showed no significant group differences in time spent in the open

zones (Fig. 3D) nor in the number of open zone entries (Fig. 3E). In addition, CA1v to LS

disconnection had no effect on general locomotor activity in the open field test (Fig. 3F). These

collective results suggest that the observed deficits in food- and water-seeking spatial memory

are not secondary to impaired nonspatial appetitive learning, general differences in anxiety-like

behavior, or locomotor impairments.

217   The medial prefrontal cortex is a collateral target of LS-projecting CA1v neurons

In addition to the LS, CA1v neurons also robustly project to the medial prefrontal cortex

(mPFC), a pathway that we have previously shown to be involved in feeding behavior (25). To

examine whether LS-projecting CA1v neurons have collateral targets in the mPFC, or in other

brain regions, we utilized a conditional dual viral neural pathway tracing approach that identifies

collateral targets of a specific monosynaptic pathway (CA1v->LS; diagram of approach in Fig.

4A, representative CA1v injection site in Fig. 4B, representative CA1v injection site in Fig. 4C).

Results revealed that LS-projecting CA1v neurons also project to the mPFC (Fig. 4D), whereas

very minimal labeling was observed in other brain regions throughout the neuraxis. Thus, it may

be the case that the impaired spatial memory for food location observed following either

reversible or chronic CA1v to LS disconnection are based, in part, on CA1v to mPFC signaling

from the same LS-projecting CA1v neurons.

CA1v to mPFC signaling does not contribute to appetitive spatial memory

We next sought to test the functional relevance to food-directed spatial memory of a CA1v-originating neural pathway (CA1v to mPFC) that is a collateral target of LS-projecting CA1v neurons. The CA1v to mPFC projections are exclusively ipsilateral (25, 43), and thus this pathway allows for a 'contralesional' approach to achieve complete and chronic functional disconnection (diagram of approach in Fig. 4E&4I, modified from (43)). This approach was utilized in favor of the dual viral approaches used for the CA1v-LS disconnection, as it provides a more complete disconnection between the structures and does not require the use of viruses and involves fewer surgical injections to achieve a similar objective. We note, however, that this contralesional approach is not a feasible option for the CA1v-LS pathway, as CA1v neurons project to the LS bilaterally (e.g., Fig. 5F). Post-surgical analyses of food intake (Supp. Fig. 4A) and body weight (Supp. Fig. 4B) found no group differences in these measures over time between sham controls and the chronic CA1v-mPFC disconnection group. Training results from the appetitive spatial memory task revealed no group differences in errors before locating the correct hole (Fig. 4F or latency to locate the food-baited hole (Fig. 4G). Importantly, memory probe results demonstrate that CA1v to mPFC disconnection did not alter the ratio of correct + adjacent / total holes explored in the first minute or the entire two minutes (Fig. 4H, 4L) compared with controls. These data collectively demonstrate that CA1v to LS mediation of foraging-related memory does not require collateral projections to the mPFC neural pathway.

LS neurons activated during appetitive spatial memory testing are primarily GABAergic

Fluorescent *in situ* hybridization analyses were used to determine the neurotransmitter phenotype cFos+ LS neurons that were activated during the appetitive food seeking spatial memory probe test. Results revealed that ~10% of c-Fos+ LS neurons express ChAT (a cholinergic marker; Fig. 5A), ~88% of cFOS+ LS neurons express GAD1 (a GABAergic; Fig. 5B), and only ~2% of cFOS+ LS neurons express VGLUT2 (a glutamatergic marker; Fig. 5C). These results suggest that LS neurons engaged during the spatial memory probe are primarily GABAergic (Fig. 5D).

The lateral hypothalamic area (LHA) is a primary second-order target of LS neurons that receive CA1v input

A dual viral tracing strategy (diagram of approach in Fig. 6A; representative LS injection

site in Fig. 6B) was utilized to identify downstream targets of LS-projecting CA1v neurons.

Results revealed that the LHA is the strongest second-order target of the CA1v to LS-projecting

neurons (Fig. 6C-D). Quantitative analyses using a custom-built data-entry platform (Axiome C,

created by JDH) are summarized graphically for a representative animal on a brain flatmap

summary diagram (Fig. 6D) for the hemisphere ipsilateral (top) and contralateral (bottom) to the

injection sites (see Supplementary Table 1 for data from complete forebrain analyses).

Retrograde transfection of the AAVs from the injection sites was not observed, as tdTomato+

cell bodies were only observed in the LS following complete forebrain histological analyses.

Additional confirmation of the LHA as a second-order target was provided via a co-injection

(COIN) strategy, from (44, 45). This approach involved injections of a monosynaptic retrograde

tracer, fluorogold (FG), in the LHA (representative injection site in Fig. 6E) and injection of a

monosynaptic anterograde tracer, AAV1-GFP, in the CA1v (representative injection site in Fig.

6F), which resulted in the observance of FG-GFP appositions in the LS to further confirm this

multisynaptic pathway (Fig. 6G).

DISCUSSION
279

Memory for the physical location of a food or water source is adaptive for maintaining

adequate energy supply for reproduction and survival. However, the neural circuits mediating

this complex behavior are not well understood as research on visuospatial memory has

predominantly used tasks with either aversive or neutral/passive reinforcement. Moreover,

whether the neural circuitry mediating spatial memory function is divergent based on the nature

(e.g., appetitive vs. aversive) or modality (e.g., food vs. water) of the reinforcement has not been

systematically investigated. The present study identified a monosynaptic CA1v to LS pathway as

a necessary substrate for food- and water-motivated spatial memory. Moreover, the selectivity of

the CA1v-LS pathway in mediating spatial memory for food and water location is supported by

results showing that disconnection of this pathway did not affect performance in an escape-

motivated spatial memory task of comparable difficulty conducted in the same apparatus,

anxiety-like behavior, locomotor activity, or olfactory and social-based appetitive memory.

Fluorescent in situ hybridization analyses also demonstrate that LS neurons engaged during

appetitive spatial memory testing are primarily GABAergic. Viral-based pathway tracing revealed that LS-projecting CA1v neurons also send collateral projections to the mPFC, however, functional disconnection of CA1v-mPFC signaling did not impair spatial memory for food location. Utilization of an anterograde multisynaptic viral tracing approach and quantitative forebrain-wide axon mapping analyses revealed that the CA1v-LS pathway sends second-order projections to various feeding and reward-relevant neural substrates, particularly the LHA. Collectively, these data establish monosynaptic CA1v to LS signaling as an essential pathway for spatial memory-based foraging behavior, and we further identify a neural network of interconnected feeding-relevant substrates that are collateral and/or downstream targets of this pathway.

Historically, the hippocampus has been divided into the dorsal and ventral subregions that have both distinct and overlapping functions, anatomical connections, and genetic and receptor expression profiles (12, 13, 17, 31, 46). The HPCd has been predominantly implicated in facilitating spatial memory, whereas the HPCv is linked more with stress responses, affective behavior, and energy balance. However, a recent study found that both the HPCd and HPCv were critical for food reward-directed spatial navigation in an obstacle-rich complex environment (47), and under certain testing conditions, spatial learning and memory in an aversive reinforcement-based water maze also requires both subregions of the HPC (48). Here we expand this literature by identifying a specific HPCv-originating pathway (CA1v->LS) that selectively mediates spatial memory for the location of appetitive (food or water) reinforcement. These novel findings suggest that the modality of the reinforcement should be considered when investigating the longitudinally-organized processing of spatial information across the dorsal-ventral HPC axis.

While HPCv neurons project widely throughout the neuraxis, projections to the LS are particularly robust (33). The LS has been implicated in food anticipatory/seeking behavior, where metabolic activation of the lateral septum peaks immediately before feeding (during food anticipatory activity) in rabbit pups (49) and gamma-rhythmic input to the LHA from the LS evokes food approach behavior (50). In addition, the LS can bidirectionally modulate food intake via a number of feeding-relevant neuropeptides, including glucagon-like peptide-1, ghrelin, and urocortin (51-53). Recent findings demonstrate that DREADDs-mediated disconnection of a HPCv to LS pathway (from CA3v and DGv) increases food intake in mice, with

opposite/hyperphagic effects observed following activation of this pathway (34). Similarly, here

we demonstrate that DREADDs-mediated acute silencing of the CA1v to LS pathway increases

food intake in rats. Together these findings support a role for HPCv-LS signaling in regulating

eating behavior. However, these results are based on analyses conducted in the home cage where

animals have free and unlimited access to food and water. Here we extend these findings by

establishing a role of CA1v to LS signaling in foraging-relevant memory processes under

conditions that are more relevant to normal mammalian behavior, where food and water are not

freely available.

The use of two different but complementary approaches for neural pathway-specific

disconnection allowed us to test at the levels of learning and memory separately. For example,

the use of a reversible (DREADDs) disconnection procedure allowed us to evaluate the effect of

CA1v to LS disconnection during the memory probe only, with circuit intact during training

(CNO was not administered at any point during training). In contrast, the chronic (caspase)

disconnection procedure was performed before training and therefore could have affected

learning if CA1v to LS pathways were required for the learning of the task. However, our results

showed that CA1v to LS disconnection did not impair learning, but rather, only memory

retention performance in the probe test conducted days after the end of training. Thus, we can

now confirm based on two complementary levels of analysis that this pathway is involved in

longer-term memory recall but not learning/acquisition, which is consistent with recent evidence

demonstrating that complete HPCv lesions impair retrieval (probe performance), but not

acquisition (learning) in a Morris water maze (54). A limitation of the present study is that we

did not evaluate 'gain of function' of CA1v->LS signaling on appetitive spatial memory.

However, we note that such gain of function approaches are more commonly utilized with

models of memory dysfunction. Thus, an important area for follow-up research is to investigate

whether enhanced CA1v->LS signaling (e.g., via excitatory pathway-specific DREADDs) can

augment memory function in models where foraging-related memory is impaired.

While the use of aversive reinforcement-based spatial memory tasks are predominant in

the field, the radial arm maze has also been used to investigate food-motivated spatial memory

(55). Common applications include win-shift tasks, where animals learn to avoid previously

visited food-baited arms (working memory, modeling spontaneous alternation and exploration in

environments with minimal resources), and win-stay tasks where animals learn to return to

previously reinforced arms modeling foraging in environments of plentiful resources (reference memory). Our experiments are analogous to the latter. However, an advantage of the present food-reinforced Barnes Maze apparatus-based approach over the radial arm maze procedure is that we've developed parallel procedures that use the same apparatus and visuospatial cues to examine water- and escape-motivated spatial memory, thus offering a unique arsenal to assess the selectivity of the circuit based on multiple reinforcement modalities (food vs. water vs. escape/aversive). Future research is needed to determine whether CA1v to LS signaling also mediates spatial working memory based on appetitive and/or aversive reinforcement.

Utilizing dual viral approaches and systematic forebrain-wide quantitative mapping analyses, our findings confirm that the CA1v-LS pathway projects downstream to other neural substrates involved in feeding and motivated behavior, including the LHA, ACB, and mPFC. This experiment was performed to generate novel neural connectivity data for the field and to demonstrate possible downstream 2nd-order targets that may inspire future research focused on these newly-identified multi-order neural circuits. We identified the LHA as the most robust downstream 2nd-order target of CA1v->LS signaling, and this multisynaptic pathway was corroborated by an additional monosynaptic tracing approach that combined retrograde tracing (FG in LHA) and anterograde tracing (AAV1-GFP in CA1v) to confirm a connection relay in the LS. While future research is required to test the functional relevance of this multisynaptic pathway to appetitive foraging-relevant memory, we hypothesize that the LHA is a functional downstream target based on recent findings linking LHA neurons with spatial memory (56).

Collective results identify a CA1v to LS pathway involved in food- and water-motivated spatial memory, but not escape-motivated spatial memory. Furthermore, the selectivity of this pathway to appetitive spatial memory is supported by data showing that neither chronic nor reversible disruption of CA1v to LS signaling influenced various other behavioral outcomes, including anxiety-like behavior, locomotor activity, and nonspatial HPC-dependent appetitive memory. We also systematically characterized collateral and second-order projections of this pathway. Present results demonstrate overlapping pathways coordinated in the higher-order control of energy balance, of which foraging behavior is an essential component given that foraging requires energy expenditure to locate food for subsequent energy intake. These data shed light on the neural systems underlying complex learned and motivated behaviors that require functional connections between cognitive and feeding-relevant substrates.

MATERIALS AND METHODS

Animals

Adult male Sprague–Dawley rats (Envigo; 250-275g on arrival) were individually housed in hanging wire cages with *ad libitum* access (except where noted) to water and chow (LabDiet 5001, LabDiet, St. Louis, MO) on a 12h:12h reverse light/dark cycle. All procedures were approved by the University of Southern California Institute of Animal Care and Use Committee.

Acute and chronic disconnection of the CA1v to LS neural pathway

Cre-dependent dual viral strategies were used to generate the following groups: [1] acute chemogenetic disconnection of the CA1v to LS neural pathway (DREADDs; diagram of approach in Fig. 2A), [2] chronic disconnection of the CA1v to LS neural pathway (caspase lesion-induced; diagram of approach in Fig. 2B), and [3] a common/shared control group for these two disconnection strategies. Regardless of group, all animals received a bilateral AAV retro-cre injection in the LS (AAV2[retro]-hSYN1-EGFP-2A- iCre-WPRE; 200nL per side) at the following stereotaxic coordinates (57): +0.84mm AP, +/-0.5mm ML, −4.8mm DV, all defined at bregma.

According to experimental group, animals received a different virus delivered to the CA1v subregion of the HPCv. All viruses were administered to the CA1v at the following stereotaxic coordinates (57): −4.9mm AP defined at bregma, +/-4.8mm ML defined at bregma, −7.8mm DV defined at skull surface at target site.

[1] DREADDs group for reversible inactivation of CA1v to LS: To allow reversible chemogenetic inactivation of the CA1v to LS neural pathway, one group of animals received a bilateral CA1v injection of a cre-dependent virus to drive expression of inhibitory designer receptors activated by designer drugs (DREADDs), (AAV-Flex-hm4Di-mCherry). This dual viral strategy drives expression of inhibitory DREADDs exclusively in LS-projecting CA1v neurons, which enables acute inactivation of these neurons by injection of the DREADDs ligand, clozapine-N-oxide (CNO), at the time of behavioral testing.

[2] Caspase group for chronic inactivation of CA1v to LS: To allow chronic disconnection of the CA1v to LS neural pathway, a second group of animals received a bilateral CA1v cre-dependent caspase virus (AAV1-Flex-taCasp3-TEVp; 200nL per side) mixed with a

cre-dependent reporter virus (AAV-flex-tdTomato; 200nL per side) for histological verification. This dual viral strategy drives expression of the apoptosis-mediator molecule caspase exclusively in LS-projecting CA1v neurons, which induces apoptotic cell death in these neurons while leaving other CA1v neurons intact.

    [3] Common control: A common control group was generated so that the DREADDs and caspase groups could both be compared to the same control group, instead of two different control groups for the DREADDs and caspase groups, respectively. This allowed us to reduce the number of animals needed for the same experimental objective. Thus, all animals received an indwelling cannula and ICV injections of CNO as described below.

    Immediately following viral injections, all animals were surgically implanted with a unilateral indwelling intracerebroventricular (ICV) cannula (26-gauge, Plastics One, Roanoke, VA) targeting the lateral ventricle (VL). Cannulae were implanted and affixed to the skull with jeweler's screws and dental cement at the following stereotaxic coordinates: −0.9mm AP defined at bregma, +1.8mm ML defined at bregma, −2.6mm DV defined at skull surface at site. Placement for the VL cannula was verified by elevation of blood glucose resulting from an injection of 210μg (2μL) of 5-thio-D-glucose (5TG) using an injector that extended 2mm beyond the end of the guide cannula (58). A post-injection elevation of at least 100% of baseline glycemia was required for subject inclusion. Animals that did not pass the 5TG test were retested with an injector that extended 2.5mm beyond the end of the guide cannula and, upon passing the 5TG test, were subsequently injected using a 2.5mm injector instead of a 2mm injector for the remainder of the study. Prior to behavioral testing where noted, all animals received an ICV 18mmol infusion (2uL total volume) of the DREADDs ligand clozapine N-oxide (CNO) (59), rendering only DREADDs animals chemogenetically inactivated. CNO injections were hand delivered using a 33-gauge microsyringe injector attached to a PE20 catheter and Hamilton syringe through the indwelling guide cannulae. Injectors were left in place for 30sec to allow for complete delivery of the CNO.

    Cre-dependent DREADD expression targeting LS-projecting neurons in CA1v was evaluated based on localization of the fluorescent reporter mCherry. Immunohistochemistry for red fluorescent protein (RFP) was conducted to amplify the mCherry signal (see Immunohistochemistry). Only animals with mCherry expression restricted within CA1v neurons were included in subsequent behavioral analyses. Representative images in Figs. 2C&D.

The general approach for neuronal apoptosis due to activation of cre-dependent caspase
targeting LS-projecting neurons in CA1v was validated in a separate cohort (n=8) based on
reduction of a cre-dependent fluorescent reporter tdTomato due to neuron cell death compared
with controls. The caspase group brains (which received a retro-cre virus in the LS in
conjunction with an injection of cre-dependent caspase virus mixed with a cre-dependent virus
that drives a tdTomato fluorescent reporter in the CA1v) were compared to control brains (that
received a retro-cre virus in the LS in conjunction with only a cre-dependent that drives a
tdTomato fluorescent reporter in the CA1v, which was equivalently diluted to match the caspase
injections). In both groups, immunohistochemistry for red fluorescent protein (RFP) was
conducted to amplify the tdTomato signal (see Immunohistochemistry). Confirmation of
successful caspase-driven lesions in the CA1v using this approach was evaluated by reduced
tdTomato fluorescence in comparison to control animals, with the expression of CA1v tdTomato
expression in the caspase group less than 15% of that observed in controls, on average
(Representative images in Supplementary Fig.2 H-I). While this cohort allowed for successful
validation of the pathway-specific disconnection caspase approach, to improve the viability of
the animals for long-term behavioral analyses, the 3rd AAV with the fluorescent reporter (cre-
dependent tdTomato AAV) was omitted for cohorts undergoing behavioral analyses. For these
groups, histological confirmation for inclusion in subsequent statistical analyses was based on
identification of injections sites confined with the LS and CA1v, as observed using darkfield
microscopy.
468
Appetitive food seeking spatial memory task
To test visuospatial learning and memory for food reinforcement, we developed a novel
spatial food seeking task modified from the traditional Barnes maze procedure (4). Throughout
this paradigm, animals were maintained at 85% free-feeding body weight. The procedure
involves an elevated gray circular platform (diameter: 122cm, height: 140cm) consisting of 18
uniform holes (9.5cm diameter) spaced every twenty degrees around the outside edge. Under one
of the holes is a hidden tunnel (38.73cm L x 11.43cm W x 7.62cm D and a 5.08cm step).
Surrounding the table are distinct spatial cues on the wall (e.g., holiday lights, colorful shapes,
stuffed unicorn) that are readily visible to the animal. In contrast to the traditional Barnes Maze
where an animal uses the spatial cues to escape mildly aversive stimuli in the environment (e.g.

bright light and loud sound), this task utilizes food as motivation, such that each hidden tunnel

contained five 45mg sucrose pellets (Bio-Serv, Flemington, NJ). Additionally, a quiet white

noise (60dB) was used to drown out background noise and floor lamps were used for low-level

ambient lighting. On the first day, each animal underwent a habituation session consisting of

1min inside a transport bin under the table, 2min underneath the start box in the center of the

table, and 3 min inside the hidden tunnel with 5 sucrose pellets. During training, each rat was

assigned a specific escape hole according to the position of the spatial cues with the location

counterbalanced across groups. Before each trial, animals were placed in the start box for 30

seconds. Animals were trained over the course of two 5-min trials per day for four days (five

days in contralesional experiments) to learn to use the spatial cues in order to locate the correct

hole with the hidden tunnel with sucrose pellets. After finding the correct hole and entering the

hidden tunnel during each 5min training session, animals were allowed to remain in the hidden

tunnel for 1 minute, and consistently consumed all 5 sucrose pellets at this time. In the event that

a rat did not find the correct hole during the 5min training session, the experimenter gently

guided the rat to the correct hole and were allowed to crawl in and then complete the training in

the same fashion as rats that found the hole on their own. Learning during training was scored

via animal head location tracking by AnyMaze Behavior Tracking Software (Stoelting Co.,

Wood Dale, IL). The incorrect hole investigations prior to finding the correct hole with sucrose

pellets ("errors before correct hole") as well as time to finding the correct hole ("latency to

correct hole") were calculated as an average of the two trials per day, and examined across days

of training.

After the conclusion of training, rats had a two-day break where no procedures occurred,

then were tested in a single 2min memory probe in which the hidden tunnel and sucrose pellets

were been removed. All animals received an ICV 18mmol infusion of CNO (2uL total volume)

1h prior to the memory probe, rendering only DREADDs animals chemogenetically inactivated

by CNO. In the memory probe, the ratio of the correct hole plus adjacent hole investigations over

the total number of hole investigations were calculated via animal head location tracking by

AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). This dependent variable was

selected a priori instead of correct hole only investigations based on recent findings

demonstrating that taste-responsive hippocampal neurons exhibit weaker spatial selectivity in

comparison to non-taste-responsive hippocampal neurons (60). When establishing and refining

the food-reinforced spatial memory procedure in multiple cohorts of rats, our preliminary data

revealed that memory probe effects in normal/control animals are either stronger in the 1st

minute, or in the 1st two minutes combined (but not the 2nd minute alone), with some cohorts

showing larger paradigm effects in minute 1, and others in minutes 1+2. Thus, for all

experiments presented we analyzed both the 1st minute and the 1st two minutes combined based

on an a priori determination from our preliminary studies. A diagram of the apparatus used for

the spatial foraging task is included in Fig. 1B.

During training sucrose pellets were taped underneath all of the holes to preclude

olfactory-based search strategies during training. Importantly, the memory probe test does not

include any sucrose pellets in the apparatus at all, and therefore there is no possibility of sucrose

odor cues to influence memory probe performance.

Appetitive water seeking spatial memory task

We modified our spatial food seeking task to utilize water reward instead of food reward.

Throughout this paradigm, animals were water restricted and received 90-min access to water

daily (given at least 1hr after the termination of behavioral procedures each day). In addition to

the habituation parameters described above, animals in this task were subjected to two 10-min

habituation sessions in the hidden tunnel with a full water dish prior to training, which allowed

them to become accustomed to reliably drinking from the water dish in the hidden tunnel.

Training procedures were the same as for the food-based task above, except instead of sucrose

pellets in the hidden tunnel, a water dish containing ~100 mL of water was placed in the back of

the hidden tunnel during training, and animals were allowed to remain in the hidden tunnel for

2min (instead of 1min, as above). Each animal consumed a minimum of 2mL of water during

this time for the training sessions. This newly-developed procedure allowed us to test spatial

learning and memory motivated by water reward using similar apparatus and stimulus conditions

to the food-reinforced task.

Aversive spatial memory escape task (Barnes maze)

To test visuospatial learning and memory for escape reinforcement, we used a modified

traditional Barnes maze procedure, which is a visuospatial-based escape task (4). Procedures

were exactly the same as above (appetitive spatial memory food seeking and water seeking tasks,

using the same apparatus, in the same room, and with the same visuospatial cues) aside from the

omission of the sucrose pellets or water dish in the hidden tunnel, the presence of mildly aversive

bright (120W) overhead lighting instead of dim ambient lighting, and a mildly aversive loud

white noise (75dB) instead of a quiet white noise (60dB). This allowed us to test spatial learning

and memory motivated by escape from aversive stimuli in a nearly-identical procedure to our

spatial foraging test for learning and memory motivated by palatable food or water consumption.

A diagram of the apparatus used for the spatial escape task is included in Fig. 1B.

General intracranial injection procedures

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine

90mg/kg body weight [BW], xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW)

followed by a pre-operative, subcutaneous injection of analgesic (ketoprofen, 5mg/kg BW). Post-

operative analgesic (subcutaneous injection of ketoprofen, 5mg/kg BW) was administered once

per day for 3 days following surgery. The surgical site was shaved and prepped with iodine and

ethanol swabs, and animals were placed in a stereotaxic apparatus for stereotaxic injections.

NMDA or viruses were delivered using a microinfusion pump (Harvard Apparatus, Cambridge,

MA, USA) connected to a 33-gauge microsyringe injector attached to a PE20 catheter and

Hamilton syringe. Flow rate was calibrated and set to 83.3nl/sec. Injectors were left in place for

2min post-injection to allow for complete delivery of the infusate. Specific viruses/drugs,

coordinates, and injection volumes for procedures are detailed below. Following the completion

of all injections, incision sites were closed using either surgical staples, or in the case of

subsequent placement of an indwelling cannula, simple interrupted sutures. Upon recovery from

anesthesia and return to dorsal recumbency, animals were returned to the home cage. All

behavioral procedures occurred 21 days after injections to allow for complete transduction and

expression of the viruses, or complete lesioning drugs. General intracranial injection procedures

were followed for all injection procedures below.

Chronic lesions of the HPCv

Lesion animals received bilateral excitotoxic HPCv lesions via intracranial injections of

N-methyl-d-aspartate (NMDA; 4ug in 200nL; 100uL per hemisphere) at the following

coordinates at three different sites along the rostrocaudal extent of the HPCv (57): [1] -4.8mm

AP, +/-5.0mm ML, -7.5mm DV, [2] -5.5mm AP, +/-4.5mm ML, -7.0mm DV, and [3] -6.1mm

AP, +/-4.8mm ML, -7.0mm DV with control animals receiving vehicle saline in the same

location. The reference points for AP and ML coordinates were defined at bregma, and the

reference point for the DV coordinate was defined at the skull surface at the target site.

Bilateral HPCv lesion brains were histologically evaluated for the correct placement of

lesions in 1 out of 5 series of brain tissue sections. Neurons were visualized using

immunohistochemistry for the neuron specific antigen NeuN (see Immunohistochemistry). The

extent of NMDA lesions was determined postmortem by immunohistochemical detection of the

neuronal marker NeuN. Rats showing pronounced (~65%) loss of NeuN labeling within the

target region compared with the representative NeuN expression following sham injections in the

control group (average HPCv count of NeuN+ cells in control group) were included in data

analysis (43). Representative images in Fig. 1A.

Contralesional disconnection of the CA1v to mPFC neural pathway

To functionally disconnect the CA1v to mPFC pathway (diagram of approach in Fig. 4E),

lesion animals received a unilateral excitotoxic CA1v lesion via an intracranial injection of

NMDA (200nL) at the following coordinates (33): −4.9mm AP defined at bregma, + or - 4.8mm

ML defined at bregma (left/right counterbalanced to be contralateral to mPFC or LHA lesion

within-animal), −7.8mm DV defined at skull surface at site, with control animals receiving

vehicle saline injections in the same location.

In addition to the CA1v lesion, CA1v to mPFC disconnect animals also received a

unilateral excitotoxic mPFC lesion via two intracranial injections of NMDA (4ug in 200nL;

100nL per injection) at the following coordinates (57): [1] +2.7mm AP, + or - 0.7mm ML

(left/right counterbalanced to be contralateral to CA1v lesion within-animal), −5.3mm DV and

[2] +3.0mm AP, + or - 0.7mm ML (left/right counterbalanced to be contralateral to CA1v lesion

within-animal), −4.7mm DV, with control animals receiving vehicle saline in the same location.

The reference points for all mPFC coordinates were defined at bregma.

Contralesional brains were histologically evaluated for the correct placement of lesions by

visualizing neurons using immunohistochemistry for the neuron specific antigen NeuN (see

Immunohistochemistry). The extent of NMDA lesions was determined postmortem by

immunohistochemical detection of the neuronal marker NeuN. Rats showing pronounced (65%)

loss of NeuN labeling within the target region compared with a sham injection in the

contralateral side were included in data analysis (43).

Food intake and body weight

For the day prior to surgery (day 0) and for two weeks thereafter, 24h chow intake was

measured daily just prior to dark cycle onset to determine effects of experimental procedures on

food intake. Spillage was accounted for daily by collecting crumbs onto Techboard paper placed

under the cages of each animal. Additionally, animals were weighed daily just prior to dark cycle

onset to determine effects of experimental procedures on body weight.

Meal pattern

Meal size, meal frequency, meal duration, and cumulative 24h food intake were

measured using Biodaq automated food intake monitors (Research Diets, New Brunswick, NJ).

Rats (control n=7, caspase n=8, and DREADDs n=8) were acclimated to the Biodaq on ad

libitum chow for 3 days. Caspase group vs. control group feeding behavior data were collected

over a 5-day period (untreated, between-subjects design). The DREADDs group was tested using

a 2-treatment within-subjects design. Rats received 18mmol infusion of ICV CNO (2uL total

volume) or vehicle (daCSF, 33% dimethyl sulfoxide in artificial cerebrospinal fluid) 1h prior to

lights off and 24h feeding behavior was measured. Meal parameters were set at minimum meal

size=0.2g and maximum intermeal interval=600s.

Social transmission of food preference (STFP)

To examine food-related memory based on social- and olfactory-based cues, we utilized

the social transmission of food preference (STFP) task and adapted protocols from (28, 39, 61,

62). Briefly, untreated normal adult rats are designated as 'Demonstrators', while experimental

groups are designated as 'Observers'. Demonstrators and Observers are habituated to a powdered

rodent chow [LabDiet 5001 (ground pellets), LabDiet, St. Louis, MO] overnight. 24h later,

Observers are individually paired with demonstrators and habituated to social interaction, where

rat pairs are placed in a social interaction arena (23.5cm W × 44.45cm L × 27cm H clear plastic

bin with Sani-chip bedding) and allowed to interact for 30min. Both Observers and

Demonstrators are returned to their home cages and food is withheld for 23hr prior to the social

interaction. For the social interaction, Demonstrators are given the opportunity to consume one

of two flavors of powdered chow (flavored with 2% marjoram or 0.5% thyme; counterbalanced

according to group assignments) for 30min in a room separate from Observers. Our pilot studies

and previous published work (62, 63) show that rats equally prefer these flavors of chow. The

Demonstrator rat is then placed in the social interaction arena with the Observer rat, and the pairs

are allowed to socially interact for 30min. Observers are then returned to their home cage and

allowed to eat *ad libitum* for 1h and then food is removed. The following day, the 23h food-

deprived Observer animals are given a home cage food preference test for either the flavor of

chow paired with the Demonstrator animal, or a novel, unpaired flavor of chow that is a flavor

that was not given to the Demonstrator animal (2% marjoram vs. 0.5% thyme; counterbalanced

according to group assignments). All animals received an ICV 18mmol infusion of CNO (2uL

total volume) 1h prior to the social interaction session, rendering only DREADDs animals

chemogenetically inactivated by CNO. Food intake (1h) was recorded with spillage accounted

for by collecting crumbs onto Techboard paper that is placed under the cages of each animal.

The % preference for the paired flavor is calculated as: 100*Demonstrator-paired flavored chow

intake/Demonstrator + Novel flavored chow intake. In this procedure, normal untreated animals

learn to prefer the Demonstrator paired flavor based on social interaction and smelling the breath

of the Demonstrator rat (28, 39, 61-63). A diagram of the STFP procedure is included in Fig. 3E.

653   Zero maze

The zero maze behavioral paradigm was used to evaluate anxiety-like behavior. The zero

maze apparatus used was an elevated circular track, divided into four equal length sections. Two

zones were open with 3 cm high curbs ('open zones'), whereas the two other zones were closed

with 17.5 cm high walls ('closed zones'). All animals received an ICV 18mmol infusion of CNO

(2uL total volume) 1h prior to testing, rendering only DREADDs animals chemogenetically

inactivated by CNO. Behavioral testing was performed during the light cycle. Animals were

placed in the maze for a single, 5min trial in which the location of the center of the animal's

body was measured by AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). The

apparatus was cleaned with 10% ethanol in between animals. During the trial, the number of

open zone entries and total time spent in open sections (defined as body center in open sections)

were measured, which are each indicators of anxiety-like behavior in this procedure. A diagram

of the zero maze apparatus is included in Fig. 3G.

Open field

668   An open field test was used to evaluate general levels of locomotor activity. The

apparatus used for the open field test was an opaque gray plastic bin (60cm × 56cm), which was

positioned on a flat table in an isolated room with a camera directly above the center of the

apparatus. Desk lamps were positioned to provide indirect lighting to all corners of the maze

such that the lighting in the box uniformly measured 30 lux throughout. All animals received an

ICV 18mmol infusion of CNO (2uL total volume) 1h prior to testing rendering only DREADDs

animals chemogenetically inactivated by CNO. Behavioral testing began at dark onset. At the

start of the 10min test, each animal was placed in the open field apparatus in the same corner

facing the center of the maze. The location of the center of the animal's body was measured with

the AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). Total distance traveled

was measured by tracking movement from the center of the animal's body throughout the test.

Identification of collateral targets of CA1v to LS projecting neurons

681   Collateral targets of the CA1v to LS neural pathway were identified using a dual viral

approach (diagram of approach in Fig. 4A) where a retrograde vector was injected into the LS

(AAV2retro-hSyn1-eGFP-2A-iCre-WPRE; 200nL per side; coordinates as above), and a Cre-

dependent anterograde vector (AAV1-CAG-Flex-tdTomato-WPRE-bGH; 200nL per side;

coordinates as above) was injected in the CA1v. This latter injection drives tdTomato transgene

expression in CA1v neurons that project to the LS, which allows for brain-wide analyses of

collateral targets. After 3 weeks of incubation time to allow for complete transduction and

expression of the viruses, brains were collected, immunohistochemically processed, and imaged

as described below.

cFOS + FISH neurotransmitter phenotyping of LS neurons engaged during appetitive

spatial memory testing

693   Activation of LS neurons from the spatial memory probe test was confirmed with c-Fos

immunoreactivity analyses. Animals receiving the DREADDs-mediated treatment for reversible

inactivation of CA1v to LS signaling underwent training in the appetitive food seeking memory

task (conducted as described above). One hour before the memory probe, animals received ICV

injection of CNO (n=4) or vehicle (n=4). Animals were tested in the memory probe and perfused

with tissue harvest 90 min later. Fluorescent *in situ* hybridization analyses for c-Fos, ChAT,

GAD1, and vGLUT2 [source] were performed, and co-localization of c-Fos positive cells and

individual neurotransmitter markers was quantitatively analyzed for the rats (n=4) in the vehicle

treated group.

Identification of second-order targets of CA1v to LS projecting neurons

To identify second order targets of CA1v to LS-projecting neurons (diagram of approach

in Fig. 5A), animals received a bilateral injection of a transsynaptic Cre-inducing anterograde

vector into CA1v (AAV1-hSyn-Cre-WPRE-hGH; 200nL per side; coordinates as above) that

drives expression of Cre in both first-order (CA1v injection site) and 2^{nd}-order (but not 3^{rd}-

order) neurons via transsynaptic virion release (45, 64). This was combined with a bilateral

injection of a Cre-dependent anterograde vector in the LS (AAV1-CAG-Flex-tdTomato-WPRE-

bGH; 200nL per side, coordinates as above). This latter injection allows for anterograde tracing

from 1^{st}-order LS targets receiving synaptic input from CA1v. After 3 weeks of incubation time

to allow for complete transduction and expression of the viruses, brains were collected,

immunohistochemically processed, and imaged as described below.

Data were entered using a custom built data-entry platform (Axiome C, created by JDH)

built around Microsoft Excel software and designed to facilitate entry of data points for all gray

matter regions across their atlas levels as described in a rat brain reference atlas: Brain Maps 4.0

(57). The Axiome C approach was used previously to facilitate the analysis of brain gene-

expression data (65). An ordinal scale, ranging from 0 (absent) to 7 (very strong), was used to

record the qualitative weight of anterograde labeling. An average value was then obtained for

each region across its atlas levels for which data were available. These data are summarized

graphically for a representative animal on a brain flatmap summary diagram (adapted from (66)).

The LHA was identified as a primary downstream target of the CA1v to LS pathway

from the analyses described above. In order to further confirm this conclusion, an additional co-

injection monosynaptic neural tracing strategy (COIN; (44, 45)) was conducted. Animals (n=6)

received unilateral pressure injections of AAV1-GFP (AAV1-hSyn-eGFP-WPRE-bGH;

Addgene) targeting the CA1v and, in the same animal, fluorogold (Fluorochrome) targeting the

LHA (injection coordinates described above). Following a 10-day incubation period, animals

were fixation-perfused and tissue was collected and processed as described below. Injection sites

were confined to the CA1v (AAV1) and LHA (FG) in the same animal ("double hits", n=4).

Appositions between FG and GFP were examined in the LS in these animals by visualizing

native GFP and GF fluorescence.

Immunohistochemistry

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine

90mg/kg BW xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW) then transcardially

perfused with 0.9% sterile saline (pH 7.4) followed by 4% paraformaldehyde (PFA) in 0.1M

borate buffer (pH 9.5; PFA). Brains were dissected from the skull and post-fixed in PFA with

15% sucrose for 24h, then flash frozen in isopentane cooled in dry ice. Brains were sectioned to

30μm thickness on a freezing microtome. Sections were collected in 5-series and stored in

antifreeze solution at −20°C until further processing. General fluorescence IHC labeling

procedures were performed as follows. The antibodies and dilutions that were used are as

follows: [1] For lesion histology using the neuron-specific protein NeuN, the rabbit anti-NeuN

primary antibody (1:1000, Abcam) was used followed by a donkey anti-rabbit conjugated to

AF488 (1:500, Jackson Immunoresearch). [2] To amplify the native tdTomato signal for

neuroanatomical tracing or DREADDs histology, the rabbit anti-RFP primary antibody (1:2000,

Rockland) was used followed by a donkey anti-rabbit conjugated to Cy3 (1:500, Jackson

Immunoresearch). [3] To amplify the native GFP signal for LS injection site histology, the

chicken anti-GFP primary antibody (1:500, Abcam) was used followed by a donkey anti-chicken

secondary antibody conjugated to AF488 (1:500, Jackson Immunoresearch). Antibodies were

prepared in 0.02M potassium phosphate buffered saline (KPBS) solution containing 0.2% bovine

serum albumin and 0.3% Triton X-100 at 4°C overnight. After thorough washing with 0.02M

KPBS, sections were incubated in secondary antibody solution. All secondary antibodies were

obtained from Jackson Immunoresearch and used at 1:500 dilution at 4°C, with overnight

incubations (Jackson Immunoresearch; West Grove, PA, USA). Sections were mounted and

coverslipped using 50% glycerol in 0.02 M KPBS and the edges were sealed with clear nail

polish. Photomicrographs were acquired using a Nikon 80i (Nikon DSQI1,1280X1024

resolution, 1.45 megapixel) under epifluorescence or darkfield illumination.

Lesions and virus expression were quantified in one out of five series of brain tissue sections and

analyses were performed in sections from Swanson Brain Atlas level 34-36 (57). The inclusion /

exclusion criteria are described above in the methods of each of these experiments.

Statistics

Data are expressed as mean +/- SEM. Differences were considered to be statistically significant

at $p<0.05$. All variables were analyzed using the advanced analytics software package Statistica

(StatSoft, Tulsa, OK, USA). For all measures of food intake, body weight, and errors/latency

during spatial foraging and spatial escape task training, differences between groups were

evaluated using two-way repeated measures ANOVAs (treatment x time). For lesion

experiments, measures during spatial foraging and spatial escape task probes, differences

between groups were evaluated using two-tailed independent two-sample Student's t-tests. For

CA1v to LS disconnection experiments, measures during spatial foraging and spatial escape task

probes, the STFP paradigm, the zero maze paradigm, and the open field test, differences between

groups were evaluated using one-way ANOVAs. Significant ANOVAs were analyzed with a

Fisher's LSD posthoc test where appropriate. Outliers were identified as being more extreme

than the median +/- 1.5 * interquartile range. For all experiments, assumptions of normality,

homogeneity of variance (HOV), and independence were met where required.

776
777   **ACKNOWLEDGEMENTS**
778

779   This work was supported by National Institute of Diabetes and Digestive and Kidney Diseases

780   grants: DK104897 to SEK, DK118944 to CML, and DK116558 to ANS. Clozapine-N-Oxide

781   was kindly provided by the National Institute of Mental Health. The authors are grateful to the

782   Kanoski lab undergraduate research assistants for their assistance in behavioral experiments and

783   histology.

784
785
786   **COMPETING INTERESTS**

788   There are not competing interests to declare.

Figure 1

Figure 1. Bilateral HPCv lesions impair spatial memory for food location. Representative HPCv lesion histology with NeuN immunohistochemistry (A; scale bars 500μm). Spatial foraging task apparatus (B). Representative navigation paths of a control animal preferentially investigating correct (filled red) and adjacent holes (outlined orange) during spatial foraging memory probe (C). Representative navigation path of HPCv lesioned animal during spatial foraging memory probe (D). Bilateral HPCv lesions did not impair learning of the spatial foraging task compared to controls, as measured by errors before locating correct hole during task training (E) and latency to locate correct hole during task training (F). Bilateral HPCv lesions impaired retention of the spatial foraging task, as measured by the ratio of investigation of correct plus adjacent holes over total investigated during the first minute of the task ($p < 0.05$; G). Dotted line indicates chance performance level (0.167). For graphs 1D-G, lesion n = 11, control n = 18. All values expressed as mean +/- SEM.

Figure 2

Figure 2. Reversible and chronic CA1v to LS neural disconnection impairs spatial memory for food and water location but not for escape location. Diagram of dual viral approach using a cre-dependent inhibitory DREADDs (AAV-hSyn-DIO-hM4D(Gi)-mCherry) approach to reversibly disconnect CA1v to LS neural pathway (with CNO injected 1hr before memory probe) (A). Diagram of dual viral approach using a cre-dependent caspase (AAV1-Flex-taCasp3-TEVp) approach to chronically disconnect CA1v to LS neural pathway (B). Representative injection site in CA1v demonstrating LS-projecting neurons infected with inhibitory DREADDs, which simultaneously drives expression of a fluorescent mCherry transgene (C, D; scale bars 500μm and 100μm, respectively). Neither reversible (DREADDs) nor chronic (caspase) disconnection of the CA1v to LS pathway impaired learning of the spatial food seeking task compared to controls, as measured by errors before correct hole during task training (E) and latency to correct hole during task training (F). Both reversible and chronic disconnection of the CA1v to LS pathway impaired retention of the food location as measured by the ratio of investigation of correct plus adjacent holes over total investigations during entire two minutes of the task ($p<0.05$; G). Likewise, reversible (DREADDs) and chronic (caspase) disconnection of the CA1v to LS pathway did not impair learning of the spatial water seeking task compared to controls, as measured by errors before correct hole during task training (H) and latency to correct hole during task training (I), but impaired memory retention of the water location during the probe (J). In contrast, disconnection of the CA1v to LS pathway either reversibly (DREADDs) or chronically (caspase) did not impair performance on the spatial escape task. There were no differences in learning as measured by errors before correct hole during task training (K) and latency to correct hole during task training (L). Unlike the spatial foraging task, retention of the spatial escape task was not impaired by reversible nor chronic disconnection of the CA1v to LS pathway (M). For graphs 2E-G (CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8. For graphs 2H-J (CA1v to LS disconnect cohort 3), DREADDs n=8, caspase n=8, control n=7). For graphs 2K-M (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control n=10. Dotted line indicates chance performance level (0.167). All values expressed as mean +/- SEM.

Figure 3

A

Day 1
Demonstrators and observers habituate to social interaction

Day 2
Demonstrators consume flavored chow (chow A) prior to social interaction

Demonstrators and observers have a 30min social interaction

Day 3
Food-restricted observers are given a 60min flavor preference test (chow A vs. novel chow B)

B
Social transmission of food preference

Preference ratio

1.0
0.8
0.6
0.4
0.2
0.0

Control   DREADDs   Caspase

C

D
Zero maze: Open zone time

Time in open zones (s)

Control   DREADDs   Caspase

E
Zero maze: Open zone entries

Entries to open zones

Control   DREADDs   Caspase

F
Open field

Distance traveled (m)

20
15
10
5

Control   DREADDs   Caspase

Figure 3. Neither reversible nor chronic CA1v to LS neural disconnection impair spatial

memory for escape location, social transmission of food preference, anxiety-like behavior,

or general locomotor activity levels. Diagram of the social transmission of food preference

(STFP) task (A). Neither reversible nor chronic disconnection of the CA1v to LS pathway impair

STFP learning compared to controls, as measured by a food preference ratio (B), with the dotted

line indicating chance preference level (0.50). Diagram of the zero maze apparatus (C). Anxiety-

like behavior was not influenced by reversible or chronic disconnection of the CA1v to LS

pathway compared to controls, as measured by performance in the zero maze task, specifically

time in open zones (D) and entries into open zones (E). Neither chronic nor reversible CA1v to

LS disconnection affected open field performance compared to controls, as measured by total

distance traveled (F). For graphs 3B, 3D, and 3E (CA1v to LS disconnect cohort 1), DREADDs

n=6, caspase n=10, control n=8. For graph 3F (CA1v to LS disconnect cohort 2), DREADDs

n=8, caspase n=12, control n=10. All values expressed as mean +/- SEM.

Figure 4

A

AAVretro-Cre-GFP

LS

AAV1-Flex-tdTomato in CA1v

CA1v

B
CA3v
CA1v
alv
alv
tdTomato (CA1v)

C
ccb
LSc
LSr
VLa
GFP (LSc)

D
PL
ILA
frf
tdTomato (mPFC)

E
mPFC — mPFC
LS   LS
CA1v — CA1v

F
Spatial foraging training (errors)
Average errors
- Control
- CA1v-mPFC disconnect
Day 1 Day 2 Day 3 Day 4 Day 5

G
Spatial foraging training (latency)
Average latency (s)
- Control
- CA1v-mPFC disconnect
Day 1 Day 2 Day 3 Day 4 Day 5

H
Memory probe
Correct + adjacent holes / total
Control   CA1v-mPFC disconnect

Figure 4. CA1v to mPFC disconnection does not impair spatial memory for food location.

Diagram of dual viral approach to identify collateral targets of the CA1v to LS neural pathway

(A). Representative CA1v injection site from collateral identification approach (B; scale bar

500μm). Representative LS injection site from collateral identification approach (C; scale bar

500μm). Representative image collateral axons of the CA1v to LS pathway located in the mPFC

(D; scale bar 50μm). Diagram of contralesional approach to functionally disconnect the CA1v to

mPFC neural pathway (E). Disconnection of the CA1v to mPFC neural pathway did not

influence learning of the spatial foraging task compared to controls, as measured by errors before

correct hole during training (F) and latency to correct hole during training (G). Disconnection of

the CA1v to mPFC pathway did not influence retention of the spatial foraging task in the

memory probe (H). For graphs 4F-H, CA1v to mPFC disconnect n=12, control n=12. All values

expressed as mean +/- SEM.

Figure 5

Figure 5. LS neurons activated during food-seeking memory probe are predominantly GABAergic. c-Fos and ChAT colocalization was observed in ~2% of LS neurons (A; scale bar 500μm), c-Fos and GAD1 colocalization was observed in ~88% of LS neurons (B; scale bar 500μm), and c-Fos and VGLUT2 colocalization was observed in ~10% of LS neurons (C; scale bar 500μm) following the spatial memory probe test (for food reinforcement). For graphs A-C, quantification from DREADDs vehicle group, n=4.

Figure 6

A

LS

Anterograde monosynaptic Cre-dependent AAV1-tdTomato

CA1v

Anterograde transsynaptic AAV2/1-hSyn-Cre

B

ccb

VLa

LSc

LSr

LS injection site

C

LHAjd

fx

LHAd

LHAjvd

LHAsfa

LHAjvv

2° tdTomato+ fibers

D

CA1v → LS → multiple regions

CEREBRAL CORTEX

CORTICAL SUBPLATE

CEREBELLAR CORTEX

CORTICAL PLATE

PALLIDUM

STRIATUM

THALAMUS

MIDBRAIN

PONS

MEDULLA

SPINAL CORD

HYPOTHALAMUS

6 5 4 3 2 1 0

Aggregate ordinal value categories
(6 = strongest; 0 = weakest / absent)

Targeted side (right)

HIPPOCAMPAL REGION

RETROHIPPOCAMPAL REGION

CA1v

THALAMUS

LATERAL HYPOTHALAMIC AREA

ORBITAL REGION

LS

SUBSTANTIA NIGRA

Contralateral side (left)

HIPPOCAMPAL REGION

LATERAL HYPOTHALAMIC AREA

E

mtt

fx

V3h

LHA FG injection site

F

CA3v

CA1v

alv

CA1v AAV1-GFP injection site

G

ccb

VLa

LSc

LSr

FG-GFP appositions

Figure 6. Identification of second-order neural projections downstream of CA1v to LS

projections. Diagram of dual viral approach to identify brain regions that are second-order (2°)

targets of the CA1v to LS neural pathway (A). Representative LS injection site from second

order identification approach (B; scale bar 100μm). Representative image of second-order fibers

of the CA1v to LS pathway within the LHA (C; scale bar 200μm). Summary of the projection

targets of LS neurons that receive input from CA1v (D; expanded information in Supplementary

Table 1). The outputs of the right side of LS neurons receiving CA1v input are represented at the

macroscale (gray matter region resolution) on a partial flatmap representation of the rat

forebrain, adapted from (66). Connection weights are represented block colors for each region

following an ordinal scale ranging from weakest (0 = very weak or absent) to strongest (6 =

strong), as there were no 7 (very strong) values. The inset at lower left represents one side of the

brain with the part represented in the upper diagrams outlined in magenta. The COIN approach

for further confirmation of second-order tracing strategy is depicted with FG injection site in the

LHA (E; scale bar 500μm), representative AAV1-GFP injection site in the CA1v (F; scale bar

100μm), and FG-GFP appositions in the LS (G; scale bar 100μm).

Supplemental Figure 1

886 **Supplementary Figure 1. Effect of bilateral HPCv lesions on food intake and body weight.**

887 There were no effects on food intake (A) or body weight (B) of bilateral HPCv lesions compared

888 to controls. Lesion n = 11, control n = 18. All values expressed as mean +/- SEM.

Supplemental Figure 2

Supplementary Figure 2. Effect of reversible and chronic disconnection of the CA1v to LS

neural pathway on food intake and body weight, and verification of the acute and chronic

approach. There were no effects on food intake (cohort 1: A; cohort 2: B) or body weight

(cohort 1: C; cohort 2: D) of reversible (DREADDs) or chronic (caspase) disconnection of the

CA1v to LS neural pathway compared to controls. Histological verification of inhibitory

DREADDs-induced acute silencing of LS-projecting CA1v neurons determines that the

appetitive spatial memory probe induces c-Fos activation in the LS in vehicle-treated animals

(E), an outcome significantly reduced in CNO-treated animals (F, G). Histological verification of

caspase-induced chronic silencing of the LS-projecting CA1v neurons demonstrates that cre-

dependent tdTomato labeling of CA1v neurons induced by LS-origin retro-cre is robust in a

control animal (H), but substantially reduced in an animal injected with cre-dependent caspase

combined with cre-dependent tdTomato due to caspase-induced cell death (I). For graphs S2A-B

(CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8. For graphs S2C-

D (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control n=10. For graphs

S2E-G, DREADDs vehicle n=4, DREADDs CNO n=4. For graphs S2H-I, caspase n=6, control

n=2). All values expressed as mean +/- SEM.

Supplemental Figure 3

Chronic Disconnection of the vCA1-LS Pathway

Supplementary Figure 3. Effect of reversible and chronic disconnection of CA1v to LS

neural pathway on feeding behavior. Compared with control animals, chronic caspase-

mediated disconnection of the CA1v to LS pathway did not significantly affect meal frequency

(A), meal size (B), meal duration (C), or 24h cumulative food intake (D) over the course of the

5-day recording period. Acute DREADDs-mediated disconnection of the CA1v to LS pathway

(via CNO infusion) resulted in non-significant trend towards increased meal frequency (E), no

differences in meal size (F) or meal duration (G), and a significant increase in 24h cumulative

food intake (H) in comparison with vehicle injection. For graphs S3A-D, caspase n=7 and

control n=7. For graphs S3E-H, DREADDs n=8 (within-subjects). All values expressed as mean

+/- SEM.

Supplemental Figure 4

A

CA1v to mPFC disconnect:
Food intake

B

CA1v to mPFC disconnect:
Body weight

921   **Supplementary Figure 4. Effect of contralesional disconnection methods on food intake and**

922   **body weight.** There were no effects on food intake (A) or body weight (B) of contralesional

923   CA1v to mPFC compared to controls. CA1v to mPFC disconnect n=12, control n=12. All values

924   expressed as mean +/- SEM.

Supplementary Table 1

927   **Supplementary Table 1.** Summary table of the forebrain projection targets of LS neurons that

928   receive input from CA1v (derived the same raw data as the brain flatmap summary diagram in

929   Fig. 6D).

REFERENCES

1.   R. G. M. Morris, U. Frey, "Hippocampal synaptic plasticity: Role in spatial learning or the automatic recording of attended experience?" in Burgess, Neil (Ed); Jeffery, Kathryn J (Ed); et al (1999) The hippocampal and parietal foundations of spatial cognition. (Oxford University Press, U Edinburgh, Ctr for Neuroscience, Edinburgh, Scotland London, 1999), pp. 220-246.

2.   D. C. Rowland, Y. Roudi, M. B. Moser, E. I. Moser, Ten Years of Grid Cells. *Annu Rev Neurosci* **39**, 19-40 (2016).

3.   R. Morris, Developments of a water-maze procedure for studying spatial learning in the rat. *Journal of neuroscience methods* **11**, 47-60 (1984).

4.   C. A. Barnes, Memory deficits associated with senescence: a neurophysiological and behavioral study in the rat. *J Comp Physiol Psychol* **93**, 74-104 (1979).

5.   J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map: preliminary evidence from unit activity in the freely-moving rat. *Brain research* (1971).

6.   T. Hafting, M. Fyhn, S. Molden, M. B. Moser, E. I. Moser, Microstructure of a spatial map in the entorhinal cortex. *Nature* **436**, 801-806 (2005).

7.   M. S. Fanselow, H.-W. Dong, Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* **65**, 7-19 (2010).

8.   A. T. Keinath *et al.*, Precise spatial coding is preserved along the longitudinal hippocampal axis. *Hippocampus* **24**, 1533-1548 (2014).

9.   K. B. Kjelstrup *et al.*, Finite scale of spatial representation in the hippocampus. *Science* **321**, 140-143 (2008).

10. L. de Hoz, J. Knox, R. G. Morris, Longitudinal axis of the hippocampus: both septal and temporal poles of the hippocampus support water maze spatial learning depending on the training protocol. *Hippocampus* **13**, 587-603 (2003).

11. J. Ferbinteanu, C. Ray, R. J. McDonald, Both dorsal and ventral hippocampus contribute to spatial learning in Long-Evans rats. *Neurosci Lett* **345**, 131-135 (2003).

12. M. B. Moser, E. I. Moser, Functional differentiation in the hippocampus. *Hippocampus* **8**, 608-619 (1998).

13. S. E. Kanoski, H. J. Grill, Hippocampus Contributions to Food Intake Control: Mnemonic, Neuroanatomical, and Endocrine Mechanisms. *Biol Psychiatry* **81**, 748-756 (2017).

14. P. G. Henke, Hippocampal pathway to the amygdala and stress ulcer development. *Brain Res Bull* **25**, 691-695 (1990).

15. K. G. Kjelstrup *et al.*, Reduced fear expression after lesions of the ventral hippocampus. *Proceedings of the National Academy of Sciences* **99**, 10825-10830 (2002).

16. S. Gaskin, A. Gamliel, M. Tardif, E. Cole, D. G. Mumby, Incidental (unreinforced) and reinforced spatial learning in rats with ventral and dorsal lesions of the hippocampus. *Behav Brain Res* **202**, 64-70 (2009).

17. M. S. Bienkowski *et al.*, Integration of gene expression and brain-wide connectivity reveals the multiscale organization of mouse hippocampal networks. *Nat Neurosci* **21**, 1628-1643 (2018).

18. R. Hannapel *et al.*, Postmeal Optogenetic Inhibition of Dorsal or Ventral Hippocampal
Pyramidal Neurons Increases Future Intake. *eNeuro* **6** (2019).
19. R. C. Hannapel, Y. H. Henderson, R. Nalloor, A. Vazdarjanova, M. B. Parent, Ventral
hippocampal neurons inhibit postprandial energy intake. *Hippocampus* **27**, 274-284
(2017).
20. B. K. Mani *et al.*, Neuroanatomical characterization of a growth hormone secretagogue
receptor-green fluorescent protein reporter mouse. *J Comp Neurol* **522**, 3644-3666
(2014).
21. J. M. Zigman, J. E. Jones, C. E. Lee, C. B. Saper, J. K. Elmquist, Expression of ghrelin
receptor mRNA in the rat and the mouse brain. *J Comp Neurol* **494**, 528-548 (2006).
22. I. Merchenthaler, M. Lane, P. Shughrue, Distribution of pre-pro-glucagon and glucagon-
like peptide-1 receptor messenger RNAs in the rat central nervous system. *J Comp*
Neurol **403**, 261-280 (1999).
23. S. E. Kanoski, S. M. Fortin, K. M. Ricks, H. J. Grill, Ghrelin Signaling in the Ventral
Hippocampus Stimulates Learned and Motivational Aspects of Feeding via PI3K-Akt
Signaling. *Biol Psychiatry* **73**, 915-923 (2013).
24. S. E. Kanoski *et al.*, Hippocampal leptin signaling reduces food intake and modulates
food-related memory processing. *Neuropsychopharmacology* **36**, 1859-1870 (2011).
25. T. M. Hsu *et al.*, A hippocampus to prefrontal cortex neural pathway inhibits food
motivation through glucagon-like peptide-1 signaling. *Molecular psychiatry* **23**, 1555-
1565 (2018).
26. T. M. Hsu, J. D. Hahn, V. R. Konanur, A. Lam, S. E. Kanoski, Hippocampal GLP-1 receptors
influence food intake, meal size, and effort-based responding for food through volume
transmission. *Neuropsychopharmacology* **40**, 327-337 (2015).
27. A. N. Suarez, C. M. Liu, A. M. Cortella, E. E. Noble, S. E. Kanoski, Ghrelin and Orexin
Interact to Increase Meal Size Through a Descending Hippocampus to Hindbrain
Signaling Pathway. *Biol Psychiatry* **87**, 1001-1011 (2020).
28. T. M. Hsu *et al.*, Hippocampus ghrelin receptor signaling promotes socially-mediated
learned food preference. *Neuropharmacology* **131**, 487-496 (2018).
29. G. D. Petrovich, N. S. Canteras, L. W. Swanson, Combinatorial amygdalar inputs to
hippocampal domains and hypothalamic behavior systems. *Brain research reviews* **38**,
247-289 (2001).
30. C. De La Rosa-Prieto *et al.*, Subicular and CA1 hippocampal projections to the accessory
olfactory bulb. *Hippocampus* **19**, 124-129 (2009).
31. L. W. Swanson, W. M. Cowan, An autoradiographic study of the organization of the
efferent connections of the hippocampal formation in the rat. *J Comp Neurol* **172**, 49-84
(1977).
32. A. Arszovszki, Z. Borhegyi, T. Klausberger, Three axonal projection routes of individual
pyramidal cells in the ventral CA1 hippocampus. *Front Neuroanat* **8**, 53 (2014).
33. P. Y. Risold, L. W. Swanson, Structural evidence for functional domains in the rat
hippocampus. *Science* **272**, 1484-1486 (1996).
34. P. Sweeney, Y. Yang, An excitatory ventral hippocampus to lateral septum circuit that
suppresses feeding. *Nature communications* **6**, 10188 (2015).

35.   R. Garcia, R. M. Vouimba, R. Jaffard, Spatial discrimination learning induces LTP-like
1018   changes in the lateral septum of mice. *Neuroreport: An International Journal for the*
1019   *Rapid Communication of Research in Neuroscience* **5**, 329-332 (1993).
1020   36.   R. Jaffard, R. M. Vouimba, A. Marighetto, R. Garcia, Long-term potentiation and long-
1021   term depression in the lateral septum in spatial working and reference memory. *J*
1022   *Physiol Paris* **90**, 339-341 (1996).
1023   37.   S. J. Terrill *et al.*, Nucleus accumbens melanin-concentrating hormone signaling
1024   promotes feeding in a sex-specific manner. *Neuropharmacology* **178**, 108270 (2020).
1025   38.   A. Carballo-Marquez, A. Vale-Martinez, G. Guillazo-Blanch, M. Marti-Nicolovius,
1026   Muscarinic receptor blockade in ventral hippocampus and prelimbic cortex impairs
1027   memory for socially transmitted food preference. *Hippocampus* **19**, 446-455 (2009).
1028   39.   R. A. Countryman, N. L. Kaban, P. J. Colombo, Hippocampal c-fos is necessary for long-
1029   term memory of a socially transmitted food preference. *Neurobiol Learn Mem* **84**, 175-
1030   183 (2005).
1031   40.   G. M. Parfitt *et al.*, Bidirectional Control of Anxiety-Related Behaviors in Mice: Role of
1032   Inputs Arising from the Ventral Hippocampus to the Lateral Septum and Medial
1033   Prefrontal Cortex. *Neuropsychopharmacology : official publication of the American*
1034   *College of Neuropsychopharmacology* **42**, 1715-1728 (2017).
1035   41.   N. L. Trent, J. L. Menard, The ventral hippocampus and the lateral septum work in
1036   tandem to regulate rats' open-arm exploration in the elevated plus-maze. *Physiology &*
1037   *behavior* **101**, 141-152 (2010).
1038   42.   J. C. Jimenez *et al.*, Anxiety Cells in a Hippocampal-Hypothalamic Circuit. *Neuron* **97**,
1039   670-683.e676 (2018).
1040   43.   T. M. Hsu *et al.*, Hippocampus ghrelin signaling mediates appetite through lateral
1041   hypothalamic orexin pathways. *eLife* **4** (2015).
1042   44.   R. H. Thompson, L. W. Swanson, Hypothesis-driven structural connectivity analysis
1043   supports network over hierarchical model of brain architecture. *Proc Natl Acad Sci U S A*
1044   **107**, 15235-15239 (2010).
1045   45.   A. N. Suarez *et al.*, Gut vagal sensory signaling regulates hippocampus function through
1046   multi-order pathways. *Nature communications* **9**, 2181 (2018).
1047   46.   M. S. Fanselow, H. W. Dong, Are the dorsal and ventral hippocampus functionally
1048   distinct structures? *Neuron* **65**, 7-19 (2010).
1049   47.   M. Contreras, T. Pelc, M. Llofriu, A. Weitzenfeld, J. M. Fellous, The ventral hippocampus
1050   is involved in multi-goal obstacle-rich spatial navigation. *Hippocampus* **28**, 853-866
1051   (2018).
1052   48.   S. L. T. Lee, D. Lew, V. Wickenheisser, E. J. Markus, Interdependence between dorsal and
1053   ventral hippocampus during spatial navigation. *Brain Behav* **9**, e01410 (2019).
1054   49.   D. Olivo, M. Caba, F. Gonzalez-Lima, J. F. Rodríguez-Landa, A. A. Corona-Morales,
1055   Metabolic activation of amygdala, lateral septum and accumbens circuits during food
1056   anticipatory behavior. *Behavioural brain research* **316**, 261-270 (2017).
1057   50.   M. Carus-Cadavieco *et al.*, Gamma oscillations organize top-down signalling to
1058   hypothalamus and enable food seeking. *Nature* **542**, 232-236 (2017).

51. S. J. Terrill *et al.*, Role of lateral septum glucagon-like peptide 1 receptors in food intake.
American journal of physiology. Regulatory, integrative and comparative physiology **311**,
R124-132 (2016).
52. S. J. Terrill, K. D. Wall, N. D. Medina, C. B. Maske, D. L. Williams, Lateral septum growth
hormone secretagogue receptor affects food intake and motivation for sucrose
reinforcement. *American journal of physiology. Regulatory, integrative and comparative*
physiology **315**, R76-r83 (2018).
53. C. Wang, C. M. Kotz, Urocortin in the lateral septal area modulates feeding induced by
orexin A in the lateral hypothalamus. *American journal of physiology. Regulatory,*
integrative and comparative physiology **283**, R358-367 (2002).
54. J. Hauser, L. H. Llano López, J. Feldon, P. A. Gargiulo, B. K. Yee, Small lesions of the dorsal
or ventral hippocampus subregions are associated with distinct impairments in working
memory and reference memory retrieval, and combining them attenuates the
acquisition rate of spatial reference memory. *Hippocampus* **30**, 938-957 (2020).
55. D. S. Olton, The radial arm maze as a tool in behavioral pharmacology. *Physiology &*
behavior **40**, 793-797 (1987).
56. X. Zhao *et al.*, Orexin-A-induced ERK1/2 activation reverses impaired spatial learning and
memory in pentylenetetrazol-kindled rats via OX1R-mediated hippocampal
neurogenesis. *Peptides* **54**, 140-147 (2014).
57. L. W. Swanson, Brain maps 4.0-Structure of the rat brain: An open access atlas with
global nervous system nomenclature ontology and flatmaps. *J Comp Neurol* **526**, 935-
943 (2018).
58. R. C. Ritter, P. G. Slusser, S. Stone, Glucoreceptors controlling feeding and blood glucose:
location in the hindbrain. *Science* **213**, 451-452 (1981).
59. E. E. Noble *et al.*, Hypothalamus-hippocampus circuitry regulates impulsivity via
melanin-concentrating hormone. *Nat Commun* **10**, 4923 (2019).
60. L. E. Herzog *et al.*, Interaction of Taste and Place Coding in the Hippocampus. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **39**, 3057-
3069 (2019).
61. R. A. Countryman, P. E. Gold, Rapid forgetting of social transmission of food preferences
in aged rats: relationship to hippocampal CREB activation. *Learn Mem* **14**, 350-358
(2007).
62. B. G. Galef, Jr., W. Y. Lee, E. E. Whiskin, Lack of interference in long-term memory for
socially learned food preferences in rats (Rattus norvegicus). *J Comp Psychol* **119**, 131-
135 (2005).
63. B. G. Galef, Jr., E. E. Whiskin, Socially transmitted food preferences can be used to study
long-term memory in rats. *Learn Behav* **31**, 160-164 (2003).
64. B. Zingg *et al.*, AAV-Mediated Anterograde Transsynaptic Tagging: Mapping
Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron* **93**, 33-
47 (2017).
65. J. D. Hahn, O. Sporns, A. G. Watts, L. W. Swanson, Macroscale intrinsic network
architecture of the hypothalamus. *Proc Natl Acad Sci U S A* **116**, 8018-8027 (2019).

66.   J. D. Hahn, Swanson, L. W,. Bowman, I., Foster, N. N., Zingg, B., Bienkowski, M. S.,
Hintiryan, H., & Dong, H.-W. , An open access mouse brain flatmap and upgraded rat
and human brain flatmaps based on current reference atlases. *J. Comp. Neurol.* (2020).
1104
1105