Hippocampus-lateral septum circuitry mediates memory for food location in rats

Elizabeth A. Davis[1*], Clarissa M. Liu[1,2*], Isabella H. Gianatiempo[1], Andrea N. Suarez[1], Alyssa M. Cortella[1], Joel D. Hahn[3], and Scott E. Kanoski[1,2]

[*]These two authors contributed equally

[1]Human and Evolutionary Biology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089; [2]Neuroscience Graduate Program, University of Southern California, 3641 Watt Way, Los Angeles, CA, USA. 90089; [3]Neurobiology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089

Corresponding author:
Scott E. Kanoski, PhD
University of Southern California
3616 Trousdale Parkway, AHF-252
Los Angeles, CA 90089-0372
Phone: 213-821-5762
Email: kanoski@usc.edu

Classification: Biological Sciences, Neuroscience

Keywords: feeding, cognition, place cells, motivation, reward, foraging, lateral hypothalamic area, prefrontal cortex

ABSTRACT

Remembering the location of a food source is essential for securing energy for survival. Here we identify a hippocampal-septal neural circuit that controls food-directed spatial memory. Both reversible and chronic disconnection of ventral hippocampus CA1 subregion (CA1v) projections to the lateral-septum (LS) using pathway-specific dual viral approaches impaired memory retention in a spatial food-seeking foraging task in rats. However, disconnection of this pathway did not affect performance in an aversive escape-motivated spatial memory task that used the same apparatus and visuospatial cues, suggesting that CA1v-LS signaling selectively mediates spatial memory for food location vs. spatial memory in general. The selectivity of this pathway in mediating foraging-related spatial memory was further supported by results showing that CA1v-LS disconnection did not affect anxiety-like behavior, locomotor activity, or social and olfactory-based appetitive learning. To examine whether CA1v-LS mediation of foraging-related spatial memory involves collateral projections of CA1v neurons, we utilized virus-based neural pathway tracing analyses to identify the mPFC as a collateral target of LS-projecting CA1v neurons. However, functional disconnection of the CA1v and mPFC did not affect spatial memory for food location, thus further supporting the selectivity of CA1v-LS signaling for this behavior. The nucleus accumbens, lateral hypothalamic area, and other brain regions associated with food motivation and reward were identified as second-order targets of CA1v-LS signaling using a multisynaptic anterograde tracing approach. Collective results reveal that CA1v to LS communication plays a critical role in remembering the environmental location of food, thus identifying a novel neural pathway regulating foraging-related memory processes.

INTRODUCTION

A survival advantage common to early humans and lower-order mammals is to accurately remember the location of a food source in the environment, and to then efficiently navigate back to a shelter or other safe location. The neurobiological substrates that regulate visuospatial navigation are therefore critical to effectively implement food-directed foraging behavior. Brain structures within the telencephalon have been identified as being essential for visuospatial mapping and egocentric orientation in the environment, including the hippocampus and the medial entorhinal cortex, respectively (1, 2). However, despite that reliably locating food in the environment is a key selection pressure driving the evolution of visuospatial navigation, the overwhelming majority of rodent model research on the substrates of spatial learning and memory have utilized procedures such as the Morris water maze and the Barnes maze that each involve escaping aversive reinforcement (3, 4). Furthermore, while single unit recordings have been used to identify specific populations of neurons that subserve distinct navigational functions (e.g., hippocampal "place cells", medial entorhinal "grid cells")(5, 6), the bulk of this research has recorded neural activity under neutral conditions void of either appetitive or aversive reinforcement. Very little research has been dedicated to identify brain regions and neural circuits that may specifically promote food-directed spatial memory, as well as the extent to which the nature of the reinforcement is a critical factor in deciphering the brain's control of visuospatial navigation.

The majority of rodent model research investigating brain regions that mediate visuospatial navigational memory has focused on the anterior and "dorsal" subregion of the hippocampus (septal pole; HPCd). However, the posterior and "ventral" hippocampus subregion (temporal pole; HPCv), while classically associated with stress- and affect-associated memory processes (7), also plays a role in visuospatial learning and memory (8-10). For example, under some testing conditions selective HPCv lesions impair spatial memory performance in the Morris water maze (10, 11). Moreover, place cells that are responsive to changes in the visuospatial environment are present within both the HPCd and HPCv pyramidal layers, with a linear increase in the scale of representation from the dorsal to the ventral pole (9). Despite a common role for the HPCd and HPCv in mediating spatial memory, there is also evidence for a functional distinction between the subregions (7, 12, 13). For instance, lesions of the HPCv but not the

HPCd alter stress responses (14) and anxiety-like behavior (15), whereas HPCd but not HPCv

lesions impair spatial memory in an incidental (nonreinforced) procedure (16). These two HPC

subregions also have distinct afferent and efferent neural connections. This disparate

neuroanatomical connectivity supports a framework for a functional diversity in which the HPCd

preferentially processes cortical-derived sensory information and the HPCv preferentially

processes metabolic and limbic-derived affective information (7, 13). The functional distinction

between these two subregions is further supported by the generation of the hippocampus gene

expression atlas, which provides a comprehensive integration of gene expression and

connectivity across the HPC axis (17). Given that both the HPCv and the HPCd participate in

spatial memory but have distinct neuroanatomical connectivity and contributions when it comes

to regulating other cognitive and mnemonic processes, it is feasible that the HPCv and HPCd

support different forms of spatial memory depending on the type of reinforcement or context

associated with the behavior.

Recent findings identify the HPCv as a critical brain substrate in regulating feeding

behavior and food-directed memory processes. Reversible inactivation of HPCv neurons after a

meal increases the size of and reduces the latency to consume a subsequent meal (18, 19). In

addition, receptors for feeding-related hormones are more abundantly expressed in the HPCv

compared to the HPCd (e.g., ghrelin receptors (20, 21), glucagon-like peptide-1 [GLP-1]

receptors (22)), and these HPCv endocrine receptor systems alter food intake and feeding-related

memory (23). For example, leptin and GLP-1 act in the HPCv (CA1v subregion) to decrease

food intake and food-motivated conditioned behaviors (24-26), whereas the orexigenic gut-

derived hormone ghrelin administered to the HPCv but not HPCd has opposite effects (23, 27).

Olfactory information, which is intimately connected with feeding behavior, is also preferentially

processed within the HPCv compared with the HPCd. The CA1v, specifically, is bidirectionally

connected to brain regions that process olfactory information (7, 28, 29), and CA1v neurons

respond more robustly to olfactory contextual cues compared with CA1d (8). Further, ghrelin

signaling in the CA1v improves olfactory- and socially-mediated memory for food preference

(30). Given the HPCv appetite-relevant neuroanatomical connections, endocrine and

neuropeptide receptor expression profiles, and functional evidence linking this subregion with

food-motivated behaviors, we hypothesize that HPCv mediation of visuospatial memory is

preferentially biased to food-reinforced foraging behavior.

HPCv pyramidal neurons have extensive projection targets throughout the brain (31), yet

the functional relevance of these pathways is poorly understood. Recently, a vHPC dentate

gyrus-CA3v to lateral septum (LS) pathway was identified that suppresses feeding (32).

Similarly, CA1v projections to the medial prefrontal cortex (mPFC) and lateral hypothalamic

area (LHA) mediate feeding-related outcomes associated with GLP-1 and ghrelin signaling,

respectively. CA1v neurons also robustly target the LS (33, 34), however, the relevance of this

pathway to feeding behavior and/or feeding-related memory is unknown. It is feasible that CA1v

to LS projections participate in regulating spatial memory for food location, a notion supported

by findings showing that neuroplastic changes occur in the LS after learning a spatial memory

task (35, 36). Thus, in addition to exploring the role of the HPCv in memory for the spatial

location of food, the present study also investigated the specific role of the CA1v to LS pathway

in foraging-relevant memory processes.

To systematically investigate the role of the HPCv and CA1v to LS signaling in

visuospatial learning and memory for food reinforcement, we developed a novel appetitive

reinforcement-based spatial foraging behavioral task that allows for direct comparison with an

aversive reinforcement-based task of similar difficulty that uses the same apparatus and spatial

cues. Performance in these tasks was assessed following pathway-specific dual viral-based

reversible (chemogenetic inhibition) or chronic (targeted apoptosis) disconnection of the CA1v

to LS pathway. To further expand neural network knowledge on CA1v to LS signaling, we used

conditional viral-based neuroanatomical tracing strategies to identify both first-order collateral

and second-order targets of LS-projecting CA1v neurons. Collective results from the present

study identify novel neural circuitry of specific relevance to foraging behavior.

RESULTS

The ventral hippocampus (HPCv) is required for remembering the spatial location of food

To examine the importance of the HPCv in memory for the spatial location of food,

animals received bilateral *N*-Methyl-D-aspartate excitotoxic lesions of the HPCv (HPCv lesion

n=12) or bilateral sham injections (control n=12) (histological analyses for the neuron-specific

NeuN antigen in Fig. 1A) and were tested rats in a novel appetitive visuospatial memory task

developed in our lab. Results revealed no significant differences in errors (incorrect hole

investigations) (Fig. 1E) or latency to locate the food source hole (Fig. 1F) during training.

However, memory probe results show that animals with HPCv lesions decreased the ratio of

correct + adjacent / total holes explored in the first minute compared with controls (Figure 1G;

$p < 0.05$, entire two minutes in Figure 1H). Post-surgical analyses of food intake (Supp. Fig. 1A)

and body weight (Supp. Fig. 1B) found no group differences in these measures over time. These

collective results indicate the HPCv is required for memory retention but not learning of the

spatial location of food in the environment.

Ventral hippocampus CA1 (CA1v) projections to the lateral septum (LS) mediate

appetitive spatial memory for food location, but not aversive spatial memory for escape

location

136   A major neuroanatomical target of HPCv CA1 neurons (CA1v) is the LS (34). We

investigated the functional relevance of the CA1v to LS signaling in learning and recalling the

spatial location of food reinforcement using conditional dual viral approaches to either reversibly

(via cre-dependent pathway-specific inhibitory chemogenetics; diagram of approach in Fig. 2A)

or permanently (via cre-dependent pathway-specific caspase-induced lesions; diagram of

approach in Fig. 2B) disconnect CA1v to LS communication. Results from the appetitive spatial

memory task reveal no significant group differences during training in either errors before

locating the correct hole (Fig. 2E) or latency to locate the food source hole (Fig. 2F). However,

memory probe results demonstrate that both acute (pathway-specific DREADDs) and chronic

(pathway-specific caspase) CA1v to LS disconnection decreased the ratio of correct + adjacent /

total holes explored in the entire two minutes of the probe compared with controls (Fig. 2G;

$p < 0.05$), with a similar trend in the first minute of the probe (Fig. 2H; $p = 0.059$). Histological

analyses confirmed successful viral transfection of LS-projecting CA1v neurons with DREADDs

(Figs. 2C&D) or caspase (Supp. Figs. 2E&F), and post-surgical analyses of food intake (Supp.

Fig 2A) and body weight (Supp. Fig 2B) found no group differences in these measures over time.

These data demonstrate that CA1v to LS communication is critical for remembering the spatial

location of food, and that these effects are unlikely to be based on differences in energy status or

food motivation.

154   To evaluate whether the CA1v to LS pathway is specifically involved in food-related

spatial memory vs. spatial memory in general, in a separate cohort of animals we tested the effect

of reversible and chronic CA1v to LS disconnection in a spatial learning task based on aversive

reinforcement rather than food. Importantly, this task uses the same apparatus and visuospatial

cues as the spatial location food memory task described above, but the animals are motivated to

locate the tunnel to escape mildly aversive stimuli (bright lights, loud noise) with no food

reinforcement in the tunnel (see Fig. 1B). Training results revealed no significant group

differences in errors before correct hole (Fig. 2I) nor latency to locate the escape hole (Fig. 2J).

During the memory probe test, there were no group differences in the ratio of correct + adjacent /

total holes investigated during the first minute (Fig. 2K) nor the entire two minutes (Fig. 2L) of

the memory probe. Histological analyses confirmed successful viral transfection of LS-

projecting CA1v neurons with DREADDs (Figs. 2C&D) or caspase (Supp. Figs. 2E&F), and

post-surgical analyses of food intake (Supp. Fig. 2C) and body weight (Supp. Fig. 2D) found no

group differences in these measures over time. These collective findings suggest that CA1v to

LS signaling specifically mediates spatial memory in a reinforcement-dependent manner.

CA1v to LS signaling does not participate in HPC-dependent nonspatial appetitive

memory, anxiety-like behavior, or levels of locomotor activity

To examine the role of CA1v to LS signaling in a nonspatial food-reinforced memory

task, we tested the effect of reversible and chronic CA1v to LS disconnection in the social

transmission of food preference (STFP) test (diagram in Fig. 3A). The HPCv plays an important

role in STFP (30, 37, 38), which tests socially-mediated food-related memory based on previous

exposure to socially-communicated olfactory cues. Results revealed no differences in preference

ratio between groups (Fig. 3B), suggesting that neither acute nor chronic CA1v to LS

disconnection impair food-related memory based on social-based olfactory stimuli.

Hippocampal-septal circuitry has been shown to play a role in mediating anxiety and

stress behavior (39-41). Thus, we sought to confirm whether the observed foraging-related

memory impairments following CA1v-LS disconnection may be based, in part, on non-specific

behavioral effects of the disconnection procedures unrelated to memory, including anxiety-like

behavior (zero maze task; diagram of apparatus in Fig. 3C) and general locomotor activity (open

field test). Results showed no significant group differences in time spent in the open zones in the

zero maze test (Fig. 3D) nor in the number of open zone entries (Fig. 3E), suggesting that neither

chronic nor reversible disconnection of the CA1v to LS circuitry influences anxiety-like

behavior. In addition, CA1v to LS disconnection had no effect on general locomotor activity in

the open field test (Fig. 3F). Overall these results indicate that that disconnection of CA1v to LS

signaling does not affect nonspatial HPC-dependent appetitive memory in STFP, general

locomotor activity, anxiety-like behavior, or locomotor activity.

The medial prefrontal cortex is a collateral target of LS-projecting CA1v neurons

In addition to the LS, CA1v neurons also robustly project to the medial prefrontal cortex

(mPFC) and the lateral hypothalamic area, two pathways that we have previously shown to be

involved in feeding behavior (25, 27, 42). To examine whether LS-projecting CA1v neurons

have collateral targets in the mPFC, LHA, or in other brain regions, we utilized a conditional

dual viral neural pathway tracing approach that identifies collateral targets of a specific

monosynaptic pathway (CA1v->LS; diagram of approach in Fig. 4A, representative CA1v

injection site in Fig. 4B, representative CA1v injection site in Fig. 4C). Results revealed that LS-

projecting CA1v neurons also project to the mPFC (Fig. 4D), whereas minimal collateral

labeling was observed in the LHA or in other brain regions (data not shown). Thus, it may be the

case that the impaired spatial memory for food location observed following either reversible or

chronic CA1v to LS disconnection are based, in part, on CA1v to mPFC signaling from the same

CA1v to LS projecting neurons.

Neither CA1v to mPFC nor CA1v to LHA signaling contribute to spatial memory for food

location

To test the functional relevance of the CA1v neural pathways that either do (CA1v to

mPFC) or do not (CA1v to LHA) collateralize from CA1 to LS projections in food-directed

spatial learning and memory, we took advantage of the exclusively ipsilateral projections from

CA1v to mPFC (25) (and to LHA (42)) and used a 'contralesional' approach to functionally

disconnect the CA1v to mPFC (or LHA) pathway (diagram of approach in Fig. 4E), while

leaving CA1v and mPFC (or LHA) communication to other brain regions intact. Post-surgical

analyses of food intake (Supp. Fig. 3A,C) and body weight (Supp. Fig. 3B,D) found no group

differences in these measures over time. Training results from the appetitive spatial memory task

revealed no group differences in errors before locating the correct hole (Fig. 4F, 4L) or latency to

locate the food-baited hole (Fig. 4G, 4L) for either the CA1v-mPFC or the CA1v-LHA

disconnection groups. Memory probe results demonstrate that neither CA1v to mPFC nor CA1v

to LHA disconnection altered the ratio of correct + adjacent / total holes explored in the first

minute (Fig. 4H, 4M) or the entire two minutes (Fig. 4I, 4N) compared with controls. These data

collectively demonstrate that CA1v to LS mediation of foraging-related memory does not require

collateral projections to the mPFC neural pathway. Further, CA1v to LHA neural signaling,

although important for other food-related conditioned appetitive behaviors (27, 42), is not

required for either learning or remembering of the spatial location of food.

Second-order targets of CA1v to LS neurons include the mPFC, the LHA, and the ACB

227   We used a dual viral tracing strategy (diagram of approach in Fig. 5A; representative LS

injection site in Fig. 5B) to identify downstream targets of LS-projecting CA1v neurons. Results

revealed that the mPFC (Fig. 5C), the LHA (Fig. 5D) and the nucleus accumbens (ACB; Fig. 5E)

are among the strongest second-order targets of the CA1v to LS-projecting neurons. Quantitative

analyses using a custom built data-entry platform (Axiome C, created by JDH) are summarized

graphically for a representative animal on a brain flatmap summary diagram (Fig. 5F) for the

hemisphere ipsilateral (top) and contralateral (bottom) to the injection sites. The data are also

summarized in tabular form in Supplementary Table 1.

DISCUSSION

237   Memory for the physical location of a food source is adaptive for maintaining adequate

energy supply for reproduction and survival. However, the neural circuits mediating this

complex behavior are not well understood as research on visuospatial memory has

predominantly used tasks with either aversive or neutral/passive reinforcement. The present

study identified a specific monosynaptic CA1v to LS pathway as a necessary substrate for food-

motivated spatial memory using a newly developed foraging task. Moreover, the selectivity of

the CA1v-LS pathway in mediating spatial memory for food location is supported by results

showing that disconnection of this pathway did not affect performance in an escape-motivated

spatial memory task of comparable difficulty conducted in the same apparatus, anxiety-like

behavior, or olfactory and social-based appetitive memory. Viral pathway tracing identified that

LS-projecting CA1v neurons also send collateral projections to the mPFC, however, functional

disconnection of CA1v-mPFC signaling did not impair spatial memory for food location.

Utilization of an anterograde multisynaptic viral tracing approach and quantitative forebrain-
wide axon mapping analyses revealed that the CA1v-LS pathway sends second-order projections
to various feeding and reward-relevant neural substrates such as the LHA, ACB, and mPFC.
Collectively, these data establish monosynaptic CA1v to LS signaling as essential for spatial
memory-based foraging behavior, and we further identify a neural network of interconnected
feeding-relevant substrates that are collateral and/or downstream targets of this pathway.

Historically, the hippocampus has been divided into the dorsal and ventral subregions
that have both distinct and overlapping functions, anatomical connections, and genetic and
receptor expression profiles (12, 13, 17, 31, 43). The HPCd has been predominantly implicated
in facilitating spatial memory, whereas the HPCv is linked with stress responses, affective
behavior, and energy balance. However, the HPCv also plays a role in spatial memory,
particularly for goal-directed navigation. Consistent with present results, a recent study found
that both the HPCd and HPCv were critical for food reward-directed spatial navigation in an
obstacle-rich complex environment (44). Under some testing conditions, spatial learning and
memory in an aversive reinforcement-based water maze also requires both subregions of the
HPC (45). Similarly, whole-brain analysis of blood flow in rats during retrieval of spatial
memory in an aversive reinforcement-based Barnes maze task revealed increased activation in
the CA1d and CA1-3v in trained animals compared to controls (46). These results indicate that
spatial processing in the hippocampus is organized longitudinally, and that both HPCd and HPCv
are important for goal-directed spatial navigation. Here we extend these findings by revealing
that the HPCv, and more specifically, CA1v to LS signaling, is preferentially biased to
processing food-reinforcement based spatial memory.

Septohippocampal circuitry, which includes extensive connections between the
hippocampus (HPCd and HPCv) and septal nuclei (LS and medial), has been studied in the
context of a variety of behaviors, including, memory, stress and anxiety. HPCv-LS connections,
in particular, are linked with anxiety-like behavior. Transient pharmacological inactivation of
ipsilateral projections from the HPCv to LS suppresses anxiety-like behavior in an elevated plus
maze task (40), while bilateral DREADDs-mediated manipulation of bilateral HPCv to LS
projections during the same task reveal distinct anxiogenic and anxiolytic populations of neurons
(39). These findings suggest that hippocampal effects on anxiety may be mediated by
functionally distinct populations in the HPCv that project to the LS, and further, are potentially

dependent on laterality. However, these studies differ from the present study in that both CA1v

and CA3v subregions of the HPCv were targeted. Though we did not find changes in anxiety-

like behavior after reversible of chronic disconnection of CA1v to LS, the inclusion of CA3v in

the work described above prevents direct comparisons to our own results. Overall, these studies

combined with present results are consistent with a framework that overlapping and divergent

functions of the HPCv to LS signaling in both feeding and emotional regulation are important for

modulating complex motivated behavior in accordance with cognitive input (47).

287   Present results identify the CA1v-LS pathway in coordinating food-motivated behavior

with learned spatial information. While the use of aversive reinforcement-based spatial memory

tasks are predominant, the radial arm maze has been used to investigate food-motivated spatial

reference memory (analogous to the spatial memory probe in the present study) and spatial

working memory using multiple consistently food-baited arms (48). Inconsistent with the present

results, a previous study utilizing a four-baited/four-unbaited version of the eight-arm radial

maze task found that NMDA-mediated lesions of the HPCd, but not HPCv, impair spatial

reference and working memory (49). Another study using the same working memory task

reported that complete HPC lesions impaired both spatial and working memory, whereas lesions

to either the HPCd, HPCv, or intermediate HPC had no effect on memory performance (50).

Future studies will need to address whether discrepancies between the present results and these

studies are based on the differences between the behavioral tasks and/or differences in loss of

function methodologies. With regards to the latter, we note that our chemogenetic and lesion

approaches offered far greater specificity (LS-projecting CA1v neurons vs. whole HPCv)

compared with these studies.

302   Our multisynaptic neuroanatomical results offer a framework for future work to examine

the broader anatomical circuitry involved in foraging-related behaviors. Utilizing dual viral

approaches and systematic forebrain-wide quantitative mapping analyses, our findings confirm

that the CA1v-LS pathway projects downstream to other neural substrates involved in feeding

and motivated behavior, including the LHA, ACB, and mPFC. Given that the LS-LHA pathway

has been shown to affect food intake (51), and that metabolic activity in the LS and ACB is

associated food anticipatory activity in rabbit pups (52), it is possible that the LHA and/or the

ACB are functional downstream targets of CA1v-LS signaling for coordinating foraging

behavior. It is unlikely, however, that the mPFC is a functional target of CA1v-LS signaling for

appetitive spatial memory control. For example, our contralesional CA1v-mPFC disconnection approach, which did not affect appetitive spatial memory, not only disconnects all ipsilateral direct HPCv to mPFC communication, but also all ipsilateral multisynaptic HPCv to mPFC connections (including via LS). Consistent with this hypothesis, anterograde labeling downstream of CA1v to LS projections was not observed in the contralateral mPFC. It is possible, however, that CA1v → LS → mPFC signaling is relevant to other motivated behaviors. While we examined food-motivated and escape-motivated spatial memory, functional analyses of this circuit may be extended to drug-motivated, social-motivated, or water-motivated spatial memory tasks.

Collectively, we identify a CA1v to LS pathway involved in foraging-related, but not escape-motivated, spatial memory. Furthermore, the selectivity of this pathway to appetitive spatial memory is supported by data showing that neither chronic nor reversible disruption of CA1v to LS signaling influenced various other behavioral outcomes. We also systematically characterized collateral and second-order projections of this pathway, which demonstrate overlapping pathways coordinated in the higher-order control of energy balance. These data shed light on the neural systems underlying complex motivated behaviors that require functional connections of cognitive and feeding-relevant substrates.

METHODS

Animals

Adult male Sprague–Dawley rats (Envigo; 250-275g on arrival) were individually housed in hanging wire cages with *ad libitum* access (except where noted) to water and chow (LabDiet 5001, LabDiet, St. Louis, MO) on a 12h:12h reverse light/dark cycle. All procedures were approved by the University of Southern California Institute of Animal Care and Use Committee.

General intracranial injection procedures

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine 90mg/kg body weight [BW], xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW) followed by a pre-operative, subcutaneous injection of analgesic (ketoprofen, 5mg/kg BW). Post-operative analgesic (subcutaneous injection of ketoprofen, 5mg/kg BW) was administered once

per day for 3 days following surgery. The surgical site was shaved and prepped with iodine and

ethanol swabs, and animals were placed in a stereotaxic apparatus for stereotaxic injections.

NMDA or viruses were delivered using a microinfusion pump (Harvard Apparatus, Cambridge,

MA, USA) connected to a 33-gauge microsyringe injector attached to a PE20 catheter and

Hamilton syringe. Flow rate was calibrated and set to 83.3nl/sec. Injectors were left in place for

2min post-injection to allow for complete delivery of the infusate. Specific viruses/drugs,

coordinates, and injection volumes for procedures are detailed below. Following the completion

of all injections, incision sites were closed using either surgical staples, or in the case of

subsequent placement of an indwelling cannula, simple interrupted sutures. Upon recovery from

anesthesia and return to dorsal recumbency, animals were returned to the home cage. All

behavioral procedures occurred 21 days after injections to allow for complete transduction and

expression of the viruses, or complete lesioning drugs. General intracranial injection procedures

were followed for all injection procedures below.

Chronic lesions of the HPCv

Lesion animals received bilateral excitotoxic HPCv lesions via intracranial injections of

N-methyl-d-aspartate (NMDA; 200nL per hemisphere) at the following coordinates at three

different sites along the rostrocaudal extent of the HPCv (53): [1] -4.8mm AP, +/-5.0mm ML, -

7.5mm DV, [2] -5.5mm AP, +/-4.5mm ML, -7.0mm DV, and [3] -6.1mm AP, +/-4.8mm ML, -

7.0mm DV with control animals receiving vehicle saline in the same location. The reference

points for AP and ML coordinates were defined at bregma, and the reference point for the DV

coordinate was defined at the skull surface at the target site.

Bilateral HPCv lesion brains were histologically evaluated for the correct placement of

lesions in 1 out of 5 series of brain tissue sections. Neurons were visualized using

immunohistochemistry for the neuron specific antigen NeuN (see Immunohistochemistry), with

a complete lesion indicated by a lack of neuronal staining in the HPCv target region. Here, n=0

animals were removed. Representative images in Fig. 1A.

Acute and chronic disconnection of the CA1v to LS neural pathway

Cre-dependent dual viral strategies were used to generate the following groups: [1] acute

chemogenetic disconnection of the CA1v to LS neural pathway (DREADDs; diagram of

approach in Fig. 2A), [2] chronic disconnection of the CA1v to LS neural pathway (caspase
lesion-induced; diagram of approach in Fig. 2B), and [3] a common/shared control group for
these two disconnection strategies.

Regardless of group, all animals received a bilateral AAV retro-cre injection in the LS
(AAV2[retro]-hSYN1-EGFP-2A- iCre-WPRE; 200nL per side) at the following stereotaxic
coordinates (53): +0.84mm AP, +/-0.5mm ML, −4.8mm DV. The reference points for all LS
coordinates were defined at bregma.

According to experimental group, animals received a different virus delivered to the
CA1v subregion of the HPCv. All viruses were administered to the CA1v at the following
stereotaxic coordinates (53): −4.9mm AP, +/-4.8mm ML, −7.8mm DV. The reference points for
AP and ML coordinates were defined at bregma, and the reference point for the DV coordinate
was defined at the skull surface at the target site.

[1] DREADDs group for reversible inactivation of CA1v to LS: To allow reversible
chemogenetic inactivation of the CA1v to LS neural pathway, one group of animals received a
bilateral CA1v injection of a cre-dependent virus to drive expression of inhibitory designer
receptors activated by designer drugs (DREADDs), (AAV-Flex-hm4Di-tdTomato). This dual
viral strategy drives expression of inhibitory DREADDs exclusively in LS-projecting CA1v
neurons, which enables acute inactivation of these neurons by injection of the DREADDs ligand,
clozapine-N-oxide (CNO), at the time of behavioral testing.

[2] Caspase group for chronic inactivation of CA1v to LS: To allow chronic
disconnection of the CA1v to LS neural pathway, a second group of animals received a bilateral
CA1v cre-dependent caspase virus (AAV1-Flex-taCasp3-TEVp; 200nL per side) mixed with a
cre-dependent reporter virus (AAV-flex-tdTomato; 200nL per side) for histological verification.
This dual viral strategy drives expression of the apoptosis-mediator molecule caspase exclusively
in LS-projecting CA1v neurons, which induces apoptotic cell death in these neurons while
leaving other CA1v neurons intact.

[3] Common control: A common control group was generated for both of these
experimental groups, with this group receiving a cre-dependent control AAV in the CA1v
(AAV-flex-tdTomato).

Immediately following viral injections, all animals were surgically implanted with a
unilateral indwelling intracerebroventricular (ICV) cannula (26-gauge, Plastics One, Roanoke,

VA) targeting the lateral ventricle (VL). Cannulae were implanted and affixed to the skull with jeweler's screws and dental cement at the following stereotaxic coordinates: −0.9mm AP, +1.8mm ML, −2.6mm DV. The reference points for AP and ML coordinates were defined at bregma, and the reference point for the DV coordinate was defined at the skull surface at the target site. Placement for the VL cannula was verified by elevation of blood glucose resulting from an injection of 210μg (2μL) of 5-thio-D-glucose (5TG) using an injector that extended 2mm beyond the end of the guide cannula (54). A post-injection elevation of at least 100% of baseline glycemia was required for subject inclusion. Animals that did not pass the 5TG test were retested with an injector that extended 2.5mm beyond the end of the guide cannula and, upon passing the 5TG test, were subsequently injected using a 2.5mm injector instead of a 2mm injector for the remainder of the study. Prior to behavioral testing where noted, all animals received an ICV 18mmol infusion (2uL total volume) of the DREADDs ligand clozapine N-oxide (CNO) (55), rendering only DREADDs animals chemogenetically inactivated. CNO injections were hand delivered using a 33-gauge microsyringe injector attached to a PE20 catheter and Hamilton syringe through the indwelling guide cannulae. Injectors were left in place for 30sec to allow for complete delivery of the CNO.

Cre-dependent DREADD expression targeting LS-projecting neurons in CA1v was evaluated based on localization of the fluorescent reporter tdTomato. Immunohistochemistry for red fluorescent protein (RFP) was conducted to amplify the tdTomato signal (see Immunohistochemistry). After histological analysis, animals were removed based on misses (n=3 for cohort 1; n=0 for cohort 2). Representative images in Figs. 2C&D.

Neuronal apoptosis due to activation of cre-dependent caspase targeting LS-projecting neurons in CA1v was evaluated based on reduction of the fluorescent reporter tdTomato due to neuron cell death compared with histological controls. Caspase brains (which received a retro-cre virus in the LS in conjunction with an injection of cre-dependent caspase virus mixed with a cre-dependent virus that drives a tdTomato fluorescent reporter in the CA1v) were compared to histological control brains (that received a retro-cre virus in the LS in conjunction with only a cre-dependent that drives a tdTomato fluorescent reporter in the CA1v, which was equivalently diluted to match the caspase injections). In both groups, immunohistochemistry for red fluorescent protein (RFP) was conducted to amplify the tdTomato signal (see Immunohistochemistry). Confirmation of successful caspase activation in the CA1v was

evaluated by reduced tdTomato fluorescence in comparison to histological control animals.

Animals were removed based on incomplete caspase lesioning (n=0 for cohort 1; n=0 for cohort

2). Representative images in Supplementary Figs 2E&F.

Identification of collateral targets of CA1v to LS projecting neurons

Collateral targets of the CA1v to LS neural pathway were identified using a dual viral

approach (diagram of approach in Fig. 4A) where a retrograde vector was injected into the LS

(AAV2retro-hSyn1-eGFP-2A-iCre-WPRE; 200nL per side; coordinates as above), and a Cre-

dependent anterograde vector (AAV1-CAG-Flex-tdTomato-WPRE-bGH; 200nL per side;

coordinates as above) was injected in the CA1v. This latter injection drives tdTomato transgene

expression in CA1v neurons that project to the LS, which allows for brain-wide analyses of

collateral targets. After 3 weeks of incubation time to allow for complete transduction and

expression of the viruses, brains were collected, immunohistochemically processed, and imaged

as described below.

Contralesional disconnection of the CA1v to mPFC neural pathway or the CA1v to LHA

neural pathway

To functionally disconnect the CA1v to mPFC pathway (diagram of approach in Fig. 4E),

or the CA1v to LHA pathway (diagram of approach in Fig. 4J) lesion animals received a

unilateral excitotoxic CA1v lesion via an intracranial injection of NMDA (200nL) at the

following coordinates (34): −4.9mm AP, + or - 4.8mm ML (left/right counterbalanced to be

contralateral to mPFC or LH lesion within-animal), −7.8mm DV, with control animals receiving

vehicle saline injections in the same location. The reference points for AP and ML coordinates

were defined at bregma, and the reference point for the DV coordinate was defined at the skull

surface at the target site.

In addition to the CA1v lesion, CA1v to mPFC disconnect animals also received a

unilateral excitotoxic mPFC lesion via two intracranial injections of NMDA (100nL per

injection) at the following coordinates (53): [1] +2.7mm AP, + or - 0.7mm ML (left/right

counterbalanced to be contralateral to CA1v lesion within-animal), −5.3mm DV and [2] +3.0mm

AP, + or - 0.7mm ML (left/right counterbalanced to be contralateral to CA1v lesion within-

animal), −4.7mm DV, with control animals receiving vehicle saline in the same location. The

reference points for all mPFC coordinates were defined at bregma.

In addition to the CA1v lesion, CA1v to LHA disconnect animals also received a

unilateral excitotoxic LHA lesion via an intracranial injection of NMDA (100nL per injection) at

the following coordinates (53): +2.9mm AP, + or – 1.1mm ML (counterbalanced to be

contralateral to CA1v lesion within-animal), −8.9mm DV, with control animals receiving vehicle

saline in the same location. The reference points for all LHA coordinates were defined at

bregma.

Contralateral brains were histologically evaluated for the correct placement of lesions by

visualizing neurons using immunohistochemistry for the neuron specific antigen NeuN (see

Immunohistochemistry). A complete lesion was indicated by a lack of neuronal labeling in the

CA1v, mPFC, or LHA target regions as necessary. Here, n=0 animals were removed.

Identification of second order targets of CA1v to LS projecting neurons

To identify second order targets of CA1v to LS-projecting neurons (diagram of approach

in Fig. 5A), animals received a bilateral injection of a transsynaptic Cre-inducing anterograde

vector into CA1v (AAV1-hSyn-Cre-WPRE-hGH; 200nL per side; coordinates as above) that

drives expression of Cre in both first-order (CA1v injection site) and 2^{nd}-order (but not 3^{rd}-

order) neurons via transsynaptic virion release (56, 57). This was combined with a bilateral

injection of a Cre-dependent anterograde vector in the LS (AAV1-CAG-Flex-tdTomato-WPRE-

bGH; 200nL per side, coordinates as above). This latter injection allows for anterograde tracing

from 1^{st}-order LS targets receiving synaptic input from CA1v. After 3 weeks of incubation time

to allow for complete transduction and expression of the viruses, brains were collected,

immunohistochemically processed, and imaged as described below.

Data were entered using a custom built data-entry platform (Axiome C, created by JDH)

built around Microsoft Excel software and designed to facilitate entry of data points for all gray

matter regions across their atlas levels as described in a rat brain reference atlas (Brain Maps 4.0,

Swanson, 2018). The Axiome C approach was used previously to facilitate the analysis of brain

gene-expression data (Hahn et al., 2019). An ordinal scale, ranging from 0 (absent) to 7 (very

strong), was used to record the qualitative weight of anterograde labeling. An average value was

then obtained for each region across its atlas levels for which data were available. These data are

summarized graphically for a representative animal on a brain flatmap summary diagram

(adapted from (58).

Food intake and body weight

For the day prior to surgery (day 0) and for two weeks thereafter, 24h chow intake was

measured daily just prior to dark cycle onset to determine effects of experimental procedures on

food intake. Spillage was accounted for daily by collecting crumbs onto Techboard paper placed

under the cages of each animal. Additionally, animals were weighed daily just prior to dark cycle

onset to determine effects of experimental procedures on body weight.

Appetitive spatial memory foraging task

To test visuospatial learning and memory for food reinforcement, we developed a novel

spatial foraging task modified from the traditional Barnes maze procedure. Throughout this

paradigm, animals were maintained at 85% free-feeding body weight. The paradigm involves an

elevated gray circular platform (diameter: 122cm, height: 140 cm) consisting of 18 uniform holes

(9.5cm diameter) spaced every twenty degrees around the outside edge. Under one of the holes is

a hidden tunnel (38.73cm L x 11.43cm W x 7.62cm D and a 5.08cm step). Surrounding the table

are distinct spatial cues on the wall (e.g. holiday lights, colorful shapes, stuffed unicorn) that are

readily visible to the animal. In contrast to the traditional Barnes Maze where an animal uses the

spatial cues to escape mildly aversive stimuli in the environment (e.g. bright light and loud

sound), this task utilizes food as motivation, such that each hidden tunnel contained five 45mg

sucrose pellets (Bio-Serv, Flemington, NJ). Additionally, a quiet white noise (60dB) was used to

drown out background noise and floor lamps were used for low-level ambient lighting. On the

first day, each animal underwent a habituation session consisting of 1min inside a transport bin

under the table, 2min underneath the start box in the center of the table, and 3min inside the

hidden tunnel with 5 sucrose pellets. During training, each rat was assigned a specific escape

hole according to the position of the spatial cues with the location counterbalanced across

groups. Animals were trained over the course of two trials per day for four days (five days in

contralesional experiments) to learn to use the spatial cues in order to locate the correct hole with

the hidden tunnel with sucrose pellets. Learning during training was scored via animal head

location tracking by AnyMaze Behavior Tracking Software (Stoelting Co., Wood Dale, IL). The

incorrect hole investigations prior to finding the correct hole with sucrose pellets ("errors before

correct hole") as well as time to finding the correct hole ("latency to correct hole") were

calculated as an average of the two trials per day, and examined across days of training. After the

conclusion of training, rats had a two-day break where no procedures occurred, then were tested

in a single 2min memory probe in which the hidden tunnel and sucrose pellets were been

removed. All animals received an ICV 18mmol infusion of CNO (2uL total volume) 1h prior to

the memory probe, rendering only DREADDs animals chemogenetically inactivated by CNO. In

the memory probe, the ratio of the correct hole plus adjacent hole investigations over the total

number of hole investigations were calculated via animal head location tracking by AnyMaze

Behavior Tracking Software (Stoelting, Wood Dale, IL). The correct holes plus adjacent over

total were compared between groups for the first minute and entire two minutes of the probe for

all tests. A diagram of the apparatus used for the spatial foraging task is included in Fig. 1B.

Aversive spatial memory escape task (Barnes maze)

To test visuospatial learning and memory for escape reinforcement, we used a modified

traditional Barnes maze procedure, which is a visuospatial-based escape task. Procedures were

exactly the same as above (Appetitive spatial memory foraging task, using the same apparatus, in

the same room, and with the same visuospatial cues) aside from the omission of the sucrose

pellets in the hidden tunnel, the presence of mildly aversive bright (120W) overhead lighting

instead of dim ambient lighting, and a mildly aversive loud white noise (75dB) instead of a quiet

white noise (60dB). This allowed us to test spatial learning and memory motivated by escape

from aversive stimuli in a nearly-identical procedure to our spatial foraging test for learning and

memory motivated by palatable food consumption. A diagram of the apparatus used for the

spatial escape task is included in Fig. 1B.

Social transmission of food preference (STFP)

To examine food-related memory based on social- and olfactory-based cues, we utilized

the social transmission of food preference (STFP) task and adapted protocols from (30, 38, 59,

60). Briefly, untreated normal adult rats are designated as 'Demonstrators', while experimental

groups are designated as 'Observers'. Demonstrators and Observers are habituated to a powdered

rodent chow [LabDiet 5001 (ground pellets), LabDiet, St. Louis, MO] overnight. 24h later,

Observers are individually paired with demonstrators and habituated to social interaction, where

rat pairs are placed in a social interaction arena (23.5cm W × 44.45cm L × 27cm H clear plastic

bin with Sani-chip bedding) and allowed to interact for 30min. Both Observers and

Demonstrators are returned to their home cages and food is withheld for 23hr prior to the social

interaction. For the social interaction, Demonstrators are given the opportunity to consume one

of two flavors of powdered chow (flavored with 2% marjoram or 0.5% thyme; counterbalanced

according to group assignments) for 30min in a room separate from Observers. Our pilot studies

and previous published work (60, 61) show that rats equally prefer these flavors of chow. The

Demonstrator rat is then placed in the social interaction arena with the Observer rat, and the pairs

are allowed to socially interact for 30min. Observers are then returned to their home cage and

allowed to eat *ad libitum* for 1h and then food is removed. The following day, the 23h food-

deprived Observer animals are given a home cage food preference test for either the flavor of

chow paired with the Demonstrator animal, or a novel, unpaired flavor of chow that is a flavor

that was not given to the Demonstrator animal (2% marjoram vs. 0.5% thyme; counterbalanced

according to group assignments). All animals received an ICV 18mmol infusion of CNO (2uL

total volume) 1h prior to the social interaction session, rendering only DREADDs animals

chemogenetically inactivated by CNO. Food intake (1h) was recorded with spillage accounted

for by collecting crumbs onto Techboard paper that is placed under the cages of each animal.

The % preference for the paired flavor is calculated as: 100*Demonstrator-paired flavored chow

intake/Demonstrator + Novel flavored chow intake. In this procedure, normal untreated animals

learn to prefer the Demonstrator paired flavor based on social interaction and smelling the breath

of the Demonstrator rat (30, 38, 59-61). A diagram of the STFP procedure is included in Fig. 3E.

Zero maze

The zero maze behavioral paradigm was used to evaluate anxiety-like behavior. The zero

maze apparatus used was an elevated circular track, divided into four equal length sections. Two

zones were open with 3 cm high curbs ('open zones'), whereas the two other zones were closed

with 17.5 cm high walls ('closed zones'). All animals received an ICV 18mmol infusion of CNO

(2uL total volume) 1h prior to testing, rendering only DREADDs animals chemogenetically

inactivated by CNO. Behavioral testing was performed during the light cycle. Animals were

placed in the maze for a single, 5min trial in which the location of the center of the animal's

body was measured by AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). The

apparatus was cleaned with 10% ethanol in between animals. During the trial, the number of

open zone entries and total time spent in open sections (defined as body center in open sections)

were measured, which are each indicators of anxiety-like behavior in this procedure. A diagram

of the zero maze apparatus is included in Fig. 3G.

Open field

An open field test was used to evaluate general levels of locomotor activity. The

apparatus used for the open field test was an opaque gray plastic bin (60cm × 56cm), which was

positioned on a flat table in an isolated room with a camera directly above the center of the

apparatus. Desk lamps were positioned to provide indirect lighting to all corners of the maze

such that the lighting in the box uniformly measured 30 lux throughout. All animals received an

ICV 18mmol infusion of CNO (2uL total volume) 1h prior to testing rendering only DREADDs

animals chemogenetically inactivated by CNO. Behavioral testing began at dark onset. At the

start of the 10min test, each animal was placed in the open field apparatus in the same corner

facing the center of the maze. The location of the center of the animal's body was measured with

the AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). Total distance traveled

was measured by tracking movement from the center of the animal's body throughout the test.

Immunohistochemistry

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine

90mg/kg BW xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW) then transcardially

perfused with 0.9% sterile saline (pH 7.4) followed by 4% paraformaldehyde (PFA) in 0.1M

borate buffer (pH 9.5; PFA). Brains were dissected from the skull and post-fixed in PFA with

15% sucrose for 24h, then flash frozen in isopentane cooled in dry ice. Brains were sectioned to

30μm thickness on a freezing microtome. Sections were collected in 5-series and stored in

antifreeze solution at −20°C until further processing. General fluorescence IHC labeling

procedures were performed as follows. The antibodies and dilutions that were used are as

follows: [1] For lesion histology using the neuron-specific protein NeuN, the rabbit anti-NeuN

primary antibody (1:1000, Abcam) was used followed by a donkey anti-rabbit conjugated to

AF488 (1:500, Jackson Immunoresearch). [2] To amplify the native tdTomato signal for

neuroanatomical tracing or DREADDs histology, the rabbit anti-RFP primary antibody (1:2000,

Rockland) was used followed by a donkey anti-rabbit conjugated to Cy3 (1:500, Jackson

Immunoresearch). [3] To amplify the native GFP signal for LS injection site histology, the

chicken anti-GFP primary antibody (1:500, Abcam) was used followed by a donkey anti-chicken

secondary antibody conjugated to AF488 (1:500, Jackson Immunoresearch). Antibodies were

prepared in 0.02M potassium phosphate buffered saline (KPBS) solution containing 0.2% bovine

serum albumin and 0.3% Triton X-100 at 4 °C overnight. After thorough washing with 0.02M

KPBS, sections were incubated in secondary antibody solution. All secondary antibodies were

obtained from Jackson Immunoresearch and used at 1:500 dilution at 4°C, with overnight

incubations (Jackson Immunoresearch; West Grove, PA, USA). Sections were mounted and

coverslipped using 50% glycerol in 0.02 M KPBS and the edges were sealed with clear nail

polish. Photomicrographs were acquired using a Nikon 80i (Nikon DSQI1,1280X1024

resolution, 1.45 megapixel) under epifluorescence or darkfield illumination.

Axon mapping

Data were entered using a custom built data-entry platform (Axiome C, created by JDH)

built around Microsoft Excel software and designed to facilitate entry of data points for all gray

matter regions across their atlas levels as described in a rat brain reference atlas (53). The

Axiome C approach was used previously to facilitate the analysis of brain gene-expression data

(62). An ordinal scale, ranging from 0 (absent) to 7 (very strong), was used to record the

qualitative weight of anterograde labeling. An average value was then obtained for each region

across its atlas levels for which data were available. The data were summarized graphically for a

representative experiment on a brain flatmap summary diagram (adapted from (58)). The data are

also summarized in tabular form in Supplementary Table 1.

Statistics

Data are expressed as mean +/- SEM. Differences were considered to be statistically

significant at p<0.05. All variables were analyzed using the advanced analytics software package

Statistica (StatSoft, Tulsa, OK, USA). For all measures of food intake, body weight, and

errors/latency during spatial foraging and spatial escape task training, differences between

groups were evaluated using two-way repeated measures ANOVAs (treatment x time). For

lesion experiments, measures during spatial foraging and spatial escape task probes, differences

between groups were evaluated using two-tailed independent two-sample Student's t-tests. For

CA1v to LS disconnection experiments, measures during spatial foraging and spatial escape task

probes, the STFP paradigm, the zero maze paradigm, and the open field test, differences between

groups were evaluated using one-way ANOVAs. Significant ANOVAs were analyzed with a

Fisher's LSD posthoc test where appropriate. Outliers were identified as being more extreme

than the median +/- 1.5 * interquartile range. For all experiments, assumptions of normality,

homogeneity of variance (HOV), and independence were met where required.

CONFLICT OF INTEREST

No conflicts of interest.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Diabetes and Digestive and Kidney Diseases

grants: DK104897 to SEK, DK118944 to CML, and DK116558 to ANS. Clozapine-N-Oxide

was kindly provided by the National Institute of Mental Health. The authors are grateful to the

Kanoski lab undergraduates for their assistance in behavioral experiments and histology.

Figure 1

A NeuN (lesion) — CA3v, CA1v, DG, alv

NeuN (sham) — CA3v, CA1v, DG, alv

E Spatial foraging training (errors)
Average errors vs Day 1, Day 2, Day 3, Day 4
Control, HPCv lesion

F Spatial foraging probe: Training errors
Average errors vs Day 1, Day 2, Day 3, Day 4
Control, HPCv lesion

Spatial foraging training (latency)
Average latency (s) vs Day 1, Day 2, Day 3, Day 4
Control, HPCv lesion

G Spatial foraging probe: Training latency
Average latency (s) vs Day 1, Day 2, Day 3, Day 4
Control, HPCv lesion

Spatial foraging probe: First minute
Correct + adjacent holes / total — Control, HPCv lesion

Memory probe (1st minute)
Correct + adjacent holes / total — Control, HPCv lesion

H Spatial foraging probe: Entire two minutes
Correct + adjacent holes / total — Control, HPCv lesion

Memory probe (2 minutes)
Correct + adjacent holes / total — Control, HPCv lesion

Figure 1. Bilateral HPCv lesions impair spatial memory for food location. Representative
HPCv lesion histology with NeuN immunohistochemistry (A; scale bars 500μm). Spatial
foraging task apparatus (B). Representative navigation paths of a control animal preferentially
investigating correct (filled red) and adjacent holes (outlined orange) during spatial foraging
memory probe (C). Representative navigation path of HPCv lesioned animal during spatial
foraging memory probe (D). Bilateral HPCv lesions did not impair learning of the spatial
foraging task compared to controls, as measured by errors before locating correct hole during
task training (E) and latency to locate correct hole during task training (F). Bilateral HPCv
lesions impaired retention of the spatial foraging task, as measured by the ratio of investigation
of correct plus adjacent holes over total investigated during the first minute of the task ($p<0.05$;
G). There were no group differences observed when evaluated over the entire two minutes of the
task (H). Dotted line indicates chance performance level (0.167). For graphs 1D-G, lesion n = 11,
control n = 18. All values expressed as mean +/- SEM.

685
686

Figure 2

A ICV
ICV CNO inhibits CA1v to LS neurons
LS
Retro-Cre
CA1v
Cre-dependent hM4Di DREADDs

B LS
Retro-Cre
CA1v
Cre-dependent caspase

C CA3v
alv
CA1v
alv
hM4Di-tdTomato

D

E Spatial foraging training (errors)
Average errors
Control
DREADDs
Caspase
Day 1 Day 2 Day 3 Day 4

F Spatial foraging training (latency)
Average latency (s)
Control
DREADDs
Caspase
Day 1 Day 2 Day 3 Day 4

G Memory probe (1st minute)
Correct + adjacent holes / total
p = 0.059
Control DREADDs Caspase

H Memory probe (2 minutes)
Correct + adjacent holes / total
*
a
b
b
Control DREADDs Caspase

I Spatial escape training (errors)
Average errors
Control
DREADDs
Caspase
Day 1 Day 2 Day 3 Day 4

J Spatial escape training (latency)
Average latency (s)
Control
DREADDs
Caspase
Day 1 Day 2 Day 3 Day 4

K Memory probe (1st minute)
Correct + adjacent holes / total
Control DREADDs Caspase

D Memory probe (2 minutes)
Correct + adjacent holes / total
Control DREADD Caspase

Figure 2. Reversible and chronic CA1v to LS neural disconnection impairs spatial memory

for food location but not for escape location. Diagram of dual viral approach using a cre-

dependent inhibitory DREADDs approach to reversibly disconnect CA1v to LS neural pathway

(A). Diagram of dual viral approach using a cre-dependent caspase approach to chronically

disconnect CA1v to LS neural pathway (B). Representative injection site in CA1v demonstrating

LS-projecting neurons infected with inhibitory DREADDs, which simultaneously drives

expression of a fluorescent tdTomato transgene (C, D; scale bars 500μm and 100μm,

respectively). Neither reversible (DREADDs) nor chronic (caspase) disconnection of the CA1v

to LS pathway impaired learning of the spatial foraging task compared to controls, as measured

by errors before correct hole during task training (E) and latency to correct hole during task

training (F). Both reversible and chronic disconnection of the CA1v to LS pathway impaired

retention of the spatial memory foraging task as measured by the ratio of investigation of correct

plus adjacent holes over total investigations during entire two minutes of the task ($p<0.05$; H).

During the first minute of the task, there was a trend toward impaired learning in the

disconnection groups, but this effect did not reach statistical significance ($p=0.059$; G). In

contrast, disconnection of the CA1v to LS pathway either reversibly (DREADDs) or chronically

(caspase) did not impair performance on the spatial escape task. There were no differences in

learning as measured by errors before correct hole during task training (I) and latency to correct

hole during task training (J). Unlike the spatial foraging task, retention of the spatial escape task

was not impaired by reversible nor chronic disconnection of the CA1v to LS pathway (K, L). For

graphs 2E-H (CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8. For

graphs 2I-L (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control n=10.

Dotted line indicates chance performance level (0.167). All values expressed as mean +/- SEM.

Figure 3

A

Day 1

Demonstrators and observers habituate to social interaction

Day 2

Demonstrators consume flavored chow (chow A) prior to social interaction

Demonstrators and observers have a 30min social interaction

Day 3

Food-restricted observers are given a 60min flavor preference test (chow A vs. novel chow B)

B

Social transmission of food preference

C

D

Zero maze: Open zone time

E

Zero maze: Open zone entries

F

Open field

Figure 3. Neither reversible nor chronic CA1v to LS neural disconnection impair spatial
memory for escape location, social transmission of food preference, anxiety-like behavior,
or general locomotor activity levels. Diagram of the social transmission of food preference
(STFP) task (A). Neither reversible nor chronic disconnection of the CA1v to LS pathway impair
STFP learning compared to controls, as measured by a food preference ratio (B), with the dotted
line indicating chance preference level (0.50). Diagram of the zero maze apparatus (C). Anxiety-
like behavior was not influenced by reversible or chronic disconnection of the CA1v to LS
pathway compared to controls, as measured by performance in the zero maze task, specifically
time in open zones (D) and entries into open zones (E). Neither chronic nor reversible CA1v to
LS disconnection affected open field performance compared to controls, as measured by total
distance traveled (F). For graphs 3B, 3D, and 3E (CA1v to LS disconnect cohort 1), DREADDs
n=6, caspase n=10, control n=8. For graph 3F (CA1v to LS disconnect cohort 2), DREADDs
n=8, caspase n=12, control n=10. All values expressed as mean +/- SEM.
725
726

Figure 4

Figure 4. Neither CA1v to mPFC nor CA1v to LHA disconnection impairs spatial memory

for food location. Diagram of dual viral approach to identify collateral targets of the CA1v to

LS neural pathway (A). Representative CA1v injection site from collateral identification

approach (B; scale bar 500μm). Representative LS injection site from collateral identification

approach (C; scale bar 500μm). Representative image collateral axons of the CA1v to LS

pathway located in the mPFC (D; scale bar 50μm). Diagram of contralesional approach to

functionally disconnect the CA1v to mPFC neural pathway (E). Disconnection of the CA1v to

mPFC neural pathway did not influence learning of the spatial foraging task compared to

controls, as measured by errors before correct hole during training (F) and latency to correct hole

during training (G). Disconnection of the CA1v to mPFC pathway did not influence retention of

the spatial foraging task, as measured by during the first minute (H) and entire two minutes (I) of

the memory probe. Diagram of contralesional approach to functionally disconnect the CA1v to

LHA neural pathway (J). Disconnection of the CA1v to LHA neural pathway did not influence

learning of the spatial foraging task compared to controls, as measured by errors before correct

hole during training (K) and latency to correct hole during training (L). Disconnection of the

CA1v to LHA pathway did not influence retention of the spatial foraging task, as measured by

correct plus adjacent holes over total holes investigated during the first minute (M) and entire

two minutes (N) of the memory probe. For graphs 4F-I, CA1v to mPFC disconnect n=12, control

n=12. For graphs 4K-N, CA1v to LHA disconnect n=12, control n=11. All values expressed as

mean +/- SEM.

Figure 5

A LS — Anterograde monosynaptic Cre-dependent AAV1-tdTomato. CA1v — Anterograde transsynaptic AAV2/1-hSyn-Cre

B ccb, VLa, LSc, LSr, LS injection site

C PL, ILA, 2° tdTomato+ fibers

D LHAjd, fx, LHAd, LHAjvd, LHAsfa, LHAjvv, 2° tdTomato+ fibers

E ACBsh, aco, 2° tdTomato+ fibers

F CA1v → LS → multiple regions

Targeted side (right)

Contralateral side (left)

Aggregate ordinal value categories
(6 = strongest; 0 = weakest / absent)

Figure 5. Identification of second-order neural projections downstream of CA1v to LS

projections. Diagram of dual viral approach to identify brain regions that are second-order (2°)

targets of the CA1v to LS neural pathway (A). Representative LS injection site from second

order identification approach (B; scale bar 100μm). Representative image of second-order fibers

of the CA1v to LS pathway within the mPFC (C; scale bar 200μm). Representative image of

second-order fibers of the CA1v to LS pathway within the LHA (D; scale bar 200μm).

Representative image of second-order fibers of the CA1v to LS pathway within the ACB (E;

scale bar 200μm). Summary of the projection targets of LS neurons that receive input from

CA1v (F). The outputs of the right side of LS neurons receiving CA1v input are represented at

the macroscale (gray matter region resolution) on a partial flatmap representation of the rat

forebrain, adapted from (58). Connection weights are represented block colors for each region

following an ordinal scale ranging from weakest (0 = very weak or absent) to strongest (6 =

strong), as there were no 7 (very strong) values. The inset at lower left represents one side of the

brain with the part represented in the upper diagrams outlined in magenta.

SUPPLEMENTARY INFORMATION

Supplemental Figure 1

768   **Supplementary Figure 1. Effect of bilateral HPCv lesions on food intake and body weight.**

769   There were no effects on food intake (A) or body weight (B) of bilateral HPCv lesions compared

770   to controls. Lesion n = 11, control n = 18. All values expressed as mean +/- SEM.

Supplemental Figure 2

A CA1v to LS disconnect cohort 1:
Food intake

B CA1v to LS disconnect cohort 2:
Food intake

C CA1v to LS disconnect cohort 1:
Body weight

D CA1v to LS disconnect cohort 2:
Body weight

E Cre-dependent tdTomato
(Control)

F Cre-dependent tdTomato
(CA1v->LS caspase)

Supplementary Figure 2. Effect of reversible and chronic disconnection of the CA1v to LS

neural pathway on food intake and body weight, and verification of the chronic approach.

There were no effects on food intake (cohort 1: A; cohort 2: B) or body weight (cohort 1: C;

cohort 2: D) of reversible (DREADDs) or chronic (caspase) disconnection of the CA1v to LS

neural pathway compared to controls. Histological verification of the dual viral approach

demonstrates that cre-dependent tdTomato labeling of CA1v neurons induced by LS-origin retro-

cre is robust in a control animal (E), but reduced in an animal injected with cre-dependent

caspase combined with cre-dependent tdTomato due to caspase-induced cell death (F). For

graphs S1A-B (CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8.

For graphs S1C-D (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control

n=10. All values expressed as mean +/- SEM.

Supplemental Figure 3

A

CA1v to mPFC disconnect: Food intake

B

CA1v to mPFC disconnect: Body weight

C

CA1v to LHA disconnect: Food intake

D

CA1v to LHA disconnect: Body weight

Supplementary Figure 3. Effect of contralesional disconnection methods on food intake and

body weight. There were no effects on food intake (A) or body weight (B) of contralesional

CA1v to mPFC compared to controls. CA1v to mPFC disconnect n=12, control n=12. There

were no effects on food intake (C) or body weight (D) of contralesional CA1v to LHA compared

to controls. CA1v to LHA disconnect n=12, control n=11. All values expressed as mean +/-

SEM.

Supplementary Table 1

794   **Supplementary Table 1.** Summary table of the forebrain projection targets of LS neurons that

795   receive input from CA1v (derived the same raw data as the brain flatmap summary diagram in

796   Fig. 5F).

REFERENCES

1. R. G. M. Morris, U. Frey, "Hippocampal synaptic plasticity: Role in spatial learning or the automatic recording of attended experience?" in Burgess, Neil (Ed); Jeffery, Kathryn J (Ed); et al (1999) The hippocampal and parietal foundations of spatial cognition. (Oxford University Press, U Edinburgh, Ctr for Neuroscience, Edinburgh, Scotland London, 1999), pp. 220-246.
2. D. C. Rowland, Y. Roudi, M. B. Moser, E. I. Moser, Ten Years of Grid Cells. *Annu Rev Neurosci* **39**, 19-40 (2016).
3. R. Morris, Developments of a water-maze procedure for studying spatial learning in the rat. *Journal of neuroscience methods* **11**, 47-60 (1984).
4. C. A. Barnes, Memory deficits associated with senescence: a neurophysiological and behavioral study in the rat. *J Comp Physiol Psychol* **93**, 74-104 (1979).
5. J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map: preliminary evidence from unit activity in the freely-moving rat. *Brain research* (1971).
6. T. Hafting, M. Fyhn, S. Molden, M. B. Moser, E. I. Moser, Microstructure of a spatial map in the entorhinal cortex. *Nature* **436**, 801-806 (2005).
7. M. S. Fanselow, H.-W. Dong, Are the dorsal and ventral hippocampus functionally distinct structures? *Neuron* **65**, 7-19 (2010).
8. A. T. Keinath *et al.*, Precise spatial coding is preserved along the longitudinal hippocampal axis. *Hippocampus* **24**, 1533-1548 (2014).
9. K. B. Kjelstrup *et al.*, Finite scale of spatial representation in the hippocampus. *Science* **321**, 140-143 (2008).
10. L. de Hoz, J. Knox, R. G. Morris, Longitudinal axis of the hippocampus: both septal and temporal poles of the hippocampus support water maze spatial learning depending on the training protocol. *Hippocampus* **13**, 587-603 (2003).
11. J. Ferbinteanu, C. Ray, R. J. McDonald, Both dorsal and ventral hippocampus contribute to spatial learning in Long-Evans rats. *Neurosci Lett* **345**, 131-135 (2003).
12. M. B. Moser, E. I. Moser, Functional differentiation in the hippocampus. *Hippocampus* **8**, 608-619 (1998).
13. S. E. Kanoski, H. J. Grill, Hippocampus Contributions to Food Intake Control: Mnemonic, Neuroanatomical, and Endocrine Mechanisms. *Biol Psychiatry* **81**, 748-756 (2017).
14. P. G. Henke, Hippocampal pathway to the amygdala and stress ulcer development. *Brain Res Bull* **25**, 691-695 (1990).
15. K. G. Kjelstrup *et al.*, Reduced fear expression after lesions of the ventral hippocampus. *Proceedings of the National Academy of Sciences* **99**, 10825-10830 (2002).
16. S. Gaskin, A. Gamliel, M. Tardif, E. Cole, D. G. Mumby, Incidental (unreinforced) and reinforced spatial learning in rats with ventral and dorsal lesions of the hippocampus. *Behav Brain Res* **202**, 64-70 (2009).
17. M. S. Bienkowski *et al.*, Integration of gene expression and brain-wide connectivity reveals the multiscale organization of mouse hippocampal networks. *Nat Neurosci* **21**, 1628-1643 (2018).

18. R. Hannapel *et al.*, Postmeal Optogenetic Inhibition of Dorsal or Ventral Hippocampal
Pyramidal Neurons Increases Future Intake. *eNeuro* **6** (2019).
19. R. C. Hannapel, Y. H. Henderson, R. Nalloor, A. Vazdarjanova, M. B. Parent, Ventral
hippocampal neurons inhibit postprandial energy intake. *Hippocampus* **27**, 274-284
(2017).
20. B. K. Mani *et al.*, Neuroanatomical characterization of a growth hormone secretagogue
receptor-green fluorescent protein reporter mouse. *J Comp Neurol* **522**, 3644-3666
(2014).
21. J. M. Zigman, J. E. Jones, C. E. Lee, C. B. Saper, J. K. Elmquist, Expression of ghrelin
receptor mRNA in the rat and the mouse brain. *J Comp Neurol* **494**, 528-548 (2006).
22. I. Merchenthaler, M. Lane, P. Shughrue, Distribution of pre-pro-glucagon and glucagon-
like peptide-1 receptor messenger RNAs in the rat central nervous system. *J Comp
Neurol* **403**, 261-280 (1999).
23. S. E. Kanoski, S. M. Fortin, K. M. Ricks, H. J. Grill, Ghrelin signaling in the ventral
hippocampus stimulates learned and motivational aspects of feeding via PI3K-Akt
signaling. *Biol Psychiatry* **73**, 915-923 (2013).
24. S. E. Kanoski *et al.*, Hippocampal leptin signaling reduces food intake and modulates
food-related memory processing. *Neuropsychopharmacology* **36**, 1859-1870 (2011).
25. T. M. Hsu *et al.*, A hippocampus to prefrontal cortex neural pathway inhibits food
motivation through glucagon-like peptide-1 signaling. *Mol Psychiatry* **23**, 1555-1565
(2018).
26. T. M. Hsu, J. D. Hahn, V. R. Konanur, A. Lam, S. E. Kanoski, Hippocampal GLP-1 receptors
influence food intake, meal size, and effort-based responding for food through volume
transmission. *Neuropsychopharmacology* **40**, 327-337 (2015).
27. A. N. Suarez, C. M. Liu, A. M. Cortella, E. E. Noble, S. E. Kanoski, Ghrelin and Orexin
Interact to Increase Meal Size Through a Descending Hippocampus to Hindbrain
Signaling Pathway. *Biol Psychiatry* **87**, 1001-1011 (2020).
28. G. D. Petrovich, N. S. Canteras, L. W. Swanson, Combinatorial amygdalar inputs to
hippocampal domains and hypothalamic behavior systems. *Brain research reviews* **38**,
247-289 (2001).
29. C. De La Rosa-Prieto *et al.*, Subicular and CA1 hippocampal projections to the accessory
olfactory bulb. *Hippocampus* **19**, 124-129 (2009).
30. T. M. Hsu *et al.*, Hippocampus ghrelin receptor signaling promotes socially-mediated
learned food preference. *Neuropharmacology* **131**, 487-496 (2018).
31. L. W. Swanson, W. M. Cowan, An autoradiographic study of the organization of the
efferent connections of the hippocampal formation in the rat. *J Comp Neurol* **172**, 49-84
(1977).
32. P. Sweeney, Y. Yang, An excitatory ventral hippocampus to lateral septum circuit that
suppresses feeding. *Nat Commun* **6**, 10188 (2015).
33. A. Arszovszki, Z. Borhegyi, T. Klausberger, Three axonal projection routes of individual
pyramidal cells in the ventral CA1 hippocampus. *Front Neuroanat* **8**, 53 (2014).
34. P. Y. Risold, L. W. Swanson, Structural evidence for functional domains in the rat
hippocampus. *Science* **272**, 1484-1486 (1996).

35. R. Garcia, R. M. Vouimba, R. Jaffard, Spatial discrimination learning induces LTP-like
changes in the lateral septum of mice. *Neuroreport: An International Journal for the*
Rapid Communication of Research in Neuroscience **5**, 329-332 (1993).
36. R. Jaffard, R. M. Vouimba, A. Marighetto, R. Garcia, Long-term potentiation and long-
term depression in the lateral septum in spatial working and reference memory. *J*
Physiol Paris **90**, 339-341 (1996).
37. A. Carballo-Marquez, A. Vale-Martinez, G. Guillazo-Blanch, M. Marti-Nicolovius,
Muscarinic receptor blockade in ventral hippocampus and prelimbic cortex impairs
memory for socially transmitted food preference. *Hippocampus* **19**, 446-455 (2009).
38. R. A. Countryman, N. L. Kaban, P. J. Colombo, Hippocampal c-fos is necessary for long-
term memory of a socially transmitted food preference. *Neurobiol Learn Mem* **84**, 175-
183 (2005).
39. G. M. Parfitt *et al.*, Bidirectional Control of Anxiety-Related Behaviors in Mice: Role of
Inputs Arising from the Ventral Hippocampus to the Lateral Septum and Medial
Prefrontal Cortex. *Neuropsychopharmacology* **42**, 1715-1728 (2017).
40. N. L. Trent, J. L. Menard, The ventral hippocampus and the lateral septum work in
tandem to regulate rats' open-arm exploration in the elevated plus-maze. *Physiology &*
behavior **101**, 141-152 (2010).
41. J. C. Jimenez *et al.*, Anxiety Cells in a Hippocampal-Hypothalamic Circuit. *Neuron* **97**,
670-683.e676 (2018).
42. T. M. Hsu *et al.*, Hippocampus ghrelin signaling mediates appetite through lateral
hypothalamic orexin pathways. *Elife* **4** (2015).
43. M. S. Fanselow, H. W. Dong, Are the dorsal and ventral hippocampus functionally
distinct structures? *Neuron* **65**, 7-19 (2010).
44. M. Contreras, T. Pelc, M. Llofriu, A. Weitzenfeld, J. M. Fellous, The ventral hippocampus
is involved in multi-goal obstacle-rich spatial navigation. *Hippocampus* **28**, 853-866
(2018).
45. S. L. T. Lee, D. Lew, V. Wickenheisser, E. J. Markus, Interdependence between dorsal and
ventral hippocampus during spatial navigation. *Brain Behav* **9**, e01410 (2019).
46. D. P. Holschneider *et al.*, Cerebral perfusion mapping during retrieval of spatial memory
in rats. *Behavioural brain research* **375**, 112116 (2019).
47. P. Sweeney, Y. Yang, Neural Circuit Mechanisms Underlying Emotional Regulation of
Homeostatic Feeding. *Trends in endocrinology and metabolism: TEM* **28**, 437-448
(2017).
48. D. S. Olton, The radial arm maze as a tool in behavioral pharmacology. *Physiology &*
behavior **40**, 793-797 (1987).
49. H. H. Pothuizen, W. N. Zhang, A. L. Jongen-Rêlo, J. Feldon, B. K. Yee, Dissociation of
function between the dorsal and the ventral hippocampus in spatial learning abilities of
the rat: a within-subject, within-task comparison of reference and working spatial
memory. *The European journal of neuroscience* **19**, 705-712 (2004).
50. L. E. Jarrard, L. P. Luu, T. L. Davidson, A study of hippocampal structure-function
relations along the septo-temporal axis. *Hippocampus* **22**, 680-692 (2012).

51.   P. Sweeney, Y. Yang, An Inhibitory Septum to Lateral Hypothalamus Circuit That
927   Suppresses Feeding. *The Journal of neuroscience : the official journal of the Society for*
928   *Neuroscience* **36**, 11185-11195 (2016).
929   52.   D. Olivo, M. Caba, F. Gonzalez-Lima, J. F. Rodríguez-Landa, A. A. Corona-Morales,
930   Metabolic activation of amygdala, lateral septum and accumbens circuits during food
931   anticipatory behavior. *Behav Brain Res* **316**, 261-270 (2017).
932   53.   L. W. Swanson, Brain maps 4.0-Structure of the rat brain: An open access atlas with
933   global nervous system nomenclature ontology and flatmaps. *J Comp Neurol* **526**, 935-
934   943 (2018).
935   54.   R. C. Ritter, P. G. Slusser, S. Stone, Glucoreceptors controlling feeding and blood glucose:
936   location in the hindbrain. *Science* **213**, 451-452 (1981).
937   55.   E. E. Noble *et al.*, Hypothalamus-hippocampus circuitry regulates impulsivity via
938   melanin-concentrating hormone. *Nat Commun* **10**, 4923 (2019).
939   56.   B. Zingg *et al.*, AAV-Mediated Anterograde Transsynaptic Tagging: Mapping
940   Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron* **93**, 33-
941   47 (2017).
942   57.   A. N. Suarez *et al.*, Gut vagal sensory signaling regulates hippocampus function through
943   multi-order pathways. *Nat Commun* **9**, 2181 (2018).
944   58.   J. D. Hahn, Swanson, L. W,. Bowman, I., Foster, N. N., Zingg, B., Bienkowski, M. S.,
945   Hintiryan, H., & Dong, H.-W. , An open access mouse brain flatmap and upgraded rat
946   and human brain flatmaps based on current reference atlases. *J. Comp. Neurol.* (2020).
947   59.   R. A. Countryman, P. E. Gold, Rapid forgetting of social transmission of food preferences
948   in aged rats: relationship to hippocampal CREB activation. *Learn Mem* **14**, 350-358
949   (2007).
950   60.   B. G. Galef, Jr., W. Y. Lee, E. E. Whiskin, Lack of interference in long-term memory for
951   socially learned food preferences in rats (Rattus norvegicus). *J Comp Psychol* **119**, 131-
952   135 (2005).
953   61.   B. G. Galef, Jr., E. E. Whiskin, Socially transmitted food preferences can be used to study
954   long-term memory in rats. *Learn Behav* **31**, 160-164 (2003).
955   62.   J. D. Hahn, O. Sporns, A. G. Watts, L. W. Swanson, Macroscale intrinsic network
956   architecture of the hypothalamus. *Proc Natl Acad Sci U S A* **116**, 8018-8027 (2019).
957