Hippocampus-lateral septum circuitry mediates foraging-related spatial memory in rats

Elizabeth A. Davis1*, Clarissa M. Liu1,2*, Isabella H. Gianatiempo1, Andrea N. Suarez1, Alyssa M. Cortella1, Joel D. Hahn3, and Scott E. Kanoski1,2

*These two authors contributed equally
1Human and Evolutionary Biology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089; 2Neuroscience Graduate Program, University of Southern California, 3641Watt Way, Los Angeles, CA, USA. 90089; 3Neurobiology Section, Department of Biological Sciences, Dornsife College of Letters, Arts and Sciences, University of Southern California, 3616 Trousdale Pkwy, Los Angeles, CA, USA, 90089

Corresponding author:
Scott E. Kanoski, PhD
University of Southern California
3616 Trousdale Parkway, AHF-252
Los Angeles, CA 90089-0372
Phone: 213-821-5762
Email: kanoski@usc.edu

Classification: Biological Sciences, Neuroscience

Keywords: feeding, cognition, place cells, motivation, reward, foraging, lateral hypothalamic area, prefrontal cortex

ABSTRACT

Remembering the location of a food or water source is essential for survival. Here we identify a hippocampal-septal neural circuit that is necessary for spatial memory based specifically on appetitive reinforcement. Both reversible and chronic disconnection of ventral hippocampus CA1 subregion (CA1v) projections to the lateral-septum (LS) using pathway-specific dual viral approaches impaired spatial memory retention for the location of either food or water reinforcement. However, disconnection of this pathway did not affect performance in an aversive escape-motivated spatial memory task that used the same apparatus and visuospatial cues. The selectivity of this pathway in mediating foraging-related spatial memory was further supported by results showing that CA1v-LS disconnection did not affect anxiety-like behavior, locomotor activity, or social and olfactory-based appetitive learning. To examine whether CA1v-LS mediation of foraging-related spatial memory involves collateral projections of CA1v neurons, we utilized virus-based neural pathway tracing analyses to identify the medial prefrontal cortex (mPFC) as a collateral target of LS-projecting CA1v neurons. However, functional disconnection of CA1v and mPFC did not affect appetitive spatial memory, thus further supporting the selectivity of CA1v-LS signaling for this behavior. The nucleus accumbens, lateral hypothalamic area, and other brain regions associated with food motivation and reward were identified as second-order targets of CA1v-LS signaling using a multisynaptic anterograde tracing approach. Collective results reveal that CA1v to LS communication plays a critical role in remembering an environmental location based on appetitive (food or water) but not aversive (escape) reinforcement, thus identifying a novel neural pathway regulating foraging-related memory processes.

INTRODUCTION

A survival advantage common to early humans and lower-order mammals is to accurately
remember the location of a food or water source in the environment, and to then efficiently
navigate back to a shelter or other safe location. The neurobiological substrates that regulate
visuospatial navigation are therefore critical to effectively implement food-directed or water-
directed foraging behavior. In rodents a number of maze tasks have been developed for studying
spatial memory and navigation, including the widely popular Morris water maze that requires
animals to swim and navigate to a submerged escape platform using distal viusospatial cues as a
guide (1). The conceptually similar Barnes maze task requires animals to learn and recall a
location for escaping mildly aversive stimuli (bright lights, loud noises) (2). These and other
tasks have been instrumental in identifying brain structures within the telencephalon that are
essential for visuospatial mapping and egocentric orientation in the environment, including the
hippocampus and the medial entorhinal cortex, respectively (3, 4). However, despite that reliably
locating food and water sources in the environment is a key selection pressure driving evolution
of visuospatial navigation, the overwhelming majority of rodent model research on the substrates
of spatial learning and memory have utilized procedures such as the Morris water maze and the
Barnes maze that each involve escaping aversive reinforcement (5, 6). Furthermore, while single
unit recordings have been used to identify specific populations of neurons that subserve distinct
navigational functions (e.g., hippocampal "place cells", medial entorhinal "grid cells") (7, 8), the
bulk of this research has recorded neural activity under neutral conditions void of either
appetitive or aversive reinforcement. Very little research has been dedicated to identify brain
regions and neural circuits that may specifically promote spatial memory based on the location of
appetitive reinforcers (e.g., food, water), as well as the extent to which the nature of the
reinforcement is a critical factor in deciphering the brain's control of visuospatial navigation.
The majority of rodent model research investigating brain regions that mediate
visuospatial navigational memory has focused on the anterior and "dorsal" subregion of the
hippocampus (septal pole; HPCd). However, the posterior and "ventral" hippocampus subregion
(temporal pole; HPCv), while classically associated with stress- and affect-associated memory
processes (9), also plays a role in visuospatial learning and memory (10-12). For example, under

some testing conditions selective HPCv lesions impair spatial memory performance in the Morris

water maze (12, 13). Moreover, place cells that are responsive to changes in the visuospatial

environment are present within both the HPCd and HPCv pyramidal layers, with a linear

increase in the scale of representation from the dorsal to the ventral pole (11). Despite a common

role for the HPCd and HPCv in mediating spatial memory, there is also evidence for a functional

distinction between the subregions (9, 14, 15). For instance, lesions of the HPCv but not the

HPCd alter stress responses (16) and anxiety-like behavior (17), whereas HPCd but not HPCv

lesions impair spatial memory in an incidental (nonreinforced) procedure (18). These two HPC

subregions also have distinct afferent and efferent neural connections. This disparate

neuroanatomical connectivity supports a framework for a functional diversity in which the HPCd

preferentially processes cortical-derived sensory information and the HPCv preferentially

processes metabolic and limbic-derived affective information (9, 15). The functional distinction

between these two subregions is further supported by the generation of the hippocampus gene

expression atlas, which provides a comprehensive integration of gene expression and

connectivity across the HPC axis (19). Given that both the HPCv and the HPCd participate in

spatial memory but have distinct neuroanatomical connectivity and contributions when it comes

to regulating other cognitive and mnemonic processes, it is feasible that the HPCv and HPCd

support different forms of spatial memory depending on the type of reinforcement and/or the

context associated with the behavior.

Recent findings identify the HPCv as a critical brain substrate in regulating feeding

behavior and food-directed memory processes. Reversible inactivation of HPCv neurons after a

meal increases the size of and reduces the latency to consume a subsequent meal (20, 21). In

addition, receptors for feeding-related hormones are more abundantly expressed in the HPCv

compared to the HPCd (e.g., ghrelin receptors (22, 23), glucagon-like peptide-1 [GLP-1]

receptors (24)), and these HPCv endocrine and neuropeptide receptor systems alter food intake

and feeding-related memory (25). For example, leptin and GLP-1 act in the HPCv (CA1v

subregion) to decrease food intake and food-motivated conditioned behaviors (26-28), whereas

the orexigenic gut-derived hormone ghrelin administered to the HPCv but not HPCd has

opposite effects (25, 29). Olfactory information, which is intimately connected with feeding

behavior, is also preferentially processed within the HPCv compared with the HPCd. The CA1v,

specifically, is bidirectionally connected to brain regions that process olfactory information (9,

30, 31), and CA1v neurons respond more robustly to olfactory contextual cues compared with

CA1d (10). Further, ghrelin signaling in the CA1v improves olfactory- and socially-mediated

memory for food preference (32). Given the HPCv appetite-relevant neuroanatomical

connections, endocrine and neuropeptide receptor expression profiles, and functional evidence

linking this subregion with food-motivated behaviors, we hypothesize that HPCv mediation of

visuospatial memory is preferentially biased to food and water-reinforced foraging behavior.

HPCv pyramidal neurons have extensive projection targets throughout the brain (33), yet

the functional relevance of these pathways is poorly understood. Recently, CA1v projections to

the medial prefrontal cortex (mPFC) and lateral hypothalamic area (LHA) were found to mediate

feeding-related outcomes associated with GLP-1 and ghrelin signaling, respectively (28, 34).

CA1v neurons also robustly target the LS (35, 36), and a HPCv dentate gyrus-CA3v to lateral

septum (LS) pathway was identified that suppresses feeding (37). However, the relevance of this

pathway to feeding-related learning and memory is unknown. It is feasible that CA1v to LS

projections participate in regulating spatial memory for food location, a notion supported by

findings showing that neuroplastic changes occur in the LS after learning a spatial memory task

(38, 39). Thus, in addition to exploring the role of the HPCv in memory for the spatial location

of food or water, the present study also investigated the specific role of the CA1v to LS pathway

in foraging-relevant memory processes.

To systematically investigate the role of the HPCv and CA1v to LS signaling in

visuospatial learning and memory for food and water reinforcement, we developed two novel

appetitive reinforcement-based spatial foraging behavioral tasks that allows for direct

comparison with an aversive reinforcement-based task of similar difficulty that uses the same

apparatus and spatial cues. Performance in these tasks was assessed following pathway-specific

dual viral-based reversible (chemogenetic inhibition) or chronic (targeted apoptosis)

disconnection of the CA1v to LS pathway. To further expand neural network knowledge on

CA1v to LS signaling, we used conditional viral-based neuroanatomical tracing strategies to

identify both first-order collateral and second-order targets of LS-projecting CA1v neurons.

Collective results from the present study identify novel neural circuitry of specific relevance to

foraging behavior.

RESULTS

The ventral hippocampus (HPCv) is required for remembering the spatial location of food

To examine the importance of the HPCv in memory for the spatial location of food, animals received bilateral *N*-Methyl-D-aspartate excitotoxic lesions of the HPCv (HPCv lesion n=12) or bilateral sham injections (control n=12) (histological analyses for the neuron-specific NeuN antigen in Fig. 1A) and were tested in a novel appetitive visuospatial memory task developed in our lab. Results revealed no significant differences in errors (incorrect hole investigations) (Fig. 1E) or latency to locate the food source hole (Fig. 1F) during training. However, memory probe results show that animals with HPCv lesions decreased the ratio of correct + adjacent / total holes explored in the first minute compared with controls (Figure 1G; $p < 0.05$, cohen's d=-1.0174). Analysis for the combined 2-minute period did not yield a statistical difference between groups.

Post-surgical analyses of food intake (Supp. Fig. 1A) and body weight (Supp. Fig. 1B) found no group differences in these measures over time. These collective results indicate the HPCv is required for memory retention but not learning of the spatial location of food in the environment.

Ventral hippocampus CA1 (CA1v) projections to the lateral septum (LS) mediate appetitive spatial memory for food and water location, but not aversive spatial memory for escape location

A major neuroanatomical target of HPCv CA1 neurons (CA1v) is the LS (36). We investigated the functional relevance of the CA1v to LS signaling in learning and recalling the spatial location of either food or water reinforcement using conditional dual viral approaches to either reversibly (via cre-dependent pathway-specific inhibitory chemogenetics; diagram of approach in Fig. 2A) or permanently (via cre-dependent pathway-specific caspase-induced lesions; diagram of approach in Fig. 2B) disconnect CA1v to LS communication. Results from the appetitive food seeking spatial memory task reveal no significant group differences during training in either errors before locating the correct hole (Fig. 2E) or latency to locate the food source hole (Fig. 2F). However, memory probe results demonstrate that both acute (pathway-specific DREADDs) and chronic (pathway-specific caspase) CA1v to LS disconnection decreased the ratio of correct + adjacent / total holes explored in the entire two minutes of the probe compared with controls

(Fig. 2G; $p < 0.05$, cohen's d=-1.142), with a similar trend in the first minute of the probe

(p=0.059).

Results from the appetitive water seeking spatial memory task in a separate cohort of rats

also reveal no significant group differences during training in either errors before locating the

correct hole (Fig. 2H) or latency to locate the food source hole (Fig. 2I). However, a significant

impairment following either acute or chronic CA1v to LS disconnection of ratio of correct +

adjacent / total holes explored in 2-min memory probe compared with controls (Fig 2J; $p < 0.05$,

cohen's d=-1.18). Analysis for the first minute alone of the probe test did not yield a statistical

difference between groups.

Histological analyses confirmed successful viral transfection of LS-projecting CA1v

neurons with DREADDs (Figs. 2C&D) or caspase (Supp. Figs. 2E&F), and post-surgical

analyses of food intake (Supp. Fig 2A) and body weight (Supp. Fig 2B) found no group

differences in these measures over time (in the absence of CNO treatment). These data

demonstrate that CA1v to LS communication is critical for remembering the spatial location of

food and water, and that these effects are unlikely to be based on differences in energy status or

food motivation.

To evaluate whether the CA1v to LS pathway is specifically involved in appetitive

reinforcement-based spatial memory (food or water) vs. spatial memory in general, in a separate

cohort of animals we tested the effect of reversible and chronic CA1v to LS disconnection in a

spatial learning task based on aversive reinforcement rather than food. Importantly, this task uses

the same apparatus and visuospatial cues as the spatial location food and water memory tasks

described above, but the animals are motivated to locate the tunnel to escape mildly aversive

stimuli (bright lights, loud noise) with no food or water reinforcement in the tunnel (see Fig. 1B).

Training results revealed no significant group differences in errors before correct hole (Fig. 2K)

nor latency to locate the escape hole (Fig. 2L). During the memory probe test, there were no

group differences in the ratio of correct + adjacent / total holes investigated during the first

minute nor the entire two minutes (Fig. 2M) of the memory probe. Histological analyses

confirmed successful viral transfection of LS-projecting CA1v neurons with DREADDs (Figs.

2C&D) or caspase (Supp. Figs. 2E&F), and post-surgical analyses of food intake (Supp. Fig. 2C)

and body weight (Supp. Fig. 2D) found no group differences in these measures over time (in the

absence of CNO treatment). These collective findings suggest that CA1v to LS signaling

specifically mediates spatial memory in a reinforcement-dependent manner, playing a role in

appetitive (food or water reward) but not aversive (escape-based) spatial memory.

Reversible, but not chronic disconnection of CA1v to LS signaling increases chow intake

under free-feeding conditions.

While the results above suggest that the impairments in food or water-based spatial

memory were not based on long-term changes in food intake or body weight, these experiments

were conducted under conditions of either chronic food or water restriction. To examine the role

of CA1v to LS signaling on feeding behavior under free-feeding conditions, we examined meal

pattern feeding analyses in the caspase vs. the control group (chronic CA1v-LS disconnection),

and in the DREADDs group following CNO or vehicle (within-subjects design; reversible

CA1v-LS disconnection). Results averaged over 5d revealed no significant effect of chronic

CA1v to LS disconnection (caspase group) on 24h meal frequency, meal size, meal duration, or

24h food intake in comparison to controls (Supp. Figs. 3A-D). However, acute and reversible

silencing of LS-projecting CA1v neurons (DREADDs group; via lateral ventricular injection of

the DREADD ligand CNO) significantly increased 24h chow intake (Supp. Fig. 3H; $p<0.05$,

cohen's d=0.10) relative to vehicle treatment, accompanied by a trend towards increased meal

frequency (Supp. Fig. 3E; $p=0.11$) and no effect on meal size or meal duration (Supp. Figs.

3F&G). These results are unlikely based on nonselective effects of the CNO, as we've recently

shown that ICV CNO injections do not affect food intake in rats when tested under similar

conditions (40). Collectively these findings show that reversible CA1v to LS disconnection

increases food intake under free-feeding conditions, whereas food intake is not affected by

chronic CA1v to LS ablation.

CA1v to LS disconnection does not affect nonspatial HPC-dependent appetitive memory

To examine the role of CA1v to LS signaling in a nonspatial HPC-dependent food-

reinforced memory task, we tested the effect of reversible and chronic CA1v to LS disconnection

in the social transmission of food preference (STFP) test (diagram in Fig. 3A). The HPCv plays

an important role in STFP (32, 41, 42), which tests socially-mediated food-related memory based

on previous exposure to socially-communicated olfactory cues. Results revealed that there were

no differences in preference ratio between groups (Fig. 3B), with all three groups performing

significantly above chance (ps<0.05). These results suggest that neither acute nor chronic CA1v

to LS disconnection impair food-related memory based on social-based olfactory stimuli.

CA1v to LS disconnection does not affect anxiety-like behavior or levels of locomotor

activity

HPCv to LS circuitry has been shown to play a role in mediating anxiety and stress behavior (43-

45). Given that altered anxiety-like behavior and/or locomotion could produce behavioral

changes that may lead to an apparent spatial memory deficit in any of the spatial learning tasks

used, we examined whether HPCv to LS disconnection yields differences in anxiety-like

behavior (zero maze task; diagram of apparatus in Fig. 3C) and/or locomotion (open field test).

Results showed no significant group differences in time spent in the open zones in the zero maze

test (Fig. 3D) nor in the number of open zone entries (Fig. 3E), suggesting that neither chronic

nor reversible disconnection of the CA1v to LS circuitry influences anxiety-like behavior. In

addition, CA1v to LS disconnection had no effect on general locomotor activity in the open field

test (Fig. 3F). These results support that observed deficits in food-seeking and water-seeking

spatial memory are not secondary to general differences in anxiety-like behavior or locomotor

impairments in the disconnection groups.

The medial prefrontal cortex is a collateral target of LS-projecting CA1v neurons

In addition to the LS, CA1v neurons also robustly project to the medial prefrontal cortex

(mPFC) and the lateral hypothalamic area, two pathways that we have previously shown to be

involved in feeding behavior (27, 29, 34). To examine whether LS-projecting CA1v neurons

have collateral targets in the mPFC, LHA, or in other brain regions, we utilized a conditional

dual viral neural pathway tracing approach that identifies collateral targets of a specific

monosynaptic pathway (CA1v->LS; diagram of approach in Fig. 4A, representative CA1v

injection site in Fig. 4B, representative CA1v injection site in Fig. 4C). Results revealed that LS-

projecting CA1v neurons also project to the mPFC (Fig. 4D), whereas no labeling was observed

in the LHA, and very minimal labeling was observed in other brain regions. Thus, it may be the

case that the impaired spatial memory for food location observed following either reversible or

chronic CA1v to LS disconnection are based, in part, on CA1v to mPFC signaling from the same

CA1v to LS projecting neurons.

Neither CA1v to mPFC nor CA1v to LHA signaling contribute to spatial memory for food location

We next sought to test the functional relevance to food-directed spatial memory of CA1v-originating neural pathways that either do (CA1v to mPFC) or do not (CA1v to LHA) collateralize from CA1 to LS projections. We also investigated the role of a pathway (CA1v to LHA) that is not a target of LS-projecting CA1v neurons, but has previously been associated with appetitive and consummatory aspects of feeding behavior (29, 34). Both the CA1v to mPFC and CA1v to LHA projections are exclusively ipsilateral (27, 34), and thus these pathways allow for a 'contralesional' approach to achieve chronic functional disconnection (diagram of approach in Fig. 4E&4I, modified from (34)). This approach was utilized in favor of the dual viral approaches used for the CA1v-LS disconnection as it does not require the use of viruses and involves fewer surgical injections to achieve a similar goal. We note that this contralesional approach is not a feasible option for the CA1v-LS pathway, as CA1v neurons project to the LS bilaterally (e.g., Fig. 5F). Post-surgical analyses of food intake (Supp. Fig. 3A,C) and body weight (Supp. Fig. 3B,D) found no group differences in these measures over time for either the CA1v-mPFC or the CA1v-LHA disconnection groups. Training results from the appetitive spatial memory task revealed no group differences in errors before locating the correct hole (Fig. 4F, 4J) or latency to locate the food-baited hole (Fig. 4G, 4K) for either the CA1v-mPFC or the CA1v-LHA disconnection groups. Importantly, memory probe results demonstrate that neither CA1v to mPFC nor CA1v to LHA disconnection altered the ratio of correct + adjacent / total holes explored in the first minute or the entire two minutes (Fig. 4H, 4L) compared with controls. These data collectively demonstrate that CA1v to LS mediation of foraging-related memory does not require collateral projections to the mPFC neural pathway. Further, CA1v to LHA neural signaling, which participates in other conditioned aspects of food intake (29, 34), is not required for either learning or remembering of the spatial location of food.

Second-order targets of CA1v to LS neurons include the mPFC, the LHA, and the ACB

We used a dual viral tracing strategy (diagram of approach in Fig. 5A; representative LS injection site in Fig. 5B) to identify downstream targets of LS-projecting CA1v neurons. Results revealed that the mPFC (Fig. 5C), the LHA (Fig. 5D) and the nucleus accumbens (ACB; Fig. 5E)

are among the strongest second-order targets of the CA1v to LS-projecting neurons. Quantitative

analyses using a custom-built data-entry platform (Axiome C, created by JDH) are summarized

graphically for a representative animal on a brain flatmap summary diagram (Fig. 5F) for the

hemisphere ipsilateral (top) and contralateral (bottom) to the injection sites. The data are also

summarized in tabular form in Supplementary Table 1.

DISCUSSION

Memory for the physical location of a food or water source is adaptive for maintaining

adequate energy supply for reproduction and survival. However, the neural circuits mediating

this complex behavior are not well understood, as research on visuospatial memory has

predominantly used tasks with either aversive or neutral/passive reinforcement. Moreover,

whether the neural circuitry mediating spatial memory function is divergent based on the nature

(e.g., appetitive vs. aversive) or modality (e.g., food vs. water) of the reinforcement has not been

systematically investigated. The present study identified a monosynaptic CA1v to LS pathway as

a necessary substrate for food- and water-motivated spatial memory. Moreover, the selectivity of

the CA1v-LS pathway in mediating spatial memory for food and water location is supported by

results showing that disconnection of this pathway did not affect performance in an escape-

motivated spatial memory task of comparable difficulty conducted in the same apparatus,

anxiety-like behavior, locomotor activity, or olfactory and social-based appetitive memory. Viral

pathway tracing identified that LS-projecting CA1v neurons also send collateral projections to

the mPFC, however, functional disconnection of CA1v-mPFC signaling did not impair spatial

memory for food location. Utilization of an anterograde multisynaptic viral tracing approach and

quantitative forebrain-wide axon mapping analyses revealed that the CA1v-LS pathway sends

second-order projections to various feeding and reward-relevant neural substrates such as the

LHA, ACB, and mPFC. Collectively, these data establish monosynaptic CA1v to LS signaling as

essential for spatial memory-based foraging behavior, and we further identify a neural network

of interconnected feeding-relevant substrates that are collateral and/or downstream targets of this

pathway.

Historically, the hippocampus has been divided into the dorsal and ventral subregions

that have both distinct and overlapping functions, anatomical connections, and genetic and

receptor expression profiles (14, 15, 19, 33, 46). The HPCd has been predominantly implicated

in facilitating spatial memory, whereas the HPCv is linked with stress responses, affective

behavior, and energy balance. However, the HPCv also plays a role in spatial memory,

particularly for goal-directed navigation. Consistent with present results, a recent study found

that both the HPCd and HPCv were critical for food reward-directed spatial navigation in an

obstacle-rich complex environment (47). Under certain testing conditions, spatial learning and

memory in an aversive reinforcement-based water maze also requires both subregions of the

HPC (48). Similarly, whole-brain analysis of blood flow in rats during retrieval of spatial

memory in an aversive reinforcement-based Barnes maze task revealed increased activation in

the CA1d and CA1-3v in trained animals compared with controls (49). These results indicate that

spatial processing in the hippocampus is organized longitudinally, and that both HPCd and HPCv

are important for goal-directed spatial navigation. Here we expand this literature by identifying a

specific HPCv-originating pathway (CA1v->LS) that selectively mediates spatial memory for the

location of appetitive (food or water) reinforcement.

While HPCv neurons project widely throughout the neuraxis, projections to the LS are

particularly robust (36). The LS has been implicated in food anticipatory/seeking behavior,

where metabolic activation of the lateral septum peaks immediately before feeding (during food

anticipatory activity) in rabbit pups (50) and gamma-rhythmic input to the LHA from the LS

evokes food approach behavior (51). In addition, the LS can bidirectionally modulate food intake

via a number of feeding-relevant neuropeptides, including glucagon-like peptide-1, ghrelin, and

urocortin (52-54). A recent finding demonstrated that DREADDs-mediated disconnection of a

HPCv to LS pathway (from CA3v and DGv) increases food intake in mice, with

opposite/hyperphagic effects observed following activation of this pathway (37). Similarly, here

we demonstrate that DREADDs-mediated acute silencing of the CA1v to LS pathway increases

food intake in rats. Together these findings support a role for HPCv-LS signaling in regulating

feeding behavior. However, these results are based on analyses conducted in the home cage

where animals have free and unlimited access to food and water. Here we extend these findings

by examining the role of CA1v to LS signaling in foraging-relevant memory processes under

conditions that are more relevant to normal mammalian behavior, where food and water are not

freely available. Not only do our results identify a novel role for CA1v to LS signaling in

processing spatial memory for food and water reward location, but we further show that this

circuit appears to selectively promote spatial memory based on appetitive and not aversive-based

reinforcement.

The use of two different but complementary approaches for neural pathway-specific

disconnection allowed us to test at the levels of learning and memory separately. For example,

the use of a reversible (DREADDs) disconnection procedure allowed us to evaluate the effect of

CA1v to LS disconnection during the memory probe only, with circuit intact during training

(CNO was not administered at any point during training). In contrast, the chronic (caspase)

disconnection procedure was performed before training and therefore could have affected

learning if CA1v to LS pathways were required for the learning of the task. However, our results

showed that CA1v to LS did not impair learning, but rather, only memory retention performance

in the probe test conducted days after the end of training. Thus, we can now confirm based on

two complementary levels of analysis that this pathway is involved in longer-term memory recall

but not learning/acquisition, which is consistent with recent evidence demonstrating that

complete HPCv lesions impair retrieval (probe performance), but not acquisition (learning) in a

Morris water maze (55).

Present results identify the CA1v-LS pathway in coordinating reward-motivated behavior

with learned spatial information. While the use of aversive reinforcement-based spatial memory

tasks are predominant, the radial arm maze has also been used to investigate food-motivated

spatial memory (56). Specifically, the radial arm maze examines different components of spatial

memory relevant to different foraging strategies. In win-shift tasks, animals learn to avoid

previously visited reward arms, which models spontaneous alternation and exploration in

environments with minimal resources. On the other hand, in win-stay tasks, animals learn to

return to previously reinforced arms modeling foraging in environments of plentiful resources.

An advantage of the present food-reinforced Barnes maze approach over the radial arm maze

procedure is that we've developed parallel procedures that use the same apparatus and

visuospatial cues to examine water- and escape-motivated spatial memory, thus offering a

powerful approach to assess the selectivity of the circuit based on multiple reinforcement

modalities (food vs. water vs. escape/aversive). While our results clearly support a role for the

HPCv in food-reinforced spatial memory, a previous study utilizing a four-baited/four-unbaited

version of the eight-arm radial maze task found that NMDA-mediated lesions of the HPCd, but

not HPCv, impair spatial reference and working memory (57). Another study using a similar

procedure reported that complete HPC lesions impaired both spatial and working memory, whereas lesions to either the HPCd, HPCv, or intermediate HPC had minimal effect on memory performance (58). Future studies will needed to address whether discrepancies between the present results and these studies are based on the differences between the behavioral tasks and/or differences in loss of function methodologies. With regards to the latter, we note that our pathway-specific chemogenetic and caspase-lesion approaches offered far greater specificity (LS-projecting CA1v neurons vs. whole HPCv) compared with these studies.

Utilizing dual viral approaches and systematic forebrain-wide quantitative mapping analyses, our findings confirm that the CA1v-LS pathway projects downstream to other neural substrates involved in feeding and motivated behavior, including the LHA, ACB, and mPFC. This experiment was performed to generate novel neural connectivity data for the field and to demonstrate possible downstream 2_{nd}-order targets that may inspire future research on these newly-identified multi-order neural circuits. Given that metabolic activity in the LS and ACB is associated with food anticipatory activity in rabbit pups (50), it is possible that the ACB is a functional downstream target of CA1v-LS signaling for coordinating foraging behavior. It is less likely, however, that the mPFC or the LHA are functional targets of CA1v-LS signaling for appetitive spatial memory control based on the lack of effects on spatial memory for food location when these pathways were disconnected. It is possible, however, that CA1v $\rightarrow$ LS $\rightarrow$ mPFC (or $\rightarrow$ LHA) signaling is relevant to other mnemonic process, such as those involved in drug-motivated or social-based memory tasks.

Collective results identify a CA1v to LS pathway involved in food- and water-motivated spatial memory, but not escape-motivated spatial memory. Furthermore, the selectivity of this pathway to appetitive spatial memory is supported by data showing that neither chronic nor reversible disruption of CA1v to LS signaling influenced various other behavioral outcomes, including anxiety-like behavior, locomotor activity, and nonspatial HPC-dependent appetitive memory. We also systematically characterized collateral and second-order projections of this pathway. Present results demonstrate overlapping pathways coordinated in the higher-order control of energy balance, of which foraging behavior is an essential component, given that foraging requires energy expenditure to locate food for subsequent energy intake. These data shed light on the neural systems underlying complex learned and motivated behaviors that require functional connections between cognitive and feeding-relevant substrates.

METHODS

Animals

Adult male Sprague–Dawley rats (Envigo; 250-275g on arrival) were individually housed in hanging wire cages with *ad libitum* access (except where noted) to water and chow (LabDiet 5001, LabDiet, St. Louis, MO) on a 12h:12h reverse light/dark cycle. All procedures were approved by the University of Southern California Institute of Animal Care and Use Committee.

General intracranial injection procedures

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine 90mg/kg body weight [BW], xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW) followed by a pre-operative, subcutaneous injection of analgesic (ketoprofen, 5mg/kg BW). Post-operative analgesic (subcutaneous injection of ketoprofen, 5mg/kg BW) was administered once per day for 3 days following surgery. The surgical site was shaved and prepped with iodine and ethanol swabs, and animals were placed in a stereotaxic apparatus for stereotaxic injections. NMDA or viruses were delivered using a microinfusion pump (Harvard Apparatus, Cambridge, MA, USA) connected to a 33-gauge microsyringe injector attached to a PE20 catheter and Hamilton syringe. Flow rate was calibrated and set to 83.3nl/sec. Injectors were left in place for 2min post-injection to allow for complete delivery of the infusate. Specific viruses/drugs, coordinates, and injection volumes for procedures are detailed below. Following the completion of all injections, incision sites were closed using either surgical staples, or in the case of subsequent placement of an indwelling cannula, simple interrupted sutures. Upon recovery from anesthesia and return to dorsal recumbency, animals were returned to the home cage. All behavioral procedures occurred 21 days after injections to allow for complete transduction and expression of the viruses, or complete lesioning drugs. General intracranial injection procedures were followed for all injection procedures below.

Chronic lesions of the HPCv

Lesion animals received bilateral excitotoxic HPCv lesions via intracranial injections of N-methyl-d-aspartate (NMDA; 4ug in 200nL; 100uL per hemisphere) at the following

coordinates at three different sites along the rostrocaudal extent of the HPCv (59): [1] -4.8mm

AP, +/-5.0mm ML,   -7.5mm DV, [2] -5.5mm AP, +/-4.5mm ML, -7.0mm DV, and [3] -6.1mm

AP, +/-4.8mm ML, -7.0mm DV with control animals receiving vehicle saline in the same

location. The reference points for AP and ML coordinates were defined at bregma, and the

reference point for the DV coordinate was defined at the skull surface at the target site.

Bilateral HPCv lesion brains were histologically evaluated for the correct placement of

lesions in 1 out of 5 series of brain tissue sections. Neurons were visualized using

immunohistochemistry for the neuron specific antigen NeuN (see Immunohistochemistry). The

extent of NMDA lesions was determined postmortem by immunohistochemical detection of the

neuronal marker NeuN. Rats showing pronounced (~65%) loss of NeuN labeling within the

target region compared with the representative NeuN expression following sham injections in the

control group (average HPCv count of NeuN+ cells in control group) were included in data

analysis (34). Representative images in Fig. 1A.

Acute and chronic disconnection of the CA1v to LS neural pathway

Cre-dependent dual viral strategies were used to generate the following groups: [1] acute

chemogenetic disconnection of the CA1v to LS neural pathway (DREADDs; diagram of

approach in Fig. 2A), [2] chronic disconnection of the CA1v to LS neural pathway (caspase

lesion-induced; diagram of approach in Fig. 2B), and [3] a common/shared control group for

these two disconnection strategies. Regardless of group, all animals received a bilateral AAV

retro-cre injection in the LS (AAV2[retro]-hSYN1-EGFP-2A- iCre-WPRE; 200nL per side) at

the following stereotaxic coordinates (59): +0.84mm AP, +/-0.5mm ML, −4.8mm DV, all

defined at bregma.

According to experimental group, animals received a different virus delivered to the

CA1v subregion of the HPCv. All viruses were administered to the CA1v at the following

stereotaxic coordinates (59): −4.9mm AP defined at bregma, +/-4.8mm ML defined at bregma,

−7.8mm DV defined at skull surface at target site.

[1] DREADDs group for reversible inactivation of CA1v to LS: To allow reversible

chemogenetic inactivation of the CA1v to LS neural pathway, one group of animals received a

bilateral CA1v injection of a cre-dependent virus to drive expression of inhibitory designer

receptors activated by designer drugs (DREADDs), (AAV-Flex-hm4Di-tdTomato). This dual

viral strategy drives expression of inhibitory DREADDs exclusively in LS-projecting CA1v
neurons, which enables acute inactivation of these neurons by injection of the DREADDs ligand,
clozapine-N-oxide (CNO), at the time of behavioral testing.

[2] Caspase group for chronic inactivation of CA1v to LS: To allow chronic
disconnection of the CA1v to LS neural pathway, a second group of animals received a bilateral
CA1v cre-dependent caspase virus (AAV1-Flex-taCasp3-TEVp; 200nL per side) mixed with a
cre-dependent reporter virus (AAV-flex-tdTomato; 200nL per side) for histological verification.
This dual viral strategy drives expression of the apoptosis-mediator molecule caspase exclusively
in LS-projecting CA1v neurons, which induces apoptotic cell death in these neurons while
leaving other CA1v neurons intact.

[3] Common control: A common control group was generated so that the DREADDs and
caspase groups could both be compared to the same control group, instead of two different
control groups for the DREADDs and caspase groups, respectively. This allowed us to reduce
the number of animals needed for the same experimental objective. Thus, all animals received an
indwelling cannula and ICV injections of CNO as described below.

Immediately following viral injections, all animals were surgically implanted with a
unilateral indwelling intracerebroventricular (ICV) cannula (26-gauge, Plastics One, Roanoke,
VA) targeting the lateral ventricle (VL). Cannulae were implanted and affixed to the skull with
jeweler's screws and dental cement at the following stereotaxic coordinates: −0.9mm AP defined
at bregma, +1.8mm ML defined at bregma, −2.6mm DV defined at skull surface at site.
Placement for the VL cannula was verified by elevation of blood glucose resulting from an
injection of 210μg (2μL) of 5-thio-D-glucose (5TG) using an injector that extended 2mm beyond
the end of the guide cannula (60). A post-injection elevation of at least 100% of baseline
glycemia was required for subject inclusion. Animals that did not pass the 5TG test were retested
with an injector that extended 2.5mm beyond the end of the guide cannula and, upon passing the
5TG test, were subsequently injected using a 2.5mm injector instead of a 2mm injector for the
remainder of the study. Prior to behavioral testing where noted, all animals received an ICV
18mmol infusion (2uL total volume) of the DREADDs ligand clozapine N-oxide (CNO) (61),
rendering only DREADDs animals chemogenetically inactivated. CNO injections were hand
delivered using a 33-gauge microsyringe injector attached to a PE20 catheter and Hamilton

syringe through the indwelling guide cannulae. Injectors were left in place for 30sec to allow for

complete delivery of the CNO.

Cre-dependent DREADD expression targeting LS-projecting neurons in CA1v was

evaluated based on localization of the fluorescent reporter tdTomato. Immunohistochemistry for

red fluorescent protein (RFP) was conducted to amplify the tdTomato signal (see

Immunohistochemistry). Only animals with tdTomato expression restricted within CA1v neurons

were included in subsequent behavioral analyses. Representative images in Figs. 2C&D.

The general approach for neuronal apoptosis due to activation of cre-dependent caspase

targeting LS-projecting neurons in CA1v was validated in a separate cohort (n=8) based on

reduction of a cre-dependent fluorescent reporter tdTomato due to neuron cell death compared

with controls. The caspase group brains (which received a retro-cre virus in the LS in

conjunction with an injection of cre-dependent caspase virus mixed with a cre-dependent virus

that drives a tdTomato fluorescent reporter in the CA1v) were compared to control brains (that

received a retro-cre virus in the LS in conjunction with only a cre-dependent that drives a

tdTomato fluorescent reporter in the CA1v, which was equivalently diluted to match the caspase

injections). In both groups, immunohistochemistry for red fluorescent protein (RFP) was

conducted to amplify the tdTomato signal (see Immunohistochemistry). Confirmation of

successful caspase-driven lesions in the CA1v using this approach was evaluated by reduced

tdTomato fluorescence in comparison to control animals, with the expression of CA1v tdTomato

expression in the caspase group less than 15% of that observed in controls, on average

(Representative images in Supplementary Figs 2E&F). While this cohort allowed for successful

validation of the pathway-specific disconnection caspase approach, to improve the viability of

the animals for long-term behavioral analyses, the 3rd AAV with the fluorescent reporter (cre-

dependent tdTomato AAV) was omitted for cohorts undergoing behavioral analyses. For these

groups, histological confirmation for inclusion in subsequent statistical analyses was based on

identification of injections sites confined with the LS and CA1v, as observed using darkfield

microscopy.

Identification of collateral targets of CA1v to LS projecting neurons

Collateral targets of the CA1v to LS neural pathway were identified using a dual viral

approach (diagram of approach in Fig. 4A) where a retrograde vector was injected into the LS

(AAV2retro-hSyn1-eGFP-2A-iCre-WPRE; 200nL per side; coordinates as above), and a Cre-

dependent anterograde vector (AAV1-CAG-Flex-tdTomato-WPRE-bGH; 200nL per side;

coordinates as above) was injected in the CA1v. This latter injection drives tdTomato transgene

expression in CA1v neurons that project to the LS, which allows for brain-wide analyses of

collateral targets. After 3 weeks of incubation time to allow for complete transduction and

expression of the viruses, brains were collected, immunohistochemically processed, and imaged

as described below.

Contralesional disconnection of the CA1v to mPFC neural pathway or the CA1v to LHA

neural pathway

To functionally disconnect the CA1v to mPFC pathway (diagram of approach in Fig. 4E),

or the CA1v to LHA pathway (diagram of approach in Fig. 4J) lesion animals received a

unilateral excitotoxic CA1v lesion via an intracranial injection of NMDA (200nL) at the

following coordinates (36): −4.9mm AP defined at bregma, + or - 4.8mm ML defined at bregma

(left/right counterbalanced to be contralateral to mPFC or LHA lesion within-animal), −7.8mm

DV defined at skull surface at site, with control animals receiving vehicle saline injections in the

same location.

In addition to the CA1v lesion, CA1v to mPFC disconnect animals also received a

unilateral excitotoxic mPFC lesion via two intracranial injections of NMDA (4ug in 200nL;

100nL per injection) at the following coordinates (59): [1] +2.7mm AP, + or - 0.7mm ML

(left/right counterbalanced to be contralateral to CA1v lesion within-animal), −5.3mm DV and

[2] +3.0mm AP, + or - 0.7mm ML (left/right counterbalanced to be contralateral to CA1v lesion

within-animal), −4.7mm DV, with control animals receiving vehicle saline in the same location.

The reference points for all mPFC coordinates were defined at bregma.

In addition to the CA1v lesion, CA1v to LHA disconnect animals also received a

unilateral excitotoxic LHA lesion via an intracranial injection of NMDA (100nL per injection) at

the following coordinates (59): +2.9mm AP, + or – 1.1mm ML (counterbalanced to be

contralateral to CA1v lesion within-animal), −8.9mm DV, with control animals receiving vehicle

saline in the same location. The reference points for all LHA coordinates were defined at

bregma.

Contralesional brains were histologically evaluated for the correct placement of lesions by

visualizing neurons using immunohistochemistry for the neuron specific antigen NeuN (see

Immunohistochemistry). The extent of NMDA lesions was determined postmortem by

immunohistochemical detection of the neuronal marker NeuN. Rats showing pronounced (65%)

loss of NeuN labeling within the target region compared with a sham injection in the

contralateral side were included in data analysis (34).

Identification of second order targets of CA1v to LS projecting neurons

To identify second order targets of CA1v to LS-projecting neurons (diagram of approach

in Fig. 5A), animals received a bilateral injection of a transsynaptic Cre-inducing anterograde

vector into CA1v (AAV1-hSyn-Cre-WPRE-hGH; 200nL per side; coordinates as above) that

drives expression of Cre in both first-order (CA1v injection site) and 2^{nd}-order (but not 3^{rd}-

order) neurons via transsynaptic virion release (62, 63). This was combined with a bilateral

injection of a Cre-dependent anterograde vector in the LS (AAV1-CAG-Flex-tdTomato-WPRE-

bGH; 200nL per side, coordinates as above). This latter injection allows for anterograde tracing

from 1^{st}-order LS targets receiving synaptic input from CA1v. After 3 weeks of incubation time

to allow for complete transduction and expression of the viruses, brains were collected,

immunohistochemically processed, and imaged as described below.

Data were entered using a custom built data-entry platform (Axiome C, created by JDH)

built around Microsoft Excel software and designed to facilitate entry of data points for all gray

matter regions across their atlas levels as described in a rat brain reference atlas: Brain Maps 4.0

(59). The Axiome C approach was used previously to facilitate the analysis of brain gene-

expression data (64). An ordinal scale, ranging from 0 (absent) to 7 (very strong), was used to

record the qualitative weight of anterograde labeling. An average value was then obtained for

each region across its atlas levels for which data were available. These data are summarized

graphically for a representative animal on a brain flatmap summary diagram (adapted from (65)).

Food intake and body weight

For the day prior to surgery (day 0) and for two weeks thereafter, 24h chow intake was

measured daily just prior to dark cycle onset to determine effects of experimental procedures on

food intake. Spillage was accounted for daily by collecting crumbs onto Techboard paper placed

under the cages of each animal. Additionally, animals were weighed daily just prior to dark cycle

onset to determine effects of experimental procedures on body weight.

Meal pattern

Meal size, meal frequency, meal duration, and cumulative 24h food intake were

measured using Biodaq automated food intake monitors (Research Diets, New Brunswick, NJ).

Rats (control n=7, caspase n=8, and DREADDs n=8) were acclimated to the Biodaq on ad

libitum chow for 3 days. Caspase group vs. control group feeding behavior data were collected

over a 5-day period (untreated, between-subjects design). The DREADDs group was tested using

a 2-treatment within-subjects design. Rats received 18mmol infusion of ICV CNO (2uL total

volume) or vehicle (daCSF, 33% dimethyl sulfoxide in artificial cerebrospinal fluid) 1h prior to

lights off and 24h feeding behavior was measured. Meal parameters were set at minimum meal

size=0.2g and maximum intermeal interval=600s.

Appetitive food seeking spatial memory task

To test visuospatial learning and memory for food reinforcement, we developed a novel

spatial food seeking task modified from the traditional Barnes maze procedure (6). Throughout

this paradigm, animals were maintained at 85% free-feeding body weight. The procedure

involves an elevated gray circular platform (diameter: 122cm, height: 140cm) consisting of 18

uniform holes (9.5cm diameter) spaced every twenty degrees around the outside edge. Under one

of the holes is a hidden tunnel (38.73cm L x 11.43cm W x 7.62cm D and a 5.08cm step).

Surrounding the table are distinct spatial cues on the wall (e.g. holiday lights, colorful shapes,

stuffed unicorn) that are readily visible to the animal. In contrast to the traditional Barnes Maze

where an animal uses the spatial cues to escape mildly aversive stimuli in the environment (e.g.

bright light and loud sound), this task utilizes food as motivation, such that each hidden tunnel

contained five 45mg sucrose pellets (Bio-Serv, Flemington, NJ). Additionally, a quiet white

noise (60dB) was used to drown out background noise and floor lamps were used for low-level

ambient lighting. On the first day, each animal underwent a habituation session consisting of

1min inside a transport bin under the table, 2min underneath the start box in the center of the

table, and 3min inside the hidden tunnel with 5 sucrose pellets. During training, each rat was

assigned a specific escape hole according to the position of the spatial cues with the location

counterbalanced across groups. Before each trial, animals were placed in the start box for 30

seconds. Animals were trained over the course of two 5-min trials per day for four days (five

days in contralesional experiments) to learn to use the spatial cues in order to locate the correct

hole with the hidden tunnel with sucrose pellets. After finding the correct hole and entering the

hidden tunnel during each 5min training session, animals were allowed to remain in the hidden

tunnel for 1 minute, and consistently consumed all 5 sucrose pellets at this time. In the event that

a rat did not find the correct hole during the 5min training session, the experimenter gently

guided the rat to the correct hole and were allowed to crawl in and then complete the training in

the same fashion as rats that found the hole on their own. Learning during training was scored

via animal head location tracking by AnyMaze Behavior Tracking Software (Stoelting Co.,

Wood Dale, IL). The incorrect hole investigations prior to finding the correct hole with sucrose

pellets ("errors before correct hole") as well as time to finding the correct hole ("latency to

correct hole") were calculated as an average of the two trials per day, and examined across days

of training.

After the conclusion of training, rats had a two-day break where no procedures occurred,

then were tested in a single 2min memory probe in which the hidden tunnel and sucrose pellets

were been removed. All animals received an ICV 18mmol infusion of CNO (2uL total volume)

1h prior to the memory probe, rendering only DREADDs animals chemogenetically inactivated

by CNO. In the memory probe, the ratio of the correct hole plus adjacent hole investigations over

the total number of hole investigations were calculated via animal head location tracking by

AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). This dependent variable was

selected a priori instead of correct hole only investigations based on recent findings

demonstrating that taste-responsive hippocampal neurons exhibit weaker spatial selectivity in

comparison to non-taste-responsive hippocampal neurons (66). When establishing and refining

the food-reinforced spatial memory procedure in multiple cohorts of rats, our preliminary data

revealed that memory probe effects in normal/control animals are either stronger in the 1_{st}

minute, or in the 1_{st} two minutes combined (but not the 2_{nd} minute alone), with some cohorts

showing larger paradigm effects in minute 1, and others in minutes 1+2. Thus, for all

experiments presented we analyzed both the 1st minute and the 1st two minutes combined based

on an a priori determination from our preliminary studies. A diagram of the apparatus used for

the spatial foraging task is included in Fig. 1B.

During training sucrose pellets were taped underneath all of the holes to preclude

olfactory-based search strategies during training. Importantly, the memory probe test does not

include any sucrose pellets in the apparatus at all, and therefore there is no possibility of sucrose

odor cues to influence memory probe performance.

Appetitive water seeking spatial memory task

We modified our spatial food seeking task to utilize water reward instead of food reward.

Throughout this paradigm, animals were water restricted and received 90-min access to water

daily (given at least 1hr after the termination of behavioral procedures each day). In addition to

the habituation parameters described above, animals in this task were subjected to two 10-min

habituation sessions in the hidden tunnel with a full water dish prior to training, which allowed

them to become accustomed to reliably drinking from the water dish in the hidden tunnel.

Training procedures were the same as for the food-based task above, except instead of sucrose

pellets in the hidden tunnel, a water dish containing ~100 mL of water was placed in the back of

the hidden tunnel during training, and animals were allowed to remain in the hidden tunnel for

2min (instead of 1min, as above). Each animal consumed a minimum of 2mL of water during

this time for the training sessions. This newly-developed procedure allowed us to test spatial

learning and memory motivated by water reward using similar apparatus and stimulus conditions

to the food-reinforced task.

Aversive spatial memory escape task (Barnes maze)

To test visuospatial learning and memory for escape reinforcement, we used a modified

traditional Barnes maze procedure, which is a visuospatial-based escape task (6). Procedures

were exactly the same as above (appetitive spatial memory food seeking and water seeking tasks,

using the same apparatus, in the same room, and with the same visuospatial cues) aside from the

omission of the sucrose pellets or water dish in the hidden tunnel, the presence of mildly aversive

bright (120W) overhead lighting instead of dim ambient lighting, and a mildly aversive loud

white noise (75dB) instead of a quiet white noise (60dB). This allowed us to test spatial learning

and memory motivated by escape from aversive stimuli in a nearly-identical procedure to our

spatial foraging test for learning and memory motivated by palatable food or water consumption.

A diagram of the apparatus used for the spatial escape task is included in Fig. 1B.

Social transmission of food preference (STFP)

To examine food-related memory based on social- and olfactory-based cues, we utilized

the social transmission of food preference (STFP) task and adapted protocols from (32, 42, 67,

68). Briefly, untreated normal adult rats are designated as 'Demonstrators', while experimental

groups are designated as 'Observers'. Demonstrators and Observers are habituated to a powdered

rodent chow [LabDiet 5001 (ground pellets), LabDiet, St. Louis, MO] overnight. 24h later,

Observers are individually paired with demonstrators and habituated to social interaction, where

rat pairs are placed in a social interaction arena (23.5cm W × 44.45cm L × 27cm H clear plastic

bin with Sani-chip bedding) and allowed to interact for 30min. Both Observers and

Demonstrators are returned to their home cages and food is withheld for 23hr prior to the social

interaction. For the social interaction, Demonstrators are given the opportunity to consume one

of two flavors of powdered chow (flavored with 2% marjoram or 0.5% thyme; counterbalanced

according to group assignments) for 30min in a room separate from Observers. Our pilot studies

and previous published work (68, 69) show that rats equally prefer these flavors of chow. The

Demonstrator rat is then placed in the social interaction arena with the Observer rat, and the pairs

are allowed to socially interact for 30min. Observers are then returned to their home cage and

allowed to eat *ad libitum* for 1h and then food is removed. The following day, the 23h food-

deprived Observer animals are given a home cage food preference test for either the flavor of

chow paired with the Demonstrator animal, or a novel, unpaired flavor of chow that is a flavor

that was not given to the Demonstrator animal (2% marjoram vs. 0.5% thyme; counterbalanced

according to group assignments). All animals received an ICV 18mmol infusion of CNO (2uL

total volume) 1h prior to the social interaction session, rendering only DREADDs animals

chemogenetically inactivated by CNO. Food intake (1h) was recorded with spillage accounted

for by collecting crumbs onto Techboard paper that is placed under the cages of each animal.

The % preference for the paired flavor is calculated as: 100*Demonstrator-paired flavored chow

intake/Demonstrator + Novel flavored chow intake. In this procedure, normal untreated animals

learn to prefer the Demonstrator paired flavor based on social interaction and smelling the breath

of the Demonstrator rat (32, 42, 67-69). A diagram of the STFP procedure is included in Fig. 3E.

Zero maze

The zero maze behavioral paradigm was used to evaluate anxiety-like behavior. The zero

maze apparatus used was an elevated circular track, divided into four equal length sections. Two

zones were open with 3 cm high curbs ('open zones'), whereas the two other zones were closed

with 17.5 cm high walls ('closed zones'). All animals received an ICV 18mmol infusion of CNO

(2uL total volume) 1h prior to testing, rendering only DREADDs animals chemogenetically

inactivated by CNO. Behavioral testing was performed during the light cycle. Animals were

placed in the maze for a single, 5min trial in which the location of the center of the animal's

body was measured by AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). The

apparatus was cleaned with 10% ethanol in between animals. During the trial, the number of

open zone entries and total time spent in open sections (defined as body center in open sections)

were measured, which are each indicators of anxiety-like behavior in this procedure. A diagram

of the zero maze apparatus is included in Fig. 3G.

Open field

An open field test was used to evaluate general levels of locomotor activity. The

apparatus used for the open field test was an opaque gray plastic bin (60cm × 56cm), which was

positioned on a flat table in an isolated room with a camera directly above the center of the

apparatus. Desk lamps were positioned to provide indirect lighting to all corners of the maze

such that the lighting in the box uniformly measured 30 lux throughout. All animals received an

ICV 18mmol infusion of CNO (2uL total volume) 1h prior to testing rendering only DREADDs

animals chemogenetically inactivated by CNO. Behavioral testing began at dark onset. At the

start of the 10min test, each animal was placed in the open field apparatus in the same corner

facing the center of the maze. The location of the center of the animal's body was measured with

the AnyMaze Behavior Tracking Software (Stoelting, Wood Dale, IL). Total distance traveled

was measured by tracking movement from the center of the animal's body throughout the test.

Immunohistochemistry

Rats were anesthetized via an intramuscular injection of an anesthesia cocktail (ketamine

90mg/kg BW xylazine, 2.8mg/kg BW and acepromazine and 0.72mg/kg BW) then transcardially

perfused with 0.9% sterile saline (pH 7.4) followed by 4% paraformaldehyde (PFA) in 0.1M

borate buffer (pH 9.5; PFA). Brains were dissected from the skull and post-fixed in PFA with

15% sucrose for 24h, then flash frozen in isopentane cooled in dry ice. Brains were sectioned to

30μm thickness on a freezing microtome. Sections were collected in 5-series and stored in

antifreeze solution at −20°C until further processing. General fluorescence IHC labeling

procedures were performed as follows. The antibodies and dilutions that were used are as

follows: [1] For lesion histology using the neuron-specific protein NeuN, the rabbit anti-NeuN

primary antibody (1:1000, Abcam) was used followed by a donkey anti-rabbit conjugated to

AF488 (1:500, Jackson Immunoresearch). [2] To amplify the native tdTomato signal for

neuroanatomical tracing or DREADDs histology, the rabbit anti-RFP primary antibody (1:2000,

Rockland) was used followed by a donkey anti-rabbit conjugated to Cy3 (1:500, Jackson

Immunoresearch). [3] To amplify the native GFP signal for LS injection site histology, the

chicken anti-GFP primary antibody (1:500, Abcam) was used followed by a donkey anti-chicken

secondary antibody conjugated to AF488 (1:500, Jackson Immunoresearch). Antibodies were

prepared in 0.02M potassium phosphate buffered saline (KPBS) solution containing 0.2% bovine

serum albumin and 0.3% Triton X-100 at 4°C overnight. After thorough washing with 0.02M

KPBS, sections were incubated in secondary antibody solution. All secondary antibodies were

obtained from Jackson Immunoresearch and used at 1:500 dilution at 4°C, with overnight

incubations (Jackson Immunoresearch; West Grove, PA, USA). Sections were mounted and

coverslipped using 50% glycerol in 0.02 M KPBS and the edges were sealed with clear nail

polish. Photomicrographs were acquired using a Nikon 80i (Nikon DSQI1,1280X1024

resolution, 1.45 megapixel) under epifluorescence or darkfield illumination.

Lesions and virus expression were quantified in one out of five series of brain tissue sections and

analyses were performed in sections from Swanson Brain Atlas level 34-36 (59). The inclusion /

exclusion criteria are described above in the methods of each of these experiments.

Statistics

Data are expressed as mean +/- SEM. Differences were considered to be statistically

significant at $p<0.05$. All variables were analyzed using the advanced analytics software package

Statistica (StatSoft, Tulsa, OK, USA). For all measures of food intake, body weight, and

errors/latency during spatial foraging and spatial escape task training, differences between

groups were evaluated using two-way repeated measures ANOVAs (treatment x time). For

lesion experiments, measures during spatial foraging and spatial escape task probes, differences

between groups were evaluated using two-tailed independent two-sample Student's t-tests. For

CA1v to LS disconnection experiments, measures during spatial foraging and spatial escape task

probes, the STFP paradigm, the zero maze paradigm, and the open field test, differences between

groups were evaluated using one-way ANOVAs. Significant ANOVAs were analyzed with a

Fisher's LSD posthoc test where appropriate. Outliers were identified as being more extreme

than the median +/- 1.5 * interquartile range. For all experiments, assumptions of normality,

homogeneity of variance (HOV), and independence were met where required.

CONFLICT OF INTEREST

No conflicts of interest.

ACKNOWLEDGEMENTS

This work was supported by National Institute of Diabetes and Digestive and Kidney Diseases

grants: DK104897 to SEK, DK118944 to CML, and DK116558 to ANS. Clozapine-N-Oxide

was kindly provided by the National Institute of Mental Health. The authors are grateful to the

Kanoski lab undergraduates for their assistance in behavioral experiments and histology.

Figure 1

Figure 1. Bilateral HPCv lesions impair spatial memory for food location. Representative HPCv lesion histology with NeuN immunohistochemistry (A; scale bars 500μm). Spatial foraging task apparatus (B). Representative navigation paths of a control animal preferentially investigating correct (filled red) and adjacent holes (outlined orange) during spatial foraging memory probe (C). Representative navigation path of HPCv lesioned animal during spatial foraging memory probe (D). Bilateral HPCv lesions did not impair learning of the spatial foraging task compared to controls, as measured by errors before locating correct hole during task training (E) and latency to locate correct hole during task training (F). Bilateral HPCv lesions impaired retention of the spatial foraging task, as measured by the ratio of investigation of correct plus adjacent holes over total investigated during the first minute of the task ($p < 0.05$; G). There were no group differences observed when evaluated over the entire two minutes of the task (H). Dotted line indicates chance performance level (0.167). For graphs 1D-G, lesion n = 11, control n = 18. All values expressed as mean +/- SEM.

Figure 2

Figure 2. Reversible and chronic CA1v to LS neural disconnection impairs spatial memory

for food and water location but not for escape location. Diagram of dual viral approach using

a cre-dependent inhibitory DREADDs approach to reversibly disconnect CA1v to LS neural

pathway (A). Diagram of dual viral approach using a cre-dependent caspase approach to

chronically disconnect CA1v to LS neural pathway (B). Representative injection site in CA1v

demonstrating LS-projecting neurons infected with inhibitory DREADDs, which simultaneously

drives expression of a fluorescent tdTomato transgene (C, D; scale bars 500μm and 100μm,

respectively). Neither reversible (DREADDs) nor chronic (caspase) disconnection of the CA1v

to LS pathway impaired learning of the spatial food seeking task compared to controls, as

measured by errors before correct hole during task training (E) and latency to correct hole during

task training (F). Both reversible and chronic disconnection of the CA1v to LS pathway impaired

retention of the food location as measured by the ratio of investigation of correct plus adjacent

holes over total investigations during entire two minutes of the task ($p<0.05$; G). Likewise,

reversible (DREADDs) and chronic (caspase) disconnection of the CA1v to LS pathway did not

impair learning of the spatial water seeking task compared to controls, as measured by errors

before correct hole during task training (H) and latency to correct hole during task training (I),

but impaired memory retention of the water location during the probe (J). In contrast,

disconnection of the CA1v to LS pathway either reversibly (DREADDs) or chronically (caspase)

did not impair performance on the spatial escape task. There were no differences in learning as

measured by errors before correct hole during task training (K) and latency to correct hole during

task training (L). Unlike the spatial foraging task, retention of the spatial escape task was not

impaired by reversible nor chronic disconnection of the CA1v to LS pathway (M). For graphs

2E-G (CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8. For graphs

2H-J (CA1v to LS disconnect cohort 3), DREADDs n=8, caspase n=8, control n=7). For graphs

2K-M (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control n=10. Dotted

line indicates chance performance level (0.167). All values expressed as mean +/- SEM.

Figure 3

Figure 3. Neither reversible nor chronic CA1v to LS neural disconnection impair spatial

memory for escape location, social transmission of food preference, anxiety-like behavior,

or general locomotor activity levels. Diagram of the social transmission of food preference

(STFP) task (A). Neither reversible nor chronic disconnection of the CA1v to LS pathway impair

STFP learning compared to controls, as measured by a food preference ratio (B), with the dotted

line indicating chance preference level (0.50). Diagram of the zero maze apparatus (C). Anxiety-

like behavior was not influenced by reversible or chronic disconnection of the CA1v to LS

pathway compared to controls, as measured by performance in the zero maze task, specifically

time in open zones (D) and entries into open zones (E). Neither chronic nor reversible CA1v to

LS disconnection affected open field performance compared to controls, as measured by total

distance traveled (F). For graphs 3B, 3D, and 3E (CA1v to LS disconnect cohort 1), DREADDs

n=6, caspase n=10, control n=8. For graph 3F (CA1v to LS disconnect cohort 2), DREADDs

n=8, caspase n=12, control n=10. All values expressed as mean +/- SEM.

Figure 4

Figure 4. Neither CA1v to mPFC nor CA1v to LHA disconnection impairs spatial memory

for food location. Diagram of dual viral approach to identify collateral targets of the CA1v to

LS neural pathway (A). Representative CA1v injection site from collateral identification

approach (B; scale bar 500μm). Representative LS injection site from collateral identification

approach (C; scale bar 500μm). Representative image collateral axons of the CA1v to LS

pathway located in the mPFC (D; scale bar 50μm). Diagram of contralesional approach to

functionally disconnect the CA1v to mPFC neural pathway (E). Disconnection of the CA1v to

mPFC neural pathway did not influence learning of the spatial foraging task compared to

controls, as measured by errors before correct hole during training (F) and latency to correct hole

during training (G). Disconnection of the CA1v to mPFC pathway did not influence retention of

the spatial foraging task in the memory probe (H). Diagram of contralesional approach to

functionally disconnect the CA1v to LHA neural pathway (I). Disconnection of the CA1v to

LHA neural pathway did not influence learning of the spatial foraging task compared to controls,

as measured by errors before correct hole during training (J) and latency to correct hole during

training (K). Disconnection of the CA1v to LHA pathway did not influence retention of the

spatial foraging task in the memory probe (L). For graphs 4F-I, CA1v to mPFC disconnect n=12,

control n=12. For graphs 4K-N, CA1v to LHA disconnect n=12, control n=11. All values

expressed as mean +/- SEM.

Figure 5

A

Anterograde monosynaptic Cre-dependent AAV1-tdTomato

LS

CA1v

Anterograde transsynaptic AAV2/1-hSyn-Cre

B

ccb

VLa

LSc

LSr

LS injection site

C

PL

ILA

2° tdTomato+ fibers

D

LHAjd

fx

LHAd

LHAjvd

LHAsfa

LHAjvv

2° tdTomato+ fibers

E

ACBsh

aco

2° tdTomato+ fibers

F

CA1v → LS → multiple regions

Targeted side (right)

HIPPOCAMPAL REGION

RETROHIPPOCAMPAL REGION

CA1v

THALAMUS

LATERAL HYPOTHALAMIC AREA

SUBSTANTIA NIGRA

ORBITAL REGION

GLOBUS PALLIDUS

OLFACTORY REGION

LS

Contralateral side (left)

HIPPOCAMPAL REGION

CA1d

CA2

CA1v

LATERAL HYPOTHALAMIC AREA

CEREBRAL CORTEX

CORTICAL SUBPLATE

CEREBELLAR CORTEX

CORTICAL PLATE

PALLIDUM

THALAMUS

STRIATUM

MIDBRAIN

PONS

MEDULLA

SPINAL CORD

HYPOTHALAMUS

| 6 | 5 | 4 | 3 | 2 | 1 | 0 |

Aggregate ordinal value categories
(6 = strongest; 0 = weakest / absent)

Figure 5. Identification of second-order neural projections downstream of CA1v to LS

projections. Diagram of dual viral approach to identify brain regions that are second-order (2°)

targets of the CA1v to LS neural pathway (A). Representative LS injection site from second

order identification approach (B; scale bar 100μm). Representative image of second-order fibers

of the CA1v to LS pathway within the mPFC (C; scale bar 200μm). Representative image of

second-order fibers of the CA1v to LS pathway within the LHA (D; scale bar 200μm).

Representative image of second-order fibers of the CA1v to LS pathway within the ACB (E;

scale bar 200μm). Summary of the projection targets of LS neurons that receive input from

CA1v (F). The outputs of the right side of LS neurons receiving CA1v input are represented at

the macroscale (gray matter region resolution) on a partial flatmap representation of the rat

forebrain, adapted from (65). Connection weights are represented block colors for each region

following an ordinal scale ranging from weakest (0 = very weak or absent) to strongest (6 =

strong), as there were no 7 (very strong) values. The inset at lower left represents one side of the

brain with the part represented in the upper diagrams outlined in magenta.

SUPPLEMENTARY INFORMATION

Supplemental Figure 1

889   **Supplementary Figure 1. Effect of bilateral HPCv lesions on food intake and body weight.**

890   There were no effects on food intake (A) or body weight (B) of bilateral HPCv lesions compared

891   to controls. Lesion n = 11, control n = 18. All values expressed as mean +/- SEM.

Supplemental Figure 2

A

CA1v to LS disconnect cohort 1: Food intake

B

CA1v to LS disconnect cohort 2: Food intake

C

CA1v to LS disconnect cohort 1: Body weight

D

CA1v to LS disconnect cohort 2: Body weight

E

F

Supplementary Figure 2. Effect of reversible and chronic disconnection of the CA1v to LS

neural pathway on food intake and body weight, and verification of the chronic approach.

There were no effects on food intake (cohort 1: A; cohort 2: B) or body weight (cohort 1: C;

cohort 2: D) of reversible (DREADDs) or chronic (caspase) disconnection of the CA1v to LS

neural pathway compared to controls. Histological verification of the dual viral approach

demonstrates that cre-dependent tdTomato labeling of CA1v neurons induced by LS-origin retro-

cre is robust in a control animal (E), but reduced in an animal injected with cre-dependent

caspase combined with cre-dependent tdTomato due to caspase-induced cell death (F). For

graphs S1A-B (CA1v to LS disconnect cohort 1), DREADDs n=6, caspase n=10, control n=8.

For graphs S1C-D (CA1v to LS disconnect cohort 2), DREADDs n=8, caspase n=12, control

n=10. All values expressed as mean +/- SEM.

Supplemental Figure 3

Chronic Disconnection of the vCA1-LS Pathway

Acute Disconnection of the vCA1-LS Pathway

Supplementary Figure 3. Effect of reversible and chronic disconnection of CA1v to LS

neural pathway on feeding behavior. Compared with control animals, chronic caspase-

mediated disconnection of the CA1v to LS pathway did not significantly affect meal frequency

(A), meal size (B), meal duration (C), or 24h cumulative food intake (D) over the course of the

5-day recording period. Acute DREADDs-mediated disconnection of the CA1v to LS pathway

(via CNO infusion) resulted in non-significant trend towards increased meal frequency (E), no

differences in meal size (F) or meal duration (G), and a significant increase in 24h cumulative

food intake (H) in comparison with vehicle injection. All values expressed as mean +/- SEM.

Supplemental Figure 4

A

**CA1v to mPFC disconnect:
Food intake**

B

**CA1v to mPFC disconnect:
Body weight**

C

**CA1v to LHA disconnect:
Food intake**

D

**CA1v to LHA disconnect:
Body weight**

Supplementary Figure 4. Effect of contralesional disconnection methods on food intake and

body weight. There were no effects on food intake (A) or body weight (B) of contralesional

CA1v to mPFC compared to controls. CA1v to mPFC disconnect n=12, control n=12. There

were no effects on food intake (C) or body weight (D) of contralesional CA1v to LHA compared

to controls. CA1v to LHA disconnect n=12, control n=11. All values expressed as mean +/-

SEM.

REFERENCES

1.   R. D'Hooge, P. P. De Deyn, Applications of the Morris water maze in the study of
learning and memory. *Brain Res Brain Res Rev* **36**, 60-90 (2001).
2.   K. Gawel, E. Gibula, M. Marszalek-Grabska, J. Filarowska, J. H. Kotlinska, Assessment of
spatial learning and memory in the Barnes maze task in rodents-methodological
consideration. *Naunyn Schmiedebergs Arch Pharmacol* **392**, 1-18 (2019).
3.   R. G. M. Morris, U. Frey, "Hippocampal synaptic plasticity: Role in spatial learning or the
automatic recording of attended experience?" in Burgess, Neil (Ed); Jeffery, Kathryn J
(Ed); et al (1999) The hippocampal and parietal foundations of spatial cognition. (Oxford
University Press, U Edinburgh, Ctr for Neuroscience, Edinburgh, Scotland London, 1999),
pp. 220-246.
4.   D. C. Rowland, Y. Roudi, M. B. Moser, E. I. Moser, Ten Years of Grid Cells. *Annu Rev*
Neurosci **39**, 19-40 (2016).
5.   R. Morris, Developments of a water-maze procedure for studying spatial learning in the
rat. *Journal of neuroscience methods* **11**, 47-60 (1984).
6.   C. A. Barnes, Memory deficits associated with senescence: a neurophysiological and
behavioral study in the rat. *J Comp Physiol Psychol* **93**, 74-104 (1979).
7.   J. O'Keefe, J. Dostrovsky, The hippocampus as a spatial map: preliminary evidence from
unit activity in the freely-moving rat. *Brain research* (1971).
8.   T. Hafting, M. Fyhn, S. Molden, M. B. Moser, E. I. Moser, Microstructure of a spatial map
in the entorhinal cortex. *Nature* **436**, 801-806 (2005).
9.   M. S. Fanselow, H.-W. Dong, Are the dorsal and ventral hippocampus functionally
distinct structures? *Neuron* **65**, 7-19 (2010).
10.   A. T. Keinath *et al.*, Precise spatial coding is preserved along the longitudinal
hippocampal axis. *Hippocampus* **24**, 1533-1548 (2014).
11.   K. B. Kjelstrup *et al.*, Finite scale of spatial representation in the hippocampus. *Science*
321, 140-143 (2008).
12.   L. de Hoz, J. Knox, R. G. Morris, Longitudinal axis of the hippocampus: both septal and
temporal poles of the hippocampus support water maze spatial learning depending on
the training protocol. *Hippocampus* **13**, 587-603 (2003).
13.   J. Ferbinteanu, C. Ray, R. J. McDonald, Both dorsal and ventral hippocampus contribute
to spatial learning in Long-Evans rats. *Neurosci Lett* **345**, 131-135 (2003).
14.   M. B. Moser, E. I. Moser, Functional differentiation in the hippocampus. *Hippocampus* **8**,
608-619 (1998).
15.   S. E. Kanoski, H. J. Grill, Hippocampus Contributions to Food Intake Control: Mnemonic,
Neuroanatomical, and Endocrine Mechanisms. *Biol Psychiatry* **81**, 748-756 (2017).
16.   P. G. Henke, Hippocampal pathway to the amygdala and stress ulcer development. *Brain*
Res Bull **25**, 691-695 (1990).
17.   K. G. Kjelstrup *et al.*, Reduced fear expression after lesions of the ventral hippocampus.
Proceedings of the National Academy of Sciences **99**, 10825-10830 (2002).

18.   S. Gaskin, A. Gamliel, M. Tardif, E. Cole, D. G. Mumby, Incidental (unreinforced) and
reinforced spatial learning in rats with ventral and dorsal lesions of the hippocampus.
Behav Brain Res **202**, 64-70 (2009).

19.   M. S. Bienkowski *et al.*, Integration of gene expression and brain-wide connectivity
reveals the multiscale organization of mouse hippocampal networks. *Nat Neurosci* **21**,
1628-1643 (2018).

20.   R. Hannapel *et al.*, Postmeal Optogenetic Inhibition of Dorsal or Ventral Hippocampal
Pyramidal Neurons Increases Future Intake. *eNeuro* **6** (2019).

21.   R. C. Hannapel, Y. H. Henderson, R. Nalloor, A. Vazdarjanova, M. B. Parent, Ventral
hippocampal neurons inhibit postprandial energy intake. *Hippocampus* **27**, 274-284
(2017).

22.   B. K. Mani *et al.*, Neuroanatomical characterization of a growth hormone secretagogue
receptor-green fluorescent protein reporter mouse. *J Comp Neurol* **522**, 3644-3666
(2014).

23.   J. M. Zigman, J. E. Jones, C. E. Lee, C. B. Saper, J. K. Elmquist, Expression of ghrelin
receptor mRNA in the rat and the mouse brain. *J Comp Neurol* **494**, 528-548 (2006).

24.   I. Merchenthaler, M. Lane, P. Shughrue, Distribution of pre-pro-glucagon and glucagon-
like peptide-1 receptor messenger RNAs in the rat central nervous system. *J Comp
Neurol* **403**, 261-280 (1999).

25.   S. E. Kanoski, S. M. Fortin, K. M. Ricks, H. J. Grill, Ghrelin signaling in the ventral
hippocampus stimulates learned and motivational aspects of feeding via PI3K-Akt
signaling. *Biol Psychiatry* **73**, 915-923 (2013).

26.   S. E. Kanoski *et al.*, Hippocampal leptin signaling reduces food intake and modulates
food-related memory processing. *Neuropsychopharmacology* **36**, 1859-1870 (2011).

27.   T. M. Hsu *et al.*, A hippocampus to prefrontal cortex neural pathway inhibits food
motivation through glucagon-like peptide-1 signaling. *Mol Psychiatry* **23**, 1555-1565
(2018).

28.   T. M. Hsu, J. D. Hahn, V. R. Konanur, A. Lam, S. E. Kanoski, Hippocampal GLP-1 receptors
influence food intake, meal size, and effort-based responding for food through volume
transmission. *Neuropsychopharmacology* **40**, 327-337 (2015).

29.   A. N. Suarez, C. M. Liu, A. M. Cortella, E. E. Noble, S. E. Kanoski, Ghrelin and Orexin
Interact to Increase Meal Size Through a Descending Hippocampus to Hindbrain
Signaling Pathway. *Biol Psychiatry* **87**, 1001-1011 (2020).

30.   G. D. Petrovich, N. S. Canteras, L. W. Swanson, Combinatorial amygdalar inputs to
hippocampal domains and hypothalamic behavior systems. *Brain research reviews* **38**,
247-289 (2001).

31.   C. De La Rosa-Prieto *et al.*, Subicular and CA1 hippocampal projections to the accessory
olfactory bulb. *Hippocampus* **19**, 124-129 (2009).

32.   T. M. Hsu *et al.*, Hippocampus ghrelin receptor signaling promotes socially-mediated
learned food preference. *Neuropharmacology* **131**, 487-496 (2018).

33.   L. W. Swanson, W. M. Cowan, An autoradiographic study of the organization of the
efferent connections of the hippocampal formation in the rat. *J Comp Neurol* **172**, 49-84
(1977).

34.   T. M. Hsu *et al.*, Hippocampus ghrelin signaling mediates appetite through lateral
hypothalamic orexin pathways. *Elife* **4** (2015).
35.   A. Arszovszki, Z. Borhegyi, T. Klausberger, Three axonal projection routes of individual
pyramidal cells in the ventral CA1 hippocampus. *Front Neuroanat* **8**, 53 (2014).
36.   P. Y. Risold, L. W. Swanson, Structural evidence for functional domains in the rat
hippocampus. *Science (New York, N.Y.)* **272**, 1484-1486 (1996).
37.   P. Sweeney, Y. Yang, An excitatory ventral hippocampus to lateral septum circuit that
suppresses feeding. *Nat Commun* **6**, 10188 (2015).
38.   R. Garcia, R. M. Vouimba, R. Jaffard, Spatial discrimination learning induces LTP-like
changes in the lateral septum of mice. *Neuroreport: An International Journal for the*
Rapid Communication of Research in Neuroscience **5**, 329-332 (1993).
39.   R. Jaffard, R. M. Vouimba, A. Marighetto, R. Garcia, Long-term potentiation and long-
term depression in the lateral septum in spatial working and reference memory. *J*
Physiol Paris **90**, 339-341 (1996).
40.   S. J. Terrill *et al.*, Nucleus accumbens melanin-concentrating hormone signaling
promotes feeding in a sex-specific manner. *Neuropharmacology* **178**, 108270 (2020).
41.   A. Carballo-Marquez, A. Vale-Martinez, G. Guillazo-Blanch, M. Marti-Nicolovius,
Muscarinic receptor blockade in ventral hippocampus and prelimbic cortex impairs
memory for socially transmitted food preference. *Hippocampus* **19**, 446-455 (2009).
42.   R. A. Countryman, N. L. Kaban, P. J. Colombo, Hippocampal c-fos is necessary for long-
term memory of a socially transmitted food preference. *Neurobiol Learn Mem* **84**, 175-
183 (2005).
43.   G. M. Parfitt *et al.*, Bidirectional Control of Anxiety-Related Behaviors in Mice: Role of
Inputs Arising from the Ventral Hippocampus to the Lateral Septum and Medial
Prefrontal Cortex. *Neuropsychopharmacology : official publication of the American*
College of Neuropsychopharmacology **42**, 1715-1728 (2017).
44.   N. L. Trent, J. L. Menard, The ventral hippocampus and the lateral septum work in
tandem to regulate rats' open-arm exploration in the elevated plus-maze. *Physiology &*
behavior **101**, 141-152 (2010).
45.   J. C. Jimenez *et al.*, Anxiety Cells in a Hippocampal-Hypothalamic Circuit. *Neuron* **97**,
670-683.e676 (2018).
46.   M. S. Fanselow, H. W. Dong, Are the dorsal and ventral hippocampus functionally
distinct structures? *Neuron* **65**, 7-19 (2010).
47.   M. Contreras, T. Pelc, M. Llofriu, A. Weitzenfeld, J. M. Fellous, The ventral hippocampus
is involved in multi-goal obstacle-rich spatial navigation. *Hippocampus* **28**, 853-866
(2018).
48.   S. L. T. Lee, D. Lew, V. Wickenheisser, E. J. Markus, Interdependence between dorsal and
ventral hippocampus during spatial navigation. *Brain Behav* **9**, e01410 (2019).
49.   D. P. Holschneider *et al.*, Cerebral perfusion mapping during retrieval of spatial memory
in rats. *Behavioural brain research* **375**, 112116 (2019).
50.   D. Olivo, M. Caba, F. Gonzalez-Lima, J. F. Rodríguez-Landa, A. A. Corona-Morales,
Metabolic activation of amygdala, lateral septum and accumbens circuits during food
anticipatory behavior. *Behav Brain Res* **316**, 261-270 (2017).

51. M. Carus-Cadavieco *et al.*, Gamma oscillations organize top-down signalling to
hypothalamus and enable food seeking. *Nature* **542**, 232-236 (2017).
52. S. J. Terrill *et al.*, Role of lateral septum glucagon-like peptide 1 receptors in food intake.
American journal of physiology. Regulatory, integrative and comparative physiology **311**,
R124-132 (2016).
53. S. J. Terrill, K. D. Wall, N. D. Medina, C. B. Maske, D. L. Williams, Lateral septum growth
hormone secretagogue receptor affects food intake and motivation for sucrose
reinforcement. *American journal of physiology. Regulatory, integrative and comparative*
physiology **315**, R76-r83 (2018).
54. C. Wang, C. M. Kotz, Urocortin in the lateral septal area modulates feeding induced by
orexin A in the lateral hypothalamus. *American journal of physiology. Regulatory,*
integrative and comparative physiology **283**, R358-367 (2002).
55. J. Hauser, L. H. Llano López, J. Feldon, P. A. Gargiulo, B. K. Yee, Small lesions of the dorsal
or ventral hippocampus subregions are associated with distinct impairments in working
memory and reference memory retrieval, and combining them attenuates the
acquisition rate of spatial reference memory. *Hippocampus* **30**, 938-957 (2020).
56. D. S. Olton, The radial arm maze as a tool in behavioral pharmacology. *Physiology &*
behavior **40**, 793-797 (1987).
57. H. H. Pothuizen, W. N. Zhang, A. L. Jongen-Rêlo, J. Feldon, B. K. Yee, Dissociation of
function between the dorsal and the ventral hippocampus in spatial learning abilities of
the rat: a within-subject, within-task comparison of reference and working spatial
memory. *The European journal of neuroscience* **19**, 705-712 (2004).
58. L. E. Jarrard, L. P. Luu, T. L. Davidson, A study of hippocampal structure-function
relations along the septo-temporal axis. *Hippocampus* **22**, 680-692 (2012).
59. L. W. Swanson, Brain maps 4.0-Structure of the rat brain: An open access atlas with
global nervous system nomenclature ontology and flatmaps. *J Comp Neurol* **526**, 935-
943 (2018).
60. R. C. Ritter, P. G. Slusser, S. Stone, Glucoreceptors controlling feeding and blood glucose:
location in the hindbrain. *Science* **213**, 451-452 (1981).
61. E. E. Noble *et al.*, Hypothalamus-hippocampus circuitry regulates impulsivity via
melanin-concentrating hormone. *Nat Commun* **10**, 4923 (2019).
62. B. Zingg *et al.*, AAV-Mediated Anterograde Transsynaptic Tagging: Mapping
Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron* **93**, 33-
47 (2017).
63. A. N. Suarez *et al.*, Gut vagal sensory signaling regulates hippocampus function through
multi-order pathways. *Nat Commun* **9**, 2181 (2018).
64. J. D. Hahn, O. Sporns, A. G. Watts, L. W. Swanson, Macroscale intrinsic network
architecture of the hypothalamus. *Proc Natl Acad Sci U S A* **116**, 8018-8027 (2019).
65. J. D. Hahn, Swanson, L. W,. Bowman, I., Foster, N. N., Zingg, B., Bienkowski, M. S.,
Hintiryan, H., & Dong, H.-W. , An open access mouse brain flatmap and upgraded rat
and human brain flatmaps based on current reference atlases. *J. Comp. Neurol.* (2020).
66. L. E. Herzog *et al.*, Interaction of Taste and Place Coding in the Hippocampus. *The*
Journal of neuroscience : the official journal of the Society for Neuroscience **39**, 3057-
3069 (2019).

67.   R. A. Countryman, P. E. Gold, Rapid forgetting of social transmission of food preferences
in aged rats: relationship to hippocampal CREB activation. *Learn Mem* **14**, 350-358
(2007).
68.   B. G. Galef, Jr., W. Y. Lee, E. E. Whiskin, Lack of interference in long-term memory for
socially learned food preferences in rats (Rattus norvegicus). *J Comp Psychol* **119**, 131-
135 (2005).
69.   B. G. Galef, Jr., E. E. Whiskin, Socially transmitted food preferences can be used to study
long-term memory in rats. *Learn Behav* **31**, 160-164 (2003).
1107