Running head: Cod destabilized by fluctuating oceans

Trade-offs of managing Arctic predator harvest instability in fluctuating oceans

Daisuke Goto[1,*], Anatoly A. Filin[2], Daniel Howell[1], Bjarte Bogstad[1], Yury Kovalev[2], and

Harald Gjøsæter[1]

[1]Institute of Marine Research, P.O. Box 1870, Nordnes, 5817 Bergen, Norway

[2]Polar Branch of the Federal State Budget Scientific Institution, Russian Federal Research

Institute of Fisheries and Oceanography ("PINRO" named after N.M. Knipovich), Akademik

Knipovich Street 6, 183038 Murmansk, Russia

[*]Corresponding author: daisuke.goto2@gmail.com

**Abstract.** Sustainable human exploitation of marine living resources stems from a delicate

balance between short-term yield stability and long-term population persistence to achieve

socioeconomic and conservation goals. However, imperfect knowledge of how oscillations in

ecosystem processes regulate fluctuations in exploited populations can obscure the risk of

missing management targets. We illustrate how the harvest policy to suppress short-term

yield fluctuation inadvertently disrupts population cycles and yield stability of exploited,

long-lived predators under stochastically fluctuating environmental forces (food availability

and regional climate) using Northeast Arctic (NEA) cod (*Gadus morhua*, an apex predatory

fish) as a case study. We use a stochastic, empirically parameterized multispecies model to

simulate NEA cod population dynamics through life-history processes; Barents Sea capelin

(*Mallotus villosus*, a pelagic forage fish) modulates cod productivity through density-

dependent cannibalism–predation dynamics, whereas sea temperature regulates cod

consumption, growth, and recruitment. We first test how capelin and sea temperature

fluctuations regulate patterns in cod yield fluctuation and then quantitatively assess how

fishing pattern designed to limit yield between-year variance (within 50–5%) perturbs cod

population–catch dynamics. Simulations suggest that capelin and temperature interactively

contribute to shifting cyclic patterns in cod yield fluctuation primarily through cod

cannibalism–predation dynamics. Wavelet analyses further show that muffling yield variance

(30 % or less) reshapes the cyclicity (shorter period and greater amplitude) of cod population

size and demography, thereby becoming progressively unsynchronized with fishing pressure.

Our work reveals unintended consequences of managing transient dynamics of fished

populations: the interworking of population cycle destabilized by inadvertently intensifying

fishing pressure, amplifying yield fluctuation and, in turn, elevating overharvest risk when

not accounting for compounded effects of stochasticity in ecologically connected processes.

These policy implications underscore the need for an ecosystem approach to designing

ecologically sound management measures to safely harvest shared living resources while

achieving socioeconomic security in increasingly more dynamic oceans in the Arctic and

elsewhere.

**Keywords**: *bottom-up control; food web; ecosystem-based management; forage fish; early*

*life history; climate anomalies; cod; cohort resonance; structural uncertainty; process-based*

*models*

43                                         INTRODUCTION

Fluctuation in wild animal populations has posed myriad socioeconomic and conservation

challenges in sustainably managing human exploitation of living natural resources for

centuries (Hjort 1914, Fryxell et al. 2010, Carpenter et al. 2015). In commercial exploitation

of oceanic species such as long-lived, apex predators, stable population dynamics provide not

only predictability to harvesters and seafood processors but also food, nutrition, and

employment security in fishing nations (Garcia and Rosenberg 2010, Hicks et al. 2019). In

modern industrial fisheries, harvest strategies are designed to maintain population abundance

of exploited species above a threshold that prevents overharvest and sustains its harvest while

suppressing yield fluctuation to achieve socioeconomic goals (Caddy and Seijo 2005, Nielsen

et al. 2018). However, oscillations in environmental forces may amplify or dampen

fluctuation in fished populations (Shelton and Mangel 2011, Rouyer et al. 2012). Further, the

demography of these populations has often been modified by long-term perturbations such as

fishing-induced selective reduction in adult survival (Hidalgo et al. 2011). Harvest strategies

that fail to account for the dynamic interplay of environmental and demographic fluctuations

may, thus, lead to exploitation intensity that is unsynchronized with natural population

cycles, potentially risking sustainability of populations and fishing operation (Hall and

Mainprize 2004, Carpenter et al. 2015) and, in turn, incurring ecological and socioeconomic

damages (Caddy and Seijo 2005, Nielsen et al. 2018).

Population fluctuation in oceanic predators can arise from stochasticity in demographic

states that reflect life-history responses (such as reproductive and migration timing) to large-

scale climate variability–an overarching force that governs dynamic habitat conditions

(temperature, pH, dissolved oxygen, etc.) over varying spatial and temporal scales (Bjørnstad

and Grenfell 2001, Ottersen et al. 2001, Cury et al. 2008). These dynamics often propagate

through species interactions to modulate productivity of fished populations (Ottersen et al.

2001, Stige et al. 2010). Environmentally driven plankton production, for example, regulates

forage fish production that support exploited apex predators (Ware and Thomson 2005).

Large between-year fluctuation can emerge particularly in early life stages of marine

populations (Leggett and Deblois 1994); seasonal to decadal variation in ocean conditions

propagates through predator–prey interactions, amplifying or attenuating variance in year-

class strength of exploited predators (Edwards and Richardson 2004). Stochasticity filtered

through such interactions can, thus, emerge as complex, nonlinear patterns in population

cycles (Bjørnstad and Grenfell 2001) and pose a challenge in designing management

measures to suppress yield variance in exploited predators (Essington et al. 2018).

Trophic interactions between cod (*Gadus morhua*) and capelin (*Mallotus villosus*) in the

North Atlantic present such dynamics that modulate short-term fluctuation in cod fishery

yield (Link et al. 2009). Productivity of cod–a long-lived, demersal (bottom-water) predatory

fish–often fluctuates from year to year with variability in both density -dependent and -

independent ecosystem (bottom-up) processes including interactions with other species

during early life stages (Lilly et al. 2008, Minto and Worm 2012). In particular, food shortage

can enhance cannibalism intensity (Yaragina et al. 2009). Capelin–a short-lived ($\leq$ 5 years),

semelparous, widely distributed pelagic planktivore–comprises a large part of adult cod diet

(Holt et al. 2019) and may influence the intensity of cod cannibalism on the young ($\leq$ 3-year-

olds) and, in turn, recruitment pulses. Moreover, owing to migratory behaviors driven by

ocean conditions and plankton distribution, time-varying capelin aggregation may reflect

changes in ecosystem processes including systemwide shifts in ice phenology, thermal

regimes, and plankton productivity (Carscadden et al. 2001, Rose 2005). Cod–capelin spatial

and temporal overlap in cod spawning and nursery grounds may, thus, modulate cod

population and yield fluctuations (Ciannelli and Bailey 2005, Johannesen et al. 2012). It

remains unclear, however, how fluctuations in bottom-up processes such as forage fish

production shape the population behavior and yield stability of intensely exploited predators

such as cod, which are concurrently regulated by annual fishing pattern designed to achieve

management goals.

Here we illustrate how the harvest policy of suppressing yield fluctuation can reshape the

population cycle of exploited predators in stochastically fluctuating environments using a

well-studied species, Northeast Arctic (NEA) cod, as a real-world case study. NEA cod,

distributed over the Barents Sea and its surrounding areas (Fig. 1a), is the world's largest

supporting two major fishing nations, Norway and Russia (Kjesbu et al. 2014). The fishery

dates back to the period when the northern coasts of these nations were likely populated after

the end of the last ice age. An international industrial trawl fishery in the Barents Sea

developed in the early 1900s and total catches by all gears peaked at more than 1.2 million t

in catch in some years in the period 1955–1975 (ICES 2019). Following the establishment of

200-mile economic zones in the late 1970s, the fishery by Norway and Russia dominated

(ICES 2019) and subsequently depleted the population prior to the establishment of a harvest

strategy by the Norwegian-Russian Fishery Commission in the early 2000s (ICES 2019).

This harvest strategy (which was in force until 2016) sets target exploitation rate with

precautionary measures while achieving short-term yield stability such as limiting between-

year catch fluctuation (to less than 10%). These harvest rules, along with favorable habitat

quality (Ottersen and Stenseth 2001, Stige et al. 2010), are suspected to allow the population

to balloon with the quota reaching over one million t in 2014, the highest in more than three

decades (Kjesbu et al. 2014).

Using a stochastic, process-based multispecies model previously developed for the Barents

Sea cod–capelin complex (Filin 2009, Howell et al. 2013), we test 1) how climatic oscillation

propagates through an interplay of cod cannibalism–capelin productivity (Hjermann et al.

2007) and 2) how fishing pattern to suppress cod yield fluctuation shapes the cyclicity of

population size and demographic fluctuations and, in turn, the potential to achieve

management targets in stochastically fluctuating environments. By using modeled oscillatory

fluctuations in sea temperature in the Kola section (Fig 1b,d)–an indicator for reginal climate

(Ingvaldsen et al. 2003) and capelin productivity (Fig. 1c,e)–proxy for the ecosystem carrying

capacity (Gjøsæter 1998), in effect, our work aims to quantitatively evaluate implications of

adapting an ecosystem approach to provisioning of annual catch targets to ensure sustainable

exploitation of marine resources while conserving ecological integrity under fluctuating

ocean conditions.

126                                    METHODS

127                                   *Study system*

The Barents Sea cod–capelin complex is among the few co-managed stocks in the North

Atlantic (Gjøsæter et al. 2012). Historically, capelin has been the major forage species

supporting exploited apex predators in the Barents Sea food web (Gjøsæter 1998); its stock

biomass has been measured by acoustic methods to reach up to seven million t in the 1970s

and 1990s (Gjøsæter 1998). Capelin–a small, pelagic, schooling fish, rarely reaching lengths

above 20 cm and ages above five-year-olds–feeds in the northern Barents Sea on meso- and

macro- zooplankton but migrates to the northern coasts of Norway and Russia to spawn in

spring. The forage fish is semelparous and thus magnifies its role of bringing energy from the

northern parts of the Barents Sea to the southern coastal areas. Although capelin supplies a

food source for many piscivores in the Barents Sea, Northeast Arctic (NEA) cod are by far

their primary predator, consuming up to more than three million t per year (Dolgov 2002).

Capelin fluctuation may, thus, regulate the intensity of cod cannibalism and year-class

strength, likely contributing to depensatory effects of depleted food supply (Hjermann et al.

2007). For example, amplified year-to-year fluctuations in capelin biomass in the 1980s and

1990s were largely responsible for the NEA cod biomass rebuilding; cod cannibalism peaked

following capelin collapses (Hjermann et al. 2007). The collapses, particularly the first one,

propagated through the rest of the Barents Sea food web to a variable degree, depending on

availability of other food (Hjermann et al. 2007, Gjøsæter et al. 2009).

*Food-web modeling*

To simulate the Barents Sea cod-capelin dynamics, we used a stochastic, food-web model

that mechanistically accounts for ecosystem processes (a model of intermediate complexity

for ecosystem, Plagányi et al. 2014)–STOCOBAR (Stock of Cod in the Barents Sea). The

model has been developed previously at Russian Federal Research Institute of Fisheries and

Oceanography (Filin 2009); the version used in this study has been described fully in Howell

et al. (2013) (Fig. 1f). For brevity, here we outline the model structure; STOCOBAR is

designed to capture age-structured population dynamics of NEA cod in Barents Sea and

adjacent areas through yearly life history processes: growth, feeding, maturation, recruitment,

natural mortality (including cannibalism on juveniles), and fishing mortality (by single

aggregate fleet fishing operation) (Howell et al. 2013). Individual fish growth is

mechanistically simulated through energetics by converting consumed food to body mass,

which, in turn, regulates maturation (Howell et al. 2013). The model projects cod population

dynamics and evaluates alternative management strategies (harvest measures and recovery

plans) by accounting for variability in ecosystem processes such as lower trophic group (one

or more prey) productivity (Howell et al. 2013). In this study, we incorporated stochasticity

explicitly in cod recruitment, capelin productivity, and sea temperature. We parameterized

the model with 30-year (1981–2010) annual bottom trawl and acoustic survey data of age-

specific life-history traits including maturity rates (proportions of adults) and body masses

(kg), and fit it to time series of cod age-specific abundance indices (catch per unit effort) and

commercial fishery catch (t), and capelin biomass (t), which were all taken from annual

assessments by the International Council for the Exploration of the Sea (ICES) Arctic

Fisheries Working Group; the survey and sample processing methods used are extensively

documented in ICES (2010).

Cod recruitment (survival of 3-yr-olds) in the model is stochastically simulated and

depends directly on 1-year-old cod abundance and cannibalism and indirectly on capelin

biomass and sea temperature. We simulated 1-yr-old cod abundance using a density- and

temperature- dependent function relating it to spawner abundance (spawning stock biomass,

SSB, t) with the Ricker model (Ricker 1958) and mortality by cannibalism (before

recruitment). SSB is computed by multiplying age-specific numbers by age-specific masses

and maturity rates. The strength of cod cannibalism fluctuates over time and was modeled

with a function of capelin biomass and cod population structure (size and age composition)

(Fig. 1f, Howell et al. 2013). Although sea temperature may affect cannibalism intensity

indirectly through consumption, we assumed that this effect is negligible within the historical

temperature range used in this study. Further, we did not include indirect temperature effects

on cannibalism through spatial distribution (Howell and Filin 2014). Based on model fitting,

we capped mortality by cannibalism at 90%, 60%, and 40% of total mortality of 1-, 2-, and 3-

yr-olds (respectively). We then added random noise to recruitment using normally distributed

residuals generated from the 1980–2007 hindcast simulations (Howell et al. 2013).

185                     *Modeling capelin and sea temperature fluctuations*

We used capelin biomass as proxy for cod carrying capacity, representing "good" and

"poor" years of habitat conditions that regulate cod growth and cannibalism. The model uses

capelin as a primary food source for cod with all other prey (young Northeast Arctic haddock

*Melanogrammus aeglefinus*, young herring *Clupea harengus*, polar cod *Boreogadus saida*,

shrimp *Pandalus borealis*, and krill Euphausiidae) being aggregated as "other food'. We

statistically modeled between-year variability in capelin production (as an aggregate of 1- to

5- yr-olds) to capture observed patterns of declining biomass 1) when cod SSB exceeding

400,000 t or 2) when capelin biomass falling below three million t in the previous year (no

feedback for cod predation on capelin), while capelin biomass is capped at six million t when

cod SSB exceeds 800,000 t. Modeled capelin biomass stochastically varies between years and

simulations (Fig. 1c), generating low-frequency oscillations (Fig. 1e)

Sea temperature regulates cod population dynamics through recruitment, growth, and

consumption to capture variable metabolic rate in the model (Fig. 1f). In this study, we

primarily focused on short-term sea temperature fluctuations (rather than multidecadal shifts),

a time scale relevant for provisioning of tactical advice in fisheries management. The model

uses a stochastically simulated thermal regime that reflects the historical fluctuation of annual

mean temperatures (AMT) in the Kola Section in the Barents Sea, which also roughly

represents climate variability in the region (Ingvaldsen et al. 2003). To create a temperature

anomaly scenario, we first split the observed AMT data (1950-2017) into three groups; cold

(AMT < 3.6°C), moderate (3.6°C ≤ AMT ≤ 4.2°C), and warm (AMT > 4.2°C) years. Then,

AMT data were randomly selected from these three groups sequentially. The modeled

thermal periods (cold, moderate, and warm) last from one to five years; the length of each

period was set randomly (effectively generating roughly decadal oscillations, Fig. 1b,d).

209                              *Cod harvest management model and scenarios*

210  We simulated dynamic cod mortality by fishing with a harvest management model based on

211 the harvest rule set for NEA cod (prior to changes in 2016, ICES 2016) and evaluated cod

212 population dynamics and between-year yield variability under alternative harvest scenarios.

213 Under the harvest rule, target exploitation rate ($F_{target}$) is set to 0.40 to project annual fishery

214 catch (or total allowable catch, TAC) when spawner abundance (SSB) remains above

215 460,000 t. These reference values ($F_{pa}$ and $B_{pa}$, respectively) are designed to take

216 precautionary measures to prevent overexploitation by accounting for uncertainty in

217 population and harvest dynamics (Rindorf et al. 2016). When SSB falls below $B_{pa}$, however,

218 $F_{target}$ is adjusted to $F_{pa}$ scaled to the proportion of SSB relative to $B_{pa}$ as:

$$F_{target} = F_{pa} \frac{SSB}{B_{pa}}$$

219 In simulations, $F_{target}$ was applied by computing age-specific fishing mortality rate as:

$$F_{ji} = \frac{F_{target\,i}}{F'} F_j^{'}$$

220 where $F_{ji}$ is instantaneous rate of fishing mortality of $j$-yr-olds in year $i$; $F_{target\,i}$ is target

221 exploitation rate in year $i$, computed in the model according to the harvest rule; $F'$ is mean

222 fishing mortality rate of adults (a fixed age range, 5- to 10- yr-olds, is used for this

223 computation) in year 1; and $F_j^{'}$ is fishing mortality rate of $j$-yr-olds in year 1.

224  The harvest policy to suppress yield fluctuation (hereafter stability constraint) is designed

225 to provide stable and predictable catch forecasts (and revenue) for harvesters and seafood

226 processors. We computed between-year variation in catch forecasts (or interannual catch

227 variability, $IAV$) as

$$IAV_i = \frac{Catch_{i+1} - Catch_i}{Catch_i} \times 100$$

228 where $IAV_i$ is a relative change in catch forecast (%) from year $i$ to $i+1$; $Catch_{i+1}$ and $Catch_i$

229 are catch forecasts in year $i+1$ and $i$ (respectively). However, the precautionary principle also

applies to this policy; the constraint is applied only when SSB equals to or exceeds $B_{pa}$. In

this study, if this rule results in *IAV* exceeding the set proportion (±%), the model algorithm

searches for $F_{target}$ until it finds the value that does not exceed the boundary set by the

constraint. The constraint was not applied in the first year after the recovery period (when

SSB ≤ $B_{pa}$); $F_{target}$ was set to $F_{pa}$ regardless of catch forecasts in the previous year.

To first evaluate how ecological uncertainty involving capelin and climate variability

contributes to between-year variance in cod catch forecasts under stability constraint, we

tested performance of the following four model configurations that represent ecological

assumptions on cod predation–cannibalism dynamics under a 10% constraint with variable

sea temperature: M1) constant capelin biomass (one million t) without cannibalism

(temperature fluctuation only), M2) varying capelin biomass without cannibalism, M3)

constant capelin biomass with cannibalism, and M4) varying capelin biomass with

cannibalism.

To evaluate how suppressing short-term variation in catch forecasts drives cod population

cycles and yield stability under variable capelin productivity and sea temperature, we

analyzed the following nine harvest scenarios of limiting between-year variation in catch

forecasts; ±0%, ±5%, ±10%, ±15%, ±20%, ±25%, ±30%, ±40%, and ±50%.

We projected 100-year population and harvest dynamics to evaluate ecological assumptions

and harvest scenarios (25 x (4+9) simulations). We analyzed each scenario based on 25

realizations because preliminary simulations showed that mean catches vary less than ±1%

with more than 25 simulations. We disregarded the first 20 yrs of model projections as a

'burn-in' to minimize effects of the initial conditions and evaluated the outputs from years

21–100. Because this study primarily focuses on implications of suppressing yield

fluctuations, we did not incorporate other sources of uncertainty such as population

estimation error and implementation error (annual yields were assumed equal to catch

forecasts). We set fishing selectivity (age-dependent gear selectivity) as time-invariant in the

first year. For clarity, we excluded the following harvest rules set for NEA cod; setting catch

forecasts for three years by applying target exploitation rate to current SSB ("three-year

forecast"), fixing the lower range of $F_{target}$, and raising $F_{target}$ when SSB becomes

exceptionally high (ICES 2019). In analyzing the harvest scenarios, we computed the

following six cod demographic state variables and management targets from model

projections: harvestable (3-yr-olds and older) population biomass (stock size, t), SSB (t),

recruitment, catch forecasts (t), mean fishing mortality, and risk (%). Risk is defined as the

probability of SSB falling below the biological limit threshold ($B_{lim}$). When SSB < $B_{lim}$, the

reproductive capacity of a population is expected to decline (Rindorf et al. 2016); harvest

rules with risk > 5% are considered unacceptable (non-precautionary) for sustainable

exploitation. For NEA cod, $B_{lim}$ is set to 220,000 t (ICES 2016).

267                                          *Spectral analyses*

To explore potential mechanisms underlying oscillatory behaviors of projected cod

population and yield fluctuations, we performed wavelet spectral analysis on model outputs

from select scenarios of environmental forces (sea temperature and capelin) and harvest

strategies (±0%, ±10%, ±20%, and ±30% stability constraints). Wavelet spectral analysis is

suited for analyzing nonlinear and non-stationary time series (Cazelles et al. 2014) such as

population dynamics of exploited species. We computed continuous Morlet wavelet power

(the square of amplitudes) spectra to transform signals in the projected time series (years 21–

100) as a function of frequency (periodicity) and time from 1000 bootstrap simulations

(randomized surrogate time series) (Roesch and Schmidbauer 2018). We derived times series

of the following five cod demographic state variables and management targets; stock size,

SSB, recruitment, catch forecasts, and mean fishing mortality. Further, to assess

(a)synchronicity between cod population and harvest dynamics, we performed cross-wavelet

analysis to test for coherency (covariation in frequency) (Cazelles et al. 2014) between stock

size and mean fishing mortality time series. All wavelet analyses were performed in R

(version 3.6.2, R Development Core Team 2019) with the WaveletComp R package v.1.1

(Roesch and Schmidbauer 2018).

284                                        RESULTS

285                        *Ecological uncertainty in cod harvest dynamics*

Under the baseline scenario, cod population dynamics reached dynamic equilibrium by year

20 with, on average, ~2.3 million t harvestable (3-yr-olds and older) biomass (stock

size),~1.1 million t spawners (SSB), 694 million recruits (3-yr-olds), and 680,000 t/yr catch

(Table 1 and Appendix 1: Fig. S1), which all aligned closely with the historical population

status and harvest of NEA cod (ICES 2017). Overall, between-year variability in cod body

masses, maturity rates, and consumption rates declined with age (from ~39% to ~7%,

Appendix 1: Fig. S1 and S2). Cod diet, on average, comprised of 10% juvenile cod (more

than 66% as 1-yr-olds), 21% capelin, and 69% others (Appendix 1: Fig. S2).

Incorporating additional ecological reality (time-varying capelin productivity and cod

cannibalism) in the model showed nonstationary patterns (varying amplitude and periodicity)

of cod catch forecasts under a 10% constraint with varying sea temperature (Fig. 2). With

constant capelin productivity and no cod cannibalism (M1), wavelet analyses showed a ~8-yr

cycle with modest amplitudes in catch forecasts (reflecting one of dominant sea temperature

signals, Fig. 2a and Appendix 1: Fig. 3a) with between-year variation of less than 2%/yr (Fig.

2e,f). With varying capelin productivity (M2), catch forecasts showed two (8 to 10- and 32-

yr) weaker cycles (roughly reflecting dominant capelin signals, Appendix 1: Fig. 3b), and

between-year variation rose only by ~4% (Fig. 2b and 2e,f). By contrast, when cod

cannibalism was introduced with constant capelin productivity (M3), catch forecasts showed

a strong, consistent 8-yr cycle and substantially higher (–56% to 70%) between-year variation

(Fig. 2c and 2e,f). When combined with varying capelin productivity (M4), however, catch

forecasts showed an 8-yr cycle with slightly weaker amplitudes and highly erratic between-

year variation (ranging from –90% to 312%, Fig. 2d and 2e,f).

*Cod population dynamics and yield stability*

Imposed catch stability control (especially 30% or less) increasingly modified the

amplitude and periodicity of not only catch forecasts but also cod population behavior (Table

1, Fig. 3, and Appendix 2: Fig. S1 and S2). Under the baseline scenario, wavelet analyses

showed modest 32-yr and weak 8-yr cycles in catch, stock size, and SSB (Fig. 3a and

Appendix 2: Fig. S2a–c) with low between-year variability (Fig. 3b and Appendix 2: Fig.

S1), whereas recruitment showed weaker, high-frequency (2- to 4- yr) cycles (Fig. 3a and

Appendix 2: Fig. S2). However, when stability control was introduced, the periodicity of

population and harvest fluctuations sharply shifted to higher (~8-yr) frequency cycles with

greater amplitudes (by, on average, 57% in catch forecast, 47% in stock size, and 160% in

SSB, Fig. 3a and Appendix 2: Fig. S2a–c) and between-year variation (by, on average, 915%

in catch forecast, 423% in stock size, 297% in SSB, and 80% in recruitment, respectively,

Fig. 3b). The stability control was also imposed increasingly more often as the constraint

became stricter (33%, 48%, and 72% of the projection years under 30%, 20%, and 10%

constraints, respectively). With variable capelin productivity, mean between-year variation in

catch forecast and fishing mortality also rose (ranging from –39% to 166% and –24% to 96%,

respectively); catch forecasts fluctuated from 142,000 to 1234,000 t/yr, whereas mean fishing

mortality fluctuated from 0.18 to 1.1/yr. These patterns were contrasted with the latest (2016)

evaluation of the harvest strategies that was based on assumptions of constant capelin

productivity and no temperature dependence, showing that mean between-year variation in

catch declines from 26.9% to 12.4% when a 10% constraint was introduced (ICES 2016).

*Population–fishery asynchrony*

Temporal fluctuations of cod stock size and mean fishing mortality became increasingly out

of phase as the strength of stability control increased (Fig. 4a). Cross-wavelet analyses

showed these unsynchronized signals (with ~4–5 yr delays) converging to shorter (~8-yr)

cycles with greater amplitudes under constraints of 30% or less (Fig. 4b and Appendix 2: Fig.

S3). With being completely out of phase with stock size under a 10% constraint, oscillations

in mean fishing mortality became progressively more amplified, peaking in year ~75 before

target exploitation rate was substantially reduced by the management model (Fig. 4a).

337                               *Stability–yield tradeoffs*

Although average catch forecasts increased (by 10–22%) as the strength of stability control

increased (Table 1), overharvest risk became increasingly greater owning primarily to high

between-year variability (Fig. 5). With constraints of 30% or more, SSB fell below the lower

biological limit threshold, $B_{lim}$, with less than 2% of simulations (Fig. 5). By contrast, with

constraints of 30% or less, SSB fell below $B_{lim}$ more frequently (up to 6.5%) because of

greater amplitudes resulting from unsynchronized cod population–harvest dynamics (Fig. 4).

These risk levels under variable capelin productivity and sea temperature were up to 600x

larger than when previously estimated under constant capelin productivity and no

temperature-dependence (ICES 2016).

347                               DISCUSSION

We demonstrate that the harvest strategy to suppress short-term yield fluctuation can

inadvertently intensify fishing pressure and destabilize population–fishery dynamics of

exploited predators if not accounting for uncertainty in how fishing pattern modifies

population cycles under fluctuating climate and ocean conditions. Our model-assisted

analyses reveal that the cyclicity of stochastic environmental forces (prey abundance and sea

temperature) can interactively shape the periodicity and amplitude of predator population and

demographic (reproduction and mortality) oscillations, which, in turn, act as a dynamic

biological filter (Bjørnstad et al. 2004, Rouyer et al. 2012) to modulate catch fluctuation. A

case study with the Northeast Arctic (NEA) cod shows that inter-cohort density-dependent

processes such as cannibalism (an additional source of juvenile survival variability, Bjørnstad

et al. 1999) can further intensify environmentally driven signals in the population cycle

through early life-history events under strict yield stability control. The undesirable (and

potentially counterintuitive) consequence of this fishing pattern–enhanced amplitude and

shortened periodicity–arises because suppressing between-year variation in adult survival

forces fishing pressure to become progressively unsynchronized with population size owing

partly to pronounced boom-and-bust cycles of exploited species (Anderson et al. 2008, Frank

et al. 2016). Although limiting catch fluctuation (30% or less for NEA cod) theoretically

allows, on average, larger yields because of more frequent, larger peaks in stock abundance

under fluctuating ocean conditions, the unsynchronized population–fishery dynamics would

also increase the probability of overharvest. Incorrect or inappropriate ecological assumptions

of predator population cycles when forecasting fishery catches can, thus, overestimate

population (and thus yield) stability, thereby underestimating (up to a few orders of

magnitude) depletion risk and likely delaying management actions.

*Ecosystem control of exploited population cycles*

Myriad ecosystem processes regulating survival of exploited predators interactively

influence how given harvest strategies perform in practice (Okamoto et al. 2016). Harvest

strategies to regulate fishing pattern unsynchronized with ecosystem carrying capacity, for

example, can amplify its variability instead (Link 2017). Our work suggests that suppressing

short-term yield fluctuation can accelerate the population cycle of fished species and, in turn,

inadvertently elevate variability not only in catch but also in population behavior in

stochastically fluctuating environments. In relatively species-poor, high-latitude systems such

as the Barents Sea, changes in ecosystem productivity likely propagate further through the

food web (less buffering) and may amplify or attenuate year-to-year fluctuation in life history

events and population size of fished predators (Bellingeri and Vincenzi 2013). Year-class

fluctuation of many fished populations is governed by temporal fluctuation in large-scale

climatic and oceanic processes (Cushing 1990) that are mediated through food-web

interactions (Lilly et al. 2008). By explicitly accounting for one of the key ecological

processes that regulate cod productivity–a density-dependent interplay of cannibalism and

food supply, our analyses reveal that the NEA cod population fluctuates with internally and

externally regulated, low-frequency (multidecadal) cycles, roughly reflecting capelin

productivity cycles when fishing pattern is not forced to suppress yield fluctuation.

Over the past half century, the Barents Sea has shown large fluctuations in productivity

(Dalpadado et al. 2012), during which capelin has collapsed three times (Gjøsæter et al.

2009). The capelin collapses subsequently intensified cod cannibalism, weakening cod

recruitment strength (Yaragina et al. 2009, Yaragina et al. 2018) and ultimately depleting the

cod stock in the late 1980s. Because Barents Sea capelin also supports (in addition to its

fishery) other apex predators as a food source, its declines may have triggered a series of

trophically mediated processes that exacerbated the cod population depletion, including

increased consumption of the young by marine mammals such as harp seals (Bogstad et al.

2015). Declined adult cod survival by overharvest can also release mesopredators from

predation (Hjermann et al. 2007), promoting resource competition with and predation on

young cod–weakening the 'cultivation' effect (Walters and Kitchell 2001). These complex,

ecological dynamics that regulate year-class strength of fished populations can have both

stabilizing and destabilizing effects under variable fishing pressure and ecosystem

productivity (Anderson et al. 2008) and need to be accounted for in designing and evaluating

management measures.

*Managing harvest stability of predators with transient cycles*

Imposing yield stability may inadvertently amplify variability of exploited species by

modifying natural population cycles owing to various sources of uncertainty often overlooked

in assessments (Botsford et al. 1997). Nearly three-quarters of assessed fish stocks worldwide

(200+) have shown irregular temporal shifts in productivity regime (Vert-pre et al. 2013);

imposing stability control without accounting for ecological uncertainty such as fluctuation in

forage species productivity may increase overharvest and depletion risks of fished predators.

In the case of NEA cod, analyses indicate that an attempt to dampen the amplitude of yield

fluctuation instead amplifies its sensitivity to environmental forces with stronger signals–sea

temperature in this study–and shorten population cycle, thereby forcing the exploited predator

to become increasingly unsynchronized with (and, in effect, inadvertently intensifying)

fishing pressure. Further, the periodicity of population and catch fluctuations progressively

shifts toward higher frequencies that reflect life-history traits such as the dominant age

groups of spawners (~6- to 8- year-olds for NEA cod, Appendix 1: Fig. 1 and Rouyer et al.

2011) as fishing pressure becomes intensified (through stricter stability control), a pattern

predicted by theory (Bjørnstad et al. 2004, Worden et al. 2010). Because industrial capture

fisheries often selectively remove larger and older individuals, biased reduction in adult

survival by fishing can further modify demography and population cycle by truncating age

structure (juvenescence, Lindegren et al. 2013, Okamoto et al. 2016). As a result, intensely

fished populations become progressively more recruitment-dependent, eroding their

resilience to environmental and demographic stochasticity (Minto et al. 2008, Hidalgo et al.

2011), confounding efforts to sustain human exploitation while conserving biological

diversity in the ecosystems.

Although large short-term variation in catch is often disfavored by harvesters and seafood

processors, oscillatory patterns in population cycle of exploited predators may provide

valuable information about the population status such as its resilience to changes in external

forcing (Anderies 2015). Magnified variance may signal impending internal structural shifts

in nature including fished populations (Brock and Carpenter 2010). The analysis of the

extensive database on global commercial marine fisheries compiled by the United Nations

Food and Agriculture Organization (FAO) suggests that imposed catch stability can hide

early warning signals that may help prevent population collapses–a potential long-term

consequence of suppressing short-term yield fluctuation (Mullon et al. 2005, Carpenter et al.

2015). Stability in exploitation of living marine resources may, thus, need to come from

promoting population resilience (restoration of age and size structure, for example) rather

than imposing apparent stability (Anderson et al. 2008, Carpenter et al. 2015).

   Our case study with the exploited Arctic piscivore–planktivore complex illustrates that the

harvest strategy to suppress yield fluctuation–a widely implemented policy tool to achieve

socioeconomic goals–can bring about counterintuitive, adverse outcomes if not accounting

for fluctuating ocean conditions. This harvest policy can inadvertently destabilize

population–fishery dynamics of exploited species as it forces fishing pattern to become

unsynchronized with the cyclicity of population size and demography under environmental

stochasticity. Because our model did not account for other sources of uncertainty such as

observation errors and model structural errors (for example, no predation feedback from adult

cod to capelin), year-to-year population and harvest fluctuations in simulations are likely

conservative estimates. Nevertheless, the findings highlight broader implications of

ecological uncertainty when designing and analyzing harvest policies for ecologically

connected populations. Evaluating a policy without accounting for variability in key

ecosystem processes–common practice in assessments of exploited species–may

fundamentally reshape population structure and behavior, thereby underestimating the risk of

population decline and possibly extinction (Gårdmark et al. 2013). With expected changes

(both mean and variance) in climatic fluctuation and, in turn, ocean productivity (Wassmann

455 et al. 2011, Cohen et al. 2014, Lewis et al. 2020), adapting ecosystem approaches tailored to

456 specific contexts (such as use of models of intermediate complexity for ecosystem, Plagányi

457 et al. 2014), as demonstrated here, will safeguard against such risk and help sustainably

458 manage marine living resources while securing socioeconomic stability.

ACKNOWLEDGEMENTS

460 We thank anonymous reviewers for comments on earlier versions of the manuscript. Some

461 figures use images from the IAN Symbols, courtesy of the Integration and Application

462 Network, University of Maryland Center for Environmental Science

463 (ian.umces.edu/symbols/). This project was partially funded by Institute of Marine

464 Research's REDUS (Reduced Uncertainty in Stock Assessments) project.

LITERTURE CITED

Anderies, J. M. 2015. Managing variance: key policy challenges for the Anthropocene.

Proceedings of the National Academy of Sciences **112**:14402-14403.

Anderson, C. N., C.-h. Hsieh, S. A. Sandin, R. Hewitt, A. Hollowed, J. Beddington, R. M.

May, and G. Sugihara. 2008. Why fishing magnifies fluctuations in fish abundance. Nature

**452**:835.

Bellingeri, M., and S. Vincenzi. 2013. Robustness of empirical food webs with varying

consumer's sensitivities to loss of resources. Journal of Theoretical Biology **333**:18-26.

Bjørnstad, O. N., J.-M. Fromentin, N. C. Stenseth, and J. Gjøsæter. 1999. Cycles and trends

in cod populations. Proceedings of the National Academy of Sciences **96**:5066-5071.

Bjørnstad, O. N., and B. T. Grenfell. 2001. Noisy clockwork: time series analysis of

population fluctuations in animals. Science **293**:638-643.

Bjørnstad, O. N., R. M. Nisbet, and J. M. Fromentin. 2004. Trends and cohort resonant

effects in age structured populations. Journal of Animal Ecology **73**:1157-1167.

Bogstad, B., H. Gjøsæter, T. Haug, and U. Lindstrøm. 2015. A review of the battle for food

in the Barents Sea: cod vs. marine mammals. Frontiers in Ecology and Evolution **3**:29.

Botsford, L. W., J. C. Castilla, and C. H. Peterson. 1997. The management of fisheries and

marine ecosystems. Science **277**:509-515.

Brock, W. A., and S. R. Carpenter. 2010. Interacting regime shifts in ecosystems: implication

for early warnings. Ecological Monographs **80**:353-367.

Caddy, J., and J. Seijo. 2005. This is more difficult than we thought! The responsibility of

scientists, managers and stakeholders to mitigate the unsustainability of marine fisheries.

Philosophical Transactions of the Royal Society of London B: Biological Sciences **360**:59-

75.

Carpenter, S. R., W. A. Brock, C. Folke, E. H. Van Nes, and M. Scheffer. 2015. Allowing

variance may enlarge the safe operating space for exploited ecosystems. Proceedings of the

National Academy of Sciences **112**:14384-14389.

Carscadden, J., K. Frank, and W. Leggett. 2001. Ecosystem changes and the effects on

capelin (*Mallotus villosus*), a major forage species. Canadian Journal of Fisheries and

Aquatic Sciences **58**:73-85.

Cazelles, B., K. Cazelles, and M. Chavez. 2014. Wavelet analysis in ecology and

epidemiology: impact of statistical tests. Journal of the Royal Society Interface

**11**:20130585.

Ciannelli, L., and K. M. Bailey. 2005. Landscape dynamics and resulting species interactions:

the cod-capelin system in the southeastern Bering Sea. Marine Ecology Progress Series

**291**:227-236.

Cohen, J., J. A. Screen, J. C. Furtado, M. Barlow, D. Whittleston, D. Coumou, J. Francis, K.

Dethloff, D. Entekhabi, and J. Overland. 2014. Recent Arctic amplification and extreme

mid-latitude weather. Nature Geoscience **7**:627-637.

Cury, P. M., Y. J. Shin, B. Planque, J. M. Durant, J. M. Fromentin, S. Kramer-Schadt, N. C.

Stenseth, M. Travers, and V. Grimm. 2008. Ecosystem oceanography for global change in

fisheries. Trends in Ecology & Evolution **23**:338-346.

Cushing, D. 1990. Plankton production and year-class strength in fish populations: an update

of the match/mismatch hypothesis. Pages 249-293 Advances in Marine Biology. Elsevier.

Dalpadado, P., R. B. Ingvaldsen, L. C. Stige, B. Bogstad, T. Knutsen, G. Ottersen, and B.

Ellertsen. 2012. Climate effects on Barents Sea ecosystem dynamics. ICES Journal of

Marine Science **69**:1303-1316.

Dolgov, A. 2002. The role of capelin (*Mallotus villosus*) in the foodweb of the Barents Sea.

ICES Journal of Marine Science **59**:1034-1045.

Edwards, M., and A. J. Richardson. 2004. Impact of climate change on marine pelagic

phenology and trophic mismatch. Nature **430**:881.

Essington, T. E., J. N. Sanchirico, and M. L. Baskett. 2018. Economic value of ecological

information in ecosystem-based natural resource management depends on exploitation

history. Proceedings of the National Academy of Sciences:201716858.

Filin, A. A. 2009. Changes in cod management Sstrategy that have resulted from warming

scenarios for The Barents Sea.*in* Theme Session on Advances in marine ecosystem

research: what we have learned from GLOBEC and what we can carry forward in future

climate related programs. International Council for the Exploration of the Sea, ICES CM

2009/C:05.

Frank, K. T., B. Petrie, W. C. Leggett, and D. G. Boyce. 2016. Large scale, synchronous

variability of marine fish populations driven by commercial exploitation. Proceedings of

the National Academy of Sciences **113**:8248-8253.

Fryxell, J. M., C. Packer, K. McCann, E. J. Solberg, and B.-E. Sæther. 2010. Resource

management cycles and the sustainability of harvested wildlife populations. Science

**328**:903-906.

Garcia, S. M., and A. A. Rosenberg. 2010. Food security and marine capture fisheries:

characteristics, trends, drivers and future perspectives. Philosophical Transactions of the

Royal Society B: Biological Sciences **365**:2869-2880.

Gårdmark, A., M. Lindegren, S. Neuenfeldt, T. Blenckner, O. Heikinheimo, B. Müller-

Karulis, S. Niiranen, M. T. Tomczak, E. Aro, and A. Wikström. 2013. Biological ensemble

modeling to evaluate potential futures of living marine resources. Ecological Applications

**23**:742-754.

Gjøsæter, H. 1998. The population biology and exploitation of capelin (*Mallotus villosus*) in

the Barents Sea. Sarsia **83**:453-496.

Gjøsæter, H., B. Bogstad, and S. Tjelmeland. 2009. Ecosystem effects of the three capelin

stock collapses in the Barents Sea. Marine Biology Research **5**:40-53.

Gjøsæter, H., S. Tjelmeland, and B. Bogstad. 2012. Ecosystem-based management of fish

species in the Barents Sea. Pages 333-352 *in* G. H. Kruse, H. I. Browman, K. L. Cochrane,

D. Evans, G. S. Jamieson, P. A. Livingston, D. Woodby, and C. I. Zhang, editors. Global

Progress in Ecosystem-Based Fisheries Management. Alaska Sea Grant, University of

Alaska Fairbanks, .

Hall, S. J., and B. Mainprize. 2004. Towards ecosystem based fisheries management. Fish

and Fisheries **5**:1-20.

Hicks, C. C., P. J. Cohen, N. A. Graham, K. L. Nash, E. H. Allison, C. D'Lima, D. J. Mills,

M. Roscher, S. H. Thilsted, and A. L. Thorne-Lyman. 2019. Harnessing global fisheries to

tackle micronutrient deficiencies. Nature **574**:95-98.

Hidalgo, M., T. Rouyer, J. C. Molinero, E. Massutí, J. Moranta, B. Guijarro, and N. C.

Stenseth. 2011. Synergistic effects of fishing-induced demographic changes and climate

variation on fish population dynamics. Marine Ecology Progress Series **426**:1-12.

Hjermann, D. Ø., B. Bogstad, A. M. Eikeset, G. Ottersen, H. Gjøsæter, and N. C. Stenseth.

2007. Food web dynamics affect Northeast Arctic cod recruitment. Proceedings of the

Royal Society of London B: Biological Sciences **274**:661-669.

Hjort, J. 1914. Fluctuations in the great fisheries of northern Europe viewed in the light of

biological research. ICES.

Holt, R. E., B. Bogstad, J. M. Durant, A. V. Dolgov, and G. Ottersen. 2019. Barents Sea cod

(*Gadus morhua*) diet composition: long-term interannual, seasonal, and ontogenetic

patterns. ICES Journal of Marine Science **76**:1641-1652.

Howell, D., and A. A. Filin. 2014. Modelling the likely impacts of climate-driven changes in

cod-capelin overlap in the Barents Sea. ICES Journal of Marine Science **71**:72-80.

Howell, D., A. A. Filin, B. Bogstad, and J. E. Stiansen. 2013. Unquantifiable uncertainty in

projecting stock response to climate change: example from North East Arctic cod. Marine

Biology Research **9**:920-931.

ICES. 2016. Report of the Second Workshop on Management Plan Evaluation on Northeast

Arctic Cod and Haddock and Barents Sea Capelin (WKNEAMP-2), 25–28 January 2016,

Kirkenes, Norway. ICES CM 2016/ACOM:47. Page 104.

ICES. 2017. Report of the Arctic Fisheries Working Group (AFWG), 19–25 April 2017,

Copenhagen, Denmark. ICES CM 2017/ACOM:06. 493 pp.

ICES. 2019. Arctic Fisheries Working Group (AFWG). ICES Scientific Reports. 1:30. 934

pp. http://doi.org/10.17895/ices.pub.5292

Ingvaldsen, R., H. Loeng, G. Ottersen, and B. Ådlandsvik. 2003. Climate variability in the

Barents Sea during the 20th century with focus on the 1990s. Pages 160-168 *in* ICES

Marine Science Symposia.

Johannesen, E., U. Lindstrøm, K. Michalsen, M. Skern-Mauritzen, P. Fauchald, B. Bogstad,

and A. Dolgov. 2012. Feeding in a heterogeneous environment: spatial dynamics in

summer foraging Barents Sea cod. Marine Ecology Progress Series **458**:181-197.

Kjesbu, O. S., B. Bogstad, J. A. Devine, H. Gjøsæter, D. Howell, R. B. Ingvaldsen, R. D.

Nash, and J. E. Skjæraasen. 2014. Synergies between climate and management for Atlantic

cod fisheries at high latitudes. Proceedings of the National Academy of Sciences **111**:3478-

3483.

Leggett, W., and E. Deblois. 1994. Recruitment in marine fishes: is it regulated by starvation

and predation in the egg and larval stages? Netherlands Journal of Sea Research **32**:119-

134.

Lewis K, van Dijken G, Arrigo K. 2020. Changes in phytoplankton concentration now drive

increased Arctic Ocean primary production. Science 369:198-202.

Lilly, G. R., K. Wieland, B. J. Rothschild, S. Sundby, K. Drinkwater, K. Brander, G.

Ottersen, J. E. Carscadden, G. B. Stenson, and G. A. Chouinard. 2008. Decline and

recovery of Atlantic cod (*Gadus morhua*) stocks throughout the North Atlantic. Pages 39–

66 *in* G. H. Kruse, K. F. Drinkwater, J. N. Ianelli, J. S. Link, D. L. Stram, V. Wespestad,

and D. Woodby, editors. Resiliency of gadic stocks to fishing and climate change. Alaska

Sea Grant, University of Alaska Fairbanks.

Lindegren, M., D. M. Checkley, T. Rouyer, A. D. MacCall, and N. C. Stenseth. 2013.

Climate, fishing, and fluctuations of sardine and anchovy in the California Current.

Proceedings of the National Academy of Sciences **110**:13672-13677.

Link, J. S. 2017. System-level optimal yield: increased value, less risk, improved stability,

and better fisheries. Canadian Journal of Fisheries and Aquatic Sciences **75**:1-16.

Link, J. S., B. Bogstad, H. Sparholt, and G. R. Lilly. 2009. Trophic role of Atlantic cod in the

ecosystem. Fish and Fisheries **10**:58-87.

Minto, C., R. A. Myers, and W. Blanchard. 2008. Survival variability and population density

in fish populations. Nature **452**:344.

Minto, C., and B. Worm. 2012. Interactions between small pelagic fish and young cod across

the North Atlantic. Ecology **93**:2139-2154.

Mullon, C., P. Fréon, and P. Cury. 2005. The dynamics of collapse in world fisheries. Fish

and Fisheries **6**:111-120.

Nielsen, J. R., E. Thunberg, D. S. Holland, J. O. Schmidt, E. A. Fulton, F. Bastardie, A. E.

Punt, I. Allen, H. Bartelings, and M. Bertignac. 2018. Integrated ecological–economic

fisheries models—Evaluation, review and challenges for implementation. Fish and

Fisheries **19**:1-29.

Okamoto, D. K., R. J. Schmitt, and S. J. Holbrook. 2016. Stochastic density effects on adult

fish survival and implications for population fluctuations. Ecology Letters **19**:153-162.

Ottersen, G., B. Planque, A. Belgrano, E. Post, P. C. Reid, and N. C. Stenseth. 2001.

Ecological effects of the North Atlantic oscillation. Oecologia **128**:1-14.

Ottersen, G., and N. C. Stenseth. 2001. Atlantic climate governs oceanographic and

ecological variability in the Barents Sea. Limnology and Oceanography **46**:1774-1780.

Plagányi, É. E., A. E. Punt, R. Hillary, E. B. Morello, O. Thébaud, T. Hutton, R. D. Pillans, J.

T. Thorson, E. A. Fulton, and A. D. Smith. 2014. Multispecies fisheries management and

conservation: tactical applications using models of intermediate complexity. Fish and

Fisheries **15**:1-22.

Ricker, W. 1958. Maximum sustained yields from fluctuating environments and mixed

stocks. Journal of the Fisheries Board of Canada **15**:991-1006.

Rindorf, A., M. Cardinale, S. Shephard, J. A. De Oliveira, E. Hjorleifsson, A. Kempf, A.

Luzenczyk, C. Millar, D. C. Miller, and C. L. Needle10. 2016. Fishing for MSY: can

"pretty good yield" ranges be used without impairing recruitment. ICES Journal of Marine

Science (topic issue on targets and limits in long-term fisheries management).

Roesch, A., and H. Schmidbauer. 2018. WaveletComp: computational wavelet analysis. R

package version 1.1. http://CRAN.R-project.org/package=WaveletComp.

Rose, G. A. 2005. Capelin (*Mallotus villosus*) distribution and climate: a sea "canary" for

marine ecosystem change. ICES Journal of Marine Science **62**:1524-1530.

Rouyer, T., A. Sadykov, J. Ohlberger, and N. C. Stenseth. 2012. Does increasing mortality

change the response of fish populations to environmental fluctuations? Ecology Letters

**15**:658-665.

Shelton, A. O., and M. Mangel. 2011. Fluctuations of fish populations and the magnifying

effects of fishing. Proceedings of the National Academy of Sciences **108**:7075-7080.

Stige, L. C., G. Ottersen, P. Dalpadado, K.-S. Chan, D. Ø. Hjermann, D. L. Lajus, N. A.

Yaragina, and N. C. Stenseth. 2010. Direct and indirect climate forcing in a multi-species

marine system. Proceedings of the Royal Society B: Biological Sciences **277**:3411-3420.

Vert-pre, K. A., R. O. Amoroso, O. P. Jensen, and R. Hilborn. 2013. Frequency and intensity

of productivity regime shifts in marine fish stocks. Proceedings of the National Academy of

Sciences **110**:1779-1784.

Walters, C., and J. F. Kitchell. 2001. Cultivation/depensation effects on juvenile survival and

recruitment: implications for the theory of fishing. Canadian Journal of Fisheries and

Aquatic Sciences **58**:39-50.

Ware, D. M., and R. E. Thomson. 2005. Bottom-up ecosystem trophic dynamics determine

fish production in the Northeast Pacific. Science **308**:1280-1284.

Wassmann, P., C. M. Duarte, S. Agusti, and M. K. Sejr. 2011. Footprints of climate change in

the Arctic marine ecosystem. Global Change Biology **17**:1235-1249.

Worden, L., L. W. Botsford, A. Hastings, and M. D. Holland. 2010. Frequency responses of

age-structured populations: Pacific salmon as an example. Theoretical Population Biology

**78**:239-249.

Yaragina, N. A., B. Bogstad, and Y. A. Kovalev. 2009. Variability in cannibalism in

Northeast Arctic cod (*Gadus morhua*) during the period 1947–2006. Marine Biology

Research **5**:75-85.

Yaragina, N. A., Y. Kovalev, and A. Chetyrkin. 2018. Extrapolating predation mortalities

back in time: an example from North-east Arctic cod cannibalism. Marine Biology

Research **14**:203-216.

TABLES

Table 1. Summary of Northeast Arctic (NEA) cod population status and management target metrics computed from stochastic, multispecies model (STOCOBAR) simulations under scenarios of varying levels of limiting yield fluctuation (stability constraints). Stock size indicates total harvestable (3-year-olds and older) biomass; SSB indicates spawning stock biomass; recruits indicates abundance of 3-year-old cod.

| population status | | stability constraints | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | baseline | 50% | 40% | 30% | 25% | 20% | 15% | 10% | 5% |
| stock size (million t) | 2.27 | 2.25 | 2.30 | 2.25 | 2.37 | 2.56 | 2.78 | 2.99 | 3.17 |
| SSB (million t) | 1.08 | 1.07 | 1.09 | 1.06 | 1.13 | 1.24 | 1.38 | 1.49 | 1.63 |
| catch (1000 t) | 684.1 | 678.5 | 691.1 | 667.7 | 701.5 | 752.7 | 805.0 | 836.2 | 800.5 |
| recruits (millions) | 700.4 | 690.2 | 704.7 | 680.4 | 707.8 | 733.9 | 743.5 | 764.1 | 755.2 |

FIGURE LEGENDS

Figure 1. Study system and food-web model. (a) Northeast Arctic (NEA) cod distribution in the Barents Sea. (b) Example of modeled sea temperature (˚C) time series based on observed annual mean temperatures in the Kola section. (c) Example of modeled capelin biomass (1000 t) time series. (d) Standardized sea temperature anomalies computed from modeled sea temperatures. (e) Standardized capelin anomalies computed from modeled capelin biomass. (f) Schematic diagram of the NEA cod–capelin stochastic, multispecies model (STOCOBAR) structure. In d and e, standardized anomalies were computed by subtracting modeled annual temperatures (b) or capelin biomass (c) from the mean and dividing by the standard deviation. Artwork: Courtesy of the Integration and Application Network, University of Maryland Center for Environmental Science (ian.umces.edu/symbols/).

Figure 2. Ecological uncertainty in Northeast Arctic (NEA) cod catch fluctuation projected with a stochastic, multispecies model (STOCOBAR) under scenarios of a ±10% stability constraint and variable sea temperature with varying model configuration of cod cannibalism–capelin productivity. (a–d) Wavelet power spectra [local (times series, left) and global (time-averaged, right)] of NEA cod catch forecasts with (a) constant capelin biomass (1000 t) without cannibalism, (b) varying capelin biomass without cannibalism, (c) constant capelin biomass with cannibalism, and (d) varying capelin biomass with cannibalism. (e) Time series (years 21–100, left) and mean year-to-year variance (%) of annual NEA cod catch (right). In wavelet spectra, red areas indicate higher power (intensity of periodicities), blue areas indicate lower power, and white areas indicate regions influenced by edge effects (outside the "cone of influence") and inferences cannot be made (Cazelles et al. 2014). In a–d, y-axis is in the logarithms to the base 2. In e (right), error bars indicate 95% confidence intervals (among-year variation over years 21–100). Artwork: Courtesy of the Integration and

Application Network, University of Maryland Center for Environmental Science (ian.umces.edu/symbols/).

Figure 3. Fluctuations in Northeast Arctic (NEA) cod population behavior and catch projected with a stochastic, multispecies model (STOCOBAR) under scenarios of 0% (baseline), ±30%, ±20%, and ±10% stability constraints. (a) Global (time-averaged) wavelet power spectra of NEA cod catch, mean (5- to 10- year-olds) fishing mortality, harvestable (3-year-olds and older) biomass (stock size), spawner abundance (SSB), and recruits (abundance of 3-year-olds). (b) Mean year-to-year variance (%) in catch, mean fishing mortality, stock size, SSB, and recruits. In a, y-axis is in the logarithms to the base 2. In b, error bars indicate 95% confidence intervals (among-year variation over years 21–100).

Figure 4. Synchrony of Northeast Arctic (NEA) cod population abundance and fishing pressure projected with a stochastic, multispecies model (STOCOBAR) under scenarios of varying strength of stability constraints. (a) Time series (years 21–100) of NEA cod harvestable (3-year-olds and older) biomass (stock size) and mean (5- to 10- year-olds) fishing mortality under 0% (baseline), ±30%, ±20%, and ±10% constraints. (b) Cross-wavelet power spectra [local (times series, left) and global (time-averaged, right)] between NEA cod stock size and mean fishing mortality under ±30%, ±20%, and ±10% constraints. In b, red areas indicate higher power (intensity of periodicities), blue areas indicate lower power, and white areas indicate regions influenced by edge effects (outside the "cone of influence") and inferences cannot be made (Cazelles et al. 2014). In b, y-axis is in the logarithms to the base 2; arrows pointing right, left, down, and up indicate the two series being in-phase, the two series being anti-phase, stock size being leading, and mean fishing mortality being leading (respectively).

Figure 5. Relationship between stability constraints (±CV%) and risk for Northeast Arctic cod computed from stochastic, multispecies model (STOCOBAR) simulations. Risk is based

on the probability of spawner abundance (SSB) falling below the lower biological limit threshold ($B_{lim}$ = 220,000 t, ICES 2016). A gray horizontal dashed line indicates the risk criterion for reproduction of the population becoming impaired (5%).

a

b