BPIFB1 loss alters airway mucus properties and diminishes mucociliary clearance

Lauren J. Donoghue[1,2], Matthew R. Markovetz[3], Cameron B. Morrison[3], Kathryn M. McFadden[1], Troy D. Rogers[3], Brian Button[3,4], Camille Ehre[3,5], David B. Hill[3,6], Barbara R. Grubb[3], and Samir N. P. Kelada[1,3]*

[1]Department of Genetics, University of North Carolina at Chapel Hill
[2]Curriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill
[3]Marsico Lung Institute, University of North Carolina at Chapel Hill
[4]Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill
[5]Division of Pediatric Pulmonology, University of North Carolina at Chapel Hill
[6]Department of Physics and Astronomy, University of North Carolina at Chapel Hill

*Corresponding author
Samir Kelada, PhD MPH
University of North Carolina
Department of Genetics
120 Mason Farm Road
Chapel Hill, NC 27599
919-962-2148
samir_kelada@med.unc.edu

Conflict of interest statement: "The authors have declared that no conflict of interest exists."

ABSTRACT

Airway mucociliary clearance (MCC) is required for host defense and often diminished in

chronic lung diseases. Effective clearance depends upon coordinated actions of the airway

epithelium and a mobile mucus layer. Dysregulation of the primary secreted airway mucin

proteins, MUC5B and MUC5AC, is associated with a reduction in the rate of MCC; however,

how other secreted proteins impact the integrity of the mucus layer and MCC remains unclear.

We previously identified the gene *Bpifb1/Lplunc1* as a regulator of airway MUC5B levels using

genetic approaches. Here, we show that BPIFB1 is required for normal mucociliary clearance *in*

vivo using *Bpifb1* knockout (KO) mice. Reduced MCC in *Bpifb1* KO mice occurred in the absence

of defects in sodium or chloride ion transport or reduced ciliary beat frequency. BPIFB1 loss

resulted in airway mucus flakes with significantly increased complex viscosity, a key biophysical

property of mucus known to impact MCC. Finally, we detected colocalization of BPIFB1 and

MUC5B in secretory granules in mice and in the protein mesh of secreted mucus in human

airway cultures. Collectively, our findings demonstrate that BPIFB1 is an important component

of the mucociliary apparatus in mice and a key component of the mucus protein network.

INTRODUCTION

The airway mucus layer plays a critical role in airway homeostasis and host defense in part by

facilitating mucociliary clearance (MCC) of inhaled particles and pathogens. Effective MCC

requires a properly hydrated airway surface, an optimal balance of components in secreted

mucus (i.e., proteins, salt), and coordinated beating of cilia. The importance of the MCC

apparatus is highlighted by the fact that mutations in genes essential to ciliary function or ion

transport cause primary ciliary dyskinesia (PCD) and cystic fibrosis (CF), respectively, in which

respiratory infections are prominent and severe. In the context of more common obstructive

airway disease such as asthma and chronic obstructive pulmonary disease (COPD), changes in

MCC have been associated with variation in the abundance of the secreted mucin proteins

MUC5B and MUC5AC, which are key contributors to the gel-like nature of healthy airway

mucus[1]. The concentration, organization, and post-translational modifications of MUC5B and

MUC5AC also have been shown to alter biophysical properties of mucus that are required for

effective MCC[2–6]. Furthermore, the ability of mucins to form a proper gel-structure is

dependent upon interactions with other secreted proteins, referred to as the "mucin

interactome"[7], although the specifics of these dependencies remain unclear.

In contrast to MUC5AC, which is typically only observed at high levels in disease states, MUC5B

is constitutively expressed in the airways of healthy individuals where it has a role in regulating

homeostasis through airway MCC and host defense[8,9]. We previously identified the gene *Bpifb1*

(also known as *Lplunc1*) as a regulator of airway MUC5B protein levels using a quantitative

genetics approach in mice and showed that loss of BPIFB1 in a knockout (KO) mouse model led

to increased levels of MUC5B protein in the airways[10]. Given that MUC5B levels have been

shown to affect MCC[9,11], we sought to determine if *Bpifb1* also had a role in regulating MCC.

We observed a substantial loss of MCC in the lower airways of *Bpifb1* KO mice. To identify

which aspects of the MCC apparatus contributed to this defect, we assessed epithelial sodium

and chloride ion transport, ciliary function, and biophysical properties of mucus. Our results

show that alterations in the mucus layer including increased mucus viscoelasticity were

associated with decreased MCC in the absence of BPIFB1 and that BPIFB1 is a component of the

mucus protein network.

RESULTS AND DISCUSSION

In order to determine if *Bpifb1* has a role in MCC, we assessed MCC in tracheas of naïve *Bpifb1*

KO and WT mice. We quantified MCC by measuring the percentage of instilled beads that were

cleared by the trachea over a fixed period of time. *Bpifb1* KO mice exhibited significantly

decreased clearance, with some mice exhibiting no tracheal bead clearance whatsoever (Figure

1A). These results indicated clearly that BPIFB1 is required for MCC.

To understand the mechanism by which BPIFB1 loss leads to defective MCC, we systematically

tested the expression and function of key components of the MCC apparatus. First, we

examined the expression of key ion transporters in the airway epithelium given their

importance in regulating airway hydration and MCC[1]. RNA sequencing of lung tissue from

Bpifb1 KO and WT mice did not reveal significant differences in expression of well-characterized

sodium and chloride ion transporter genes including *Cftr*, *Scnn1b*, and *Tmem16a* (Figure S1).

However, given that a lack of difference in ion transporter gene expression does not rule out a

difference in functionality, we also measured bioelectric capacities in *Bpifb1* KO and WT

tracheas. We found no differences in short-circuit current (I_{sc}) at baseline or following

stimulation with compounds targeting the major ion channels in the airway epithelium (i.e.

CFTR, ENaC, CaCC), (Figure 1B). This indicated that the loss of MCC in *Bpifb1* KO mice was likely

not driven by altered ion transport. Additionally, as ciliary beating is essential for MCC, we

directly measured ciliary beat frequency (CBF) in intact tracheas *in situ*. In comparison to WT

mice, *Bpifb1* KO mice had a slightly elevated CBF, albeit not statistically significant at $P < 0.05$

(Fig 1C). These data indicated that reduced MCC was not associated with reduced ciliary

beating.

Next, to determine if *Bpifb1* loss influenced the expression of other genes or biological

processes that could underlie the defect in MCC, we examined global gene expression in lung

tissue from *Bpifb1* KO and WT mice. Excluding *Bpifb1*, only 4 protein coding genes and 2 long

non-coding RNAs were differentially expressed by genotype (|fold change| $\geq$ 2 and adjusted *P*-

value $\leq$ 0.05, Figure S1). Of these genes, none have been previously annotated for a direct role

in MCC or related pathways. Furthermore, genes related to mucin production, including *Muc5b*,

Muc5ac, *Spdef*, and *Foxa3,* were not differentially expressed. These gene expression results

suggested the reduction in MCC in *Bpifb1* KO mice was not mediated by differential expression

of key MCC or mucus production genes.

Given the lack of differences in hallmark causes of reduced MCC, we investigated if there were

differences in the airway mucus *per se* that could be contributing to the observed loss of MCC.

Using microbead rheology, we measured the complex viscosity (η*), a measure of both viscosity

and elasticity, of mucus flakes (i.e. the insoluble portions of airway mucus) in bronchoalveolar

lavage (BAL) fluid from *Bpifb1* KO vs WT mice. For these measurements, mice were intranasally

exposed to house dust mite allergen for three weeks to increase the levels of mucus in BAL, and

there were no differences in the degree of eosinophilic inflammation by genotype

(Supplementary Figure S2). We found that the complex viscosity of mucus flakes was

significantly increased in *Bpifb1* KO mice compared to WT mice (Figure 2A, B). By imaging of

whole BAL, we determined this increase was independent of the size and number of mucus

flakes, and there were no differences in these metrics by genotype (Supplementary Figure S3A-

E). Consistent with these findings and our previous report of increased MUC5B in BAL from

naïve *Bpifb1* KO mice[10], we found that MUC5B levels were elevated by genotype in the

supernatant fraction of centrifuged BAL from naïve mice, but not in the pelleted (i.e. insoluble)

fraction (Figure 2C). Thus, loss of BPIFB1 results in altered biophysical properties of mucus

flakes and increased levels of soluble MUC5B.

To gain insight into whether BPIFB1 colocalizes with mucin protein in the airways and, if so,

whether this association occurs before mucin granule release, we assessed the spatial

relationship between BPIFB1 and mucin protein using fluorescent immunostaining. By

colocalization analysis of labeled BPIFB1 and MUC5B protein in large airways of mice challenged

with allergen, we determined that the vast majority of MUC5B positive cells were also positive

for BPIFB1 (Figure 3A, B), indicating regulated cellular co-expression of MUC5B and BPIFB1.

Colocalization was also present in the airways of mice not challenged with allergen

(Supplementary Figure S4). Furthermore, MUC5B positive granules were also positive for

BPIFB1 and luminal mucus contained both MUC5B and BPIFB1 (Figure 3C), suggesting that

packaging and secretion of these proteins is likely coupled.

Finally, we performed immunogold-labeled scanning electron microscopy for BPIFB1 and

MUC5B in human bronchial epithelial cells (HBECs) cultured at air-liquid-interface to determine

if BPIFB1 closely associated with extracellular mucus. We detected BPIFB1 exclusively in mucus

strands coating the cellular surface (Figure 4). This finding confirmed that BPIFB1 is a

component of the mucus protein network produced by human airway cells.

Significantly altered levels of either *BPIFB1* gene or protein expression have been observed in

CF, COPD, asthma, and IPF[12–15], indicating a strong connection between *BPIFB1* and muco-

obstructive diseases. After identifying *Bpifb1* as a regulator of airway mucus levels through an

unbiased genome-wide search in the mouse[10], we sought to interrogate the specific function of

BPIFB1 in the airways. We found that the loss of BPIFB1 reduced MCC dramatically in naïve

Bpifb1 KO mice, demonstrating the critical role of this gene in regulating MCC. Thus, in addition

to the strong connection between *BPIFB1* gene/protein expression and muco-obstructive

diseases, our findings indicate BPIFB1 may have an important role in airway host defense by

regulating MCC. Studies in humans have shown that there is significant interindividual variation

in baseline MCC, and analysis of MCC in monozygotic and dizygotic twins has implicated genetic

variation as a modifier of MCC[16]. Given the identification of common genetic variation

associated with the levels of *BPIFB1* gene expression[17], genetic variation in *BPIFB1* regulation

could be one such factor regulating baseline MCC in humans. Furthermore, BPIFB1 was one of

the top 20 proteins detected in endotracheal tube mucus from individuals without a history of

lung disease or smoking, highlighting its abundance in non-disease states[18].

Due to the lack of differences in basal or stimulated transport of sodium and chloride ions,

ciliary beating, and MCC-related gene expression between *Bpfib1* KO and WT mice, our data

point toward changes to the airway mucus layer as the cause of reduced MCC. More

specifically, the increased complex viscosity of mucus, as we observed in mucus flakes from

Bpifb1 KO mice and may be the case for soluble mucus as well, is suggestive of mucus that is

more difficult to clear from the airways. *In vitro* studies with human airway cultures[19–22], as well

as earlier studies in model organisms[23–25], have demonstrated that mucus transport is directly

related to the rheological properties of mucus. The increased CBF that we observed in *Bpifb1*

KO mice, although not statistically significant, may be indicative of an attempt to compensate

for altered airways mucus[26]. Such a phenomenon has been demonstrated in human bronchial

epithelial cultures where CBF increased following a moderate increase in mucus viscosity or

mucus load[27].

The increased complex viscosity of mucus flakes in *Bpifb1* KO mice may be attributable to a

change in mucin network structure, which is a function of the density and/or nature of mucin-

mucin/mucin-protein interactions[7]. In muco-obstructive airway disease contexts, for example,

heightened mucus viscoelasticity due to oxidation-dependent mucin cross-linking[5] or increased

mucin concentration[28] has been described. Direct measurement of mucin concentration was

not possible in our experiments due to the high dilution factor of mucus in lavage fluid.

However, the lack of increased MUC5B in pelleted BAL from *Bpifb1* KO mice provides some

evidence against the hypothesis that mucin concentration is increased in mucus flakes and is

the driver of changes in mucus rheology. That said, we detected elevated levels of soluble

MUC5B in *Bpifb1* KO mice, which raises the question of whether MUC5B levels are higher in

Bpifb1 KO mice simply due to decreased MCC or increased mucin concentration. *In vitro*

studies, in which mucus can be collected directly from the surface of epithelial cells, will be

required to determine specifically how BPIFB1 impacts the mucus layer.

Although the polymeric nature of gel-forming mucins is essential for the biophysical properties

of mucus, studies have shown that mucins alone cannot create complex networks, and thus,

other factors, including secreted globular proteins, are necessary for mucus formation[29]. Our

detection of BPIFB1 in secreted mucus from HBECs confirms BPFIB1 is indeed a component of

the mucus protein network and supports the conclusion that this non-mucin protein plays an

important role in the mucus layer. Further evidence for a direct role of BPIFB1 in the mucus

network comes from a finding that BPIFB1 has strong physical interactions with mucin proteins

in HEBC secretions. *Radicioni et al*. found that after treating mucus from HBEC cultures with

guanidinium hydrochloride and detergents (i.e. dissociative conditions), BPIFB1 was one of few

globular proteins interacting with mucins[7]. Additionally, our data demonstrating shared

regulatory mechanisms between BPIFB1 and MUC5B suggest strongly coordinated roles. We

previously found that *Bpifb1* and *Muc5b* gene expression was strongly correlated in mouse lung

tissue[10], and here we have shown that BPIFB1 and MUC5B protein are colocalized in airway

epithelial cells and present in secreted mucus. Cellular co-expression between *BPIFB1* and

MUC5B mRNA has also been demonstrated in human airways by single cell RNA sequencing

studies[30,31] and between BPIFB1 and MUC5AC protein in COPD[32]. Lastly, our finding that BPIFB1

is present in mucin secretory granules, which are tightly packed and only contain the necessary

proteins for mucus, implicates an additional regulatory mechanism coupling these two proteins.

Whether there is an optimal stoichiometry between levels of secreted BPIFB1 and mucin

proteins for MCC, as may be the case between MUC5B and MUC5AC[2,33], remains to be

determined.

In summary, we demonstrate that BPIFB1 is critical for MCC *in vivo*. These findings establish

that mucociliary dysfunction can be driven by non-mucin protein components of airway mucus,

highlighting their importance in health and disease.

METHODS

A detailed descriptions of the methods used in this study can be found in the Supplementary

Methods (Supplementary File 1).

Study Approval.

All experiments conducted with mice in this study were compliant with an Institutional Animal

Care and Use Committee protocol at an animal facility approved and accredited by the

Association for Assessment and Accreditation of Laboratory Animal Care International. Primary

cells were provided by the Cystic Fibrosis (CF) Center Tissue Core Facility of the University of

North Carolina at Chapel Hill under the auspices of protocols approved by the Institutional

Committee on the protection of the rights of human subjects.

Data availability. Complete raw data and counts generated by RNA sequencing were deposited

in the NCBI's Gene Expression Omnibus (GEO) database under entry GSE150452.

AUTHOR CONTRIBUTIONS

208 **ACKNOWLEDGEMENTS**

209 We thank Aiman Abzhanova for assistance with immunostaining. This work was supported in

210 part by NIH HL122711 (SNPK). LJD was supported by NIH F31HL143853 and a grant from the

211 National Institute of General Medical Sciences under award 5T32 GM007092.

212
213
REFERENCES

1.   Fahy J V., Dickey BF. Airway mucus function and dysfunction. *N Engl J Med*.

2011;364(10):978. doi:10.1056/NEJMc1014719

2.   Kesimer M, Ford AA, Ceppe A, et al. Airway mucin concentration as a marker of chronic

bronchitis. *N Engl J Med*. 2017;377(10):911-922. doi:10.1056/NEJMoa1701632

3.   Bonser LR, Zlock L, Finkbeiner W, Erle DJ. Epithelial tethering of MUC5AC-rich mucus

impairs mucociliary transport in asthma. *J Clin Invest*. 2016;126(6):2367-2371.

doi:10.1172/JCI84910

4.   Kesimer M, Makhov AM, Griffith JD, Verdugo P, Sheehan JK. Unpacking a gel-forming

mucin : a view of MUC5B organization after granular release. 2018:15-22.

doi:10.1152/ajplung.00194.2009.

5.   Yuan S, Hollinger M, Lachowicz-Scroggins ME, et al. Oxidation increases mucin polymer

cross-links to stiffen airway mucus gels. *Sci Transl Med*. 2015;7(276):1-10.

doi:10.1126/scitranslmed.3010525

6.   Welsh KG, Rousseau K, Fisher G, et al. MUC5AC and a Glycosylated Variant of MUC5B

Alter Mucin Composition in Children With Acute Asthma. *Chest*. 2017;152(4):771-779.

doi:10.1016/j.chest.2017.07.001

7.   Radicioni G, Cao R, Carpenter J, et al. The innate immune properties of airway mucosal

surfaces are regulated by dynamic interactions between mucins and interacting proteins:

the mucin interactome. *Mucosal Immunol*. 2016;9(6):1442-1454.

doi:10.1038/mi.2016.27

8.   Roy MG, Livraghi-Butrico A, Fletcher AA, et al. Muc5b is required for airway defence.

Nature. 2014;505(7483). doi:10.1038/nature12807

9.   Hancock LA, Hennessy CE, Solomon GM, et al. Muc5b overexpression causes mucociliary

dysfunction and enhances lung fibrosis in mice. *Nat Commun*. 2018;9(1):1-10.

doi:10.1038/s41467-018-07768-9

10.   Donoghue L, Livraghi-Butrico A, McFadden K, et al. Identification of trans Protein QTL for

Secreted Airway Mucins in Mice and a Causal Role for Bpifb1. *Genetics*.

2017;207(October):801-812. doi:10.1534/genetics.117.300211/-/DC1.1

243 11. Evans CM, Raclawska DS, Ttofali F, et al. The polymeric mucin Muc5ac is required for

244   allergic airway hyperreactivity. *Nat Commun*. 2015;6(May 2014).

245   doi:10.1038/ncomms7281

246 12. Bingle L, Wilson K, Musa M, et al. BPIFB1 (LPLUNC1) is upregulated in cystic fibrosis lung

247   disease. *Histochem Cell Biol*. 2012;138(5):749-758. doi:10.1007/s00418-012-0990-8

248 13. Gao J, Ohlmeier S, Nieminen P, et al. Elevated sputum BPIFB1 levels in smokers with

249   chronic obstructive pulmonary disease: a longitudinal study. *Am J Physiol Lung Cell Mol*

250   *Physiol*. 2015;309(1):L17-26. doi:10.1152/ajplung.00082.2015

251 14. Wu J, Kobayashi M, Sousa EA, et al. Differential Proteomic Analysis of Bronchoalveolar

252   Lavage Fluid in Asthmatics following Segmental Antigen Challenge. *Mol Cell Proteomics*.

253   2005;4(9):1251-1264. doi:10.1074/mcp.M500041-MCP200

254 15. Bingle CD, Araujo B, Wallace W a, Hirani N, Bingle L. What is top of the charts?

255   BPIFB1/LPLUNC1 localises to the bronchiolised epithelium in the honeycomb cysts in UIP.

256   *Thorax*. 2013;68(12):1167-1168. doi:10.1136/thoraxjnl-2013-204179

257 16. Camner P, Philipson K, Friberg L. Tracheobronchial Clearance in Twins. 1972:Vol 24.

258 17. Saferali A, Obeidat M, Bérubé J-C, et al. Polymorphisms associated with expression of

259   BPIFA1/BPIFB1 and lung disease severity in cystic fibrosis. *Am J Respir Cell Mol Biol*.

260   2015;53(5):607-614. doi:10.1165/rcmb.2014-0182OC

261 18. Markovetz MR, Subramani DB, Kissner WJ, et al. Endotracheal tube mucus as a source of

262   airway mucus for rheological study. *Am J Physiol Lung Cell Mol Physiol*. 2019;317(4):L498-

263   L509. doi:10.1152/ajplung.00238.2019

264 19. King M. Experimental models for studying mucociliary clearance. *Eur Respir J*.

1998;11(1):222-228. doi:10.1183/09031936.98.11010222

20.   Hill DB, Vasquez PA, Mellnik J, et al. A Biophysical Basis for Mucus Solids Concentration

as a Candidate Biomarker for Airways Disease. 2014;9(2):1-11.

doi:10.1371/journal.pone.0087681

21.   Button B, Cai LH, Ehre C, et al. A periciliary brush promotes the lung health by separating

the mucus layer from airway epithelia. *Science (80-)*. 2012;337(6097):937-941.

doi:10.1126/science.1223012

22.   Carpenter J, Lynch SE, Cribb JA, Kylstra S, Hill DB, Superfine R. Buffer drains and mucus is

transported upward in a tilted mucus clearance assay. *Am J Physiol - Lung Cell Mol*

Physiol. 2018;315(5):L910-L918. doi:10.1152/ajplung.00274.2018

23.   Shih CK, Litt M, Khan MA, Wolf DP. Effect of nondialyzable solids concentration and

viscoelasticity on ciliary transport of tracheal mucus. *Am Rev Respir Dis*. 1977;115(6):989-

995. doi:10.1164/arrd.1977.115.6.989

24.   Lorenzi G, Bohm GM, Guimaraes ET, Costa Vaz MA, King M, Saldiva PHN. Correlation

between rheologic properties and in vitro ciliary transport of rat nasal mucus.

Biorheology. 1992;29(4):433-440. doi:10.3233/BIR-1992-29406

25.   Yu DM, Amidon GL, Weiner ND, Fleisher D, Goldberg AH. The Role of Rheological

Properties in Mucociliary Transport by Frog Palate Ciliated Model. *Pharm Res*.

1994;11(12).

26.   Button B, Okada SF, Frederick CB, Thelin WR, Boucher RC. Mechanosensitive ATP release

maintains proper mucus hydration of airways. *Sci Signal*. 2013;6(279):1-10.

doi:10.1126/scisignal.2003755

27. Liu L, Shastry S, Byan-Parker S, et al. An autoregulatory mechanism governing mucociliary

transport is sensitive to mucus load. *Am J Respir Cell Mol Biol*. 2014;51(4):485-493.

doi:10.1165/rcmb.2013-0499MA

28. Esther CR, Muhlebach MS, Ehre C, et al. Mucus accumulation in the lungs precedes

structural changes and infection in children with cystic fibrosis. *Sci Transl Med*.

2019;11(486). doi:10.1126/scitranslmed.aav3488

29. Thornton DJ, Sheehan JK. From mucins to mucus: toward a more coherent understanding

of this essential barrier. *Proc Am Thorac Soc*. 2004;1(1):54-61. doi:10.1513/pats.2306016

30. Braga FAV, Kar G, Berg M, et al. Cell States in Health and in Asthma. *Nat Med*. 2019.

31. Goldfarbmuren KC, Jackson ND, Sajuthi SP, et al. Dissecting the cellular specificity of

smoking effects and reconstructing lineages in the human airway epithelium. *Nat*

Commun. 2020;11(1). doi:10.1038/s41467-020-16239-z

32. De Smet EG, Seys LJM, Verhamme FM, et al. Association of innate defense proteins

BPIFA1 and BPIFB1 with disease severity in COPD. *Int J COPD*. 2018;13:11-27.

doi:10.2147/COPD.S144136

33. Kirkham S, Kolsum U, Rousseau K, Singh D, Vestbo J, Thornton DJ. MUC5B is the major

mucin in the gel phase of sputum in chronic obstructive pulmonary disease. *Am J Respir*

Crit Care Med. 2008;178(10):1033-1039. doi:10.1164/rccm.200803-391OC

 FIGURE LEGENDS

Figure 1. Decreased mucociliary clearance in *Bpifb1* KO mice. (A) Percent bead clearance in

tracheal mucociliary clearance assay in naive WT and *Bpifb1* KO mice. Mean ± SEM depicted.

n=6/genotype. Significance of two-sided Student's *t*-test denoted as **, $P < 0.01$. (B) Ion

transport measured in Ussing chambers. Mean ± SEM depicted. n=6/genotype. No differences

between genotypes by two-sided Student's *t*-test. (C) Ciliary beat frequency by *in situ* tracheal

assay. Mean ± SEM depicted. n=7-8/genotype. Difference between genotypes not significant by

two-sided Welch's *t*-test, P = 0.1.

Figure 2. Biophysical properties of airway mucus are altered in *Bpifb1* KO mice.

(A) Distribution of rheological signal in BAL from a representative allergen exposed WT and

Bpifb1 KO mouse. (B) Complex viscosity of BAL mucus fraction. Boxplots depict median and

interquartile ranges. n=13/genotype. Significant difference between genotypes by ANCOVA

modeling denoted as *, $P < 0.05$. (C) MUC5B in pellet and supernatant fractions of BAL from

naive WT and *Bpifb1* KO mice by mucin western blotting. Mean ± SEM depicted. Significant

difference between genotypes in supernatant fraction by two-sided Welch's *t*-test denoted as

**, $P < 0.01$. n=5/genotype.

Figure 3: BPIFB1 and MUC5B colocalize in murine airway epithelial cells. (A) BPIFB1 (green)

and MUC5B (red) immunostaining in airway epithelial cells from allergen challenged mice. Scale

bar = 10 μm. (B) Percentage of cells positively stained for BPIFB1 only, MUC5B only, or both in

allergen challenged mice. n = 7 mice. (C) BPIFB1 (green) and MUC5B (red) immunostaining in

large airways of allergen challenged mice demonstrating BPIFB1 staining in luminal mucus.

Scale bar = 50 µm.

Figure 4: BPIFB1 is part of mucus network in human bronchial epithelial cell cultures.

Immunogold-labeled scanning electron microscopy for (A) negative control, (B) MUC5B, and (C)

BPIFB1. Scale bar = 1 µm.

Supplementary Figure Legends provided in Supplementary File 1.

Figure 1. Decreased mucociliary clearance in *Bpifb1* KO mice.
(A) Percent bead clearance in tracheal mucociliary clearance assay in naive WT and *Bpifb1* KO mice. Mean ± SEM depicted. n=6/genotype. Significance of two-sided Student's *t*-test denoted as **, $P < 0.01$. (B) Ion transport measured in Ussing chambers. Mean ± SEM depicted. n=6/genotype. No differences between genotypes by two-sided Student's *t*-test. (C) Ciliary beat frequency by *in situ* tracheal assay. Mean ± SEM depicted. n=7-8/genotype. Difference between genotypes not significant by two-sided Welch's *t*-test, $P = 0.1$.

Figure 2. Biophysical properties of airway mucus are altered in *Bpifb1* KO mice.
(A) Distribution of rheological signal in BAL from a representative allergen exposed WT and *Bpifb1* KO mouse. (B) Complex viscosity of BAL mucus fraction. Boxplots depict median and interquartile ranges. n=13/genotype. Significant difference between genotypes by ANOVA modeling denoted as *, $P < 0.05$. (C) MUC5B in pellet and supernatant fractions of BAL from naive WT and *Bpifb1* KO mice by mucin western blotting. Mean ± SEM depicted. Significant difference between genotypes in supernatant fraction by two-sided Welch's *t*-test denoted as **, $P < 0.01$. n=5/genotype.

Figure 3: BPIFB1 and MUC5B colocalize in murine airway epithelial cells.
(A) BPIFB1 (green) and MUC5B (red) immunostaining in airway epithelial cells from allergen challegned mice. Scale bar = 10 μm. (B) Percentage of cells positively stained for BPIFB1 only, MUC5B only, or both in allergen challegned mice. n=7 mice. (C) BPIFB1 (green) and MUC5B (red) immunostaining in large airways of allergen challegned mice demonstrating BPIFB1 staining in luminal mucus. Scale bar = 50 μm.

Figure 4: BPIFB1 is part of mucus netwrok in human bronchial epithelial cell culutures.
Immunogold-labeled scanned electron microscopy for (A) negative control, (B) MUC5B, and (C) BPIFB1. Scale bar = 1 μm.