Single cell RNA sequence analysis of human bone marrow samples reveals new targets for isolation of skeletal stem cells using DNA-coated gold nanoparticles.

Elloise Matthews[1]*, Stuart Lanham[1]*, Maria-Eleni Kyriazi[2], Konstantina Alexaki[2], Afaf H. El-Sagheer[3], Tom Brown[3], Antonios G. Kanaras[2], Jonathan West[4,5], Ben D. MacArthur[1,5,6], Patrick S. Stumpf[1,7], and Richard O.C. Oreffo[1].

[1] Centre for Human Development, Stem Cells and Regeneration, Human Development and Health, Institute of Developmental Sciences, Faculty of Medicine, University of Southampton, Southampton, SO16 6YD, UK

[2] Physics and Astronomy, Faculty of Physical Sciences and Engineering, University of Southampton, Southampton SO17 1BJ, UK

[3] Department of Chemistry, University of Oxford, Chemistry Research Laboratory, 12 Mansfield Road, Oxford OX1 3TA, UK

[4] Cancer Sciences, Faculty of Medicine, University of Southampton, Southampton, SO16 6YD, UK

[5] Institute for Life Sciences, University of Southampton, Southampton SO17 1BJ, UK

[6] Mathematical Sciences, University of Southampton, Southampton, SO17 1BJ, UK

[7] Joint Research Center for Computational Biomedicine, RWTH Aachen University, Aachen, 52074, Germany

* These authors contributed equally

Article Type: Original Article.

Running Title: Drop-Seq of human bone marrow samples reveals new targets for isolation of skeletal stem cells using DNA-coated gold nanoparticles.

Author's Email:	E. Matthews	E.Z.Matthews@soton.ac.uk
	S. Lanham	S.A.Lanham@soton.ac.uk
	M.E. Kyriazi	M.Kyriazis@soton.ac.uk
	K. Alexaki	K.Alexaki@soton.ac.uk
	A.G. Kanaras	A.Kanaras@soton.ac.uk
	A.H. El-Sagheer	afaf.el-sagheer@chem.ox.ac.uk
	T. Brown	tom.brown@chem.ox.ac.uk
	J. West	J.J.West@soton.ac.uk
	B.D. MacArthur	B.D.Macarthur@soton.ac.uk
	P.S. Stumpf	pstumpf@ukaachen.de
	R.O.C. Oreffo	roco@soton.ac.uk

Corresponding Author: Patrick Stumpf or Richard Oreffo

Bone and Joint Research Group, MP887, Institute of Developmental Sciences, Southampton General Hospital, Tremona Road, SO16 6YD, UK

Fax: +44 2381 205255; Email: pstumpf@ukaachen.de or roco@soton.ac.uk

Abstract Word Count: 165

Manuscript Word Count: 4409

Number of Figures: 6

Number of Tables: 1

Keywords: Spherical nucleic acid, human bone marrow, Drop-Seq, skeletal stem cells

Abstract

There is a wealth of data indicating human bone marrow derived stromal cells (HBMSCs) contain the skeletal stem cell (SSC) with the potential to differentiate along the stromal osteogenic, adipogenic and chondrogenic lineages. However, despite these advances, current methods to isolate skeletal stem cells (SSCs) from human tissues are difficult as no single specific marker has been identified limiting understanding of SSC fate, immunophenotype and widespread clinical application of these cells. While a number of cell surface markers can enrich for SSCs, none of the proposed markers, alone, provide a platform to isolate single cells with the ability to form bone, cartilage, and adipose tissue in humans. The current study details the application of oligonucleotide-coated nanoparticles, spherical nucleic acids (SNAs), to rapidly isolate human cells using mRNAs signatures detected in SSC in real time, to identify stem and progenitor skeletal populations using single cell RNA sequencing. The current approach provides new targets and a platform to advance SSC isolation, enrichment with significant therapeutic impact therein.

INTRODUCTION

Cell fate in culture, tissues, and organisms can be followed using a variety of techniques, although cell phenotype information is typically limited to cell surface epitopes [1]. The detection of a specific mRNA responsible for the expression of a certain protein can be used to determine and characterise the phenotype of the cell. However, traditional methods to assess specific mRNAs such as in-situ hybridisation, northern blot, or quantitative-PCR require cell fixation or lysis to isolate the RNA, resulting in loss of the cells for further experiments.

Bone has the capacity to regenerate, evidence of the presence of a stem cell in bone. While this regenerative capacity has long been recognized, the *in vivo* identity of the skeletal population has only recently been confirmed [2-4]. There is a wealth of data indicating human bone marrow derived stromal cells (HBMSCs) contain the skeletal stem cell (SSC) with the potential to differentiate along the osteogenic, adipogenic and chondrogenic lineages. The availability of a SSC pool has garnered significant interest across the regenerative medicine community given the potential clinical application [5]. Despite this, current methods to isolate SSCs from human tissues remain challenging in the absence of a single specific marker for the SSC.

The current inability, to date, to isolate a homogenous skeletal stem cell population has hampered understanding of SSC fate, immuno-phenotype and simple selection criteria; all limiting factors in the widespread clinical application of these cells. While a range of cell surface markers can enrich for SSCs [2, 6, 7], none of the proposed markers, alone, holds the potential to isolate single cells with the ability to form bone, cartilage, and adipose tissue in humans.

Gold nanoparticles (AuNPs) are easy to synthesise, tuneable and show potential for surface ligand functionalisation with, for example, oligonucleotides. These oligonucleotide functionalised AuNPs, termed spherical nucleic acids (SNAs), display enhanced stability, excellent cell uptake and a degree of resistance to enzyme degradation. Critically, the nanoparticles can be tailored to enhance selectivity and specificity towards the detection of intracellular targets such as RNA.

We have recently developed a protocol using AuNPs to rapidly isolate human cells enriched in SSCs without affecting long term cell viability [8].

In the current studies, we have harnessed SNAs to rapidly isolate human cells using mRNAs signatures detected in SSC in real time, to identify stem and progenitor skeletal populations using single cell RNA sequencing (Drop-Seq). Drop-Seq provides a single cell RNA sequencing platform that allows full transcriptome sequencing of upwards of tens of thousands of individual cells within a single experiment [9]. We have used Drop-Seq to generate mono-disperse nanoliter-sized droplets at >2 kHz for single cell capture. These droplets are stochastically loaded with single cells and, following cell lysis, poly-adenylated transcripts from individual cells are captured on functionalised micro-particles.

In this study, we have applied Drop-Seq to uncover the transcriptome signatures of enriched SSC populations and identify transcripts that distinguish SSCs from other human bone marrow cell types. Our scRNA-seq data reveals candidate markers which highly complement the AuNP cell sorting approach to enrich SSC populations, validated through *in vitro* clonogenic assessment. We present *Decorin, Transferrin* and

Caldesmon1 as novel markers which serve as targets for isolation of SSC populations. The current approach provides new targets and a platform to advance SSC isolation, enrichment with significant therapeutic impact therein.

METHODS

Cell Culture
Bone marrow samples were obtained from haematologically healthy patients undergoing hip replacement surgery under local ethics committee approval local ethics committee approval (LREC194/99/1 and 18/NW/0231 and 210/01). In brief, bone marrow was washed in α-MEM medium and passed through a 70 μm cell strainer. Only samples intended for incubation with SNAs were treated with RosetteSep Human Granulocyte Depletion Cocktail (StemCell Technologies, Cambridge, UK) adding 100 μL to bone marrow resuspended in 5 ml α-MEM medium for 20 minutes. All samples were subjected to density centrifugation using Lymphoprep™ (Lonza). The buffy coat layer, containing bone marrow mononuclear cells was washed in basal medium (α-MEM containing 10% FBS and 100 U mL^{-1} penicillin and 100 μg mL^{-1} streptomycin; Lonza). Cells were subsequently either sorted using MACS or incubated with SNAs.

Enrichment of CD45-/CD146+, CD144+ and CD144-/CD106+ cell populations
Whole bone marrow was diluted 1:1 with α-MEM (Gibco) digested with collagenase IV (20U/mL, Thermo Fisher, 17104019) for 30 min at 37 °C under continuous rotation. Subsequently, bone marrow mononuclear cells were isolated as described by Kanczler et al. 2019 (PMID: 30729460). Where indicated, magnetic cell sorting (MACS) was conducted to enrich or deplete cells. Up to a total of 1x10^8 BM-MNCs were used for each separation. Cells expressing CD45 (Miltenyi Biotec, 130-045-801), CD146 (Miltenyi Biotec, 130-093-596), CD144 (Miltenyi Biotec, 130-097-857), CD106 (Miltenyi Biotec, 130-104-123 and 130-048-102) were isolated using LS columns (Miltenyi Biotec, 130-042-401) according to manufacturer's instructions. Each of three patient samples were sorted into three distinct populations; CD45-/CD146+, CD144+ and CD144-/CD106+ cells.

Stro-1 Enrichment of CFU-Fs
Marrow cells were initially incubated with blocking buffer (α-MEM, 10 % human serum, 5 % FCS and 1 % bovine serum albumin) and subsequently with primary Stro-1 antibody (undiluted hybridoma culture supernatant) [10]. Following washes in buffer (2 mM ethylenediaminetetraacetic acid (EDTA) and 1 % BSA in phosphate buffered saline), cells were incubated with magnetic bead-conjugated secondary antibody (Miltenyi Biotec). After further washes, Stro-1 positive cells were collected by magnetic-activated cell sorting (MACS) according to manufacturer's instructions (Miltenyi Biotec).

Sythesis of spherical nucleic acids
SNAs were manufactured as previously detailed [11] and sequences used detailed in Table 1. All sense strands were modified with FAM dye at the 5' end and a thiol at the 3' end. All flare strands were functionalised with a Cy5 dye at 5' end. For dual-SNA selection, *SPARCL1* SNAs were modified with a JOE fluorophore instead of Cy5 at the 5' end of flare strands. SNAs were added to the cell suspension in basal medium (at 20x10^6 cells per ml) at 0.2 nM for 1 hour at 37 °C with rotation mixing. Cells were

then washed in basal medium and resuspended in FACS solution prior to FACS sorting.

FACS

SNA treated cells were collected from human bone marrow using a FACS Aria cytometer (Becton Dickinson, Wokingham, UK). Cells were gated for monocytes, single cells, and Cy5 fluorescence, with the top 20% of fluorescent cells collected. Positive samples were determined to be within the top half of these Cy5 fluorescent cells and negative samples were considered to be within the bottom half of the collected cells. For dual-SNA studies, a four-way gating system sorted Cy5+/JOE+, Cy5-/JOE+, Cy5+/JOE-, Cy5-/JOE- cells, with only the top 10% of Cy5+/JOE+ collected for plating. Data were analysed on Flowing Software version 2.5 (http://www.flowingsoftware.com). Collected cells were subsequently used for CFU-F quantification, cellular differentiation assays, or Drop-Seq.

Colony Forming Units-Fibroblast (CFU-F) Assay

CFU-F assessment was conducted to demonstrate colony growth [5, 12]. Between 1000 and 2500 positive or negative collected cells were placed into each well of 12 or 6-well tissue culture plates. Cells were grown for 14 days, with a medium change after 7 days. On day 14, wells were washed with PBS and then fixed with 95% ethanol for 10 minutes. The wells were then air dried and 1 mL 0.05% crystal violet solution added to each well for 1 minute. The wells were washed twice with distilled water and the number of stained colonies determined.

Microscopy

Cells were cultured at 37 °C in 5% CO_2 in 24 well plates and imaged using a Zeiss Axiovert 200 inverted microscope with an Axiocam MR camera for fluorescent imaging and Axiovert HR camera for brightfield imaging operated by Zeiss Axiovision software version 4.7.

Statistics

Wilcoxon-Mann-Whitney statistical analysis and ANOVA were performed where appropriate using the SPSS for Windows program version 23 (IBM Corp, Portsmouth, Hampshire, UK). Data is presented as mean ± 95% confidence limits and significance was determined with a p-level of 0.05 or lower.

Drop-Seq of Human Bone Marrow Cells

Drop-Seq was performed as previously described [9] and outlined in the online Drop-Seq Laboratory Protocol version 3.1 (http://mccarrolllab.org/dropseq/) with any adjustments described below. Drop-Seq samples were processed as two separate experiments, with the first including the CD45-/CD146+, CD144+ and CD144-/CD106+ populations and the second including the SNA-Cy5+ and Stro1+ populations (which for the purpose of clarity, will be known as Drop-Seq1 and Drop-Seq2 respectively). In brief, droplet microfluidic devices, according to Macosko et al. [9] were fabricated in poly(dimethylsiloxane) (PDMS) and functionalised to an oleophilic state by incubating channels with 1% trichloro(1H,1H,2H,2H-perfluoro-octyl)silane (Sigma-Aldrich, 448931) in HFE-7500 (3M) for 5–10 min at RT. Cells were co-encapsulated with Drop-Seq beads (ChemGenes: lot 083117) in 1 nL droplets using 15000 mL/hr oil (QX200, Biorad), 4000 mL/hr cell suspension and 4000 mL/hr bead flow rates

(droplet generation frequency 2 kHz). Following cell lysis, the droplet emulsion was broken using perflouroctanol (Sigma-Aldrich) to generate single-cell transcriptomes attached to microparticles (STAMPs). cDNA synthesis was conducted using Maxima H Minus Reverse Transcriptase (Thermo Fisher Scientific). This was followed by PCR amplification (Kapa HiFi Hotstart ReadyMix) using 4+15 PCR cycles for Drop-Seq1 and 4+14 cycles for Drop-Seq2 (95 °C 3 minutes - 4 cycles of: 98 °C 20 s; 65 °C 45 s; 72 °C 3min – 15/14 cycles of: 98 °C 20 s; 67 °C 20 s; 72 °C 3 min - 72 °C 5 min; 4 °C hold). Libraries were tagmented (Nextera XT DNA Library Preparation Kit, Illumina) and amplified before pooling samples for paired-end sequencing using a NextSeq 500/550 High Output Kit v2.5 (Illumina, 20024906) and NextSeq 500 system (Illumina).

Sequence alignment

Raw sequencing reads were aligned following the Drop-Seq Core Computational Protocol (Drop-Seq tools v1.0, http://mccarrolllab.org/dropseq/). Sequencing data was aligned to human hg19 reference genome (GSM1629193) using STAR software [13]. Raw sequencing reads were demultiplexed, grouping reads by cell barcode to generate a digital gene expression (DGE) matrix for downstream analysis, using Drop-Seq tools (v1.0). A modified multi-mapper pipeline was executed to correct multiple alignment [14].

Data clustering

Analysis of the Drop-Seq1 and Drop-Seq2 datasets was conducted using the software R (version3.5.0) according to the standard pipeline of functions included in the R Seurat package (version 3.0) (http://satijalab.org/seurat/). Drop-Seq1 and Drop-Seq2 datasets were processed and analysed separately.

Drop-Seq1

The Drop-Seq1 dataset, comprising 26,657 cells, was subjected to various quality control measures; filtering out of low quality cells (cells expressing <200 genes or >10% mitochondrial genes) and potential cell doublets (cells expressing >4,500 genes). Cells were then normalised using the function; *sctransform*. Linear dimensionality reduction was performed using Principle Component Analysis (PCA) and non-linear dimension reduction using Uniform Manifold Approximation and Projection (UMAP). Clustering analysis was conducted using the Seurat function, *FindClusters*, at a resolution of 1.1 and default parameters. To assign identifies to the clusters, the function, *FindAllMarkers*, was performed to characterise significant cluster-specific gene expression for comparison against previously described biomarkers for bone marrow cell populations. This revealed 15 distinct cell types. Pericyte and endothelial cell clusters were defined by expression of *VWF* and *LEPR* respectively.

Drop-Seq2

For the Drop-Seq2 dataset, comprising 1,637 cells, quality control thresholds were more refined due to the reduced number of cells. Cells expressing <8% mitochondrial RNA, or <200/>2,500 unique reads were filtered from the dataset. Cells were then normalised using the function; *sctransform*. Linear dimensionality reduction was performed using Principle Component Analysis (PCA) and non-linear dimension reduction using Uniform Manifold Approximation and Projection (UMAP). Clustering analysis was conducted using the Seurat function, *FindClusters*, at a resolution of 1.3 and default parameters. This revealed 28 clusters which were broadly classified into

8 cell types. Skeletal stem cell cluster was defined by expression of *CXCL12, LEPR* and *SPARCL1.*

Table 1. Sense and flare oligonucleotide sequences. X: thiol modifier 6 S-S (CPG resin from Glen Research)

SNA Target	Sense Strand Sequence (5'to 3') with modifications	Flare Strand Sequence (5' to 3') with modifications
CALD1	FAM – TATTGCTGCTTGATGGGTCGATAAAAAAAA – X	Cy5 – ATCGACCCATCAAGC
CD73	FAM – GTTGCGTTCATCAATGGGCGACAAAAAAAA – X	Cy5 – GTCGCCCATTGA
CD164	FAM – GTCTCATGCCTGCGATAGGGACAAAAAAAA – X	Cy5 – GTCCCTATCGCA
CD200	FAM – GGTCCTGATTCCGGTGACGTTTAAAAAAAA – X	Cy5 – AAACGTCACCGG
Ceruloplasmin	FAM – TATATTCCATGTACATCGGCCTAAAAAAAA – X	Cy5 – AGGCCGATGTAC
Col1a2	FAM – CCTCCAACTTAGCCGAAACCTCAAAAAAAA – X	Cy5 – GAGGTTTCGGCTAAG
CXCL12	FAM – GAATTTTGAGATGCTTGACGTTAAAAAAAA – X	Cy5 – AACGTCAAGCAT
Decorin	FAM – GCACTTTGTCCAGACCCAAATCAAAAAAAA – X	Cy5 – GATTTGGGTCTGG
FMO3	FAM – GGTTCCAAATCGAGTGACGAGCAAAAAAAA – X	Cy5 – GCTCGTCACTCGAT
HSPA8	FAM – AGCAGTACGGAGGCGTCTTACAAAAAAAA – X	Cy5 – TGTAAGACGCCTC
Osterix	FAM – ACTCAGGCTCCGGTCCTACAGTAAAAAAAA – X	Cy5 – ACTGTAGGACCG
Podoplanin	FAM – ACCAATGAAGCCGATGGCTAGTAAAAAAAA – X	Cy5 – ACTAGCCATCGG
RUNX2	FAM – TGTGGTTGTTTGTGAGGCGAATAAAAAAAA – X	Cy5 – ATTCGCCTCACA
Scrambled	FAM – ATGGTATACCGAAAGACTGTTAAAAA – X	Cy5 – AACAGTCTTTCG
SPARCL1	FAM – GTTGGCACCAGAGAGTATTTCAAAAAAAAA – X	Cy5 or JOE – TGAAATACTCTCTGG
Transferrin	FAM – GACGCTTTTCATATGGTCGCGGAAAAAAAA – X	Cy5 – CCGCGACCATATG

RESULTS AND DISCUSSION

Drop-Seq of whole bone marrow reveals SPARCL1 as a potential SSC marker

A sub-endothelial perivascular origin of the SSCs in bone marrow is widely acknowledged [3, 15-17]. To characterise the cellular heterogeneity within human adult bone marrow and identify potential markers for SSC enrichment, scRNA-seq was performed, using the Drop-Seq methodology [9]. Three cell populations per patient were fractioned for sequencing; the previously described CD45-/CD146+ skeletal progenitor population [3], CD144+ endothelial cells [18] and CD144-/CD106+ pericytes [19]. In total, **9,795** cells were profiled, reduced to **7,385** following quality control processing. To further increase the cell pool for identification of skeletal progenitors, this data was combined with sc-RNAseq data for unsorted bone marrow populations from 3 patients [14](data available from ArrayExpress under accession number E-MTAB-8630), enabling comparison between depleted and unfractioned bone marrow. In total, the combined data, shown in figure 1A, comprised 17,527 cells with an average of 2,082 transcripts per cell.

An unsupervised clustering strategy was employed to detect distinct cell subtypes with unique gene expression signatures [20]. Briefly, 9 cell clusters were identified, corresponding to hematopoietic and non-hematopoietic cell types (Figure 1B). Guided by the localised expression of established cell type markers, clusters were labelled as haematopoietic progenitors (*CD34*+), granulocytes (*MPO+,LTF*+), erythroblasts (*HBD*+), B lymphocytes (*MZB1*+), monocytes (*CD14*+), T lymphocytes (*CD8A*+), dendrocytes (*FCGERA*+), endothelial cells (*VWF*+) and pericytes (*LEPR+,CXCL12*+) (Supplementary figure 1).

We next sought to identify candidate SSC markers, through examination of localised gene expression in pericytes and endothelial cell clusters, representing 233 and 460 cells respectively. Analysis of differential gene expression between pericytes/endothelial cells, termed 'Skeletal progenitors', and other cell types indicated *SPARCL1,* as a potential SSC marker due to elevated expression (Figure 1C*). FABP4, CXCL12, JUN* and *SPARCL1* were among the 4 most differentially expressed genes in pericytes and endothelial cells (Figure 1D-G). However, from these mRNA, *SPARCL1* expression was strongly localised to the SSC region, with expression observed in 55.5% of skeletal progenitors and only 1.3% in other cell types. Given that *SPARCL1* has been previously shown to activate BMP/TGF-β signalling to promote the differentiation of C2C12 cells in mice [21], is highly homogenous to bone regulatory protein SPARC/OCN [22], new SNAs were designed against *SPARCL1* mRNA to investigate *SPARCL1* as a new target for SSC enrichment.

Figure 1: Unsupervised clustering and UMAP plots, with each point representative of a single cell. A) Cell populations were mapped onto the UMAP representation to visualise spread of cells across clusters**. B**) Identification of lineage biomarkers characterised the clusters into 9 cell types. C) Top ten differentially expressed markers for skeletal progenitor clusters when compared to all other cell types. Embedding heatmap plots shown localised expression of D) *FABP4 E*) *CXCL12* F) *JUN* G) *SPARCL1.*

SNAs targeting *SPARCL1, CD200* and *CD146* mRNA enrich for CFU-F

To test suitable candidate SNAs for SSC isolation, SNAs were designed targeting a range of mRNAs; these included SPARCL1, as identified by scRNA-seq, and other SNAs including: *HSPA8 (*encodes antigen detected by STRO-1 antibody)[23], *RUNX2* [24]*,* and scrambled control. Other potential mRNA targets were *CD164, CD146* [3], *SP7* (Osterix) [24] and *CXCL12* [25]. Recently, Chan et al., 2018 and Debnath et al., 2018 have suggested likely candidate protein markers for SSCs [6, 26], and thus, SNAs were designed targeting *CD73, CD200*, and *PODOPLANIN (PDPN).* The top 20% of fluorescent cells were collected following incubation with the various SNAs, and positive and negative fractions were deemed the top half and bottom half of the collected cells, respectively. Each SNA was tested against a minimum of 3 different patients, and up to 3 SNAs were tested for each patient. In total, bone marrow samples from 18 different patients were tested.

To determine the clonogenic capacity of SNA-sorted populations, cells were plated at limiting dilution, whereby each fibroblastic colony formed, is derived from a single SSC or progenitor, termed CFU-F [27]. The normalised CFU-F counts as a percentage of the unsorted cells are shown in figure 2. Enrichment of CFU-F in comparison to scramble control and unsorted cells was observed in all positive fractions, while the negative fractions displayed minimal CFU-F. The SNAs which showed the most consistent CFU-F enrichment in comparison to unsorted cells, were *CD200, SPARCL1, CD146* and *osterix.* Interestingly, the Osterix SNA also demonstrated regular CFU-F enrichment in the negative cell fraction. In agreement with the scRNA-seq data, *SPARCL1 mRNA* sorts cells with enhanced clonogenic functionality. Similarly, *CD200+* and *CD146+* were defined as populations of interest for further scRNA-seq analysis, consistent with previous findings. These results are consistent with previous findings which identify *CD200*+ [6, 28] and *CD146*+ [3, 17, 29] as viable targets for SSC isolation. However, limitations to both populations have been identified; *CD200 mRNA* displays poor SSC-specificity [30, 31], while SSCs have been identified in both *CD146+* and *CD146-/low* fractions [32, 33]. Therefore, these populations do not likely encompass a purified pool encompassing all SSCs, though will generate a valuable resource, together with *SPARCL1+,* for further investigation and characterisation of SSCs.

Figure 2: Cells enriched for *CD146, CD200*, and *SPARCL1* targets demonstrate enhanced CFU-F capacity in comparison to unselected cells. SNAs were designed to target a wide range of molecular markers, identified from scRNA-Seq libraries and surrounding literature. The positive (red) and negative (black) populations, together with unsorted cells (green), were plated at 5,000 cells per well of a 12-well plate and colonies were counted after two weeks of culture. For each SNA, each point represents the mean CFU-F count from a different patient, displayed as a percentage of unsorted CFU-F counts. Horizontal bars represent mean with SD error bars.

SPARCL1 SNA displays enhanced specificity for CFU-F formation compared to traditional Stro1+ MACS populations

Following the enrichment in CFU-F counts, studies were undertaken to evaluate if the *SPARCL1+* SNA-selected cells displayed enhanced CFU-F enrichment compared to the current SSC enrichment methodology using MACS to select Stro-1+ cells. BM-MNCs were divided into two parts and processed separately. The first fraction was incubated with SNAs targeting *SPARCL1 mRNA* and cells were collected using FACS. Meanwhile, the remainder of the sample was incubated with Stro-1 antibody and Stro-1+ cells were collected using the MACS methodology [34]. The Stro-1+ cells were sequentially sorted using the *SPARCL1* SNA to collect Stro1+*SPARCL1+* populations. Positive and negative fractions were plated to determine CFU-F enrichment.

Evaluation of data from patients 39 and 40 (Figure 3) demonstrated *SPARCL1+* SNA-selected cells produced more CFU-F than unsorted and *SPARCL1-* fractions (black bars), although this did not reach significance. Additionally, no significant difference was observed between *SPARCL1+* SNA-selected cells and Stro-1+ MACS-only collected cells for patients 38 and 40, indicating *SPARCL1+* cells contain the same number of CFU-F as Stro-1+ cells. It was previously shown that all CFU-F reside within Stro-1+ fraction [35], therefore the current study suggests *SPARCL1* is a comparable target for SSC enrichment, with the additional benefits of the SNA cell isolation protocol being less time-consuming and therefore preferred. Furthermore, when Stro-1+ were subsequently sorted using *SPARCL1* SNAs, significantly enhanced CFU-F enrichment was observed in Stro-1+*SPARCL1+* cells compared to Stro1+*SPARCL1-* (p<0.001 for all patients) and *Stro-1+* (only) (p≤0.03 for all patients) populations, indicating the *SPARCL1* SNA holds enhanced specificity for CFU-Fs than Stro-1 antibody.

Figure 3: Comparison of SPARCL1 SNA to Stro-1 antibody for CFU-F isolation. A. Schematic of experimental protocol for different cell population collected. B. Cells were either unselected (black bars) or Stro-1+ selected cells (green bars) from the same patient sample. Only selected statistics are shown for clarity. ns - not significant, * p<0.05, *** p<0.001. Bars show mean with 95% confidence limits. n=3 for each timepoint for each patient.

Drop-Seq of SNA-selected populations reveal novel targets for SSC isolation

The scarcity of SSC within human bone marrow, <1 in 100,000 of bone mononuclear cells [36], presents a significant obstacle for transcriptomic profiling of SSCs, as limitations of scRNA-seq protocols in detecting cell types with a frequency less than 1% have been recently reported as these cells typically appear as outliers [37]. Therefore, pre-sorting cells prior to downstream scRNA-seq can increase read depth by reducing competition for sequencing capacity between cells and genes [38]. Importantly, reducing the frequency of redundant cell types increases the allocation of

sequencing power for the characterisation of rare cells and identification of unique biomarkers. In order to increase the SSC pool for scRNA-seq, Drop-Seq of MACS-collected Stro-1+ cells, and cells enriched using SNAs targeting *CD200, SPARCL1* and *CD146 mRNAs* were performed (Figure 4A). In total the transcriptomes of 1,637 cells were sequenced, producing a dataset of 1,363 cells with an average of 1,923 transcripts per cell following quality control filtering. Unsupervised clustering produced 15 clusters which, following analysis of differential gene expression, were classified as 8 distinct cell types; Skeletal stem cells/progenitors, monocytes, erythroblasts, granulocytes, B lymphocytes, dendritic cells, T lymphocytes and Haematopoietic stem cells (Figure 5B). For example, clusters 0,3,6 and 11 showed high expression of the markers; *Glycoprotein-A* (GYPA) and Haemoglobin subunit alpha 1 (HBA1) (Supplementary figure 2), therefore identifying these cell clusters as erythroblasts.

Skeletal progenitor identities were assigned based on expression of stromal markers; *CXCL12* [25] and *LEPR* [39], and the expression of *SPARCL1*. The skeletal progenitor cluster consisted of 18.75% *SPARCL1+* cells and 81.25% Stro-1+ cells, supporting the presence of SSCs within these enriched populations. No cells belonging to the CD146+ and CD200+ sequenced populations were represented within the cluster of skeletal progenitors. Overall, the sequenced populations maintained a degree of heterogeneity and included cells across the classified cell-types; *SPARCL1+* cells were represented in all 8 cell-type subpopulations (Figure 4C). Despite this observed heterogeneity, the SNA methodology was found to enrich for monocytes; with *CD200+, CD146+* and *SPARCL1+* populations consisting of 66%, 42% and 56% monocytes, respectively (Figure 4C), therefore demonstrating the efficiency of the SNA protocol in filtering undesirable cell types. Similarly, the MACS Stro-1+ population contained a high proportion of monocytes (37%), though Stro1+ cells were largely erythroblasts (52%), consistent with the findings of Simmons and Torok-Strob who demonstrated that CFU-F predominantly resided within the Stro-1+/GYPA- fraction [35].

Figure 5: Drop-Seq analysis of 1,363 selected HBMSCs. UMAP was conducted for visualisation purposes to reduce the data to two dimensions. Each point represents a single cell. A) Cells were sorted using SNAs targeting CD146, CD200 and SPARCL1 and sequenced, together with a MACS Stro-1+ population, following the Drop-Seq methodology. B) Unsupervised clustering at a resolution of 1.3 identified 15 subpopulations based on gene expression profiles. Clusters

were classified into 8 distinct cell subpopulations, corresponding to haematopoietic and non-haematopoietic cell types. C) The proportion of cells from each sequenced population represented across the cell-type clusters reveals enrichment of monocytes within SNA-sorted populations (CD200+, CD146+ and SPARCL1+). The Stro1+ population is predominantly erythroblasts and monocytes.

Novel SSC targets; *DCN, TF* and *CALD1* show consistent enrichment of CFU-F and enhanced enrichment of CFU-F in comparison to *SPARCL1+* cells.

To further characterise the molecular signature of SSCs, differential gene analysis was performed to identify markers with localised expressed in skeletal progenitor cell types. The top ten differentially expressed genes in the SSC clusters are shown in figure 5A, all potential candidate SSC markers given the extremely low representation within other cell types; >0.75%. The most highly expressed differential marker was *Decorin* (*DCN*), expressed by 63% of skeletal progenitors with an avg_LogFC in gene expression of ~1.5. In addition to *DCN,* 4 other markers; *Collagen Type I Alpha 2 Chain (COL1A2), Flavin-containing monoxygenase-3 (FMO3), Transferrin (TF)* and *Caldesmon (CALD1)* were identified as potential candidates for SSC enrichment based on exclusivity and level of expression in skeletal progenitors compared to other cell types (figure 5A).

To assess the functionality of the novel markers, for isolation of SSCs, SNAs targeting *DCN, TF, COL1A2, CALD1,* and *FMO3* were designed. HBMSCs were incubated with the SNAs for 1 hour before positive (top 10 % brightest Cy5+) and negative (next 10 % Cy5+) cell fractions were sorted. SNAs were tested on a minimum of two different patient samples per target. For each sample, positive, negative and unsorted cells were plated at 5,000 cells per well of a 12 well plate to assess SSC enrichment within each fraction. Following crystal violet staining, visible clusters were counted, indicative of the number of SSC colonies. The number of CFU-F for each selected population was expressed as a percentage of unsorted cells (Figure 5B). Furthermore, the CFU-F value for *SPARCL1+* and *CD146+* cells (previously showed the highest CFU-F enrichment), was included to allow comparison with the new SNA populations. For each SNA-target, enrichment for CFU-F was observed in all positive fractions. *DCN+, TF+* and *CALD1+* cells showed enhanced enrichment in comparison to *SPARCL1+* cells and comparable enrichment to *CD146+* cells, and therefore defined as candidate SSC biomarkers. A role of *DCN* in skeletal development has been identified previously; high expression has been reported in a variety of cells, including osteoblasts, perivascular chondrocytes and throughout the bone periosteum [40]. Additionally, DCN can function in signalling cascades; regulating cell differentiation, angiogenesis and bone mineralisation [41-46]. Similarly, the essential role of *TF* in iron-delivery and consequently, cell growth and survival, correlates with highly proliferative cell types and anti-apoptotic activity [47-50]. Finally, *CALD1* functions in calcium-mediated signalling pathways in bone [51], and its expression in BMMNCs has been reported previously [52]. In summary, there is emerging evidence to substantiate *DCN, TF* and *CALD1* as useful targets to explore for SSC isolation.

MACS provides a less time-consuming approach to cell sorting of large cell numbers than the SNA-based FACS strategy applied in this study, although MACS relies on the detection of surface epitopes [53]. It was therefore of interest to investigate whether the same level of SSC enrichment, observed following SNA-based cell sorting, could

be achieved using MACS, with *DCN* and *TF* as antigen targets. Negative and positive fractions were collected for three samples and cells were plated at 50,000 cells per 3 wells of a 12-well plate for CFU-F assessment. In marked contrast to the CFU-F enrichment observed in *DCN+* and *TF+* populations following AuNP mRNA detection, cells sorted using DCN and TF antibodies did not display CFU-F capacity in positive fractions following MACS (figure 5C). This result indicates the molecular targets identified by scRNA-seq for SNA cell sorting are not suitable for surface epitope-based strategies, confirming mRNA levels do not directly correlate into protein synthesis [54-56]. This limitation was previously described by Fitter et al who identified no correlation between *HSPA8* (mRNA) and Stro-1 (surface protein) [23].

Figure 5: A) **Proportion of 'Skeletal Progenitors' and 'Other cell types' expressing the top 10 differentially expressed genes.** The ten most differentially expressed genes were plotted, with data-point size indicative of the average logarithm fold change (avg_logFC) in gene expression. B**) SNAs were designed to target DCN, TF, CALD1, COL1A2 and FMO3; markers identified following Drop-Seq of Stro-1+, SPARCL1+, CD200+ and CD146+ populations.** The positive and negative populations, together with unsorted cells, were plated at 5,000 cells per well of a 12-well plate and colonies were counted after two weeks of culture. For each SNA, each point represents the mean CFU-F count from a different patient, displayed as a percentage of unsorted CFU-F counts. Horizontal bars represent mean with SD error bars. C) **Average CFU-F count for positive and negative fractions for DCN and TF MACS collected selected populations, together with CFU-F counts in unsorted cells of the same samples.** *MACS was performed to sort HBMSCs based on expression of Decorin and Transferrin. Positive and negative fractions, together with unsorted HBMSCs, were plated at 50,000 cells per well*

in a 12 well plate (5000 cells for unsorted). After two weeks of culture, colonies were stained and CFU-F counted.

Dual-SNA incubation

Following identification of SNA targets that sort populations with an enhanced CFU-F capacity, 2nd generation SNAs targeting *SPARCL1 were* designed using JOE fluorescent dye in the place of Cy5 on the 5' end of flare strands. The choice of dye made no difference to the CFU-F count isolated (data not shown). Cells were incubated with *SPARCL1* JOE SNAs, together with Cy5 SNAs targeting either *CD146, CALD1, DCN, COL1A2, DCN, FMO3* or *TF*. The top 10% Cy5+/JOE+ fluorescent cells were collected and plated for CFU-F assessment. *SPARCL1+TF+* cells demonstrated the most enhanced CFU-F enrichment compared to unsorted cells (figure 6).

Figure 6: CFU-F following dual-SNA selection. Cells were incubated with *SPARCL1* JOE-tagged AuNPs and Cy5-tagged AuNPs targeting *CD146, CALD1, DCN, COL1A2, DCN, FMO3* or *TF*. The top 10% Cy5+/JOE+ fluorescent cells were collected and plated at 5000 cells per well on a 12 well plate. For each SNA combination, each point represents the mean CFU-F count from a different patient, displayed as a percentage of unsorted CFU-F counts. Horizontal bars represent mean with SD error bars.

CONCLUSION

In summary, we demonstrate single cell RNA sequencing provides a powerful tool for the parallel single-cell sequencing of human adult bone marrow utilizing DNA-coated gold nanoparticles. This approach revealed vast heterogeneity within unsorted and enriched bone marrow populations. Initial profiling of pericyte, endothelial, skeletal progenitor and unsorted cells revealed *SPARCL1+* as a potential SSC biomarker.

When targeted using AuNP mRNA detection, *SPARCL1+* cells displayed a 3-fold enhanced capacity for CFU-F in comparison to unsorted cells and improved specificity for CFU-F enrichment than the widely accepted Stro-1+ pool of SSCs [35]. Similarly, *CD146* and *CD200* were identified as mRNA targets for AuNP cell sorting to enrich CFU-F in SSC populations. Delivery of *SPARCL1+, CD146+, CD200+,* and Stro-1+ populations into the Drop-Seq methodology revealed a second generation of markers, which demonstrated improved capacity for CFU-F enrichment. *DCN, TF* and *CALD1* were identified as novel markers that increased CFU-F capacity when used both alone or in conjunction to *SPARCL1*-targeted SNAs.

The current studies demonstrate, for the first time, that Drop-Seq can reveal novel markers for SSC isolation, which sort cells with clonogenic functionality *in vitro*. Furthermore, we demonstrate scRNA-seq approaches identify markers that highly complementary for cell sorting strategies harnessing mRNA detection, like the AuNP protocol applied in this study, in contrast to epitope expression. Overall, the reported findings describe novel markers that enable the isolation of an enriched pool of SSCs from human adult bone marrow, a valuable resource for future development of SSC-based therapies in the treatment of skeletal damage and disease.

Acknowledgements

ROCO and AGK acknowledge financial support from the Biotechnology and Biological Sciences Research Council (BB/P017711/1). ROCO acknowledges support from the UK Regenerative Medicine Platform "Acellular / Smart Materials – 3D Architecture" (MR/R015651/1), the Rosetrees Trust, Wessex Medical Research as well as Institute for Life Sciences and University of Southampton. PSS, BDM and JJW acknowledge funding from the Medical Research Council (MC_PC_15078).

References

1. Specht, E.A., E. Braselmann, and A.E. Palmer, *A Critical and Comparative Review of Fluorescent Tools for Live-Cell Imaging.* Annu Rev Physiol, 2017. **79**: p. 93-117.
2. Chan, C.K.F., et al., *Identification of the Human Skeletal Stem Cell.* Cell, 2018. **175**(1): p. 43-56 e21.
3. Sacchetti, B., et al., *Self-renewing osteoprogenitors in bone marrow sinusoids can organize a hematopoietic microenvironment.* Cell, 2007. **131**(2): p. 324-36.
4. Mendez-Ferrer, S., et al., *Mesenchymal and haematopoietic stem cells form a unique bone marrow niche.* Nature, 2010. **466**(7308): p. 829-34.
5. Dawson, J.I., et al., *Concise review: bridging the gap: bone regeneration using skeletal stem cell-based strategies - where are we now?* Stem Cells, 2014. **32**(1): p. 35-44.
6. Debnath, S., et al., *Discovery of a periosteal stem cell mediating intramembranous bone formation.* Nature, 2018. **562**(7725): p. 133-139.
7. Gothard, D., et al., *Prospective isolation of human bone marrow stromal cell subsets: A comparative study between Stro-1-, CD146- and CD105-enriched populations.* J Tissue Eng, 2014. **5**: p. 2041731414551763.
8. S.A. Lanham, M.X., M.E. Kyriazi, K. Alexaki, A.H. El-Sagheer, T. Brown, A.G. Kanaras, R.O.C. Oreffo, *Optimising DNA-Coated Gold Nanoparticles to Enrich Skeletal Stem Cells from Human Bone Marrow Samples.* Biorxiv preprint, 2020.

9.   Macosko, E.Z., et al., *Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets.* Cell, 2015. **161**(5): p. 1202-1214.

10.   Howard, D., et al., *Immunoselection and adenoviral genetic modulation of human osteoprogenitors: in vivo bone formation on PLA scaffold.* Biochemical and Biophysical Research Communications, 2002. **299**(2): p. 208-215.

11.   Kyriazi, M.-E., et al., *Multiplexed mRNA Sensing and Combinatorial-Targeted Drug Delivery Using DNA-Gold Nanoparticle Dimers.* ACS Nano, 2018. **12**(4): p. 3333-3340.

12.   Sengers, B.G., J.I. Dawson, and R.O. Oreffo, *Characterisation of human bone marrow stromal cell heterogeneity for skeletal regeneration strategies using a two-stage colony assay and computational modelling.* Bone, 2010. **46**(2): p. 496-503.

13.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-21.

14.   Stumpf, P.S., et al., *Mapping biology from mouse to man using transfer learning.* bioRxiv, 2019: p. 2019.12.26.888842.

15.   Ambrosi, T.H., M.T. Longaker, and C.K.F. Chan, *A Revised Perspective of Skeletal Stem Cell Biology.* Frontiers in Cell and Developmental Biology, 2019. **7**(189).

16.   Crisan, M., et al., *A perivascular origin for mesenchymal stem cells in multiple human organs.* Cell Stem Cell, 2008. **3**(3): p. 301-13.

17.   Shi, S. and S. Gronthos, *Perivascular niche of postnatal mesenchymal stem cells in human bone marrow and dental pulp.* J Bone Miner Res, 2003. **18**(4): p. 696-704.

18.   Lampugnani, M.G., et al., *A novel endothelial-specific membrane protein is a marker of cell-cell contacts.* J Cell Biol, 1992. **118**(6): p. 1511-22.

19.   Gronthos, S., et al., *Molecular and cellular characterisation of highly purified stromal stem cells derived from human bone marrow.* J Cell Sci, 2003. **116**(Pt 9): p. 1827-35.

20.   Satija, R., et al., *Spatial reconstruction of single-cell gene expression data.* Nat Biotechnol, 2015. **33**(5): p. 495-502.

21.   Wang, Y., et al., *SPARCL1 promotes C2C12 cell differentiation via BMP7-mediated BMP/TGF-beta cell signaling pathway.* Cell Death Dis, 2019. **10**(11): p. 852.

22.   Mansergh, F.C., et al., *Osteopenia in Sparc (osteonectin)-deficient mice: characterization of phenotypic determinants of femoral strength and changes in gene expression.* Physiol Genomics, 2007. **32**(1): p. 64-73.

23.   Fitter, S., et al., *The Mesenchymal Precursor Cell Marker Antibody STRO-1 Binds to Cell Surface Heat Shock Cognate 70.* STEM CELLS, 2017. **35**(4): p. 940-951.

24.   Nakashima, K. and B. de Crombrugghe, *Transcriptional mechanisms in osteoblast differentiation and bone formation.* Trends Genet, 2003. **19**(8): p. 458-66.

25.   Sugiyama, T., et al., *Maintenance of the hematopoietic stem cell pool by CXCL12-CXCR4 chemokine signaling in bone marrow stromal cell niches.* Immunity, 2006. **25**(6): p. 977-88.

26.   Chan, C.K.F., et al., *Identification of the Human Skeletal Stem Cell.* Cell, 2018. **175**(1): p. 43-56.e21.

27.   Friedenstein, A.J., R.K. Chailakhjan, and K.S. Lalykina, *The development of fibroblast colonies in monolayer cultures of guinea-pig bone marrow and spleen cells.* Cell Tissue Kinet, 1970. **3**(4): p. 393-403.

28.   Delorme, B., et al., *Specific lineage-priming of bone marrow mesenchymal stem cells provides the molecular framework for their plasticity.* Stem Cells, 2009. **27**(5): p. 1142-51.

29.   Sorrentino, A., et al., *Isolation and characterization of CD146+ multipotent mesenchymal stromal cells.* Exp Hematol, 2008. **36**(8): p. 1035-46.

30. Rostovskaya, M. and K. Anastassiadis, *Differential expression of surface markers in mouse bone marrow mesenchymal stromal cell subpopulations with distinct lineage commitment.* PLoS One, 2012. **7**(12): p. e51221.

31. Barclay, A.N., et al., *CD200 and membrane protein interactions in the control of myeloid cells.* Trends Immunol, 2002. **23**(6): p. 285-90.

32. Tormin, A., et al., *CD146 expression on primary nonhematopoietic bone marrow stem cells is correlated with in situ localization.* Blood, 2011. **117**(19): p. 5067-77.

33. Harkness, L., et al., *CD146/MCAM defines functionality of human bone marrow stromal stem cell populations.* Stem Cell Research & Therapy, 2016. **7**(1): p. 4.

34. Kanczler, J., et al., *Isolation, Differentiation, and Characterization of Human Bone Marrow Stem Cells In Vitro and In Vivo.* Methods Mol Biol, 2019. **1914**: p. 53-70.

35. Simmons, P.J. and B. Torok-Storb, *Identification of stromal cell precursors in human bone marrow by a novel monoclonal antibody, STRO-1.* Blood, 1991. **78**(1): p. 55-62.

36. Pittenger, M.F., et al., *Multilineage Potential of Adult Human Mesenchymal Stem Cells.* Science, 1999. **284**(5411): p. 143-147.

37. Zheng, G.X., et al., *Massively parallel digital transcriptional profiling of single cells.* Nat Commun, 2017. **8**: p. 14049.

38. Haque, A., et al., *A practical guide to single-cell RNA-sequencing for biomedical research and clinical applications.* Genome Med, 2017. **9**(1): p. 75.

39. Zhou, B.O., et al., *Leptin-receptor-expressing mesenchymal stromal cells represent the main source of bone formed by adult bone marrow.* Cell Stem Cell, 2014. **15**(2): p. 154-68.

40. Bianco, P., et al., *Expression and localization of the two small proteoglycans biglycan and decorin in developing human skeletal and non-skeletal tissues.* J Histochem Cytochem, 1990. **38**(11): p. 1549-63.

41. Santra, M., et al., *Ectopic decorin expression up-regulates VEGF expression in mouse cerebral endothelial cells via activation of the transcription factors Sp1, HIF1α, and Stat3.* Journal of Neurochemistry, 2008. **105**(2): p. 324-337.

42. Grant, D.S., et al., *Decorin suppresses tumor cell-mediated angiogenesis.* Oncogene, 2002. **21**(31): p. 4765-77.

43. Cabello-Verrugio, C. and E. Brandan, *A novel modulatory mechanism of transforming growth factor-beta signaling through decorin and LRP-1.* J Biol Chem, 2007. **282**(26): p. 18842-50.

44. Bi, Y., et al., *Extracellular matrix proteoglycans control the fate of bone marrow stromal cells.* J Biol Chem, 2005. **280**(34): p. 30481-9.

45. Barry, F., et al., *Chondrogenic differentiation of mesenchymal stem cells from bone marrow: differentiation-dependent gene expression of matrix components.* Exp Cell Res, 2001. **268**(2): p. 189-200.

46. Egusa, H., et al., *Downregulation of extracellular matrix-related gene clusters during osteogenic differentiation of human bone marrow- and adipose tissue-derived stromal cells.* Tissue Eng, 2007. **13**(10): p. 2589-600.

47. Nunez, M.T., et al., *Transferrin receptors in developing murine erythroid cells.* Br J Haematol, 1977. **36**(4): p. 519-26.

48. Hamilton, T.A., H.G. Wada, and H.H. Sussman, *Identification of transferrin receptors on the surface of human cultured cells.* Proc Natl Acad Sci U S A, 1979. **76**(12): p. 6406-10.

49. Lesnikov, V., M. Lesnikova, and H.J. Deeg, *Pro-apoptotic and anti-apoptotic effects of transferrin and transferrin-derived glycans on hematopoietic cells and lymphocytes.* Exp Hematol, 2001. **29**(4): p. 477-89.

50. Netz, D.J., et al., *Eukaryotic DNA polymerases require an iron-sulfur cluster for the formation of active complexes.* Nat Chem Biol, 2011. **8**(1): p. 125-32.

51. Zhang, L.Y., et al., *Fibroblasts play a potential role in bone destruction via osteopontin related caldesmon expression and polymerization in human non-functioning pituitary adenomas.* Sci Rep, 2017. **7**(1): p. 17523.

52. Salasznyk, R.M., et al., *Comparing the protein expression profiles of human mesenchymal stem cells and human osteoblasts using gene ontologies.* Stem Cells Dev, 2005. **14**(4): p. 354-66.

53. Sutermaster, B.A. and E.M. Darling, *Considerations for high-yield, high-throughput cell enrichment: fluorescence versus magnetic sorting.* Scientific Reports, 2019. **9**(1): p. 227.

54. Gygi, S.P., et al., *Correlation between Protein and mRNA Abundance in Yeast.* Molecular and Cellular Biology, 1999. **19**(3): p. 1720-1730.

55. Vogel, C. and E.M. Marcotte, *Insights into the regulation of protein abundance from proteomic and transcriptomic analyses.* Nat Rev Genet, 2012. **13**(4): p. 227-32.

56. Koussounadis, A., et al., *Relationship between differentially expressed mRNA and mRNA-protein correlations in a xenograft model system.* Sci Rep, 2015. **5**: p. 10775.

Supplementary figure 1. Use of lineage biomarkers to characterise Drop-Seq data clusters from CD45-/CD146+ skeletal progenitor population, CD144+ endothelial cells, and CD144-/CD106+ pericytes into 9 cell types.

Supplementary figure 2. Use of lineage biomarkers to characterise Drop-Seq data clusters from Stro-1+ cells, and cells enriched using SNAs targeting CD200, SPARCL1 and CD146 mRNAs into 8 cell types.