Title: Nutritional status and fecundity are synchronised by muscular exopheresis

Authors: Michał Turek[1,2*], Małgorzata Piechota[3†], Nilesh Shanmugam[3†], Marta Niklewicz[3],

Konrad Kowalski[3], Agnieszka Chacińska[1,2], Wojciech Pokrzywa[3*]

Affiliations:

[1]*Laboratory of Mitochondrial Biogenesis, Centre of New Technologies, University of*

Warsaw, Warsaw, Poland.

[2]*ReMedy International Research Agenda Unit, University of Warsaw, Warsaw, Poland.*

[3]*Laboratory of Protein Metabolism in Development and Aging, International Institute of*

Molecular and Cell Biology, Warsaw, Poland.

* Co-corresponding authors

[†] These authors contributed equally to this work

Correspondence should be directed to:

WP: wpokrzywa@iimcb.gov.pl

MT: m.turek@cent.uw.edu.pl

Abstract

Organismal functionality and reproduction depend on metabolic rewiring and balanced energy resources. However, the crosstalk between organismal homeostasis and fecundity, and the associated paracrine signaling mechanisms are still poorly understood. Using the *Caenorhabditis elegans* we discovered that secretory vesicles termed exophers, attributed in neurons to the removal of neurotoxic components, are released by body wall muscles to support embryonic growth. We found that exopher formation (exopheresis) is a non-cell autonomous process regulated by egg formation in the uterus. Our data suggest that exophers serve as transporters for muscle-generated yolk proteins used for nourishing and improving the growth rate of the next generation. We propose that the primary role of muscular exopheresis is to stimulate the reproductive capacity, thereby influencing the adaptation of worm populations to the current environmental conditions.

Main

The proper cellular function relies on the removal of unwanted contents by proteolysis or degradation mainly via the ubiquitin-proteasome system (UPS) and autophagy[1]. Recently a complementary mechanism was described in *Caenorhabditis elegans*. Under proteotoxic stress, worms' neurons can remove protein aggregates and damaged mitochondria via large membrane-surrounded vesicles called exophers[2]. Waste excretion by exophers in mouse neuronal HT22 cells under oxidative stress conditions was also reported[3]. This extrusion phenomenon could constitute a significant but still poorly explored metabolic waste management pathway. However, not only neurons can remove cellular content via exophers. Using worms expressing fluorescent reporters in the body wall muscle cells (BWM), we identified muscle-derived exophers; these reporters allow simultaneous tracking of proteasomes, closely associated with proteostasis, and mitochondria, which can be extruded in

neuronal tissue[2]. Muscular exophers are filled with proteasomes (both 19S and 20S subunits)

(**Fig. 1a**), and approximately 12% of them contain mitochondria (**Extended Data Fig. 1**).

Muscular exophers are generally more abundant and prominent (with diameters ranging from

2 to 15 μm) (**Fig. 1b**) than neuronal exophers[2], and the majority are generated by adult

hermaphrodite mid-body muscles (**Fig. 1c**). Exophers are formed in the muscle cell and

expelled outside via a pinching off mechanism (**Fig. 1d**). Some exophers that bud off remain

connected with the extruding BWM via a thin but elastic tube that permits continued transfer

of a large amount of cellular material into the extruded vesicle (**Fig. 1e, Supplementary Video**

1), similar to neuronal exophers[2]. Proteostasis impairment significantly increases neuronal

exopher output[2,3]. In contrast, the number of muscular exophers did not change in response to

depletion of the central proteostasis transcription factor HSF-1 (via *hsf-1* RNAi) or to heat

stress, and the number increased slightly under conditions of oxidative stress (**Fig. 1f-g**). These

observations suggest that proteostasis regulation might be not the core function of muscle

exopheresis.

Next, we assessed the number of exophers at different time points of the *C.*

elegans hermaphrodite life cycle. Reminiscent of neuronal exophers, muscular exophers are

not produced during the larval stages, and their maximum level is reached around the second

and third days of hermaphrodite adulthood (**Fig. 2a**). Because this time point coincides with

the worm's maximum reproductive rate, we wondered if reproduction could influence exopher

formation. To examine this possibility, we followed exopheresis in males. For the first three

days of adulthood, males did not produce any exophers (**Fig. 2a**). This finding suggests that

germ cell maturation in the reproductive system of hermaphrodite worms, the process of oocyte

fertilization, or embryonic development might regulate muscle exopheresis.

To test these hypotheses, we took advantage of a thermosensitive *C. elegans fem-1* mutant

strain that does not produce viable sperm at the restrictive temperature of 25 °C. At the

permissive temperature of 15 °C, some animals can reproduce as well as wild type

hermaphrodites, whereas the rest of the population is sterile[4]. The offspring-producing *fem-*

1 mutant animals grown at 15 °C generated a high number of muscular exophers. Contrarily,

animals raised either at 15 °C or 25 °C that were unable to fertilize oocytes did not activate

muscular exopheresis (**Fig. 2b-c**), indicating that neither the presence of the female gonad nor

the temperature itself were sufficient to trigger exopher release. Notably, hermaphrodites

sterilized via fluorodeoxyuridine (FUdR) treatment[5] extruded no exophers or only a few per

animal (**Fig. 2d**), suggesting that the occurrence of developing embryos could indeed stimulate

muscular exopheresis. We also found that FUdR-treated hermaphrodites often contained

exopher-like structures in their BWM (**Fig. 2e**, middle and right panels). Interestingly, we

detected objects resembling non-extruded exophers in the BWM of males (**Fig. 2e**, left panel),

indicating that males are devoid of mechanisms triggering their expulsion.

The above results suggest that the occurrence of developing embryos could induce muscular

exopheresis. Consistently, we observed a positive correlation between the number of exophers

released and the number of eggs present in the worm uterus (**Fig. 3a**). To further explore this

link, we depleted the mRNA of genes responsible for various processes associated with egg-

laying. RNAi depletion of the G-protein signalling gene *goa-1* leads to hyperactive egg-laying

behaviour, resulting in the presence of fewer early-stage eggs within the uterus[6]. As expected,

goa-1 knockdown caused a significant drop in the level of exopher release. In contrast, the egg-

laying defects induced by *egl-1* and *egl-4* RNAi, which lead to egg retention in the uterus[7],

increased exopher formation by muscle cells (**Fig. 3b**). In the absence of food, worms halt egg-

laying and retain fertilized eggs in the uterus[8,9]; therefore, we anticipated that worms would

generate more exophers when experiencing a food shortage. Indeed, the accumulation of

developing eggs in the uterus caused by the transfer of adult worms to food-free plates resulted

in a significant increase in muscle exopher secretion (**Fig. 3c**). Next, we tested whether worm

embryos could directly induce exopher production. We exposed young adult worms to extract

from developing eggs from the wild-type strain (**Fig. 3d**). Intriguingly, worms placed in contact

with material derived from lysed embryos increased exopheresis by approximately 20% (**Fig.**

3e). This observation suggests that molecules that diffuse from embryos *in utero* are

responsible for exopheresis induction. Finally, we decided to confirm that even in the face of

disturbed proteostasis specifically in the BWM, a signal associated with the developing

embryos in the uterus would be the primary regulator of muscular exopheresis. To this end, we

knocked down the myosin-directed chaperone UNC-45, which results in severe defects in the

organization of thick muscle filaments, as well as embryonic defects in cytokinesis and polarity

determination[10-12]. We initiated *unc-45* depletion in L4 larvae, which first leads to a disturbance

of proteostasis in BWM, and later, in young gravid adults, to inhibition of embryonic

development. Despite the dysfunction of a myosin-chaperone network in muscle cells, which

was indicated by complete paralysis of worms, we observed a dramatic inhibition of exopher

formation (**Fig. 3f**), highlighting the predominant role of maturating eggs in muscular

exopheresis.

Production of a single exopher involves the elimination of a substantial amount of the muscle

cell content. The high number of exophers produced during the *C. elegans* lifespan by BWM

leads to the removal of a relatively significant portion of cellular mass by a single tissue. We

hypothesized that this process should have substantial effects on worm muscle functionality

and healthspan. To this end, we selected three types of worms from the synchronized

population of gravid adults based on the number of extruded exophers, i.e., few (< 2), many (>

20), and control (6-8) animals (**Fig. 4a**), and analyzed their locomotory behaviour. In neurons,

exophers production is correlated with improved cell functionality; however, worms with

intensified exopheresis did not show enhanced mobility (**Fig. 4b**, **Extended Data Fig. 2**). On

the contrary, these animals displayed a reduction in exploratory behaviour (**Fig. 4c-d**).

Furthermore, the active use of muscular exopheresis was correlated with prolonged lifespan,

revealing a long-term beneficial effect (**Fig. 4e**).

Because, developing embryos stimulate exopher release from the mother BWM, we next

wondered whether this process benefited the offspring. Previous work showed that neuronal

exopher content can be transported through the worm body to reach distant scavenger cells

(coelomocytes)[2]. Moreover, muscle-specific transcriptomic analysis revealed the presence of

significant levels of vitellogenin mRNAs (i.e., *vit-2*, *-5* and *-6*)[13]. Therefore, we hypothesized

that the components of the yolk from BWM are transported through exophers to be used as a

source of raw materials for maturating eggs. To address this possibility, we used RNAi

knockdown of *vit-1* (a vitellogenin-coding gene) to reduce the level of the principal yolk

protein[14] and investigated whether it led to an increase in exopher biogenesis as a possible

compensatory mechanism. Indeed, the number of accumulated muscle-released exophers

nearly doubled in response to yolk protein depletion (**Fig. 4f**). Moreover, intensification of

exopheresis in the mother increased the amount of vitellogenin in developing eggs (**Fig. 4g**).

Next, we followed the localization of vitellogenin-2 (VIT-2) fused to GFP and muscle exopher

markers in the hermaphrodite. We detected the presence of VIT-2::GFP in the BWM of day-2

adult worms, as well as a significant accumulation in many muscle exophers (**Fig. 4h,**

Extended Data Fig. 3). These results suggest that exophers can mediate the transport of

additional portions of muscle-produced vitellogenin, which is ultimately deposited in oocytes

from the body cavity[15]. Finally, we followed the growth of the progeny of hermaphrodites

exhibiting different levels of exopheresis (**Fig. 4a**). We found that offspring from mothers with

a high number of muscle-released exophers (which showed no change in egg-laying behaviour)

grew faster (**Fig. 4i**). This is in line with a previous reports showing that yolk-reach eggs

support animals post-embryonic survival and development[14,16,17].

Discussion

We have shown here that muscular exopheresis in *C. elegans* represents a previously

uncharacterized nutrient management program associated with nourishing the next generation

of worms. The availability of food for mothers affects the number of developing eggs in the

uterus and the developmental stage at which the eggs are laid[18]. Activation of exopheresis

occurs with the first appearance of developing eggs in the uterus and intensifies in situations

of environmental food depletion. Likewise, disturbance of yolk synthesis increases exopher

biosynthesis. Consequently, in mothers with highly active exopheresis, the volume of yolk

content in eggs increased. Our results show that yolk protein produced in BWM is transferred

to exophers and ultimately delivered to oocytes. Therefore, the use of exophers for the transport

of vitellogenin represents an elegant mechanism by which remote cells can help to enrich the

nourishment of developing embryos. This process leads to the production of larvae that are

better prepared to thrive in the current environmental conditions. Thus, muscular exopheresis

is likely an adaptive mechanism that affects the dynamics of population growth. The impact of

exopheresis on early reproduction may be particularly crucial for wild worms, given the

significantly shortened life expectancy observed under more natural conditions[19].

Neuronal exopher formation occurs in higher animals[3]. In addition, yolk protein can be

synthesized in the muscles of oviparous animals like zebrafish[20], and is supplemented from the

mother to the intraovarian embryo in viviparous animals[21]. Hence, it is tempting to speculate

that the role of muscular expheresis in supporting progeny development could be evolutionarily

conserved. However, the exact mechanisms by which oocyte fertilization and subsequent

embryonic development initiate exopher formation and how this exopheresis is executed at the

molecular level require further studies.

References

Dikic, I. Proteasomal and Autophagic Degradation Systems. *Annu Rev Biochem* **86**, 193-224, doi:10.1146/annurev-biochem-061516-044908 (2017).

Melentijevic, I. *et al.* C. elegans neurons jettison protein aggregates and mitochondria under neurotoxic stress. *Nature* **542**, 367-371, doi:10.1038/nature21362 (2017).

Hualin, F., Jilong, L., Peng, D., Weilin, J. & Daxiang, C.   (bioRxiv, 2019).

Nelson, G. A., Lew, K. K. & Ward, S. Intersex, a temperature-sensitive mutant of the nematode Caenorhabditis elegans. *Dev Biol* **66**, 386-409, doi:10.1016/0012-1606(78)90247-6 (1978).

Hosono, R. Sterilization and growth inhibition of Caenorhabditis elegans by 5-fluorodeoxyuridine. *Exp Gerontol* **13**, 369-374, doi:10.1016/0531-5565(78)90047-5 (1978).

Bany, I. A., Dong, M. Q. & Koelle, M. R. Genetic and cellular basis for acetylcholine inhibition of Caenorhabditis elegans egg-laying behavior. *J Neurosci* **23**, 8060-8069 (2003).

Hirose, T. *et al.* Cyclic GMP-dependent protein kinase EGL-4 controls body size and lifespan in C elegans. *Development* **130**, 1089-1099, doi:10.1242/dev.00330 (2003).

Daniels, S. A., Ailion, M., Thomas, J. H. & Sengupta, P. egl-4 acts through a transforming growth factor-beta/SMAD pathway in Caenorhabditis elegans to regulate multiple neuronal circuits in response to sensory cues. *Genetics* **156**, 123-141 (2000).

Dong, M. Q., Chase, D., Patikoglou, G. A. & Koelle, M. R. Multiple RGS proteins alter neural G protein signaling to allow C. elegans to rapidly change behavior when fed. *Genes Dev* **14**, 2003-2014 (2000).

Kachur, T., Ao, W., Berger, J. & Pilgrim, D. Maternal UNC-45 is involved in cytokinesis and colocalizes with non-muscle myosin in the early Caenorhabditis elegans embryo. *J Cell Sci* **117**, 5313-5321, doi:10.1242/jcs.01389 (2004).

Gazda, L. *et al.* The myosin chaperone UNC-45 is organized in tandem modules to support myofilament formation in C. elegans. *Cell* **152**, 183-195, doi:10.1016/j.cell.2012.12.025 (2013).

Pokrzywa, W. & Hoppe, T. Chaperoning myosin assembly in muscle formation and aging. *Worm* **2**, e25644, doi:10.4161/worm.25644 (2013).

Blazie, S. M. *et al.* Comparative RNA-Seq analysis reveals pervasive tissue-specific alternative polyadenylation in Caenorhabditis elegans intestine and muscles. *BMC Biol* **13**, 4, doi:10.1186/s12915-015-0116-6 (2015).

Perez, M. F. & Lehner, B. Vitellogenins - Yolk Gene Function and Regulation in. *Front Physiol* **10**, 1067, doi:10.3389/fphys.2019.01067 (2019).

Hall, D. H. *et al.* Ultrastructural features of the adult hermaphrodite gonad of Caenorhabditis elegans: relations between the germ line and soma. *Dev Biol* **212**, 101-123, doi:10.1006/dbio.1999.9356 (1999).

Van Rompay, L., Borghgraef, C., Beets, I., Caers, J. & Temmerman, L. New genetic regulators question relevance of abundant yolk protein production in C. elegans. *Sci Rep* **5**, 16381, doi:10.1038/srep16381 (2015).

Perez, M. F., Francesconi, M., Hidalgo-Carcedo, C. & Lehner, B. Maternal age generates phenotypic variation in Caenorhabditis elegans. *Nature* **552**, 106-109, doi:10.1038/nature25012 (2017).

18 Muschiol, D., Schroeder, F. & Traunspurger, W. Life cycle and population growth rate
of Caenorhabditis elegans studied by a new method. *BMC Ecol* **9**, 14,
doi:10.1186/1472-6785-9-14 (2009).
19 Van Voorhies, W. A., Fuchs, J. & Thomas, S. The longevity of Caenorhabditis elegans
in soil. *Biol Lett* **1**, 247-249, doi:10.1098/rsbl.2004.0278 (2005).
20 Zhong, L. *et al.* Distribution of vitellogenin in zebrafish (Danio rerio) tissues for
biomarker analysis. *Aquat Toxicol* **149**, 1-7, doi:10.1016/j.aquatox.2014.01.022 (2014).
21 Iida, A. *et al.* Mother-to-embryo vitellogenin transport in a viviparous teleost. *Proc Natl*
Acad Sci U S A **116**, 22359-22365, doi:10.1073/pnas.1913012116 (2019).
220

Figures

Figure 1. *C. elegans* **muscles expel cellular content via exophers. a**, Muscular exophers contain organelles and large protein complexes. Arrows: white – exopher, blue – muscle cell, green – mitochondria, red – proteasome foci, yellow – unidentified vesicle. MOM – mitochondrial outer membrane. **b**, BMW actively releases significant amounts of exopher. The image shows the middle part of the worm's body with muscles marked with dashed lines. Arrows indicate representative exophers, and the asterisk indicates the position of the vulva. **c**, Production of muscular exophers is not evenly distributed across all muscle cells. The highest number of exophers is produced by the muscles adjacent to the vulva. n = 46 animals; three biological replicates. **d**, Exophers are formed via a pinching-off mechanism. Arrows: white – exopher, blue – muscle cell, red – distorted muscle cell membrane during exopher formation. **e**, Exophers may remain connected to the sending BWM cells via thin elastic tubes that allow further transfer of cellular material. Arrows: white – exopher, blue – muscle cell, red – cellular material transferred to exopher via elastic tube. **f**, Proteostasis disruption by *hsf-1* knockdown does not increase exopher production. n = 60 and 55 animals, two biological replicates. **g**, Proteostasis induced via oxidative stress but not via heat stress increases exopher production. n = 30 animals; three biological replicates.

Scale bars are 10 μm. Data are shown as mean ± SEM. ns – not significant, ** P < 0.01, **** P < 0.0001; **c**, **f**, **g**, two-tailed Welch's t-test.

Figure 2. Exopher formation is sex-specific and fertility-dependent. a, The highest number of exophers is produced during the hermaphrodite reproductive period and in aging animals. Males do not produce exophers during the first days of adulthood and begin to generate a small number of exophers later in life. Starting n = 90 hermaphrodites and 150 males; three biological replicates. **b**, Feminized hermaphrodites of thermosensitive a *fem-1* mutant strain do not produce exophers regardless of growth temperature. n = 10 - 26 animals; one biological replicate. **c**, Representative images of the middle part of the worm body in panel b. **d**, Hermaphrodites sterilized via FUdR treatment produce no exophers or only a few per animal. n = 17 and 20 animals; two biological replicates. **e**, Males and sterile hermaphrodites (via FUdR treatment) show the formation of spherical structures in the BWM that resemble mature exophers. MOM – mitochondrial outer membrane.

Scale bars are 10 μm. Data are shown as mean ± SEM; *** P < 0.001, **** P < 0.0001; **b**, two-tailed Welch's t-test.

Figure 3. Muscular exopheresis is a non-cell autonomous process regulated by *in utero* developing embryos. a, The number of produced exophers positively correlates with the number of *in utero* embryos. The violet line is a linear regression line, and each orange point represents one animal. All animals were 1 - 3 days old; n = 54 animals; three biological replicates. **b**, RNAi knockdown of genes regulating the egg laying rate and their presence in the uterus influences exopher production. n = 50 - 60 animals; two biological replicates. **c**, Egg retention in the uterus caused by starvation increases exopher production. n = 20 animals; one biological replicate. **d**, Schematic representation of the experimental setup for investigating the influence of egg lysate on exopher production. **e**, Exposure of young-adult worms to embryo-derived substances increases exopher production. n = 17 and 20 animals; one biological replicate. **f**, RNAi knockdown of the essential muscular co-chaperone *unc-45* reduces exopher production. n = 28 and 17 animals; two biological replicates.

Data are shown as mean ± SEM; * P < 0.05, ** P < 0.01, **** P < 0.0001; **b**, **c**, **f**, two-tailed Welch's t-test; **e**, one-tailed Welch's t-test.

Figure 4. Worms with overactive muscular exopheresis live longer and their offspring develop faster. a, Schematic representation of the experimental setup for investigating the influence of overactive exopheresis on F0 and F1 worm generation. **b**, Exopher production does not improve muscle functionality. n = 11 - 38; three biological replicates. **c-d**, Animals with overactive exopheresis have reduced exploratory behaviour presented as a reduction in straight line distance travelled (c) and by representative worms traces on the plate (d). n = 11 - 34; three biological replicates. **e**, Production of a high number of exophers increases lifespan. n = 63 - 88 animals; three biological replicates. **f**, RNAi knockdown of the egg yolk precursor protein VIT-1 increases the number of muscular exophers. n = 20 animals; two biological replicates. **g**, Embryos from hermaphrodite mothers that produce a high number of exophers contain more egg yolk precursor protein VIT-2. Representative images of embryos with VIT-2::GFP levels from mothers with different exopheresis activity are show above the graph. n = 15 and 14 eggs, five and six biological replicates (animals). **h**, Muscle-produced VIT-2 is released from muscles via exophers. The image shows the midbody of worms expressing the proteasome subunit RPN-5 tagged with RFP in BWM and VIT-2::GFP endogenous expression. Arrows: white – exopher, blue – muscle cell. Dashed lines mark eggs present in the uterus. Scale bar is 10 μm. **i**, Offspring of worms with overactive exopheresis develop faster. YA – young adult stage, L4 – last larval stage. n = 317 – 372 animals; two biological replicates. Data are shown as mean ± SEM. * P < 0.05, ** P < 0.01, *** P < 0.001, **** P < 0.0001; **e**, Log-rank (Mantel-Cox) test; **f**, **g**, two-tailed Welch's t-test; **i**, Fisher's exact test.

Extended Data Figure 1. Small fraction of muscular exophers contain mitochondria. n = 60; six biological replicates. Data are shown as mean ± SEM.

Extended Data Figure 2. Active exophoresis affects worm mobility. Graph shows the track lengths of animals exhibiting different exopheresis activities. n = 11 - 38; three biological replicates. Data are shown as mean ± SEM.

Extended Data Figure 3. Exophers contain muscle-produced vitellogenin. Images show the formation of an exopher filled with the proteasome and vitellogenin. Images were captured before and after RPN-5::RFP photobleaching, confirming that the high signal from VIT-2::GFP in forming exophers is not an imaging artefact. The white circle marks the area that

304   was bleached using a 555 nm laser and the position of developing exopher. Dashed lines mark

305   different tissue borders: m – muscle, i – intestine, g – gonad. Scale bar is 10 µm.

Methods

Data reporting

No statistical methods were used to predetermine the sample size. The experiments were not randomized. The investigators were not blinded to allocation during experiments and outcome assessment except for lifespan measurement and exploratory behaviour experiments.

Worm maintenance and strains

Worms were maintained on Nematode Growing Medium (NGM) plates seeded with OP50 *Escherichia coli* bacteria at 20 °C unless otherwise stated[1]. A list of all strains used in the study, together with the information in which experiments they were used is provided in Supplementary Table 1.

Generation of plasmids

All constructs were cloned using the SLiCE method[2] and were sequenced for verification. Construct for the expression of GFP on the mitochondrial outer membrane in the worm's body wall muscles was prepared as follows. First, destination vector pMT26 containing pCG150 vector backbone with the *myo-3* promoter and *unc-54* 3'UTR sequences separated by KspAI restriction site was made. Next, codon-optimized sequences[3] for GFP, linker containing attB5 sequence, and the sequence of the first 55 amino acids from TOMM-20 protein[4] (from a plasmid that was synthesized for this study) were PCR amplified and inserted into linearized pMT26 vector. To generate construct for expression of RPN-5 or PAS-7 tagged with RFP in worm's body wall muscles, sequences encoding respective proteins were PCR amplified and inserted into pMT26 linearized destination vector. As a template for the *rpn-5*-coding sequence, we used plasmid bearing codon-optimized *rpn-5* cDNA with 3 artificial introns, which was synthesized for this study. The sequence of *pas-7* gene was directly amplified from

N2 worms genomic DNA, and the sequence of RFP (wrmScarlet) was amplified from pSEM89

plasmid[5].

Transgenic strains generation

Worms transgenic strains were created by microparticle bombardment using *unc-*

119(ed3) rescue as a selection marker[6]. After phenotypic confirmation of successful plasmid

insertion, transformants were backcrossed two times against N2 strain to remove the *unc-*

119(ed3) background.

Scoring exophers and Microscopy

For scoring of exophers, a confocal microscope or a stereomicroscope was used.

When using the confocal microscope, animals were transferred onto 3 % agarose pads prepared

in H_2O and formed on a microscope slide. Next, animals were immobilized on the pad using 6

μL of PolySciences 0.05 μm polystyrene microspheres or 25 μM tetramisole and covered with

a glass coverslip. Immediately afterward, animals were imaged using an inverted Zeiss

LSM800 laser-scanning confocal microscope with 40x or 63x oil immersion objectives. The

488 nm and 561 nm lasers were used for excitation of GFP and RFP fluorescent proteins,

respectively. Z-stacks, which covered the whole animal, were taken and the number of

exophers released by muscles was counted. Finally, data were normalized to the average

number of exophers present in control animals and compared between conditions.

For scoring exophers with the stereomicroscope, we used Leica M165FC stereomicroscope

equipped with Leica EL6000 lamp and standard Texas Red and GFP filter sets. Age

synchronized, freely moving, day-2 adult animals were directly visualized on culturing NGM

plates and the number of visible exophers released by muscles were counted. Finally, data were

normalized to the average number of exophers present in control animals and compared

between conditions.

Representative pictures of exophers presented in the manuscript were taken with the usage of

an inverted Zeiss 700 laser-scanning confocal microscope equipped with a 40x oil objective.

The 488 nm and 555 nm lasers were used for excitation of GFP and RFP fluorescent proteins,

respectively. To investigate the presence and distribution of exophers, z-stacks were taken and

processed with ZEN software.

RNA interference

RNA interference in *C. elegans* was performed using the standard RNAi feeding method and

RNAi clones[7]. For experiments, we used NGM plates supplemented with 1 mM IPTG and 25

µg/µl carbenicillin seeded with HT115 *E. coli* bacteria expressing double-stranded RNA

(dsRNA) against the gene of interest or, as a control, with bacteria without a vector. Worms

were placed on freshly prepared RNAi plates, either as age-synchronized pretzel-stage eggs,

L1 larvae, or L4 larvae. The number of exophers was counted on day 2 adult worms using a

confocal microscope or a stereomicroscope.

Stress influence on exophers production

Worms were age-synchronized using alkaline hypochlorite solution (bleaching procedure), as

previously described[8]. The harvested eggs were incubated overnight at 16 ˚C for hatching.

Approx. 1000 L1 larvae were transferred on NGM plates and incubated at 20 °C till they

reached day-2 of adulthood. The worms were further channeled to the respective stress

treatments.

Oxidative stress

Approx. 100, day-2 adult worms were washed-off from the NGM plates and rinsed 3 times

with M9 buffer. The worms to be stressed were suspended in 1 ml of 5 mM hydrogen peroxide

solution prepared in M9 buffer, whereas control worms were suspended in M9 buffer. The

tubes were incubated on a shaker at 20 ˚C for 60 minutes.

Heat stress

Similarly, approx. 100, day-2 adult worms were washed-off form NGM plates and rinsed 3

times with M9 buffer. The worms were further suspended in a 1 ml M9 buffer. The worms to

be heat stressed were incubated in a shaker at 33 °C for 60 minutes, whereas the control animals

were incubated at 20 °C for 60 minutes.

Exophers quantification

From each stress/control treatment, 30 worms were picked onto agarose pad slides individually

for the quantification of exophers. Exophers were quantified using a confocal microscope.

Obtained data were normalized to the average number of exophers present in control animals

and compared between conditions.

Number of exophers on consecutive days

For each biological replicate, 30 L4 larvae hermaphrodites or 50 L4 larvae males were

transferred to fresh NGM plates (5 hermaphrodites or 10 males per plate). For the next

consecutive 15 days, the number of exophers in each animal was counted using a

stereomicroscope. Hermaphrodites were transferred to fresh plates every 2 – 3 days. Males

were kept on the same plate until the end of the experiment. All animals that died during the

experiment time course were removed from the plate.

Measuring exophers in *fem-1* mutant

Approx. 50 L1 larvae from *fem-1(hc17)ts* mutant strain expressing GFP in BWM were

transferred to 2 fresh NGM plates. During the time course of the experiment, one of the plates

was kept at 15 °C while the second one was kept at 25 °C. When worms reached L4 larval

stage, each worm was transferred to a separate plate and grown at the same temperature as

before. After 48 hours at 25 °C or 72 hours at 15 °C, using a stereomicroscope, a number of

exophers released from worms muscles was counted and plates were scored for F1 offspring

to assign all worms as fertile or infertile.

FUdR assay

Age-synchronized animals were placed on NGM plates seeded with OP50 *E. coli* bacteria as a

food source until they reached young adulthood (day 0). Next, animals were selected and

moved to test plates containing 25 μM fluorodeoxyuridine (FUdR) to prevent embryonic

development and egg hatching[9] or control plates without FUdR. The number of exophers was

counted on adult day-2 using confocal microscopy.

Exophers and *in utero* eggs correlation

For correlating the number of exophers with the number of eggs present in worms uterus, 1 to

3-days old animals were used. First, muscular exophers for a single worm were counted using

the stereomicroscope. Next, worm was transferred to a 10 μl drop of 1.8 % hypochlorite

solution put on a microscope slide. Finally, after approx. 5 minutes when the hermaphrodite

mother was fully bleached, the number of eggs released from worm's uterus was counted. Data

analysis was performed using GraphPad Prism 8 software.

Starvation assay

To assess the influence of worms starvation on the exophers production, day-2 adult worms

were moved to bacteria-free NGM plates. After 24 hours of food deprivation, the number of

exophers was counted using a stereomicroscope. As a control, day-3 adult worms grown for

the whole time on NGM plates seeded with bacteria were used. Obtained data were normalized

to the average number of exophers present in control animals and compared between

conditions.

Eggs lysate assay

To obtain eggs, age-synchronized N2 gravid hermaphrodites were bleached using an alkaline

hypochlorite solution. Harvested eggs were suspended in appropriate volume of M9 buffer to

reach the concentration of approx. 200 eggs per µl. Next, eggs were flash-frozen in liquid

nitrogen, thawed on ice, and sonicated 3 times for 10 seconds to obtain eggs lysate. 150 µl of

eggs lysate was mixed with 150 µl of concentrated OP50 *E. coli* bacteria and placed in a 0.5

ml Eppendorf tube. Approx. 30 day-1 young adult hermaphrodites were transferred to

Eppendorf tube containing a mixture of eggs lysate and bacteria and were placed on the rotator

for 2 hours at room temperature. After 2 hours, the content of the tube was placed on the fresh

NGM plate seeded with OP50 bacteria. 18 hours later the number of exophers in each worm

was counted. For control animals, the whole protocol was the same except that for bleaching

step no worms were used.

Worm exploratory behaviour

Age-synchronized, day-2 adult worms which had an average (6-8), few (< 2), or many (> 20)

exophers were sorted to separate NGM plates. Approximately 10 worms per replicate were

brought onto NGM plates and the worm movement was recorded for 2 minutes using the

WormLab system (MBF Bioscience). The frame rate, exposure time, and gain were set to 7.5

frames per second, 0.0031 s, and 1, respectively. The track length, straight-line distance, center

point speed, and the overall track pattern of individual worms were analyzed using the

WormLab software (MBF Bioscience).

Lifespan assay

For each biological replicate, approx. 300 late L4 larvae were transferred to fresh NGM plate.

On day 2 of worms' adulthood, the number of exophers in each animal was counted and worms

with less than 2 or more than 20 visible exophers were transferred to new plates (25 – 30

animals). Additional 25 – 30 worms were picked blindly from the same original population,

moved to a fresh plate, and served as a control. Lifespan measurements were performed on

these three respective populations. During lifespan measurements, worms were scored daily

for movement and pharyngeal pumping until their death. All worms were moved to fresh plates

every 1 – 2 days during the eggs-laying period and every 3 – 5 days afterward. Animals that

crawled off the plate or exhibited baggy phenotype were censored from the experiment. Data

analysis was performed using GraphPad Prism 8 software.

Vitellogenin levels in embryos

Vitellogenin levels in embryos were measured based on the GFP signal from fluorescently

tagged VIT-2 protein. Approx. 200 late L4 larvae were transferred to fresh NGM plate. On day

2 of worms adulthood, the number of exophers in each animal was counted and worms with

less than 2 or more than 20 visible exophers were transferred to new plates. Next, animals from

each group were individually transferred to a 10 µl drop of M9 buffer which was placed on a

microscope slide. Using a sharp injection needle, worm was cut open to release eggs from the

uterus. Fluorescent signal from 2-cell embryo stage was captured using Leica M165FC

stereomicroscope equipped with Leica EL6000 lamp, standard GFP filter set, and Leica

DFC365 FX CCD camera. Magnification used for recording pictures was set to 192x. Exposure

time and gain were set to 600 ms and 2, respectively. The fluorescent signal was quantified

using Leica Las X software and was normalized to the average signal from eggs, which were

obtained from animals with less than 2 exophers.

Worms development assay

25 – 30 age-synchronized, day-2 adult worms which had an average (6-8), few (< 2), or many

(> 20) exophers were sorted to separate NGM plates. Gravid adults were allowed to lay eggs

for 4 hours, afterward removed from the plates, and the development of their offspring was

followed. 46 hours later, using the stereomicroscope, developmental stage of each animal was

checked, and the proportion between L4 larvae stage worms and young adult worms was

calculated.

Statistical analysis

Statistical tests used in this study were two-tailed Welch's t-test, one-tailed Welch's t-test, Log-

rank (Mantel-Cox) test, and Fisher's exact test. P-value < 0.05 was considered significant.

Methods references

1.   Brenner, S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71–94 (1974).

2.   Zhang, Y., Werling, U. & Edelmann, W. SLiCE: a novel bacterial cell extract-based

DNA cloning method. *Nucleic Acids Res.* **40**, e55–e55 (2012).

3.   Redemann, S. *et al.* Codon adaptation-based control of protein expression in C.

elegans. *Nat. Methods* **8**, 250–252 (2011).

4.   Watanabe, S. *et al.* Protein localization in electron micrographs using fluorescence

nanoscopy. *Nat. Methods* **8**, 80–84 (2011).

5.   El Mouridi, S. *et al.* Reliable CRISPR/Cas9 Genome Engineering in

Caenorhabditis elegans Using a Single Efficient sgRNA and

an Easily Recognizable Phenotype. *G3 Genes|Genomes|Genetics* **7**, 1429 LP – 1437

(2017).

6.   Praitis, V., Casey, E., Collar, D. & Austin, J. Creation of low-copy integrated

transgenic lines in Caenorhabditis elegans. *Genetics* **157**, 1217–1226 (2001).

7.   Kamath, R. S. & Ahringer, J. Genome-wide RNAi screening in Caenorhabditis

elegans. *Methods* **30**, 313–321 (2003).

8.   Porta-de-la-Riva, M., Fontrodona, L., Villanueva, A. & Cerón, J. Basic Caenorhabditis

elegans methods: synchronization and observation. *J. Vis. Exp.* e4019–e4019 (2012).

doi:10.3791/4019

9.   Mitchell, D. H., Stiles, J. W., Santelli, J. & Sanadi, D. R. Synchronous Growth and

Aging of Caenorhabditis elegans in the Presence of Fluorodeoxyuridine1. *J. Gerontol.*

34, 28–36 (1979).

Acknowledgements

We thank the Caenorhabditis Genetics Center (funded by the NIH National Center for Research

Resources, P40 OD010440) for strains; T. Hoppe for expert advice and Carl Kutzner for RNAi

clones; T. Wegierski for confocal microscopy assistance; B. Uszczyńska-Ratajczak, K.

Szczepanowska, H. Bringmann, and members of Chacińska and Pokrzywa laboratories for

discussions and comments on the manuscript.

Funding

Work in the W.P. laboratory was supported by the Foundation for Polish Science co-financed by the European Union under the European Regional Development Fund (grant POIR.04.04.00-00-5EAB/18-00 to W.P.), the European Molecular Biology Organization (EMBO Installation Grant No. 3916 to M.P., K.K., and W.P.), the Polish National Science Center (grant UMO-2016/23/B/NZ3/00753 to N.S. and W.P.), and the Deutsche Forschungsgemeinschaft (DFG FOR 2743 to W.P.). Work in the A.C. laboratory was funded by "Regenerative Mechanisms for Health" project MAB/2017/2 (M.T. and A.C.) carried out within the International Research Agendas programme of the Foundation for Polish Science co-financed by the European Union under the European Regional Development Fund and was supported by a POLONEZ Fellowship of National Science Centre, Poland, 2016/21/P/NZ3/03891 (M.T.), within European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 665778.

Author contributions

M.T., W.P., M.P., N.S., M.N., and K.K. conducted and designed experiments, M.T. and W.P. conceived the project and supervised the study. W.P., A.C. and M.T. secured the funding. W.P. and M.T. wrote the manuscript with input from A.C. and M.P

Author Information

The authors declare no competing financial interests. Correspondence and requests for materials should be addressed to M.T., (m.turek@cent.uw.edu.pl) and W.P. (wpokrzywa@iimcb.gov.pl).

Supplementary Information

Supplementary Video 1

The video shows the cellular content being transfer from BWM cell to exopher via elastic tube.

Supplementary Table 1

The file contains a table with the list of all *C. elegans* strains used in this study.

Supplementary Table 2

The file contains a table with the statistical summary of lifespan experiments